The recovery of 12,789 genomes revealed the diversity, function, and microbial interactions of the geothermal spring microbiome

Yu-Xian Li[1,2,†], Yang-Zhi Rao[1,3,†], Ze-Wei Li[1], Shao-Peng Li[4], Yan-Ni Qu[1], Jia-Liang Kuang[5], Yong-Jian Chen[6,7], Yan-Ling Qi[1], Qi-Jun Xie[1], Jian-Yu Jiao[9], Wen-Sheng Shu[8], Wen-Jun Li[9,10,*], Zheng-Shuang Hua[1,*]

[1] Chinese Academy of Sciences Key Laboratory of Urban Pollutant Conversion, Department of Environmental Science and Engineering, University of Science and Technology of China, 230026 Hefei, PR China.

[2] Southern Marine Science and Engineering Guangdong Laboratory (Guangzhou), No. 1119 Haibin Road, Guangzhou 511458, China.

[3] Centre for Infection and Immunity Study (CIIS), School of Medicine (Shenzhen), Sun Yat-sen University, Shenzhen 518107, China.

[4] Zhejiang Tiantong Forest Ecosystem National Observation and Research Station, Institute of Eco-Chongming, School of Ecological and Environmental Sciences, East China Normal University, Shanghai, 200241, PR China.

[5] The Key Lab of Pollution Control and Ecosystem Restoration in Industry Clusters, Ministry of Education, School of Environment and Energy, South China University of Technology, Guangzhou, Guangdong 510006, RP China

[6] Department of Ecology, College of Natural Resources and Environment, and Guangdong Laboratory for Lingnan Modern Agriculture, Guangdong Provincial Key Laboratory of Eco-Circular Agriculture, South China Agricultural University, Guangzhou 510642, China

[7] Guangdong Engineering Technology Research Centre of Modern Eco-Agriculture and Circular Agriculture, Guangzhou 510642, China

[8] School of Life Sciences, South China Normal University, Guangzhou 510275, PR China

[9] State Key Laboratory of Biocontrol, Guangdong Provincial Key Laboratory of Plant Resources and Southern Marine Science and Engineering Guangdong Laboratory (Zhuhai), School of Life Sciences, Sun Yat-Sen University, Guangzhou 510275, PR China

[10] State Key Laboratory of Desert and Oasis Ecology, Xinjiang Key Laboratory of Biodiversity Conservation and Application in Arid Lands, Xinjiang Institute of Ecology and Geography, Chinese Academy of Sciences, Urumqi, PR

China

†These authors contributed equally to this work: Yu-Xian Li, Yang-Zhi Rao.

*Correspondence and requests for materials should be addressed to W.J.L. (email: liwenjun3@mail.sysu.edu.cn) and

Z.S.H (email: hzhengsh@ustc.edu.cn).

Authors' e-mail address list:

Y.X.L. (li_yuxian@gmlab.ac.cn)

Y.Z.R. (raoyzh7@mail.sysu.edu.cn)

Z.W.L. (lizw@mail.ustc.edu.cn)

S.P.L. (spli@des.ecnu.edu.cn)

Y.N.Q. (quyanni@ustc.edu.cn)

J.L.K. (kjialiang@scut.edu.cn)

C.Y.J. (chenyj@scau.edu.cn)

Y.L.Q. (qiyanling7@foxmail.com)

Q.J.X. (qjxie@mail.ustc.edu.cn)

J.Y.J. (jiaojy5@mail.sysu.edu.cn)

W.S.S. (shuwensheng@m.scnu.edu.cn)

W.J.L. (liwenjun3@mail.sysu.edu.cn)

Z.S.H. (hzhengsh@ustc.edu.cn)

Abstract

Geothermal springs, characterized by extreme physicochemical conditions, represent ecologically and evolutionarily

significant habitats that foster unique microbial communities and drive adaptive evolutionary processes. Despite their

importance, the complex microbial interactions and underlying mechanisms governing community assembly in these

environments remain poorly understood. In this study, we conducted systematic sampling across 49 geothermal

springs in Tengchong, Yunnan, over a six-year period (2016-2021), and performed metagenomic sequencing on 152

samples. We successfully reconstructed 12,789 non-redundant microbial genomes, revealing an exceptionally high

level of phylogenetic and functional diversity within the spring microbiomes. Our analyses demonstrate that pH and

temperature are the primary deterministic drivers shaping both microbial species composition and functional potential,

thereby segregating the communities into three distinct groups: acidic, hyperthermal, and thermal. Furthermore,

ecological network analysis revealed that extreme environmental conditions significantly alter network topology,

resulting in less complex but more efficient microbial interaction networks. Collectively, this study provides a

comprehensive resource and mechanistic insights into the microbial diversity, community structure, and species

interactions in geothermal spring ecosystems.

Introduction

Microorganisms are important participants in the biogeochemical process across all ecosystems on Earth[1–4].

Consequently, investigating microbial species and functional diversity is vital to understanding both microbial

community dynamics and the biogeochemical cycle within ecosystems. Geothermal springs, as typical high-

temperature terrestrial environments with distinct geochemical characteristics, form confined "islands" that harbor

specialized and potentially ancient microbes[5–7]. Previous cultivation studies and molecular surveys have revealed

geothermal springs featured bacteria such as Aquificae, Proteobacteria, Cyanobacteria, Chloroflexi, and Thermus[8,9];

as well as archaea such as Crenarchaeota, Korarchaeota, and Nanoarchaeota[8–11]. Due to their geological similarity to

early Earth and relatively simple microbial community structures, geothermal springs serve as ideal model systems

for understanding microbial community structure, function, interactions, and assembly processes in ancient

environments[8,10,12–14].   In particular, geothermal springs in Tengchong, Yunnan, China, which are magmatic-

hydrothermal systems with a legacy of Cenozoic volcanic activities[15], nurture a notable diversity of acidic geothermal

springs and unique microbial diversity compared to other high-temperature systems[15–17]. Despite these advances, the

microbial interactions in geothermal springs remained largely unexplored.

Disentangling ecological drivers that govern microbial community establishment and maintain biodiversity is

both essential and demanding. It is well documented that both biotic and abiotic factors influence the distribution of

microbes, as they interact not only with other sympatric microbes but also with the environments they reside in[18,19].

Constructing co-occurrence networks is an effective approach for biologically investigating relationships and

interactions among microorganisms within complex environments[20,21]. Interpreting the network structure through

various parameters and identifying key species and functional groups can provide insights into how these microbes

stabilize and support community establishment[22,23]. Previously, network inference largely relied on single-marker

gene-based amplicon sequencing. While this approach is simple and cost-effective, it has notable limitations, such as

limited taxonomic resolution, amplification biases, and a lack of functional information. The advent of massively

parallel DNA sequencing has revitalized microbial ecology, enabling more in-depth studies of microbial interactions

and ecological processes that govern the community assembly. This advancement allows for a more comprehensive

investigation of microbial communities at both the species and functional levels.

Based on the 12,789 MAGs reconstructed from in-depth genome-resolved metagenomic sequencing, here we

uncovered unexpectedly high microbial diversity in geothermal springs, spanning 110 archaeal and bacterial phyla,

with approximately 93% of identified genera being newly discovered. Strong environmental heterogeneity caused the

segregation of samples into three distinct groups, where the dominant microbes in each group were also the key

functional components. By considering the environmental physicochemical characteristics and the distribution

patterns of microbial communities, we performed an in-depth analysis integrating molecular ecological networks

(MENs). This integrative framework disclosed how biotic and abiotic factors sustain the biodiversity in geothermal

spring ecosystems. Our study offers comprehensive insights into the diversity, function, and microbial interactions in

hot spring ecosystems, bridging gaps in understanding these complex microbial species-functional dynamics.

Results and discussion

Diversity and novelty of the hot spring microbiome. A total of 152 samples covering 49 hot springs located in

Tengchong (Yunnan, China) were collected and metagenomically sequenced, spanning six years (2016-2021) (Fig.

S1a). The sampling sites displayed tremendous environmental heterogeneity across space and time in physiochemical

parameters (Fig. S1b, Supplementary Table 1), especially covering a wide range of temperatures (23-99°C) and pH

values (2.0-9.73) (Fig. S1c). With manual curation, we successfully reconstructed 20,849 MAGs from all 152

metagenomes (Fig. 1a). Among them, 5,990 (28.7%) high-quality (completeness $\geq$ 90%, contamination < 5%) and

6,799 medium-quality (completeness $\geq$ 50%, contamination < 10%) were kept for further analyses. Genomic

information statistics showed the high reliability of the final set of 12,789 MAGs (Fig. S2). Remarkably, our

tremendously diverse genome set was unexpectedly comparable to the genome set recovered from over 20-fold of soil

metagenomes (3,304) in a recent study[24]. Regarding genomes with completeness >90%, and contamination <5%, the

number of genomes in our study (5,990) even slightly surpassed (5,184). Taxonomic assignment based on the Genome

Taxonomy Database (GTDB r207)[25] identified 2,944 archaeal and 9,845 bacterial genomes (Supplementary Table 2),

clustering them into 3,176 non-redundant species at 95% average nucleotide identity (ANI)[26,27]. The phylogenetic tree

of these 3,176 RMAGs revealed unexpected diversity and novelty among microbes inhabiting the geothermal spring

ecosystem (Fig. 1b)[28]. All RMAGs spanned 12 archaeal and 98 bacterial phyla, covering 63.2% of all known microbial

phyla (GTDB r207). Among them, five phyla, including Thermoproteota, Patescibacteria, Bacteroidota, Chloroflexota,

and Proteobacteria, each contained more than 200 RMAGs. Surprisingly, the phylogenetic diversity of geothermal

spring samples also exceeded the phylogenetic diversity of soil at the phylum level (11 archaeal phyla and 88 bacterial

phyla) in a previous study[24]. This could be a result of the immense environmental heterogeneity in our geothermal

spring sampling sites (Fig. S1b and c), and the "nested" microbial diversity of non-extreme environments from the

long evolutionary history of the terrestrial geothermal spring microbiome[13]. Moreover, RMAGs in this study exhibit

remarkable taxonomic novelty, identifying two new phyla, 11 new classes, 62 new orders, 213 new families, 721 new

genera, and 1,558 new species which account for 85.1% of the total diversity of the species level (Fig. 1c). Overall,

this dataset represents a tremendous and comprehensive prokaryotic genomic resource from geothermal spring

ecosystems, revealing a large amount of uncharted "microbial dark matter".

rpS3 gene or 16S ribosomal RNA gene was often used to represent the composition of the microbial

community[26,22]. However, the short sequence length and potential misassignment of the *rpS3* sequence to the

associated bin limit the classification power of *rpS3*. In this study, we implemented a genome-coverage-based

approach to calculate the relative abundance of microbes (Fig. S1), directly linking community compositions with

genome-resolved microbial functions and thereby avoiding less accurate predictive methodologies (e.g., PICRUST)[29].

The reads mapped to 3,176 RMAGs covered an average of 88% of the entire microbial community across 152 samples

(Fig. 1d, Supplementary Table 3), indicating that reconstructed RMAGs in this study are highly representative of the

in situ microbial communities. Apparently, microbial community compositions differ at the phylum level (Fig. 2a,

Supplementary Table 4). Archaea were significantly predominant (Mann-Whitney U test, P < 2.2e-16, Supplementary

Table 5) in geothermal springs with ultra-low pH (< 5) or ultra-high temperature (> 75 °C), including DeReTiYan

(DRTY), GuMingQuan (GMQ), ShuiReBaoZha (SRBZ), ZiMeiQuan (ZMQ), and ZhenZhuQuan (ZZQ) (Fig. 2a).

When pH < 5, samples such as DRTY-3, DRTY-6, and DRTY-9, were dominated by Thermoplasmatota and

Thermoproteota (Fig. 2b). In hyperthermal geothermal springs (GMQ, SRBZ, ZMQ, and ZZQ), Thermoproteota and

Aquificota were the most dominant archaea and bacteria, respectively (Fig. 2b and c). As for the rest of the geothermal

springs with less extreme environmental conditions, communities were mainly composed of bacteria from a wide

variety of phyla, with Chloroflexota, Proteobacteria, Acidobacteriota, Bipolaricaulota, and Armatimonadota being the

most abundant (Fig. 2c). Despite discernible differences in community compositions among geothermal springs,

metabolic profiles based on KEGG Orthology (KO) categories remain remarkably consistent across all 152 samples

(Fig. 2d, Supplementary Table 6 and 7). Such patterns were coherent with previous notions that higher-level functional

turnover was relatively stable compared to changes in species compositions[30,31]. Principal Coordinates Analysis

(PCoA) using dissimilarity matrices of RMAG and KO abundance revealed that all samples were segregated into three

distinct groups based on the pH and temperature (Figs. 2e-h, Supplementary Table 1). In comparison, the microbial

communities varied little over temporal and seasonal scales (winter or summer) (Fig. S3). Therefore, the community

structure of geothermal springs showed greater variations over physiochemical changes than temporally, with pH and

temperature being the most vital environmental drivers. Additionally, although different samples remain consistent at

higher-level functional turnover (Fig. 2d), the lower-level functional genes were more sensitive to environmental

changes (Fig. 2h).

pH and temperature drove the species and functional community structure. To further unveil the major

environmental determinants shaping community structure, we examined the relationship between environmental

variables and microbial diversity and metabolic function using the Mantel test and redundancy analysis (RDA) (Fig.

3a, Fig. S4, Supplementary Table 8). Results showed that pH, Na^+, Ca^{2+}, temperature, and K^+ were the five most

prominent factors significantly correlated with community composition, with pH being the most deterministic.

Combined with the PCoA results, we, therefore, divided the 152 samples into three groups based on their pH and

temperature: 1) acidic group (AG): geothermal springs with pH ≤ 5; 2) hyperthermal group (HG): geothermal springs

with pH > 5 and temperatures > 75˚C; 3) thermal group (TG): geothermal springs with pH > 5 and temperatures ≤

75˚C (Supplementary Table 1). Such grouping pattern was further corroborated by the ANalysis Of Similarities

(ANOSIM) test (R=0.679, P=0.001, Fig. 3b, Fig. S5). Interestingly, principal component analysis (PCA) based on

environmental variables of these sites showed a similar grouping pattern (Fig. S6). This alignment suggests a strong

ecological relationship between environmental conditions and microbial life, where specific factors consistently shape

microbial communities across different samples. Comparisons of the three groups showed that microbial communities

with extreme physicochemical conditions (AG and HG) exhibited significantly lower species and functional diversity

(Fig. S7 and S8, Supplementary Table 9). Among the 3,176 RMAGs, only 351 were found to be shared among all

three groups (Fig. S9a). Most of them were classified into Thermoproteota, Chloroflexota, and Proteobacteria (Fig.

S9b). TG represents a relatively benign habitat and harbors the highest number of unique taxa (1,341), most of which

belong to bacterial phyla, including 197 from Proteobacteria, 151 from Patescibacteria, and 149 from Chloroflexota

(Fig. S9b). Although HG contains far fewer taxa, the majority (1,515 of the 1,598, or 98%) were also present in TG,

demonstrating that HG represents a subset of the taxa detected in TG. In contrast, Thermoplasmatota, Micrarchaeota,

Nanoarchaeota, and Thermotogota appeared to be more abundant in samples belonging to AG, while Aquificota were

favored in hyperthermal habitats, and Thermoproteota and Aenigmatarchaeota inhabited both AG and GH (Fig. 3c).

In comparison, other bacterial phyla such as KSB1, Myxococcota, and Desulfobacterota_B distributed more

abundantly in TG.

To better compare the intraspecific functional diversity of three geothermal spring groups, functional annotation

was conducted on all MAGs, classifying all genes in each genome family into core and accessory genes according to

their functions (see Methods, Supplementary Table 10). Among the three groups, TG had the highest number of core

and accessory gene families, followed by HG and AG (Fig. 3d). This pattern likely reflects the species diversity of the

three groups (Fig. S7), as higher microbial diversity typically corresponds to a larger gene pool. However, the

proportion of accessory gene families was significantly higher in AG, followed by HG and TG (Fig. 3e). It is well

documented that accessory genes are more dynamic, frequently entering and leaving the genome through mechanisms

such as horizontal gene transfer, which enhances genetic diversity and adaptability[32]. This dynamism is likely

associated with the need to acquire genes for adaptation to acidic or hyperthermal environments. By looking into the

detailed functions of accessory genes, AG and HG possessed significantly more genes involved in genetic information

processing (Fig. S10). Notably, approximately 73.2% (AG) and 74.5% (HG) of accessory gene families were

unannotated or unclassified (Supplementary Table 11). We conjecture that species in AG and HG may rely on these

unknown accessory genes to adapt to their extreme environments, necessitating further investigation into their

functions.

Species and functional network patterns differed in three environmental groups. Based on the relative

abundances of 3,176 RMAGs in all samples, Molecular Ecological Networks (MENs) were built separately for the

three groups to reveal the interactions among microorganisms[22,33,34] (Fig. 4a, b, and c). Generally, all three networks

had "non-random", "scale-free" and "small-world" properties and modular structures (Supplementary Table 12)[22]. The

network size (total number of nodes) and network connectivity (total number of links)[35,22] in the three MENs

demonstrated a decreasing trend as environmental conditions became harsher, with TG (343 nodes linked by 1,243

edges) having the largest MEN but AG (84 nodes linked by 321 edges) and HG (172 nodes linked by 553 edges)

possessing much smaller MENs. Despite having a smaller MEN, AG exhibited a higher average degree (or average

connectivity, i.e., average links per node), average clustering coefficient (the extent to which nodes are clustered), and

connectedness (the proportion of realized links in all possible ones), indicating a higher level of network clustering

within the community (Fig. 4d, e, and g, Supplementary Table 13). Therefore, the neutral process might be less

prevalent when pH decreases[35]. Compared to TG, the higher magnitude of local clustering and shorter average path

distance in AG and HG suggested a faster response to environmental disturbances in geothermal springs with

physiochemical extremes (Fig. 4e and f), which are also hallmarks of small-world networks. All MENs in the three

groups were highly modular, with 19 large modules (modules with $\geq$ 5 nodes) accounting for 88-97.6% of the nodes.

This demonstrated the higher stability and robustness of MENs in terrestrial geothermal ecosystems. It also reflected

that the entire community may be organized into distinct sub-communities, with possible functional specialization

occurring within each sub-community. This was especially true for the HG group, which exhibited the highest

modularity (Fig. 4b). Interestingly, the physiochemical extremes (low pH and high temperature) have a pronounced

effect on the composition of MENs. In AG, nodes with higher degrees were mostly assigned to archaea, including

Thermoplasmatota, Micrarchaeota, and Thermoproteota. These nodes exhibited strong interactions with those from

other phyla, such as Aenigmatarchaeota and Desulfobacterota (Fig. 4a, Supplementary Table 10). In HG, high-degree

nodes were mainly classified into both bacteria and archaea, including Thermoproteota, Aquificota, and Deinococcota,

with each module consistently dominated by microbes from the same phylum (Fig. 4b). This also corroborated the

presence of possible functional specialization within different modules. The MEN of TG showed higher microbial

diversity and complexity, with Bacteria playing more significant roles in the relatively benign environments (Fig. 4c).

To unravel the functional specialization performed by microbes in each module and potential functional

interactions among different modules, as well as to establish links between microbes at both the species and function

levels, we further constructed functional networks for the three groups based on the relative abundance of core

functional genes (KOs) (Fig. S11-13, Supplementary Table 14). The top 50 functional genes with the highest node

degree were selected to represent the core metabolic features of certain communities (Supplementary Table 15 and

16). Overall, the key metabolic processes of the three groups appeared to be markedly distinct. We observed that

dissimilarity sulfate reduction (DSR) and some carbon metabolism pathways, including tricarboxylic acid cycle (TCA),

pyruvate metabolism, and fermentation, played a significant role in AG, and sulfur oxidation and carbon fixation

pathways, such as reverse tricarboxylic acid cycle (rTCA), dicarboxylate/4-hydroxybutyrate or

dicarboxylate/hydroxybutyrate (DC-HB) cycle, and Wood-Ljungdahl (WL) pathway, were crucial in HG. As for TG,

the WL pathway, rTCA, electron transport chain (ETC), TCA, glycolysis, pyruvate metabolism, fermentation, and

dissimilatory nitrate reduction (DNR) were the most important functions. Such metabolic distinction in the three

groups could be attributed to functional cooperation performed by microbes from different modules.

Conclusion

This study is the first to systematically sample geothermal spring ecosystems at both spatial and temporal scales, albeit

locally, to explore how environmental parameters influence the composition and function of microbial communities.

From 12,789 bacterial and archaeal MAGs reconstructed from 152 metagenomes, we uncovered immense

phylogenetic diversity among microbes inhabiting terrestrial geothermal springs, with both species and functional

diversity primarily shaped by pH and temperature, which were divided into three groups: acid group (AG),

hyperthermal group (HG), and thermal group (TG). By combining species and functional MENs, we found that

networks in samples with extreme physicochemical conditions (AG and HG) were less complex but more efficient

than those in relatively benign environments (TG). Microbial communities in HG formed highly modular structures

to enhance cooperation, with most modules highly independent of the others. Dominant microbes define both the

composition and function of the geothermal spring ecosystem, with archaea prevailing in extreme conditions,

especially in AG. Altogether, our study comprehensively and thoroughly elucidates the microbial species and

functional diversity, as well as microbial interactions of geothermal spring ecosystems.

Methods

Study sites, sampling, DNA extraction, and sequencing. Sediment samples were collected twice a year, once in

summer and once in winter, from 49 geothermal springs located in Tengchong, Yunnan, China from 2016 to 2021. A

total of 152 samples were collected and physicochemical parameters were measured. Community DNA was extracted

and subsequently conducted for metagenomic sequencing. The detailed procedures for sample collection, DNA

extraction, and metagenomic sequencing were described in the previous study[36]. Sample information, including

sample images, sampling time, geographical locations, and physicochemical parameters were provided in Fig. S1 and

Table S1. The total genomic DNA was sequenced using the Illumina Hiseq 4000 instrument at Beijing Novogene

Bioinformatics Technology Co., Ltd (Beijing, China). Raw sequencing data for each sample was generated with an

average amount of 30 Gbp (2×150 bp).

Metagenomic assembly, genome binning, and curation. Raw sequencing reads were preprocessed as described in

detail in the previous study[37]. The quality reads of each sample were *de novo* assembled using SPAdes[38] (version

3.15.2) with parameters: -k 21,33,55,77,99,127 --meta. Scaffolds with lengths < 2500 bp in each assembly were

eliminated for subsequent analyses. BBMap[39] (version 38.92, http://sourceforge.net/projects/bbmap/) was used to

calculate the coverage information of each scaffold with the following parameters: k=15 minid=0.97 build=1. Multiple

steps were performed to obtain high-quality genomes from assembled sequences. First, genome binning was

conducted by using MetaBAT[40] (version 2.12.1), CONCOCT[41] (version 1.1.0), and MaxBin 2[42] (version 2.2.7)

separately. DAS Tool[43] was subsequently applied to identify the most accurate and non-redundant bins from the

combined bins generated by the three tools. Reads were recruited by mapping the respective metagenomes to the

selected bins using BBMap. Reads recruited from each bin underwent further reassembly using SPAdes with

parameters: -k 21,33,55,77,99,127 --careful. Finally, a total of 20,849 MAGs with high- and medium-quality was

retained.

Genome quality, including completeness, contaminations, and strain heterogeneity were estimated using

CheckM[44] (version 1.1.3). For bins associated with DPANN and CPR, genome quality was also assessed by calculating

the occurrence frequency of 48 and 43 single-copy genes (SCGs) respectively[26]. Bins with completeness < 50% were

discarded. Genome bins with contamination > 3% were decontaminated by identifying scaffolds with multiple SCGs

and discordant coverage. Transfer RNAs were identified using tRNAscan-SE[45] (version 2.0). Ribosomal RNA genes

including 5S, 16S, and 23S were detected using RNAmmer[46] (version 1.2) and BLAST[47] (version 2.12.0+) searches

against the RDP database[48]. To further improve the quality of genomes, all 16S rRNA genes were aligned using the

online tool SINA (version 1.2.12) (https://www.arb-silva.de/aligner/) with default parameters for taxonomy

assignments. Based on the taxonomy assignment of all genomes, archaeal genomes containing 16S rRNA assigned to

bacteria or vice versa, and genomes with over four 16S rRNA gene copies were removed. After completing all the

screening steps mentioned above, a total of 12790 genomes were subjected to further analyses. Dereplication was

conducted on these MAGs using dRep[49] (version 3.2.2) at a 95% ANI threshold (species level)[26], resulting in the

generation of 3,176 representative MAGs (RMAGs).

Taxonomic, functional annotation, and pan-genome analysis of RMAGs, and. Taxonomic information of all

12,789 MAGs was determined by GTDB-Tk[28] (version 2.0.0; GTDB release 207) with the "classify_wf" command.

Taxonomy of MAGs did not match the taxonomy of the RMAG they were readjusted according to the RMAGs within

each cluster. Putative protein-coding genes (CDSs) of all 12,789 MAGs were predicted using Prodigal[50] v2.6.3 under

the "-p single" mode with standard genetic code and annotated against the KEGG database with parameters: -E 1e-5

using kofam_scan[51] (version 1.3.0). To investigate the microbial pangenome, gene families were identified from

RMAGs with more than one genome. CDS predicted by Prodigal were first subject to all-to-all blast. Only blast hits

with identity ≥ 90%, aligned region ≥ 100bp, and covering ≥ 50% of subject and query genes were kept and

subsequently clustered into gene families using MCL[52] v21-257 with parameter: -I 1.4[53]. The functions of each gene

family were determined according to the voting of each annotated protein sequence.

Phylogenetic analysis. The 3,176 RMAGs representing all microbes inhabiting Tengchong geothermal springs were

used to reconstruct the phylogeny. Based on the recommendation by GTDB-Tk, 120 and 53 conserved marker genes

were selected for reconstructing the bacteria and archaea phylogenies, respectively. Alignment profiles of

concatenated marker genes generated by GTDB-Tk were extracted, and poorly aligned regions were trimmed using

TrimAl[54] (version 1.4.rev22) with parameters: -automated1. IQ-TREE[55] (version 1.6.12) was applied to reconstruct

the maximal-likelihood phylogenies of bacteria and archaea, with the best models determined to be LG+F+R10 for

both bacterial and archaeal trees.

Relative abundance calculation. The relative abundances of each RMAG in each site were calculated. Reads from

each metagenome were mapped onto all 3,176 RMAGs and assembled scaffolds with lengths $\geq$ 500 bp using BBMap,

respectively. The relative abundance of each RMAG in the corresponding sample was calculated as the proportion of

reads recruited from the RMAG to the total reads recruited from all scaffolds with lengths $\geq$ 500 bp. Considering the

bias of genome binning and mapping, false positive reads may be mismatched in mapping and generate an extremely

low relative abundance of RMAGs. By comparing to the *rpS3* data, we replaced the relative abundance below 1E-4

with 0 to remove potential false positives (Supplementary Table 4). Finally, a total of 37,523 microbes with relative

abundances > 0 were identified in all 152 metagenomes, which is comparable to the identified *rpS3* genes(27,436) in

all metagenomes using AMPHORA[56]. This improved method may provide a reference for future microbial community

ecological analysis based on metagenomic data. The relative abundance of gene i (RA_{Gi}) annotated with KOs in one

sample can be calculated by:

$$RA_{Gi} = \sum (No._{Gi} * RA_{RMAGj})$$

where $No._{Gi}$ is the number of *gene i* in *RMAG j*, RA_{RMAGj} is the relative abundance of *RMAG j* in a geothermal spring

sample. The relative abundance of each KO category was calculated as the sum of the relative abundances of all

individual KOs within the category.

Network construction and analysis. All 152 samples were divided into three groups (acidic, thermal, and

hyperthermal) according to their environmental parameters. MENs for each group were constructed based on

community composition and function using iNAP[57] (https://inap.denglab.org.cn). For species MENs, we used the

relative abundance of 3,176 RMAGs as input. Species present in over 25% of samples within each group were kept

for network construction. Then, three MENs were constructed based on Spearman correlation coefficients, and the

correlation cut-off threshold was automatically determined by an RMT-based approach[22,58]. Various network

properties and node properties were calculated based on the adjacent matrix using the MENA Pipeline[57]. As for

functional MENs of the three groups, we selected functional genes occupied in core metabolic pathways to construct

the MENs, based on the relative abundance of selected genes of RMAGs present in each species MENs. Functional

genes that occupied at least 75% of samples within each group were retained for subsequent analysis. Other details

regarding the construction of functional MENs were consistent with the procedures applied for species MENs. All

networks were visualized using Cytoscape[59] (v3.9.1).

Statistical analyses. All statistical analyses were performed in the R software (v4.2.3). Microbial community

dissimilarity (beta diversity) was calculated by the Bray-Curtis index via the vegdist() function in the vegan R package

(v2.6-6.1)[60]. Mann-Whitney U test in R package stats (v4.2.2)[61] was used to test the difference in archaeal community

between extreme geothermal springs and normal geothermal springs. Boxplot, staked bar, Venn diagram, and ternary

diagram were all generated by the "ggplot2" R package (v3.4.1)[62]. The ratio differences of accessory gene families of

three geochemical groups were tested by ANOVA with function aov() in R package stats (v4.2.2)[61], and HSD post hoc

all-pairwise comparisons test[63] was conducted by function HSD.test in R package agricolae (v1.3-2)[64]. Heatmap was

generated by the "pheatmap" R package (v1.0.12)[65].

Code availability

All workflows for phylogeny, annotations, and comparative/evolutionary genomic analyses were deposited at Figshare

(10.6084/m9.figshare.26942083). We also used the following published codes:

quality_control.pl (https://github.com/hzhengsh/qualityControl),

concatenate_multiple_genes.pl (https://github.com/hzhengsh/phylogeny).

Data availability

The datasets analyzed or generated in this study can be found in our Figshare repository

(10.6084/m9.figshare.26942083). All MAGs described in this study have been deposited at NCBI under the BioProject

ID PRJNA544494 (the accession ID of each MAG can be found in Supplementary Table 2).

Acknowledgments

We thank Guangdong Magigene Biotechnology Co., Ltd. China for the assistance in data analysis and the entire staff

from Yunnan Tengchong Volcano and Spa Tourist Attraction Development Corporation for strong support. This work

was financially supported by the National Natural Science Foundation of China (Nos. 32200002，32400002,

91951205, and 32170014).

Author contributions

Y.X.L., Y.Z.R., and Z.S.H. conceived the study. Y.X.L., Y.Z.R., Y.L.Q., Y.N.Q., and Q.J.X. measured physicochemical

parameters and DNA extraction. Y.X.L., Y.Z.R., Z.W.L., S.P.L., J.L.K., Y.J.C., Z.S.H., Y.L.Q., J.Y.J., and Y.N.Q.

performed the metagenomic analysis, genome binning, functional annotation, and ecological analysis. Y.X.L., Y.Z.R.,

Z.W.L, S.P.L., Z.S.H., W.J.L., and W.S.S. wrote the manuscript. All authors discussed the results and commented on

the manuscript.

Competing interests

The authors declare no competing interests.

Reference

1.   Kuang, J. *et al.* Predicting taxonomic and functional structure of microbial communities in acid mine drainage.

ISME J **10**, 1527–1539 (2016).

2.   Ma, B. *et al.* Genetic correlation network prediction of forest soil microbial functional organization. *ISME J* **12**,

2492–2505 (2018).

3.   Sunagawa, S. *et al.* Structure and function of the global ocean microbiome. *Science* **348**, 1261359 (2015).

4.   Tully, B. J., Wheat, C. G., Glazer, B. T. & Huber, J. A. A dynamic microbial community with high functional

redundancy inhabits the cold, oxic subseafloor aquifer. *ISME J* **12**, 1–16 (2018).

5.   Weiss, M. C. *et al.* The physiology and habitat of the last universal common ancestor. *Nat Microbiol* **1**, 16116

(2016).

6.   Brock, T. D. Life at High Temperatures. *Science* **230**, 132–138 (1985).

7.   Des Marais, D. J. & Walter, M. R. Terrestrial Hot Spring Systems: Introduction. *Astrobiology* **19**, 1419–1432

(2019).

8.   Power, J. F. *et al.* Microbial biogeography of 925 geothermal springs in New Zealand. *Nat Commun* **9**, 2876

(2018).

9.   Cole, J. K. *et al.* Sediment microbial communities in Great Boiling Spring are controlled by temperature and

distinct from water communities. *The ISME Journal* **7**, 718–729 (2013).

10.   Barns, S. M., Fundyga, R. E., Jeffries, M. W. & Pace, N. R. Remarkable archaeal diversity detected in a

Yellowstone National Park hot spring environment. *Proc Natl Acad Sci U S A* **91**, 1609–1613 (1994).

11.   Huber, H. *et al.* A new phylum of Archaea represented by a nanosized hyperthermophilic symbiont. *Nature* **417**,

63–67 (2002).

12.   Shu, W.-S. & Huang, L.-N. Microbial diversity in extreme environments. *Nat Rev Microbiol* **20**, 219–235 (2022).

13.   Damer, B. & Deamer, D. The Hot Spring Hypothesis for an Origin of Life. *Astrobiology* **20**, 429–452 (2020).

14.   Colman, D. R., Lindsay, M. R. & Boyd, E. S. Mixing of meteoric and geothermal fluids supports hyperdiverse

chemosynthetic hydrothermal communities. *Nat Commun* **10**, 681 (2019).

15.   Guo, Q., Liu, M., Li, J., Zhang, X. & Wang, Y. Acid hot springs discharged from the Rehai hydrothermal system

of the Tengchong volcanic area (China): formed via magmatic fluid absorption or geothermal steam heating?

Bull Volcanol **76**, 868 (2014).

16.   Hedlund, B. P. *et al.* A review of the microbiology of the Rehai geothermal field in Tengchong, Yunnan Province,

China. *Geoscience Frontiers* **3**, 273–288 (2012).

17.   Lai, D. *et al.* Resource partitioning and amino acid assimilation in a terrestrial geothermal spring. *The ISME*

Journal **17**, 2112–2122 (2023).

18.   HilleRisLambers, J., Adler, P. B., Harpole, W. S., Levine, J. M. & Mayfield, M. M. Rethinking Community

Assembly through the Lens of Coexistence Theory. *Annu. Rev. Ecol. Evol. Syst.* **43**, 227–248 (2012).

19.   Hanson, C. A., Fuhrman, J. A., Horner-Devine, M. C. & Martiny, J. B. H. Beyond biogeographic patterns:

processes shaping the microbial landscape. *Nat Rev Microbiol* **10**, 497–506 (2012).

20.   Toju, H. *et al.* Species-rich networks and eco-evolutionary synthesis at the metacommunity level. *Nat Ecol Evol*
1, 1–11 (2017).

21.   Zamkovaya, T., Foster, J. S., de Crécy-Lagard, V. & Conesa, A. A network approach to elucidate and prioritize
microbial dark matter in microbial communities. *ISME J* **15**, 228–244 (2021).

22.   Yuan, M. M. *et al.* Climate warming enhances microbial network complexity and stability. *Nat. Clim. Chang.*
11, 343–348 (2021).

23.   Thébault, E. & Fontaine, C. Stability of Ecological Communities and the Architecture of Mutualistic and
Trophic Networks. *Science* **329**, 853–856 (2010).

24.   Ma, B. *et al.* A genomic catalogue of soil microbiomes boosts mining of biodiversity and genetic resources. *Nat*
Commun **14**, 7318 (2023).

25.   Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically
consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Research* **50**, D785–D794
(2022).

26.   He, C. *et al.* Genome-resolved metagenomics reveals site-specific diversity of episymbiotic CPR bacteria and
DPANN archaea in groundwater ecosystems. *Nat Microbiol* **6**, 354–365 (2021).

27.   Parks, D. H. *et al.* A complete domain-to-species taxonomy for Bacteria and Archaea. *Nat Biotechnol* **38**, 1079–
1086 (2020).

28.   Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk v2: memory friendly classification
with the genome taxonomy database. *Bioinformatics* **38**, 5315–5316 (2022).

29.   Sun, S., Jones, R. B. & Fodor, A. A. Inference-based accuracy of metagenome prediction tools varies across
sample types and functional categories. *Microbiome* **8**, 46 (2020).

30.   Wang, Y. *et al.* Successional dynamics and alternative stable states in a saline activated sludge microbial
community over 9 years. *Microbiome* **9**, 199 (2021).

31.   Louca, S. *et al.* Function and functional redundancy in microbial systems. *Nat Ecol Evol* **2**, 936–943 (2018).

32.   Brockhurst, M. A. *et al.* The Ecology and Evolution of Pangenomes. *Current Biology* **29**, R1094–R1103 (2019).

33.   Ya, T. *et al.* Successional Dynamics of Molecular Ecological Network of Anammox Microbial Communities
under Elevated Salinity. *Water Research* **188**, 116540 (2021).

34.   Shi, S. *et al.* The interconnected rhizosphere: High network complexity dominates rhizosphere assemblages.

Ecology Letters **19**, 926–936 (2016).

35.   Deng, Y. *et al.* Molecular ecological network analyses. *BMC Bioinformatics* **13**, 113 (2012).

36.   Li, Y.-X. *et al.* Deciphering Symbiotic Interactions of "Candidatus Aenigmarchaeota" with Inferred Horizontal

Gene Transfers and Co-occurrence Networks. *mSystems* **6**, e00606-21 (2021).

37.   Hua, Z.-S. *et al.* Ecological roles of dominant and rare prokaryotes in acid mine drainage revealed by

metagenomics and metatranscriptomics. *ISME J* **9**, 1280–1294 (2015).

38.   SPAdes: A New Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing | Journal of

Computational Biology. https://www.liebertpub.com/doi/abs/10.1089/cmb.2012.0021.

39.   Bushnell, B. *BBMap: A Fast, Accurate, Splice-Aware Aligner*. https://www.osti.gov/biblio/1241166 (2014).

40.   Kang, D. D., Froula, J., Egan, R. & Wang, Z. MetaBAT, an efficient tool for accurately reconstructing single

genomes from complex microbial communities. *PeerJ* **3**, e1165 (2015).

41.   Alneberg, J. *et al.* CONCOCT: Clustering cONtigs on COverage and ComposiTion. Preprint at

https://doi.org/10.48550/arXiv.1312.4038 (2013).

42.   MaxBin 2.0: an automated binning algorithm to recover genomes from multiple metagenomic datasets |

Bioinformatics | Oxford Academic. https://academic.oup.com/bioinformatics/article/32/4/605/1744462.

43.   Sieber, C. M. K. *et al.* Recovery of genomes from metagenomes via a dereplication, aggregation and scoring

strategy. *Nat Microbiol* **3**, 836–843 (2018).

44.   CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes.

https://genome.cshlp.org/content/25/7/1043.short.

45.   tRNAscan-SE: A Program for Improved Detection of Transfer RNA Genes in Genomic Sequence | Nucleic

Acids Research | Oxford Academic. https://academic.oup.com/nar/article-abstract/25/5/955/5133591.

46.   RNAmmer: consistent and rapid annotation of ribosomal RNA genes | Nucleic Acids Research | Oxford

Academic. https://academic.oup.com/nar/article/35/9/3100/2401119?login=false.

47.   Ye, J., McGinnis, S. & Madden, T. L. BLAST: improvements for better sequence analysis. *Nucleic Acids*

Research **34**, W6–W9 (2006).

48.   Cole, J. R. *et al.* Ribosomal Database Project: data and tools for high throughput rRNA analysis. *Nucl. Acids*

Res. **42**, D633–D642 (2014).

49.   Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate genomic comparisons

that enables improved genome recovery from metagenomes through de-replication. *ISME J* **11**, 2864–2868

(2017).

50.   Prodigal: prokaryotic gene recognition and translation initiation site identification | BMC Bioinformatics | Full

Text. https://bmcbioinformatics.biomedcentral.com/articles/10.1186/1471-2105-11-119?report=reader.

51.   Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive score

threshold. *Bioinformatics* **36**, 2251–2252 (2020).

52.   Enright, A. J. An efficient algorithm for large-scale detection of protein families. *Nucleic Acids Research* **30**,

1575–1584 (2002).

53.   Qu, Y.-N. *et al.* Panguiarchaeum symbiosum, a potential hyperthermophilic symbiont in the TACK superphylum.

Cell Reports **42**, 112158 (2023).

54.   Capella-Gutiérrez, S., Silla-Martínez, J. M. & Gabaldón, T. trimAl: a tool for automated alignment trimming in

large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

55.   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: A Fast and Effective Stochastic

Algorithm for Estimating Maximum-Likelihood Phylogenies. *Molecular Biology and Evolution* **32**, 268–274

(2015).

56.   Wu, M. & Scott, A. J. Phylogenomic analysis of bacterial and archaeal sequences with AMPHORA2.

Bioinformatics **28**, 1033–1034 (2012).

57.   Feng, K. *et al.* iNAP: An integrated network analysis pipeline for microbiome studies. *iMeta* **1**, (2022).

58.   Zhou, J. *et al.* Functional Molecular Ecological Networks. *mBio* **1**, e00169-10 (2010).

59.   Shannon, P. *et al.* Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction

Networks. *Genome Res.* **13**, 2498–2504 (2003).

60.   Oksanen, J. *et al. Vegan: Community Ecology Package. R Package (Version 2.5–4)[Computer Software].* (2019).

61.   Team, R. C. R: A language and environment for statistical computing [Computer software manual]. *Vienna,*

Austria (2016).

62.   Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis.* (Springer International Publishing : Imprint:

Springer, Cham, 2016). doi:10.1007/978-3-319-24277-4.

63. Simes, R. J. An improved Bonferroni procedure for multiple tests of significance. *Biometrika* **73**, 751–754

(1986).

64. Fd, M. Agricolae: statistical procedures for agricultural research. *R package version* **1**, 1–6 (2016).

65. Kolde, R. *Pheatmap: Pretty Heatmaps*. (2019).

Fig. 1 | Metagenomic genome reconstruction of geothermal springs in Yunnan. a, Genomic quality of all 20,849 genomes reconstructed from 152 geothermal spring metagenomes. **b**, Phylogenetic distribution of 3,176 RMAGs recovered in this study. The best-fit model of the archaeal and bacterial phylogenetic tree was LG+F+R10. Clades with a bootstrap value ≥ 90% were marked by green and purple dots for the archaeal and bacterial trees respectively. Archaeal classes and bacterial phyla without nomenclature were shown in red text. The orange circles represented the number of RMAGs of each archaeal class and bacterial phylum. **c**, Stacked bar plot for novelty quantification of all archaeal and bacterial genomes. **d**, Boxplot showed the relative abundance of 3,176 RMAGs in 152 samples. The bar plot counted the number of sample sites with different ranges of relative abundance of RMAGs.

Fig. 2 | Community composition and functional profiles of 152 samples. a, The relative abundance of archaea and bacteria in 152 samples. **b**, The proportion of all archaea in all samples. Bars in different colors represented different classes of archaea. **c**, The proportion of all bacteria in all samples. Bars in different colors represented different phyla of bacteria. **d**, the relative abundance of KEGG standard categories (level B) in all samples. **e-f**, Principal coordinate analysis (PCoA) of all RMAGs, archaea, bacteria, and KOs in different sites based on Bray–Curtis distances.

Fig. 3 | Relationship between environmental variables and species and functional composition. a, The heatmap

showed the pairwise correlations of multiple environmental variables with the color gradient denoting Pearson's

correlation coefficients. The edges represented the correlations between community structure and environmental

variables. Edge width corresponds to Mantel's r value, and edge color corresponds to Mantel's P value. Abbreviation:

Temp., Temperature; Cl$^-$, Salinaty; TOM, Total organic matter; TOC, Total organic carbon; TN, Total nitrogen; NH$_4^+$,

Ammonia; NO$_3^-$, Nitrate; NO$_2^-$, Nitrite; TP, Total phosphorus; AP, Available phosphorus; AS, Available sulfur; SO$_4^{2-}$,

Sulfate; K$^+$, Exchangeable potassium; Na$^+$, Exchangeable sodium; Ca^{2+}, Exchangeable calcium; Mg2+, Exchangeable

 magnesium; Si^{2+}, Effective silicon; TCu, Total bronze; TZn, Total zinc; TPb, Total lead; TFe, Total iron; TCr, Total

 chrome. **b**, Analysis of similarities (ANOSIM) plot showed the dissimilarity of the three groups. The width of the bar

 is directly proportional to the sample size. The "**" represented the significant difference (P = 0.001) between every

 two groups based on the pairwise ANOSIM analysis of the three groups. **c**, The ternary diagram of the relative

 abundance of microorganisms in three groups. The top 25 microorganisms with high average relative abundance in

 all 152 samples were shown in the ternary diagram. **d**, The number of core and accessory gene families of each group.

 e, Boxplot showed the proportion of accessory gene families of each group. The "**" represented the P value < 0.001

 in the Wilcoxon rank sum test.

 Fig. 4 | Co-occurrence microbial networks of three geothermal spring groups. a-c, Visualization of constructed

 MENs of **a**, AG, **b**, HG, and **c**, TG. The nodes were colored according to the phylum taxonomy of RMAGs. The circle

 size represented the nodes' degrees, and the values listed in Supplementary Data 10. A connection represents a strong

 (Spearman's correlation coefficient ≥ 0.78) and significant (P-value < 0.01) correlation. **d-h**, Important network

533   properties of three groups, including **d**, Average degree, **e**, Average clustering coefficient, **f**, Average path distance, **g**,

534   Connectedness, **h**, Modularity. Details of all network properties are listed in Supplementary Data 9.

535