PP2A-dependent internalisation of GABA$_B$ receptors in somatostatin interneurons regulates function and plasticity.

Nithya Sethumadhavan[1], Max A Wilson[2,4], Anna Sumera[2,3,4], Desiree Loreth[1], Rita M Loureiro[4], Imre Vida[4], Akos Kulik[1$], Sam A Booker [2, 3, 4$],

[1] – Institute of Physiology II, University of Freiburg, Freiburg, Germany

[2] – Simons Initiative for the Developing Brain, University of Edinburgh, Edinburgh, UK

[3] – Patrick Wild Centre for Autism Research, University of Edinburgh, Edinburgh, UK

[4] – Centre for Discovery Brain Sciences, University of Edinburgh, Edinburgh, UK

[5] – Charité Universitätmedizin, Berlin, Germany;

Co-corresponding authors ([$])

Sam A Booker (Lead contact)

Simons Initiative for the Developing Brain,

Centre for Discovery Brain Sciences

University of Edinburgh

Hugh Robson Building, George Square,

Edinburgh, EH8 9XD, UK

sbooker@ed.ac.uk

Akos Kulik

Institute of Physiology II,

Faculty of Medicine,

University of Freiburg,

Hermann-Herder-Str. 7,

Freiburg, 79104, Germany

Tel.: +49 761 203 67305

akos.kulik@physiologie.uni-freiburg.de

Figures: 8 (4 supplementary)

Tables: 0 (2 supplementary)

Abstract:

Cortical circuits rely on a precise balance of inhibitory and excitatory neurotransmission to
encode information reliably and prevent pathology. Metabotropic GABA$_B$ receptors
(GABA$_B$Rs) are key regulators of inhibitory signalling in mammalian neurons. In
GABAergic interneurons (INs), GABA$_B$R activation reduces inhibition overall, leading to
disinhibitory mechanisms. In the hippocampus, somatostatin-expressing (SST) INs form a
major subtype that provides feedback inhibition to the distal dendrites of principal cells
(PCs) and other INs. Plasticity of SST INs is well established as a mechanism controlling
hippocampal circuit function through both inhibitory and disinhibitory pathways and
depends on metabotropic glutamate receptors (mGluRs) and GABA$_B$Rs. However,
whether activation of GABABRs induces metaplastic changes in SST INs, and how this
influences circuit function and behaviour, remains unclear.

Here, we combined quantitative SDS-digested freeze-fracture replica immunoelectron
microscopy, ex vivo electrophysiology, in vivo behavioural testing, and pharmacological
manipulation of GABA$_B$Rs. We show that receptor activation directly regulates SST IN
plasticity via protein phosphatase 2A (PP2A)-dependent internalisation. GABA$_B$R
activation not only controls its own surface expression but also regulates membrane levels
of mGluR1α and high-voltage-activated Ca$_v$1.2 (L-type) Ca^{2+} channels. This GABA$_B$R-
dependent metaplasticity shifts circuit plasticity toward greater enhancement of long-range
inputs to the CA1 region and disrupts contextual memory formation. These findings
demonstrate that receptor-mediated surface dynamics in SST INs are critical for
maintaining physiological neurotransmission and proper hippocampal microcircuit function.

Introduction

The appropriate function of cortical microcircuits relies on dynamically balanced excitatory
and inhibitory synaptic transmission, which maintains the neural code and contributes to
memory formation[1-3]. How these circuits respond to changing activity level, or indeed
sustained pharmacological activity, is crucial to understand how the brain functions at rest
and how neuropathological conditions lead to altered behaviour and cognition.

Synaptic inhibition is produced, in large part, by a heterogenous population of local
interneurons (INs). These INs release GABA from their presynaptic terminals, which
activates both synaptic and extrasynaptic ionotropic GABA$_A$ receptors (GABA$_A$Rs) and

metabotropic GABA$_B$ receptors (GABA$_B$Rs), on excitatory principal cells and INs alike [4].

Such inhibition of INs reduces their activity, leading to reduced local GABA release, and

thus a reduction of inhibitory control – known as disinhibition. Dynamic interactions

between inhibition and disinhibition are critical to ensure the timing and strength of

neuronal activity leading to complex behaviour. To date, most studied forms of disinhibition

focus on direct synaptic GABA$_A$R-mediated effects, which can regulate circuit level

plasticity in hippocampus [5] and neocortex [6]. While the direct effect of GABA$_B$R signalling

on INs has been well described by ourselves and others [7-14], how their activation shapes

the long-term excitability of inhibitory cells and influence circuit activity remains less well

understood.

In CA1 of the hippocampus, a major type of IN are those which express the neuropeptide

somatostatin (SST)[15] which have their somatodendritic domain within *stratum*

oriens/alveus (str. O/A) receiving inputs from local pyramidal cells (PCs)[10,16,17], with axons

ramifying in *str. lacunosum-moleculare* (*Str. L-M*). Thus, SST INs provide powerful

feedback inhibition into the CA1 circuit[18]. Therefore, the activation of SST INs leads to

complex circuit functions. First, they directly inhibit distal dendrites of CA1 PCs co-

terminating with entorhinal cortex inputs to strongly inhibit spatial information[19-21]. Second,

SST INs disinhibit proximal dendritic domains and thus gate contextual inputs from CA3[22].

As such, long-term potentiation of SST INs leads to a functional re-arrangement of the

hippocampal information transfer, favouring CA3 inputs [22,23] and are potentially able to

alter the circuit dynamics of the hippocampus over behaviourally relevant time-scales [24].

We have shown that GABA$_B$Rs on SST INs acutely inhibit the induction of long-term

potentiation (LTP)[11] and their local circuit integration [10], but do not directly hyperpolarise

SST INs. Given the high affinity of GABA for GABA$_B$Rs and their plethora of accessory

proteins, they undergo rapid desensitization [8,25-29] and agonist dependent internalisation

[30,31]. Not only this, but GABA$_B$Rs are known to interact with metabotropic glutamate

receptors (mGluRs), preferentially with type 1α receptors (mGluR1α), in other GABAergic

cells in the cerebellum [32,33]. Considering the high density of both mGluR1α, and

GABA$_B$Rs on SST INs [34,35] it is plausible that receptor cross-talk also exists in these cells.

Whether GABA$_B$R signalling is susceptible to rapid internalisation in SST INs, and how this

affects cell function and the local circuit remains unknown.

We hypothesised that GABA$_B$R activation leads to internalisation of itself, favouring greater

plasticity in SST INs. To determine this, we performed quantitative SDS-digested freeze-

fracture replica immunogold labelling (SDS-FRL) electron microscopy, *ex vivo*

electrophysiological recordings from CA1 of the mouse hippocampus following prolonged
application of the selective $GABA_BR$ agonist baclofen, and behavioural characterisation.
We found, in opposition to our hypothesis, that after sustained $GABA_BR$ activation,
synaptic plasticity of SST INs was reduced, independent of direct receptor activation. This
reduction in plasticity was due to concomitant protein phosphatase 2A (PP2A)-mediated
internalisation of $GABA_BRs$, $Ca_v1.2$ (L-type) voltage-gated Ca^{2+} channels (VGCCs) and
mGluR1α – both of which are required for LTP of SST INs. Finally, we show that baclofen
administration leads to circuit level strengthening of long-range inputs to CA1, which are
dependent on SST IN plasticity, and an inability to form contextual memories.

Results

In this study we tested the hypothesis that sustained baclofen application leads to internalisation of $GABA_B$Rs, which favours enhanced plasticity of SST INs following receptor activation. For this, we performed detailed quantitative immunoelectron microscopy, electrophysiology, and behaviour in adult mice.

Native $GABA_B$Rs, mGluR1α and L-type calcium channels are down-regulated in SST IN membranes following baclofen pre-application, which requires PP2A

We have previously shown that mouse SST IN dendrites possess a high density of $GABA_{B1}$, $Ca_V1.2$, and mGluR1α[11]. The latter two proteins are required for associative theta-burst stimulus (aTBS) LTP of SST INs [36,37], and is inhibited by $GABA_B$Rs [11]. To determine whether pre-application of the $GABA_B$R agonist baclofen (20 µM) altered membrane localization of $GABA_B$Rs as compared to mGluR1α and $Ca_V1.2$, we performed SDS-FRL immunoelectron microscopy analysis from acute mouse hippocampal slices pre-treated with baclofen (n=5 mice), compared slices treated with vehicle (n=5 mice) or the $GABA_B$R antagonist CGP-55,845 (5 µM, n=3 mice; Figure 1A). All SDS-FRL data are shown as animal average values, with statistics shown based on linear mixed-effects modelling to account for inter-animal variability. Data from each dendritic profile within animal is shown in Figure S1.

Under control conditions (Figure 1B), putative SST IN dendrites were identified based on high levels of surface labelling for mGluR1α [34], with an average density of 37.6 ± 4.7 particles/µm^2 (Figure1B, E). Consistent with our previous data in rats [11], mGluR1α-positive dendrites displayed high-density $GABA_{B1}$ subunit labelling of 35.1 ± 2.9 particles/µm^2 and moderate expression of the L-type channel subunit $Ca_V1.2$ at 5.46 ± 0.98 particles/µm^2 (Figure 1B, E). Compared to control slices, baclofen pre-application led to a 49% reduction in density to 17.9 ± 2.5 particles/µm^2 for $GABA_{B1}$ ($t_{(5,5)}$=5.07, *p*=0.004, Tukey test; Figure 1C, E). We observed a 58% reduction in $Ca_V1.2$ to 2.3 ± 0.3 particles/µm^2 ($t_{(5,5)}$=5.36, *p*=0.0009, Tukey test) and a 70% reduction of mGluR1α labelling 11.2 ± 1.8 particles/µm^2 ($t_{(5,5)}$=6.36, *p*=0.0004, Tukey test) (Figure 1C, E). Application of the $GABA_B$R antagonist CGP-55,845 did not decrease density of $GABA_{B1}$ ($t_{(5,3)}$=1.97, *p*=0.18, Tukey test), $Ca_V1.2$ ($t_{(5,3)}$=2.01, *p*=0.17, Tukey test) or mGluR1α ($t_{(5,3)}$=0.69, *p*=0.78, Tukey test) relative to control levels (Figure 1D, E). This effect on effector channels appeared confined to SST INs, as measurement of CA1 PC dendrites in the same replicas revealed that while baclofen pre-application reduced $GABA_{B1}$ density by 49.9% ($F_{(5,5)}$=45.22, p=4.57x10^{-10},

LMM, Supplementary Figure 2) Ca$_V$1.2 was unchanged ($F_{(3,3)}$=0.009, p=0.93, LMM;
Supplementary Figure 2). mGluR1α was not tested as this protein is not expressed
abundantly on CA1 PCs. Together, these data reveal that GABA$_B$R activation leads to its

own reduction from the membrane of SST INs, but also leads to down-regulation of Ca$_V$1.2
and mGluR1α.

Figure 1 Sustained GABA$_B$R activation leads to reduction of membrane associated
GABA$_{B1,}$ Ca$_V$1.2, and mGluR1α proteins from putative SST-INs. **A** Schematic of the
experimental overview, indicating treatment and experimental sample times. **B** Example
electron micrographs of SDS-FRL for mGluR1α (15 nm immunoparticles), Ca$_V$1.2 (10 nm,
blue overlay) and GABA$_{B1}$ (5 nm, red overlay) on a putative SST IN dendrite in *str. O/A* of
CA1 in a vehicle slice. The same labelling but performed in brain slices following 20
minutes baclofen pre-application (20 µM, **C**) or CGP-55,845 (5 µM, **D**). **E** Quantification of
labelling density for GABA$_{B1}$ (left), Ca$_V$1.2 (middle) and mGluR1α (right), in slices treated
with vehicle (Ctrl., black, 5 mice), baclofen pre-application (Bacl., red, 5 mice), or CGP-
55,845 (CGP, magenta, 3 mice). All data are shown as boxplots depicting the 25-75%
range with median, maximum range and data from individual mice overlaid as open

circles. Statistics shown as: ns – p>0.05, * - p<0.05, ** - p<0.01, *** - p<0.001; all from

post hoc Tukey tests following LMM analysis. Scale bar represents 200 nm.

Baclofen pre-application leads to reduced Cav1.2 and group 1 mGluR-mediated currents

in SST INs.

To determine whether baclofen pre-application regulates the function of $Ca_v1.2$ and

mGluR1α in SST INs, we next performed whole-cell patch clamp recordings from identified

cells using targeted pharmacology to interrogate receptor internalisation (Figure 2A), as

$GABA_BR$ internalisation has been shown to be dependent on increased activity of

$PP2A$[26,31,38].

As $Ca_v1.2$ containing VGCCs are a major target of $GABA_BR$ mediated inhibition in SST

INs[11], we first performed whole-cell recordings using a Cs-gluconate based internal

solution containing QX-314 (5 mM) to block voltage activated K^+ and Na^+ channels,

respectively, in the presence of CNQX (10 µM), DL-AP5 (50 µM) and gabazine (10 µM) to

block ionotropic receptor signalling. High-voltage activated Ca^{2+} responses were

measured from -60 mV voltage-clamp, in which 5 mV depolarising current steps were

applied, followed by bath application of the selective L-type VGCC blocker nifedipine

(10 µM; Figure 2B). The average subtracted L-type current measured across a range of

depolarising stimuli revealed large amplitude currents in SST INs recorded under control

conditions, which were substantially reduced, but with a similar voltage-dependency in

slices pre-application with baclofen. We found that baclofen pre-application reduced the

peak L-type current measured by 37% ($U_{(16,11)}$= 50; P=0.032, Mann-Whitney test, Figure

2C). We performed the same recordings, but in slices that had been incubated with the

PP2A inhibitor okadaic acid (OA, 1 µM; Figure 2D). Blocking PP2A activity had minimal

effect on current-voltage response or peak L-type currents following baclofen pre-

application ($U_{(5,5)}$= 0; P=0.008, Mann-Whitney; Figure 2E).

For mGluR1 signalling, we recorded SST INs in whole-cell configuration then puff applied

the selective, high-affinity group I mGluR agonist S-DHPG (50 µM, 200 ms, 10 PSI) within

100-200 µm of the cell body. S-DHPG puffs resulted in large and slow depolarising

currents in SST INs, which were partially sensitive to the mGluR1α selective antagonist

LY367,385 (100 µM). Measurement of the same S-DHPG induced currents in slices that

were pre-application with baclofen possessed typically smaller currents (Figure 2F).

Quantification of peak S-DHPG currents revealed that baclofen pre-application reduced

the amplitude (F(1,22)=7.99, P=0.01, 2-way ANOVA [interaction], Figure 2G) and the charge transfer (F(1,22)=9.59, P=0.005, 2-way ANOVA [interaction], Figure 2H), effects which were not present when slices were incubated with OA prior to baclofen pre-application.

Figure 2: Ca$_V$1.2 and group 1 mGluR-mediated currents are reduced following baclofen pre-application. **A** Schematic of recording conditions, highlighting treatment periods and recording times. **B** Example nifedipine-sensitive VGCC current responses recorded in SST INs following depolarising stimulation from -60 mV. Dashed line indicates control response amplitude for comparison. **C** Quantification of voltage-dependency of nifedipine-sensitive VGCC currents (left) and peak nifedipine-sensitive currents (right). **D** Example nifedipine-subtracted isolated VGCC current responses in SST INs pre-application with 1 µM okadaic acid. **E** Quantification of voltage-dependency of nifedipine-sensitive VGCC currents (left) and peak nifedipine-sensitive currents (right) in okadaic acid pre-application SST INs. **F** Example voltage responses following 50 µM S-DHPG puff application to SST INs, recorded under control conditions (black) or following baclofen pre-application (red), following either vehicle or 1 µM okadaic acid pre-application. **G** Quantification of absolute current amplitudes following S-DHPG puff. **H** Quantification of the area-under-curve (AUC) for S-DHPG puff induced transients. Data are shown as either mean ± SEM (current-voltage plots) or box-plots showing 25-75% box with median, with maximum range shown.

Individual cell data is shown overlaid. Statistics shown as: ns – p>0.05, * - p<0.05; from 2-

way ANOVA and Mann Whitney non-parametric tests

Baclofen pre-application dependent reductions in $Ca_V1.2$ and mGluR1α are mediated by

PP2A.

To confirm surface down-regulation of $GABA_BRs$ was indeed dependent on PP2A in SST

INs, we utilised a mouse line expressing a Channelrhodopsin2 fused to yellow

fluorescence protein (ChR2/YFP) under the SST promotor to allow selective SDS-FRL

from identified SST IN dendrites [10]. Initially, we prepared slices from WT mice treated with

the PP2A inhibitor OA. However, in these slices we observed minimal to no labelling for

mGluR1α, $GABA_{B1}$ or $Ca_V1.2$ labelling in putative SST IN and CA1 PC dendritic profiles

(Figure S3). This effect was observed across >5 independent experiments.

As such, we next used the the small molecule PP2A inhibitor fostriecin (50 nM) which

slices were incubated with for 1 hour prior to baclofen pre-application (Figure 3A). In *str.*

O/A from the ChR2/YFP mice, we observed strong immunolabeling for YFP (33.7 ± 1.07

particles/$μm^2$, Figure 3B). In vehicle slices, we observed a 45% reduction of $GABA_{B1}$

labelling following baclofen pre-treatment (16.7± 1.02 particles/$μm^2$, $t_{(3,2)}$=3.45, *p*=0.037,

Kenward-Roger test, Figure 3B, D), consistent with observations from mGluR1α labelled

dendrites (Figure 1). Incubation with fostriecin prevented the baclofen pre-treatement

mediated reduction in $GABA_{B1}$ surface density (28.3 ± 3.2 particles/$μm^2$ $t_{(3,2)}$=0.88, *p*=0.44,

Kenward-Roger test, Figure 3C, D), confirming that PP2A inhibition prevents $GABA_BR$

internalisation.

Figure 3: Inhibition of PP2A prevents internalisation of GABA$_B$Rs in SST INs following their activation. **A** Schematic of experimental conditions, highlighting treatment periods and fixation times **B** Example electron micrographs of SDS-FRL for SST-cre dependent expression of ChR2/YFP (18 nm immunoparticles) and GABA$_{B1}$ (6 nm, red overlay) in *str. O/A* of CA1 of slices with vehicle or baclofen pre-application (20 μM). **C** The same labelling but performed in slices incubated with 50 nM fostriecin for 1 hour prior to baclofen pre-application. **D** Quantification of GABA$_{B1}$ density in vehicle slices (Ctrl., black, 3 mice) or those pre-application with baclofen (Bacl., red, 2 mice) with and without fostriecin

incubation. All data are shown as boxplots depicting the 25-75% range with median, maximum range shown, and average density calculated from individual mice results overlaid as open circles. Scale bar represents 200 nm.

To determine whether GABA$_B$R-dependent internalisation of L-type calcium channels also requires PP2A-dependent mechanisms, we performed the same experiments but labelling for Ca$_V$1.2 (Figure 4A). In replicas from vehicle treated slices we observed Ca$_V$1.2 labelling of 3.7± 0.3 particles/µm^2, which was 46% reduced in baclofen pre-application slices (1.8 ± 0.3 particles/µm^2 t$_{(3,2)}$=3.32, *p*=0.0497, Kenward-Roger test, Figure 4B, D), consistent with with mGluR1α labelling experiments (Figure 1). Fostriecin pre-incubation (Figure 3C) prevented this reduction in Ca$_V$1.2 labelling following baclofen treatment (3.5 ± 0.4 particles/µm^2 t$_{(3,2)}$=1.06, *p*=0.326, Kenward-Roger test, Figure 3C, D), confirming that PP2A inhibition prevents Ca$_V$1.2 channel internalisation.

These data confirm that baclofen induced GABA$_B$R internalisation requires PP2A in SST INs in *str. O/*A, and that inhibiting this internalisation also prevents loss of Ca$_V$1.2 from their dendritic membranes.

Figure 4: Inhibition of PP2A prevents internalisation of Ca_V1.2 in SST INs following GABA_BR activation. **A** Schematic of experimental conditions, highlighting treatment periods and fixation times. **B** Example electron micrographs of SDS-FRL for SST-cre dependent expression of ChR2/YFP (18 nm immunoparticles) and Ca_V1.2 (6 nm, blue overlay) in *str. O/A* of CA1 of slices with vehicle or baclofen (20 µM). **C** The same labelling but performed in slices pre-application with 50 nM fostriecin for 1 hour prior to baclofen

application. **D** Quantification of $Ca_V1.2$ density in slices treated with vehicle (Ctrl., black, 3

mice) or baclofen (Bacl., red, 2 mice) with and without fostriecin pre-application. All data

are ~~is~~ shown as boxplots depicting the 25-75% range with median, maximum range

shown, and average density calculated from individual mice results overlaid ~~as~~ open

circles. Scale bar represents 200 nm.

Sustained $GABA_BR$ activation leads to reduced plasticity of SST INs that is mediated by

receptor internalisation

We have previously shown that aTBS induced LTP in SST INs can be prevented by

acutely activating $GABA_BRs$[11]. However, our data suggests that application of baclofen

leads to reduced LTP machinery from SST INs. To determine whether sustained activation

of $GABA_BRs$ shape the long-term plasticity landscape of SST INs, we first performed

whole-cell patch-clamp recordings in *str. O/A* from adult male and female transgenic mice

expressing YFP (Venus variant) under the vGAT promoter (Figure 5A). We recorded 108

str. O/A INs for the current study, those that were recovered (n=70) typically had horizontal

dendrites confined to *str. Oriens* (*str. O*), with axons projecting to *Str. L-M* and were

uniformly SST immunoreactive (Figure 5B). Recorded SST INs displayed prominent I_h-

mediated sag potentials, and possessed a medium to high rate of action potential (AP)

discharge (Figure 5C), which was lower after baclofen pre-application; due to reduced

input resistance and elevated rheobase (Supplementary Table 1).

Under vehicle control conditions, we observed excitatory postsynaptic currents (EPSCs)

resulting from *alveus* stimulation of 197.8 ± 17.6 pA (13.3 ± 3.9 V stimulation, n=11 SST

INs), which following aTBS-LTP induction led to a facilitation of 68.5 ± 12.9% above

baseline levels (p=0.001, Wilcoxon matched-pairs test). In baclofen pre-application slices,

we observed baseline EPSCs with an average amplitude of 195.6 ± 54.6 pA (8.5 ± 2.9 V

stimulation, n= 8 SST INs), which did not differ from vehicle recordings in amplitude ($U_{(11,}$

$_{8)}$= 28, P=0.21, Mann-Whitney) or stimulation strength ($U_{(11, 8)}$= 33, P=0.38, Mann-

Whitney). An aTBS-LTP induction in baclofen pre-application slices failed to increase

EPSC amplitude above baseline levels (11.3 ± 9.6% above baseline, p=0.38, Wilcoxon

matched-pairs test), and which was lower than for vehicle conditions ($U_{(11,8)}$= 10; p=0.004,

Mann-Whitney). To confirm that this loss of aTBS-LTP was not due to residual direct

activation of $GABA_BRs$ we performed the same experiments, but in which the $GABA_BR$

antagonist CGP-55,845 (CGP, 5 µM) was applied to the bath through-out recording

(Figure 5E). In the presence of CGP, we still failed to induce aTBS-LTP in baclofen pre-
application slices ($31.3 \pm 19.6\%$ below baseline, p=0.31, Wilcoxon matched-pairs test),
despite persistent induction in vehicle treated slices ($54.5 \pm 28.0\%$ above baseline
p=0.031, Wilcoxon matched-pairs test; $U_{(6,6)}$=5; p=0.041 Mann Whitey); confirming that
aTBS-LTP inhibition was not due to direct receptor activation.

As postsynaptic $GABA_BRs$ undergo PP2A dependent internalisation (Figure 3) [26,30,31,38],
we next asked if the impaired aTBS-LTP in SST INs was due to internalisation of the
receptor. For these experiments, we incubated slices with OA (1 µM) a for 60-120 minutes
prior to vehicle or baclofen pre-application. Following OA incubation, we continued to
observe reliable aTBS-LTP in both vehicle slices ($112.1 \pm 26.4\%$ above baseline, p=0.016
Wilcoxon paired test) and baclofen pre-application slices in SST INs ($140.1 \pm 31.3\%$ above
baseline, p=0.029 Wilcoxon paired test), which were not different ($U_{(7,5)}$ = 15; p=0.76,
Mann-Whitney test; Figure 5F). As our SDS-FRL labelling was not possible with OA
(Supplemetary Figure 3), we next confirmed that fostriecin (50 nM) produced similar
effects to OA. After fostriecin incubation (60-90 minutes), we observed a tendency towards
aTBS-LTP in vehicle slices ($81.8 \pm 45.6\%$ above baseline, p=0.063 Wilcoxon paired test)
and robust LTP in baclofen pre-application slices ($69.0 \pm 17.0\%$ of baseline, p=0.016
Wilcoxon paired test), which was not different from vehicle ($U_{(5,7)}$=16, p=0.876, Mann-
Whitney test; Figure 5G).

Combined, these data indicate that sustained $GABA_BR$ activation impairs aTBS-LTP
induction in SST INs, which is independent of direct receptor activation and which relies on
internalisation of the receptor via PP2A-dependent phosphorylation.

Figure 5: LTP of local inputs to SST INs is abolished by prolonged baclofen application. **A**

Schematic of recording conditions, highlighting treatment periods and recording times. **B**

Example reconstruction of a SST IN recorded from *str. O/A* of the adult mouse CA1 region,

showing somatodendritic (black) and axonal (red) ramifications. Inset immunoreactivity for

SST (green) and biocytin (purple). **C** Voltage responses in response hyper- to depolarising
current steps (-500 to +500 pA, 100 pA steps, 500 ms duration) from SST INs recorded in
following vehicle (black) and baclofen-pre-application (Bacl.$_{pre-app}$; red). Right,
quantification of the action potential output of SST INs recorded under both conditions. **D**
Example EPSC traces from vehicle (upper, black) and 20 µM baclofen pre-application
(lower, red) measured under control conditions; for aTBS-LTP traces baseline EPSCs are
shown underlain (grey and pink). Middle, time-course of plot of EPSC amplitude before
and after aTBS-LTP induction for vehicle (black) and baclofen pre-application (red); 100%
baseline level is shown for reference (grey dashed line). Right, bar-chart of EPSC
amplitude change post aTBS-LTP (25-30 minutes) compared to baseline for vehicle
(black) and baclofen pre-application (red); 100% baseline level indicated (grey dashed
line). **E** aTBS-LTP recordings from SST INs performed in the presence of 5 µM CGP-
55,845 (CGP) according to the same scheme as **D**. **F** aTBS-LTP following pre-application
of slices with 1 µM okadaic acid for 1 hour with subsequent vehicle or baclofen pre-
application. **G** aTBS-LTP in slices pre-application with 50 nM fostriecin for 1 hour prior to
vehicle or baclofen pre-application. All data are shown as either mean ± SEM (time-course
plots) or box-plots showing 25-75% box with median, with maximum range shown and
individual cell results. Statistics shown as: ns – p>0.05, * - p<0.05, ** - p<0.01from Mann-
Whitney non-parametric tests.

Prolonged activation of GABA$_B$Rs impairs presynaptic release of GABA, but not
postsynaptic expression of GABA$_B$Rs in CA1 PCs

Assessing the wider function of GABA$_B$R signalling following sustained pharmacological
activation is critical to infer circuit wide effects. In CA1 PCs, following baclofen pre-
application we observed a tendency towards reduced GABA$_{B1}$, but unchanged Ca$_V$1.2
surface expression. We next determined if these effects altered inhibitory signalling in and
onto CA1 PCs. Therefore, we recorded GABA$_B$R-mediated slow IPSCs in the presence of
CNQX (10 µM), DL-AP5 (50 µM) and gabazine (10 µM) to block ionotropic receptor
signalling (Figure 6A). We found that slow IPSCs were substantially reduced following
baclofen pre-application (U$_{(10,7)}$=3, p=0.0007, Mann-Whitney test; Figure 6B). However,
baclofen-mediated whole-cell currents (I$_{WC}$) were unaffected by baclofen pre-application
(Figure 6C), with normal peak current-density (U$_{(9,7)}$= 15.5, p=0.10, Mann-Whitney test;
Figure 6D). One explanation for this disconnect between synaptic and pharmacological
activation of GABA$_B$Rs could be impaired GABA release. As such, we measured the

input/output (I/O) relationship of ionotropic GABA$_A$R-mediated IPSCs following *str. L-M*
stimulation as a proxy for synaptic strength. We found that for the same stimulus, fast
IPSCs were consistently lower in amplitude (Figure 6E), which upon quantification
revealed 46% lower I/O slopes ($U_{(7,6)}$=4, p=0.014, Mann-Whitney test, Figure 6F). To
determine whether baclofen pre-application alters presynaptic GABA$_B$R function, we next
bath applied 10 µM baclofen to recordings of fast IPSCs (Figure 6G). Slices with pre-
application of baclofen tended to have lower GABA$_B$R presynaptic inhibition of IPSCs in
str. L-M ($U_{(7,6)}$=7, p=0.051, Mann-Whitney test, Figure 6H). Finally, we compared the effect
of baclofen pre-application on CA1 PC intrinsic excitability. Contrary to what we observed
in SST INs, we find that CA1 PCs are largely unaffected by baclofen pre-application, with
subtle effects on membrane capacitance and action potential kinetics observed (Table S2).

Together these data suggest that prolonged GABA$_B$R activation has limited effects on the
postsynaptic function of CA1 PCs, in particular we find no evidence for reduced functional
GABA$_B$R expression. However, due to strong reductions in presynaptic GABA release,
postsynaptic IPSCs are functionally reduced.

Figure 6: Prolonged GABA$_B$R activation decreases GABA release in str. L-M, specific for
presynaptic but not postsynaptic GABA$_B$R-mediated inhibition. **A** Experimental schematic
of recordings from CA1 PCs and example slow IPSCs recorded following 5x 200 Hz
stimulation of *str. L-M* in the presence of CNQX (10 µM), DL-AP5 (50 µM) and gabazine

(10 µM) from -65 mV voltage-clamp from control slices (black) and those pre-application

with 20 µM baclofen (red). **B** quantification of slow IPSC amplitudes from control and

baclofen pre-application slices. **C** Time course of whole-cell current (I_{WC}) recorded from -

65 mV following bath application of 10 µM baclofen in control and baclofen pre-application

slices. **D** Quantification of I_{WC} current density, normalised to cell capacitance. **E** Fast

IPSCs evoked following paired stimulation of *str. L-M* from 0 mV using Cs-gluconate

internal solution in the presence of CNQX (10 µM), DL-AP5 (50 µM) from control and

baclofen pre-application slices. Traces recorded following bath application of 10 µM

baclofen are shown underlain (grey/pink). Right, input-output (I/O) plot of fast IPSC

amplitude from both groups. **F** Quantification of I/O slope for fast IPSCs. **G** Time course of

fast IPSC amplitude before, and following bath application of 10 µM baclofen and 5 µM

CGP-55,845 (CGP) in control and baclofen pre-application slices. **H** Quantification of

baclofen mediated inhibition of fast IPSC amplitudes, expressed as % of baseline. Data is

shown as either mean ± SEM (**C, E, G**) or box-plots showing 25-75% box with median,

with maximum range (**B, D, F, H**). Individual cell data is shown overlaid. Statistics shown

as: ns – p>0.05, * - p<0.05, *** - p<0.001; from Mann Whitney non-parametric tests.

GABA$_B$R-dependent internalisation of LTP proteins in SST INs leads to enhanced
temporoammonic plasticity onto CA1 PCs

SST INs are known to control the ability of CA1 PCs to undergo plasticity, notably in their

distal dendrites in *str. L-M* aligned with entorhinal cortex inputs [23]. To determine whether

prolonged activation of GABA$_B$Rs, and subsequent loss of SST IN plasticity alter

temporoammonic LTP we performed extracellular field recordings from *str. L-M* of CA1, in

which we induced plasticity at SST INs via repetitive stimulation of the alveus, followed by

high frequency stimulation (HFS, 1x 100 Hz) to the temporoammonic pathway in vehicle or

those with pre-application of 20 µM baclofen for 20 minutes (Figure 7A). Following HFS to

str. L-M we observed increased fEPSP amplitudes when measured in all recordings from

both treatment groups (Figure 7B), which when measured at 50-60 minutes post induction

were 25.6 ± 7.8% above baseline in vehicle control recordings, and which did not differ in

baclofen pre-application slices (20.5 ± 10.4%; $U_{(15,19)}$=104, p=0.19, Mann-Whitney test,

Figure 7C). However, we noted that a higher proportion of slices pre-application with

baclofen failed to undergo LTP (>10% above baseline) as compared to vehicle controls

(χ^2=7.5, p=0.0062 Chi square test; Figure 7D). Therefore, we next analysed only those

recordings in which LTP was observed (Figure 7E). Further quantification of successful

LTP recordings (Figure 7F) revealed that post-tetanic potentiation (0-1 minute post HFS) was 34% higher in baclofen pre-application slices ($U_{(11,10)}$= 25, p=0.018, Mann-Whitney test) and fEPSPs measured at 20-30 minutes (consistent with whole-cell recordings) were also 24% higher than control slices ($U_{(12,10)}$=27, p=0.015, Mann-Whitney test), while LTP measured at 50-60 minutes was higher, albeit not significantly so ($U_{(12,10)}$=37, p=0.07, Mann-Whitney test). We observed no difference in fEPSP slope relative to stimulus strength between control and baclofen pre-application slices (Figure 7G). These effects were likely specific to SST INs, as performing the same recordings but in the presence of the mGluR1α specific antagonist LY367,385 (100 µM), prevented any baclofen pre-application effects on temporoammonic LTP (Figure S4). This also excludes the effects of baclofen pre-application on GABAergic release and CA1 PC excitability as being causal to this effect.

These data confirm that prolonged GABA$_B$R activation leading to a down-regulation in surface expression of GABA$_{B1}$, mGluR1α, and Ca$_V$1.2 in SST INs has profound effects on

the ability of the CA1 microcircuit to encode inputs in *str. L-M*.

Figure 7: Prior GABA$_B$R activation abolishes SST IN inhibition of short-term temporoammonic LTP. **A** Example fEPSP responses recorded in *str. L-M* following temporoammonic stimulation, from control (black) and 20 µM baclofen pre-application (red) slices. Traces are shown for initial baseline (solid line), following *alveus* LTP induction (dashed lines) and following *str. L-M* 1x 100 Hz HFS (grey/pink lines). Right,

summary time-course of all fEPSP recordings normalised to the initial baseline, indicating LTP induction points (lightning bolts). **B** Time-course of LTP induced in *str. L-M* (SLM, lightning bolt), normalised to the 20-30 minutes baseline following *alveus* stimulation in all recordings. **C** Magnitude of fEPSP facilitation at 50-60 minutes post str. L-M HFS in all recordings. **D** Proportion of recordings in both control and baclofen pre-application that successfully induced LTP (>10% facilitation at 50-60 minutes post HFS). **E** Time-course of LTP induced in *str. L-M* normalised to the 20-30 minutes baseline following *alveus* stimulation in recordings where LTP was induced. **F** Quantification of fEPSP facilitation in control and baclofen pre-application slices at 0-1 (left), 20-30 (middle), and 50-60 (right) minutes post *str. L-M* HFS. **G** comparison of fEPSP slope and stimulus strength for all recordings in both groups. Data is shown as either mean ± SEM (**A, B, E**) or box-plots showing 25-75% box with median, with maximum range (**C, F**), proportions (**D**), or as individual data (**g**). Individual cell data is shown overlaid. Statistics shown as: ns – p>0.05, * - p<0.05; from Mann Whitney non-parametric tests.

Activation of GABA$_B$Rs prevents acquisition of contextual fear memories

Finally, as SST IN plasticity has been causally linked to contextual fear conditioning when administered during training[39], we asked if earlier baclofen administration impairs fear learning, For this, we administered mice with 2 mg/kg baclofen (or saline controls, both intraperitoneal injection) 1 hour prior to fear conditioning. Mice were then administered 3 foot-shocks (0.4 nA, 1-minute intervals, 5 minutes total) in the conditioning arena, and the proportion of time spent freezing measured. To determine contextual memory was retained by mice, they were returned to the same arena 24 hours later for 5 minutes (Figure 8A). Overall, saline control mice froze 42 ± 2% of the time during the initial test, increasing to 65 ± 4% freezing upon recall (p=0.004, Holm-Sidak test). By comparison, mice treated with baclofen 1 hour before training froze 17 ± 3% of the time, which did not increase upon recall (20 ± 7%, p=0.63, Holm-Sidak test), but which was markedly lower than saline controls (p<0.0001, Holm Sidak test; Figure 8B). These data confirm that prior administration of baclofen produces sustained impairments in contextual fear conditioning.

This data suggests that prolonged GABA$_B$R activation – as experience with in vivo administration - likely has lasting effects on the ability of hippocampal circuits to perform complex behavioural tasks and may provide important considerations for the use of baclofen therapeutically.

Figure 8: Prior GABA_BR activation impairs contextual memory formation. **A** Schematic of
contextual fear conditioning experiments, in which mice were dosed with 2 mg/kg baclofen
1 hour prior to training. **B** Quantification of % time freezing during the test and recall
phases of conditional fear conditioning. Data is shown as box-plots showing 25-75% box
with median, with maximum range. Individual cell data is shown overlaid. Statistics shown
as: ns – p>0.05, ** - p<0.01, **** - p<0.0001; from 2-way ANOVA with Holm-Sidak post-
tests.

Discussion

In this study we tested the hypothesis that sustained GABA_BR activation leads to receptor
internalisation in hippocampal SST INs, releasing inhibition of synaptic plasticity. We
disproved this hypothesis and show baclofen administration leads to internalisation of
GABA_BRs, Ca_V1.2 and also mGluR1α in SST INs, which is mediated by PP2A-dependent
phosphorylation. This internalisation impairs the long-term ability of SST INs to undergo
synaptic plasticity, which in turn shifts the balance of plasticity at temporoammonic inputs
and impairs contextual fear memory. These data reveal a novel mechanism of long-term
disinhibition that has the potential to shape the plasticity landscape of the CA1 region over
behaviourally relevant time-scales.

PP2A-dependent GABA_BR internalisation regulates SST IN activity

SST INs are a critical element in cortical circuits, gating the strength of inputs, and their
plasticity, arriving from entorhinal cortex [19], and associative inputs within the hippocampus
itself [22,24]. These effects are mediated by a complex interplay of direct inhibitory actions on
postsynaptic CA1 PCs, via presynaptic inhibition of glutamatergic and GABAergic inputs
and through regulation of astrocyte function [40]. Indeed, SST IN activity leads to profound
disinhibition of the hippocampal circuit to strengthen CA3 inputs [22] and limit CA1 PC

dendritic function [24]. Thus, plasticity of SST INs themselves may act to shift the balance of

CA1 function [23], which has a direct effect on CA3-dependent behaviours involving

contextual information [41]. Our work has shown that GABA$_B$Rs acutely inhibit L-type

VGCCs comprising Ca$_V$1.2 subunits on SST INs to inhibit LTP [11], which led to our

hypothesis that GABA$_B$R internalisation would promote LTP induction, as this key

inhibitory mechanism was lost. We have failed to prove this hypothesis, with the opposite

appearing to be the case, due to coincident internalisation of the key LTP machinery,

namely mGluR1α and L-type VGCCs; and which depends on activation of PP2A and that

has been implicated in GABA$_B$R phosphorylation in other brain areas [26,30,31,38].

Following GABA$_B$R activation, adenylyl cyclase activity is inhibited, leading to lower cAMP

levels and reduced PKA activation, which in turn limits the phosphorylation of Serine 892

on the GABA$_{B2}$ subunit, which destabilises the receptor at the cell surface [42]. Additionally,

PP2A dephosphorylates Serine 783 on the GABA$_{B1}$ subunit, which also destabilises the

receptor leading to internalisation [26]. mGluR1α is also a Type C G-protein coupled

receptor (GPCR), thus it is highly plausible that these receptors are targeted for

internalisation through the same intracellular cascade [43], particularly given PP2A has been

shown to regulate recycling of mGluR1 isoforms [44]. For Ca$_V$1.2, the situation is more

complex, as it is internalised upon GABA$_B$R activation which appears to prevented by

PP2A inhibition, while functional currents are not restored. Given the close proximity of

GABA$_{B1}$ to Ca$_V$1.2 in SST IN dendrites[11] it may form part of the GABA$_B$R interactome in

SST INs, as is the case for other Ca$_V$ proteins[29], thus function is impaired when GABA$_B$Rs

are lost. However, Ca$_V$1.2 conductance is regulated by PKA phosphorylation[45], thus, loss

of GABA$_B$R-mediated inhibition may counterbalance a numerical loss of channels. Beyond

VGCCs, native GABA$_B$Rs interact with a variety of transmembrane and cytosolic proteins

in pyramidal cells [28,29], if the same interactome is present in SST INs and whether these

proteins are similarly lost from these cells remain unexplored. Indeed, the functional

association of GABA$_B$Rs with other postsynaptic effectors (e.g. Kir3 channels) may differ in

diverse INs [8,9,12]. As INs make up only approximately 10% of neurons in the hippocampus

[46], efforts to identify the GABA$_B$R interactome using bulk protein samples[29] have likely

underestimated cell-type specific effects. Future recent advances in single-cell proteomic

analysis [47] may allow detection of cell-type specific GABA$_B$R interactomes, crucial to

determine such function. Our data provides limited evidence of impaired postsynaptic

GABA$_B$R function in CA1 pyramidal cells under the same conditions, further suggesting

cell-type specificity.

GABA$_B$R internalisation on SST INs impairs hippocampal circuit function

It has long been appreciated that acute GABA$_B$R activation leads to paradoxical functions at the circuit level, notably enhanced LTP [48,49] and increase seizure activity[50], despite directly hyperpolarising membranes [51,52] and inhibiting pre- and postsynaptic Ca^{2+} channels [53,54] on PCs. These effects point to a function beyond direct inhibition (favouring a disinhibitory model), exemplified by GABA$_B$R inhibition on multiple IN types across the hippocampus [4]. Our data extends this, providing evidence that prolonged GABA$_B$R activation fundamentally shifts the ability of SST INs to undergo LTP on behaviourally relevant timescales. Given that SST IN LTP has been well associated with numerous behavioural functions, including spatial and contextual memory [24,41,55-57], this indicates a potential role of GABA$_B$Rs in regulating this activity. We confirm that LTP in SST INs is crucial to maintain transfer of information to CA1 via the TA path, as previously reported [19], the control of which is impaired when baclofen is pre-applied. Furthermore, we show that GABA$_B$R activation prior to contextual fear conditioning is sufficient to prevent memory encoding. While baclofen has been shown to impair fear conditioning previously [39], these studies relied on administering the drug immediately (<15 minutes) before fear acquisition, suggesting direct inhibitory effects. The pharmacokinetics of baclofen suggest peak brain concentrations with 30 minutes of IP administration, which remain high for several hours [58]. Therefore, it is nigh on impossible to estimate when (and if) loss of SST IN LTP supersedes inhibition driven by GABA$_B$R inhibition. Future studies investigating GABA$_B$R surface localisation on SST INs over hours and days post baclofen injection may reveal the temporal resolution of this relationship, but are outside the scope of this project.

Translational importance

Baclofen is a common and important medicine, primarily prescribed for muscle spasm, but also tested for alcohol dependency, neurodevelopmental disorders, among other conditions (reviewed in [59]), up to 100 mg/day; but is associated with many common neurological side effects. Thus, understanding how baclofen affects cortical circuits may inform its best use, particularly in conditions where GABAergic inhibition has also been implicated.

One of the most common clinical features associated with baclofen administration is the paradoxical induction of seizure activity, both in patients [60] and in rodent models [50].

Originally proposed as a potential anti-seizure medication [61], effects of baclofen on
hippocampal circuits has previously been described, leading to seizure like activity [62]. The
baclofen-dependent loss of SST IN excitatory signalling we observe may favour long-term
reduction in local SST IN inhibition, which would be predicted to cause seizures [63].
Interestingly, we recently showed higher GABA$_B$R signalling in PCs in patients who have
experienced seizures [64]. It is tempting to speculate that such increases may result from
reduced inhibitory tone, leading to compensatory upregulation of receptor expression.
Further study is required to determine if this is the case.

A recent off-label use for high-dose baclofen has been in treating addiction, particularly
alcoholism use disorder [65]. In rodents, GABA$_B$Rs have been shown to be lost in the cortex
following chronic alcohol administration [66] and ventral tegmental area (VTA) following
administration of amphetamine or cocaine [26], albeit not tested in SST INs. However, it has
been shown that acute alcohol administration bidirectionally affects SST INs, with low
doses increasing activity and high doses attenuating activity [67]. SST INs are present in
cortical [15,68] and VTA [69] circuits, and their activity is strongly downregulated in chronic
alcohol consuming mice, which may be corrected by inhibiting them directly[70]. Our data
suggests that prolonged baclofen administration may prevent SST IN activity, akin to
chemogenetic approaches used previously, thus may reflect a mechanism by which
baclofen could produce therapeutic benefit.

Limitations:

We have identified some key limitations of our current study. First, while we show that
internalisation of GABA$_B$Rs from SST IN dendritic membranes occurs following 20 minutes
of baclofen application, using our approach we were not able to identify the specific
temporal dynamics of this loss. If such effects occur rapidly (within seconds) or require
longer (minutes to hours) remains unclear. Nevertheless, baclofen as a therapy is often
administered at moderate to high doses (up to 1g/day) and has a CSF half-life of several
hours [58], thus such temporal dynamics – whilst mechanistically interesting – are somewhat
negated by these pharmacokinetics. Further study identifying whether GABA$_B$R
internalisation on SST INs displays dose- and time-dependence would provide useful
insight to baclofen's therapeutic use. Furthermore, we have not identified how long it takes
SST INs to recover LTP following baclofen administration, which may also provide
important therapeutic information.

One unusual observation in our data was that okadaic acid appeared to occlude labelling

of $GABA_{B1}$ in SDS-FRL experiments, both on SST INs and CA1 PCs (Figure S3). We do

not have a good explanation for this effect. One possibility is that SDS-FRL relies on

immunolabelling of receptors [71] and okadaic acid as a large polypeptide interacts with an

epitope on PP2A close to its interaction site with $GABA_{B1}$. Such an interaction may mask

or occlude antibody binding, that would otherwise not be observed with the use of small

molecule pharmacology, such as fostriecin. A similar phenomenon has been proposed for

the binding of the amyloid precursor protein (APP) to $GABA_{B1}$, which has been indicated in

SDS-FRL to be diminished in APP overexpressing mice[72], but in which mice $GABA_BR$

function is unaltered [73]. These data indicate the importance of providing multiple lines of

evidence to support conclusions based on physiology or anatomy alone, which we have

provided through comparative use of okadaic acid and fostriecin that both selectively

inhibit PP2A [74].

Finally, our data suggest that in SST INs $GABA_BRs$ likely interact with themselves, and

influence the function of – mGluR1α, $Ca_V1.2$, and PP2A. Such proteins do not appear on

previously published interactomes of $GABA_{B1}$ or $GABA_{B2}$ in the hippocampus [29]. Our data

suggest the existence of cell-type-specific interactomes, which may not be detectable at a

whole brain, or even region-specific level. Such an endeavour would almost certainly

identify cell-type specific differences even within neurochemically defined classes, as

highlighted by previous work [4,12]. However, such analyses are beyond the scope of the

current study. Future work should seek to identify which proteins $GABA_BRs$ interact with, in

a cell-type specific manner, and potentially if such interactions change with age or

neuropathology.

Summary and outlook:

Here we show in hippocampal SST INs that prolonged activation of $GABA_BRs$ is sufficient

to cause internalisation of the receptor through PP2A-dependent mechanisms, which also

leads to loss of mGluR1α and $Ca_V1.2$ to prevent synaptic plasticity at CA1 PC inputs.

These data provide compelling evidence of how the inhibitory molecule baclofen exerts a

paradoxical disinhibitory effect by reducing SST IN function, which has relevance for

hippocampal function and disease.

Acknowledgements: We wish to thank Natalie Wernet and Julia Bank for technical
support, BVS technicians for supporting animal husbandy, and members of the Booker,
Vida, and Kulik labs for discussions of the data. Funding was provided by: UK Medical
Research Council (MR/Y014529/1, SAB); DFG (FOR 2134, IV, AK), BIOSS-2 (AK), CRC-
TRR 384/1 (IV, AK) the Patrick Wild Centre (SAB), and the Simons Initiative for the
Developing Brain (SFARI: 529085 - SAB)

Author Contributions: Conceptualisation: SAB, AK; Methodology: NS, SAB, MW, AS,
DL, AK; Validation: SAB, AK; Formal Analysis: SAB, AK, NS, MW, AS; Investigation: SAB,
NS, MW, DL, AS, RL; Writing original draft: SAB; Writing – review and editing: SAB, NS,
IV, AK; Supervision: SAB, IV, AK; Funding acquisition: SAB, IV, AK.

Declaration of interests*:* The authors declare no competing interests.

 Materials and methods:

Animals:

All *ex vivo* electrophysiology was performed in brain slices prepared from male and female
mice (2-4 months old), with all procedures performed according to Home Office (ASPA,
2013) and The University of Edinburgh Ethical Board guidelines. Mice were maintained on
a C57/Bl6J[CRL] background and expressed yellow-fluorescent protein (YFP) under the
vesicular GABA transporter (vGAT)[75]. Mice were housed on a 12h light/dark cycle, with *ad*
libitum access to food and water.

For SDS-FRL, male mice (8-week-old, n=12) either wild-type or in which SST-INs
selectively expressed Channelrhodopsin2(ChR2)-YFP fusion protein were derived from
crossing SST-Cre and Ai32[RCL-ChR2(H134R)/EYFP] transgenic mice. These mice and male
Wistar rats (8-week-old, n = 3) were used for quatitative immuno electron microscopic
analysis. For behavioural analysis, male C57/Bl6J mice (8–12 weeks old, n = 6 per group)
were used. Care and handling of the animals prior to and during the experimental
procedures followed European Union and national regulations (German Animal Welfare
Act) and all experiments were performed in accordance with institutional guidelines
(University of Freiburg, Germany) with permission from local authority (Freiburg, X21/04B).

Acute brain slice preparation:

Brain slices were prepared as previously described [76]. Mice were terminally anaesthetised
with isoflurane, decapitated, then their brains rapidly dissected into ice-cold sucrose-
modified artificial cerebrospinal fluid (sucrose-ACSF; ACSF; in mM: 87 NaCl, 2.5 KCl, 25
NaHCO$_3$, 1.25 NaH$_2$PO$_4$, 25 glucose, 75 sucrose, 7 MgCl$_2$, 0.5 CaCl$_2$) which was
saturated with carbogen (95% O$_2$/5% CO$_2$). Brains were glued to a vibratome stage (Leica
VT1200S, Leica, Germany), then 300 μm (for whole-cell recordings) or 500 μm (for
extracellular recordings) horizontal slices containing the hippocampi were prepared.
Following slicing, brain slices were placed in either: a submerged holding chamber
containing sucrose-ACSF warmed to 35 °C for 30 min, then at room temperature; or on
small squares of filter paper placed in a liquid/gas interface chamber, containing recording
ACSF (in mM: 125 NaCl, 2.5 KCl, 25 NaHCO$_3$, 1.25 NaH$_2$PO$_4$, 25 glucose, 1 MgCl$_2$, 2
CaCl$_2$) and bubbled with carbogen.

Whole-cell patch clamp recordings:

Briefly, 400 μm horizontal hippocampal slices were prepared from 60-90 day-old C57/Bl6J mice as previously described[77]. For recording, slices were transferred to a submerged chamber maintained at 31 ± 1°C. Whole-cell recordings were made using pipettes filled with either K-gluconate or Cs-gluconate based solutions. Voltage-clamp recordings were performed from a potential of either -65 mV or -60 mV, while current-clamp recordings were made from resting membrane potential. Intrinsic properties of recorded neurons were characterized on their voltage response to hyper- to depolarizing current steps, in current-clamp. HVA calcium transients, mGluR EPSCs, and GABA$_B$R-mediated currents were measured in the presence of 10 μM CNQX, 50 μM DL-AP-5, and 50 μM picrotoxin. GABA$_A$R IPSCs were elicited by placing a bipolar stimulating electrode in str. lacunosum-moleculare approximately 200-300 μm rostral to the recorded neuron, in the presence of DL- 10 μM CNQX, 50 μM DL-AP-5. LTP was induced at alveus inputs to SST INs with EPSCs elicited by a bipolar stimulating electrode placed in the alveus, in the presence of 50 μM picrotoxin. Following baseline, associative LTP was induced by theta-burst stimulation paired with a depolarization to -20 mV, repeated 3 times at 30 second intervals. EPSCs were measured for 25 minutes following induction. For all stimulation, the pulse duration was 200 μs delivered via a constant voltage generator. For HVA VGCC activation, once a whole-cell patch-clamp recording was obtained using a Cs-Gluconate internal solution (containing QX-314), the cell was allowed to dialyse for 5 minutes. Then at -60 mV voltage-clamp, 3 families of depolarizing steps were applied (-60 to 20 mV, 5 mV steps, 500 ms duration). The L-type VGCC blocker nifedipine (20 μM) was then applied to the bath for 3 minutes and 3 further families of depolarizing steps obtained. All traces were then P/N subtracted based on the 5 mV step and the peak current measured. The peak current under nifedipine was then subtracted from control. For mGluR EPSC experiments, slow-EPSCs were elicited by focal puff application of the Group 1 mGluR agonist DPHG (100 μM, 500 ms, 20 psi, diluted from stock in 150 mM NaCl) delivered via a patch pipette placed ~50 μm from the cell body. The specificity of the mGluR current was assessed by subsequent bath application of the mGluR1α-specific antagonist LY-367,385 (100 μM).

For pre-application experiments, slices were transferred to a holding chamber containing recording ACSF and 20 μM R-baclofen for 20 minutes, slices were then placed in the recording chamber and rinsed with ACSF without R-baclofen for 4-6 minutes prior to recording. To block PP2A activity, slices were transferred to a chamber containing ACSF and 1 μM Okadaic Acid for at least 3 hours prior to recording or baclofen pre-application. All other drugs were applied directly to the perfusing ACSF.

Field recordings:

For field excitatory postsynaptic potential (fEPSP) recordings, hippocampal slices were transferred to an interface recording chamber perfused with carbogenated artificial cerebrospinal fluid (ACSF) at 2–3 mL/min and maintained at 30 ± 1 °C. Recording electrodes (1–3 MΩ) were pulled from borosilicate glass capillaries (1.5 mm outer / 0.86 mm inner diameter; Harvard Apparatus, UK) using a horizontal puller and filled with recording ACSF. Slices were visualised with a wide-field microscope (Leica, Germany), and electrodes were positioned in the str. L-M of CA1.

Extracellular fEPSPs were evoked using a paired-pulse protocol delivered via a bipolar stimulating electrode placed in *str. L-M*, approximately 500 µm to 1 mm from the recording electrode, to activate the temporoammonic pathway. A second stimulating electrode was positioned in parallel within the alveus of CA1. To activate GABA¬B receptors, slices were perfused with 20 µM baclofen for 20 min, followed by a 20-min washout period.

A 10-min baseline was recorded under control conditions or following baclofen washout. LTP was first induced via alveus stimulation using a high-frequency stimulation (HFS) protocol consisting of three 100 Hz trains of 20 pulses delivered 30 s apart, or a protocol consisting of five 100 Hz trains of 5 pulses, delivered 30 s apart. LTP was then induced in str. L-M using an HFS protocol of 1x 100 Hz train of 100 pulses delivered 30 s apart.

Potentiation was monitored for 60 min following induction. LTP magnitude was quantified as the mean fEPSP slope measured 50–60 min post-induction, normalised to the 10-min baseline. LTP was considered successful if the 50–60 min slope exceeded baseline by >10%. Recordings were excluded if baseline stability, assessed by comparing mean slopes at 1–2 min and 9–10 min, varied ±10%. Signals were filtered online (1 Hz high-pass, 500 Hz low-pass) and digitised at 10 kHz. Data were acquired and analysed offline using WinLTP (v3.01, University of Bristol, UK).

Neuronal visualization and immunohistochemistry:

All neurons were filled with biocytin during recording, fixed overnight in 4% PFA and labelled with streptavidin and antibodies to SST according to previous methods[78]. Slices were washed in PBS then blocked for 1 hour at room temperature (10% NGS, 0.3% Triton X-100 and 0.05% NaN_3 in PBS). Following blocking slices were incubated with primary antibodies at 4°C for 48h (1:1000 rabbit SST-14, T-4103, BMA Biomedicals, Switzerland;

5% NGS, 0.3% Triton X-100 and 0.05% NaN$_3$ in PBS), washed in PBS then transferred to

a secondary antibody solution containing fluorescent-conjugated streptavidin (1:500 goat

anti-rabbit AlexaFluor 488, 1:500 Streptavidin AlexaFluor 633, 3% NGS, 0.1% Triton X-

100, 0.05% NaN$_3$). Following labeling the slices were washed in 0.1M PB and mounted on

glass slides with Vectashield HardSet mounting medium (Vector Labs, UK). To identify

SST Ins Confocal image stacks of recorded neurons were taken on a Leica SP8 confocal

microscope using a 63x (1.4 NA) objective at 1024 × 1024 resolution (Z step size 1 μm).

Electron microscopy:

Acute slice preparation and pharmacology for SDS-FRL

Acute hippocampal slices were prepared as previously described [76]. Animals were sedated

with isoflurane then they were decapitated and the brains were rapidly dissected and

chilled in semi-frozen carbogenated (95% O2/5% CO2) sucrose-substituted artificial

cerebrospinal fluid (sucrose-ACSF, in mM: 87 NaCl, 2.5 KCl, 25 NaHCO3, 1.25 NaH2PO4,

25 glucose, 75 sucrose, 7 MgCl2, 0.5 CaCl2, 1 Na-pyruvate, 1 Na-ascorbate). Transverse

hippocampal slices (200 μm thick) were cut on a vibratome (VT1200s, Leica, Germany)

and stored submerged in sucrose-ACSF warmed to 35°C for at least 30 min and

subsequently at room temperature (RT). Both solutions were equilibrated with 95% O2 and

5% CO2 gas mixture throughout experiments. Acute slices were divided into four groups

for pharmacological treatment: 1, incubated in ACSF (control, 30 min), 2, ACSF + GABA$_B$R

agonist baclofen (Bac, 20 μM, 20 min), 3, ACSF + protein phosphatase 2A inhibitor

fostriecin (FS, 50 nM, 90 min), 4, ACSF + fostriecin + baclofen (FS/Bac, FS 50 nM 90 min

+ Bac 20 μM, 20 min). After pharmacological manipulations, slices were transferred into

fixative containing 1% paraformaldehyde and 15% saturated picric acid made up in 0.1 M

phosphate buffer (PB) overnight (O/N) at 4 °C.

SDS-FRL immuneoelctron microscopy

After fixation, slices were cryoprotected in 30% glycerol in 0.1M PB O/N at 4°C [10]. Blocks

containing stratum oriens-alveus were trimmed from the slices and frozen under high-

pressure (HPM 100, Leica). The frozen samples were fractured at -140°C and the

fractured faces were coated with carbon, (5 nm), platinum-carbon (2 nm) and carbon (18

nm) in a freeze-fracture replica machine (ACE 900, Leica). Replicas were digested at 80°C

in a solution containing 2.5% SDS and 20% sucrose diluted in 15 mM Tris buffer (TB, pH

8.3) for 20 hr. Subsequently, replicas were washed in washing buffer containing 0.05%

bovine serum albumin (BSA, Roth, Germany) and 0.1% Tween 20 in 50 mM Tris-buffered saline (TBS) and then blocked in a solution containing 5% BSA and 0.1% Tween 20 in TBS for 1 hr at RT. Afterwards, replicas prepared from SST-cre mice were incubated in the following mixtures of primary antibodies in a solution containing 1% BSA and 0.1% Tween 20 made up in TBS for 3 days at 15°C: (i) $GABA_{B1}$ (B17, rabbit, 10 μg/ml; Kulik et al., 2002) and green fluorescence protein (GFP-1010, chicken 0.4 μg/ml, Aves Labs, Oregon; Booker et al., 2020) or (ii) $Ca_v1.2$ (mouse, 50 μg/ml, NeuroMab Facility, California; Booker et al., 2018) and GFP. Replicas were washed then reacted with 6 nm ($GABA_{B1}$ and $Ca_v1.2$) and 18 nm (GFP) gold particle-conjugated secondary antibodies (1:30, Jackson ImmunoResearch Europe, Cambridgeshire) O/N at 15°C. The replicas prepared from WT mice were incubated in $Ca_v1.2$ (mouse, 50 μg/ml, NeuroMab Facility, California), GABAB1 (B17, rabbit, 10 μg/ml) and metabotropic glutamate receptor 1α-subunit (mGluR1α, Guinea pig, 0.4 μg/ml, Frontier Institute, Hokkaido; Booker et al., 2018) in a solution containing 1% BSA and 0.1% Tween 20 made up in TBS for 3 days at 15°C. Replicas were washed then reacted with 5 nm ($GABA_{B1}$) 10nm ($Ca_v1.2$) and 18 nm (mGluR1α) gold particle-conjugated secondary antibodies (1:30, Jackson ImmunoResearch Europe, Cambridgeshire) O/N at 15°C. Finally, replicas were washed in TBS then distilled water and mounted on 100-mesh grids.

Electron microscopy and quantitative analysis of protein density

Labeled replicas were analyzed under an electron microscope (JEM 2100 Plus). Since all antibodies target intracellular protein epitopes, immunoreactivity can be detected on the plasma membrane's protoplasmic face (P-face) but not the exoplasmic face (E-face). The surface density of receptors and ion channels was determined by dividing the absolute number of particles labeling molecules of interest by the surface of the respective segments of the dendritic shafts of GFP-positive SST-INs on replicas.

Behavioural analysis:

To investigate the role of $GABA_BR$ activation in fear learning, we performed a fear conditioning experiment. Male C57/Bl6J mice (8–12 weeks old, n = 6 per group) were handled and habituated. On the day of the test, mice were given a two-minute baseline period before receiving three-foot shocks (0.06mA) spaced one minute apart. Following the last shock, the mice remained in the fear conditioning arena for an additional minute, then they were transferred to the home cage. The entire training session lasted for approx.

five minutes. Mice were divided into test and control groups. Test group were injected with

baclofen (i.p., 2 mg/kg), while the control group received a vehicle (0.9% NaCl). One hour

following the baclofen or vehicle injection, freezing time was measured in the same context

as recall in order to access fear memory.

Statistical analysis:

For *ex vivo* electrophysiology, one or two cells or slices were recorded per animal per

treatment, limiting the ability to assess intra-animal variability. As such cells and slices

were treated as independent replicates and statistical analyses were conducted on these

data. Parametric and non-parametric tests were applied as appropriate, including two-way

ANOVA, Student's t-test, Mann–Whitney U-tests, and Wilcoxon signed-rank tests. For

SDS-FRL, where many dendritic profiles were sampled per animal replicate, we performed

linear-mixed effects modelling (LMM) of using the "lmer" function to limit the potential

inclusion of pseudoreplication. Animal was assigned as random effects, then type III

ANOVA (Satterthwaites method) was performed on the model to assess main effects (e.g.

drug treatment). If a statistical interaction between main effects was observed, then post-

hoc analysis was performed (Tukey post hoc tests). For all comparisons data is shown as

box-plots with median and 25–75% percentiles displayed in the box; minimum and

maximum extents are shown as whiskers, with individual data points shown from cell or

animal averages. For timecourses and current-voltage plots, data is shown as

mean ± SEM. Statistical significance was defined as $p < 0.05$. Statistical tests and

graphing were performed using either GraphPad Prism (GraphPad Software v10.4.1, San

Diego, CA, United States) or R-Studio.

References:

References:

1 Bartos, M., Vida, I. & Jonas, P. Synaptic mechanisms of synchronized gamma
oscillations in inhibitory interneuron networks. *Nature reviews neuroscience* **8**, 45-56
(2007).

2 Burgess, N. & O'Keefe, J. Neuronal computations underlying the firing of place cells and
their role in navigation. *Hippocampus* **6**, 749-762 (1996).

3 Feldman, D. E. The spike-timing dependence of plasticity. *Neuron* **75**, 556-571 (2012).

4 Kulik, Á., Booker, S. A. & Vida, I. Differential distribution and function of GABABRs in
somato-dendritic and axonal compartments of principal cells and interneurons in cortical
circuits. *Neuropharmacology* **136**, 80-91 (2018).

5 Guet-McCreight, A., Skinner, F. K. & Topolnik, L. Common principles in functional
organization of VIP/calretinin cell-driven disinhibitory circuits across cortical areas.
Frontiers in neural circuits **14**, 32 (2020).

6 Williams, L. E. & Holtmaat, A. Higher-order thalamocortical inputs gate synaptic long-
term potentiation via disinhibition. *Neuron* **101**, 91-102. e104 (2019).

7 Booker, S. A. *et al.* Differential surface density and modulatory effects of presynaptic
GABAB receptors in hippocampal cholecystokinin and parvalbumin basket cells. *Brain*
Structure and Function **222**, 3677-3690 (2017).

8 Booker, S. A. *et al.* KCTD12 auxiliary proteins modulate kinetics of GABAB receptor-
mediated inhibition in cholecystokinin-containing interneurons. *Cerebral Cortex* **27**, 2318-
2334 (2017).

9 Booker, S. A. *et al.* Differential GABAB-receptor-mediated effects in perisomatic-and
dendrite-targeting parvalbumin interneurons. *Journal of Neuroscience* **33**, 7961-7974
(2013).

10 Booker, S. A. *et al.* Presynaptic GABAB receptors functionally uncouple somatostatin
interneurons from the active hippocampal network. *Elife* **9**, e51156 (2020).

11 Booker, S. A. *et al.* Postsynaptic GABABRs inhibit L-type calcium channels and abolish
long-term potentiation in hippocampal somatostatin interneurons. *Cell reports* **22**, 36-43
(2018).

12 Degro, C. E., Vida, I. & Booker, S. A. Postsynaptic GABAB-receptor mediated currents
in diverse dentate gyrus interneuron types. *Hippocampus* **34**, 551-562 (2024).

13 Mott, D. D., Li, Q., Okazaki, M. M., Turner, D. A. & Lewis, D. V. GABAB-receptor–
mediated currents in interneurons of the dentate-hilus border. *Journal of neurophysiology*
82, 1438-1450 (1999).

14 Price, C. J., Scott, R., Rusakov, D. A. & Capogna, M. GABAB receptor modulation of
feedforward inhibition through hippocampal neurogliaform cells. *Journal of Neuroscience*
28, 6974-6982 (2008).

15 Booker, S. A. & Vida, I. Morphological diversity and connectivity of hippocampal
interneurons. *Cell and tissue research* **373**, 619-641 (2018).

16 Ali, A. B. & Thomson, A. M. Facilitating pyramid to horizontal oriens-alveus interneurone
inputs: dual intracellular recordings in slices of rat hippocampus. *The Journal of physiology*
507, 185-199 (1998).

17 Lacaille, J.-C., Mueller, A. L., Kunkel, D. D. & Schwartzkroin, P. A. Local circuit
interactions between oriens/alveus interneurons and CA1 pyramidal cells in hippocampal
slices: electrophysiology and morphology. *Journal of Neuroscience* **7**, 1979-1993 (1987).

18 Müller, C. & Remy, S. Dendritic inhibition mediated by O-LM and bistratified
interneurons in the hippocampus. *Frontiers in synaptic neuroscience* **6**, 23 (2014).

19 Yanovsky, Y., Sergeeva, O., Freund, T. & Haas, H. Activation of interneurons at the
stratum oriens/alveus border suppresses excitatory transmission to apical dendrites in the
CA1 area of the mouse hippocampus. *Neuroscience* **77**, 87-96 (1997).

20 Lovett-Barron, M. *et al.* Regulation of neuronal input transformations by tunable
dendritic inhibition. *Nature neuroscience* **15**, 423-430 (2012).

21 Singh, S. & Topolnik, L. Inhibitory circuits in fear memory and fear-related disorders.
Frontiers in neural circuits **17**, 1122314 (2023).

22 Leão, R. N. *et al.* OLM interneurons differentially modulate CA3 and entorhinal inputs to
hippocampal CA1 neurons. *Nature neuroscience* **15**, 1524-1530 (2012).

23 Udakis, M., Pedrosa, V., Chamberlain, S. E., Clopath, C. & Mellor, J. R. Interneuron-
specific plasticity at parvalbumin and somatostatin inhibitory synapses onto CA1 pyramidal
neurons shapes hippocampal output. *Nature communications* **11**, 4395 (2020).

24 Udakis, M., Claydon, M. D., Zhu, H. W., Oakes, E. C. & Mellor, J. R. Hippocampal OLM
interneurons regulate CA1 place cell plasticity and remapping. *Nature Communications* **16**,
9912 (2025).

25 Dinamarca, M. C. *et al.* Complex formation of APP with GABAB receptors links axonal
trafficking to amyloidogenic processing. *Nature communications* **10**, 1331 (2019).

26 Padgett, C. L. *et al.* Methamphetamine-evoked depression of GABAB receptor signaling
in GABA neurons of the VTA. *Neuron* **73**, 978-989 (2012).

27 Raveh, A., Turecek, R. & Bettler, B. in *Advances in Pharmacology* Vol. 73   145-165
(Elsevier, 2015).

28 Schwenk, J. *et al.* Native GABAB receptors are heteromultimers with a family of
auxiliary subunits. *Nature* **465**, 231-235 (2010).

29 Schwenk, J. *et al.* Modular composition and dynamics of native GABAB receptors
identified by high-resolution proteomics. *Nature neuroscience* **19**, 233-242 (2016).

30 Hearing, M. *et al.* Repeated cocaine weakens GABAB-Girk signaling in layer 5/6
pyramidal neurons in the prelimbic cortex. *Neuron* **80**, 159-170 (2013).

31 Li, X. *et al.* Direct interaction of PP2A phosphatase with GABAB receptors alters
functional signaling. *Journal of Neuroscience* **40**, 2808-2816 (2020).

32 Hirono, M., Yoshioka, T. & Konishi, S. GABAB receptor activation enhances mGluR-
mediated responses at cerebellar excitatory synapses. *Nature neuroscience* **4**, 1207-1216
(2001).

33 Tabata, T. *et al.* Ca2+ activity at GABAB receptors constitutively promotes metabotropic
glutamate signaling in the absence of GABA. *Proceedings of the National Academy of*
Sciences **101**, 16952-16957 (2004).

34 Baude, A. *et al.* The metabotropic glutamate receptor (mGluRIα) is concentrated at
perisynaptic membrane of neuronal subpopulations as detected by immunogold reaction.
Neuron **11**, 771-787 (1993).

35 Blasco-Ibáñez, J. & Freund, T. Synaptic input of horizontal interneurons in stratum
oriens of the hippocampal CA1 subfield: structural basis of feed-back activation. *European*
Journal of Neuroscience **7**, 2170-2180 (1995).

36 Topolnik, L., Azzi, M., Morin, F., Kougioumoutzakis, A. & Lacaille, J. C. mGluR1/5
subtype-specific calcium signalling and induction of long-term potentiation in rat
hippocampal oriens/alveus interneurones. *The Journal of physiology* **575**, 115-131 (2006).

37 Vasuta, C. *et al.* Metaplastic regulation of CA1 Schaffer collateral pathway plasticity by
Hebbian MGluR1a-mediated plasticity at excitatory synapses onto somatostatin-
expressing interneurons. *eneuro* **2** (2015).

38 Terunuma, M. *et al.* Prolonged activation of NMDA receptors promotes
dephosphorylation and alters postendocytic sorting of GABAB receptors. *Proceedings of*
the National Academy of Sciences **107**, 13918-13923 (2010).

39 Heaney, C. F. *et al.* Baclofen administration alters fear extinction and GABAergic
protein levels. *Neurobiology of learning and memory* **98**, 261-271 (2012).

40 Matos, M. *et al.* Astrocytes detect and upregulate transmission at inhibitory synapses of
somatostatin interneurons onto pyramidal cells. *Nature communications* **9**, 4254 (2018).

41 Honoré, E. & Lacaille, J.-C. Object location learning in mice requires hippocampal
somatostatin interneuron activity and is facilitated by mTORC1-mediated long-term
potentiation of their excitatory synapses. *Molecular Brain* **15**, 101 (2022).

42 Couve, A. *et al.* Cyclic AMP–dependent protein kinase phosphorylation facilitates
GABAB receptor–effector coupling. *Nature neuroscience* **5**, 415-424 (2002).

43 Pin, J.-P. & Bettler, B. Organization and functions of mGlu and GABAB receptor
complexes. *Nature* **540**, 60-68 (2016).

44 Sistiaga, A. & Sánchez-Prieto, J. Protein phosphatase 1 and 2A inhibitors prolong the
switch in the control of glutamate release by group I metabotropic glutamate receptors:
characterization of the inhibitory pathway. *Journal of neurochemistry* **75**, 1566-1574
(2000).

45 Minobe, E. *et al.* A new phosphorylation site in cardiac L-type Ca2+ channels (Cav1. 2)
responsible for its cAMP-mediated modulation. *American Journal of Physiology-Cell*
Physiology **307**, C999-C1009 (2014).

46 Freund, T. F. & Buzsáki, G. Interneurons of the hippocampus. *hippocampus* **6**, 347-470
(1996).

47 Karlsson, F. *et al.* Single-Cell Protein Interactomes by the Proximity Network Assay.
bioRxiv, 2025.2006.2019.660329 (2025). https://doi.org/10.1101/2025.06.19.660329

48 Davies, C. H., Starkey, S. J., Pozza, M. F. & Collingridge, G. L. GABAB autoreceptors
regulate the induction of LTP. *Nature* **349**, 609-611 (1991).

49 Mott, D. D. & Lewis, D. V. Facilitation of the induction of long-term potentiation by
GABAB receptors. *Science* **252**, 1718-1720 (1991).

50 Mott, D. D., Bragdon, A. C., Lewis, D. V. & Wilson, W. A. Baclofen has a proepileptic
effect in the rat dentate gyrus. *The Journal of Pharmacology and Experimental*
Therapeutics **249**, 721-725 (1989).

51 Lüscher, C., Jan, L. Y., Stoffel, M., Malenka, R. C. & Nicoll, R. A. G protein-coupled
inwardly rectifying K+ channels (GIRKs) mediate postsynaptic but not presynaptic
transmitter actions in hippocampal neurons. *Neuron* **19**, 687-695 (1997).

52 Sodickson, D. L. & Bean, B. P. GABAB receptor-activated inwardly rectifying potassium
current in dissociated hippocampal CA3 neurons. *Journal of Neuroscience* **16**, 6374-6385
(1996).

53 Chalifoux, J. R. & Carter, A. G. GABAB receptor modulation of voltage-sensitive calcium
channels in spines and dendrites. *Journal of Neuroscience* **31**, 4221-4232 (2011).

54 Thompson, S. M. & Gähwiler, B. Comparison of the actions of baclofen at pre-and
postsynaptic receptors in the rat hippocampus in vitro. *The Journal of physiology* **451**, 329-
345 (1992).

55 Grigoryan, G. *et al.* Synaptic plasticity at the dentate gyrus granule cell to somatostatin-
expressing interneuron synapses supports object location memory. *Proceedings of the*
National Academy of Sciences **120**, e2312752120 (2023).

56 Hainmueller, T., Cazala, A., Huang, L.-W. & Bartos, M. Subfield-specific interneuron
circuits govern the hippocampal response to novelty in male mice. *Nature communications*
15, 714 (2024).

57 Yuan, M. *et al.* Predictive goal coding by dentate gyrus somatostatin-expressing
interneurons in male mice. *Nature Communications* **16**, 5382 (2025).

58 Balerio, G. N. & Rubio, M. C. Pharmacokinetic-pharmacodynamic modeling of the
antinociceptive effect of baclofen in mice. *European journal of drug metabolism and*
pharmacokinetics **27**, 163-169 (2002).

59 Romito, J. W. *et al.* Baclofen therapeutics, toxicity, and withdrawal: a narrative review.
SAGE open medicine **9**, 20503121211022197 (2021).

60 Kofler, M., Kronenberg, M., Rifici, C., Saltuari, L. & Bauer, G. Epileptic seizures
associated with intrathecal baclofen application. *Neurology* **44**, 25-25 (1994).

61 Ault, B. & Nadler, J. V. Anticonvulsant-like actions of baclofen in the rat hippocampal
slice. *British journal of pharmacology* **78**, 701-708 (1983).

62 Dugladze, T. *et al.* GABAB autoreceptor-mediated cell type-specific reduction of
inhibition in epileptic mice. *Proceedings of the National Academy of Sciences* **110**, 15073-
15078 (2013).

63 Drexel, M., Rahimi, S. & Sperk, G. Silencing of Hippocampal Somatostatin Interneurons
Induces Recurrent Spontaneous Limbic Seizures in Mice. *Neuroscience* **487**, 155-165
(2022). https://doi.org/https://doi.org/10.1016/j.neuroscience.2022.02.007

64 Wilson, M. A. *et al.* Phylogenetic divergence of GABAB receptor signaling in neocortical
networks over adult life. *Nature Communications* **16**, 4194 (2025).

65 de Beaurepaire, R. *et al.* The Use of Baclofen as a Treatment for Alcohol Use Disorder:
A Clinical Practice Perspective. *Frontiers in Psychiatry* **Volume 9 - 2018** (2019).
https://doi.org/10.3389/fpsyt.2018.00708

66 Peng, S.-Y. *et al.* Reduced motor cortex GABABR function following chronic alcohol
exposure. *Molecular Psychiatry* **26**, 383-395 (2021). https://doi.org/10.1038/s41380-020-
01009-6

67 Li, M. *et al.* Alcohol reduces the activity of somatostatin interneurons in the mouse
prefrontal cortex: A neural basis for its disinhibitory effect? *Neuropharmacology* **188**,
108501 (2021). https://doi.org/https://doi.org/10.1016/j.neuropharm.2021.108501

68 Rudy, B., Fishell, G., Lee, S. & Hjerling-Leffler, J. Three groups of interneurons account
for nearly 100% of neocortical GABAergic neurons. *Developmental Neurobiology* **71**, 45-
61 (2011). https://doi.org/https://doi.org/10.1002/dneu.20853

69 Nagaeva, E. *et al.* Heterogeneous somatostatin-expressing neuron population in mouse
ventral tegmental area. *eLife* **9**, e59328 (2020). https://doi.org/10.7554/eLife.59328

70 Dao, N. C., Brockway, D. F., Suresh Nair, M., Sicher, A. R. & Crowley, N. A.
Somatostatin neurons control an alcohol binge drinking prelimbic microcircuit in mice.
Neuropsychopharmacology **46**, 1906-1917 (2021). https://doi.org/10.1038/s41386-021-
01050-1

71 Masugi-Tokita, M. & Shigemoto, R. High-resolution quantitative visualization of
glutamate and GABA receptors at central synapses. *Current opinion in neurobiology* **17**,
387-393 (2007).

72 Martín-Belmonte, A. *et al.* Nanoscale alterations in GABAB receptors and GIRK channel
organization on the hippocampus of APP/PS1 mice. *Alzheimer's research & therapy* **14**,
136 (2022).

73 Meftah, S. *et al.* GABAB Receptor signaling in CA1 Pyramidal Cells is not Regulated by
Aging in the APP/PS1 Mouse Model of Amyloid Pathology. *eneuro* **13** (2026).

74 Neale, D. A., Morris, J. C., Verrills, N. M. & Ammit, A. J. Understanding the regulatory
landscape of protein phosphatase 2A (PP2A): Pharmacological modulators and potential
therapeutics. *Pharmacology & Therapeutics* **269**, 108834 (2025).

75 Wang, Y. *et al.* Fluorescent labeling of both GABAergic and glycinergic neurons in
vesicular GABA transporter (VGAT)–Venus transgenic mouse. *Neuroscience* **164**, 1031-
1043 (2009).

76Oliveira, L. S., Sumera, A. & Booker, S. A. Repeated whole-cell patch-clamp recording
from CA1 pyramidal cells in rodent hippocampal slices followed by axon initial segment
labeling. *STAR protocols* **2** (2021).

77Oliveira, L. S., Sumera, A. & Booker, S. A. Repeated whole-cell patch-clamp recording
from CA1 pyramidal cells in rodent hippocampal slices followed by axon initial segment
labeling. *STAR protocols* **2**, 100336 (2021).

78Yang, Y. *et al.* Identifying foetal forebrain interneurons as a target for monogenic autism
risk factors and the polygenic 16p11. 2 microdeletion. *BMC neuroscience* **24**, 5 (2023).