Demographic history shapes forest tree vulnerability to climate change

Thomas Francisco[1,*], Isabelle Lesur-Kupin[1], Carlos Guadaño-Peyrot[2], Sanna Olsson[2], Marija Kravanja[3], Marjana Westergren[3], Sara Pinosio[4], Thibaut Capblancq[5], Giovanni G. Vendramin[4], Katharina B. Budde[6], Lene R. Nielsen[7], James M. Doonan[7], Delphine Grivet[2], Elia Vajana[4], Juliette Archambeau[8], Andrea Piotti[4], Santiago C. González-Martínez[1,*]

[1]National Research Institute for Agriculture, Food and the Environment (INRAE), University of Bordeaux, UMR 1202 BIOGECO, 33610 Cestas, France

[2]Department of Forest Ecology and Genetics, Institute of Forest Sciences (ICIFOR-INIA), CSIC, Madrid, Spain

[3]Slovenian Forestry Institute, Večna pot 2, 1000, Ljubljana, Slovenia

[4]Institute of Biosciences and BioResources (IBBR), National Research Council (CNR), 50019 Sesto Fiorentino (Firenze), Italy

[5]Laboratoire d'Écologie Alpine, Université Grenoble Alpes, Université Savoie Mont-Blanc, CNRS, 38058 Grenoble, France

[6]Northwest German Forest Research Institute, Professor-Oelkers-Strasse 6, 34346 Hann. Münden, Germany

[7]Department of Geosciences and Natural Resource Management, University of Copenhagen, 1958 Frederiksberg C, Copenhagen, Denmark

[8]National Research Institute for Agriculture, Food and the Environment (INRAE), UMR 1391 ISPA, 88140 Villenave d'Ornon, France

*Corresponding authors:

Thomas Francisco (thomas.francisco@inrae.fr) and Santiago C. González-Martínez

(santiago.gonzalez-martinez@inrae.fr)

INRAE UMR BIOGECO 1202,

DOMAINE DE L'HERMITAGE

69 Route d'Arcachon

33612 Cestas, France

ORCID Number

Thomas Francisco: 0009-0005-8618-0747, thomas.francisco@inrae.fr

Isabelle Lesur-Kupin: 0000-0001-7632-3955, isabelle.lesurkupin@inrae.fr

Carlos Guadaño-Peyrot: 0009-0007-8663-5216, carlospeyrot77@gmail.com

Sanna Olsson: 0000-0002-1199-4499, sanna.olsson@inia.csic.es

Marija Kravanja: 0009-0002-1478-6457, marija.kravanja@gozdis.si

Marjana Westergren: 0000-0002-4204-0161, marjana.westergren@gozdis.si

Sara Pinosio: 0000-0002-1820-3637, sara.pinosio@cnr.it

Thibaut Capblancq: 0000-0001-5024-1302, thibaut.capblancq@univ-grenoble-alpes.fr

Giovanni G. Vendramin: 0000-0001-9921-7872, giovannigiuseppe.vendramin@cnr.it

Katharina B. Budde: 0000-0002-5361-2815, katharina.budde@nw-fva.de

Lene R. Nielsen: 0000-0002-7214-8691, lron@ign.ku.dk

James Doonan: 0000-0002-0523-5808, jd@ign.ku.dk

Delphine Grivet: 0000-0001-8168-4456, dgrivet@inia.csic.es

Elia Vajana: 0000-0003-1340-3389, eliavajana@cnr.it

Juliette Archambeau: 0000-0003-0155-9538, juliette.archambeau@inrae.fr

Andrea Piotti: 0000-0002-3324-5325, andrea.piotti@cnr.it

Santiago C. González-Martínez: 0000-0002-4534-3766, santiago.gonzalez-martinez@inrae.fr

Abstract

Demographic history is expected to play a central role in shaping population vulnerability to climate

change through its lasting effects on effective population sizes and genetic connectivity. However,

existing studies report contrasting outcomes, and the consequences of alternative demographic

histories have seldom been assessed concurrently across multiple taxa. Here, we analysed population

genomic data from six of the major European forest tree species to address this gap. Across species,

genetic isolation reduced adaptive potential, as reflected in lower standing genetic diversity. Greater

genetic differentiation was also associated with increased maladaptation to climate, whereas the

purge or accumulation of deleterious mutations depended on the severity of past demographic

events. Finally, the beneficial effects of gene flow were evidenced across the six species by more

optimal climate adaptation in highly connected populations. Altogether, our results provide valuable

insights into how genetic differentiation, reflecting the combined effects of genetic drift and limited

historical gene flow, influences current vulnerability of forest tree populations to climate change.

Introduction

Trees are foundation species in many terrestrial ecosystems, supporting remarkable biodiversity, regulating the climate and providing other vital ecosystem services on a global scale[1]. Due to their long generation time and sessile nature, trees are particularly exposed to environmental hazards and gradual changes. An increasing body of literature indicates that they will not be able to adapt and/or disperse at rates matching projected climate change[2-4] and forests worldwide are already displaying signs of stress and decline[5,6]. The way in which populations respond to environmental changes depends on the intensity of the stress they experience, their sensitivity to this stress and their capacity to adapt to the new conditions, which together determine their vulnerability to a changing environment[7,8].

Genetic metrics are increasingly being used to assess vulnerability to climate change, as they capture impacts and responses at the intraspecific level[9-11]. Genetic diversity originates diverse phenotypes, some of which may confer advantages in shifting climate, thus forming the basis of the evolutionary potential of populations and, possibly, different patterns of local adaptation[12]. Although local adaptation is common in plants[13], the degree of local adaptation can be population-specific[14]. Both adaptation lags to current climate and pre-adaptation to future climate can influence the extent to which a population will be affected by climate change[15,16]. Recent advances in ecological genomics have introduced new metrics to quantify the extent of climate adaptation in populations (e.g., the genomic discrepancy index, GDI[17]). Genetic variation between populations resulting from different rates of accumulation and purging of deleterious mutations, can also shape their sensitivity to climate change. Indeed, levels of genetic load, defined as the reduction of fitness in a population due to deleterious mutations, can further constrain a population's ability to persist under shifting climatic pressures[18]. Additionally, inbreeding depression has been reported to be particularly high in gymnosperms compared to angiosperms[19,20], as evidenced, for example, by the high number of lethal equivalents affecting tree fitness observed in conifers[20-24]. Overall, regardless of their impact on

current fitness or adaptive potential, population attributes such as genetic diversity, climate

adaptation, or genetic load are shaped by demographic processes, and are therefore neither static

through time nor uniform across populations[23-27].

Over evolutionary timescales, tree populations have been profoundly shaped by repeated glacial

cycles, large-scale environmental shifts and, more recently, land-use changes. These events triggered

large distributional shifts that came with important variation in population size and connectivity, and

thereby shaped the genetic makeup of populations. The impact of demographic history on wild

populations is widely recognised, primarily through its effects on effective population sizes, gene flow

and range limits[26,28,29]. However, its overall impact on tree species' genetic diversity remains unclear

and requires broader investigation across species with contrasting demographic histories[30].

Demographic history can also affect a population's potential for climate adaptation by influencing the

amount of standing genetic variation available for natural selection and the patterns of adaptive or

maladaptive gene flow[31-33]. Furthermore, genetic load is also impacted by demographic historical

events, through the purging or accumulation of deleterious mutations[24,25]. Although well-established

theoretical predictions on the link between demographic history and genetic load exist[34,35], empirical

studies have reported contrasting results[36-39], highlighting the need for further research, particularly

in forest trees.

In this study, we investigated how historical demographic processes may have shaped key

components of population vulnerability, namely sensitivity and adaptive potential, in six ecologically

and economically important tree species: four conifers (*Taxus baccata* L., *Pinus pinea* L., *Pinus*

pinaster Ait. and *Pinus sylvestris* L.) and two angiosperms (*Fraxinus excelsior* L. and *Fagus sylvatica* L.).

By integrating genomic data from 6,435 trees across 326 populations spanning most of the species'

distributions in Europe and surrounding regions, we estimated genetic diversity, genetic load, and

current climate adaptation, and related these metrics to population-level genetic differentiation, used

here as a proxy for demographic history. Finally, we interpreted these associations in the light of how

historical demographic processes, including genetic drift and population isolation, may have

influenced present-day sensitivity and adaptive potential to ongoing climate change.

Results

Population genetic differentiation reflects the species' demographic history

The six tree species included in this study were chosen to follow a range of typical population sizes

and levels of connectivity, likely resulting from the interplay between life-history traits and

contrasting demographic histories. Population-level genetic differentiation (F_{ST}) reflected well these

patterns (Fig. 1). Populations of *T. baccata* and *P. pinea* consistently showed intermediate to high F_{ST}

(median values of 0.151 and 0.380, respectively; higher than any other species), which may have

arisen from their known reduced effective population size and connectivity, resulting from recurrent

historical bottlenecks and long-term habitat contraction[40-42]. In contrast, *P. pinaster*, displayed

greater variability in F_{ST} across populations, reflecting the presence of continuous and well-connected

populations in parts of its range, as well as smaller and fragmented ones in other parts, likely

resulting from contractions during glacial periods followed by post-glacial range expansions and

reduced gene flow across gene pools[30,43]. Finally, *F. excelsior*, *F. sylvatica* and *P. sylvestris* populations

displayed very low F_{ST}, with only a few populations showing higher genetic differentiation (e.g., F_{ST} >

0.2 only in two *F. excelsior* and one *P. sylvestris* populations). Most populations of these species are

known to be well-connected due to a combination of factors such as multiple glacial refugia,

extensive post-glacial expansion and admixture, notable cold tolerance and efficient long-distance

gene flow[19,30,43-51]. Nevertheless, smaller, fragmented populations can sporadically be found in few,

mainly southern, parts of the range of these species due to environmental constraints or

geographical barriers (*F. excelsior*[48]; *F. sylvatica*[52]; reviewed in Pyhäjärvi *et al.*[19] for *P. sylvestris*).

In addition, the geographical distribution of population-level F_{ST} varied among species. Highly

differentiated populations of *T. baccata* and *P. pinea* showed no clear spatial pattern (Fig. 1A, B),

with high F_{ST} even in populations located within the species' core range. In contrast, the other species

exhibited higher F_{ST} towards the range edges compared to their core populations (Fig. 1C-F; see also

Theraroz et al.[39] and Guadaño-Peyrot et al.[53]).

Key intraspecific components of climate change vulnerability vary across populations and

species

Genetic diversity (Hs) varied considerably within species, with distinct patterns emerging across

species (Fig. 2A; but notice that direct comparisons of genetic diversity among species are not

appropriate due to the use of different genotyping platforms; see Methods). At the within-species

level, *P. pinea* exhibited the greatest among-population variability in Hs, as evidenced by its relatively

flat cumulative curve compared with the other species. The other two pine species also had higher

among-population variability in Hs than *T. baccata* and the two broadleaves, for which the cumulative

curves were steeper. Hs changed significantly with longitude in most species, increasing eastward in

T. baccata, *F. sylvatica* and *P. sylvestris*, decreasing in *P. pinea* and *P. pinaster* and showing no

significant trend in *F. excelsior* (Fig. S2A; Table S1). Additionally, Hs increased significantly with

latitude in *P. pinaster* and *P. sylvestris*, with southern range-edge populations of both species

showing lower genetic diversity (Fig. S2B). These patterns likely reflect known glacial and post-glacial

forest dynamics as reconstructed from genetic data and pollen records (e.g.,[39,42,44,54-58]).

We also observed marked differences in realised genetic load (i.e., estimated by the number of

deleterious mutations in the homozygous state; hereafter referred to as 'genetic load'; see Methods)

among species (Fig. 2B). Interestingly, *T. baccata* and *P. pinea* exhibited the highest genetic load

values, whereas *P. pinaster* and *P. sylvestris* had comparable lower medians. Furthermore, among-

population variability in genetic load was greatly reduced in *P. sylvestris* compared to the other

species (e.g., 0.101 to 0.111 in *P. sylvestris* compared to 0.074 to 0.140 in *P. pinaster*).

Using the best genotype-environment association (GEA) models, we selected candidate single

nucleotide polymorphisms (SNPs) to estimate climate adaptation via the genomic discrepancy index

(GDI; corresponding to the deviation of a population from the expected GEA models; see Methods),

retaining between 100 (*T. baccata*) and 2,957 (*F. excelsior)* SNPs depending on the species (Figs. 2C,

S3; Table S2). The distribution of GDI (Fig. 2D) showed considerable variability within and among

species. The distributions for *F. sylvatica* and *P. pinaster* were nearly Gaussian, whereas *F. excelsior*

and P. sylvestris were still roughly Gaussian but exhibited long tails with higher GDI values. In

contrast, *T. baccata* and *P. pinea* displayed very long tails of higher GDI values, approaching a

bimodal distribution, with many populations exhibiting elevated GDI. Populations of *T. baccata* with

high GDI are primarily found in southern regions, including Greece, Bosnia-Herzegovina, Spain and

Italy[17].

Population genetic differentiation as a major factor explaining key components of climate

change vulnerability

Genetic diversity (*H*s) decreased with increasing population-level genetic differentiation (F_{ST}) in all

species (Table 1), a pattern that was further confirmed by using unbiased Jost's D (Table S3; see

Methods). Associations remained highly significant and consistent across species when atypical

population-level F_{ST} values were removed and when geography was considered in the models (except

for *F. excelsior*; Table 1; Fig. S4; see Methods).

All conifer species, except *P. pinea*, showed a lower genetic load (i.e., a reduction of deleterious

mutations in homozygous form) in populations with higher F_{ST}. In contrast, the increase in genetic

load with F_{ST} observed in *P. pinea* is likely linked to its unique demographic history, characterised by a

relatively recent population collapse[42]. These relationships were consistent whether populations with

atypically high population-level F_{ST} values were included or not in the analysis (Figs. 3A, S5A). Genetic

load was not computed for the two broadleaves (*F. excelsior* and *F. sylvatica*) as its impact is

expected to be lower in angiosperm species (see Introduction).

Climate adaptation, assessed via GDI, also varied among species in ways that reflected the

population size and connectivity gradient illustrated in Fig. 1G, with higher genetic differentiation

having from negative to negligible effects on climate adaptation along this gradient (Table 1; Fig. 3B,

S5B). In *T. baccata* and *P. pinea*, GDI was higher in populations exhibiting greater F_{ST}. A similar

pattern was observed in *P. pinaster*, *F. excelsior* and *F. sylvatica*, although for the latter two species it

was only evident when populations with atypically high population-level F_{ST} were included, likely

reflecting higher thresholds of isolation required to affect climate adaptation. Finally, no significant

association was detected between population-level F_{ST} and GDI in the widespread, weakly

differentiated *P. sylvestris*.

Discussion

Our study suggests that, beyond macro-glacial dynamics, local historical events and demography, as

reflected in present-day genetic differentiation, explain a substantial portion of the variability in

genetic diversity among populations. We consistently observed a reduction in genetic diversity with

increasing population-level F_{ST}. This result remained robust after accounting for geographic effects

(Table 1), except for *F. excelsior*. Since geography has previously been linked to glacial-postglacial

dynamics[59], these findings support the additional role of local genetic drift and limited historical gene

flow in shaping the observed patterns. Overall, our work suggests that even species capable of long-

distance dispersal and therefore substantial gene flow may still experience long-term isolation. This

isolation likely led to reduced genetic diversity through a combination of ecological or environmental

specialisation, genetic drift, and inbreeding[30,60,61]. Since rapid adaptation mostly relies on standing

genetic variation[62], the general reduction in genetic diversity in highly differentiated populations

could limit their adaptive potential, making them particularly vulnerable to climate change[12,39].

Genetic load varied among species, reflecting differences in their life-history traits and demographic

history. A reduction in deleterious mutations with increasing genetic differentiation was observed

across all species except *P. pinea*. Historical isolation is expected to intensify genetic drift, reducing

effective population size and increasing inbreeding. This can increase the fixation of deleterious

alleles, thereby strengthening purifying selection and ultimately reducing realised genetic load[63,65].

Similar patterns have been reported in other long-lived organisms, where increased genetic drift and

isolation have led to the purging of deleterious mutations, likely increasing their adaptive

potential[37,66]. The unique pattern observed in *P. pinea* (positive association with genetic

differentiation) can also be explained by demographic history. Present-day gene pools of this species

likely persisted through near-extinction in a small number of Iberian populations with fossil records

dating back to the last Neanderthals (c. 49 kyr BP)[42,67]. Under such prolonged and extreme isolation,

reduced effective population size and limited gene flow likely promoted the accumulation, rather

than the purging, of deleterious mutations[65,68].

Across species, genetic load tracked gradients of population size and connectivity (Fig. 1G), with

higher values observed in *T. baccata* and *P. pinea* than in *P. pinaster* and *P. sylvestris*. This pattern

suggests that species with smaller and more fragmented populations tend to accumulate more

deleterious mutations, regardless of within-species dynamics. Consistent with this, demographic

history and the resulting patterns of population differentiation account for much of the variation in

genetic load across conifer populations. In particular, the accumulation of deleterious mutations

appears to be shaped by historical effective population size and connectivity, underscoring the

importance of considering these factors when assessing the consequences of genetic load for

population persistence under climate change.

Increasing genetic differentiation was associated with suboptimal climate adaptation in all species

except *P. sylvestris*. Alongside the parallel reduction in genetic diversity with increasing

differentiation, these results highlight the potential beneficial role of gene flow in enhancing

adaptation by maintaining standing genetic variation, a key driver of adaptive evolution[33,69].

Furthermore, both theoretical predictions and empirical evidence also showed that, by introducing

pre-adapted alleles, gene flow can further facilitate climate tracking, particularly in leading-edge

populations[32,70,71]. We observed marked interspecific differences in climate adaptation in our study.

Species with a prevalence of small and fragmented populations (e.g., *T. baccata* and *P. pinea*)

showed stronger negative associations between population genetic differentiation and climate

adaptation than species with larger, more continuous populations (e.g., *F. sylvatica* and *P. sylvestris*).

These differences may, at least in part, reflect variation in historical effective population size, as the

benefits of increased standing genetic diversity and enhanced climate tracking via gene flow are

likely to be more pronounced in smaller populations[32,71,72].

Interestingly, across the six tree species, populations with low genetic differentiation consistently

showed no signs of reduced climate adaptation. These results provide further empirical support for

the positive role of genetic connectivity in enhancing climate adaptation in small, fragmented

populations, while showing no detectable negative effects in larger, more continuous populations.

However, while biologically meaningful, these insights need to be taken with caution. For instance,

climate adaptation models used here rely on several assumptions, including that genotype-

environment associations accurately capture adaptive signals[73]. Furthermore, higher Euclidean

distances between observed and predicted genomic compositions (GDI) were interpreted here as

signatures of suboptimal climate adaptation, therefore neglecting other potential explanations, such

as alternative adaptive pathways or pre-adaptation to future climate.

Conclusion

Genomic analyses of 6,435 trees from 326 populations across six ecologically and economically

important forest tree species revealed major effects of demographic history, as summarised by

population-level genetic differentiation, on shaping population vulnerability to climate change.

Population adaptive potential, reflected in within-population genetic diversity, declined consistently

with historical isolation. At the same time, sensitivity to climate change was likely shaped by reduced

effective population size and limited historical gene flow, leading either to the purging or the

accumulation of deleterious mutations, and increasing maladaptation to contemporary climates.

Overall, this work provides valuable insights into how genetic differentiation, reflecting genetic drift

and limited historical gene flow, influences current vulnerability of forest tree populations to climate

change. Future empirical work could aim to link genetic metrics to effective population size,

phenotypic performance, or reproductive fitness, to better assess the contribution of each factor to

climate vulnerability. Our findings can also inform management strategies, including assisted gene

flow, by helping to identify populations most at risk from climate change and those likely to benefit

from increased genetic variance in fitness-related traits[74]. Assessing current gene flow dynamics will

be crucial to determine whether rapid environmental change is already reshaping historical patterns

of connectivity and genetic differentiation.

Materials and methods

Sampling and genotyping

Our study focused on six European tree species, namely *Taxus baccata*, *Pinus pinea*, *Pinus pinaster*,

Fraxinus excelsior, Fagus sylvatica and *Pinus sylvestris.* Species were selected to represent

contrasting demographic histories, including different phylogenetic groups, stand formation and

population sizes, as well as levels of connectivity and population genetic structure[30,43] (Fig. 1G; see

Results). For each species, 29-86 populations spanning most of their ranges were sampled, with total

sample sizes per species ranging from 490 to 1,729 mature trees (except *for F. excelsior* where young

trees were sampled; see Supplementary Files 1 and 2). Particular attention was paid to sample

populations with different degrees of genetic differentiation, including both core and range-edge

populations.

Genomic data for 490 trees representing 29 populations of *T. baccata* were obtained from a previous

study[17]. This dataset comprised single nucleotide polymorphisms (SNPs) generated by combining

gene-capture and single-primer enrichment technology (SPET) data.

Genomic data for 461 trees from 31 natural populations of *F. excelsior* were compiled from one

previous data paper and one study: 286 trees sequenced using SPET[75] and 175 using low-coverage

whole genome sequencing[76], following Doonan *et al.*[77]. SNPs were called against the same reference

genome, FRAX_0001_PL[78].

Genomic data for 1,224 trees from 51 populations of *F. sylvatica* were generated by combining two

previous datasets: 370 trees from a data paper[75] and 854 trees from a published dataset[79], both of

which produced using SPET. SNP quality filtering and the removal of outliers and duplicates were

performed on the merged dataset following the procedure described in Pinosio *et al.*[75].

Genomic data for the three pine species (*P. pinea*: 952 trees from 48 populations; *P. pinaster*: 1,579

trees from 86 populations; *P. sylvestris*: 1,729 trees from 81 populations) were retrieved: from a

previous study[80] for *P. pinaster* and from a data paper[81] for *P. pinea* and *P. sylvestris*. SNP genotypes

were generated using the multispecies 4TREE (*P. pinea*, *P. pinaster*) and the PiSy50k (*P. sylvestris*)

Axiom arrays (Thermo Fisher Scientific, USA) at IGA Technology Services (Udine, Italy). Following

Olsson *et al.*[81], we applied the quality thresholds for markers CR ≥ 85, FLD ≥ 3.2, HetSO ≥ -0.3 and

samples DQC ≥ 0.4 with QC call rate ≥ 85, in order to maximise SNP retention while ensuring high

data quality.

Overall, genomic datasets included 6,435 trees from 326 populations, mainly in Europe and

bordering regions in North Africa and Asia (see details in Supplementary Files 1 and 2). For each

species, we performed population genetic structure analyses to ensure that merged datasets were

biologically consistent and appropriate for downstream analyses (Figs. S6, S7). Two of the 326

populations (COM and FUEbis of *P. pinaster*) were excluded because they were duplicated or had too

few samples. For each species, two datasets were then generated after removing monomorphic SNPs

and variants that failed additional filtering thresholds (coverage depth < 7 or > 150, and allele

balance at heterozygous sites < 0.2 or > 0.8). The first dataset (*dataset 1*) retained SNPs and

individuals with less than 15% missing data and SNPs with a minor allele count of at least three

(except for *T. baccata*, see Francisco *et al.*[17]). This dataset was used for population genetic structure

(PCA, STRUCTURE) and differentiation (population-level F_{ST}) analyses, as well as for calculating

genetic diversity (Hs), as these analyses are sensitive to high rates of missing data and low-frequency

SNPs[82]. *Dataset 1* comprised between 3,563 (*P. pinea*) and 34,760 SNPs (*P. sylvestris*), depending on

the species, with a total of 6,262 trees genotyped across the 324 populations (Table S4). The second

dataset (*dataset 2*), used for computing climate adaptation and genetic load statistics, was filtered

out for missing data > 30% (both for SNPs and individuals) and for minor allele counts lower than

twice the number of individuals in the smallest population before filtering (Table S4). This latter filter

was applied to remove low-frequency alleles that could otherwise increase false-positive rates, while

retaining potential climate-associated SNPs or deleterious mutations that are fixed in small

populations. *Dataset 2* included between 3,790 (*P. pinea*) and 34,848 SNPs (*P. sylvestris*), depending

on the species, with a total of 6,326 trees genotyped across the 324 populations (Table S4). From

these datasets, SNPs from coding regions that could be polarised and annotated using the *Pinus*

tabuliformis or *Taxus chinensis* reference genomes[83,84] were used to estimate genetic load (see

below). This yielded 4,700; 1,195; 2,219; and 12,808 SNPs for *T. baccata*, *P. pinea*, *P. pinaster*, and *P.*

sylvestris, respectively. Finally, for climate adaptation analyses, missing data in *dataset 2* were

imputed using the most common genotype within each gene pool.

Climatic data

Climatic data were extracted at high spatial resolution (30 arc-seconds) from Climate Downscaling

Tool (ClimateDT[85]). The 1901-1950 period was defined as the reference climate to which the

populations are currently adapted, except for *F. excelsior*, for which the 1991-2020 period was used

instead. This more recent time interval may better reflect the climate to which populations are

adapted in this species, given the relatively young age of the sampled trees and the species' known

capacity for rapid adaptation[86]. For each species, the retained climatic variables (Table S5) were

selected following the approach described in Francisco *et al.*[17]: (i) a pre-selection step based on

species life-history traits and the available literature, (ii) forward selection analysis based on genomic

importance, and (iii) removal of correlated variables (Pearson's $|r| > 0.75$) to ensure that

multicollinearity remained low and the variance inflation factor below 10 across variables (see

Supplementary Material).

Population genetic structure and differentiation

For each species, principal component analysis (PCA) was used to assess population genetic

structure, as implemented in the VEGAN R package[87] (Fig. S6). Genetic clusters were also examined

based on previously[17,88] or newly run Bayesian clustering analyses, as implemented in STRUCTURE

v2.3.4[89] (Fig. S7). For newly run STRUCTURE analyses, ten independent MCMC runs were performed

for each number of potential clusters (K = 1-10), with 100,000 iterations per run and a burn-in of

10,000 iterations. The most probable number of genetic clusters was determined using the ΔK

method[90].

Population-level genetic differentiation (F_{ST}), used here as a proxy for population-specific

demographic history, encompassing historical population size fluctuation, gene flow and connectivity

(e.g.,[91,92]), was calculated using BayeScan v2.1[93]. In this approach, F_{ST} represents the genetic

differentiation of each population relative to a predicted ancestral gene pool, making it less

influenced by local outliers and more reflective of broad historical patterns of divergence. Given the

comprehensive sampling effort, including both core and range-edge populations for each species,

populations with high F_{ST} can be considered more historically isolated than the species average.

Genetic metrics of vulnerability to climate change

Genetic differences among individuals of the same species can produce diverse phenotypes within

populations, some of which may confer advantages in shifting climate. The extent of this genetic

diversity can determine the population's evolutionary potential[12]. Genetic variation within

populations was quantified using Nei's genetic diversity (Hs), as implemented in the HIERFSTAT R

package[94]. Due to the differences in experimental design to obtain genomic data (e.g., gene capture,

SPET designs and different genotyping arrays), genetic diversity values cannot be compared across

species.

Genetic load was estimated only for the conifer species in this study (*T. baccata*, *P. pinea*, *P. pinaster*,

and *P. sylvestris*), as the accumulation of deleterious mutations in these species has been shown to

be greater than in broadleaves and linked to fitness-related traits (e.g.,[95]; see Introduction). First,

SNPs were mapped onto the *Pinus tabuliformis* reference genome[83] (*P. pinea*, *P. pinaster* and *P.*

sylvestris; see Supplementary Material) or onto the *Taxus chinensis* reference genome[84] (*T. baccata*).

SnpEff software (version 5.1) was then used to identify SNPs causing amino acid changes in protein-

coding sequences[96]. Finally, the functional impact of mutations was predicted using PROVEAN

(version 1.1.5), a software that assesses the impact of mutations by evaluating sequence

conservation and alignment scores across homologous proteins[97]. For each SNP, PROVEAN predicted

its impact on the biological function of the corresponding protein and assigned a 'deleteriousness'

score. Mutations were considered deleterious when the PROVEAN score was lower than -2.5[97].

Realised genetic load was then computed following González-Martínez *et al.*[27], as the ratio of the

expected number of deleterious mutations in homozygosis to the number of expected neutral

mutations. As most deleterious alleles are thought to be recessive, the realised genetic load is

therefore expected to have a stronger immediate impact on fitness than the absolute number of

deleterious mutations[27,65].

Population climate adaptation was quantified using a two-step procedure. First, we investigated the

presence of genomic markers potentially involved in climate adaptation by carrying out genotype-

environment association (GEA) analyses. For each species, six GEA methods (correcting or not for

population genetic structure) were applied to identify candidate loci: redundancy analysis (RDA),

partial redundancy analysis (pRDA), latent factor mixed models (LFMM), BayPass, gradient forest (GF)

uncorrected for population genetic structure (GF-raw) and GF corrected for population genetic

structure (GF-corrected; see Table S6 and Supplementary Material for details). We then defined an

all-outlier SNP set comprising all candidate SNPs identified across the GEA analyses. Following

Archambeau *et al.*[98], an *outlier* SNP set was further defined, including only unlinked candidate SNPs

identified by at least two GEA methods. For this, we used a linkage disequilibrium threshold of R^2 <

|0.7| within contigs or 2,000 bp, depending on the available information for each species, and

filtered the datasets for linked loci using the GENETICS R package[99]. For each species, a *random* SNP set

was also generated by randomly selecting SNPs not identified as candidates by any GEA method,

matching both the number and allele frequency distribution of the *outlier* set (see Supplementary

Material).

Second, climate adaptation was quantified for each population within species by computing the

genomic discrepancy index (GDI[17]) using the RDA method. GDI computes the Euclidean distances

between the observed and predicted population scores along the retained RDA axes. Higher

population GDI indicates a greater deviation from the established GEA relationship relative to other

populations. For most species, the *all-outlier* SNP set was used to build the reference GEA model for

GDI estimates. However, for *T. baccata* and *P. pinaster*, we applied SNP sets (*random* and *outlier*,

respectively) previously identified as most suitable through model evaluation with fitness proxies[17,98].

For all species, GDI estimates based on alternative SNP sets were highly correlated with those

obtained from the selected sets (Pearson's correlation > 0.64; Table S2).

Linking demographic history to vulnerability to climate change

To assess the effects of demographic history on population vulnerability to climate change, we

examined, for each species, the relationship between population-level F_{ST} and genetic diversity,

genetic load and GDI (see Supplementary File 3). Associations were evaluated using linear models

with F-test p-values, as implemented in the STATS R package[100]. To avoid interpreting associations

driven solely by extreme values, linear models were investigated with and without populations

exhibiting atypically high F_{ST}, defined as those falling outside 2.4 times the interquartile range of

population-level F_{ST} within each species.

Author Contributions

Designed research: TF, AP, SCG-M; acquired and pretreated genomic data: SO, MK, MW, SP, GGV, KBB, LRN, JMD, DG, EV; analysed data: TF, IL-K, SCG-M; wrote the paper: TF, SCG-M; obtained funding: MW, GGV, DG, AP, SCG-M, LRN; all authors participated in discussions and revised multiple versions of the manuscript, including the final version.

Acknowledgments

TF was funded by a PhD grant of the INRAE Scientific Department ECODIV. EV was supported by the MSCA European fellowship MedForAct (GA 101107604). MK was supported by the Slovenian Research Agency (research core funding no. P4-0107 and Young Researcher grant no. 58170). This research was funded by EU H2020 FORGENIUS (grant agreement No. 862221) and Horizon Europe OptFORESTS (grant agreement No. 101081774) projects. Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union. Neither the European Union nor the granting authority can be held responsible for them. Additional *F. excelsior* data was generated by the AshAdapt project supported by the Independent Research Fund Denmark (DFF Technology and Production Sciences) under grant no. 8022-00355B. We are grateful to the Genotoul bioinformatics platform Toulouse Occitanie (Bioinfo Genotoul, https://bioinfo.genotoul.fr/) for providing computing and storage resources.

Data Archiving Statement

Genomic data for *Taxus baccata* from Francisco *et al*.[17] is available at Data INRAE (doi: https://doi.org/10.57745/JJ0ZEI); for *Pinus pinaster* from Olsson *et al*.[80] at Zenodo (doi: 10.5281/zenodo.14950394); for *Pinus pinea* and *Pinus sylvestris* from Olsson *et al*.[81] at Zenodo (doi: 10.5281/zenodo.17455887 and 10.5281/zenodo.17462035, respectively). Newly combined genomic

438   data for *Fraxinus excelsior* and *Fagus sylvatica* are openly available at Data INRAE (doi:

439   10.57745/OG2FBY).

441   *Code availability*

442   Code used for data processing and analysis will be made publicly available upon publication at a

443   GitHub public repository.

 Table and Figures

Table 1. **Association between population genetic differentiation (F_{ST}) and genetic diversity (Hs).**

Linear model coefficient signs (β), explained variance (R^2), and F-test p-values, summarising the

direction and strength of the relationship between population-level F_{ST}, used here as a proxy for

demographic history, and genetic diversity (Hs). Analyses were run including and excluding

populations exhibiting atypically high F_{ST} values and accounting or not for geographical variation in Hs

(longitude 'Lon' and latitude 'Lat').

Species	Excluding atypical populations		Including atypical populations	
	Hs ~ F_{ST}	Hs ~ F_{ST} | Lon + Lat	Hs ~ F_{ST}	Hs ~ F_{ST} | Lon + Lat
T. baccata	(-) 0.59***	(-) 0.51***	(-) 0.59***	(-) 0.51***
P. pinea	(-) 0.87***	(-) 0.70***	(-) 0.87***	(-) 0.70***
P. pinaster	(-) 0.46***	(-) 0.24***	(-) 0.52***	(-) 0.28***
F. excelsior	(-) 0.13^m	(-) 0.12^m	(-) 0.43***	(-) 0.33***
F. sylvatica	(-) 0.56***	(-) 0.50***	(-) 0.88***	(-) 0.71***
P. sylvestris	(-) 0.33***	(-) 0.31***	(-) 0.48***	(-) 0.40***

453 ^{m}: $0.10 < p < 0.05$; ***: $p < 0.001$

Figure 1. **Description of the six tree species**. **A-F** Geographical range and population-level genetic

differentiation (F_{ST}). Each circle represents a population, with colour intensity indicating the

magnitude of F_{ST}. Inserts show the density distribution of population-level F_{ST} values, represented on

a common scale to facilitate visual comparison across species. Shaded areas depict the approximate

species range[101] for the geographic extent of latitude 30°N-60°N and longitude 10°W-37°E. Sampled

populations of *P. sylvestris* occurring outside this range are not shown but can be found in Fig. S1.

Notice that species such as *T. baccata*, despite having a large geographical range, have a scattered

distribution while others (e.g., *P. sylvestris*) constitute large, stand-forming populations. **G** Conceptual

illustration of the gradient in population size and connectivity represented by the six species under

study[30,40-43,46,50], which spans from species with predominantly small and isolated populations to

species with large and well-connected populations. The schematic is inspired by[43].

Figure 2. **Genetic metrics of population vulnerability to climate change across the studied species**.

A Cumulative distribution of scaled genetic diversity (*H*s; calculated by subtracting the mean and

dividing by the standard deviation) across populations. A stepped curve indicates low variability in *H*s

across populations (e.g., *F. sylvatica*), whereas a flatter curve indicates high variability (e.g., *P. pinea*).

B Comparison of realised genetic load across the four conifers species, with Tukey post-hoc groups

indicated by lowercase letters. **C** Manhattan plots showing the retained candidate SNPs (coloured)

used for GDI calculation, plotted against the total number of SNP, which varied across species (Table

S2). *P*-values were obtained from the RDA GEA models. **D** GDI density plot across populations for

each species. Black line represents the median GDI values. Note that values are not directly

comparable among species.

Figure 3. **Population-level genetic differentiation (F_{ST}) as a key factor explaining vulnerability to climate change.** Linear relationships between (**A**) realised genetic load (only conifers) or (**B**) GDI, and population-level F_{ST}, after excluding populations with atypically high F_{ST} values (see Methods and Fig. S5 for relationships including atypical populations). The coefficient of determination (R^2) and F-test p-values are shown for each model.

References

1. Brockerhoff, E. G. *et al.* Forest biodiversity, ecosystem functioning and the provision of ecosystem services. *Biodivers. Conserv.* **26**, 3005-3035 (2017).

2. Fréjaville, T., Vizcaíno-Palomar, N., Fady, B., Kremer, A. & Benito Garzón, M. Range margin populations show high climate adaptation lags in European trees. *Glob. Change Biol.* **26**, 484-495 (2020).

3. Petit-Cailleux, C. *et al.* Tree mortality risks under climate change in Europe: Assessment of silviculture practices and genetic conservation networks. *Front. Ecol. Evol.* **9**, 706414 (2021).

4. Martinez del Castillo, E. *et al.* Climate-change-driven growth decline of European beech forests. *Commun. Biol.* **5**, 163 (2022).

5. Forzieri, G. *et al.* Emergent vulnerability to climate-driven disturbances in European forests. *Nat. Commun.* **12**, 1081 (2021).

6. Taccoen, A., Piedallu, C., Seynave, I., Gégout-Petit, A. & Gégout, J.-C. Climate change-induced background tree mortality is exacerbated towards the warm limits of the species ranges. *Ann. For. Sci.* **79**, 23 (2022).

7. *Climate Change 2007: The Physical Science Basis*. (Cambridge university press, Cambridge, 2007).

8. Foden, W. B. *et al.* Climate change vulnerability assessment of species. *WIREs Clim. Change* **10**, e551 (2019).

9. Fitzpatrick, M. C. & Keller, S. R. Ecological genomics meets community-level modelling of biodiversity: Mapping the genomic landscape of current and future environmental adaptation. *Ecol. Lett.* **18**, 1-16 (2015).

10. Barratt, C. D. *et al.* Life on the edge: A new toolbox for population-level climate change vulnerability assessments. *Methods Ecol. Evol.* **15**, 2038-2058 (2024).

11. Lazic, D. *et al.* Genomic variation of European beech reveals signals of local adaptation despite high levels of phenotypic plasticity. *Nat. Commun.* **15**, 8553 (2024).

12. Ørsted, M., Hoffmann, A. A., Sverrisdóttir, E., Nielsen, K. L. & Kristensen, T. N. Genomic variation predicts adaptive evolutionary responses better than population bottleneck history. *PLoS Genet.* **15**, e1008205 (2019).

13. Leimu, R. & Fischer, M. A meta-analysis of local adaptation in plants. *PLoS ONE* **3**, e4010 (2008).

14. Marcora, P. I. *et al.* Are populations of *Polylepis australis* locally adapted along their elevation gradient? *Neotrop. Biodivers.* **7**, 246-256 (2021).

15. Browne, L., Wright, J. W., Fitz-Gibbon, S., Gugger, P. F. & Sork, V. L. Adaptational lag to temperature in valley oak (*Quercus lobata*) can be mitigated by genome-informed assisted gene flow. *Proc. Nat. Acad. Sci. USA* **116**, 25179-25185 (2019).

16. De Kort, H. *et al.* Pre-adaptation to climate change through topography-driven phenotypic plasticity. *J. Ecol.* **108**, 1465-1474 (2020).

17. Francisco, T. *et al.* Genomic signatures of climate-driven (mal)adaptation in an iconic conifer, the english yew (*Taxus baccata* L.). *Evol. Appl.* **18**, e70160 (2025).

18. Kardos, M., Taylor, H. R., Ellegren, H., Luikart, G. & Allendorf, F. W. Genomics advances the study of inbreeding depression in the wild. *Evol. Appl.* **9**, 1205-1218 (2016).

19. Pyhäjärvi, T., Kujala, S. T. & Savolainen, O. 275 years of forestry meets genomics in *Pinus sylvestris*. *Evol. Appl.* **13**, 11-30 (2020).

20. Cheptou, P.-O. The evolutionary ecology of inbreeding depression in wild plant populations and its impact on plant mating systems. *Front Plant Sci.* **15**, 1359037 (2024).

21. Lande, R., Schemske, D. W. & Schultz, S. T. High inbreeding depression, selective interference among loci, and the threshold selfing rate for purging recessive lethal mutations. *Evolution* **48**, 965-978 (1994).

22. Williams, C. G. & Savolainen, O. Inbreeding depression in conifers: Implications for breeding strategy. *For. Sci.* **42**, 102-117 (1996).

23. Olson-Manning, C. F., Wagner, M. R. & Mitchell-Olds, T. Adaptive evolution: Evaluating empirical support for theoretical predictions. *Nat. Rev. Genet.* **13**, 867-877 (2012).

24. Peischl, S., Dupanloup, I., Kirkpatrick, M. & Excoffier, L. On the accumulation of deleterious mutations during range expansions. *Mol. Ecol.* **22**, 5972-5982 (2013).

25. Peischl, S. & Excoffier, L. Expansion load: Recessive mutations and the role of standing genetic variation. *Mol. Ecol.* **24**, 2084-2094 (2015).

26. Ellegren, H. & Galtier, N. Determinants of genetic diversity. *Nat. Rev. Genet.* **17**, 422-433 (2016).

27. González-Martínez, S. C., Ridout, K. & Pannell, J. R. Range expansion compromises adaptive evolution in an outcrossing plant. *Curr. Biol.* **27**, 2544-2551.e4 (2017).

28. Hewitt, G. The genetic legacy of the Quaternary ice ages. *Nature* **405**, 907-913 (2000).

29. Star, B. & Spencer, H. G. Effects of genetic drift and gene flow on the selective maintenance of genetic variation. *Genetics* **194**, 235-244 (2013).

30. Milesi, P. *et al.* Resilience of genetic diversity in forest trees over the Quaternary. *Nat. Commun.* **15**, 8538 (2024).

31. Atkins, K. E. & Travis, J. M. J. Local adaptation and the evolution of species' ranges under climate change. *J. of Theor. Biol.* **266**, 449-457 (2010).

32. Kremer, A. *et al.* Long-distance gene flow and adaptation of forest trees to rapid climate change. *Ecol. Lett.* **15**, 378-392 (2012).

33. Tigano, A. & Friesen, V. L. Genomics of local adaptation with gene flow. *Mol. Ecol.* **25**, 2144-2164 (2016).

34. Bataillon, T. & Kirkpatrick, M. Inbreeding depression due to mildly deleterious mutations in finite populations: Size does matter. *Genet. Res.* **75**, 75-81 (2000).

35. Glémin, S. How are deleterious mutations purged? Drift versus nonrandom mating. *Evolution* **57**, 2678-2687 (2003).

36. Rogers, R. L. & Slatkin, M. Excess of genomic defects in a woolly mammoth on Wrangel island. *PLoS Genet.* **13**, e1006601 (2017).

37. Grossen, C., Guillaume, F., Keller, L. F. & Croll, D. Purging of highly deleterious mutations through severe bottlenecks in Alpine ibex. *Nat. Commun.* **11**, 1001 (2020).

38. Hasselgren, M. *et al.* Genomic and fitness consequences of inbreeding in an endangered carnivore. *Mol. Ecol.* **30**, 2790-2799 (2021).

39. Theraroz, A. *et al.* The genetic consequences of population marginality: A case study in maritime pine. *Divers. Distrib.* **30**, e13910 (2024).

40. Dubreuil, M. *et al.* Genetic effects of chronic habitat fragmentation revisited: Strong genetic structure in a temperate tree, *Taxus baccata* (Taxaceae), with great dispersal capability. *Am. J. Bot.* **97**, 303-310 (2010).

41. Burgarella, C. *et al.* Recent population decline and selection shape diversity of taxol-related genes. *Mol. Ecol.* **21**, 3006-3021 (2012).

42. Jaramillo-Correa, J. P. *et al.* Evolutionary rate and genetic load in an emblematic Mediterranean tree following an ancient and prolonged population collapse. *Mol. Ecol.* **29**, 4797-4811 (2020).

43. Alberto, F. J. *et al.* Potential for evolutionary responses to climate change - evidence from tree populations. *Glob. Change Biol.* **19**, 1645-1661 (2013).

44. Magri, D. Patterns of post-glacial spread and the extent of glacial refugia of European beech (*Fagus sylvatica*). *J. Biogeogr.* **35**, 450-463 (2008).

45. Magri, D. *et al.* A new scenario for the Quaternary history of European beech populations: Palaeobotanical evidence and genetic consequences. *New Phytol.* **171**, 199-221 (2006).

46. Heuertz, M. *et al.* Nuclear microsatellites reveal contrasting patterns of genetic structure between western and southeastern European populations of the common ash (*Fraxinus excelsior* L.). *Evolution* **58**, 976-988 (2004).

47. Heuertz, M. *et al.* Chloroplast DNA variation and postglacial recolonization of common ash (*Fraxinus excelsior* L.) in Europe. *Mol. Ecol.* **13**, 3437-3452 (2004).

48. Tollefsrud, M. M. *et al.* Genetic structure in the northern range margins of common ash, *Fraxinus excelsior* L. *PLoS ONE* **11**, e0167104 (2016).

49. Postolache, D. *et al.* Genetic signatures of divergent selection in European beech (*Fagus sylvatica* L.) are associated with the variation in temperature and precipitation across its distribution range. *Mol. Ecol.* **30**, 5029-5047 (2021).

50. Tyrmi, J. S. *et al.* Genomics of clinal local adaptation in *Pinus sylvestris* under continuous environmental and spatial genetic setting. *G3 (Bethesda)* **10**, 2683-2696 (2020).

51. Hurme, P., Repo, T., Savolainen, O. & Pääkkönen, T. Climatic adaptation of bud set and frost hardiness in Scots pine (*Pinus sylvestris*). *Can. J. For. Res.* **27**, 716-723 (1997).

52. Sánchez de Dios, R. *et al. Fagus sylvatica* L. Peripheral populations in the Mediterranean Iberian Peninsula: Climatic or anthropic relicts? *Ecosystems* **24**, 211-226 (2021).

53. Guadaño-Peyrot, C. *et al.* Genetic signatures of population marginality in four European conifers. In preparation.

54. Gómez, A., Vendramin, G. G., González-Martínez, S. C. & Alía, R. Genetic diversity and differentiation of two Mediterranean pines (*Pinus halepensis* Mill. and *Pinus pinaster* Ait.) along a latitudinal cline using chloroplast microsatellite markers. *Divers. Distrib.* **11**, 257-263 (2005).

55. Cheddadi, R. *et al.* Imprints of glacial refugia in the modern genetic diversity of *Pinus sylvestris*. *Glob. Ecol. Biogeogr.* **15**, 271-282 (2006).

56. Naydenov, K., Senneville, S., Beaulieu, J., Tremblay, F. & Bousquet, J. Glacial vicariance in Eurasia: Mitochondrial DNA evidence from Scots pine for a complex heritage involving genetically distinct refugia at mid-northern latitudes and in Asia Minor. *BMC Evol. Biol.* **7**, 233 (2007).

57. Mayol, M. *et al.* Adapting through glacial cycles: Insights from a long-lived tree (*Taxus baccata*). *New Phytol.* **208**, 973-986 (2015).

58. Wachowiak, W., Perry, A., Zaborowska, J., González-Martínez, S. C. & Cavers, S. Admixture and selection patterns across the European distribution of Scots pine, *Pinus sylvestris* (Pinaceae). *Bot. J. Linn. Soc.* **200**, 416-432 (2022).

59. Petit, R. J. *et al.* Glacial refugia: Hotspots but not melting pots of genetic diversity. *Science* **300**, 1563-1565 (2003).

60. Temunović, M. *et al.* Environmental heterogeneity explains the genetic structure of continental and mediterranean populations of *Fraxinus angustifolia* Vahl. *PLoS ONE* **7**, e42764 (2012).

61. Potter, K. M., Campbell, A. R., Josserand, S. A., Nelson, C. D. & Jetton, R. M. Population isolation results in unexpectedly high differentiation in Carolina hemlock (*Tsuga caroliniana*), an imperiled southern Appalachian endemic conifer. *Tree Genet. Genomes* **13**, 105 (2017).

62. Barrett, R. D. H. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol. Evol.* **23**, 38-44 (2008).

63. Hedrick, P. W. & Garcia-Dorado, A. Understanding inbreeding depression, purging, and genetic rescue. *Trends Ecol. Evol.* **31**, 940-952 (2016).

64. García-Dorado, A. Understanding and predicting the fitness decline of shrunk populations: Inbreeding, purging, mutation, and standard selection. *Genetics* **190**, 1461-1476 (2012).

65. Dussex, N., Morales, H. E., Grossen, C., Dalén, L. & Oosterhout, C. van. Purging and accumulation of genetic load in conservation. *Trends Ecol. Evol.* **38**, 961-969 (2023).

66. Xue, Y. *et al.* Mountain gorilla genomes reveal the impact of long-term population decline and inbreeding. *Science* **348**, 242-245 (2015).

67. Mutke, S., Vendramin, G. G., Fady, B., Bagnoli, F. & González-Martínez, S. C. Molecular and Quantitative Genetics of Stone Pine (*Pinus pinea*). in *Genetic Diversity* in Horticultural Plants (ed. Nandwani, D.) 61-84 (Springer International Publishing, Cham, 2019). doi:10.1007/978-3-319-96454-6_3.

68. Feng, Y. *et al.* Genome sequences and population genomics provide insights into the demographic history, inbreeding, and mutation load of two 'living fossil' tree species of Dipteronia. *Plant J.* **117**, 177-192 (2024).

69. Chen, J., Glémin, S. & Lascoux, M. Genetic diversity and the efficacy of purifying selection across plant and animal species. *Mol. Biol. Evol.* **34**, 1417-1428 (2017).

70. Sexton, J. P., Strauss, S. Y. & Rice, K. J. Gene flow increases fitness at the warm edge of a species' range. *Proc. Nat. Acad. Sci. USA* **108**, 11704-11709 (2011).

71. Bontrager, M. & Angert, A. L. Gene flow improves fitness at a range edge under climate change. *Evol. Lett.* **3**, 55-68 (2019).

72. Frankham, R. Genetic rescue of small inbred populations: Meta-analysis reveals large and consistent benefits of gene flow. *Mol. Ecol.* **24**, 2610-2618 (2015).

73. Rellstab, C., Dauphin, B. & Exposito-Alonso, M. Prospects and limitations of genomic offset in conservation management. *Evol. Appl.* **14**, 1202-1212 (2021).

74. Aitken, S. N. & Bemmels, J. B. Time to get moving: Assisted gene flow of forest trees. *Evol. Appl.* **9**, 271-290 (2016).

75. Pinosio, S. *et al.* The FORGENIUS Genomic Resources: New Genotyping Tools and Genomic Data for 23 Forest Tree Species and Their Genetic Conservation Units. *Mol. Ecol. Resour.* **26**, e70115 (2026).

76. Budde, K. B., Doonan, J.M., Hanika, T., Verbylaite, R., Ulaszewski, B. , Kjær E.D., Heuertz, M., Nielsen, L.R. Evolutionary consequences of ash dieback in natural common ash populations in Europe. In preparation.

77. Doonan, J. M. *et al.* Multiple, single trait GWAS and supervised machine learning reveal the genetic architecture of *Fraxinus excelsior* tolerance to ash dieback in Europe. *Plant Cell Environ.* **48**, 3793-3809 (2025).

78. Meger, J., Ulaszewski, B., Pałucka, M., Kozioł, C. & Burczyk, J. Genomic prediction of resistance to Hymenoscyphus fraxineus in common ash (*Fraxinus excelsior* L.) populations. *Evol. Appl.* **17**, e13694 (2024).

79. Bajc, M. *et al. Fagus sylvatica* BU2012: A genetic dataset for trial trees from the international beech provenance trial 'Kamenski hrib/Straža (SI) 1996/1998'. https://dirros.openscience.si/IzpisGradiva.php?id=27722&lang=eng (2026) doi:10.20315/Data.0014.

80. Olsson, S. *et al.* Genetic assignment at different geographical levels: A case study in a forest tree species (*Pinus pinaster* Ait.) using SNP markers. *Evol. Appl.* **18**, e70145 (2025).

81. Olsson, S. *et al.* Range-wide SNP reference data and filtering effects in three major European pines. In preparation.

82. Linck, E. & Battey, C. J. Minor allele frequency thresholds strongly affect population structure inference with genomic data sets. *Mol. Ecol. Resour.* **19**, 639-647 (2019).

83. Niu, S. *et al.* The Chinese pine genome and methylome unveil key features of conifer evolution. *Cell* **185**, 204-217.e14 (2022).

84. Xiong, X. *et al.* The *Taxus* genome provides insights into paclitaxel biosynthesis. *Nat. Plants* **7**, 1026-1036 (2021).

85. Marchi, M., Bucci, G., Iovieno, P. & Ray, D. ClimateDT: A global scale-free dynamic downscaling portal for historic and future climate data. *Environments* **11**, 82 (2024).

86. Metheringham, C. L. *et al.* Rapid polygenic adaptation in a wild population of ash trees under a novel fungal epidemic. *Science* **388**, 1422-1425 (2025).

87. Oksanen, J. *et al.* vegan: Community Ecology Package. 2.6-8 https://doi.org/10.32614/CRAN.package.vegan (2025).

88. Olsson, S.*,* Jaramillo-Correa, J. P., Vajana, E., Pinosio, S., Macaya-Sanz, D., Bagnoli, F., Avanzi, C., Piotti, A., Alía, R., Mutke, S., Vendramin, G. G., Grivet, D., González-Martínez, S. C. Thriving with low diversity: demographic history and evolutionary dynamics of stone pine (*P. pinea* L). In preparation.

89. Pritchard, J. K., Stephens, M. & Donnelly, P. Inference of population structure using multilocus genotype data. *Genetics* **155**, 945-959 (2000).

90. Evanno, G., Regnaut, S. & Goudet, J. Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. *Mol. Ecol.* **14**, 2611-2620 (2005).

91. Savolainen, O., Pyhäjärvi, T. & Knürr, T. Gene flow and local adaptation in trees. *Annu. Rev. Ecol., Evol. Syst.* **38**, 595-619 (2007).

92. Kitada, S., Nakamichi, R. & Kishino, H. Understanding population structure in an evolutionary context: Population-specific FST and pairwise FST. *G3: Genes Genomes Genet.* **11**, jkab316 (2021).

93. Foll, M. & Gaggiotti, O. A genome-scan method to identify selected loci appropriate for both dominant and codominant markers: A bayesian perspective. *Genetics* **180**, 977-993 (2008).

94. Nei, M. *Molecular Evolutionary Genetics*. (Columbia University Press, 1987). doi:10.7312/nei-92038.

95. Piñeiro, R. *et al.* A higher load of deleterious mutations has a detrimental effect on tree growth in maritime pine. 2025.11.09.687477 Preprint at https://doi.org/10.1101/2025.11.09.687477 (2025).

96. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff. *Fly* **6**, 80-92 (2012).

97. Choi, Y. & Chan, A. P. PROVEAN web server: A tool to predict the functional effect of amino acid substitutions and indels. *Bioinformatics* **31**, 2745-2747 (2015).

98. Archambeau, J. *et al.* Evaluating genomic offset predictions in a forest tree with high population genetic structure. *Am. Nat.* 000-000 (2025).

99. Warnes, G. genetics: Population Genetics. 1.3.8.1.3 https://doi.org/10.32614/CRAN.package.genetics (2012).

100. R Core Team. R: A language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria* (2023).

101. Caudullo, G., Welk, E. & San-Miguel-Ayanz, J. Chorological maps for the main European woody species. *Data Br.* **12**, 662-666 (2017).