Utility and validity of group atlas versus personalized functional network approaches for depressive constructs

Ellyn R. Butler, MS[1], Lauren B. Alloy, PhD[2], Damon D. Pham, BS[3], Noelle I. Samia, PhD[4, 5], Robin Nusslock, PhD[1, 6], and Amanda F. Mejia, PhD[3]

[1]Department of Psychology, Northwestern University, Evanston, IL 60208
[2]Department of Psychology and Neuroscience, Temple University, Philadelphia, PA 19122
[3]Department of Statistics, Indiana University, Bloomington, IN 47405
[4]Department of Statistics and Data Science, Northwestern University, Evanston, IL 60208
[5]NSF-Simons AI Institute for the Sky (SkAI), Chicago, IL 60611
[6]Institute for Policy Research, Northwestern University, Evanston, IL 60208

Co-author e-mail addresses: lalloy@temple.edu, ddpham@iu.edu, n-samia@northwestern.edu, nusslock@northwestern.edu, afmejia@iu.edu
Corresponding author: Ellyn Butler
Address: Northwestern University, Dept. of Psychology, 2029 Sheridan Ave, Evanston, IL 60201
Phone: (773) 255-5739
E-mail: ellynbutler2027@u.northwestern.edu

Running title: Group versus personalized functional network approaches

Abstract

Background: To understand the neurobiology underlying psychopathology, we need valid measurements of brain function. Group atlases for brain functional connectivity (FC) allow for efficient comparisons, but they fail to account for inter-individual variability in network topography, a problem that personalized methods address. We assess the validity and predictive utility of group and personalized approaches of quantifying FC by 1) comparing effect sizes of associations with clinical metrics; and 2) accounting for spatial features of brain networks when examining the association between FC and clinical metrics.

Methods: 324 teens ages 13-16 participated. Personalized networks were estimated using a hierarchical Bayesian model. Effect size comparisons were done by comparing the correlations between FC and clinical metrics (depression, ruminative coping style, and sensitivity to punishment/reward) with Steigler's Z-test. We also conducted regressions, with clinical metrics as the dependent variables. Those models included FC and spatial features, together and alone.

Results: The effect size comparisons did not survive FDR correction. However, exploratory permutation tests show that 1) the magnitude of the correlations with depression are larger on average for the intersection estimates of FC than the group estimates; and 2) the magnitude of the correlations with a ruminative coping style are larger on average for the intersection estimates of FC than the personalized estimate. The other comparisons conducted using permutation tests are not significant. Multiple regression analyses demonstrated that only spatial features of networks, not FC, are associated with sensitivity to reward.

Discussion: These results imply that the intersection estimates are more valid than the group estimates, and that the intersection estimates have greater predictive utility than personalized estimates. Further, spatial features of functions networks may be useful in and of themselves in certain contexts. Therefore, researchers in psychiatry should take into consideration functional network topography in order to gain a better understanding of the neurobiology underlying psychopathology.

1   Acknowledgements

E.R.B. conceptualized the work, conducted data analysis, and wrote the manuscript. A.F.M. supervised project conceptualization, execution, and data analysis. A.F.M. and D.D.P. assisted with image processing. N.I.S. provided feedback on data analysis. L.B.A. and R.N. obtained the funding for data collection, collected the data, and provided guidance on using the available data. All authors reviewed and edited the manuscript.

This work was supported by the National Institute of Mental Health [Grant No. R01 MH123473 (to L.B.A. and R.N.) and R01 MH126911 (to L.B.A. and R.N.)]; the National Institute on Aging [Grant No. R01 AG083919 (to A.F.M.)]; the National Institute of Biomedical Imaging and Bioengineering [Grant No. R01 EB027119 (to A.F.M.)]; the NSF-Simons AI-Institute for the Sky (SkAI) [Grant No. AST-2421845 (to N.I.S.)]; the Simons Foundation [Grant No. MPS-AI-00010513 (to N.I.S.)]; the National Science Foundation Graduate Research Fellowship [Grant No. DGE-2234667 (to E.R.B.)]; and the National Institute of Mental Health's National Research Service Award [Grant No. F31MH140552 (to E.R.B.)]. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

This research was supported in part through the computational resources and staff contributions provided for the Quest high performance computing facility at Northwestern University which is jointly supported by the Office of the Provost, the Office for Research, and Northwestern University Information Technology.

2   Disclosure

All authors declare no relevant conflicts of interests.

3   Introduction

Understanding the pathophysiology underlying psychopathology is a central goal of researchers and physicians alike [1]. In order to do this well, we must have valid measurements of key biological features [2], including aspects of brain function. Measures are valid when they tap into the construct as it has been defined. For instance, if we define depression as low mood or loss of interest or pleasure [3], then a valid measure would ask about aspects of these symptoms, as opposed to, for instance, one's intelligence.

In the field of cognitive and affective neuroscience, the focus has historically been on the reliability of different metrics of brain function, as opposed to the validity of the metrics. This hinders our ability to understand the pathophysiology and biological mechanisms under diseases and disorders [4]. Therefore, we must work towards establishing and ultimately improving the validity of brain metrics that show promise in helping us better understand pathophysiology and mechanisms in order to eventually impact health outcomes.

Functional connectivity (FC) derived from functional magnetic resonance imaging (fMRI) data has been used to study human brain networks for over 30 years [5], and is increasingly investigated in the context of human health and disease [6, 7, 8, 9, 10, 11]. Recognizing that there are many common features in functional brain networks across people, researchers have developed atlases, which group locations in the brain into a fixed number of networks [12, 13, 14]. These atlases then can be used by researchers to summarize high-dimensional functional data, improving our ability to compare results across studies, and to build theories about the nature of different networks.

While there are common features in brain networks across people, there is also substantial inter-individual variation in the placement, size, and shape of networks – the topography of networks [15]. This suggests that analyses using FC-based group atlases mix in effects across different brain networks into any given network [16]. Therefore, results finding differences in FC across groups or along continuous dimensions using FC-based group atlases may be finding such results due to differences in functional topography across groups and dimensions,

rather than true differences in FC [16, 17].

This problem has led to the development of personalized functional network approaches, which classify locations in the brain as members of certain networks for each individual person. Using these methods, researchers have shown that functional brain networks are dominated by stable group and individual factors [15, 18], with brain network topography being largely stable across time and cognitive states [15]. Further, the proportion of the cortical surface devoted to a particular network – or the *expansion* of the network – has been shown to be associated with related cognitive functions. For instance, the salience network has been shown to be more expansive among those who are more depressed [19], and greater fronto-parietal expansion has been shown to be associated with better cognitive function [20]. Therefore, it is important that the field increasingly move towards studying functional brain networks using personalized approaches.

Quantifying aspects of personalized functional networks is complicated by the fact that this is an emerging field, and as such, there are a variety of methods that differ substantially, with no single gold standard. One key consideration when choosing a method is whether or not to assume that each location in the brain belongs to exactly one network, or if each point should be allowed to belong to any number of networks. Previous work demonstrates that there are regions in the brain known as hubs that are critical for communication between networks, and as such, can be considered to belong to multiple networks [21, 22, 18]. However, popular methods such as Infomap and the multi-session hierarchical Bayesian model (MS-HBM) constrain points to belong to exactly one network [23, 15, 24, 25]. To address this problem, researchers have developed personalized functional network methods that allow points in the brain to contribute to any number of networks [26, 27, 28, 18, 29]. The network maps that emerge provide evidence in favor of relaxing the assumption that locations in the brain must belong to exactly one network. Using methods that allow for networks to overlap, researchers have found that there are in fact hubs at the individual level [18]. To provide the most valid representation of topography, it is important to allow for

each location in any given individual's brain to belong to any number of networks.

Nevertheless, most prior studies examining the relationship between cognitive or clinical metrics and brain functional organization are based on methods that constrain each point in the brain to belong to a single network. These studies have shown mixed results on the predictive utility of and effects sizes associated with personalized functional network metrics in comparison to group atlas approaches [25, 30]. Some have found larger effects for functional connectivity estimates based on personalized functional network approaches [25], whereas others have found larger effects based on group atlas approaches [30]. The larger effects sometimes found using group atlas approaches have at least two possible explanations: effects of mixing in connectivity from other networks, or of noise introduced by personalized functional network approaches. For instance, considering people who are depressed have more expansive salience networks than those who are not [19], if the Yeo17 group atlas [12] was used to estimate FC, signals from adjacent networks such as the control networks actually could be originating from the salience network among depressed individuals. Therefore, any associations between depression and control network connectivity may be driven by group differences in salience network topography. This signal mixing effect may explain why bigger effects are sometimes found with group atlas approaches to estimating FC. Alternatively, the group atlas approaches simply may have less noise in the topography underlying FC estimates, as group atlases are typically an amalgamation of hundreds or thousands of participants' data, whereas network membership derived from personalized approaches relies heavily on a single participant's data.

It is important to keep in mind that the value of the greater validity provided by personalized approaches depends on the research goals. If the goal of the researcher is simply to predict some phenotype, a group atlas approach to estimating FC could be preferable to a personalized network approach if it has greater predictive utility. But if the goal is to *understand* something about the network, a personalized approach likely is preferable, as it does not mix signals across networks, and consequently, we are able to interpret connectivity

estimates as originating from the selected network. Personalized approaches also can be expected to produce larger effect sizes in cases where there are true group differences in connectivity within a given network, barring excessive noise, given that they do not mix signals between networks. In other words, connectivity estimated using personalized approaches is more valid than connectivity estimated using group atlases, and as such, provides more interpretable findings.

In the current work, we evaluate whether signal mixing due to topographic differences drives the higher predictive utility sometimes found with group atlas-based approaches to estimating FC. To do so, we estimate FC within and between networks using three different network-defining approaches 2: 1) the canonical Yeo17 group atlas [12]; 2) personalized functional networks mapped onto the Yeo17 networks [27]; and 3) the *intersection* of each individual's personalized functional networks and the group atlas. The intersection approach allows one to mask out locations in the brain that belong to other networks according to the group atlas, and potential noise from the individual networks, while excluding locations that are not part of a specific network for a given individual. Therefore, the intersection approach only retains features specific to the individual, while benefiting from the greater reliability of group atlases.

By estimating FC using the intersection between the group atlas and each individual's personalized functional networks, we are able to determine if the greater predictive utility observed with FC estimates using a group atlas is due to the mixing in of FC from other networks due to individual differences in network topography. Specifically, we test the following hypotheses: Hypothesis #1 – if effect sizes with clinical metrics are greater for group-based estimates than intersection estimates, then what is useful about the group-based estimates of FC is that they are mixing in FC across networks within a single network FC estimate. Therefore, group-based estimates of FC would have higher predictive utility and low validity. Hypothesis #2 – if effect sizes with clinical metrics are greater for the intersection estimates than the personalized estimates, then the personalized estimates have excessive

noise. Therefore, personalized estimates of FC would have higher validity and low predictive utility. To provide greater clarity on the similarity in results across methods of quantifying FC, exploratory analyses are conducted to compare the effect sizes between the group-based and personalized estimates of FC.

Finally, we examine the extent to which FC estimated using a group atlas is associated with clinical metrics above and beyond spatial features. This directly tests if the group atlas effects typically conducted in the literature are truly FC effects, or if they are driven by spatial features of networks. We use the following clinical metrics across analyses: depression, ruminative coping style, sensitivity to punishment, and sensitivity to reward. These were selected to maximize power in the current study, as these measures are available for the most participants. While we focus on clinical metrics in the current work, these questions are relevant to brain-behavior associations using other phenotypes.

4 Materials and Methods

4.1 Participants

For this project, we use the Reward and Immune Systems in Emotion (RISE) [31] and Circadian, Reward, and Emotion Systems in Teens (CREST) datasets [32]. These datasets include fMRI and self-report data, in addition to a broad array of other metrics that are not the focus of this paper. Since July 2021, adolescents have been being recruited from the Philadelphia area to participate in the studies. To be eligible for RISE, participants must be 13-16 years old and not have experienced a major depressive episode. Further, participants are excluded from RISE if they have a history of cancer, heart disease, surgery, an ongoing autoimmune disease or disorder involving chronic inflammation (e.g., Crohn's Disease, Type 1 Diabetes, Lupus, asthma, severe allergies), if they are HIV positive, or if they took immunosuppressant medications in the past three months.

To be eligible for CREST, participants must be 13-16 years old and not have experienced

a manic or hypomanic episode or a bipolar spectrum disorder. Further, participants are excluded from CREST if they endorse any of the following: 1) use of melatonin or photo-sensitizing medications, 2) current or past treatment with bright light, or 3) shift work or travel across multiple time zones within the past month. These exclusion criteria are assessed in a phone interview and are designed to ensure that there are no contraindications or major confounding factors for the dim light melatonin onset (DLMO) procedures conducted in CREST. If a participant has traveled across time zones, but is otherwise eligible, participation is delayed for at least one month. Participants are excluded from both RISE and CREST if they endorse current psychotic symptoms, or have MRI contraindications (e.g., metal in the body, traumatic brain injury, pregnancy, severe claustrophobia; assessed with a standard MRI screening form). Participants in RISE are oversampled from the low end of the reward responsiveness dimension, whereas participants in CREST are oversampled from the high end [33, 34]. Temple University's Institutional Review Board approved the protocol.

The final sample consists of adolescents who have complete depression, ruminative coping style, sensitivity to punishment, sensitivity to reward, and neuroimaging data as of June 18, 2025. Neuroimaging data are considered complete for a given participant if they have a T1-weighted structural scan, at least five minutes worth of degrees of freedom remaining following confound regression from the combined functional scans, and have a mean framewise displacement less than 0.5mm. 473 adolescents at least began the screening process (i.e., answered at least the first question from the Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ)). 82 adolescents were missing some data from at least one of the clinical measures (40 missing depression; 32 missing ruminative coping style; and 34 missing sensitivity to punishment or reward). Four adolescents had some neuroimaging data but had not started the SPSRQ (i.e., are not part of the 473). 62 adolescents did not participate in the neuroimaging session. No adolescents had fewer than five minutes of data remaining after confound regression (range = 8.58 - 29.04 minutes; mean = 27.53 minutes;

median = 28.55 minutes). 29 adolescents had a mean framewise displacement greater than 0.5mm. 149 adolescents did not have sufficient clinical or neuroimaging data, resulting in a final sample of 324 adolescents.

Table 1: Sample characteristics by sex

	Female (N=192)	Male (N=132)	Overall (N=324)
Age			
Mean (SD)	15.5 (1.04)	15.4 (1.09)	15.5 (1.06)
Median [Min, Max]	15.5 [13.0, 18.1]	15.4 [13.1, 17.5]	15.5 [13.0, 18.1]
Race			
African American or Black	57 (29.7%)	22 (16.7%)	79 (24.4%)
Asian-American or Asian	7 (3.6%)	7 (5.3%)	14 (4.3%)
Biracial/Multiracial	10 (5.2%)	13 (9.8%)	23 (7.1%)
Caucasian/White	116 (60.4%)	89 (67.4%)	205 (63.3%)
Native Hawaiian or Other Pacific Islander	1 (0.5%)	0 (0%)	1 (0.3%)
Other	1 (0.5%)	1 (0.8%)	2 (0.6%)
Ethnicity			
Hispanic or Latino/a	13 (6.8%)	19 (14.4%)	32 (9.9%)
Not Hispanic or Latino/a	177 (92.2%)	111 (84.1%)	288 (88.9%)
Missing	2 (1.0%)	2 (1.5%)	4 (1.2%)
Average Caregiver Years of Education			
Mean (SD)	16.5 (2.32)	17.1 (2.45)	16.7 (2.39)
Median [Min, Max]	17.0 [3.00, 22.5]	17.0 [12.0, 24.0]	17.0 [3.00, 24.0]
Missing	7 (3.6%)	6 (4.5%)	13 (4.0%)
Family's Gross Total Income			
$0 - $4,999	1 (0.5%)	0 (0%)	1 (0.3%)
$5,000 - $19,999	2 (1.0%)	0 (0%)	2 (0.6%)
$20,000 - $34,999	6 (3.1%)	4 (3.0%)	10 (3.1%)
$35,000 - $49,999	11 (5.7%)	3 (2.3%)	14 (4.3%)
$50,000 - $74,999	22 (11.5%)	7 (5.3%)	29 (9.0%)
$75,000 - $99,999	18 (9.4%)	8 (6.1%)	26 (8.0%)
$100,000 - $149,999	25 (13.0%)	25 (18.9%)	50 (15.4%)
$150,000 - $199,999	28 (14.6%)	24 (18.2%)	52 (16.0%)
$200,000 - $249,999	22 (11.5%)	15 (11.4%)	37 (11.4%)
$250,000 - $299,999	16 (8.3%)	15 (11.4%)	31 (9.6%)
$300,000 and higher	25 (13.0%)	15 (11.4%)	40 (12.3%)
Not known	2 (1.0%)	2 (1.5%)	4 (1.2%)
Missing	14 (7.3%)	14 (10.6%)	28 (8.6%)

Note: 12 years of education corresponds to having completed high school. Average Caregiver Years of Education is the average across two caregivers if the child has two caregivers, and is the years of education of a single caregiver if they have one caregiver.

4.2 Assessment

Depression

We use the Beck Depression Inventory - II (BDI-II) to assess depression severity over the preceding two weeks [35]. The BDI-II taps into constructs of sadness, pessimism, suicidal thoughts or wishes, loss of interest, worthlessness, and difficulty with concentration, among other depression-related constructs. Historically, the BDI-II has been shown to have a two-factor structure [36, 37] – comprising factors tapping into cognitive-affective and somatic-vegetative symptoms of depression. Exploratory factor analyses on the current sample demonstrate a largely unidimensional structure ($\lambda_1 = 4.65$, $\lambda_2 = 1.15$, $\alpha = 0.86$, $\omega_H = 0.69$), which implies good internal consistency and justifies using a sum score of the items [38]. The BDI-II has demonstrated good psychometic properties, including high test-retest reliability ($r = 0.73$ to 0.96), and convergent validity (i.e., high correlations with other measures of depression) [37].

Ruminative coping style

The 10-item Ruminative Responses Scale (RRS) is used to assess tendency to use a ruminative coping style [39]. The RRS taps into constructs of thinking about how alone one feels, how fatigued one is, how one wishes a situation had gone better, and how angry one is with oneself, among other rumination-related constructs. Historically, the RRS has been shown to have a two-factor structure [39] – reflective pondering and brooding. Exploratory factor analyses on the current sample demonstrate a largely unidimensional structure ($\lambda_1 = 8.48$, $\lambda_2 = 1.27$, $\alpha = 0.92$, $\omega_H = 0.75$), which implies good internal consistency and justifies using a sum score of the items [38]. The RRS has demonstrated decent psychometric properties, including moderate test-retest reliability ($r = 0.60$) [39].

Sensitivity

We use the Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ) to assess sensitivity to punishments and rewards [34]. The SPSRQ taps into constructs such as refraining from doing illegal activities, being motivated by money, doing things to be

praised, and showing off in a group. Historically, the SPSRQ has been shown to have a two-factor structure [34] – sensitivity to punishment and sensitivity to reward. Exploratory factor analyses on the current sample demonstrate a largely two-dimensional structure ($\lambda_1 = 6.36$, $\lambda_2 = 4.56$, $\alpha = 0.83$, $\omega_H = 0.52$), which justifies using separate sum score of the items for sensitivity to punishment and sensitivity to reward, respectively [38]. The SPSRQ has demonstrated good psychometric properties, with high test-retest reliability at three months for the punishment ($r = 0.89$) and reward ($r = 0.87$) subscales [34].

4.3   Neuroimaging

Acquisition

Imaging data were collected at the Temple University Brain Research & Imaging Center (TUBRIC) on a Siemens MAGNETOM Prisma 3-Tesla scanner with a 64-channel phased-array parallel transmit and receive RF head/neck coil. Structural images were acquired using an MPRAGE sequence to acquire 208 axial slices (voxel size = 0.8mm^3; TR=2300ms; TE=2.99ms; FOV=256x256; Matrix=320x320; Flip Angle=7°). Three functional scans were used: The Chatroom Interact task [40], and two runs of the Monetary Incentive Delay (MID) task [41]. The Chatroom Interact task uses a slice-accelerated multiband EPI sequence (multiband acceleration factor: 2. GRAPPA acceleration factor: 2) covering 64 axial slices (voxel size = 2.0mm^3; TR=2050ms; TE=25ms; FOV=208x208mm; Matrix=104x104; Flip Angle=76°). The MID task also uses a slice-accelerated multiband EPI sequence with identical scan parameters. A total of 390 volumes were collected for the Chatroom Interact task, and 289 for each of the runs of MID.

These tasks can be concatenated to increase reliability of functional network metrics [42]. Further, functional network metrics can be validly and reliably estimated using task data [43, 44], and functional network metrics derived from tasks tend to have greater predictive utility than rest even when the task is not aligned with the phenotype of interest [43]. We therefore concatenate these three functional scans after processing for use in all analyses.

Processing

We use fMRIPrep version 23.2.0 [45], Connectome Workbench [46], and R packages [47, 48] to process the magnetic resonance imaging data. Processing steps include correcting T1-weighted images for intensity non-uniformity and skull-stripping; slice-time correcting the functional images, resampling them into their native space, and then projecting them into the fsLR-32k surface space (resampled to 10k per hemisphere for computational purposes). We use simultaneous nuisance regression (translation (x, y, z), rotation (yaw, pitch, roll), and global signal; the derivatives, squares, and squared derivatives of each of those signals; outlier flagging with DVARS (dual cutoff method) [49]; and .01 Hz high-pass filter using 12 discrete cosine transform bases) [50]. DVARS is estimated on otherwise nuisance-regressed data to reduce redundancy in the covariates [48].

Network metrics

To construct individual-level network engagement maps, we use a Bayesian Brain Mapping (BBM) technique [51] (Figure 1). BBM employs a hierarchical Bayesian framework with group-derived priors on spatial functional topography. The role of the priors is to mitigate noise and produce reliable maps of individual vertex-level network engagement and membership [27]. In the current work, the priors are derived from the data used in the downstream analyses, resulting in an empirical Bayesian approach. Since the priors are established before model fitting, the BBM model can be fit for each individual separately for efficient computation. We use the Yeo17 atlas [12] as a template instead of group independent component analysis (ICA) maps as in previous work [27], to allow comparisons to FC estimates derived using the Yeo17 group atlas. The Yeo17 atlas was used as a template to guide the prior, such that it was assumed that there are 17 networks, which should resemble the topographical layout of the Yeo17 atlas, while being allowed to overlap.

We derive prior mean and variance maps from the Yeo17 group atlas in combination with the processed functional data from each participant in the sample [51], similarly to how we have done previously [52]. Briefly, each functional scan is split down the middle to

produce pseudo test-retest data, and for each pseudo-session we produce a map of network engagement for each of the 17 networks using a variation of dual regression [53]. In the first "regression" we compute the median time course across the vertices within each parcel, and in the second regression we regress those time courses against the fMRI time series to obtain maps of network engagement. Using the resulting set of test-retest network engagement maps for all participants, we estimate the prior mean and between-subject variance maps using variance decomposition [27]. We then use these prior mean and variance maps as priors in the BBM model for individual network maps.

After fitting the BBM model, we identify vertices belonging to each of the 17 networks by performing the Bayesian equivalent of a hypothesis test at every vertex, based on the estimated engagement level and its posterior standard deviation. While in hypothesis testing it is common to test against a null hypothesis of zero engagement, here to avoid statistically significant but negligible effect sizes, we test against a null hypothesis of negligible-or-smaller engagement. For any vertices where the null hypothesis is rejected, we can conclude that they exceed a predetermined minimum effect size. Specifically, we set the minimum effect size to correspond to a z-score of $z = 1$ of the prior mean map. To avoid false positives, for each network we apply False Discovery Rate (FDR) correction [54] to control the FDR at 0.01 across all 18566 vertices.

Figure 1: Pipeline for estimating personalized networks using the BayesBrainMap R package [51]. Functions that produce the surface outputs in the blue boxes are listed below their corresponding boxes. The Default A network is highlighted for illustration purposes. Features from all Yeo17 networks were estimated.

After identifying vertices with significant engagement in each network, we compute inter- and intra-network connectivity. *Inter*-network connectivity is computed as the correlation between the time series for each network. The network time series are computed across all significant vertices in the network, weighted by the surface area that each vertex occupies in fsLR-32k space and the estimated engagement of the vertex for a given network. Specifically, the following equation was used to compute connectivity (C_{ghi}) between networks g and $h = 1, \ldots, 17$ for subject i:

$$C_{ghi} = \mathrm{Corr}\left(\frac{\sum_{v \in \mathcal{V}_g} a_v s_{giv} \mathbf{t}_{iv}}{\sum_{v \in \mathcal{V}_g} a_v s_{giv}}, \frac{\sum_{v \in \mathcal{V}_h} a_v s_{hiv} \mathbf{t}_{iv}}{\sum_{v \in \mathcal{V}_h} a_v s_{hiv}} \right) \tag{1}$$

where $\mathcal{V}_g$ indexes the vertices belonging to networks $g = 1, \ldots, 17$, and $h = 1, \ldots, 17$, respectively; $a_v =$ the surface area of the vertex v in fs-LR32k space; $s_{giv} =$ the engagement level for subject i in network g at vertex v; and $\mathbf{t}_{iv} =$ the time series for subject i at vertex v.

The intersection estimates of inter-network connectivity are computed in a similar manner, except vertices that are not belonging to a given network in the Yeo17 atlas were excluded (Figure 2). *Intra*-network connectivity for each network g, D_{gi}, is computed as the average correlation in the BOLD time series between all pairs of vertices in the network, weighted by the surface area that each vertex occupies in fsLR-32k space and the engagement of the vertex in network g:

$$D_{gi} = \frac{\sum_{v_1 \in \mathcal{V}_g} \sum_{v_2 > v_1} a_{v_1} s_{giv_1} a_{v_2} s_{giv_2} \mathrm{Corr}(\mathbf{t}_{iv_1}, \mathbf{t}_{iv_2})}{\sum_{v_1 \in \mathcal{V}_g} \sum_{v_2 > v_1} a_{v_1} s_{giv_1} a_{v_2} s_{giv_2}}. \tag{2}$$

For group-based FC, the correlations between pairs of vertices were only weighted by the surface area of the vertices, as all vertices have equal engagement in the Yeo17 group atlas.

Figure 2: Group, personalized, and intersection maps. (A) shows the Default A and Somato-motor B networks isolated from the Yeo17 group atlas; (B) shows personalized thresholded engagement maps for an individual; and (C) shows intersection maps for this same individual. Black arrows in (B) and (C) point out differences between the personalized and intersection thresholded engagement maps.

Finally, we derive spatial and FC regressors for the multiple regression models. For the spatial covariates, we perform singular value decomposition on the engagement maps for each network separately. To reduce background noise, we first threshold the engagement maps such that negligible engagements are set to zero. Specifically, we apply a threshold of

$z = 2$ on the prior mean map. Singular vectors that explain at least 1% of the variance in engagement were retained for the regressions. For the FC regressors, we compute intra- and inter-network FC using the Yeo17 atlas and perform singular value decomposition to reduce the dimensionality, and retain the singular vectors that explain at least 1% of the variance in FC.

4.4 Statistical Analyses

We conduct two sets of analyses to evaluate the utility and validity of group atlas versus personalized functional network approaches: 1) effect sizes comparisons of the association between intra- and inter-network FC estimated using group, personalized, and intersection network maps and a series of clinical metrics (effect size comparisons); and 2) ability of intra- and inter-network FC estimated using a group atlas to predict clinical metrics, above and beyond network topography features (multiple regressions).

Effect Size Comparisons

We perform comparisons of correlations between clinical metrics and FC estimated using a group atlas, personalized networks, and the intersection between the two. We use Steiger's Z-test for equality of correlations [55], as it accounts for dependence between the observations across the two correlations. Specifically, for the kth clinical metric, $k = 1, \ldots, 4$, and jth FC estimate (inter- and intra-network FC), $j = 1, \ldots, 153$, r_{1jk} = the correlation between the metric and group-based FC; r_{2jk} = the correlation between the metric and personalized FC; and r_{3jk} = the correlation between the metric and intersection FC.

Dropping the j and k subscripts for simplicity momentarily, the test statistic Z_{AB} for comparing correlations r_A and r_B, where $A \neq B$, $A, B = 1, 2, 3$, takes the following form ():

$$Z = \frac{z_A - z_B}{\sqrt{\frac{2(1-z_{AB})}{n-3}}}, \ Z \sim \mathbb{N}(0, 1) \text{ asymptotically under } H_0 \tag{3}$$

where z_A and z_B, are the Fisher-transformed correlations r_A and r_B, respectively, and z_{AB} is

the Fisher-transformed correlation between the two FC metrics of interest. To evaluate Hyp. #1, we test if r_{1jk} and r_{2jk} differ for each measure of inter- and intra-network connectivity j and metric k. That is, $r_A = r_{1jk}$, $r_B = r_{2jk}$, and $r_{AB} = r_{12j}$. To evaluate Hyp. #2 using the continuous self-report measures, we test if r_{2jk} and r_{3jk} differ for each network j and metric k. That is, $r_A = r_{2jk}$, $r_B = r_{3jk}$, and $r_{AB} = r_{23j}$. The null hypotheses are rejected based on the p-values from the Steiger test, controlling the False Discovery Rate at 0.01 across all 2142 tests to account for multiple comparisons [54].

Multiple Regressions

For each clinical metric $k = 1, \ldots, 4$, we fit a series of linear models using observations from n participants, where the dependent variable of interest is the clinical metric (Y_{ik}), with error terms $\epsilon_{ik} \overset{iid}{\sim} N(0, \sigma_k^2)$, $i = 1, \ldots, n$. Independent variables include (a) the SVD singular vectors derived from the FC estimates based on the Yeo17 group atlas $(\boldsymbol{w}_i)$ and (b) the SVD singular vector scores from the spatial network engagement maps $(\boldsymbol{u}_{ij})$. Recall that we perform a separate SVD on the engagement maps for each network, so $\boldsymbol{u}_{ij}$ is indexed by both subjects i and spatial singular vectors $j = 1, \ldots, J$. We fit four separate sets of multiple regression models, one set for each clinical metric. For each metric, the full model takes the following form:

$$Y_{ik} = \beta_{0k} + \boldsymbol{w}_i^T \boldsymbol{\beta}_k + \left[\sum_{j=1}^{J} \boldsymbol{u}_{ij}^T \boldsymbol{\beta}_{jk}^{S2} \right] + \epsilon_{ik}. \tag{4}$$

The first reduced model includes only the spatial singular vectors and takes the following form. Comparing the full model to this reduced model allows us to determine if the group atlas-derived FC estimates $(\boldsymbol{u}_i)$ explain any additional variance in the clinical metrics above and beyond the spatial singular vectors $(\boldsymbol{u}_{ij})$, and how much:

$$Y_{ik} = \beta_{0k} + \left[\sum_{j=1}^{J} \boldsymbol{u}_{ij}^T \boldsymbol{\beta}_{jk}^{S2} \right] + \epsilon_{ik}. \tag{5}$$

The second reduced model includes only the FC predictors and takes the following form.

Comparing the full model to this reduced model allows us to determine if the spatial singular vectors explain a portion of the predictive power of the group atlas-derived FC estimates, and how much:

$$Y_{ik} = \beta_{0k} + \boldsymbol{W}_i^T \boldsymbol{\beta}_k + \epsilon_{ik}. \tag{6}$$

We use nested model F tests to compare the reduced and full models. We only perform these nested model tests for clinical metrics where the full model explained a significant amount of variance in the metric.

5   Results

In this section, we summarize the results of the effect size comparisons and the multiple regressions model comparisons to assess the validity and predictive utility of personalized and group-based approaches to quantifying FC.

The effect size comparison analyses show that, after applying FDR correction, none of the individual effect size comparisons remain significant. However, a different picture emerges when examining the average difference in magnitude of the associations across all of the FC variables using permutation tests (see Supplement for details). Figure 3 shows that 1) the magnitude of the correlations with depression are larger on average for the intersection estimates of FC than the group estimates ($p = 0.038$); and 2) the magnitude of the correlations with a ruminative coping style are larger on average for the intersection estimates of FC than the personalized estimates ($p = 0.010$). These results imply that 1) the intersection estimates may remove signal from non-target networks that are included in the group estimates, which for the group estimates reduces observed effect sizes; and 2) the intersection estimates may remove noise that is often characteristic of personalized methods. The other comparisons conducted using permutation tests are not significant.

Turning now to the multiple regression models, 10 FC singular vectors and 115 spatial

Figure 3: Histograms of the absolute values of the correlations between clinical metrics and different ways of quantifying FC. Comparisons were made to test if the absolute values of the correlations between FC quantified using one of the three methods (group, personalized, intersection) and the clinical metrics differed across methods. None of the comparisons survived FDR correction. Permutation tests were then used to determine if, on average, differences in the absolute values of the correlations differ on average across methods of quantifying FC for each clinical metric. Significant results are indicated with an asterisk. A.) Comparison of the correlations with clinical metrics for the group and intersection methods; B.) Comparison of the correlations with clinical metrics for the personalized and intersection methods; and C.) Comparison of the correlations with clinical metrics for the group and personalized methods.

singular vectors are retained using the 1% variance explained cutoff for inclusion of singular vectors from the SVDs. The multiple regression analyses showed that neither FC nor spatial features of networks explained a significant portion of variance in depression, ruminative coping style, or sensitivity to punishment, based on the full model. As such, we do not consider the nested models for these clinical measures.

Spatial features of networks, but not FC, explain a significant portion of variance in sensitivity to reward, based on the reduced models. Using nested model comparisons, FC did not explain additional variance in sensitivity to reward above and beyond spatial features ($F(198, 208) = 0.84$, $p = 0.588$). As a sensitivity analysis, we repeat the multiple regression models using a 2% variance explained cutoff for inclusion of singular vectors from the SVDs. Using this criterion, 5 FC singular vectors and 28 spatial singular vectors are retained. Results of the full and reduced models using these features are consistent with those reported above.

	Depression		Rumination		Punishment		Reward	
	FC	Spatial	FC	Spatial	FC	Spatial	FC	Spatial
F	0.334	0.896	0.685	0.743	1.055	1.200	0.956	**1.687**
p-value	0.970	0.741	0.738	0.961	0.398	0.129	0.482	**< 0.0001**
R^2	0.011	0.331	0.021	0.291	0.033	0.399	0.030	**0.483**
R^2_{adj}	−0.021	−0.038	−0.010	−0.101	0.002	0.066	−0.001	**0.196**

Table 2: Full model results for functional connectivity (FC) and spatial predictors, for each of the four clinical measures we consider (depression, ruminative coping style, sensitivity to punishment, and sensitivity to reward). To test each collection of predictors, we perform an F-test on all the corresponding coefficients. The first row shows the value of the $F(10, 313)$ test statistic, and the second row shows the corresponding p-value.

6 Discussion

There is increasing evidence and support for the idea that accounting for the personalized nature of functional topography may result in more valid estimates of FC. Yet, both personalized and group atlas-based approaches to estimating FC may have good predictive utility, depending on the outcome of interest. In the current study, we interrogate specific hypothe-

ses around the advantages and disadvantages of group versus personalized approaches by comparing the predictive utility of three different methods for quantifying FC. In addition to group atlas-based and personalized network-based FC, we introduce a novel intersection-based approach designed to realize the advantages of both. Further, we interrogate whether the predictive utility of group atlas-based FC is derived, at least in part, from aliased-in spatial features of the networks.

After controlling for multiple comparisons over the 1836 comparisons of correlations between FC and clinical metrics, we find no statistically significant differences in the magnitude of the correlations between various ways of quantifying FC. However, after conducting a series of permutations tests, we see that 1) the magnitude of the correlations with depression are larger on average for the intersection estimates of FC than the group estimates; and 2) the magnitude of the correlations with a ruminative coping style are larger on average for the intersection estimates of FC than the personalized estimate. These results imply that 1) the intersection estimates may remove signal from non-target networks that are included in the group estimates, which for the group estimates reduces observed effect sizes, suggesting that the intersection estimates are more valid than the group estimates; and 2) the intersection estimates may remove noise that is often characteristic of personalized methods, suggesting that intersection estimates have greater predictive utility than personalized estimates. The other comparisons conducted using permutation tests are not significant. These results suggest that, with respect to some of the clinical outcome metrics used in the current study, the intersection estimates may be the best in terms of both predictive utility and validity.

Using multiple regression models we find that, among the four clinical metrics tested, only spatial features explain a significant portion of the variance in sensitivity to reward, and for no other clinical metrics. The fact that spatial network features are associated with sensitivity to reward suggests that there are functional brain features that do meaningfully relate to clinical outcomes. Therefore, future studies may benefit from deriving spatial features of networks when using personalized methods, instead of simply considering FC.

This represents an opportunity unique to personalized methods, as group atlases assume that spatial features of networks are invariant across people.

The current study is limited by a small set of outcome metrics and a relatively small number of subjects, particularly given the high-dimensional nature of our functional network features. With only four clinical measures to serve as dependent variables, we are restricted in our ability to determine the relative predictive utility and validity of our FC metrics. Other potentially relevant dependent variables include performance on vocabulary, reading, and crystallized cognition tasks [56], and behaviors such as mobility as captured by smartphones [57]. The study is also limited by the fact that there are 324 subjects in the current work, while we are trying to draw inferences about thousands of functional brain network features. Notably, this is a common problem in neuroimaging studies, with many studies having far fewer subjects [58, 59], and we did conduct a variety of dimensionality reduction and multiple comparison correction procedures. Nevertheless, this remains a concern, as it is possible that with a more focused, theory-driven analysis we may have been able to find significant results.

Our study has several notable strengths. First, we propose a novel intersection network mapping approach to realize the advantages of both group atlases and personalized network maps. This allows us to distinguish the features specific to the group and personalized approaches. The intersection improves over the group atlas by excluding features that do not map onto an individual's functional topography. It also may improve over the personalized network maps by capitalizing on the noise reduction properties of group atlases, though it runs the risk of missing important signal from hubs and true features of subjects' networks. Therefore, our intersection-based approach may be useful for future studies examining the validity of methods for quantifying FC. Further, intersection network maps may provide a reasonable compromise between noisy personalized network maps and rigid group atlases for quantifying FC.

Finally, a strength of our study is our use of a Bayesian brain mapping (BBM) method that meaningfully improves over popular methods for estimating personalized networks [27].

Popular methods include dual regression [53] and Infomap [15, 42]. BBM improves over dual regression by reducing noise with population-derived priors on network topography. BBM also improves over methods like Infomap by allowing vertices to be classified as belonging to multiple networks. This represents an important advantage of BBM, as we know that there are hubs – regions that affiliate with multiple networks – in the brain [60, 61, 22]. Notably, work that has compared personalized methods that allow for network overlap to those that do not has found that allowing for network overlap produces more reliable maps than constraining locations in the brain to belong to a single network [29].

Future research examining the utility and validity of different methods of quantifying FC should include outcomes metrics that have been shown to be more robustly associated with FC. Generally speaking, FC is more strongly associated with cognitive metrics than clinical metrics [56], so it will be important to conduct similar analyses in a large dataset with subjects who have cognitive measures. In such a dataset, we may gain greater clarity on the nature of what is useful about group and personalized estimates of FC.

Our findings support the idea that future studies that seek to uncover the neurobiological bases of psychopathology should continue to investigate spatial features of functional networks, as these features are emerging as potentially important biological substrates of psychopathology [62, 19, 63, 52]. Therefore, whether or not personalized approaches provide more valid estimates of FC than group atlases, it will continue to be important to utilize personalized network approaches as they allow us to quantify spatial features of functional networks. More broadly, if we are able to validly measure these spatial features, we will be able to gain a better understanding of the neurobiology underlying psychopathology, opening the doors to identifying potentially effective targets for intervention.

References

[1] Bruce N Cuthbert. "The RDoC framework: facilitating transition from ICD/DSM to dimensional approaches that integrate neuroscience and psychopathology". In: *World psychiatry* 13.1 (2014), pp. 28–35.

[2] George A Morgan et al. "Measurement validity". In: *Journal of the American Academy of Child & Adolescent Psychiatry* 40.6 (2001), pp. 729–731.

[3] American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* 5th. 2013. DOI: 10.1176/appi.books.9780890425596.

[4] Brian Kraus et al. "Beyond Reliability: The Impact of Group-Defined Atlases on Neuroimaging Measurement Validity". In: *Biological Psychiatry* 97.9 (2025), S65.

[5] Bharat Biswal et al. "Functional connectivity in the motor cortex of resting human brain using echo-planar MRI". In: *Magnetic resonance in medicine* 34.4 (1995), pp. 537–541.

[6] Damien A Fair et al. "Functional brain networks develop from a "local to distributed" organization". In: *PLoS computational biology* 5.5 (2009), e1000381.

[7] Dardo Tomasi and Nora D Volkow. "Aging and functional brain networks". In: *Molecular psychiatry* 17.5 (2012), pp. 549–558.

[8] Janis Brakowski et al. "Resting state brain network function in major depression–depression symptomatology, antidepressant treatment effects, future research". In: *Journal of psychiatric research* 92 (2017), pp. 147–159.

[9] Cedric Huchuan Xia et al. "Linked dimensions of psychopathology and connectivity in functional brain networks". In: *Nature communications* 9.1 (2018), p. 3003.

[10] Valerie J Sydnor et al. "Neurodevelopment of the association cortices: Patterns, mechanisms, and implications for psychopathology". In: *Neuron* 109.18 (2021), pp. 2820–2846.

[11]   Matej Perovnik et al. "Functional brain networks in the evaluation of patients with neurodegenerative disorders". In: *Nature Reviews Neurology* 19.2 (2023), pp. 73–90.

[12]   BT Thomas Yeo et al. "The organization of the human cerebral cortex estimated by intrinsic functional connectivity". In: *Journal of neurophysiology* (2011).

[13]   Jonathan D Power et al. "Functional network organization of the human brain". In: *Neuron* 72.4 (2011), pp. 665–678.

[14]   Alexander Schaefer et al. "Local-global parcellation of the human cerebral cortex from intrinsic functional connectivity MRI". In: *Cerebral cortex* 28.9 (2018), pp. 3095–3114.

[15]   Caterina Gratton et al. "Functional brain networks are dominated by stable group and individual factors, not cognitive or daily variation". In: *Neuron* 98.2 (2018), pp. 439–452.

[16]   Janine Diane Bijsterbosch et al. "The relationship between spatial configuration and functional connectivity of brain regions". In: *elife* 7 (2018), e32992.

[17]   Janine Diane Bijsterbosch et al. "The relationship between spatial configuration and functional connectivity of brain regions revisited". In: *Elife* 8 (2019), e44890.

[18]   Janine D Bijsterbosch et al. "Evaluating functional brain organization in individuals and identifying contributions to network overlap". In: *Imaging Neuroscience* 1 (2023), pp. 1–19.

[19]   Charles J Lynch et al. "Frontostriatal salience network expansion in individuals in depression". In: *Nature* (2024), pp. 1–10.

[20]   Arielle S Keller et al. "Personalized functional brain network topography is associated with individual differences in youth cognition". In: *Nature communications* 14.1 (2023), p. 8411.

[21]   Martijn P Van den Heuvel and Olaf Sporns. "Network hubs in the human brain". In: *Trends in cognitive sciences* 17.12 (2013), pp. 683–696.

[22] Stuart Oldham and Alex Fornito. "The development of brain network hubs". In: *Developmental cognitive neuroscience* 36 (2019), p. 100607.

[23] Martin Rosvall and Carl T Bergstrom. "Maps of information flow reveal community structure in complex networks". In: *arXiv preprint physics.soc-ph/0707.0609* 3 (2007).

[24] Benjamin A Seitzman et al. "Trait-like variants in human functional brain networks". In: *Proceedings of the National Academy of Sciences* 116.45 (2019), pp. 22851–22861.

[25] Ru Kong et al. "Individual-specific areal-level parcellations improve functional connectivity prediction of behavior". In: *Cerebral cortex* 31.10 (2021), pp. 4477–4500.

[26] Samuel J Harrison et al. "Large-scale probabilistic functional modes from resting state fMRI". In: *NeuroImage* 109 (2015), pp. 217–231.

[27] Amanda F Mejia et al. "Template independent component analysis: targeted and reliable estimation of subject-level brain networks using big data population priors". In: *Journal of the American Statistical Association* 115.531 (2020), pp. 1151–1177.

[28] Amanda F Mejia et al. "Template independent component analysis with spatial priors for accurate subject-level brain network estimation and inference". In: *Journal of Computational and Graphical Statistics* 32.2 (2023), pp. 413–433.

[29] Robert JM Hermosillo et al. "A precision functional atlas of personalized network topography and probabilities". In: *Nature neuroscience* 27.5 (2024), pp. 1000–1013.

[30] Alexis Grace Porter. "Understanding Cognitive and Psychopathological Variability: The Role of Individual Differences in Machine Learning Applications". PhD thesis. Northwestern University, 2024.

[31] Lauren B Alloy et al. "Reward and Immune Systems in Emotion (RISE) prospective longitudinal study: Protocol overview of an integrative reward-inflammation model of first onset of major depression in adolescence". In: *Brain, Behavior, & Immunity-Health* 30 (2023), p. 100643.

[32] Lauren B Alloy et al. "Circadian, Reward, and Emotion Systems in Teens prospective longitudinal study: protocol overview of an integrative reward-circadian rhythm model of first onset of bipolar spectrum disorder in adolescence". In: *BMC psychiatry* 23.1 (2023), p. 602.

[33] Charles S Carver and Teri L White. "Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: the BIS/BAS scales." In: *Journal of personality and social psychology* 67.2 (1994), p. 319.

[34] Rafael Torrubia et al. "The Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ) as a measure of Gray's anxiety and impulsivity dimensions". In: *Personality and individual differences* 31.6 (2001), pp. 837–862.

[35] Aaron T Beck, Robert A Steer, Gregory K Brown, et al. "Beck depression inventory". In: (1996).

[36] Mark A Whisman, John E Perez, and Wiveka Ramel. "Factor structure of the Beck Depression Inventory—Second Edition (BDI-II) in a student sample". In: *Journal of clinical psychology* 56.4 (2000), pp. 545–551.

[37] Yuan-Pang Wang and Clarice Gorenstein. "Psychometric properties of the Beck Depression Inventory-II: a comprehensive review". In: *Brazilian Journal of Psychiatry* 35 (2013), pp. 416–431.

[38] Keith F Widaman and William Revelle. "Thinking thrice about sum scores, and then some more about measurement and analysis". In: *Behavior Research Methods* 55.2 (2023), pp. 788–806.

[39] Wendy Treynor, Richard Gonzalez, and Susan Nolen-Hoeksema. "Rumination reconsidered: A psychometric analysis". In: *Cognitive therapy and research* 27 (2003), pp. 247–259.

[40] Jennifer S Silk et al. "Peer acceptance and rejection through the eyes of youth: pupillary, eyetracking and ecological data from the Chatroom Interact task". In: *Social cognitive and affective neuroscience* 7.1 (2012), pp. 93–105.

[41] Brian Knutson et al. "Distributed neural representation of expected value". In: *Journal of Neuroscience* 25.19 (2005), pp. 4806–4812.

[42] Brian T Kraus et al. "Network variants are similar between task and rest states". In: *Neuroimage* 229 (2021), p. 117743.

[43] Emily S Finn. "Is it time to put rest to rest?" In: *Trends in cognitive sciences* 25.12 (2021), pp. 1021–1032.

[44] Jingnan Du et al. "Within-Individual Precision Mapping of Brain Networks Exclusively Using Task Data". In: *bioRxiv* (2025), pp. 2025–02.

[45] Oscar Esteban et al. "fMRIPrep: a robust preprocessing pipeline for functional MRI". In: *Nature methods* 16.1 (2019), pp. 111–116.

[46] Daniel S Marcus et al. "Human Connectome Project informatics: quality control, database services, and data visualization". In: *Neuroimage* 80 (2013), pp. 202–219.

[47] Damon D Pham, John Muschelli, and Amanda F Mejia. "ciftiTools: A package for reading, writing, visualizing, and manipulating CIFTI files in R". In: *NeuroImage* 250 (2022), p. 118877.

[48] Damon D Pham et al. "Less is more: balancing noise reduction and data retention in fMRI with data-driven scrubbing". In: *NeuroImage* 270 (2023), p. 119972.

[49] Soroosh Afyouni and Thomas E Nichols. "Insight and inference for DVARS". In: *NeuroImage* 172 (2018), pp. 291–312.

[50] C Büchel et al. "Characterizing stimulus–response functions using nonlinear regressors in parametric fMRI experiments". In: *Neuroimage* 8.2 (1998), pp. 140–148.

[51] Nohelia Da Silva Sanchez et al. "Bayesian brain mapping: population-informed individualized functional topography and connectivity". In: *arXiv preprint arXiv:2602.01551* (2026).

[52] Ellyn R Butler et al. "Sex differences in response to violence: role of salience network expansion and connectivity on depression". In: *Translational Psychiatry* 15.1 (2025), p. 427.

[53] Christian F Beckmann et al. "Group comparison of resting-state FMRI data using multi-subject ICA and dual regression". In: *Neuroimage* 47.Suppl 1 (2009), S148.

[54] Yoav Benjamini and Yosef Hochberg. "Controlling the false discovery rate: a practical and powerful approach to multiple testing". In: *Journal of the Royal statistical society: series B (Methodological)* 57.1 (1995), pp. 289–300.

[55] James H Steiger. "Tests for comparing elements of a correlation matrix." In: *Psychological bulletin* 87.2 (1980), p. 245.

[56] Jianzhong Chen et al. "Shared and unique brain network features predict cognitive, personality, and mental health scores in the ABCD study". In: *Nature communications* 13.1 (2022), p. 2217.

[57] Cedric Huchuan Xia et al. "Mobile footprinting: linking individual distinctiveness in mobility patterns to mood, sleep, and brain functional connectivity". In: *Neuropsychopharmacology* 47.9 (2022), pp. 1662–1671.

[58] Denes Szucs and John PA Ioannidis. "Sample size evolution in neuroimaging research: An evaluation of highly-cited studies (1990–2012) and of latest practices (2017–2018) in high-impact journals". In: *NeuroImage* 221 (2020), p. 117164.

[59] Colin G DeYoung et al. "Beyond increasing sample sizes: Optimizing effect sizes in neuroimaging research on individual differences". In: *Journal of Cognitive Neuroscience* 37.6 (2025), pp. 1023–1034.

[60] Olaf Sporns, Christopher J Honey, and Rolf Kötter. "Identification and classification of hubs in brain networks". In: *PloS one* 2.10 (2007), e1049.

[61] Jonathan D Power et al. "Evidence for hubs in human functional brain networks". In: *Neuron* 79.4 (2013), pp. 798–813.

[62] Zaixu Cui et al. "Linking individual differences in personalized functional network topography to psychopathology in youth". In: *Biological psychiatry* 92.12 (2022), pp. 973–983.

[63] Daniel Mamah et al. "Size and topography of the brain's functional networks with psychotic experiences, schizophrenia, and bipolar disorder". In: *Biological Psychiatry Global Open Science* 4.6 (2024), p. 100386.

[64] Hongming Li et al. "Uncovering functional connectivity patterns predictive of cognition in youth using interpretable predictive modeling". In: *Proceedings of the National Academy of Sciences* 122.42 (2025), e2505600122.