Constrained Diffusion as a Paradigm for Evolution

Daniel Lazarev[a,b,c,d,e], Anna Sappington[b,f,g], Grant Chau[h], Ruochi Zhang[b,c], and
Bonnie Berger[*a,b,c,f]

[a]Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, USA
[b]Computer Science and Artificial Intelligence Laboratory, Massachusetts Institute of
Technology, Cambridge, MA, USA
[c]Broad Institute of MIT and Harvard, Cambridge, MA, USA
[d]Analytic and Translational Genetics Unit, Massachusetts General Hospital, Cambridge,
MA, USA
[e]Renaissance School of Medicine at Stony Brook University, Stony Brook, NY, USA
[f]Harvard Medical School, Boston, MA
[g]Department of Electrical Engineering and Computer Science, Massachusetts Institute of
Technology, Cambridge, MA, USA
[h]Department of Computer Science, Dartmouth College, Hanover, NH, USA

Abstract

One of the most basic questions in computational biology is to describe the mechanisms driving evolutionary dynamics. Here, we newly view evolution through the lens of a diffusion process that is constrained by the physical, biological, and environmental viability of organisms at a given time. Building on analogies from statistical mechanics and information theory, we introduce `DiffEvol`, a framework that models evolution as constrained diffusion over a discrete genotype space, in contrast to classical diffusive systems where all states are accessible. Using frequency and sequence data alone, `DiffEvol` estimates these constraints by inverting the diffusion dynamics to recover the constrained subspace representing the viable genotype manifold, as well as its evolution over time. Applied to SARS-CoV-2 sequence data spanning 2020–2024, `DiffEvol` reconstructs constraint functions that recapitulate known viral fitness trends, including a pronounced "phase transition" that occurred following the SARS-CoV-2 vaccine rollout. Using the constraint subspace representation of the data characterizes such features and trends more clearly. This framework can be used to not only improve forecasting of emergent strains, but also to produce more accurate reverse time analyses of the evolutionary dynamics to identify ancestral variants and the forces that shaped the modern virus's evolutionary trajectory. More generally, this formulation provides an interpretable mathematical framework linking stochastic mutation processes to evolving fitness landscapes, unifying statistical, mechanistic, and information-theoretic views of evolution. Thus, constrained diffusion offers a mathematical language for evolutionary dynamics in any system where random variation interacts with slowly-changing structural or global constraints, and can be applied to more complex evolutionary phenomena such as vaccine resistance, viral escape, and protein evolution.

---

[*]To whom correspondence should be addressed: `bab@mit.edu`

1 Introduction

More than describing the "fact of evolution," as Bertrand Russell would later call it [18], Darwin's most important contribution was arguably advancing a simple but compelling mechanism driving this evolution. Nevertheless, *On the Origin of Species* contains no substantial mathematical expressions, spurring many attempts to precisely describe how random genetic mutations differentially influence the expression of biological traits, and consequently, the natural selection of organisms that possess more favorable traits.

In the wake of the COVID-19 pandemic and the modern deluge of large-scale genomic and epidemiological data, this question has simultaneously taken on new urgency while being afforded the resources needed to address it. With respect to viral evolution, recent computational approaches have sought to learn genotype–phenotype maps and evolutionary dynamics directly from data using statistical and machine-learning models [15, 8], notably introducing protein and viral language models [11, 12]. Although such black-box methods can achieve impressive predictive performance, they do not generally illuminate the mechanisms or the forces shaping evolution, which can limit their biological interpretability and generalizability to contexts that differ substantially from their training distribution.

Here, we take a complementary, mechanistic approach: we use the simple mathematical language of diffusion, or heat flow, to describe evolution. In the classical kinetic theory of gases in $\mathbb{R}^3$, the kinetic energy distribution of particles evolves via random collisions. By analogy, we consider the space of genetic, or genomic, sequences of length L, $\mathcal{G}_L = \{A, T, C, G\}^L$, and view the evolution of the frequency distribution of points in this space (each representing a possible organism with that genetic sequence) as driven by random mutations. A key distinction, however, is that although all points in $\mathbb{R}^3$ are accessible to gas particles, not all points in $\mathcal{G}_L$ are biologically viable: Most possible genetic sequences correspond to nonfunctional or nonviable organisms and thus remain unoccupied. Unlike gas diffusion, evolution thus is not an unconstrained diffusion but one restricted to a dynamically evolving subspace of $\mathcal{G}_L$ that consists of viable or favored sequences. Hard constraints delimit unviable regions of $\mathcal{G}_L$, while softer, continuously evolving constraints—reflecting fitness, epidemiological, or environmental factors—shape the directions in which diffusion occurs. These constraints themselves can shift over time, as with the introduction of vaccines or changes in population immunity, producing evolutionary trajectories that may never reach steady state (the diffusion may not settle to a uniform distribution on the constrained subspace).

This diffusion-based formulation builds upon a broad theoretical lineage connecting stochastic evolutionary processes to physical and mathematical frameworks. Studies of adaptive dynamics describe how discrete mutations converge to a continuous diffusion process in the small-mutation limit [3]. Diffusion maps and related spectral methods [4] have shown how geometric structure and low-dimensional manifolds can emerge from local random-walk dynamics, while diffusion-based graph learning approaches [10] provide practical means of capturing global constraints from local transition information. Similarly, recent methods for Bayesian gene constraint estimation [26] infer selective limits with a statistical model using data such as mutational burden and depletion patterns, a conceptual analog of the constraint function estimated here. At a broader physical level, the evolution of such constrained systems parallels gradient-flow formulations of free-energy minimization, as in the Jordan–Kinderlehrer–Otto framework [13, 14], and thermodynamic perspectives on replication and selection [9].

Based on our constrained diffusion model, we newly propose `DiffEvol`, which uses observed sequence frequencies f_t over time to invert the diffusion process and estimate the evolving constraint

function, $k(t)$, which encodes the feasible geometry of the genotype subspace and its evolution. The resulting model offers a unified, interpretable description of evolutionary dynamics that deconvolves the diffusion aspect of evolution, driven by random mutation, from the dynamic constraints, which encode fitness, functional, and epidemiological structure. Using four years of SARS-CoV-2 sequence data from GISAID, we demonstrate how `DiffEvol` recovers evolving fitness constraints and reveals phase transitions in viral adaptation, with the potential for both improved predictive forecasting of emergent strains—for example, by leveraging the constraint information into predictive frameworks —and reverse-time reconstruction of ancestral variants.

This conceptual shift connects naturally to a wide range of empirical and theoretical work on evolutionary dynamics. Network representations of genotype spaces have demonstrated that their connectivity profoundly affects molecular evolution and extinction [1, 23], while network-propagation models have highlighted how local perturbations in mutational landscapes can amplify through gene and protein networks [7]. High-resolution studies of mutation rates and constraint landscapes across the human genome and tumors [19], as well as information-theoretic analyses of large biological datasets [22], underscore the growing role of quantitative constraint modeling in understanding evolution. In the viral context, recent work on SARS-CoV-2 and other pathogens has leveraged such ideas to predict future variants [8], map mutational drivers [11], and forecast viral escape from immune or therapeutic pressures [20].

`DiffEvol` is complementary to existing approaches for modeling evolutionary dynamics. Classical phylogenetic and phylodynamic models describe lineage relationships and population-level parameters over time [15], while fitness-landscape inference methods estimate selection coefficients from mutational patterns [26]. More recently, protein and viral language models have demonstrated impressive predictive power by learning high-dimensional genotype–phenotype mappings directly from sequence data [11, 12, 20]. In contrast, `DiffEvol` focuses on recovering a mechanistically interpretable, time-resolved constraint geometry by inverting mutation-driven diffusion. Its goal is not to replace predictive models, but to provide an invertible dynamical framework that deconvolves mutation from evolving structural constraints and offers a mathematical language for understanding the forces shaping evolutionary trajectories.

While we demonstrate `DiffEvol` using SARS-CoV-2 as a densely sampled, real-world case study with well-characterized selective shifts, the framework is not specific to this pathogen. SARS-CoV-2 exhibits substantial adaptive restructuring under immune and vaccination pressures [8, 2, 21], making it a suitable system for illustrating constraint dynamics. However, the formulation applies more generally to rapidly evolving RNA viruses such as influenza or HIV, as well as to protein evolution or other adaptive systems.

More broadly, constrained diffusion provides a unifying language for evolutionary dynamics in which random variation interacts with slowly changing structural constraints, offering a principled and interpretable mathematical paradigm that can improve evolutionary inference and the performance of downstream predictive models.

Put differently, constrained diffusion provides not merely a modeling technique but a structural decomposition of evolutionary dynamics into mutation-driven diffusion and constraint-driven geometric selection.

Figure 1: Schematic showing diffusion of a ball of gas for three different setups. **(A)** unconstrained diffusion; **(B)** constrained diffusion; **(C)**: constrained diffusion with evolving constraints.

2   Methods

2.1   Overview of `DiffEvol`

For a sequence of bases of length L, the *genotype space* is the discrete set,

$$\mathcal{G} = \mathcal{G}_L = \{\text{A}, \text{T}, \text{G}, \text{C}\}^L$$

representing all possible sequences of that length (we will omit the subscript L when unambiguous). (We replace U with T for RNA viruses.) We define the constrained genotype subspace, $\mathcal{G}_c \subset \mathcal{G}$, corresponding to the feasible, likely, or biologically realizable sequences, given the various internal (*e.g.*, biochemical or structural) and external (*e.g.*, environmental or epidemiological) constraints on the genotype space. These constraints are represented by a *constraint function*

$$k : \mathcal{G} \to [0, 1] \,,$$

which assigns to each $g \in \mathcal{G}$ a value indicating the degree to which it belongs to the feasible subset [24, 17]. Intuitively, k encodes the equilibrium or limiting distribution of genotypes for the current environment, while the entropy of k quantifies the effective (logarithmic) volume of $\mathcal{G}_c$ [5, 6]. We can alternatively view $\mathcal{G}_c$ as the support of the constraint function.

We posit that evolution can be viewed as a diffusion process restricted to $\mathcal{G}_c$. Random mutations drive the diffusive motion across genotype space, but the geometry and topology of $\mathcal{G}_c$—as determined by the constraint function—govern the accessible directions and the long-term behavior of this diffusion. These geometric features, in turn, emerge from the combined effects of molecular fitness landscapes, interactions with phenotypes and environments, population-level epidemiological pressures, and potential self-interactions among genes.

Because biological and environmental conditions change, the constraint function itself evolves over time. Let $k(t)_{t \in [0,T]}$ denote the family of time-indexed constraint functions, each defining the geometry of the feasible subspace $\mathcal{G}_c(t)$ at time t. Given empirical frequency data that allow

inference of $k(t)$ for $t \in [0,T]$, one can estimate the most likely constraint function at a future time. The resulting evolutionary process is therefore a diffusion on a dynamic manifold $\mathcal{G}_c(t)$, permitting both forward-time prediction of emergent variants and reverse-time reconstruction of ancestral ones.

The current model, described in the next section, for the above theoretical paradigm assumes (1) a constant mutation rate equal to the mean per-site rate, (2) random and independent mutations, and (3) fixed sequence length L across variants. In practice, these assumptions can be relaxed—for example, by introducing a blank or gap character as a fifth nucleotide state or by replacing the Hamming distance with the Levenshtein distance to account for insertions and deletions.

We emphasize as well that assuming a constant mean per-site mutation rate is not essential to our framework, but is included in order to simplify applications with real data, where per-site mutation rates would often not be available. Significantly, this assumption does not impose a molecular clock at the level of observable dynamics. All non-clock-like behaviors—including accelerations, slowdowns, and punctuated evolutionary shifts—are captured by the time-varying constraint function $k(t)$ inferred from data. In this way, `DiffEvol` explicitly decouples stochastic mutation from evolving selection and environmental structure. As we explicitly note in the following section, the framework naturally generalizes to time- or site-dependent mutation rates. The constant-rate assumption is therefore a modeling convenience rather than a biological restriction.

2.2 Constrained Diffusion

Let m_{ij} be the mutation rate between g_i and g_j, where $g_i, g_j \in \mathcal{G}$, and $d(\cdot,\cdot)$ the Hamming distance. We let $\mathbf{f}_n \in \mathbb{R}^G$ be the normalized frequency vector at discrete time $n \in [0,T]$, where $G = |\mathcal{G}|$. Although $G = 4^L$ is astronomically large for realistic viral genomes, the set of biologically and epidemiologically viable sequences is vastly smaller [8], and evolution is effectively confined to this much lower-dimensional region of genotype space. Moreover, we can focus on the segment of the viral genome encoding the virulent, most quickly evolving component of the virus (*e.g.*, the spike protein for SARS-CoV-2).

Discrete-time mutation dynamics. We define the mutation adjacency matrix M, with entries given by,

$$M_{ij} = (1 - m_{ij})^{L - h_{ij}} m_{ij}^{h_{ij}}, \qquad h_{ij} = d(g_i, g_j), \tag{1}$$

which is symmetric because $h_{ij} = h_{ji}$. Assuming a constant per-site mutation rate m allows us to rewrite Eq. (1) as

$$M_{ij} = (1 - m)^{L - h_{ij}} m^{h_{ij}}. \tag{2}$$

Evidently, the mathematical derivations below and in the Appendix are entirely equivalent regardless of whether we use Eqs. (1) or (2). Its degree matrix $D = \text{diag}(M\mathbf{1}_G)$, where $\mathbf{1}_G$ is the column vector of ones of size G, yields the row-stochastic transition matrix,

$$W = D^{-1}M.$$

The *unconstrained* discrete diffusion equation, or random walk with transition matrix, W, is given by the difference equation,

$$\mathbf{f}_{n+1} - \mathbf{f}_n = (W - I)\mathbf{f}_n. \tag{3}$$

Its solution is given by,

$$\mathbf{f}_n = \mathbf{f}_0 W^n, \tag{4}$$

which diffuses mass over genotype space at a rate determined by m and converges to the uniform distribution. We note that in the limit of the step size going to zero, Eq. (3) becomes the standard heat equation (see, *e.g.*, [16]),

$$\partial_t \mathbf{f} = \Delta \mathbf{f}. \tag{5}$$

Continuous-time embedding and the heat kernel. The discrete-time semigroup $\{W^n\}_{n=1}^T$ admits a natural continuous-time extension. If we view each replication cycle as a time step Δt, then

$$W = I + \Delta t \, \mathcal{L},$$

for some generator $\mathcal{L}$. With $t = n\Delta t$, we take $\Delta t \to 0$, or equivalently, $n \to \infty$, which gives,

$$W^n \longrightarrow e^{t\mathcal{L}},$$

where $\{e^{t\mathcal{L}}\}_{t\in[0,T]}$ is the continuous-time mutation-diffusion semigroup. Thus, both W^n and $e^{t\mathcal{L}}$ represent the same mutation dynamics under either discrete or continuous time, respectively.

The singular value decomposition (SVD) of $W = USU^\top$ gives,

$$e^{tW} = U e^{tS} U^\top, \tag{6}$$

mirroring classical diffusion maps and heat-kernel methods. This provides a continuous-time, smoothed mutation operator that remains stable for large t and noisy data.

Time-dependent constraints. Evolution does not explore all of $\mathcal{G}$ but instead remains confined to a dynamically evolving constrained subspace,

$$\mathcal{G}_c(t) = \{g_i : k_i(t) > 0\},$$

encoded by a constraint function

$$\mathbf{k}(t) = (k_1(t), \ldots, k_G(t)), \qquad 0 \le k_i(t) \le 1.$$

Hard constraints correspond to permanently nonviable sequences ($k_i(t) = 0$), while soft constraints vary slowly with biological, epidemiological, or environmental pressures [17, 24, 25]. The constraint function is often considered to evolve on a slower timescale than mutation, which allow mutation and constraint dynamics to be separated.

Discrete constrained diffusion. Under constraints, the observed distribution at time n is modeled as

$$\mathbf{f}_n = \frac{\mathbf{k}(n) \circ (\mathbf{f}_0 W^n)}{\sum_i k_i(n) \, [\mathbf{f}_0 W^n]_i} = \frac{\mathbf{k}(n) \circ (\mathbf{f}_0 W^n)}{\mathbf{1}_G^\top \, \mathbf{k}(n) \circ (\mathbf{f}_0 W^n)}. \tag{7}$$

Here $\mathbf{f}_0 W^n$ represents diffusion under mutation alone, and the element-wise product with $\mathbf{k}(n)$ restricts this distribution to viable genotypes before renormalization. If $\mathbf{k}(n)$ is constant over time, the sequence converges to a stationary distribution proportional to $\mathbf{k}$. If $\mathbf{k}(n)$ evolves, diffusion occurs on a time-varying, slowly drifting constraint manifold $\mathcal{G}_c(n)$.

Continuous-time constrained diffusion. Using the continuous semigroup $e^{t\mathcal{L}}$, the analogous continuous-time model is

$$\mathbf{f}(t) = \frac{\mathbf{k}(t) \circ (\mathbf{f}_0 e^{t\mathcal{L}})}{\sum_i k_i(t) \, [\mathbf{f}_0 e^{t\mathcal{L}}]_i}. \tag{8}$$

As in discrete time, $\mathbf{k}(t)$ selects the viable region of genotype space, and the heat-kernel operator $e^{t\mathcal{L}}$ propagates mutation continuously. The two formulations are interchangeable under the standard discretization $t = n\Delta t$, with the continuous formulation used for its smoothing effects and favorable properties in the numerical implementation of the method.

Instead of solving the usual heat equation, Eq. (5), Eq. (8) is the solution to the *constrained diffusion equation*,

$$\partial_t \mathbf{f}(t) = \mathbf{f} \circ (\mathcal{M} - \overline{\mathcal{M}} \, \mathbf{1}_G) + \mathbf{f} \circ (\mathcal{S} - \overline{\mathcal{S}} \, \mathbf{1}_G) \tag{9}$$

where $\mathcal{M} = (\mathcal{L} \, \mathbf{f}_0 e^{t\mathcal{L}}) \oslash (\mathbf{f}_0 e^{t\mathcal{L}})$ represents the contribution due to the random mutations, while $\mathcal{S} = \partial_t \mathbf{k}(t) \oslash \mathbf{k}(t)$ represents that due to the evolving constraints, with $\overline{\mathcal{S}} = \mathcal{S}^\top \mathbf{f}$ and $\overline{\mathcal{M}} = \mathcal{M}^\top \mathbf{f}$ their mean values taken with respect to the frequencies. Eq. (9) reduces to Eq. (5) if $\mathbf{k}(t) = \mathbf{1}_G$. (See Appendix for the derivation of Eq. (9).)

2.3 The `DiffEvol` Method

Having formulated mutation as a constrained diffusion on genotype space, we now describe how this framework yields a practical method for learning the time-evolving constraint functions directly from observed frequency data. The key idea is that the constrained diffusion model can be *inverted*: given the observed frequencies $\{\mathbf{f}_n\}$ and the mutation kernel W, we can solve for the constraint function $\mathbf{k}(n)$ at each time point. The resulting time series $\mathbf{k}(0), \ldots, \mathbf{k}(N)$ encodes the geometry of the dynamically evolving feasible subspace $\mathcal{G}_c(n)$.

Direct inversion in discrete time. The discrete constrained diffusion equation, given by Eq. (7), implies the proportionality $\mathbf{k}(n) \propto \mathbf{f}_n \oslash (\mathbf{f}_0 W^n)$, with $\oslash$ denoting elementwise (Hadamard) division. Normalizing yields,

$$\mathbf{k}(n) = \frac{\mathbf{f}_n \oslash (\mathbf{f}_0 W^n)}{\sum_i \left[\mathbf{f}_n \oslash (\mathbf{f}_0 W^n) \right]_i}. \tag{10}$$

Computing Eq. (10) for each $n \in [0, N]$ produces the *constraint matrix* $K \in \mathbb{R}^{N \times G}$, whose nth row describes the feasible region of genotype space at time n.

Continuous-time analogue and the heat kernel. Mutation can also be modeled in continuous time via the semigroup $g(t) = \mathbf{f}_0 e^{tW}$, which is the solution to the continuous time and discrete space diffusion heat equation, $\partial_t g(t) = W g(t)$. The constrained continuous-time equation, given by Eq. (8), implies, exactly as in the discrete case, that $\mathbf{k}(t) \propto \mathbf{f}(t) \oslash (\mathbf{f}_0 e^{tW})$, and thus the continuous-time inverse solution is given by,

$$\mathbf{k}(t) = \frac{\mathbf{f}(t) \oslash (\mathbf{f}_0 e^{tW})}{\sum_i \left[\mathbf{f}(t) \oslash (\mathbf{f}_0 e^{tW}) \right]_i}. \tag{11}$$

In practice, continuous-time inversion is computed spectrally, using the SVD formulation in Eq. (6). This gives a smoothed, continuous-time mutation operator that is well-conditioned for large t.

Figure 2: (A) The ground truth constraint matrix, K_{true}; (B) the frequency matrix, F_n, with added noise; and (C) the estimated constraint matrix, K_{est}, using `DiffEvol`.

Estimating the temporal dynamics of constraints. The sequence $\{\mathbf{k}(n)\}_{n=1}^{T}$ typically evolves on a slower timescale than mutation. We can therefore treat the rows of K as a time series that lies near a low-dimensional submanifold in $\mathbb{R}^G$. This makes possible:

- **Forward prediction:** estimating future constraint vectors $\mathbf{k}(T{+}1), \mathbf{k}(T{+}2), \ldots$ and thereby future prevalence distributions via Eqs. (7) or (8).

- **Backward reconstruction:** inferring likely ancestral constraint states and ancestral prevalence distributions, provided the explored region of W is invertible.

- **Quantifying selective shifts:** identifying abrupt or gradual changes in $\mathbf{k}(n)$ corresponding to selective sweeps, immune or therapeutic pressures, or major epidemiological transitions.

Summary. The `DiffEvol` method is thus a principled inversion of the constrained diffusion process. It decouples mutation from constraint, reconstructs a time-resolved viability landscape from empirical frequencies, and provides a coherent forward–backward dynamical model for predicting future variants and inferring ancestral ones. In this sense, $\mathbf{k}(n)$ is not merely a fitted parameter but a mechanistically meaningful description of the evolving constraint geometry that steers viral evolution.

3 Results

3.1 Toy model

To test the performance of our method and some of the assumptions that underlie it, we built a toy model. Let K_{true} be the ground truth constraint matrix, F_n the frequency matrix with added noise and K_{est} the estimated constraint matrix using `DiffEvol`. We added noise drawn from a normal distribution $\mathcal{N}(0, 10^{-5})$, where the standard deviation was chosen so that it is approximately two orders of magnitude smaller than the mean value of the frequency matrix entries.

To better simulate the real data, and to test whether we are justified in using the mean mutation rate in the transition matrix used to estimate the constraint matrix (Methods), we also used a variable mutation rate in the ground truth transition matrix used to generate the simulated frequencies. We generated a vector of G mutation rates by taking the absolute value of numbers drawn from $\mathcal{N}(m, m)$. We used $m = 10^{-5}$ for the simulations shown in plots in Fig. 2 and Table 1.

To assess the performance of our method on simulated data, we use the weighted absolute percent error (WAPE), given by,

$$WAPE(X_{ij}, \hat{X}_{ij}) = \frac{\sum_{i,j} |X_{ij} - \hat{X}_{ij}|}{\sum_{i,j} |X_{ij}|},$$

where X_{ij} is the real data and $\hat{X}_{ij}$ is the predicted data.

Table 1: Performance of `DiffEvol` in reconstructing the constraint matrix from the frequency matrix with added noise. WAPE was measured relative to K_{true} and U is the uniform matrix with entries given by $1/G$ for which the entropy attains its maximum. Used variable mutation rates in generating the frequency matrix.

Matrix	Entropy	WAPE
K_{true}	10.389	——
F_n	10.230	50.02%
K_{est}	10.437	2.46%
U	11.556	130.61%

Note that the WAPE for U in Table 1 is greater than 100% because of the large number of entries for which $U_{ij} > K_{\text{true}}$.

3.2 SARS-CoV-2 model

Data acquisition and preprocessing. SARS-CoV-2 genomic sequences were obtained from the GISAID database spanning January 1, 2020 to May 29, 2023. FASTA files were parsed to extract sequence data, collection dates, and geographic locations. For each record, we extract the collection date, location, and full nucleotide sequence. Unique sequences were identified and assigned sequential indices, and this resulted in a map between the original 47,550 filtered sequences and a compressed set of 3,262 unique variants. This alignment step guarantees that sequences are matched with their corresponding temporal frequency profiles.

Occurrence matrix. Temporal occurrence patterns were encoded in a sparse matrix where rows represent time points (823 unique dates) and columns represent sequence variants. Sequences were filtered based on observational coverage, retaining variants with at least 10 total occurrences across the study period. This removed rare or potentially artifactual sequences while preserving biologically relevant variants.

Frequency normalization. Raw occurrence counts were converted to normalized frequencies through a two-stage process. Firstly, we normalize counts by time point (column-wise) and compute the relative frequencies that sum to 1 at each time point. Next, frequencies were normalized by variant (row-wise) to standardize the temporal profile of each variant across time. This was to ensure that variants with different absolute abundances could be compared on a similar scale.

To denoise the raw frequency data and extract underlying evolutionary constraints, we apply `DiffEvol`, which models evolution as constrained diffusion over the discrete genotype space defined by our dataset. The constraint matrix K consists of $T = 823$ time points and $G = 3,262$

Figure 3: (A) UMAP plot and (B) heat map based on frequency data of the SARS-CoV-2 virus nucleotide genotypes over three years. (C) Corresponding constraint matrix using `DiffEvol`. Variants are ordered by their mean emergence time.

genotypes. For comparison, we report the following metrics for both the raw frequency matrix and the constraint matrix K:

1. **Mean Evolution.** Temporal trajectories of mean constraint or frequency values (averaged across all genotypes at each time point). This depicts overall population-level dynamics and responses to selective pressures.

2. **Entropy.** Shannon entropy of the distribution at each time point, quantifying genotypic diversity. The effective number of genotypes was computed as the inverse Simpson index: $N_{\text{eff}} = \frac{1}{\sum_i p_i^2}$, where p_i represents normalized frequencies or constraints.

3. **Diversity.** Number of dominant variants at a given time.

4. **Rate of Change.** First-order temporal derivatives ($\frac{dK}{dt}$ and $\frac{d\text{Gini}}{dt}$) to identify periods of rapid evolutionary change and potential phase transitions.

5. **Gini Coefficient.** Measures concentration of the frequency or constraint distribution, with values near 1 indicating dominance (high concentration) by few variants and values near 0 indicating uniform distribution (random chance). Temporal changes in the Gini coefficient ($\frac{d\text{Gini}}{dt}$) depict shifts in competitive dynamics.

6. **Variance.** Temporal variance across genotypes, indicating the spread of the distribution and the stability of the constraint landscape.

7. **Top** $n(= 15)$ **sequence constraints.** Depicts which n variants have the highest mean constraint (best "fitness" on average).

To examine the impact of vaccination on viral evolution, we identified the vaccine rollout period (December 15, 2020 to March 15, 2021) and compared pre-vaccine, vaccine rollout, and post-vaccine periods. Statistical distributions of constraints before and after this period were compared to quantify the shifts in the evolutionary landscape. Comparison of 4 shows that `DiffEvol` is effective at extracting evolutionary signals from noisy observational data. Notably, the vaccine rollout period marked a clear evolutionary regime shift and directional selection where previously successful variants were systematically eliminated (F) and the selective sweep in the middle of the vaccination period converged rapidly to low diversity (E) with the dominance of a single variant. This pattern is consistent with strong immune selection following vaccination that overwhelms other evolutionary pressures (transmission fitness, geographic effects, stochastic drift).

Finally, we also applied `DiffEvol` to Inflluenza A virus (Subtype H1N1). The corresponding plots depicting the frequency and constraint matrices can be found in Appendix.

4   Discussion

We introduced `DiffEvol`, a framework that models evolution as a diffusion process constrained by time-varying biological and environmental constraint functions. By inverting the diffusion dynamics on genotype space, `DiffEvol` infers the evolving constraint functions that reflect the various forces driving evolutionary change. Applied to SARS-CoV-2, it reconstructs constraint shifts corresponding to major fitness transitions, including those following vaccine rollout.

In future work, we hope to explicitly analyze both forward-time forecasting of variants and reverse-time reconstruction of ancestral states using `DiffEvol`'s constraint functions. We also aim to enable

Figure 4: **Comparison of raw frequency dynamics and `DiffEvol`-derived constrained diffusion for SARS-CoV-2 (2020–2023).**

Figure 4: `DiffEvol` denoises raw frequency data (blue) into interpretable constraint dynamics (red), revealing punctuated increases in evolutionary constraint around the vaccine rollout (gray). (A–E) show smoother, biologically meaningful trends in constraint intensity, diversity, rate of change, variance, and selection concentration compared to raw data. (F i–ii) illustrate that `DiffEvol` compresses thousands of sequences into a small set of coherent evolutionary trajectories capturing dominant adaptive modes.

the incorporation of multiple data modalities and multiscale data (e.g., cell-level, protein, and gene expression data) to further inform the model.

Taken together, our formulation unifies stochastic mutation and structured selection within a single diffusion–constraint framework, providing an interpretable alternative to black-box predictive models. It offers a general mathematical language for adaptive processes where random variation interacts with evolving structural constraints, from viral evolution to molecular and cellular systems.

Acknowledgments

This work was partially funded by NIH grant R35GM141861 (to BB).

References

[1] Aguirre, J., Catalán, P., Cuesta, J., Manrubia, S.: On the networked architecture of genotype spaces and its critical effects on molecular evolution. Open Biol. **8**, 180069 (2018)

[2] Bradley, C.C., Wang, C., Gordon, A.J.E., Wen, A.X., Luna, P.N., Cooke, M.B., Kohrn, B.F., Kennedy, S.R., Avadhanula, V., Piedra, P.A., Lichtarge, O., Shaw, C.A., Ronca, S.E., Herman, C.: Targeted accurate rna consensus sequencing (tarc-seq) reveals mechanisms of replication error affecting sars-cov-2 divergence. Nature Microbiology **9**(5), 1382–1392 (2024)

[3] Champagnat, N., Lambert, A.: Evolution of discrete populations and the canonical diffusion of adaptive dynamics. The Annals of Applied Probability **17**(1), 102 – 155 (2007)

[4] Coifman, R.R., Lafon, S., Lee, A.B., Maggioni, M., Nadler, B., Warner, F., Zucker, S.W.: Geometric diffusions as a tool for harmonic analysis and structure definition of data: Diffusion maps. Proceedings of the National Academy of Sciences **102**(21), 7426–7431. https://doi.org/10.1073/pnas.0500334102, `https://www.pnas.org/doi/abs/10.1073/pnas.0500334102`

[5] Coifman, R.R., Wickerhauser, M.V.: Entropy-based algorithms for best basis selection. IEEE Transactions on Information Theory **38**(2), 713–718 (1992)

[6] Cover, T., Thomas, J.: Elements of Information Theory. John Wiley (1991)

[7] Cowen, L., Ideker, T., Raphael, B.J., Sharan, R.: Network propagation: a universal amplifier of genetic associations. Nature Reviews Genetics **18**(9), 551–562 (2017)

[8] Cyrus Maher, M., Bartha, I., Weaver, S., di Iulio, J., Ferri, E., Soriaga, L., Lempp, F.A., Hie, B.L., Bryson, B., Berger, B., Robertson, D.L., Snell, G., Corti, D., Virgin, H.W., Pond, S.L.K.,

Telenti, A.: Predicting the mutational drivers of future SARS-CoV-2 variants of concern. Science Translational Medicine **14**(633), eabk3445 (2022)

[9] England, J.L.: Statistical physics of self-replication. J. Chem. Phys. **139**, 121923 (2013)

[10] Gasteiger, J., Weißenberger, S., Günnemann, S.: Diffusion Improves Graph Learning. In: Wallach, H., Larochelle, H., Beygelzimer, A., d'Alché-Buc, F., Fox, E., Garnett, R. (eds.) Advances in Neural Information Processing Systems. vol. 32. Curran Associates, Inc. (2019)

[11] Hie, B., Zhong, E.D., Berger, B., Bryson, B.: Learning the language of viral evolution and escape. Science **371**(6526), 284–288 (2021). https://doi.org/10.1126/science.abd7331

[12] Hie, B.L., Yang, K.K., Kim, P.S.: Evolutionary velocity with protein language models predicts evolutionary dynamics of diverse proteins. Cell Systems **13**(4), 274–285.e6 (2022)

[13] Jordan, R., Kinderlehrer, D., Otto, F.: Free energy and the Fokker-Planck equation. Physica D **107**, 265–271 (1997)

[14] Jordan, R., Kinderlehrer, D., Otto, F.: Variational formulation of the Fokker-Planck equation. SIAM J. Math Anal. **29**(1), 1–17 (1998)

[15] Koelle, K., Cobey, S., Grenfell, B., Pascual, M.: Epochal Evolution Shapes the Phylodynamics of Interpandemic Influenza A (H3N2) in Humans. Science **314**(5807), 1898–1903 (2006). https://doi.org/10.1126/science.1132745, `https://www.science.org/doi/abs/10.1126/science.1132745`

[16] Lawler, G.F.: Random Walk and the Heat Equation. American Mathematical Society, Providence, Rhode Island (2010)

[17] Lazarev, D.: Information measures for entropy and symmetry. arXiv:2211.14857 (2023)

[18] Russell, B.: Religion and Science. Oxford University Press, Oxford, second edn. (1997)

[19] Sherman, M.A., Yaari, A.U., Priebe, O., Dietlein, F., Loh, P.R., Berger, B.: Genome-wide mapping of somatic mutation rates uncovers drivers of cancer. Nature Biotechnology **40**(11), 1634–1643 (2022)

[20] Thadani, N.N., Gurev, S., Notin, P., Youssef, N., Rollins, N.J., Ritter, D., Sander, C., Gal, Y., Marks, D.S.: Learning from prepandemic data to forecast viral escape. Nature **622**(7984), 818–825 (2023)

[21] Willett, J.D.S., Gravel, A., Dubuc, I., Gudimard, L., dos Santos Pereira Andrade, A.C., Lacasse, É., Fortin, P., Liu, J.L., Cervantes, J.A., Galvez, J.H., Djambazian, H.H.V., Zwaig, M., Roy, A.M., Lee, S., Chen, S.H., Ragoussis, J., Flamand, L.: Sars-cov-2 rapidly evolves lineage-specific phenotypic differences when passaged repeatedly in immune-naïve mice. Communications Biology **7**(1), 191 (2024)

[22] Yu, Y.W., Daniels, N.M., Danko, D.C., Berger, B.: Entropy-scaling search of massive biological data. Cell Systems **1**(2), 130–140 (2023/06/11 2015). https://doi.org/10.1016/j.cels.2015.08.004, `https://doi.org/10.1016/j.cels.2015.08.004`

[23] Yubero, P., Manrubia, S., Aguirre, J.: The space of genotypes is a network of networks: implications for evolutionary and extinction dynamics. Sci. Rep. **7**, 13813 (2017)

[24] Zadeh, L.A.: Fuzzy sets. Inf. Control **8**, 338–353 (1965)

[25] Zadeh, L.A.: Probability measures of fuzzy events. J. Math. Anal. Appl. **23**, 421–427 (1968)

[26] Zeng, T., Spence, J.P., Mostafavi, H., Pritchard, J.K.: Bayesian estimation of gene constraint from an evolutionary model with gene features. Nature Genetics **56**(8), 1632–1643 (2024)