The Brain Age Gap as a Predictor of Alcohol Initiation in Adolescence

Byrne, H., Ph.D.[1], Visontay, R., Ph.D.[1], Devine, E. K. Ph.D.[1], Wade, N. E., Ph.D.[2], Jacobus,

J., Ph.D.[2], Moore, A. J.[3], Ph.D., Squeglia, L. M., Ph.D.[3]*, Mewton, L., Ph.D.[1]*

*Joint senior authors

[1] The Matilda Centre for Research in Mental Health and Substance Use, The University of

Sydney, Sydney, NSW, Australia

[2] Department of Psychiatry, University of California San Diego, La Jolla, CA, United States

[3] Department of Psychiatry and Behavioral Sciences, Medical University of South Carolina,

Charleston, SC, United States

Corresponding author: Hollie Byrne

The Matilda Centre for Research in Mental Health and Substance Use, The University of

Sydney, Sydney, NSW, Australia

Abstract

Background: Growing evidence suggests regional and network-level brain imaging features

in late childhood are predictive of alcohol use in adolescence. However, the directionality of

these effects (i.e. whether they reflect accelerated or delayed neuromaturation) are mixed. We

applied a Brain Age Gap Estimation (BrainAGE) model to examine whether overall

deviations from typical brain aging trajectories are predictive of (1) alcohol initiation and (2)

use behaviour (experimentation versus binge drinking) in adolescence.

Methods: Data from the Adolescent Brain Cognitive Development study release 6.0 were

used. Baseline (ages 9-11) structural imaging features (cortical volume, area, and subcortical

volume) were used to estimate BrainAGE. Alcohol use was determined using self-report data

from the Substance Use Interview and Timeline Follow-Back across follow-ups (waves 1-6;

ages 10-17). Logistic generalized mixed effects models examined whether BrainAGE

predicted group status between (1) non-initiators (n=3,639) and initiators (n=1,176), and; (2)

experimentation (at least one full drink, no binge episodes; n=461) and binge drinking (at

least one episode; n=438).

Results: When adjusting for age, sex, and pubertal status, a one-standard-deviation decrease

in BrainAGE (equivalent to 1.64 years) at baseline was associated with a 9.5% increase in

odds of alcohol initiation in adolescence. However, this effect did not survive adjustment for

sociodemographic and prior alcohol exposure covariates. Further, BrainAGE did not

discriminate between experimentation and binge drinking.

Conclusions: Findings suggest BrainAGE in late childhood may reflect potential risk for

alcohol initiation, but not behaviours, in adolescence. However, this association likely reflects

complex interactions between brain structure and contextual factors, warranting further

investigation.

Introduction

Alcohol use in adolescence poses a significant public health concern. Recent data from the World Health Organization shows 57% of adolescents reporting alcohol use before age 16, and 20% reporting having experienced significant drunkenness (more than one binge drinking episode) by this age (Lorena Charrier et al., 2024). Earlier alcohol initiation is associated with increased risk of adverse outcomes across the lifespan, including subsequent heavy drinking and substance use disorders (Gardner et al., 2024; Lee et al., 2024), poorer academic performance (Hjarnaa et al., 2023), greater risk-taking behaviours (Clare et al., 2026), and increased mental health and social difficulties (Tackaberry-Giddens et al., 2023). Because adolescence is a sensitive period for neurodevelopment, alcohol exposure during this time can also disrupt maturation of neural systems involved in executive function, reward processing, and emotion regulation (Spear, 2018), making adolescent alcohol use particularly consequential. As a result, clarifying the risk factors and mechanisms underlying alcohol initiation is critical for informing prevention efforts and early intervention during this age period.

While the impacts of alcohol use on the brain have been well documented, evidence increasingly suggests distinct neurobiological characteristics may predate alcohol use. Specifically, research suggests aberrant development of subcortical and frontal systems prospectively predicts early initiation and escalation of alcohol use in adolescence (Lees et al., 2021), aligning with hypotheses suggesting neurobiological correlates of addiction may reflect both causes and consequences (Bogdan et al., 2023; Miller et al., 2024). This is supported by recent studies conducted in large, population imaging datasets such as the Adolescent Brain Cognitive Development (ABCD) study that have identified structural morphometric properties at both the regional (Green et al., 2024; Miller et al., 2024) and network (Byrne et al., 2026) level in late childhood (ages 9-11) which are predictive of alcohol use behaviours in early adolescence (up to age 15). However, the directionality of these effects – specifically, whether these observations reflect 'accelerated' or 'delayed' neuromaturation – have yet to be determined. Further, it remains unclear the extent to which alcohol use behaviours, particularly experimentation compared to binge drinking, are differentially related to structural brain properties.

Currently, few analytical methods are available to robustly assess maturation in the developing adolescent brain. One approach is brain age gap estimation (often abbreviated as

'BrainAGE' or 'BAG'), a neuroimaging-based metric that computes the difference between

an individual's predicted brain age – derived from machine learning models applied to brain-

imaging measures – and their chronological age. Notably, while these models do not provide

an accurate prediction of maturation, BrainAGE can offer a general indicative measure of

deviations from typical brain aging trajectories (Whitmore & Beck, 2025). In this context, a

positive gap is interpreted as the brain appearing "older" than expected in comparison to the

normative or control sample, while a negative gap points to the brain appearing "younger".

BrainAGE has been frequently applied within the adolescent and adult alcohol literature to

investigate the impacts of alcohol use on deviations from normative developmental

trajectories, which have typically identified a positive (higher) BrainAGE following chronic

and/or heavy alcohol use (Scheffler et al., 2024). However, the predictive utility of BrainAGE

in the initiation and type of alcohol use behaviour (experimentation vs. binge drinking)

remains unknown.

Using data from the ABCD study, this work aims to investigate whether BrainAGE in late

childhood (baseline; ages 9-11) can predict alcohol initiation in mid adolescence (wave 6;

ages 15-17). Next, this work aims to examine whether baseline BrainAGE can differentiate

between adolescents who have experimented with alcohol (defined here as consumption of a

full drink but no binge drinking) compared to those who report at least one binge drinking

episode.

Methods and Materials

Participants

Data from the ABCD Study release 6.0 were used, which includes behavioral and biological

data from 11,868 participants aged 9-11 years at baseline. Briefly, participants were recruited

from 21 sites across the United States between September 2016 and November 2018, with

follow-up visits conducted at yearly intervals. Parents/caregivers provided signed informed

consent and all participants gave assent. The ABCD protocol was approved by the centralized

institutional review board (IRB) at the University of California, San Diego and by the IRBs at

the 21 sites. Release 6.0 contains data from baseline through to wave 6 follow-up, with

varying numbers of missed visits across follow-ups and data collection still ongoing at this

stage. As data at the final available follow-up (wave 6; ages 15-17) aligns with the critical

window for adolescent alcohol initiation (Swendsen et al., 2012), we included all available

data through to wave 6 (n = 5,056). Participants who had not initiated alcohol use were

retained only if they had data at wave 6, whereas initiators were included if initiation

occurred at any point prior to or at wave 6. To evaluate whether non-initiators had sufficient

time to initiate, we compared their age distribution at wave 6 with the age-at-first-use

distribution among initiators (*Figure S1*). The overlap in these distributions suggests that the

non-drinkers included in these analyses were comparable in age to the alcohol initiation

group, supporting the use of this approach for classification.

For BrainAGE calculation, participants were excluded if they were missing structural MRI

data at baseline (n = 72), did not meet neuroimaging quality-control criteria (n = 472), or

lacked complete covariate information required for harmonization of imaging data across

scanners (n = 841; see *Image Processing*). For group classification, participants were

excluded if they reported consuming a full alcoholic drink at baseline (n = 21); if non-

drinking status within the non-initiator group could not be confirmed at wave 6 (i.e. missing

data/not yet tested at wave 6; n = 5,647; see *Alcohol use classification*); if binge drinking

status within the experimenter subgroup could not be determined at wave 6 (n = 277; see

Alcohol use classification). Of the 11,868 participants recruited at baseline, 4,815 met

inclusion criteria (*Figure 1*; n = 1,176 initiators).

Measures

Image processing

Preprocessed imaging data acquired at baseline (ages 9-11) were used (Hagler et al., 2019).

Cortical parcellation and subcortical segmentation of T1-weighted structural imaging data

were performed using FreeSurfer (v7.1.1; https://surfer.nmr.mgh.harvard.edu/). Postprocessed

cortical volume and area measurements mapped to the Desikan-Killiany atlas included 34

cortical parcellations per hemisphere, for a total of 68 regions of interest (ROIs) (Desikan et

al., 2006), respectively, and 53 bilateral global and subcortical volumetric measures derived

from the FreeSurfer output.

To remove unwanted technical variability while preserving biological signals, neuroimaging

measures were harmonized across scanners using ComBat (v1.0.13; Fortin et al., 2018).

Scanner characteristics were modeled as batch covariates, while age, sex, and pubertal status

(derived from parent-report Pubertal Development Scale scores; see *Pubertal status*) were

preserved as biological covariates of interest.

BrainAGE model

To estimate BrainAGE, we used the Drobinin et al. (2022) model, which has demonstrated

good external validity for use in adolescent datasets (Michael et al., 2026). The Drobinin

model was trained on 1,299 participants aged 9–19 years, drawn from six independent non-

ABCD samples. Age and sex distributions for each sample are reported in Drobinin et al.

(2022). Participants were included in model training if they fell within the 9–19-year age

range, passed automated MRI quality-control procedures, had no psychiatric diagnoses, and

had an IQ greater than 75.

In Drobinin et al. (2022), model training was conducted within the *tidymodels* (version 0.1.1)

framework, using the eXtreme Gradient Boosting (XGBoost) machine learning algorithm

(version 1.0.0.2). Scan age in the Drobinin model was predicted from 189 features, including

cortical gray matter volume and surface area measurements, as well as bilateral global and

subcortical volume measurements. Cortical features were obtained using the FreeSurfer

default Desikan-Killiany atlas, while bilateral global and subcortical volume measures were

obtained using FreeSurfer segmentation output (aseg). The Drobinin model reported a mean

absolute error (MAE; the average absolute difference between predicted and chronological

age, where values of 1-2 reflect good model prediction accuracy in this context [Whitmore &

Beck, 2025]) of 1.53 years in typically developing adolescents; generalized to the validation

dataset (mean absolute error = 1.55 years, 1.98 bias adjusted); and to the independent at-risk

sample (mean absolute error = 1.49 years, 1.86 bias adjusted).

While the Drobinin et al. (2022) model was originally trained on 189 structural features, only

175 of these features were available in the ABCD dataset. Therefore, for the current study, the

model was used to predict BrainAGE based on the 175 available features, replicating the

methods by Whitmore et al. (2023). BrainAGE was estimated on the full harmonized sample

(n = 10,483) prior to group classification.

Bias correction

Consistent with prior work (Le et al., 2018; Smith et al., 2019; Whitmore et al., 2023), we

applied bias correction to account for systematic over- or underestimation related to age.

Specifically, a linear model was fit to the validation data to derive a slope and intercept,

which were then used to adjust predicted ages in the test set. Age was also included as a

covariate in all subsequent statistical analyses to further control for residual age-related bias,

as recommended in other work (Peng et al., 2021; Whitmore & Beck, 2025), and age outliers

(participants aged ≤8.9 or ≥11.1; n = 3) were removed.

Alcohol use classification

Alcohol use

Participants were classified into alcohol use groups using a hierarchical approach based on

longitudinal self-reported data from the Substance Use Interview (SUI) and Timeline Follow-

Back (TLFB) modules. Participants lacking complete SUI or TLFB data as per our exclusion

criteria were excluded.

Primary model classification (non-initiators vs. initiators): 'Non-initiators' (n = 3,639) were

defined as participants who did not report consumption of a full alcoholic drink in any

session prior to and including wave 6. Participants with insufficient data at wave 6 to confirm

non-initiation status at this timepoint were excluded. 'Initiators' (n = 1,176) were defined as

participants who reported consuming at least one full alcoholic drink in any follow-up session

before wave 6. This was assessed using binary 'Yes'/'No' participant responses to the

question: "Have you used since we last saw you: A full drink of beer, wine or liquor (rum,

vodka, gin, whiskey)," as self-reported during the SUI.

Secondary model classification (experimenters vs. binge drinkers): Among participants

classified as 'Initiators', 'Bingers' (n = 438) were defined as participants who reported at

least one binge drinking episode at any timepoint during follow-up. Binge episodes were

defined as ≥4 standard drinks for females or ≥5 for males in a single drinking session, as self-

reported in the TLFB. 'Experimenters' (n = 461) were defined as participants who reported a

full drink of alcohol by wave 6 with no reported binge drinking episodes at any session prior

to wave 6. To ensure Experimenters were appropriately characterized, Experimenters were

required to have data available at wave 6 confirming non-binge drinking status at this

timepoint (n = 277 excluded).

Covariates

Previous research has identified robust associations between age, sex, and pubertal status,

respectively, in BrainAGE estimates (Holm et al., 2023; Whitmore et al., 2023). Therefore,

these variables were included as primary covariates in the study. Chronological age and sex at

baseline were determined from parent-reported date of child's birth and sex assigned at birth,

respectively. Baseline pubertal status was assessed using the parent-report Pubertal

Development Scale (PDS), which evaluates physical markers of pubertal maturation such as

growth of body hair, skin changes, and development of secondary sexual characteristics for

males and females, respectively (Petersen et al., 1988). An overall summary score ranging

from 1 to 5 is calculated to represent each participant's pubertal stage, approximating the

Tanner stages of pubertal development (Tanner & Whitehouse, 1976), with higher scores

indicating more advanced pubertal development. Parent-report scores were used in this study

based on recommendations for use of the PDS in baseline ABCD data (Cheng et al., 2021).

Statistical analysis

Primary analysis

A logistic generalized linear mixed effects model (GLMM) with a binomial distribution and

logit link was used to predict alcohol use status (non-initiators vs. initiators) from baseline

BrainAGE, controlling for baseline age, sex, and pubertal status (fixed effects), and random

intercepts accounting for family nested within data collection site. Numeric predictors

(BrainAGE, age, pubertal status) were z-standardized (mean = 0, SD = 1) for interpretability

and effect size comparability. Models were fitted via maximum likelihood with Laplace

approximation using the *lme4* (v 1.1.37; Bates et al., 2015) package in R (*bobyqa* optimizer).

Effects are reported as odds ratios (ORs) with 95% confidence intervals. Significance was

assessed with two-sided tests at $\alpha = .05$.

Secondary analysis

The primary model was repeated using the same parameters to predict alcohol behaviour

subgroup classification (experimentation vs. binge drinking) from baseline BrainAGE.

Sensitivity analyses

Two sensitivity analyses were performed using the primary model classification (non-

initiators vs. initiators).

Sociodemographics and prior alcohol exposure

To assess whether observed differences in BrainAGE were independent of sociodemographic

differences and/or prior alcohol exposure, the primary analysis was repeated with adjustment

for covariates associated with brain structure and/or early alcohol use in the ABCD sample.

This included key sociodemographic variables (race, ethnicity, highest parental education,

religious attitudes against alcohol), as well as additional variables related to prenatal exposure

and early alcohol use (maternal alcohol use during pregnancy, participant reported sipping at

baseline), as performed in previous work (Byrne et al., 2026). These variables were not

interpreted as biological traits but included to reduce confounding related to structural and

sociocultural factors known to affect neurodevelopmental trajectories and substance use risk.

Further information on the variables used are reported in *Table S4*.

Exclusion of other substance use from non-initiator group

Inclusion of participants in the non-initiating group who initiate other substances during

follow-up may confound results, as initiation of these substances may reflect shared high-risk

neural substrates with alcohol (Miller et al., 2024). Therefore, to assess whether BrainAGE

differs at baseline between 'true' non-initiators and those who later initiated other substances,

the GLMM was repeated excluding non-initiators who initiated cigarette, e-cigarette, or

cannabis smoking at baseline and/or during follow-up, as reported during the SUI (see *Table

S4* for further detail on these measures).

Results

Participant characteristics

At baseline, participants who initiated alcohol use at or by wave 6 (n = 1,176) differed on

nearly all characteristics compared to non-initiators (n = 3,639). Specifically, initiators were

older than non-initiators, and a higher proportion were female, had been exposed to alcohol *in*

utero, and reported alcohol sipping at baseline. In the non-initiating group, a greater

proportion reported religious (negative) attitudes against alcohol compared to alcohol

initiators, and groups differed in distribution of pubertal status, race, ethnicity, and highest

caregiver education. Participant characteristics are reported in *Table 1*.

Figure 1. Flowchart for participant inclusion/exclusion criteria.

To assess whether baseline age differences could bias follow-up comparisons, where alcohol

use might be skewed in the initiator group because of older age at baseline, age distributions

for non-initiators at wave 6 and age at first reported drink in the initiator group were

examined (*Figure S1*). No meaningful skew toward older ages was observed in the initiator

group relative to non-initiators, suggesting that older age in the initiator group was unlikely to

bias the overall findings.

Table 1. Participant characteristics of study sample.

Characteristic	Non-initiators (n = 3,639)	Drinkers (n = 1,176)	*p* value
BrainAGE[1,3]	0.52 (-0.56-1.70)	0.30 (-0.76-1.52)	<0.001**
Age at baseline (years)[1,3]	9.94 (9.41-10.48)	10.35 (9.81-10.75)	<0.001**
Sex[2,4]			<0.001**
Female	1,768 (49%)	638 (54%)	
Pubertal status at baseline[2,4]			<0.001**
1	2,040 (56%)	565 (48%)	
2	771 (21%)	271 (23%)	
3	779 (21%)	318 (27%)	
4	48 (1.3%)	21 (1.8%)	
5	<10	<10	
Race[2,5]			<0.001**
American Indian/Alaska Native	18 (0.5%)	<10	
Asian	87 (2.4%)	11 (0.9%)	
Black (African American)	454 (13%)	55 (4.8%)	
More than one race	380 (10%)	118 (10%)	
Native Hawaiian/Other Pacific Islander	<10	<10	
White	2,610 (73%)	944 (83%)	
Missing/Other	87	38	
Ethnicity (Hispanic)[2,4]	663 (18%)	215 (18%)	0.9
Missing	46	10	
Caregiver education (highest)[2,5]			<0.001**
Up to high school (no diploma)	131 (3.6%)	25 (2.1%)	
High school diploma/GED	236 (6.5%)	70 (6.0%)	
Some college	896 (25%)	260 (22%)	
Bachelor's degree	1,045 (29%)	288 (24%)	
Graduate school or professional degree	1,326 (36%)	533 (45%)	
Missing	<10	0	
Religious attitudes against alcohol[2,4]	2,127 (59%)	517 (44%)	<0.001**
Missing	46	14	
Prenatal alcohol exposure[2,4]	840 (24%)	427 (39%)	<0.001**
Missing	203	61	
Alcohol sipping at baseline[2,4]	741 (20%)	517 (44%)	<0.001**
Missing	0	14	

259   [1] Median (IQR); [2] n (%); [3] Wilcoxon rank sum test; [4] Pearson's Chi-squared test; [5] Fisher's exact test. Counts
260   less than 10 have been suppressed to protect participant confidentiality. * < 0.05; ** <0.001.

262　In contrast, when comparing subgroup characteristics, Experimenters (n = 461) and Bingers

263　(n = 438) only differed on a small number of measures (*Table 2*).

264　**Table 2.** Participant characteristics within Experimenter and Binger subgroups.

Characteristic	Experimenters (n = 461)	Bingers (n = 438)	*p* value
BrainAGE[1,3]	0.32 (-0.78-1.55)	0.19 (-0.83-1.41)	0.5
Age at baseline (years)[1,3]	10.29 (9.74-10.73)	10.42 (9.94-10.80)	0.009*
Sex[2,4]			0.2
Female	247 (54%)	252 (58%)	
Pubertal status at baseline[2,4]			0.5
1	218 (47%)	203 (46%)	
2	102 (22%)	100 (23%)	
3	127 (28%)	127 (29%)	
4	14 (3.0%)	<10	
5	0	<10	
Race[2,5]			0.4
American Indian/Alaska Native	<10	<10	
Asian	<10	<10	
Black (African American)	23 (5.1%)	16 (3.8%)	
More than one race	50 (11%)	40 (9.5%)	
Native Hawaiian/Other Pacific Islander	<10	<10	
White	373 (82%)	283 (86%)	
Missing/Other	<10	16	
Ethnicity (Hispanic)[2,4]	63 (14%)	87 (20%)	0.013*
Missing	<10	<10	
Caregiver education (highest)[2,5]			0.068
Up to high school (no diploma)	<10	14 (3.2%)	
High school diploma/GED	23 (5.0%)	25 (5.7%)	
Some college	90 (20%)	109 (25%)	
Bachelor's degree	108 (23%)	110 (25%)	
Graduate school or professional degree	231 (50%)	180 (41%)	
Missing	0	0	
Religious attitudes against alcohol[2,4]	237 (52%)	195 (45%)	0.039*
Missing	<10	<10	
Prenatal alcohol exposure[2,4]	169 (40%)	164 (40%)	0.9
Missing	38	27	
Alcohol sipping at baseline[2,4]	185 (40%)	204 (47%)	0.051
Missing	0	0	

[1] Median (IQR); [2] n (%); [3] Wilcoxon rank sum test; [4] Pearson's Chi-squared test; [5] Fisher's exact test. Counts

less than 10 have been suppressed to protect participant confidentiality. * < 0.05; ** <0.001.

BrainAGE model performance

In the original BrainAGE model validation, the model performed with a corrected (bias

adjusted) MAE of 1.98 (Drobinin et al., 2022), aligning with the 1–2-year range typically

reported by other adolescent BrainAGE models (Franke et al., 2012). In the current study, the

model achieved an MAE of 2.28 prior to correction for age bias, and an MAE of 1.43 years

after linear bias correction. These findings correspond closely to the study by Whitmore et al.

using the same model in ABCD data (2.32 uncorrected, 1.45 uncorrected), with minor

differences likely attributable to differences in harmonization following exclusion of

participants with missing pubertal data. Similar to the findings by Whitmore et al. (2023),

participants from the ABCD study also exhibited, on average, a marginally higher (more

positive) BrainAGE compared to those included in the original Drobinin et al. (2022) model

(*Table 2*).

Baseline BrainAGE as a predictor of early alcohol initiation

Baseline BrainAGE (Figure 1) significantly predicted alcohol use status by/at wave 6 (*Table*

3). Specifically, a one-standard-deviation increase (equivalent to 1.64 years) in BrainAGE

was associated with an 8.7% decrease in the odds of early alcohol use, holding other

covariates constant. This equates to a 9.5% increase in the odds of early alcohol use with a

one-standard-deviation decrease in BrainAGE.

Figure 1. Scan age compared to BrainAGE between the initiation and non-initiation groups.

Table 3. Results from logistic GLMM predicting early alcohol use status (non-initiation vs. initiation) from baseline BrainAGE.

Predictor	OR	95% CI	*p* value
BrainAGE (std)	0.913	0.844-0.987	0.023*
Age (std)	1.7	1.55-1.86	<.001**
Sex (female)	1.28	1.06-1.54	0.011*
Pubertal status (std)	1.04	0.95-1.14	0.377

OR = odds ratio; 95% CI = confidence intervals (odds ratio); std = standardized values. * < 0.05; ** <0.001.

 Baseline BrainAGE as a predictor of alcohol use behaviours

Contrary to the primary model, BrainAGE was not identified as a significant predictor of

drinking behaviour (experimentation versus binge drinking; *Figure 2*; *Table 4*).

Figure 2. Scan age compared to BrainAGE between the experimenter and binge drinking

groups.

Table 4. Results from logistic GLMM predicting early alcohol use behaviour (experimentation vs.

binge drinking) from baseline BrainAGE.

Predictor	OR	95% CI	*p* value
BrainAGE (std)	0.999	0.862-1.158	0.989
Age (std)	1.235	1.048-1.455	0.012*
Sex (female)	1.498	1.042-2.155	0.029*
Pubertal status (std)	0.82	0.671-1.003	0.053

OR = odds ratio; 95% CI = confidence intervals (odds ratio); std = standardized values. * < 0.05.

Sensitivity analyses

Adjustment for secondary covariates

Following inclusion of secondary covariates in the primary model, no significant association
between baseline BrainAGE and alcohol initiation status was identified (*Table S2*).

Exclusion of participants of initiating other substances

Following exclusion of participants in the non-initiating group who initiated other substances
at baseline and reported use at any time during follow-up, BrainAGE was still identified as a
significant predictor of early alcohol use (*Table S4*).

Discussion

The present study investigated whether BrainAGE in late childhood could predict subsequent alcohol initiation and use behaviour in adolescence. Overall, ABCD participants showed a higher BrainAGE on average (i.e. more "mature appearing" brains) relative to chronological age compared to the model's reference population. However, lower BrainAGE (a one-standard-deviation decrease) compared to the non-initiating group was associated with a 9.5% increased likelihood of alcohol initiation at follow-up, independent of chronological age, sex, and pubertal status. However, these effects were not robust to adjustment for sociodemographic characteristics and prior alcohol exposure. Additionally, within the initiation group, BrainAGE did not differentiate between experimentation and more harmful binge drinking behaviours. Overall, these findings suggest lower BrainAGE may predate alcohol initiation in adolescence, however, this relationship likely reflects complex, context-dependent risk rather than a direct effect of brain structure alone, warranting further investigation.

Results from our primary analysis suggest that lower BrainAGE ("less mature appearing" brains) in late childhood relative to non-initiators is predictive of subsequent alcohol initiation in adolescence. This stands in contrast to prior work identifying markers of accelerated neuromaturation, such as prefrontal cortical thinning (Brumback et al., 2016; Squeglia et al., 2017) and regional volumetric increases (Miller et al., 2024), as predictors of alcohol use. This apparent discrepancy may reflect the limitations of region-of-interest approaches, given that brain development across childhood and adolescence is characterized by coordinated, network-level reorganization and structural co-maturation (Liang et al., 2024). By contrast, BrainAGE integrates cortical and subcortical features into a unified framework, where deviations reflect imbalances within and between these systems. This suggests that lower BrainAGE in the initiation group aligns with neurodevelopmental models proposing that such imbalances heighten vulnerability to substance use (Casey & Jones, 2010; Steinberg, 2017). However, collapsing these features into a summary score from a single timepoint precludes determination of how individual regions or systems contribute to the observed differences, and how these contributions evolve across development. Future work incorporating longitudinal and complementary network-level models alongside BrainAGE (such as structural covariance networks or connectome-based approaches) would help clarify whether disruption is coordinated or system-specific, more precisely mapping findings onto neurodevelopmental models of substance use vulnerability.

The attenuation of BrainAGE group differences following adjustment for sociodemographic

characteristics and prior alcohol exposure suggests these factors may partially account for the

observed relationship between BrainAGE and alcohol initiation, and warrants cautious

interpretation of the primary findings. Both BrainAGE and alcohol initiation have been

linked to a variety of environmental factors and life experiences, including socioeconomic

status and childhood adversity (Beck et al., 2026; Cohen et al., 2024; Kendler et al., 2014;

Lui et al., 2023), while the impacts of prenatal alcohol exposure on both regional and global

brain development have been well-documented (Lees et al., 2020). This finding underscores

the need to consider the contribution of biopsychosocial factors when examining

neuroanatomical predictors of alcohol use, where future mediation and causal inference

models will be essential for disentangling the independent and interactive contributions of

these variables.

While BrainAGE was predictive of alcohol initiation, it did not differentiate between

experimental and binge drinking behaviours, despite prior evidence that these may be

neurobiologically distinct. For example, Rane et al. (2022) demonstrated that structural brain

features in early adolescence (age 14) are more strongly predictive of later binge drinking

than experimental use, suggesting the neural correlates underlying escalation may be

dissociable from and more pronounced than those associated with initiation alone. It is

possible that a composite measure such as BrainAGE, when indexed at ages 9-11, may lack

the sensitivity to capture such nuances, particularly given the reduced statistical power from a

smaller binge drinking subsample. Additional factors including family history of substance

use, impulsivity, and psychopathology have been linked to both binge drinking independently

of experimentation (Englund et al., 2008; Tschorn et al., 2021) and to BrainAGE in

adolescence (Cropley et al., 2021; Dunlop et al., 2021), thus potentially obscuring group

differences. Future work examining the relative contributions of such factors alongside

BrainAGE would assist in determining its predictive utility across the spectrum of adolescent

drinking behaviours.

Limitations

Strengths of this work include the use of a large, demographically diverse prospective

sample, consideration of multiple sociodemographic and alcohol exposure variables, and

stringent alcohol use criteria. However, interpreting BrainAGE estimates in children and

adolescents remains a significant challenge (Whitmore and Beck, 2025). As BrainAGE

collapses hundreds of regionally distinct and asynchronously maturing features into a single

global deviation score, such measures may obscure tissue- and network-specific patterns

associated with genetic, environmental, and/or psychosocial factors. As these limitations are

exacerbated within cross-sectional frameworks, normative BrainAGE trajectories in the

paediatric population will be essential for informing interpretation of such metrics. While

some longitudinal BrainAGE models are currently available (e.g. Beck et al., 2026;

Whitmore et al., 2023), the development of an externally validated, comprehensive model

integrating multimodal imaging features will be pertinent to realizing this goal (Michael et

al., 2026).

Conclusion

The current findings suggest that BrainAGE in late childhood may reflect potential risk for

subsequent alcohol use initiation, but not use behaviours, in adolescence. However, this

association likely reflects complex, context-dependent risk rather than a direct effect of brain

structure alone. Further research tracking longitudinal trajectories of BrainAGE within a

latent change score framework to examine bidirectional relationships between other genetic,

environmental, and psychosocial variables will be essential for determining the role of

BrainAGE as a predictor of alcohol use in adolescence.

Acknowledgements: Data used in the preparation of this article were obtained from

the Adolescent Brain Cognitive Development™ (ABCD) Study, held in the NIH Brain

Development Cohorts Data Sharing Platform. This is a multisite, longitudinal study designed

to recruit more than 10,000 children age 9-10 and follow them over 10 years into early

adulthood. The ABCD Study® is supported by the National Institutes of Health and

additional federal partners under award numbers U01DA041048, U01DA050989,

U01DA051016, U01DA041022, U01DA051018, U01DA051037, U01DA050987,

U01DA041174, U01DA041106, U01DA041117, U01DA041028, U01DA041134,

U01DA050988, U01DA051039, U01DA041156, U01DA041025, U01DA041120,

U01DA051038, U01DA041148, U01DA041093, U01DA041089, U24DA041123,

U24DA041147. A full list of supporters is available at https://abcdstudy.org/federal-

partners.html. A listing of participating sites and a complete listing of the study investigators

can be found at https://abcdstudy.org/consortium_members/. ABCD consortium investigators

designed and implemented the study and/or provided data but did not necessarily participate

410 in the analysis or writing of this report. This manuscript reflects the views of the authors and

411 may not reflect the opinions or views of the NIH or ABCD consortium investigators. This

412 work was supported by funding from the NIH (01AA030575, Mewton/Squeglia; and

413 K24AA031052, Squeglia).

References

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software*, *67*(1), 1–48.

https://doi.org/10.18637/jss.v067.i01

Beck, D., Carrick, C., Aksnes, E. R., MacSweeney, N., Westlye, L. T., Fuhrmann, D., &

Tamnes, C. K. (2026). Synchrony between brain age and internalising and

externalising symptoms across adolescence. In *medRxiv: The preprint server for*

health sciences (p. 2025.12.31.25343265).

https://doi.org/10.64898/2025.12.31.25343265

Bogdan, R., Hatoum, A. S., Johnson, E. C., & Agrawal, A. (2023). The Genetically Informed

Neurobiology of Addiction (GINA) model. *Nature Reviews Neuroscience*, *24*(1), 40–

57. https://doi.org/10.1038/s41583-022-00656-8

Brumback, T. Y., Worley, M., Nguyen-Louie, T. T., Squeglia, L. M., Jacobus, J., & Tapert, S.

F. (2016). Neural predictors of alcohol use and psychopathology symptoms in

adolescents. *Development and Psychopathology*, *28*(4pt1), 1209–1216.

https://doi.org/10.1017/S0954579416000766

Byrne, H., Visontay, R., Devine, E. K., Wade, N. E., Jacobus, J., Squeglia, L. M., & Mewton,

L. (2026). Brain network features predating early alcohol initiation in adolescence.

Translational Psychiatry. https://doi.org/10.1038/s41398-026-03906-w

Casey, B. J., & Jones, R. M. (2010). Neurobiology of the Adolescent Brain and Behavior:

Implications for Substance Use Disorders. *Journal of the American Academy of Child*

& Adolescent Psychiatry, *49*(12), 1189–1201.

https://doi.org/10.1016/j.jaac.2010.08.017

Cheng, T. W., Magis-Weinberg, L., Guazzelli Williamson, V., Ladouceur, C. D., Whittle, S.

L., Herting, M. M., Uban, K. A., Byrne, M. L., Barendse, M. E. A., Shirtcliff, E. A., &

Pfeifer, J. H. (2021). A Researcher's Guide to the Measurement and Modeling of

Puberty in the ABCD Study® at Baseline. *Frontiers in Endocrinology*, *Volume 12-*

2021. https://doi.org/10.3389/fendo.2021.608575

Clare, P. J., Yuen, W. S., Henderson, A., Kypri, K., Bruno, R., Slade, T., Hutchinson, D.,

McBride, N., Wodalowski, M., McCambridge, J., Degenhardt, L., Boland, V. C.,

Mattick, R. P., & Peacock, A. (2026). Does alcohol use and related harm differ based

on the age of initiation to alcohol? Results from a prospective cohort study. *Addiction*,

121(1), 44–57. https://doi.org/10.1111/add.70183

Cohen, J. W., Ramphal, B., DeSerisy, M., Zhao, Y., Pagliaccio, D., Colcombe, S., Milham, M.

P., & Margolis, A. E. (2024). Relative brain age is associated with socioeconomic

status and anxiety/depression problems in youth. *Developmental Psychology*, *60*(1),

199–209. https://doi.org/10.1037/dev0001593

Cropley, V. L., Tian, Y., Fernando, K., Mansour L., S., Pantelis, C., Cocchi, L., & Zalesky, A.

(2021). Brain-Predicted Age Associates With Psychopathology Dimensions in Youths.

Biological Psychiatry: Cognitive Neuroscience and Neuroimaging, *6*(4), 410–419.

https://doi.org/10.1016/j.bpsc.2020.07.014

Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner,

R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J.

(2006). An automated labeling system for subdividing the human cerebral cortex on

MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968–980.

https://doi.org/10.1016/j.neuroimage.2006.01.021

Drobinin, V., Van Gestel, H., Helmick, C. A., Schmidt, M. H., Bowen, C. V., & Uher, R.

(2022). The Developmental Brain Age Is Associated With Adversity, Depression, and

Functional Outcomes Among Adolescents. *Biological Psychiatry: Cognitive*

Neuroscience and Neuroimaging, *7*(4), 406–414.

https://doi.org/10.1016/j.bpsc.2021.09.004

Dunlop, K., Victoria, L. W., Downar, J., Gunning, F. M., & Liston, C. (2021). Accelerated

brain aging predicts impulsivity and symptom severity in depression.

Neuropsychopharmacology: Official Publication of the American College of

Neuropsychopharmacology, *46*(5), 911–919. https://doi.org/10.1038/s41386-021-

00967-x

Englund, M. M., Egeland, B., Oliva, E. M., & Collins, W. A. (2008). Childhood and

adolescent predictors of heavy drinking and alcohol use disorders in early adulthood:

A longitudinal developmental analysis. *Addiction*, *103*(s1), 23–35.

https://doi.org/10.1111/j.1360-0443.2008.02174.x

Fortin, J.-P., Cullen, N., Sheline, Y. I., Taylor, W. D., Aselcioglu, I., Cook, P. A., Adams, P.,

Cooper, C., Fava, M., McGrath, P. J., McInnis, M., Phillips, M. L., Trivedi, M. H.,

Weissman, M. M., & Shinohara, R. T. (2018). Harmonization of cortical thickness

measurements across scanners and sites. *NeuroImage*, *167*, 104–120.

https://doi.org/10.1016/j.neuroimage.2017.11.024

Franke, K., Luders, E., May, A., Wilke, M., & Gaser, C. (2012). Brain maturation: Predicting

individual BrainAGE in children and adolescents using structural MRI. *NeuroImage*,

63(3), 1305–1312. https://doi.org/10.1016/j.neuroimage.2012.08.001

Gardner, L. A., Stockings, E., Champion, K. E., Mather, M., & Newton, N. C. (2024).

Alcohol initiation before age 15 predicts earlier hazardous drinking: A survival

analysis of a 7-year prospective longitudinal cohort of Australian adolescents.

Addiction, *119*(3), 518–529. https://doi.org/10.1111/add.16376

Green, R., Wolf, B. J., Chen, A., Kirkland, A. E., Ferguson, P. L., Browning, B. D., Bryant, B.

E., Tomko, R. L., Gray, K. M., Mewton, L., & Squeglia, L. M. (2024). Predictors of

Substance Use Initiation by Early Adolescence. *The American Journal of Psychiatry*, *181*(5), 423–433. https://doi.org/10.1176/appi.ajp.20230882

Hagler, D. J. J., Hatton, S., Cornejo, M. D., Makowski, C., Fair, D. A., Dick, A. S., Sutherland, M. T., Casey, B. J., Barch, D. M., Harms, M. P., Watts, R., Bjork, J. M., Garavan, H. P., Hilmer, L., Pung, C. J., Sicat, C. S., Kuperman, J., Bartsch, H., Xue, F., … Dale, A. M. (2019). Image processing and analysis methods for the Adolescent Brain Cognitive Development Study. *NeuroImage*, *202*, 116091. https://doi.org/10.1016/j.neuroimage.2019.116091

Hjarnaa, L., Møller, S. P., Curtis, A. B., Becker, U., Andersen, O., Torvik, F. A., & Tolstrup, J. S. (2023). Alcohol Intake and Academic Performance and Dropout in High School: A Prospective Cohort Study in 65,233 Adolescents. *Journal of Adolescent Health*, *73*(6), 1083–1092. https://doi.org/10.1016/j.jadohealth.2023.07.008

Holm, M. C., Leonardsen, E. H., Beck, D., Dahl, A., Kjelkenes, R., de Lange, A.-M. G., & Westlye, L. T. (2023). Linking brain maturation and puberty during early adolescence using longitudinal brain age prediction in the ABCD cohort. *Developmental Cognitive Neuroscience*, *60*, 101220. https://doi.org/10.1016/j.dcn.2023.101220

Kendler, K. S., Gardner, C. O., Hickman, M., Heron, J., Macleod, J., Lewis, G., & Dick, D. M. (2014). Socioeconomic Status and Alcohol-Related Behaviors in Mid- to Late Adolescence in the Avon Longitudinal Study of Parents and Children. *Journal of Studies on Alcohol and Drugs*, *75*(4), 541–545. https://doi.org/10.15288/jsad.2014.75.541

Le, T. T., Kuplicki, R. T., McKinney, B. A., Yeh, H.-W., Thompson, W. K., Paulus, M. P., Tulsa 1000 Investigators, Aupperle, R. L., Bodurka, J., Cha, Y.-H., Feinstein, J. S., Khalsa, S. S., Savitz, J., Simmons, W. K., & Victor, T. A. (2018). A Nonlinear Simulation Framework Supports Adjusting for Age When Analyzing BrainAGE.

Frontiers in Aging Neuroscience, *Volume 10-2018*.

https://www.frontiersin.org/journals/aging-

neuroscience/articles/10.3389/fnagi.2018.00317

Lee, Y. Y., Slade, T., Chatterton, M. L., Le, L. K.-D., Perez, J. K., Faller, J., Chapman, C.,

Newton, N. C., Sunderland, M., Teesson, M., & Mihalopoulos, C. (2024). Age at first

drink and its influence on alcohol use behaviours in young adulthood: Evidence from

an Australian household-based panel study. *Preventive Medicine*, *181*, 107898.

https://doi.org/10.1016/j.ypmed.2024.107898

Lees, B., Garcia, A. M., Debenham, J., Kirkland, A. E., Bryant, B. E., Mewton, L., &

Squeglia, L. M. (2021). Promising vulnerability markers of substance use and misuse:

A review of human neurobehavioral studies. *Neuropharmacology*, *187*, 108500.

https://doi.org/10.1016/j.neuropharm.2021.108500

Lees, B., Mewton, L., Stapinski, L. A., Teesson, M., & Squeglia, L. M. (2020). Association of

prenatal alcohol exposure with preadolescent alcohol sipping in the ABCD study®.

Drug and Alcohol Dependence, *214*, 108187.

https://doi.org/10.1016/j.drugalcdep.2020.108187

Liang, X., Sun, L., Liao, X., Lei, T., Xia, M., Duan, D., Zeng, Z., Li, Q., Xu, Z., Men, W.,

Wang, Y., Tan, S., Gao, J.-H., Qin, S., Tao, S., Dong, Q., Zhao, T., & He, Y. (2024).

Structural connectome architecture shapes the maturation of cortical morphology

from childhood to adolescence. *Nature Communications*, *15*(1), 784.

https://doi.org/10.1038/s41467-024-44863-6

Lorena Charrier, Saskia van Dorsselaer, Natale Canale, Tibor Baska, Biljana Kilibarda,

Rosanna Irene Comoretto, Tommaso Galeotti, Judith Brown, & Alessio Vieno. (2024).

A focus on adolescent substance use in Europe, central Asia and Canada. Health

Behaviour in School-aged Children international report from the 2021/2022 survey.

Volume 3. World Health Organization. Policy Commons.

https://policycommons.net/artifacts/12374985/a-focus-on-adolescent-substance-use-

in-europe-central-asia-and-canada/

Lui, C. K., Witbrodt, J., Li, L., Tam, C. C., Williams, E., Guo, Z., & Mulia, N. (2023).

Associations between early childhood adversity and behavioral, substance use, and

academic outcomes in childhood through adolescence in a U.S. longitudinal cohort.

Drug and Alcohol Dependence, *244*, 109795.

https://doi.org/10.1016/j.drugalcdep.2023.109795

Michael, C., Jones, N. S., Hanson, J. L., Westerman, H. B., Klump, K. L., Mitchell, C.,

Monk, C. S., Burt, S. A., & Hyde, L. W. (2026). A Systematic Evaluation of the

Performance of Multiple Brain Age Algorithms in Two Cohorts of Youth. *Human*

Brain Mapping, *47*(2), e70458. https://doi.org/10.1002/hbm.70458

Miller, A. P., Baranger, D. A. A., Paul, S. E., Garavan, H., Mackey, S., Tapert, S. F., LeBlanc,

K. H., Agrawal, A., & Bogdan, R. (2024). Neuroanatomical Variability and Substance

Use Initiation in Late Childhood and Early Adolescence. *JAMA Network Open*, *7*(12),

e2452027–e2452027. https://doi.org/10.1001/jamanetworkopen.2024.52027

Peng, H., Gong, W., Beckmann, C. F., Vedaldi, A., & Smith, S. M. (2021). Accurate brain age

prediction with lightweight deep neural networks. *Medical Image Analysis*, *68*,

101871. https://doi.org/10.1016/j.media.2020.101871

Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of

pubertal status: Reliability, validity, and initial norms. *Journal of Youth and*

Adolescence, *17*(2), 117–133. https://doi.org/10.1007/BF01537962

Rane, R. P., de Man, E. F., Kim, J., Görgen, K., Tschorn, M., Rapp, M. A., Banaschewski, T.,

Bokde, A. L., Desrivieres, S., Flor, H., Grigis, A., Garavan, H., Gowland, P. A., Brühl,

R., Martinot, J.-L., Martinot, M.-L. P., Artiges, E., Nees, F., Papadopoulos Orfanos,

D., … IMAGEN consortium. (2022). Structural differences in adolescent brains can predict alcohol misuse. *eLife*, *11*, e77545. https://doi.org/10.7554/eLife.77545

Scheffler, F., Ipser, J., Pancholi, D., Murphy, A., Cao, Z., Ottino-González, J., Thompson, P. M., Shoptaw, S., Conrod, P., Mackey, S., Garavan, H., & Stein, D. J. (2024). Mega-analysis of the brain-age gap in substance use disorder: An ENIGMA Addiction working group study. *Addiction (Abingdon, England)*, *119*(11), 1937–1946. https://doi.org/10.1111/add.16621

Smith, S. M., Vidaurre, D., Alfaro-Almagro, F., Nichols, T. E., & Miller, K. L. (2019). Estimation of brain age delta from brain imaging. *NeuroImage*, *200*, 528–539. https://doi.org/10.1016/j.neuroimage.2019.06.017

Spear, L. P. (2018). Effects of adolescent alcohol consumption on the brain and behaviour. *Nature Reviews Neuroscience*, *19*(4), 197–214. https://doi.org/10.1038/nrn.2018.10

Squeglia, L. M., Ball, T. M., Jacobus, J., Brumback, T., McKenna, B. S., Nguyen-Louie, T. T., Sorg, S. F., Paulus, M. P., & Tapert, S. F. (2017). Neural Predictors of Initiating Alcohol Use During Adolescence. *The American Journal of Psychiatry*, *174*(2), 172–185. https://doi.org/10.1176/appi.ajp.2016.15121587

Steinberg, L. (2017). A social neuroscience perspective on adolescent risk-taking. In *Biosocial theories of crime* (pp. 435–463). Routledge.

Swendsen, J., Burstein, M., Case, B., Conway, K. P., Dierker, L., He, J., & Merikangas, K. R. (2012). Use and Abuse of Alcohol and Illicit Drugs in US Adolescents: Results of the National Comorbidity Survey–Adolescent Supplement. *Archives of General Psychiatry*, *69*(4), 390–398. https://doi.org/10.1001/archgenpsychiatry.2011.1503

Tackaberry-Giddens, L., Henderson, J., & Hawke, L. D. (2023). Early alcohol onset: Clinical and demographic characteristics of service-seeking youth. *Journal of the Canadian*

Academy of Child and Adolescent Psychiatry = Journal de l'Academie Canadienne

de Psychiatrie de l'enfant et de l'adolescent, *32*(3), 150–160.

Tanner, J. M., & Whitehouse, R. H. (1976). Clinical longitudinal standards for height, weight,

height velocity, weight velocity, and stages of puberty. *Archives of Disease in*

Childhood, *51*(3), 170. https://doi.org/10.1136/adc.51.3.170

Tschorn, M., Lorenz, R. C., O'Reilly, P. F., Reichenberg, A., Banaschewski, T., Bokde, A. L.

W., Quinlan, E. B., Desrivières, S., Flor, H., Grigis, A., Garavan, H., Gowland, P.,

Ittermann, B., Martinot, J.-L., Artiges, E., Nees, F., Papadopoulos Orfanos, D.,

Poustka, L., Millenet, S., … IMAGEN Consortium. (2021). Differential predictors for

alcohol use in adolescents as a function of familial risk. *Translational Psychiatry*,

11(1), 157. https://doi.org/10.1038/s41398-021-01260-7

Whitmore, L. B., Weston, S. J., & Mills, K. L. (2023). BrainAGE as a measure of maturation

during early adolescence. *Imaging Neuroscience (Cambridge, Mass.)*, *1*.

https://doi.org/10.1162/imag_a_00037

Whitmore, L., & Beck, D. (2025). Current challenges and future directions for brain age

prediction in children and adolescents. *Nature Communications*, *16*(1), 7771.

https://doi.org/10.1038/s41467-025-63222-7

604