Bacterial Stress Responses Lower mRNA-Protein Level Correlations

Sena Gizem Suer[1,2,3,7], Yi Yang Lim[1,2], Ganeshwari Dhurve[1,4], Rabia Şen[5,7], Jérôme Arnoux[1,2,3,7], Cemal Erdem[5,7], André Mateus[3,4,6,7] and Kemal Avican[1,2,3,7,8*]

[1]Department of Molecular Biology, Umeå University, Umeå, 90187, Sweden.

[2]Integrated Science Lab (IceLab), Umeå University, Umeå, 90187, Sweden.

[3]Umeå Centre for Microbial research (UCMR), Umeå University, Umeå, 90187, Sweden.

[4]Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå, 90187, Sweden.

[5]Department of Medical Biosciences, Science for Life Laboratory, Umeå University, Umeå, 90187, Sweden.

[6]Department of Chemistry, Umeå University, Umeå, 90187, Sweden.

[7] Science for Life Laboratory, Umeå University, Umeå, 90187, Sweden.

[8]Lead contact

[*]Correspondence: kemal.avican@umu.se (K.A.)

Abstract

Diverse bacterial pathogens have evolved complex regulatory mechanisms to adapt to various environmental stresses during infection. The uncertainty in mRNA-protein levels in response to environmental stressors complicates our understanding of bacterial physiology and their adaptation to stressful environments. To examine this issue, we have integrated transcriptomics and proteomics data on three human bacterial pathogens *Salmonella enterica* Typhimurium, *Yersinia pseudotuberculosis*, and *Staphylococcus aureus* under ten infection-relevant stress conditions. We observed positive correlations between mRNA and protein levels, which were decreased under different stress conditions. Essential genes exhibited higher expression levels with lower variation across the conditions and stronger mRNA-protein correlations compared to non-essential genes, highlighting their critical role in bacterial adaptability and survival. Moreover, we identified a substantial number of genes with stress-induced non-correlating mRNA-protein levels, particularly under conditions triggering strong stress responses. Particularly this level was dramatically lowered for osmotic stress specific genes affected by impaired translational activity under osmotic stress. Our findings highlight the prevalence of non-correlating mRNA-protein levels and the potential role of post-translational modifications in modulating protein levels in response to environmental stressors during infection. This study provides a comprehensive framework for integrating transcriptomics and proteomics data and identifies potential gene products that might significantly impact the ability of diverse bacterial pathogens to adapt to hostile infection environments.

Introduction

Bacterial pathogens have evolved complex mechanisms to adapt to different environmental stresses in the host such as nutrient limitations, temperature or pH shifts, and antimicrobial challenges during infection. They respond to deleterious effects of such stressors via stress response mechanisms, involving complex gene regulatory networks, for survival. Therefore, comprehensive understanding of bacterial stress response mechanisms is critical for development of novel therapeutic approaches to resolve bacterial infections. One useful approach is through investigation of global transcriptomes of diverse bacterial pathogens under a variety of stress conditions that mimic host environments. This provides information on regulation across diverse stress conditions that could be used in attempts to reveal complex gene regulations and identify key stress response genes. We previously performed a large-scale transcriptomic study of 32 human bacterial pathogens, PATHOgenex RNA Atlas[1], aiming to understand bacterial stress responses under 11 *in vivo* mimicking stress conditions. The study revealed a dynamic regulation of gene expression under different stress conditions and suggested a set of genes that has potential to be candidate as new antimicrobial targets. Co-PATHOgenex[2], the continuation of the PATHOgenex study, identifies stress-specific stimulons by generating gene co-expression modules. These stimulons represent sets of genes with unique expression patterns under specific stressors, providing insights into the stress adaptation to different stressors. Yet the main functional units shaping the cellular response are proteins and there is, generally, a weak correlation between mRNA and protein levels[3–11]. This highlights the complex nature of gene regulation and the diverse post-transcriptional factors affecting protein stability, and abundance[12].

Linking protein expression to gene expression can help to identify key proteins in response to specific stimuli or conditions and highlight mechanisms that drive bacterial adaptation, evasion of the immune system, and disease development[13–17]. This is crucial for developing more effective diagnostic tools, treatment strategies, and vaccines for bacterial infections[18,19]. Therefore, integrated transcriptomic and proteomic studies have emerged as powerful tools for understanding the complex molecular mechanisms underlying bacterial pathogenesis[15]. These studies measure mRNA and proteins levels from the same source and combine this information for comprehensive understanding of the interplay between pathogenic bacteria and their host environments, revealing gene products that are essential for bacterial survival, virulence, and resistance[13,20–22]. These efforts have provided valuable insights into the complex mechanisms that bacteria employ to adapt and survive under challenging conditions. Despite the numerous studies integrating transcriptome and proteome data, efforts to identify genes where mRNA and protein levels *do not* correlate are relatively limited[23,24]. Focusing on

such genes could offer valuable insight into the complex post-transcriptional and post-
translational regulatory mechanisms that escape detection by analyses addressing correlated
gene-expression patterns.

In this study, we employed an integrative transcriptomic and proteomic analysis of *Salmonella*
enterica Typhimurium, *Yersinia pseudotuberculosis*, and *Staphylococcus aureus* under ten
infection-relevant stress conditions to investigate mRNA-protein level correlations. We
revealed that mRNA-protein correlations significantly varied under different stress conditions
for the three species. Essential genes consistently showed higher mRNA and protein
expression levels and stronger correlations, emphasizing their importance in bacterial survival.
Stress conditions such as hypoxia and nutritional downshift exhibited the lowest correlations,
suggesting that post-transcriptional and post-translational regulation play a significant role
under these conditions. We identified plentiful genes that showed non-correlating mRNA-
protein levels in the three species, as well as sets of genes that were species specific,
particularly under conditions eliciting strong stress responses. Among all conditions tested,
osmotic stress stood out as exhibiting the most pronounced mRNA-protein decoupling. Using
the MOBILE[25] framework, we characterized the osmotic stress stimulon and linked the
observed mRNA-protein decoupling to a specific impairment of translation initiation.

Materials and Methods

Bacterial strains and growth conditions

Salmonella enterica subsp. enterica serovar Typhimurium SL1344, *Staphylococcus aureus*
subsp. aureus MSSA476, and *Yersinia pseudotuberculosi*s serotype O:3 (strain YPIII) species
were selected from PATHOgenex database[1] to analyze transcriptomic and proteomic data.

Salmonella enterica SL1344 and *Staphylococcus aureus* MSSA476 strains used in this study
did not carry antibiotic resistance markers and were grown in Luria–Bertani (LB) medium (BD
Difco™ LB Agar, Lennox) at 37 °C with shaking. The kanamycin-resistant wt *Y.*
pseudotuberculosis (YPIII/pIBX) strain was used in this study. Unless otherwise stated, the
wild-type strain was grown in LB medium supplemented with 50 µg/mL kanamycin at 26°C
with shaking.

Stress exposure and sample collection

Polysome profiling

The wt *Y. pseudotuberculosis* strain was grown overnight in LB, subcultured to an initial optical
density (OD_{600}) of 0.05 in 200 mL of LB. When the culture reached an OD_{600} of 0.3, the culture

was divided into three 50 mL aliquots, each subjected to one of the following conditions for 10
minutes: (1) unexposed control, (2) osmotic stress (0.5 M NaCl), or (3) low-iron stress (250
µM 2,2′-Bipyridyl; Sigma-Aldrich). Cells were harvested by filtration, immediately flash-frozen
in liquid nitrogen, and stored at -80°C until lysis.

Frozen cells were lysed by cryogenic grinding in liquid nitrogen using aluminum oxide. Lysis
buffer was added during grinding, and the lysate was allowed to thaw gradually throughout
the process. The mixture was transferred into a pre-cooled 1.5 mL reaction tube and
immediately placed on ice. Lysates were centrifuged at 18,000 x g for 5 minutes at 4°C. The
resulting supernatants (~400 µL) were divided into 50 µL aliquots in new, pre-cooled 1.5 mL
tubes and stored at −80°C. Total RNA concentration was quantified using the Qubit
fluorometer. Each aliquot was used only once for downstream polysome profiling.

Linear 15–45% sucrose gradients were prepared using 10% and 50% (w/v) sucrose solutions
in 50 mM Tris-HCl (pH 7.5) and supplemented with 100 mM NH_4Cl, 10 mM $MgCl_2$, and 6 mM
β-mercaptoethanol. Solutions were sterile filtered and gradients generated using the Gradient
Master (BioComp Instruments) following the manufacturer's preprogrammed settings
(SW60Ti rotor, short sucrose program, w/v 1st). Prepared gradients were stored at 4°C
overnight prior to use.

For polysome analysis, 50 µg of cell lysate was carefully layered onto each gradient without
disturbing the interface. Samples were centrifuged at 40,000 rpm for 2 hours and 30 minutes
at 4°C in an Optima L-100K ultracentrifuge (Beckman Coulter) equipped with an SW60Ti rotor.
Immediately following centrifugation, the sucrose gradients were carefully placed on ice to
prevent disturbance, and continuous polysome absorbance profiles at 254 nm were recorded
using the BioComp gradient fractionation system complemented with a TRIAX flow cell. 14
discrete fractions were sequentially collected from the top to the bottom of each gradient using
the automated fraction collector, following the manufacturer's instructions and stored at −80°C
until further analysis.

Puromycin assay

To assess active translation, the wt strain was cultured as described above for overnight and
subculture preparation. Upon reaching exponential phase (OD_{600} of 0.4), the culture was
distributed into four separate flasks representing the following treatment conditions: (1)
chloramphenicol (50 µg/mL), (2) unexposed control, (3) osmotic stress (0.5 M NaCl), or (4)
low-iron stress (250 µM 2,2′-Bipyridyl; Sigma-Aldrich). Cultures grew with their respective
conditions for 10 minutes. Subsequently, to abruptly freeze and capture the translational
landscape, both 10 µg/mL puromycin and 200 µg/mL chloramphenicol were simultaneously

added to the culture, followed by an additional 10-minute incubation period prior to cell lysis.
The experiment was performed in four independent biological replicates.

To extract proteins, 20 mL of bacterial culture was collected from each treatment group and
cells were immediately harvested by centrifugation at 4,000 × g for 10 minutes at 4°C. The
resulting cell pellets were washed twice with phosphate-buffered saline and resuspended in 1
ml of lysis buffer (100 mM Tris-HCl, 10 mM EDTA, 1x cOmplete™ protease inhibitor cocktail
(Roche)). The cell suspensions were lysed on ice using sonication for nine 20 second cycles,
each cycle comprised 20 second sonication step separated by 30 second intervals with 10%
amplitude. Following centrifugation at 13,000 x g for 10 minutes at 4°C, the supernatant was
collected and stored at -20°C. Protein concentrations were defined using QuBit Protein Assay
Kit (Catalog Number: Q33211).

Immunoblotting

Protein concentrations were fixed across samples, and proteins were mixed in 1x sample
buffer (2% SDS, 62.5 mM Tris-HCl, 20% glycerol, 0.05% bromophenol blue, 1% β-
mercaptoethanol). 17 µl of proteins were separated by sodium dodecyl sulfate (SDS)-PAGE
on 4-15% Mini-PROTEAN TGX gels using 80V for 20 minutes, followed by 120V for 1 hour.
SDS gel was transferred onto methanol activated Immobilon-P PVDF membrane (Millipore)
by semidry electroblotting using Bio-Rad Trans-Blot SD Semi-Dry Transfer Cell (70V for 1
hour 10 minutes). Membranes were blocked with 5% milk in 1x Tris-buffered saline with Tween
(TBST) and probed with Anti-Puromycin at 1:10,000 dilution (Sigma-Aldrich, MABE343, clone
12D10) primary antibody in 1x TBST, overnight at 4°C. Following each antibody incubation,
the membranes were washed three times for 10 minutes with 1x TBST. Membranes were
incubated with secondary antibody: monoclonal Sheep Anti-Mouse IgG - Horseradish
Peroxidase at 1:7,000 dilution (Cytiva, NA931V) in 1x TBST for 1 hour at room temperature.
The final blot was developed by chemiluminescence (Immobilon Western Chemiluminescent
HRP Substrate, Millipore) for 5 minutes, and the bands were detected with Amersham Imager
680. Band intensities were quantified using ImageJ (version 1.54p).

Transcriptomic data retrieval and proteomic data generation

The transcriptome data published for *S. enterica*, *Y. pseudotuberculosis*, and *S. aureus* were
obtained from Gene Expression Omnibus (GEO)[26] with accession number GSE152295.

Proteomics data was obtained by exposing the three bacterial species to the same stress
conditions as mentioned in Avican *et al.*[1], with unexposed control condition and excluding
virulence inducing condition. After stress exposures, 3 ml of bacterial cultures were pelleted
with 9000 RPM for 2 minutes centrifugation and cell pellets were lysed in 200 µl 2% SDS.

Thereafter, 5 µg of each sample were reduced with 20 mM TCEP. Samples were digested

with a modified sp3 protocol[27], as previously described[28]. Briefly, samples were added to a

bead suspension (10 µg of beads (Sera-Mag Speed Beads, 4515-2105-050250, 6515-2105-

050250) in 10 µl 15% formic acid and 30 µl ethanol) and incubated shaking for 15 min at room

temperature. Beads were then washed four times with 70% ethanol. Proteins were digested

overnight by adding 40 µl of 5 mM chloroacetamide, 1.25 mM TCEP, and 200 ng trypsin in

100 mM HEPES pH 8.5. Peptides were eluted from the beads and were dried under vacuum.

Peptides were then labelled with TMT11plex (Thermo Fisher Scientific), pooled and desalted

with solid-phase extraction using a Waters OASIS HLB µElution Plate (30 µm). Samples were

fractionated into 48 fractions on a reversed-phase C18 system running under high pH

conditions, with every sixth fraction being pooled together. Samples were analyzed by LC-

MS/MS using a data-dependent acquisition strategy on a Thermo Fisher Scientific Vanquish

Neo LC coupled with an Thermo Fisher Scientific Orbitrap Exploris 480. Raw files were

processed with MSFragger using standard settings for TMT[29] against FASTA databases of

each organism obtained from NCBI; for *S. enterica* Typhimurium SL1344 accession numbers

NC_016810, NC_017718, NC_017719, NC_017720; for *Y. pseudotuberculosis* accession

number NC_010465; for *S. aureus* (MSSA476) accession number NC_002952. The mass

spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via

the PRIDE partner repository[28] with the dataset identifier PXD055243.

Data processing and normalization

Processing of transcriptomic data

Transcriptomic data were imported from PATHOgenex, including statistical values such as

LogFC, p-value, and adjusted p-value. Moreover, the differential expression analysis was also

already conducted, and thus no further pre-processing was done here.

Processing of proteomic data

Raw protein files containing intensity data from TMT runs were imported into R[30] (version

4.4.2). Data were pre-processed to retain only protein locus tag, intensities values, and razor

peptide count -defined as peptides shared across multiple proteins. Proteins were then filtered

to retain only those quantified by at least two unique peptides and exhibiting non-zero intensity

values; entries with zero intensity were considered missing data and excluded. This

processing yielded one matrix per species per run, in which each row contained a protein

identifier, its total intensity, and its corresponding razor peptide count.

To estimate protein abundance, each protein was mapped to its corresponding treatment
condition. The relative reporter ion abundance of a given protein for its treatment was divided
by the total relative reporter ion abundance as a first estimate of protein abundance. Since
longer proteins tend to generate a higher number of razor peptides — and thus more ions —
this estimate was further divided by the razor peptide count to obtain the Relative Intensity
(RI), ensuring that RI reflects protein abundance independently of protein size. Matrices from
the same species were subsequently merged across runs, and proteins absent across all
replicates were excluded, yielding three species-specific matrices each containing RI values
for every detected protein across the three biological replicates and their associated treatment
conditions.

To identify the most appropriate normalization strategy, 8 normalization methods were
evaluated using the NormalyzerDE[31] package (version 1.23.2): Variance Stabilizing
Normalization (VSN), Log2-transformation, Quantile, Global Intensity, Mean, Median, Linear
Regression, and Cyclic Loess. Normalized outputs were visualized using principal component
analysis (PCA) plots generated with ggplot2[32] (version 4.0.2) to assess replicate clustering
within each stress condition. VSN consistently produced the tightest replicate clustering
across condition and was therefore applied to the three species-specific matrices to generate
VSN-normalized RIs values for downstream analysis.

Differential abundance analysis was performed on the normalized proteome data using the
limma[33] package (version 3.58.1) with default parameters, applying empirical Bayes
moderation to stabilize variance estimates across proteins. Differentially abundant proteins
were identified for each species to characterize the proteomic response to every stress
condition, and highlight proteins potentially involved in key regulatory mechanisms.

Data analysis

Correlation analysis

215   To assess the correlation between mRNA and protein levels, we used the log2-transformed
216   mean TPM values for mRNAs and the log2-transformed mean RI values for proteins from
217   three replicates of each condition. The Pearson correlation coefficient was calculated for each
218   condition to determine the strength of the correlation. Statistical significance p-value < 0.05
219   was evaluated using Student's t-test.

Identification of gene clusters with discordant mRNA and protein level regulation

To identify genes with discordant mRNA and protein level regulation under 10 stress
conditions, the LogFC values of mRNA from control data were combined with the LogFC

values of protein, calculated using limma[33] differential expression analysis to create a single
data table and visualize using scatter plots separately for each species. Thereafter, the
combined transcriptome and proteome data were filtered with p-value < 0.05 to select genes
whose mRNA or protein level changes upon exposure to stress conditions.

Genes were then grouped in 6 different clusters based on the expression patterns as
described in Table 1. Cluster 1; mRNA downregulated in at least one stress condition but
protein level is not changed, Cluster 2; mRNA downregulated in at least one condition but
protein is upregulated, Cluster 3; mRNA level is not changed in any condition but protein is
downregulated, Cluster 4; mRNA level is not changed in any condition but protein is
upregulated in at least one condition, Cluster 5; mRNA is upregulated in at least one condition
but protein is downregulated, and Cluster 6; mRNA is upregulated in at least one condition but
protein level is not changed.

Integrative omics analysis with MOBILE

MOBILE pipeline[25] is used for three species (SA, SE, and YP) with 11 samples including both
transcriptomic and proteomic datasets. The RI datasets were log2-transformed and
normalized. First, they were filtered to only retain the most highly variant features (top 10% for
transcripts and 20% for proteins). Next, the MOBILE pipeline (MATLAB version R2023a) is
run, using the glmnet package[34] and $Y = \beta.X + \delta$ formalism, where Y matrix is the proteomics
dataset and X is the transcriptomics. The objective is to obtain the coefficients matrix β. δ is
the y-intercept values, which are almost zero by design. The regression process is repeated
10000 times with randomly selected random number generation seeds to yield an ensemble
of Lasso matrices. By identifying coefficients that appeared in at least half of the models, a
robust Lasso coefficient matrix is selected. This matrix represents the stable associations
between analytes across omics layers.

To find stress-specific interactions, MOBILE employs a "Leave-One-Group-Out" (LOGO)
strategy where each sample exposed to the stress is excluded from the input one-at-a-time.
Here, both the osmotic stress and control condition samples were subtracted, and
corresponding final matrices were named as OSS-IAN and CTRL-IAN matrices. IANs are the
Integrated Association Networks that are coalesced gene-level networks, where nodes
represent genes of the matrix analytes and edges stand for the non-zero coefficients.

Functional annotation

Enrichment analysis

Gene Ontology (GO) enrichment analyses were conducted using the clusterProfiler[35] package v4.10.1 in R. Most significant enriched terms were identified using a cut-off criterion of p-value<0.05. Plots showing the most significant enriched Biological Process (BP) GO terms were generated using the ggplot2 package version 4.0.2.

Over representation analysis (ORA)

The subsequent step involves comparing OSS-IAN and FULL-IAN (all condition matrix), to obtain OSS-dependent interactions. Similarly, comparing CTRL-IAN to FULL-IAN yields CTRL-dependent interactions. Next, pathway enrichment analysis (clusterProfiler[35] version 4.14.6) was adopted for the genes list obtained by comparing OSS-dependent list to CTRL-dependent gene list. Gene Ontology gene sets curated for three species (*SA, SE, YP)* were used in gmt format with a minimum gene set size of 15 and a maximum of 500. Significantly enriched gene sets were filtered by q-value < 0.2 and p-value < 0.05.

Results

Transcriptomic and proteomic analysis of three human bacterial pathogens under ten infection relevant stress conditions

We analyzed transcriptomic and proteomic datasets of three human bacterial pathogens including Gram-negative *Salmonella enterica* Typhimurium SL1344, *Yersinia pseudotuberculosis* YPIII and Gram-positive *Staphylococcus aureus* MSSA476. We used transcriptomes of the three pathogens under 10 infection relevant stress conditions together with unexposed control condition from PATHOgenex RNA Atlas[1] and complemented it with mass-spectrometry proteomic profiling under the same stressors (**Figure 1A**). As mass spectrometry-based proteomics often yields a lower coverage than RNA-seq, we compared detected transcripts and proteins across all conditions in three pathogens, focusing on protein coding sequences (CDSs) in the analysis. While almost all CDSs were detected by RNA-seq for the three pathogens, proteomics could detect between 48% of the CDSs for *S. enterica* Typhimurium and *Y. pseudotuberculosis*, and 32% for *S. aureus* (**Figure 1B**). Detected proteins covered nearly the full spectrum of mRNA expressions, with a bias towards highly expressed mRNAs across the conditions for the three pathogens (**Figure 1C**).

mRNA and protein level correlation under stress

The correlation between mRNA and protein levels has been previously explored in bacteria under individual growth conditions[24,36–38]. Our data allows us to assess whether mRNA levels can serve as an accurate indicator of protein levels in three human bacterial pathogens. To evaluate how different stressors affect the mRNA-protein level correlation, we calculated the intensity relative to razor peptide (RI) **(Supplementary Table 1)** values for protein levels and compared to mRNA levels in Transcript Per Million (TPM)[39] for every stress condition including the unexposed control condition. We have previously shown that the three biological replicates of the RNA-seq samples clustered together[1]. For the proteome dataset generated in this study, we conducted pairwise comparisons of the three replicates of each condition and computed Pearson correlation coefficient values (R-value). The R-value for pairwise comparison within the replicates of the same condition varied from 0.86-0.96 for the three pathogens **(Supplementary Figures 1-3)** showing strong similarities between the replicates. We then measured the correlation between mRNA and protein levels by calculating Pearson correlation coefficient values of the log2 transformed mean TPM and log2 transformed mean RI of each condition for the three pathogens **(Figure 2A, B, and C)**. For these measurements, we limited the analysis to the genes that are detected as mRNA and protein across all tested conditions for each of the pathogens. In the present study, we observed that correlations between mRNA and protein levels under 11 conditions ranged from 0.46-0.65 for *S. enterica* Typhimurium, 0.51-0.71 for *S. aureus* and 0.55-0.72 for *Y. pseudotuberculosis* **(Figure 2A, B, and C)**. These correlation coefficients are comparable to those reported in previous studies across diverse biological systems[24,38,40–42]. The variation in mRNA–protein correlation across stress conditions suggests that distinct molecular mechanisms, activated by each stressor, might modulate translational or post-translational regulation.

Essential genes have higher mRNA-protein level correlations in the three species

The positive correlation between mRNA-protein levels for all detected genes prompted us to investigate subsets of genes with different correlations levels. Essential genes have been shown to have stronger correlations in comparison to non-essentials for *Pseudomonas aeruginosa* during different growth phases[23]. Therefore, we sought to investigate mRNA-protein level correlation profiles of essential genes in the three species. Of the 354, 341, and 445 previously identified essential genes for *S. enterica* Typhimurium[43] *S. aureus*[44] and *Y. pseudotuberculosis*[45] respectively, we used 300 (85%), 278 (82%), and 335 (75%) essential genes detected in both our transcriptome and proteome datasets. We found that mean expression levels of mRNAs and proteins were significantly higher for essential genes in

comparison to non-essentials for the three species (**Figure 3A and B).** This was true for the expression levels in every condition (**Supplementary Figures 4 and 5**). Moreover, we observed that essential genes have higher mRNA-protein level correlation than all detected genes for all three species (**Figure 3C**), except for *S. enterica* Typhimurium and *Y. pseudotuberculosis* under nutritional downshift (Nd) stress condition. As previously observed in other species[28,46], essential genes showed significantly lower variance on both mRNA and protein expression across the 11 conditions for the three species (**Figure 3D and E**). These findings show that essential genes of *S. enterica* Typhimurium, *S. aureus*, and *Y. pseudotuberculosis* have high and stable expression at both mRNA and protein level under a wide range of stress conditions with robust correlations between mRNA and protein levels. This underlines the critical role these genes play in the organisms' adaptability and survival under different stress conditions.

Stress responses differentially affect mRNA-protein correlation

The variation in correlation under different stress conditions prompted us to investigate possible underlaying reasons. The mRNA-protein levels in Hypoxia (Hyp) and Nutritional downshift (Nd) showed the lowest correlation across three species **(Figure 2A, B, and C)**. Those stress conditions differ from others in the length of the exposure time: 4 hours for Hyp and 30 minutes for Nd, against 10 minutes for others **(Figure 1A)**. To determine whether exposure duration influences mRNA-protein correlation, we examined the relationship between correlation and exposure time across all conditions. Despite a slight negative trend between the mRNA-protein level correlation and the exposure time (R=-0.33; *p-value*=0.078), this relationship was not statistically significant **(Figure 4A).** Additionally, multiple conditions did not follow this trend.

To assess whether the degree of stress responses through differentially regulated mRNAs and proteins has an impact on mRNA-protein level correlations, we used differentially expressed genes (DEGs) and differentially expressed proteins (DEPs) for each condition in the three species by using unexposed samples as control. Before identifying DEPs, we evaluated 8 normalization methods from NormalyzerDE[31], on the RI protein abundance data and found that Variance Stabilizing Normalization (VSN) was the most effective in clustering the condition together (**Supplementary Figure 6**). Then we used limma[33] differential expression package[33] for the identification of DEPs with VSN normalized data. For DEGs, we used previously reported differential expression analysis of PATHOgenex RNA Atlas[1]. We employed *p*-value < 0.05 as the threshold to identify DEGs and DEPs for each condition. Notably, the number of DEGs were significantly higher compared to DEPs in all conditions for the three species **(Figure 4B)**. This can be linked to the stronger regulation of mRNA levels

than of protein levels in humans[47,48]. Moreover, percentages of DEGs and DEPs were highest under Hyp, Nd, and stationary phase (Sp) conditions for the three pathogens (**Figure 4B**). Next, we tested whether the percentages of DEGs and DEPs have an impact on mRNA-protein level correlations. To do this, we measured the association between the percentages of DEGs and DEPs and the mRNA-protein level correlations with Pearson correlation coefficient. Our results revealed a strong negative correlation between the percentages of DEGs and DEPs and the mRNA-protein level correlation (**Figure 4C**). This suggests that other factors, such as post-transcriptional and post-translational regulation triggered by different stressors, might be influencing protein levels alongside the mRNA expression.

Identification of genes whose mRNA-protein levels are discordantly regulated under stress conditions

The decreased correlation between mRNA-protein levels with increased number of DEGs and DEPs under specific stress conditions led us to investigate the genes whose mRNA-protein levels are discordantly regulated when the bacteria are exposed to stress conditions. We clustered genes into 6 groups (Table 1: Classification of differentially expressed genes based on mRNA and protein regulation patterns.**Table 1**) corresponding to distinct combinations of differential expression patterns at the mRNA and protein levels (**Figure 5A and B and Supplementary Table 2**). In agreement with lowest correlation between mRNA-protein levels observed under Hyp, Nd, and Sp conditions (**Figure 2A-C**), discordantly regulated genes were mostly found under those conditions for the three species (**Supplementary Figure 7**). Noteworthy, Clusters 1 and 6, where mRNAs were regulated but not proteins, had the highest number of discordantly regulated genes. This is in accordance with higher number of differentially regulated mRNAs (DEGs) compared to differentially regulated proteins (DEPs) in all conditions for the three species (**Figure 4B**).

Table 1: Classification of differentially expressed genes based on mRNA and protein regulation patterns.

Cluster	mRNA regulation	Protein regulation	Description
1	▼ Downregulated	● Not changed	mRNA is downregulated in at least one stress condition, while protein level remains unchanged
2	▼ Downregulated	▲ Upregulated	mRNA is downregulated in at least one condition, while protein level is upregulated
3	● Not changed	▼ Downregulated	mRNA level remains unchanged across all conditions, while protein is downregulated
4	● Not changed	▲ Upregulated	mRNA level remains unchanged across all conditions, while protein is upregulated in at least one condition
5	▲ Upregulated	▼ Downregulated	mRNA is upregulated in at least one condition, while protein level is downregulated
6	▲ Upregulated	● Not changed	mRNA is upregulated in at least one condition, while protein level remains unchanged

Discordantly regulated genes are enriched in translation

To investigate the underlying mechanisms leading to lowered mRNA-protein level correlation under stress conditions, we performed Gene Ontology[49] (GO) enrichment analysis for genes exhibiting discordant mRNA and protein levels. Interestingly, we found that biological processes such as translation, RNA processing, RNA modification, proteolysis, and cell division, among other biological processes were commonly enriched for the three species, particularly when mRNA levels were changed but there was no change at the protein level (**Figure 5C and Supplementary Table 3**). This suggests that genes encoding subunits of translation machinery together with other components of translation such as tRNA and rRNA were discordantly regulated under stress conditions for the three pathogens. In detail, proteins in Cluster 1 have stable levels despite mRNA being downregulated, likely due to increased protein stability. In contrast, proteins in Cluster 6 have stable levels despite mRNA being upregulated, suggesting reduced translation or increased proteolysis, which is enriched for *S. enterica* Typhimurium (**Figure 5C**). Moreover, genes encoding proteins involved in central metabolism such as tricarboxylic acid cycle were commonly enriched for the three species in Cluster 5, where mRNAs were upregulated and proteins downregulated. This indicates presence of post-transcriptional regulations such as regulation via small regulatory RNAs involved in carbon metabolism[50,51]or post-translational modifications which have high prevalence in bacterial metabolic enzymes[52].

mRNA-protein level stress-specific stimulon genes under stress

We previously identified stress-specific stimulon genes as sets of genes that are co-regulated in response to distinct stress conditions[2]. To assess whether these stimulon genes exhibit distinct mRNA-protein level correlations compared to other genes, we analyzed mRNA and protein differential expression data across all genes and stress-specific stimulons. No significant differences in mRNA-protein level correlations were observed between stimulon genes and other genes across most conditions except in the case of the osmotic stress-specific stimulon (**Figure 6A**). Notably, genes within the osmotic stimulon showed a markedly lower mRNA-protein level correlation across all three pathogens studied.

To investigate this further, we used MOBILE framework[25] to identify osmotic stress specific associations between mRNA and protein levels. The variation in mRNA–protein correlation across stress conditions suggests that distinct molecular mechanisms, differentially activated by each stressor, modulate translational or post-translational regulation. Through repeated Lasso regression between paired omics layers, MOBILE enables the identification of condition-specific associations without requiring prior knowledge. The transcriptomic and

proteomic datasets with all conditions were used as an input for MOBILE, yielding a list of statistically robust associations between proteins and mRNAs. Next, the osmotic stress (Oss) condition data from both datasets were excluded from the input, and MOBILE is re-run to obtain condition-dependent associations list by comparing the all-data run to the latter. Similarly, the control (Ctrl) condition data is held out, and MOBILE re-run yielded another list of associations to identify CTRL-data dependent associations (**Figure 6B**). By comparing the osmotic stress-dependent and control-dependent association lists, we determined by over representation analysis (ORA), that the OSS-specific associations can be used to generate a gene-level network and a weighted genes list based on connectivity of each gene. Using that gene list, we ran a GO enrichment analysis. Interestingly, translation-related processes and components of the translational machinery were enriched only for *Y. pseudotuberculosis* (**Figure 6C**). This suggests a functional link between osmotic stress-specific mRNA-protein decoupling and translational regulation.

Based on this finding, we hypothesized that the reduced mRNA-protein correlation observed for osmotic stress-specific stimulon genes might result from impaired translation under stress. To test this, we performed polysome profiling on *Y. pseudotuberculosis* exposed to osmotic stress for 10 minutes, using unstressed bacteria collected at matched time points as controls. The polysome profiles showed no notable differences between osmotic stress and control condition (**Figure 6D**), indicating that global translation remains active under osmotic stress. However, global translational activity may be insufficient to meet the heightened translational demand triggered by the strong transcriptional induction of osmotic stress-specific stimulon genes.

To further characterize the translational response, translation efficiency was assessed using a puromycin incorporation assay across osmotic stress, low iron, and unexposed control conditions. Chloramphenicol-treated bacteria served as a translationally inactive control. Since puromycin causes a premature translation termination, its incorporation serves as a direct proxy for active translation. The results showed that translational activity was significantly reduced under osmotic stress compared to both unexposed control and low iron conditions, indicating that osmotic stress specifically impairs translation activity (**Figure 6E**). Therefore, the reduced mRNA-protein correlation for osmotic stress-specific stimulons is not surprising given the overall reduction in translational activity. These stimulon genes exhibit significant changes in mRNA abundance during stress. However, because the translational machinery is impaired, it cannot efficiently process or respond to these altered transcript levels. As a result, the correlation between mRNA and protein decreases. Notably, this decoupling is specific to osmotic stress and was not observed under low iron conditions.

Discussion

Bacterial species continuously sense and respond to changes in stressful host conditions and environments. Such stress responses include a series of regulatory mechanisms at both transcriptional and translational levels leading to changes on mRNA and subsequently protein level. However, it is known that transcription and translation could be uncoupled constitutively or under certain conditions, such as stress conditions, at a global or at a gene-specific level[53,54] Therefore, the relationship between mRNA and protein levels can vary depending on the conditions, and this variability has been the subject of extensive research and ongoing debate[12,48,54,55]. Nevertheless, mRNA levels have been shown to be positively correlated with protein levels in many different organisms[8,23,36,40,41,56–60]. Revealing relationship between mRNA and protein level in pathogenic bacteria is therefore critical for understanding bacterial pathogenesis and adaptation mechanism to host conditions during infection. Integrative transcriptome and proteome analyses have previously shown lowered mRNA-protein correlation levels under stress conditions such as stationary phase, heat shock, osmotic stress, and oxidative stress in *P. aeruginosa* and *Mycoplasma pneumoniae*[23,59].

We conducted a comprehensive analysis by integrating transcriptomic and proteomic datasets of *S. enterica* Typhimurium, *Y. pseudotuberculosis*, and *S. aureus* under 10 infection-relevant stress conditions, as well as a control condition during exponential growth. The inclusion of diverse pathogens, which are distantly related yet possess conserved genes, enabled us to identify both common and species-specific mRNA-protein relationships across a range of conditions mimicking host environments. Consistent with prior findings in other organisms, our analyses indicate a positive correlation between mRNA and protein levels across the three bacterial pathogens. However, this correlation is significantly reduced under stress conditions, particularly those that trigger a strong stress response.

This reduction is not surprising, as stress response proteins might be involved in post-transcriptional or post-translational regulations that can decouple mRNA and protein levels. Such regulations can occur through the stabilization or destabilization of mRNA and proteins as a response to the stress. RNA-binding proteins such as CsrA and ProQ can bind to target mRNAs under specific conditions, either stabilizing or destabilizing them, or preventing their translation[61–66]. Furthermore, the discoordination of transcription and translation elongation rates under stress conditions may contribute to the decrease in mRNA-protein level correlation[67]. For example, in *Escherichia coli* exposed to oxidative stress, there is a significant induction of mRNA expression of the OxyR regulon, which is crucial for survival under oxidative stress, while the translation elongation rate significantly slows down[68,69].

In contrast, under nutrient-limited conditions, there has been tight coordination between

transcription and translation[70,71]. This suggests that the reduced mRNA-protein level

correlation in our nutritional downshift condition might not be solely due to the discoordination

of transcription and translation but rather other regulatory mechanisms. Conformational

changes in riboswitches upon a change in the environment can also lead to reprogrammed

mRNA translation and degradation, as is the case for the lysC riboswitch in the presence and

absence of lysine[72].

Even though mostly studied in eukaryotes, post-transcriptional nucleotide modifications

triggered by stress conditions can also have an impact on mRNA-protein level as it is shown

for increased level of m^6A methylation under hypoxic for human cancer cells[73]. On the other

hand, m^6A methylation was shown to be lowered under increased temperature for *P.*

aeruginosa[74]. Similarly post-translational modifications such as methylation, phosphorylation,

acetylation, and sumoylation can directly impact protein stability or indirectly through triggered

signaling cascade in both eukaryotes and prokaryotes[75,76]. Even though protein stability was

not specifically investigated, it was shown that cellular protein phosphorylation was increased

during *Streptomyces* differentiation to form spores, which is induced by stress conditions[77].

To assess whether the mRNA-protein discrepancies observed for DEGs are a common

feature across species, we identified conserved DEGs across the three pathogens and

compared their expression patterns. These conserved genes exhibited similar non-correlating

expression patterns under the same stress conditions, suggesting that post-transcriptional and

post-translational modifications affecting these genes may be conserved across species. This

highlights the evolutionary significance of these regulatory mechanisms in bacterial adaptation

to stress.

Among all conditions investigated for stress stimulons, osmotic stress exhibited the most

pronounced mRNA-protein decoupling. This impairment may stem from structural remodeling

of the cell envelope. Osmotic stress induces rapid cell depolymerization to manage turgor

pressure[78,79], which can physically disrupt membrane integrity and the efficiency of

transmembrane transporters. In case of deficient methionine uptake, translation initiation

would be affected due to the need of formylmethionine at the start of bacterial protein

synthesis[56,57]

Our study underscores the importance of integrating transcriptomic and proteomic data to gain

a holistic understanding of bacterial responses to infection-relevant stressors. The observed

discrepancies between mRNA and protein levels emphasize the need for caution when

inferring protein abundance solely from transcriptomic data The precise post-transcriptional

and post-translational mechanisms contributing to the observed mRNA-protein discrepancies

remain to be elucidated, as the currently available datasets do not allow their direct
identification, representing an important avenue for future investigation.

In conclusion, our findings provide a comprehensive overview of mRNA-protein relationships
under stress conditions in 3 bacterial pathogens, laying the groundwork for further
investigations into the regulatory mechanisms mediating bacterial adaptation and survival in
hostile host environments. The multi-species, multi-condition nature of this dataset
furthermore represents a valuable resource for training predictive models of protein
abundance from mRNA levels, particularly under stress conditions where the mRNA-protein
relationship deviates substantially from baseline.

Funding

This work was supported by Swedish Research Council (No. 2021-02466), Kempestiftelserna
(JCK22-0017), and the Medical Faculty at Umeå University (FS 2.1.6-281-22) to K. Avican, by
Swedish Research Council Excellence Center grant (No. 2022-06543) for the Center for
Modeling Adaptive Mechanisms in Living Systems Under Stress to K. Avican

CE is funded by the Knut and Alice Wallenberg Foundation under the SciLifeLab and
Wallenberg Data Driven Life Science Program (KAW 2020.0239).

References

1. Avican, K. *et al.* RNA atlas of human bacterial pathogens uncovers stress dynamics linked to
infection. *Nat. Commun.* **12**, 3282 (2021).
2. Fernandez, L., Rosvall, M., Normark, J., Fallman, M. & Avican, K. Co-PATHOgenex web
application for assessing complex stress responses in pathogenic bacteria. *Microbiol Spectr* **12**,
e02781-23 (2024).
3. Lee, M. J. & Yaffe, M. B. Protein Regulation in Signal Transduction. *Cold Spring Harb Perspect*
Biol **8**, a005918 (2016).
4. Bathke, J., Konzer, A., Remes, B., McIntosh, M. & Klug, G. Comparative analyses of the variation
of the transcriptome and proteome of Rhodobacter sphaeroides throughout growth. *BMC*
Genomics **20**, 358 (2019).
5. Zhu, Y. *et al.* Integrative analysis of transcriptome and proteome provides insights into adaptation
to cadmium stress in Sedum plumbizincicola. *Ecotoxicol Env. Saf* **230**, 113–149 (2022).
6. Liu, S. *et al.* Transcriptome and Proteome of Methicillin-Resistant Staphylococcus aureus Small-
Colony Variants Reveal Changed Metabolism and Increased Immune Evasion. *Microbiol Spectr*
11, e0189822 (2023).
7. Grunberger, F. *et al.* Uncovering the temporal dynamics and regulatory networks of thermal stress
response in a hyperthermophile using transcriptomics and proteomics. *mBio* **14**, e0217423 (2023).
8. Edfors, F. *et al.* Gene-specific correlation of RNA and protein levels in human cells and tissues.
Mol Syst Biol **12**, 883 (2016).
9. Feng, X. & Ma, Q. Transcriptome and proteome profiling revealed molecular mechanism of
selenium responses in bread wheat (Triticum aestivum L.). *BMC Plant Biol* **21**, 584 (2021).
10. Backman, M. *et al.* Multi-omics insights into functional alterations of the liver in insulin-deficient
diabetes mellitus. *Mol Metab* **26**, 30–44 (2019).

11. Chen, G. *et al.* Discordant protein and mRNA expression in lung adenocarcinomas. *Mol Cell*
Proteomics **1**, 304–13 (2002).

12. Vogel, C. & Marcotte, E. M. Insights into the regulation of protein abundance from proteomic and
transcriptomic analyses. *Nat Rev Genet* **13**, 227–32 (2012).

13. Wang, J. *et al.* Combined proteomic and transcriptomic analysis of the antimicrobial mechanism of
tannic acid against Staphylococcus aureus. *Front Pharmacol* **14**, 1178177 (2023).

14. Kavela, S. *et al.* Use of an Integrated Multi-Omics Approach To Identify Molecular Mechanisms and
Critical Factors Involved in the Pathogenesis of Leptospira. *Microbiol Spectr* **11**, e0313522 (2023).

15. Huang, L. *et al.* Integration of Transcriptomic and Proteomic Approaches Reveals the Temperature-
Dependent Virulence of Pseudomonas plecoglossicida. *Front Cell Infect Microbiol* **8**, 207 (2018).

16. Kocharunchitt, C., King, T., Gobius, K., Bowman, J. P. & Ross, T. Integrated transcriptomic and
proteomic analysis of the physiological response of Escherichia coli O157:H7 Sakai to steady-state
conditions of cold and water activity stress. *Mol Cell Proteomics* **11**, M111.009019 (2012).

17. Xiong, Q. *et al.* Integrated transcriptomic and proteomic analysis of the global response of
Synechococcus to high light stress. *Mol Cell Proteomics* **14**, 1038–53 (2015).

18. Perez-Llarena, F. J. & Bou, G. Proteomics As a Tool for Studying Bacterial Virulence and
Antimicrobial Resistance. *Front Microbiol* **7**, 410 (2016).

19. Yang, X. *et al.* RNA markers for ultra-rapid molecular antimicrobial susceptibility testing in
fluoroquinolone-treated Klebsiella pneumoniae. *J Antimicrob Chemother* **75**, 1747–1755 (2020).

20. Ballen, V. *et al.* Antibiotic Resistance and Virulence Profiles of Klebsiella pneumoniae Strains
Isolated From Different Clinical Sources. *Front Cell Infect Microbiol* **11**, 738223 (2021).

21. Chaffin, D. O., Taylor, D., Skerrett, S. J. & Rubens, C. E. Changes in the Staphylococcus aureus
transcriptome during early adaptation to the lung. *PLoS One* **7**, e41329 (2012).

22. Liou, G. G., Chao Kaberdina, A., Wang, W. S., Kaberdin, V. R. & Lin-Chao, S. Combined
Transcriptomic and Proteomic Profiling of E. coli under Microaerobic versus Aerobic Conditions:
The Multifaceted Roles of Noncoding Small RNAs and Oxygen-Dependent Sensing in Global Gene
Expression Control. *Int J Mol Sci* **23**, 2570 (2022).

23. Zhang, M. *et al.* Impact of Growth Rate on the Protein-mRNA Ratio in Pseudomonas aeruginosa.
mBio **14**, e0306722 (2023).

24. Maier, T. *et al.* Quantification of mRNA and protein and integration with protein turnover in a
bacterium. *Mol. Syst. Biol.* **7**, MSB201138 (2011).

25. Erdem, C., Gross, S. M., Heiser, L. M. & Birtwistle, M. R. MOBILE pipeline enables identification of
context-specific networks and regulatory mechanisms. *Nat. Commun.* **14**, 3991 (2023).

26. Barrett, T. *et al.* NCBI GEO: archive for functional genomics data sets—update. *Nucleic Acids Res.*
41, D991–D995 (2013).

27. Hughes, C. S. *et al.* Ultrasensitive proteome analysis using paramagnetic bead technology. *Mol*
Syst Biol **10**, 757 (2014).

28. Mateus, A. *et al.* The functional proteome landscape of Escherichia coli. *Nature* **588**, 473–478
(2020).

29. Kong, A. T., Leprevost, F. V., Avtonomov, D. M., Mellacheruvu, D. & Nesvizhskii, A. I. MSFragger:
ultrafast and comprehensive peptide identification in mass spectrometry-based proteomics. *Nat*
Methods **14**, 513–520 (2017).

30. R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation for
Statistical Computing, Vienna, Austria, 2024).

31. Willforss, J., Chawade, A. & Levander, F. NormalyzerDE: Online Tool for Improved Normalization
of Omics Expression Data and High-Sensitivity Differential Expression Analysis. *J Proteome Res*
18, 732–740 (2019).

32. Wickham, H. *Ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag New York, 2016).

33. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

34. Friedman, J. H., Hastie, T. & Tibshirani, R. Regularization Paths for Generalized Linear Models via
Coordinate Descent. *J. Stat. Softw.* **33**, 1–22 (2010).

35. Wu, T. *et al.* clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. *Innov. Camb*
2, 100141 (2021).

36. Caglar, M. U. *et al.* The E. coli molecular phenotype under different growth conditions. *Sci. Rep.* **7**,
45303 (2017).

37. Kwon, T., Huse, H. K., Vogel, C., Whiteley, M. & Marcotte, E. M. Protein-to-mRNA Ratios Are
Conserved between Pseudomonas aeruginosa Strains. *J. Proteome Res.* **13**, 2370–2380 (2014).
38. Jayapal, K. P. *et al.* Uncovering Genes with Divergent mRNA-Protein Dynamics in Streptomyces
coelicolor. *PLOS ONE* **3**, e2097 (2008).
39. Li, B. & Dewey, C. N. RSEM: accurate transcript quantification from RNA-Seq data with or without
a reference genome. *BMC Bioinformatics* **12**, 323 (2011).
40. Choi, Y. W., Park, S. A., Lee, H. W., Kim, D. S. & Lee, N. G. Analysis of growth phase-dependent
proteome profiles reveals differential regulation of mRNA and protein in Helicobacter pylori.
PROTEOMICS **8**, 2665–2675 (2008).
41. Chen, W.-H. *et al.* Integration of multi-omics data of a genome-reduced bacterium: Prevalence of
post-transcriptional regulation and its correlation with protein abundances. *Nucleic Acids Res.* **44**,
1192–1202 (2016).
42. Lu, P., Vogel, C., Wang, R., Yao, X. & Marcotte, E. M. Absolute protein expression profiling
estimates the relative contributions of transcriptional and translational regulation. *Nat. Biotechnol.*
25, 117–124 (2007).
43. Barquist, L. *et al.* A comparison of dense transposon insertion libraries in the Salmonella serovars
Typhi and Typhimurium. *Nucleic Acids Res.* **41**, 4549–4564 (2013).
44. Chaudhuri, R. R. *et al.* Comprehensive identification of essential Staphylococcus aureus genes
using Transposon-Mediated Differential Hybridisation (TMDH). *BMC Genomics* **10**, 291 (2009).
45. Willcocks, S. J., Stabler, R. A., Atkins, H. S., Oyston, P. F. & Wren, B. W. High-throughput analysis
of Yersinia pseudotuberculosis gene essentiality in optimised in vitro conditions, and implications
for the speciation of Yersinia pestis. *BMC Microbiol.* **18**, 46 (2018).
46. Sigalova, O. M., Shaeiri, A., Forneris, M., Furlong, E. E. & Zaugg, J. B. Predictive features of gene
expression variation reveal mechanistic link with differential expression. *Mol. Syst. Biol.* **16**,
MSB209539 (2020).
47. Robles, M. S., Cox, J. & Mann, M. In-Vivo Quantitative Proteomics Reveals a Key Contribution of
Post-Transcriptional Mechanisms to the Circadian Regulation of Liver Metabolism. *PLOS Genet.*
10, e1004047 (2014).
48. Jovanovic, M. *et al.* Dynamic profiling of the protein life cycle in response to pathogens. *Science*
347, 1259038 (2015).
49. Ashburner, M. *et al.* Gene Ontology: tool for the unification of biology. *Nat. Genet.* **25**, 25–29 (2000).
50. Durica-Mitic, S., Göpel, Y. & Görke, B. Carbohydrate Utilization in Bacteria: Making the Most Out
of Sugars with the Help of Small Regulatory RNAs. *Microbiol. Spectr.* **6**, 10.1128/microbiolspec.rwr-
0013–2017 (2018).
51. Raina, M. *et al.* Dual-function AzuCR RNA modulates carbon metabolism. *Proc. Natl. Acad. Sci.*
119, e2117930119 (2022).
52. Pisithkul, T., Patel, N. M. & Amador-Noguez, D. Post-translational modifications as key regulators
of bacterial metabolic fluxes. *Curr. Opin. Microbiol.* **24**, 29–37 (2015).
53. Gebauer, F. & Hentze, M. W. Molecular mechanisms of translational control. *Nat. Rev. Mol. Cell
Biol.* **5**, 827–835 (2004).
54. Maier, T., Güell, M. & Serrano, L. Correlation of mRNA and protein in complex biological samples.
FEBS Lett. **583**, 3966–3973 (2009).
55. Schwanhäusser, B. *et al.* Global quantification of mammalian gene expression control. *Nature* **473**,
337–342 (2011).
56. Becker, K. *et al.* Quantifying post-transcriptional regulation in the development of Drosophila
melanogaster. *Nat. Commun.* **9**, 4970 (2018).
57. Corbin, R. W. *et al.* Toward a Protein Profile of Escherichia coli: Comparison to Its Transcription
Profile. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 9232–9237 (2003).
58. Gygi, S. P., Rochon, Y., Franza, B. R. & Aebersold, R. Correlation between Protein and mRNA
Abundance in Yeast. *Mol. Cell. Biol.* **19**, 1720–1730 (1999).
59. Jeacock, L., Faria, J. & Horn, D. Codon usage bias controls mRNA and protein abundance in
trypanosomatids. *eLife* **7**, e32496 (2018).
60. Ponnala, L., Wang, Y., Sun, Q. & van Wijk, K. J. Correlation of mRNA and protein abundance in
the developing maize leaf. *Plant J.* **78**, 424–440 (2014).

61. Yakhnin, H. *et al.* CsrA Represses Translation of sdiA, Which Encodes the N-Acylhomoserine-l-
Lactone Receptor of Escherichia coli, by Binding Exclusively within the Coding Region of sdiA
mRNA. *J. Bacteriol.* **193**, 6162–6170 (2011).

62. Pourciau, C., Lai, Y.-J., Gorelik, M., Babitzke, P. & Romeo, T. Diverse Mechanisms and Circuitry
for Global Regulation by the RNA-Binding Protein CsrA. *Front. Microbiol.* **11**, (2020).

63. Wang, X. *et al.* CsrA post-transcriptionally represses pgaABCD, responsible for synthesis of a
biofilm polysaccharide adhesin of Escherichia coli. *Mol. Microbiol.* **56**, 1648–1663 (2005).

64. Dubey, A. K. *et al.* CsrA Regulates Translation of the Escherichia coli Carbon Starvation Gene,
cstA, by Blocking Ribosome Access to the cstA Transcript. *J. Bacteriol.* **185**, 4450–4460 (2003).

65. Holmqvist, E., Li, L., Bischler, T., Barquist, L. & Vogel, J. Global Maps of ProQ Binding *In Vivo*
Reveal Target Recognition via RNA Structure and Stability Control at mRNA 3′ Ends. *Mol. Cell* **70**,
971-982.e6 (2018).

66. Smirnov, A., Wang, C., Drewry, L. L. & Vogel, J. Molecular mechanism of mRNA repression in trans
by a ProQ-dependent small RNA. *EMBO J.* **36**, 1029–1045 (2017).

67. Irastortza-Olaziregi, M. & Amster-Choder, O. Coupled Transcription-Translation in Prokaryotes: An
Old Couple With New Surprises. *Front. Microbiol.* **11**, (2021).

68. Zhu, M. & Dai, X. Maintenance of translational elongation rate underlies the survival of Escherichia
coli during oxidative stress. *Nucleic Acids Res.* **47**, 7592–7604 (2019).

69. Zhu, M. & Dai, X. Bacterial stress defense: the crucial role of ribosome speed. *Cell. Mol. Life Sci.*
77, 853–858 (2020).

70. Vogel, U. & Jensen, K. F. The RNA chain elongation rate in Escherichia coli depends on the growth
rate. *J. Bacteriol.* **176**, 2807–2813 (1994).

71. Zhu, M., Mori, M., Hwa, T. & Dai, X. Disruption of transcription–translation coordination in
Escherichia coli leads to premature transcriptional termination. *Nat. Microbiol.* **4**, 2347–2356
(2019).

72. Caron, M.-P. *et al.* Dual-acting riboswitch control of translation initiation and mRNA decay. *Proc.*
Natl. Acad. Sci. **109**, E3444–E3453 (2012).

73. Scheithauer, B. W. & Bruner, J. M. Central nervous system tumors. *Clin. Lab. Med.* **7**, 157–179
(1987).

74. Deng, X. *et al.* Widespread occurrence of N6-methyladenosine in bacterial mRNA. *Nucleic Acids*
Res. **43**, 6557–6567 (2015).

75. Lee, J. M., Hammarén, H. M., Savitski, M. M. & Baek, S. H. Control of protein stability by post-
translational modifications. *Nat. Commun.* **14**, 201 (2023).

76. Cain, J. A., Solis, N. & Cordwell, S. J. Beyond gene expression: The impact of protein post-
translational modifications in bacteria. *J. Proteomics* **97**, 265–286 (2014).

77. Manteca, A., Ye, J., Sánchez, J. & Jensen, O. N. Phosphoproteome Analysis of Streptomyces
Development Reveals Extensive Protein Phosphorylation Accompanying Bacterial Differentiation.
J. Proteome Res. **10**, 5481–5492 (2011).

78. Bremer, E. & Krämer, R. Responses of Microorganisms to Osmotic Stress. *Annu. Rev. Microbiol.*
73, 313–334 (2019).

79. Mueller, E. A. & Levin, P. A. Bacterial Cell Wall Quality Control during Environmental Stress. *mBio*
11, 10.1128/mbio.02456-20 (2020).

Figure 1. Transcriptomic and proteomic analysis of three human bacterial pathogens
under ten infection relevant stress conditions. A) Schematic presentation of data
collection and data generation for gene expression on mRNA and protein level under stress
conditions. **B)** Percentages of CDSs detected by transcriptomics and proteomics for the three
species. **C**) Distribution of transcript abundance across all conditions (light blue) in the three
species; yellow indicates the fraction of proteins detected, while dark blue represents the
transcripts for which no corresponding protein was detected.

Figure 2. mRNA-protein levels show differing degree of statistically significant positive
correlation under different stress conditions and control condition in A) *S. enterica*
Typhimurium, **B)** *S. aureus* and **C)** *Y. pseudotuberculosis*. mRNA and protein expression
levels were taken as log2 transformed TPM and RI values, respectively. The expression
values of genes that are detected in all conditions, as mRNA and protein, were used in this
analysis. Pearsson correlation coefficient (R) shows positive correlation under all conditions
in the three species. *p* values were calculated using Student's t-test.

725
726

 Figure 3. Essential genes of the three pathogens have higher mRNA and protein expressions with higher correlation and lower variance across the 11 conditions in comparison to all genes. A) Mean mRNA expression frequency of essential and non-essential genes in the 11 conditions. **B)** Mean protein expression frequency of essential and non-essential genes in the 11 conditions. **C)** Essential genes and all genes mRNA-protein level correlation under 11 conditions for the three species. **D)** Frequency of standard deviation of mRNA expression under 11 conditions for the three species. **E)** Frequency of standard deviation of protein expression under 11 conditions for the three species. mRNA-protein level correlations were measeured with Pearsson correlation coefficient (R). p values were calculated using a Wilcoxon rank sum test.

Figure 4. Degree of stress response lowers mRNA-protein level correlations in the three species. A) mRNA-protein level correlations for conditions exposed to stressors for 10 minutes, 20 minutes, 30 minutes, 4 hours, and longer than 4 hours in the three species. **B)** Percentages of differentially expressed genes (DEGs) and proteins (DEPs) under the 10 stress conditions in comparison to unexposed control conditions for the three species. **C)** Scatter plot showing negative correlation between percentages of DEGs (top) and DEPs (bottom) and mRNA-protein level correlation. Correlations were measured with Pearsson correlation coefficient (R). *p*-values were calculated using Student's t-test.

Figure 5. Stress response induces different patterns of discordant mRNA and protein level regulation highlighting key role for translation. A) Scatter plots with fold changes of mRNA and protein levels for each the 10 stress conditions in comparison to control for the three species. The 6 gene clusters whose mRNA and protein level have different discordant regulation patterns are shown in different colors. **B)** mRNA and protein log2 fold change patterns of the 6 discordantly regulated gene clusters under 10 stress conditions for the three species. The number of discordantly regulated genes in each cluster are indicated with 'n' **C)** GO biological process enrichment analysis of 6 discordantly regulated gene clusters for the three species.