Bacterial Stress Responses Lower mRNA-Protein Level Correlations

Sena G. Süer[1,2,3,4†], Jérôme Arnoux[1,2,3,4†], Yi Y. Lim[1,2], Ganeshwari Dhurve[1,5], Rabia Şen[4,6], Cemal Erdem[4,6], André Mateus[3,4,5,7] and Kemal Avican[1,2,3,4,8]*

1. Department of Molecular Biology, Umeå University, Umeå, 90187, Sweden.
2. Integrated Science Lab (IceLab), Umeå University, Umeå, 90187, Sweden.
3. Umeå Centre for Microbial research (UCMR), Umeå University, Umeå, 90187, Sweden.
4. Science for Life Laboratory, Umeå University, Umeå, 90187, Sweden.
5. Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå, 90187, Sweden.
6. Department of Medical Biosciences, Science for Life Laboratory, Umeå University, Umeå, 90187, Sweden.
7. Department of Chemistry, Umeå University, Umeå, 90187, Sweden.
8. Lead contact

†These authors share first authorship.

*Correspondence: Kemal Avican

Email: kemal.avican@umu.se

Author Contributions: K.A. designed research; S.G.S., JA, Y.Y.L., G.D., R.Ş., C.E., A.M and K.A performed research; S.G.S., and G.D. contributed new reagents/analytic tools; S.G.S., J.A., Y.Y.L., R.Ş., C.E., A.M and K.A. analyzed data; and S.G.S., J.A., and K.A. wrote the paper with input from G.D., R.Ş., C.E., A.M

Competing Interest Statement: The authors declare no competing interests.

Classification: BIOLOGICAL SCIENCES, Microbiology; Systems Biology

Keywords: mRNA-protein correlation, infection, stress response, pathogen.

This PDF file includes:
Main Text
Figures 1 to 6
Tables 1 to 1

1 Abstract

Diverse bacterial pathogens have evolved complex regulatory mechanisms to adapt to various
environmental stresses during infection. The uncertainty in mRNA-protein levels in response to
environmental stressors complicates our understanding of bacterial physiology and their adaptation
to stressful environments. To examine this issue, we have integrated transcriptomics and
proteomics data on three human bacterial pathogens *Salmonella enterica* Typhimurium, *Yersinia*
pseudotuberculosis, and *Staphylococcus aureus* under ten infection-relevant stress conditions. We
observed positive correlations between mRNA and protein levels, which were decreased under
different stress conditions. Essential genes exhibited higher expression levels with lower variation
across the conditions and stronger mRNA-protein correlations compared to non-essential genes,
highlighting their critical role in bacterial adaptability and survival. Moreover, we identified a
substantial number of genes with stress-induced non-correlating mRNA-protein levels, especially
under conditions triggering strong stress responses. Particularly this level was dramatically lowered
for osmotic stress specific genes affected by impaired translational activity under osmotic stress.
Our findings highlight the prevalence of non-correlating mRNA-protein levels and the potential role
of post-translational modifications in modulating protein levels in response to environmental
stressors during infection. This study provides a comprehensive framework for integrating
transcriptomics and proteomics data and identifies potential gene products that might significantly
impact the ability of diverse bacterial pathogens to adapt to hostile infection environments.

20 Significance Statement

Understanding how bacteria adapt to host environments is crucial for combating infections. We
employed an integrative transcriptomics and proteomics approach to investigate mRNA-protein
correlations in three clinically relevant pathogens under ten infection-relevant stress conditions. We
identified genes whose mRNA-protein relationships are significantly disrupted by specific stressors.
This study provides a deeper understanding of mRNA-protein level uncertainty, coupling it to stress
responses known to trigger critical post-transcriptional and post-translational regulations. These
findings help reveal novel mechanisms of bacterial adaptation and pathogenesis. By providing a
comprehensive, multi-species dataset, this study serves as a foundational resource for infection
biology, driving future advancements in understanding complex regulatory networks and identifying
potential antimicrobial targets.

31 Main Text
32 Introduction

Bacterial pathogens have evolved complex mechanisms to adapt to different environmental
stresses in the host such as nutrient limitations, temperature or pH shifts, and antimicrobial
challenges during infection. They respond to deleterious effects of such stressors via stress
response mechanisms, involving complex gene regulatory networks, for survival. Therefore,
comprehensive understanding of bacterial stress response mechanisms is critical for development
of novel therapeutic approaches to resolve bacterial infections. One useful approach is through
investigation of global transcriptomes of diverse bacterial pathogens under a variety of stress
conditions that mimic host environments. This provides information on regulation across diverse
stress conditions that could be used in attempts to reveal complex gene regulations and identify
key stress response genes. We previously performed a large-scale transcriptomic study of 32
human bacterial pathogens, PATHOgenex RNA Atlas (1), aiming to understand bacterial stress
responses under 11 *in vivo* mimicking stress conditions. The study revealed a dynamic regulation
of gene expression under different stress conditions and suggested a set of genes that has potential
to be candidate as new antimicrobial targets. Co-PATHOgenex (2), the continuation of the
PATHOgenex study, identifies stress-specific stimulons by generating gene co-expression

modules. These stimulons represent sets of genes with unique expression patterns under specific
stressors, providing insights into the stress adaptation to different stressors.

Yet the main functional units shaping the cellular response are proteins and there is, generally, a
weak correlation between mRNA and protein levels (3–11). This highlights the complex nature of
gene regulation and the diverse post-transcriptional factors affecting protein stability, and
abundance (12). It is known that transcription and translation could be uncoupled constitutively or
under certain conditions, such as stress conditions, at a global or at a gene-specific level (13, 14)
Therefore, the relationship between mRNA and protein levels can vary depending on the
conditions, and this variability has been the subject of extensive research and ongoing debate (12,
14–16). Nevertheless, mRNA levels have been shown to be positively correlated with protein levels
in many different organisms (8, 17–25). Linking protein expression to gene expression can help to
identify key proteins in response to specific stimuli or conditions and highlight mechanisms that
drive bacterial adaptation, evasion of the immune system, and disease development (26–30). This
is crucial for developing more effective diagnostic tools, treatment strategies, and vaccines for
bacterial infections (31, 32). Therefore, integrated transcriptomic and proteomic studies have
emerged as powerful tools for understanding the complex molecular mechanisms underlying
bacterial pathogenesis (28). Despite the numerous studies integrating transcriptome and proteome
data, efforts to identify genes where mRNA and protein levels *do not* correlate are relatively limited
(17, 33). Focusing on such genes could offer valuable insight into the complex post-transcriptional
and post-translational regulatory mechanisms that escape detection by analyses addressing
correlated gene-expression patterns.

In this study, we employed an integrative transcriptomic and proteomic analysis of *Salmonella*
enterica Typhimurium, *Yersinia pseudotuberculosis*, and *Staphylococcus aureus* under ten
infection-relevant stress conditions to investigate mRNA-protein level correlations. We revealed
that mRNA-protein correlations significantly varied under different stress conditions for the three
species. Essential genes consistently showed higher mRNA and protein expression levels and
stronger correlations, emphasizing their importance in bacterial survival. Stress conditions such as
hypoxia and nutritional downshift exhibited the lowest correlations, suggesting that post-
transcriptional and post-translational regulation play a significant role under these conditions. We
identified plentiful genes that showed non-correlating mRNA-protein levels in the three species, as
well as sets of genes that were species specific, particularly under conditions eliciting strong stress
responses. Among all conditions tested, osmotic stress stood out as exhibiting the most
pronounced mRNA-protein decoupling. Using the MOBILE (34) framework, we characterized the
osmotic stress stimulon and linked the observed mRNA-protein decoupling to a specific impairment
of translation initiation.

Results

Transcriptomic and proteomic analysis of three human bacterial pathogens under ten
infection relevant stress conditions

We analyzed transcriptomic and proteomic datasets of three human bacterial pathogens including
Gram-negative *S. enterica* Typhimurium SL1344, *Y. pseudotuberculosis* YPIII and Gram-positive
S. aureus MSSA476. We used transcriptomes of the three pathogens under 10 infection relevant
stress conditions together with unexposed control condition from PATHOgenex RNA Atlas (1) and
complemented it with mass-spectrometry proteomic profiling under the same stressors (**Figure**
1A). As mass spectrometry-based proteomics often yields a lower coverage than RNA-seq, we
compare detected transcripts and proteins across all conditions in three pathogens, focusing on
protein coding sequences (CDSs) in the analysis. While almost all CDSs were detected by RNA-
seq for the three pathogens (98% for *S. enterica* and *Y. pseudotuberculosis*, and 90% for *S.*
aureus), proteomics could detect between 48% of the CDSs for *S. enterica* Typhimurium and *Y.*

pseudotuberculosis, and 32% for *S. aureus* (**Figure 1B**). Detected proteins covered nearly the full
spectrum of mRNA expressions, with a bias towards highly expressed mRNAs across the
conditions for the three pathogens (**Figure 1C**).

mRNA-protein level correlation under stress

The correlation between mRNA and protein levels has been previously explored in bacteria under
individual growth conditions (18, 33, 35, 36). Our data allows us to assess whether mRNA levels
can serve as an accurate indicator of protein levels in three human bacterial pathogens. To
evaluate how different stressors affect the mRNA-protein level correlation, we calculated the
intensity-Based Absolute Quantification (iBAQ) (15) **(Supplementary Table 1)** values for protein
levels and compared to mRNA levels in Transcript Per Million (TPM) (37) for every stress condition
including the unexposed control condition (1). We have previously shown that the three biological
replicates of the RNA-seq samples clustered together (1). For the proteome dataset generated in
this study, we conducted pairwise comparisons of the three replicates of each condition and
computed Pearson correlation coefficient values (R-value). The R-value for pairwise comparison
within the replicates of the same condition varied from 0.89-0.96 for the three pathogens
(Supplementary Figures 1-3) showing strong similarities between the replicates. We then
measured the correlation between mRNA and protein levels by calculating Pearson correlation
coefficient values of the log2 transformed mean TPM and log2 transformed mean iBAQ of genes
for each condition and pathogens **(Figure 2A, B, and C)**. For these measurements, we limited the
analysis to the genes that are detected as mRNA and protein across all tested conditions. We
observed that correlations between mRNA and protein levels under 11 conditions ranged from
0.48-0.68 for *S. enterica* Typhimurium, 0.54-0.72 for *S. aureus* and 0.59-0.76 for *Y.*
pseudotuberculosis **(Figure 2A, B, and C)**. These correlation coefficients are comparable to those
reported in previous studies across diverse biological systems (19, 20, 33, 36, 38). The variation in
mRNA–protein correlation across stress conditions suggests that distinct molecular mechanisms,
activated by each stressor, might modulate translational or post-translational regulations leading to
differences.

Essential genes have higher mRNA-protein level correlations in the three species

The positive correlation between mRNA-protein levels for all detected genes prompted us to
investigate subsets of genes with different correlations levels. Essential genes have been shown
to have stronger correlations in comparison to non-essentials for *Pseudomonas aeruginosa* during
different growth phases (17). Therefore, we sought to investigate mRNA-protein level correlation
profiles of essential genes in the three species. Of the 354, 341, and 445 previously identified
essential genes for *S. enterica* Typhimurium (39) *S. aureus* (40) and *Y. pseudotuberculosis* (41)
respectively, we detected 300 (85%), 280 (82%), and 335 (75%) essential genes in our
transcriptome and proteome datasets. We found that mean expression levels of mRNAs and
proteins were significantly higher for essential genes in comparison to non-essentials for the three
species (**Figure 3A and B).** This was true for the expression levels in every condition
(Supplementary Figures 4 and 5). Moreover, we observed that essential genes have higher
mRNA-protein level correlation than all detected genes for all three species (**Figure 3C**), except for
S. enterica Typhimurium and *Y. pseudotuberculosis* under nutritional downshift (Nd) stress
condition. This distinct uncoupling in the Gram-negative species hints at divergent regulatory
priorities during sudden starvation. As previously observed in other species (42, 43), essential
genes showed significantly lower variance on both mRNA and protein expression across the 11
conditions for the three species (**Figure 3D and E**). These findings show that essential genes of *S.*
enterica Typhimurium, *S. aureus*, and *Y. pseudotuberculosis* have high and stable mRNA and
protein expression under a wide range of stress conditions with robust mRNA-protein level
correlations.

Stress responses differentially affect mRNA-protein correlation

The variation in correlation under different stress conditions prompted us to investigate possible underlaying reasons. The mRNA-protein levels in Hypoxia (Hyp) and Nd showed the lowest correlation across three species **(Figure 2A, B, and C)**. Those stress conditions differ from others in the length of the exposure time: 4 hours for Hyp and 30 minutes for Nd, against 10 minutes for others **(Figure 1A)** besides condition itself. To determine whether exposure duration influences mRNA-protein level correlation, we examined the relationship between correlation and exposure time across all conditions. We observed a decline in mRNA-protein level correlation associated with longer exposure times (R = -0.53; p = 0.002**; Figure 4A)**. However, this effect may not be solely dependent on time; robust, stressor-specific responses triggered by conditions like Nd, stationary phase (Sp), and Temperature (Tm) likely act as confounding factors that drive this decrease.

To assess whether the degree of stress responses through differentially regulated mRNAs and proteins has an impact on mRNA-protein level correlations, we used differentially expressed genes (DEGs) and differentially expressed proteins (DEPs) for each condition in the three species by using unexposed samples as control. Before identifying DEPs, we evaluated 8 normalization methods from NormalyzerDE (44), on the iBAQ protein abundance data and found that Variance Stabilizing Normalization (VSN) was the most effective in clustering the condition together (**Supplementary Figure 6**). Then we used limma (45) differential expression package for the identification of DEPs with VSN normalized data. For DEGs, we used previously reported differential expression analysis of PATHOgenex RNA Atlas (1). We employed a p-value < 0.05 and a log2-Fold Change > |0.5| as thresholds to identify DEGs and DEPs for each condition. Notably, the number of DEGs were significantly higher compared to DEPs in all conditions for the three species **(Figure 4B)**. This can be linked to the stronger regulation of mRNA levels than of protein levels as previously observed in humans (16, 46). Moreover, percentages of DEGs and DEPs were highest under Hyp, Nd, and Sp conditions for the three pathogens (**Figure 4B**). Next, we tested whether the differential expressions have an impact on mRNA-protein level correlations. To do this, we measured the association between the percentages of DEGs and DEPs and the mRNA-protein level correlations with Pearson correlation coefficient. Our results revealed a strong negative correlation between differential expressions and mRNA-protein level correlation (**Figure 4C**). This suggests that other factors, such as post-transcriptional and post-translational regulation triggered by different stressors, might be influencing protein levels alongside the mRNA expression.

Identification of genes whose mRNA-protein levels are discordantly regulated under stress conditions

The decreased correlation between mRNA-protein levels with increased number of DEGs and DEPs under specific stress conditions led us to investigate the genes whose mRNA-protein levels are discordantly regulated when the bacteria are exposed to stress conditions. We clustered genes into 6 groups (Error! Reference source not found.Error! Reference source not found.) corresponding to distinct combinations of differential expression patterns at the mRNA and protein levels in at least one stress condition (**Figure 5A and B and Supplementary Table 2**). In agreement with lowest correlation between mRNA-protein levels observed under Hyp, Nd, and Sp conditions (**Figure 2A-C**), discordantly regulated genes were mostly found under those conditions for the three species (**Supplementary Figure 7**). Noteworthy, Clusters 1 and 6, where mRNAs were regulated but not proteins, had the highest number of discordantly regulated genes. This is in accordance with higher number of differentially regulated mRNAs (DEGs) compared to differentially regulated proteins (DEPs) in all conditions for the three species (**Figure 4B**).

Discordantly regulated genes are enriched in translation

To investigate whether the genes exhibiting discordant mRNA and protein levels under stress conditions are enriched in certain feature, we performed Gene Ontology (GO) (47) enrichment analysis. Interestingly, we found that biological processes such as translation, RNA processing, RNA modification, phosphorylation, and cell division, among other biological processes were commonly enriched for the three species, particularly when mRNA levels were changed but there was no change at the protein level (**Supplementary Figure 8 and Supplementary Table 3**). This suggests that genes encoding subunits of translation machinery together with other components of translation such as tRNA and rRNA were discordantly regulated under stress conditions for the three pathogens. In detail, proteins in Cluster 1 have stable levels despite mRNA being downregulated, likely due to increased protein stability. Supporting this, we observed a cross-species enrichment of proteins involved in protein folding and post-translational processing pathways. In contrast, proteins in Cluster 6 have stable levels despite mRNA being upregulated, suggesting reduced translation or increased proteolysis, which is enriched for *S. enterica* Typhimurium. Furthermore, we observed an enrichment in protein transport and secretion pathway for all species within this cluster. This indicates that the observed protein stability is not only due to translation and degradation activity, but also to a rapid export from the cell. Moreover, genes encoding proteins involved in protein phosphorylation were commonly enriched for the three species in Cluster 4 and 5. In cluster 4, protein levels were upregulated despite stable mRNA abundance, whereas Cluster 5 exhibited mRNAs upregulation and proteins downregulation (**Supplementary Figure 8)**. Notably, Cluster 4 was also commonly enriched for lipid metabolism across the three species. This indicates presence of post-transcriptional regulations such as regulation via small regulatory RNAs involved in carbon metabolism (48, 49) or post-translational modifications which have high prevalence in bacterial metabolic enzymes (50).

mRNA-protein level stress-specific stimulon genes under stress

We previously identified stress-specific stimulon genes as sets of genes that are uniquely co-regulated in response to distinct stress conditions (2). To assess whether these stimulon genes exhibit distinct mRNA-protein level correlations compared to other genes, we analyzed mRNA and protein differential expression data across all genes and stress-specific stimulons. No significant differences in mRNA-protein level correlations were observed between stimulon genes and other genes across most conditions except in the case of the osmotic stress-specific stimulon (**Figure 6A**). Notably, genes within this stimulon showed a markedly lower mRNA-protein level correlation across all three pathogens studied.

To investigate this further, we used MOBILE framework (34) to identify osmotic stress specific associations between mRNA and protein levels. Through repeated Lasso regression between paired omics layers, MOBILE enables the identification of condition-specific associations without requiring prior knowledge. The transcriptomic and proteomic datasets with all conditions were used as an input for MOBILE, yielding a list of statistically robust associations between proteins and mRNAs. Next, the osmotic stress (Oss) condition data from both datasets were excluded from the input, and MOBILE is re-run to obtain condition-dependent associations list by comparing the all-data run to the latter. Similarly, the control (Ctrl) condition data is held out, and MOBILE re-run yielded another list of associations to identify CTRL-data dependent associations (**Figure 6B**). We identified Oss specific associations by over representation analysis (ORA), comparing the osmotic stress-dependent and control-dependent association lists. Using Oss specific associations gene list, we ran a GO enrichment analysis. Interestingly, translation-related processes and components of the translational machinery were enriched only for *Y. pseudotuberculosis* (**Figure 6C**). This suggests a functional link between osmotic stress-specific mRNA-protein decoupling and translational regulation. In contrast, *S. aureus* showed no significant enrichment in osmotic-specific

pathways. This might reflect the inherent physiological resilience of Gram-positive bacteria and
metabolic adaptation to osmotic stress (51).

Based on this finding, we hypothesized that the reduced mRNA-protein correlation observed for
osmotic stress-specific stimulon genes might result from impaired translation under osmotic stress.
To test this, we performed polysome profiling on *Y. pseudotuberculosis* exposed to Oss for 10
minutes, using unexposed bacteria collected at matched time points as controls The polysome
profiles showed no notable differences between the osmotic stress and control conditions, with a
clear retention of polysome peaks (**Figure 6D**). This indicates that under osmotic stress, ribosomes
are still successfully recruited to and load onto transcripts. However, intact polysomes do not
necessarily equate to active protein synthesis, as ribosomes may be stalled mid-translation.
Therefore, global translational activity may be insufficient to meet the heightened translational
demand triggered by the strong transcriptional induction of osmotic stress-specific stimulon genes.

To further characterize the translational response, translation efficiency was assessed using a
puromycin incorporation assay across osmotic stress, low iron, and unexposed control conditions.
Chloramphenicol-treated bacteria served as a translationally inactive control. Since puromycin
causes a premature translation termination, its incorporation serves as a direct proxy for active
translation. The results showed that translational activity was significantly reduced under osmotic
stress compared to both unexposed control and low iron conditions, indicating that osmotic stress
specifically impairs translation activity (**Figure 6E and Supplementary Figure 9**). Therefore, the
reduced mRNA-protein correlation for osmotic stress-specific stimulons is not surprising given the
overall reduction in translational activity. These stimulon genes exhibit significant changes in mRNA
abundance during stress. However, because the translational machinery is impaired, it cannot
efficiently process or respond to these altered transcript levels. As a result, the correlation between
mRNA and protein decreases. Notably, this decoupling is specific to osmotic stress and was not
observed under low iron conditions.

Discussion

We conducted a comprehensive analysis by integrating transcriptomic and proteomic datasets
of *S. enterica* Typhimurium, *Y. pseudotuberculosis*, and *S. aureus* under 10 infection-relevant
stress conditions, as well as a control condition during exponential growth. The inclusion of diverse
pathogens, which are distantly related yet possess conserved genes, enabled us to identify both
common and species-specific mRNA-protein relationships across a range of conditions mimicking
host environments. Consistent with prior findings in other organisms, our analyses indicate a
positive correlation between mRNA and protein levels across the three bacterial pathogens.
However, this correlation is significantly reduced under stress conditions, particularly those that
trigger a strong stress response.

This reduction is not surprising, as stress response proteins might be involved in post-
transcriptional or post-translational regulations that can decouple mRNA and protein levels. Such
regulations can occur through the stabilization or destabilization of mRNA and proteins as a
response to the stress. RNA-binding proteins such as CsrA and ProQ can bind to target mRNAs
under specific conditions, either stabilizing or destabilizing them, or preventing their translation (52–
57). Furthermore, the discoordination of transcription and translation elongation rates under stress
conditions may contribute to the decrease in mRNA-protein level correlation (58). For example, in
Escherichia coli exposed to oxidative stress, there is a significant induction of mRNA expression of
the OxyR regulon, which is crucial for survival under oxidative stress, while the translation
elongation rate significantly slows down (59, 60).

In contrast, under nutrient-limited conditions, there has been tight coordination between
transcription and translation (61, 62). This suggests that the reduced mRNA-protein level

correlation in our nutritional downshift condition might not be solely due to the discoordination of
transcription and translation but rather other regulatory mechanisms. Conformational changes in
riboswitches upon a change in the environment can also lead to reprogrammed mRNA translation
and degradation, as is the case for the lysC riboswitch in the presence and absence of lysine (63).

Even though mostly studied in eukaryotes, post-transcriptional nucleotide modifications triggered
by stress conditions can also have an impact on mRNA-protein level as it is shown for increased
level of m^6A methylation under hypoxic for human cancer cells (64). On the other hand, m^6A
methylation was shown to be lowered under increased temperature for *P. aeruginosa* (65). Similarly
post-translational modifications such as methylation, phosphorylation, acetylation, and sumoylation
can directly impact protein stability or indirectly through triggered signaling cascade in both
eukaryotes and prokaryotes (66, 67). Even though protein stability was not specifically investigated,
it was shown that cellular protein phosphorylation was increased during *Streptomyces*
differentiation to form spores, which is induced by stress conditions (68).

To assess whether the mRNA-protein discrepancies observed for DEGs are a common feature
across species, we identified conserved DEGs across the three pathogens and compared their
expression patterns. These conserved genes exhibited similar non-correlating expression patterns
under the same stress conditions, suggesting that post-transcriptional and post-translational
modifications affecting these genes may be conserved across species. This highlights the
evolutionary significance of these regulatory mechanisms in bacterial adaptation to stress.

Among all conditions investigated for stress stimulons, osmotic stress exhibited the most
pronounced mRNA-protein decoupling. This impairment likely stems from a compounding effect on
both translation initiation and elongation. Initially, the rapid structural remodeling and cell
depolymerization required to manage turgor pressure (69, 70) can physically disrupt membrane
integrity and the efficiency of transmembrane transporters. This disruption can lead to deficient
amino acid uptake, such as methionine, which would directly impair translation initiation due to the
requirement of formylmethionine at the start of bacterial protein (21, 22). Concurrently, our
experimental validation points to a secondary bottleneck during active synthesis. The maintenance
of intact polysomes alongside significantly reduced puromycin incorporation indicates that
ribosomes also stall during elongation. Together, these dual impairments in initiation and elongation
prevent the translational machinery from efficiently processing the highly induced stress transcripts,
resulting in the observed mRNA-protein decoupling.

Our study underscores the importance of integrating transcriptomic and proteomic data to gain a
holistic understanding of bacterial responses to infection-relevant stressors. The observed
discrepancies between mRNA and protein levels emphasize the need for caution when inferring
protein abundance solely from transcriptomic data. The precise post-transcriptional and post-
translational mechanisms contributing to the observed mRNA-protein discrepancies remain to be
elucidated, as the currently available datasets do not allow their direct identification, representing
an important avenue for future investigation.

In conclusion, our findings provide a comprehensive overview of mRNA-protein relationships under
stress conditions in three bacterial pathogens, laying the groundwork for further investigations into
the regulatory mechanisms mediating bacterial adaptation and survival in hostile host
environments. The multi-species, multi-condition nature of this dataset furthermore represents a
valuable resource for training predictive models of protein abundance from mRNA levels,
particularly under stress conditions where the mRNA-protein relationship deviates substantially
from baseline.

Materials and Methods

Bacterial strains and growth conditions

Salmonella enterica subsp. enterica serovar Typhimurium SL1344, *Staphylococcus aureus*
MSSA476, and *Yersinia pseudotuberculosis* serotype O:3 (strain YPIII) species were selected from
PATHOgenex database[1] to analyze transcriptomic and proteomic data.

Salmonella enterica SL1344 and *Staphylococcus aureus* MSSA476 strains used in this study did
not carry antibiotic resistance markers and were grown in Luria–Bertani (LB) medium (BD Difco™
LB Agar, Lennox) at 37 °C with shaking. The kanamycin-resistant wt *Y. pseudotuberculosis*
(YPIII/pIBX) strain was used in this study. Unless otherwise stated, the wild-type strain was grown
in LB medium supplemented with 50 µg/mL kanamycin at 26°C with shaking.

Stress exposure and sample collection

Polysome profiling

The wt *Y. pseudotuberculosis* strain was grown overnight in LB, subcultured to an initial optical
density (OD_{600}) of 0.05 in 200 mL of LB. When the culture reached an OD_{600} of 0.3, the culture was
divided into three 50 mL aliquots, each subjected to one of the following conditions for 10 minutes:
(1) unexposed control, (2) osmotic stress (0.5 M NaCl), or (3) low-iron stress (250 µM 2,2′-Bipyridyl;
Sigma-Aldrich). Cells were harvested by filtration, immediately flash-frozen in liquid nitrogen, and
stored at -80°C until lysis.

Frozen cells were lysed by cryogenic grinding in liquid nitrogen using aluminum oxide. Lysis buffer
was added during grinding, and the lysate was allowed to thaw gradually throughout the process.
The mixture was transferred into a pre-cooled 1.5 mL reaction tube and immediately placed on ice.
Lysates were centrifuged at 18,000 x *g* for 5 minutes at 4°C. The resulting supernatants (~400 µL)
were divided into 50 µL aliquots in new, pre-cooled 1.5 mL tubes and stored at −80°C. Total RNA
concentration was quantified using the Qubit fluorometer. Each aliquot was used only once for
downstream polysome profiling.

Linear 15–45% sucrose gradients were prepared using 10% and 50% (w/v) sucrose solutions in
50 mM Tris-HCl (pH 7.5) and supplemented with 100 mM NH_4Cl, 10 mM $MgCl_2$, and 6 mM β-
mercaptoethanol. Solutions were sterile filtered and gradients generated using the Gradient Master
(BioComp Instruments) following the manufacturer's preprogrammed settings (SW60Ti rotor, short
sucrose program, w/v 1st). Prepared gradients were stored at 4°C overnight prior to use.

For polysome analysis, 50 µg of cell lysate was carefully layered onto each gradient without
disturbing the interface. Samples were centrifuged at 40,000 rpm for 2 hours and 30 minutes at
4°C in an Optima L-100K ultracentrifuge (Beckman Coulter) equipped with an SW60Ti rotor.
Immediately following centrifugation, the sucrose gradients were carefully placed on ice to prevent
disturbance, and continuous polysome absorbance profiles at 254 nm were recorded using the
BioComp gradient fractionation system complemented with a TRIAX flow cell. 14 discrete fractions
were sequentially collected from the top to the bottom of each gradient using the automated fraction
collector, following the manufacturer's instructions and stored at −80°C until further analysis.

Puromycin assay

To assess active translation, the wt strain was cultured as described above for overnight and
subculture preparation. Upon reaching exponential phase (OD_{600} of 0.4), the culture was distributed
into four separate flasks representing the following treatment conditions: (1) chloramphenicol (50
µg/mL), (2) unexposed control, (3) osmotic stress (0.5 M NaCl), or (4) low-iron stress (250 µM 2,2′-

Bipyridyl; Sigma-Aldrich). Cultures grew with their respective conditions for 10 minutes.
Subsequently, to abruptly freeze and capture the translational landscape, both 10 µg/mL puromycin
and 200 µg/mL chloramphenicol were simultaneously added to the culture, followed by an
additional 10-minute incubation period prior to cell lysis. The experiment was performed in four
independent biological replicates.

To extract proteins, 20 mL of bacterial culture was collected from each treatment group and cells
were immediately harvested by centrifugation at 4,000 × g for 10 minutes at 4°C. The resulting cell
pellets were washed twice with phosphate-buffered saline and resuspended in 1 ml of lysis buffer
(100 mM Tris-HCl, 10 mM EDTA, 1x cOmplete™ protease inhibitor cocktail (Roche)). The cell
suspensions were lysed on ice using sonication for nine 20 second cycles, each cycle comprised
20 second sonication step separated by 30 second intervals with 10% amplitude. Following
centrifugation at 13,000 x g for 10 minutes at 4°C, the supernatant was collected and stored at -
20°C. Protein concentrations were defined using QuBit Protein Assay Kit (Catalog Number:
Q33211).

Immunoblotting

Protein concentrations were fixed across samples, and proteins were mixed in 1x sample buffer
(2% SDS, 62.5 mM Tris-HCl, 20% glycerol, 0.05% bromophenol blue, 1% β-mercaptoethanol). 17
µl of proteins were separated by sodium dodecyl sulfate (SDS)-PAGE on 4-15% Mini-PROTEAN
TGX gels using 80V for 20 minutes, followed by 120V for 1 hour. SDS gel was transferred onto
methanol activated Immobilon-P PVDF membrane (Millipore) by semidry electroblotting using Bio-
Rad Trans-Blot SD Semi-Dry Transfer Cell (70V for 1 hour 10 minutes). Membranes were blocked
with 5% milk in 1x Tris-buffered saline with Tween (TBST) and probed with Anti-Puromycin at
1:10,000 dilution (Sigma-Aldrich, MABE343, clone 12D10) primary antibody in 1x TBST, overnight
at 4°C. Following each antibody incubation, the membranes were washed three times for 10
minutes with 1x TBST. Membranes were incubated with secondary antibody: monoclonal Sheep
Anti-Mouse IgG - Horseradish Peroxidase at 1:7,000 dilution (Cytiva, NA931V) in 1x TBST for 1
hour at room temperature. The final blot was developed by chemiluminescence (Immobilon
Western Chemiluminescent HRP Substrate, Millipore) for 5 minutes, and the bands were detected
with Amersham Imager 680. Band intensities were quantified using ImageJ (version 1.54p).

Transcriptomic data retrieval and proteomic data generation

The transcriptome data published for *S. enterica*, *Y. pseudotuberculosis*, and *S. aureus* were
obtained from Gene Expression Omnibus (GEO) (71) with accession number GSE152295.

Proteomics data was obtained by exposing the three bacterial species to the same stress
conditions as mentioned in Avican *et al.* (1), with unexposed control condition and excluding
virulence inducing condition. After stress exposures, 3 ml of bacterial cultures were pelleted with
9000 RPM for 2 minutes centrifugation and cell pellets were lysed in 200 µl 2% SDS. Thereafter, 5
µg of each sample were reduced with 20 mM TCEP. Samples were digested with a modified sp3
protocol (72), as previously described (42). Briefly, samples were added to a bead suspension (10
µg of beads (Sera-Mag Speed Beads, 4515-2105-050250, 6515-2105-050250) in 10 µl 15% formic
acid and 30 µl ethanol) and incubated shaking for 15 min at room temperature. Beads were then
washed four times with 70% ethanol. Proteins were digested overnight by adding 40 µl of 5 mM
chloroacetamide, 1.25 mM TCEP, and 200 ng trypsin in 100 mM HEPES pH 8.5. Peptides were
eluted from the beads and were dried under vacuum. Peptides were then labelled with TMT11plex
(Thermo Fisher Scientific), pooled and desalted with solid-phase extraction using a Waters OASIS
HLB µElution Plate (30 µm). Samples were fractionated into 48 fractions on a reversed-phase C18
system running under high pH conditions, with every sixth fraction being pooled together. Samples
were analyzed by LC-MS/MS using a data-dependent acquisition strategy on a Thermo Fisher
Scientific Vanquish Neo LC coupled with an Thermo Fisher Scientific Orbitrap Exploris 480. Raw

files were processed with MSFragger using standard settings for TMT (73) against FASTA
databases of each organism obtained from NCBI; for *S. enterica* Typhimurium SL1344 accession
numbers NC_016810, NC_017718, NC_017719, NC_017720; for *Y. pseudotuberculosis* accession
number NC_010465; for *S. aureus* (MSSA476) accession number NC_002952. The mass
spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the
PRIDE partner repository (74) with the dataset identifier PXD055243.

Data processing and normalization

Processing of transcriptomic data

429 Transcriptomic data were imported from PATHOgenex, including statistical values such as LogFC,
430 *p*-value, and adjusted *p*-value. Moreover, the differential expression analysis was also already
431 conducted, and thus no further pre-processing was done here.

Processing of proteomic data

Raw protein files containing intensity data from TMT runs were imported into R (75) (version 4.4.2).
Data were pre-processed to retain only protein locus tag, intensities values, and razor peptide count
-defined as peptides shared across multiple proteins. Proteins were then filtered to retain only those
quantified by at least two unique peptides and exhibiting non-zero intensity values; entries with zero
intensity were considered missing data and excluded. This processing yielded one matrix per
species per run, in which each row contained a protein identifier, its total intensity, and its
corresponding razor peptide count.

To estimate protein abundance, each protein was mapped to its corresponding treatment condition.
The relative reporter ion abundance of a given protein for its treatment was divided by the total
relative reporter ion abundance as a first estimate of protein abundance. Given the large variation
in protein length observed across the dataset — ranging from 10 to 3,000 amino acids in *S. enterica*
and *Y. pseudotuberculosis*, and from 10 to approximately 2,000 amino acids in *S. aureus*
(**Supplementary Figure 10**), we applied a correction to account for size-dependent differences in
peptide yield. Since longer proteins produce more tryptic peptides upon digestion, and
consequently more ions, the initial estimate was divided by the number of expected tryptic peptides
to obtain iBAQ (15). This ensures that iBAQ reflects protein abundance independently of protein
size. Matrices from the same species were subsequently merged across runs, and proteins absent
across all replicates were excluded, yielding three species-specific matrices each containing iBAQ
values for every detected protein across the three biological replicates and their associated
treatment conditions.

To identify the most appropriate normalization strategy, 8 normalization methods were evaluated
using the NormalyzerDE (44) package (version 1.23.2): VSN, Log2-transformation, Quantile,
Global Intensity, Mean, Median, Linear Regression, and Cyclic Loess. Normalized outputs were
visualized using principal component analysis (PCA) plots generated with ggplot2 (76) (version
4.0.2) to assess replicate clustering within each stress condition. VSN consistently produced the
tightest replicate clustering across condition and was therefore applied to the three species-specific
matrices to generate VSN-normalized iBAQ values for downstream analysis.

Differential abundance analysis was performed on the normalized proteome data using the limma
(45) package (version 3.58.1) with default parameters, applying empirical Bayes moderation to
stabilize variance estimates across proteins. Differentially abundant proteins were identified for
each species to characterize the proteomic response to every stress condition, and highlight
proteins potentially involved in key regulatory mechanisms.

465	**Data analysis**
466	**Correlation analysis**
467	To assess the correlation between mRNA and protein levels, we used the log2-transformed mean
468	TPM values for mRNAs and the log2-transformed mean iBAQ values for proteins from three
469	replicates of each condition. The Pearson correlation coefficient was calculated for each condition
470	to determine the strength of the correlation. Statistical significance p-value < 0.05 was evaluated
471	using Student's t-test.
472	**Identification of gene clusters with discordant mRNA and protein level regulation**
473	To identify genes with discordant mRNA and protein level regulation under 10 stress conditions,
474	the Z-score values of mRNA from control data were combined with the Z-score values of protein,
475	calculated using limma differential expression analysis to create a single data table and visualize
476	using scatter plots separately for each species. Thereafter, the combined transcriptome and
477	proteome data were filtered with *p*-value < 0.05 to select genes whose mRNA or protein level
478	changes upon exposure to stress conditions.
479	Genes were then grouped in 6 different clusters based on the expression patterns as described in
480	Table 1. Cluster 1; mRNA downregulated in at least one stress condition but protein level is not
481	changed, Cluster 2; mRNA downregulated in at least one condition but protein is upregulated,
482	Cluster 3; mRNA level is not changed in any condition but protein is downregulated, Cluster 4;
483	mRNA level is not changed in any condition but protein is upregulated in at least one condition,
484	Cluster 5; mRNA is upregulated in at least one condition but protein is downregulated, and Cluster
485	6; mRNA is upregulated in at least one condition but protein level is not changed.
486	**Integrative omics analysis with MOBILE**
487	MOBILE pipeline (34) is used for three species (SA, SE, and YP) with 11 samples including both
488	transcriptomic and proteomic datasets. The iBAQ datasets were log2-transformed and normalized.
489	First, they were filtered to only retain the most highly variant features (top 10% for transcripts and
490	20% for proteins). Next, the MOBILE pipeline (MATLAB version R2023a) is run, using the glmnet
491	package (77) and $Y = \beta.X + \delta$ formalism, where Y matrix is the proteomics dataset and X is the
492	transcriptomics. The objective is to obtain the coefficients matrix β. δ is the y-intercept values, which
493	are almost zero by design. The regression process is repeated 10000 times with randomly selected
494	random number generation seeds to yield an ensemble of Lasso matrices. By identifying
495	coefficients that appeared in at least half of the models, a robust Lasso coefficient matrix is
496	selected. This matrix represents the stable associations between analytes across omics layers.
497	To find stress-specific interactions, MOBILE employs a "Leave-One-Group-Out" (LOGO) strategy
498	where each sample exposed to the stress is excluded from the input one-at-a-time. Here, both the
499	osmotic stress and control condition samples were subtracted, and corresponding final matrices
500	were named as OSS-IAN and CTRL-IAN matrices. IANs are the Integrated Association Networks
501	that are coalesced gene-level networks, where nodes represent genes of the matrix analytes and
502	edges stand for the non-zero coefficients.
503	**Functional annotation**
504	**Enrichment analysis**
505	Gene Ontology (GO) enrichment analyses were conducted using the clusterProfiler (78) package
506	v4.10.1 in R. Most significant enriched terms were identified using a cut-off criterion of p-
507	value<0.05. Plots showing the most significant enriched Biological Process (BP) GO terms were
508	generated using the ggplot2 package.

Over representation analysis (ORA)

The subsequent step involves comparing OSS-IAN and FULL-IAN (all condition matrix), to obtain OSS-dependent interactions. Similarly, comparing CTRL-IAN to FULL-IAN yields CTRL-dependent interactions. Next, pathway enrichment analysis (clusterProfiler version 4.14.6) was adopted for the genes list obtained by comparing OSS-dependent list to CTRL-dependent gene list. Gene Ontology gene sets curated for three species (*SA, SE, YP)* were used in gmt format with a minimum gene set size of 15 and a maximum of 500. Significantly enriched gene sets were filtered by q-value < 0.2 and p-value < 0.05.

Funding

This work was supported by Swedish Research Council (No. 2021-02466), Kempestiftelserna (JCK22-0017), and the Medical Faculty at Umeå University (FS 2.1.6-281-22) to K. Avican, by Swedish Research Council Excellence Center grant (No. 2022-06543) for the Center for Modeling Adaptive Mechanisms in Living Systems Under Stress to K. Avican

CE is funded by the Knut and Alice Wallenberg Foundation under the SciLifeLab and Wallenberg Data Driven Life Science Program (KAW 2020.0239).

Mass spectrometry analysis was enabled by support from Kempestiftelserna (grant number: JCK3126 to A.M.).

Data availability

All data necessary to reproduce the findings of this study are publicly available without restriction.

The complete dataset is available from Zenodo (79) (https://zenodo.org/records/19488690). Transcriptomic data were obtained from PATHOgenex RNA Atlas (1); Protein sequences and Mass spectrometry results are provided in the Zenodo repository.

Code Availability

All code and commands to reproduce the analysis are included in a publicly available.

A repository containing a Rnotebook with all the code to reproduce the figures and analyzes presented in this manuscript, is available on GitHub at https://github.com/avicanlab/RNAProtCorr.

All packages and versions are listed in the Renv file available in the GitHub repository.

References

1. K. Avican, *et al.*, RNA atlas of human bacterial pathogens uncovers stress dynamics linked to infection. *Nat Commun* **12**, 3282 (2021).

2. L. Fernandez, M. Rosvall, J. Normark, M. Fallman, K. Avican, Co-PATHOgenex web application for assessing complex stress responses in pathogenic bacteria. *Microbiol Spectr* **12**, e02781-23 (2024).

3. M. J. Lee, M. B. Yaffe, Protein Regulation in Signal Transduction. *Cold Spring Harb Perspect Biol* **8**, a005918 (2016).

4.   J. Bathke, A. Konzer, B. Remes, M. McIntosh, G. Klug, Comparative analyses of the
variation of the transcriptome and proteome of Rhodobacter sphaeroides throughout
growth. *BMC Genomics* **20**, 358 (2019).

5.   Y. Zhu, *et al.*, Integrative analysis of transcriptome and proteome provides insights into
adaptation to cadmium stress in Sedum plumbizincicola. *Ecotoxicol Environ Saf* **230**, 113–
149 (2022).

6.   S. Liu, *et al.*, Transcriptome and Proteome of Methicillin-Resistant Staphylococcus aureus
Small-Colony Variants Reveal Changed Metabolism and Increased Immune Evasion.
Microbiol Spectr **11**, e0189822 (2023).

7.   F. Grunberger, *et al.*, Uncovering the temporal dynamics and regulatory networks of thermal
stress response in a hyperthermophile using transcriptomics and proteomics. *mBio* **14**,
e0217423 (2023).

8.   F. Edfors, *et al.*, Gene-specific correlation of RNA and protein levels in human cells and
tissues. *Mol Syst Biol* **12**, 883 (2016).

9.   X. Feng, Q. Ma, Transcriptome and proteome profiling revealed molecular mechanism of
selenium responses in bread wheat (Triticum aestivum L.). *BMC Plant Biol* **21**, 584 (2021).

10.   M. Backman, *et al.*, Multi-omics insights into functional alterations of the liver in insulin-
deficient diabetes mellitus. *Mol Metab* **26**, 30–44 (2019).

11.   G. Chen, *et al.*, Discordant protein and mRNA expression in lung adenocarcinomas. *Mol
Cell Proteomics* **1**, 304–13 (2002).

12.   C. Vogel, E. M. Marcotte, Insights into the regulation of protein abundance from proteomic
and transcriptomic analyses. *Nat Rev Genet* **13**, 227–32 (2012).

13.   F. Gebauer, M. W. Hentze, Molecular mechanisms of translational control. *Nat Rev Mol Cell
Biol* **5**, 827–835 (2004).

14.   T. Maier, M. Güell, L. Serrano, Correlation of mRNA and protein in complex biological
samples. *FEBS Letters* **583**, 3966–3973 (2009).

15.   B. Schwanhäusser, *et al.*, Global quantification of mammalian gene expression control.
Nature **473**, 337–342 (2011).

16.   M. Jovanovic, *et al.*, Dynamic profiling of the protein life cycle in response to pathogens.
Science **347**, 1259038 (2015).

17.   M. Zhang, *et al.*, Impact of Growth Rate on the Protein-mRNA Ratio in Pseudomonas
aeruginosa. *mBio* **14**, e0306722 (2023).

18.   M. U. Caglar, *et al.*, The E. coli molecular phenotype under different growth conditions. *Sci
Rep* **7**, 45303 (2017).

19.   Y. W. Choi, S. A. Park, H. W. Lee, D. S. Kim, N. G. Lee, Analysis of growth phase-
dependent proteome profiles reveals differential regulation of mRNA and protein in
Helicobacter pylori. *PROTEOMICS* **8**, 2665–2675 (2008).

20. W.-H. Chen, *et al.*, Integration of multi-omics data of a genome-reduced bacterium:
Prevalence of post-transcriptional regulation and its correlation with protein abundances.
Nucleic Acids Res **44**, 1192–1202 (2016).

21. K. Becker, *et al.*, Quantifying post-transcriptional regulation in the development of
Drosophila melanogaster. *Nat Commun* **9**, 4970 (2018).

22. R. W. Corbin, *et al.*, Toward a Protein Profile of Escherichia coli: Comparison to Its
Transcription Profile. *Proceedings of the National Academy of Sciences of the United*
States of America **100**, 9232–9237 (2003).

23. S. P. Gygi, Y. Rochon, B. R. Franza, R. Aebersold, Correlation between Protein and mRNA
Abundance in Yeast. *Molecular and Cellular Biology* **19**, 1720–1730 (1999).

24. L. Jeacock, J. Faria, D. Horn, Codon usage bias controls mRNA and protein abundance in
trypanosomatids. *eLife* **7**, e32496 (2018).

25. L. Ponnala, Y. Wang, Q. Sun, K. J. van Wijk, Correlation of mRNA and protein abundance
in the developing maize leaf. *The Plant Journal* **78**, 424–440 (2014).

26. J. Wang, *et al.*, Combined proteomic and transcriptomic analysis of the antimicrobial
mechanism of tannic acid against Staphylococcus aureus. *Front Pharmacol* **14**, 1178177
(2023).

27. S. Kavela, *et al.*, Use of an Integrated Multi-Omics Approach To Identify Molecular
Mechanisms and Critical Factors Involved in the Pathogenesis of Leptospira. *Microbiol*
Spectr **11**, e0313522 (2023).

28. L. Huang, *et al.*, Integration of Transcriptomic and Proteomic Approaches Reveals the
Temperature-Dependent Virulence of Pseudomonas plecoglossicida. *Front Cell Infect*
Microbiol **8**, 207 (2018).

29. C. Kocharunchitt, T. King, K. Gobius, J. P. Bowman, T. Ross, Integrated transcriptomic and
proteomic analysis of the physiological response of Escherichia coli O157:H7 Sakai to
steady-state conditions of cold and water activity stress. *Mol Cell Proteomics* **11**,
M111.009019 (2012).

30. Q. Xiong, *et al.*, Integrated transcriptomic and proteomic analysis of the global response of
Synechococcus to high light stress. *Mol Cell Proteomics* **14**, 1038–53 (2015).

31. F. J. Perez-Llarena, G. Bou, Proteomics As a Tool for Studying Bacterial Virulence and
Antimicrobial Resistance. *Front Microbiol* **7**, 410 (2016).

32. X. Yang, *et al.*, RNA markers for ultra-rapid molecular antimicrobial susceptibility testing in
fluoroquinolone-treated Klebsiella pneumoniae. *J Antimicrob Chemother* **75**, 1747–1755
(2020).

33. T. Maier, *et al.*, Quantification of mRNA and protein and integration with protein turnover in
a bacterium. *Mol Syst Biol* **7**, MSB201138 (2011).

34. C. Erdem, S. M. Gross, L. M. Heiser, M. R. Birtwistle, MOBILE pipeline enables
identification of context-specific networks and regulatory mechanisms. *Nat Commun* **14**,
3991 (2023).

35. T. Kwon, H. K. Huse, C. Vogel, M. Whiteley, E. M. Marcotte, Protein-to-mRNA Ratios Are
Conserved between Pseudomonas aeruginosa Strains. *J. Proteome Res.* **13**, 2370–2380
(2014).

36. K. P. Jayapal, *et al.*, Uncovering Genes with Divergent mRNA-Protein Dynamics in
Streptomyces coelicolor. *PLOS ONE* **3**, e2097 (2008).

37. B. Li, C. N. Dewey, RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* **12**, 323 (2011).

38. P. Lu, C. Vogel, R. Wang, X. Yao, E. M. Marcotte, Absolute protein expression profiling
estimates the relative contributions of transcriptional and translational regulation. *Nat
Biotechnol* **25**, 117–124 (2007).

39. L. Barquist, *et al.*, A comparison of dense transposon insertion libraries in the Salmonella
serovars Typhi and Typhimurium. *Nucleic Acids Res* **41**, 4549–4564 (2013).

40. R. R. Chaudhuri, *et al.*, Comprehensive identification of essential Staphylococcus aureus
genes using Transposon-Mediated Differential Hybridisation (TMDH). *BMC Genomics* **10**,
291 (2009).

41. S. J. Willcocks, R. A. Stabler, H. S. Atkins, P. F. Oyston, B. W. Wren, High-throughput
analysis of Yersinia pseudotuberculosis gene essentiality in optimised in vitro conditions,
and implications for the speciation of Yersinia pestis. *BMC Microbiol* **18**, 46 (2018).

42. A. Mateus, *et al.*, The functional proteome landscape of Escherichia coli. *Nature* **588**, 473–
478 (2020).

43. O. M. Sigalova, A. Shaeiri, M. Forneris, E. E. Furlong, J. B. Zaugg, Predictive features of
gene expression variation reveal mechanistic link with differential expression. *Mol Syst Biol*
16, MSB209539 (2020).

44. J. Willforss, A. Chawade, F. Levander, NormalyzerDE: Online Tool for Improved
Normalization of Omics Expression Data and High-Sensitivity Differential Expression
Analysis. *J Proteome Res* **18**, 732–740 (2019).

45. M. E. Ritchie, *et al.*, limma powers differential expression analyses for RNA-sequencing and
microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

46. M. S. Robles, J. Cox, M. Mann, In-Vivo Quantitative Proteomics Reveals a Key Contribution
of Post-Transcriptional Mechanisms to the Circadian Regulation of Liver Metabolism. *PLOS
Genetics* **10**, e1004047 (2014).

47. M. Ashburner, *et al.*, Gene Ontology: tool for the unification of biology. *Nat Genet* **25**, 25–29
(2000).

48. S. Durica-Mitic, Y. Göpel, B. Görke, Carbohydrate Utilization in Bacteria: Making the Most
Out of Sugars with the Help of Small Regulatory RNAs. *Microbiology Spectrum* **6**,
10.1128/microbiolspec.rwr-0013–2017 (2018).

49. M. Raina, *et al.*, Dual-function AzuCR RNA modulates carbon metabolism. *Proceedings of
the National Academy of Sciences* **119**, e2117930119 (2022).

50.   T. Pisithkul, N. M. Patel, D. Amador-Noguez, Post-translational modifications as key
659   regulators of bacterial metabolic fluxes. *Current Opinion in Microbiology* **24**, 29–37 (2015).

660   51.   M. Hecker, J. Pané-Farré, V. Uwe, SigB-Dependent General Stress Response in Bacillus
661   subtilis and Related Gram-Positive Bacteria. *Annual Review of Microbiology* **61**, 215–236
662   (2007).

663   52.   H. Yakhnin, *et al.*, CsrA Represses Translation of sdiA, Which Encodes the N-
664   Acylhomoserine-l-Lactone Receptor of Escherichia coli, by Binding Exclusively within the
665   Coding Region of sdiA mRNA. *Journal of Bacteriology* **193**, 6162–6170 (2011).

666   53.   C. Pourciau, Y.-J. Lai, M. Gorelik, P. Babitzke, T. Romeo, Diverse Mechanisms and
667   Circuitry for Global Regulation by the RNA-Binding Protein CsrA. *Front. Microbiol.* **11**
668   (2020).

669   54.   X. Wang, *et al.*, CsrA post-transcriptionally represses pgaABCD, responsible for synthesis
670   of a biofilm polysaccharide adhesin of Escherichia coli. *Molecular Microbiology* **56**, 1648–
671   1663 (2005).

672   55.   A. K. Dubey, *et al.*, CsrA Regulates Translation of the Escherichia coli Carbon Starvation
673   Gene, cstA, by Blocking Ribosome Access to the cstA Transcript. *Journal of Bacteriology*
674   **185**, 4450–4460 (2003).

675   56.   E. Holmqvist, L. Li, T. Bischler, L. Barquist, J. Vogel, Global Maps of ProQ Binding *In Vivo*
676   Reveal Target Recognition via RNA Structure and Stability Control at mRNA 3′ Ends.
677   *Molecular Cell* **70**, 971-982.e6 (2018).

678   57.   A. Smirnov, C. Wang, L. L. Drewry, J. Vogel, Molecular mechanism of mRNA repression in
679   trans by a ProQ-dependent small RNA. *EMBO J* **36**, 1029–1045 (2017).

680   58.   M. Irastortza-Olaziregi, O. Amster-Choder, Coupled Transcription-Translation in
681   Prokaryotes: An Old Couple With New Surprises. *Front. Microbiol.* **11** (2021).

682   59.   M. Zhu, X. Dai, Maintenance of translational elongation rate underlies the survival of
683   Escherichia coli during oxidative stress. *Nucleic Acids Res* **47**, 7592–7604 (2019).

684   60.   M. Zhu, X. Dai, Bacterial stress defense: the crucial role of ribosome speed. *Cell. Mol. Life
685   Sci.* **77**, 853–858 (2020).

686   61.   M. Zhu, M. Mori, T. Hwa, X. Dai, Disruption of transcription–translation coordination in
687   Escherichia coli leads to premature transcriptional termination. *Nat Microbiol* **4**, 2347–2356
688   (2019).

689   62.   U. Vogel, K. F. Jensen, The RNA chain elongation rate in Escherichia coli depends on the
690   growth rate. *Journal of Bacteriology* **176**, 2807–2813 (1994).

691   63.   M.-P. Caron, *et al.*, Dual-acting riboswitch control of translation initiation and mRNA decay.
692   *Proceedings of the National Academy of Sciences* **109**, E3444–E3453 (2012).

693   64.   B. W. Scheithauer, J. M. Bruner, Central nervous system tumors. *Clin Lab Med* **7**, 157–179
694   (1987).

695   65.   X. Deng, *et al.*, Widespread occurrence of N6-methyladenosine in bacterial mRNA. *Nucleic*
696   *Acids Res* **43**, 6557–6567 (2015).

697   66.   J. M. Lee, H. M. Hammarén, M. M. Savitski, S. H. Baek, Control of protein stability by post-
698   translational modifications. *Nat Commun* **14**, 201 (2023).

699   67.   J. A. Cain, N. Solis, S. J. Cordwell, Beyond gene expression: The impact of protein post-
700   translational modifications in bacteria. *Journal of Proteomics* **97**, 265–286 (2014).

701   68.   A. Manteca, J. Ye, J. Sánchez, O. N. Jensen, Phosphoproteome Analysis of Streptomyces
702   Development Reveals Extensive Protein Phosphorylation Accompanying Bacterial
703   Differentiation. *J. Proteome Res.* **10**, 5481–5492 (2011).

704   69.   E. Bremer, R. Krämer, Responses of Microorganisms to Osmotic Stress. *Annu. Rev.*
705   *Microbiol.* **73**, 313–334 (2019).

706   70.   E. A. Mueller, P. A. Levin, Bacterial Cell Wall Quality Control during Environmental Stress.
707   *mBio* **11**, 10.1128/mbio.02456-20 (2020).

708   71.   T. Barrett, *et al.*, NCBI GEO: archive for functional genomics data sets—update. *Nucleic*
709   *Acids Res* **41**, D991–D995 (2013).

710   72.   C. S. Hughes, *et al.*, Ultrasensitive proteome analysis using paramagnetic bead technology.
711   *Mol Syst Biol* **10**, 757 (2014).

712   73.   A. T. Kong, F. V. Leprevost, D. M. Avtonomov, D. Mellacheruvu, A. I. Nesvizhskii,
713   MSFragger: ultrafast and comprehensive peptide identification in mass spectrometry-based
714   proteomics. *Nat Methods* **14**, 513–520 (2017).

715   74.   E. W. Deutsch, *et al.*, The ProteomeXchange consortium at 10 years: 2023 update. *Nucleic*
716   *Acids Res* **51**, D1539–D1548 (2023).

717   75.   R Core Team, *R: A Language and Environment for Statistical Computing* (R Foundation for
718   Statistical Computing, 2024).

719   76.   H. Wickham, *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag New York,
720   2016).

721   77.   J. H. Friedman, T. Hastie, R. Tibshirani, Regularization Paths for Generalized Linear
722   Models via Coordinate Descent. *Journal of Statistical Software* **33**, 1–22 (2010).

723   78.   T. Wu, *et al.*, clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.
724   *Innovation (Camb)* **2**, 100141 (2021).

725   79.   European Organization For Nuclear Research, OpenAIRE, Zenodo: Research. Shared.
726   [Preprint] (2013). Available at: https://www.zenodo.org/ [Accessed 11 December 2025].

727   Figures and Tables

728
Figure 1. Transcriptomic and proteomic analysis of three human bacterial pathogens under ten
infection relevant stress conditions. **(A)** Schematic presentation of data collection and data
generation for gene expression on mRNA and protein levels under stress conditions. **(B)**
Percentages of CDSs detected by transcriptomics and proteomics for the three species. **(C)**
Distribution of transcript abundance across all conditions (light blue) in the three species; yellow
indicates the fraction of proteins detected, while dark blue represents the transcripts for which no
corresponding protein was detected.
736

737
Figure 2. mRNA-protein levels show differing degrees of statistically significant positive correlation
under different stress conditions and control conditions in **(A)** *S. enterica* Typhimurium, **(B)** *S.*
aureus and **(C)** *Y. pseudotuberculosis*. mRNA and protein expression levels were taken as $\log_2$
transformed TPM and iBAQ values, respectively. The expression values of genes that are detected
in all conditions as mRNA and protein were used in this analysis. Pearson correlation coefficient
(R) shows positive correlation under all conditions in the three species. *p* values were calculated
using Student's t-test.

Figure 3. Essential genes of the three pathogens have higher mRNA and protein expressions with higher correlation and lower variance across the 11 conditions in comparison to all genes. **(A)** Mean mRNA expression frequency of essential and non-essential genes in the 11 conditions. **(B)** Mean protein expression frequency of essential and non-essential genes in the 11 conditions. **(C)** Essential genes and all genes mRNA-protein level correlation under 11 conditions for the three species. **(D)** Frequency of standard deviation of mRNA expression under 11 conditions for the three species. **(E)** Frequency of standard deviation of protein expression under 11 conditions for the three species. mRNA-protein level correlations were measured with Pearson correlation coefficient (R). *p* values were calculated using a Wilcoxon rank sum test.

Figure 4. mRNA-protein level correlation lowers with increasing degree of stress response. **(A)** mRNA-protein level correlations for conditions exposed to stressors for 10 minutes, 20 minutes, 30 minutes, 4 hours, and longer than 4 hours in the three species. **(B)** Percentages of differentially expressed genes (DEGs) and proteins (DEPs) under the 10 stress conditions in comparison to unexposed control conditions for the three species. **(C)** Scatter plots showing negative correlation between percentages of DEGs (top) and DEPs (bottom) and mRNA-protein level correlation. Correlations were measured with Spearman correlation (R) for **(A)** and Pearson correlation (R) for **(C)**. *p* values were calculated using Student's t-test.

Figure 5. Stress response induces different patterns of discordant mRNA and protein level regulation. **(A)** Scatter plots with Z-scores of mRNA and protein levels for each of the 10 stress conditions in comparison to control for the three species. The 6 gene clusters whose mRNA and protein level have different discordant regulation patterns are shown in different colors. **(B)** Z-score patterns of mRNA and protein levels across 10 stress conditions for the 6 discordantly regulated gene clusters in each species. The number of discordantly regulated genes per cluster is indicated with "*n*".

Figure 6. Impaired translation efficiency lowers mRNA-protein levels under osmotic stress. A) Stress specific stimulon genes and all genes mRNA-protein level correlation under 10 conditions for the three species. B) Schematic representation of MOBILE workflow for identifying Oss-specific mRNA-protein associations. The diagram outlines MOBILE analysis applied to transcriptomics and proteomics datasets from 11 conditions. Initial data from the control condition (LOGO Ctrl), the complete dataset (FULL), and the Oss condition (LOGO Oss) are used to construct corresponding base coefficient matrices. The FULL matrix is then compared against the Ctrl and Oss matrices to generate condition-dependent coefficient matrices, which highlight specific mRNA-protein connections that are either gained (red squares) or lost (blue squares). Finally, a comparison between the Ctrl-dependent and Oss-dependent matrices isolates the Oss-specific coefficient matrix, resulting in a targeted Oss-specific connections gene list. C) Over representation analysis (ORA) of Oss specific connections across three species, representing enrichment across biological process, molecular function, and cellular component. D) Polysome profiles of wt *Y. pseudotuberculosis* cells under control (Ctrl) and osmotic stress (0.5M NaCl) conditions. E) Western blot membrane (upper panel) and Coomassie blue-stained gel (lower panel) of total proteins extracted from wt *Y. pseudotuberculosis* cells under negative control (Ch), positive control

792 (Ctrl), osmotic stress (Oss), and low iron (Li). Active translation of nascent proteins was detected
793 using a monoclonal anti-puromycin antibody.

797 **Table 1.** Classification of differentially expressed genes based on mRNA and protein regulation
798 patterns.

Cluster	mRNA regulation	Protein regulation	Description
1	▼ Downregulated	● Not changed	mRNA is downregulated in at least one stress condition, while protein level remains unchanged
2	▼ Downregulated	▲ Upregulated	mRNA is downregulated in at least one condition, while protein level is upregulated
3	● Not changed	▼ Downregulated	mRNA level remains unchanged across all conditions, while protein is downregulated
4	● Not changed	▲ Upregulated	mRNA level remains unchanged across all conditions, while protein is upregulated in at least one condition
5	▲ Upregulated	▼ Downregulated	mRNA is upregulated in at least one condition, while protein level is downregulated
6	▲ Upregulated	● Not changed	mRNA is upregulated in at least one condition, while protein level remains unchanged

25