Quantitative Imaging of the Heterogeneity of Brain Potassium Depletion in Experimental Focal Ischemia

Alexander Kharlamov[3], Victor E. Yushmanov[3], Kirk A. Easley[4], Boris Yanovski[2,3], Stephen C. Jones[1,3]

1. CerebroScope, the dba entity of SciencePlusPlease LLC, 4165 Blair St., Pittsburgh, PA 15207-1508, USA.

2. Department of Anesthesiology, B'nai Zion Medical Center, Sderot Eliyahu Golomb 47, Haifa, Israel

3. Previously Department of Anesthesiology, Allegheny-Singer Research Institute, 320 E. North Ave., Pittsburgh, PA, 15212

4. Department of Biostatistics and Bioinformatics, Rollins School of Public Health, Emory University, 1518 Clifton Road NE, Atlanta, GA, 30322, USA

Correspondence to:
Stephen C. Jones, PhD
CerebroScope
4165 Blair St.
Pittsburgh, PA 15207-1508
USA
412-208-3397
412-849-9584 (cell)
sjones0123@alum.mit.edu
sjones@cerebro-scope.com

Number of text pages: 41

Word count: 6017 words
Number of Figures: 6
Number of Tables: 3

Running Head: Lower Potassium in the Peripheral Ischemic Core

Abstract

With few exceptions, pathological progression in ischemic stroke is presumed to occur uniformly within the ischemic core region. These exceptions include edema formation, brain tissue $[Na^+]$ increase, and the qualitative visually-observed decrease of brain tissue $[K^+]$, $[K^+]_{br}$, all of which occur in peripheral regions of the ischemic core. We hypothesize that $[K^+]_{br}$ within these peripheral regions are heterogeneous (with lower $[K^+]_{br}$ in the peripheral compared to the central ischemic core) and are not associated with neuronal degradation.

Permanent focal ischemia in 13 rats was produced for 2.5-5 h. Brain sections were quantitatively stained for K^+ to assess $[K^+]_{br}$ variations between the peripheral and central ischemic core. Regions within the cortical ribbon were used to explore differing rates of K^+-depletion expressed as the slopes of $[K^+]_{br}$ vs. time relations. Adjacent sections were observed for reflective change and stained for microtubule-associated protein 2 (MAP2) to identify the ischemic region and to relate neuronal pathology to $[K^+]_{br}$ variations.

The mean value of normal cortex (NC) $[K^+]_{br}$ was 96 mEq/kg and of K^+-depletion in all ischemic regions over time was 12.2 mEq/kg/h, consistent with measurements from other studies. Exaggerated K^+-depletion occurred in 56% of the peripheral ischemic core regions classed as depleted peripheral ischemic core (ICp-DP) regions. These were clearly separated (p<0.001) from the non-depleted peripheral ischemic core (ICp-ND) regions. The normal cortex (NC) regions show stability of $[K^+]_{br}$ with a slope near zero. However, the 13.6 mEq/kg/h slopes of the central ischemic core (ICc) and ICp-ND regions were similar (p=0.99) and showed a significant decrease over time. The 6.2 mEq/kg/h slope of the ICp-DP regions was significantly different from that of the ICc (p=0.010) and the ICp-ND (p=0.0071). This lower slope of the ICp-DP curve 2.5 h after stroke onset is due to the accelerated K^+-efflux from 0 to 2.5 h, as its value at stroke onset must be ~100 mEq/kg. However, these differential K^+

losses were not reflected in the homogeneous peripheral ischemic core MAP2 immunoreactivity losses. Unlike $[K^+]_{br}$, there was no difference between the MAP2 immunoreactivity in K^+-depleted and non-K^+-depleted peripheral ischemic core regions (ICp-ND vs ICp-DP, ICp-ND vs ICp-DP, unpaired t-test, p=0.83, p=0.16, respectively).

While confirming previous results of quantitative regional losses of $[K^+]_{br}$ in the ischemic core, we show that K^+ dynamics within the peripheral and the central ischemic core are heterogeneous and not related to MAP2-assessed neuronal structural integrity: the K^+-depleted regions in the peripheral ischemic core regions are presumably closer to glymphatic system and other K^+-efflux pathways. Such differing K^+ dynamics at the edge of the ischemic core in the hyper-acute period in first hours after ischemic onset possibly relate to the spreading depolarization-mediated expansion of the infarct during the period of secondary brain injury. Peripheral ischemic core regions with less K^+ might limit spreading depolarization initiation and propagation if there is insufficient K^+ for depolarization to occur and make restoration of parenchymal membrane potential improbable even if the functionality of the Na^+,K^+-ATPase is restored. Further study of differing K^+-dynamics within the ischemic core might lead to a better understanding of ischemic stroke pathophysiology.

Keywords: acute focal ischemic stroke; potassium imaging; spreading depolarization; peri-infarct depolarizations

Introduction

Sodium and K^+ dynamics in ischemic stroke have been viewed as important issues because they are the major cations that maintain membrane potential via the Na^+,K^+-ATPase and whose transient intra-intercellular disequilibrium subserve ischemic depolarization. Many experimental studies have used increasingly more sophisticated methods for their determination (Grochowski et al., 2019), including atomic absorption spectroscopy (Pappius et al., 1956; Walsh, 1955), ion-sensitive electrodes (Octeau et al., 2018), non-hydrogen-based MRI (Atkinson et al., 2014; Elabyad et al., 2014; Yushmanov et al., 2007; Yushmanov et al., 2009b), ion-sensitive fluorochrome imaging (Wu et al., 2022), and genetically encoded fluorescent probes (Bischof et al., 2017). Although Na^+ and K^+ can be viewed as complementary cations, with K^+ being intracellular and Na^+ extracellular, the vast majority of experimental effort has been directed at Na^+ dynamics. Na^+ accumulation and subsequent edema is considered one of the principal mechanisms of ischemic stroke pathophysiology (O'Donnell et al., 2004; Simard et al., 2012) and is the major pathological factor of brain-edema instigated tentorial herniation (Treadwell et al., 2010; Wen et al., 2020), the most egregious event in the progression of ischemic pathology. This focus on Na^+ has left the consideration of K^+ dynamics in the background. However, although Na^+ is important in the immediate pathological consequences of ischemic stroke, K^+ is equally important if recovery issues are to be considered. Because K^+ moves from intra- to extra-cellular during depolarization, it is exposed to the brain's exit systems, in contrast to Na^+'s cellular entrapment. Thus post-ischemic measurements of K^+ (of course almost always made in concert with Na^+) show a consistent and excessive loss of K^+ from the brain (Betz et al., 1990; Betz et al., 1994; Dickinson et al., 1992; Ito et al., 1979; Lo et al., 1987; Menzies et al., 1990; Schuier et al., 1980; Shibata et al., 1974; Yang et al., 1992; Young et al., 1987) as presented in Table 1.

These measurements of K^+ in acute ischemic stroke were made with the assumption that the ischemic core $[K^+]_{br}$ was homogeneous. However, there have been reports of intra-core heterogeneity in ischemic stroke. These include observations in the peripheral regions of ischemic core of increased

edema, lowered brain tissue [K^+], [K^+]$_{br}$, and increased [Na^+]$_{br}$ (Kato et al., 1987), a maximal rate of [Na^+]$_{br}$ increase (Jones et al., 2006; Yushmanov et al., 2009b), transient blood-brain barrier (BBB) breakdown to gadolinium diethylenetriaminepentaacetic acid (Gd-DTPA) (Yushmanov et al., 2009a), decreased reflective change (Kharlamov et al., 2001), decreased MAP2 immunoreactivity (IMR) (Dawson et al., 1996; Kharlamov et al., 2001; Pettigrew et al., 1996), and decreasing [K^+]$_{ex}$ (Sick et al., 1998). These measurements suggest that the peripheral ischemic core has a special and unique significance during this acute period that relate to the SDs that occur repetitively around the edge of the ischemic core (Kumagai et al., 2011; Nakamura et al., 2010).

To expand our understanding of K^+ dynamics to include a more intra-ischemic core view, we propose to use quantitative K^+-imaging (Yushmanov et al., 2013). Previous methods used for the measurements in Table 1 cannot address the regionality within the ischemic region needed to assemble a complete picture of K^+-related ischemic core pathology. However quantitative K^+-imaging (Yushmanov et al., 2013) permits the examination of our hypotheses concerning the regional and temporal differences in the clearance of K^+ from ischemic tissue: 1) That at early times after ischemia, K^+ efflux from the peripheral and central ischemic core are different; and 2) That these differences in K^+-dynamics are not based on neuronal integrity assessed by calpain-mediated microtubule disassembly via MAP2 IMR changes.

Answers to these hypotheses could help us understand and mitigate K^+ related stroke issues. As SDs continue to occur in the initial several hours after stroke onset more K^+ becomes extracellular, and more and more K^+ is lost from the brain. There are at least two consequences of this K^+ loss, that are equally important when compared to Na^+ accumulation: K^+ dependent 1) neuronal recovery; and 2) SD ignition and propagation. Even if energy supply can be resupplied to marginally ischemic tissue by the reestablishment of Na^+,K^+-ATPase activity, without K^+, the membrane potential cannot be reestablished. If appropriate [K^+]$_{in}$ levels cannot be restored to allow SD's depolarization to occur even though K^+ is not a primary factor in SD initiation (Andrew et al., 2022; Ricks et al., 2025) then eventually SDs will become less likely.

Methods

Animal procedures

Animal procedures were performed in compliance with the National Institutes of Health "Guide for the Care and Use of Laboratory Animals" and were approved by the Institutional Animal Care and Use Committee, DHS/OLAW Assurance #A3693-01. Thirteen Sprague/Dawley male rats (285 ± 7 g, mean ± sem) were anesthetized with isoflurane (2.5% induction and 1.0 ± 0.5% maintenance), 30% oxygen and balance nitrous oxide, intubated, and artificially ventilated (Harvard Apparatus, model 683, South Natick, MA). During maintenance anesthesia, the concentration of isoflurane was titrated by monitoring the arterial blood pressure response to tail-pinch. Body temperature was maintained at 37°C by a rectal probe and servo-controlled heating lamp (YSI 74, Yellow Springs, OH). Venous and arterial femoral catheters (PE50) were inserted. Arterial blood gases ($PaCO_2$, PaO_2), and pH were determined with a portable blood analyzer (i-STAT, Heska, Loveland, CO). Arterial blood pressure was continuously monitored. Pancuronium (Gensia Sicor Pharmaceuticals, Irvine, CA) was administered hourly (0.7 mg/kg, IM). Physiological variables and laser Doppler flowmetry (LDF)-determined cerebral blood flow (CBF) were digitally recorded with Windaq software (Dataq, Inc., Columbus OH).

Model of brain ischemia

The animals were subjected to permanent focal ischemia produced by right middle cerebral artery transection (MCAt) midway between the olfactory tract and inferior cerebral vein followed by permanent bilateral common carotid artery occlusion (biCCAo) (Chen et al., 1986; Jones et al., 2006) or, in one animal, MCA permanent intraluminal suture occlusion (Yushmanov et al., 2009c). CBF was monitored by LDF (λ = 780 nm, LaserFlo BPM403A, Vasamedics, Inc., St. Paul, MN) with the probe firmly placed in a 1-mm diameter skull bone window over the right MCA territory -4 mm from bregma and 5 mm from the midline (see Fig. 1A). CBF was expressed as the percent of pre-ischemic control after correction for biological zero obtained at the end of the experiment (Kharlamov et al., 2004). CBF was not recorded in two animals due to technical issues. Animals were killed at 15 minute intervals between 2.5 and 5 h

after stroke onset (two each at 2.75 and 4.5 h, and one at each of the other 9 times). The brains were removed and frozen over dry ice and stored at -70°C. In the MCA suture animal, the whole head was frozen and the brain chipped out.

Brain processing

Brain tissue samples were collected in the microtome-cryostat (Minotome, IEC, Needham Heights, MA) at -7°C using a micropuncher (2 x 0.5 mm ID) for flame photometry analysis of $[K^+]_{br}$ (Hu et al., 2000; Yushmanov et al., 2013). Micropunch samples (depth ~2 mm, 0.545 ± 0.005 mg, 14 punches/level x 2 levels/animal x 13 animals) from ischemic and contralateral (normal) brain areas were taken at two different rostro-caudal levels as shown in Fig. 1A: L1 (+0.7 to -1.3 mm from bregma) and L2 (-1.3 to -3.3 mm from bregma). The K^+-content of brain tissue samples was determined by emission flame photometry at 766 nm (IL943 flame photometer, Instrumentation Laboratory, Lexington, MA) and expressed as K^+ brain concentration, $[K^+]_{br}$, in mEq/kg (Yushmanov et al., 2013). The change of surface reflectivity, or reflective change (Kharlamov et al., 2001), on adjacent dried coronal sections (35 µm, Fig. 2A) provided unequivocal differentiation of ischemic from normal cortex during micropunch sampling and confirmed the extent of focal ischemic damage. The images of the cut-face of the brain (Fig. 2B) at each level (S1, S2, and S3, Fig. 1A) were digitized for later alignment of the micropunch locations with adjacent histo-K^+-stained sections (Fig. 2C). Additional adjacent sections were immunoreactivity (IMR) stained for the cytoskeletal protein MAP2 (Kharlamov et al., 2001), as shown in Fig. 2G.

Histological K^+-staining and $[K^+]_{br}$ calibration

The quantitative histochemical method for brain K^+-staining described previously (Yushmanov et al., 2013) was used. This method is based on the chemical reaction of K^+ with sodium cobalt hexanitrite, $Na_3Co_2(NO)_6$, with formation of potassium cobaltinitrite crystals. The barely visible yellow crystalline precipitates of potassium cobaltinitrite transform into easily detectable black cobalt sulfide crystals by adding ammonium sulfide in a cold bath kept less than 4°C (Mies et al., 1984; Silberstein et al., 2004;

Yushmanov et al., 2013). The sections histochemically stained for K^+ (histo-K^+-stained) were digitized and calibrated using flame photometry-determined $[K^+]_{br}$ values from micro-punch samples (Fig. 2D). The optical density's of each histo-K^+-stained section were separately calibrated against flame photometry determined $[K^+]_{br}$ from micropunch samples taken directly from the cut-face left after that microtome section was taken.

Time-course analysis of cortical-ribbon K^+-depletion

In the regions of reflective change and depressed MAP2 IMR, $[K^+]_{br}$ dynamics were analyzed in the acute period after experimental ischemic stroke in three coronal sections (S1, S2, and S3 in Fig. 1A). Striking differences in $[K^+]_{br}$ between the central ischemic core, ICc, and the dorsal and ventral peripheral ischemic core regions, ICpd and ICpv (Fig. 2F), were apparent from visual assessment of the histo-K^+-stained sections: $[K^+]_{br}$ was depressed in some but not all peripheral ischemic core regions. Further $[K^+]_{br}$ analysis was driven by this unusual pattern of excessively depressed $[K^+]_{br}$, termed K^+-depletion, in the peripheral ischemic core in relation to the central ischemic core.

The calibrated histo-K^+ images (Fig. 2E) were analyzed for spatial changes of $[K^+]_{br}$ using the ribbon tool of the MCID imaging system (Version 7, InterFocus Imaging Ltd., Linton, United Kingdom). The ribbon tool produced a plot of the average $[K^+]_{br}$ within the cortical rim from the dorsal hemispheric fissure to the ventral midline and allows the placement of within-ribbon regions-of-interest (ROIs). The width of the ribbon was defined by the plateau of the GV profile of the density across the cortex layers indicating uniform staining. Using this width of 0.9 mm, the cortical profiles of $[K^+]_{br}$ along the ischemic and non-ischemic hemispheres were obtained (see Fig. 2F and 2H).

The 39 $[K^+]_{br}$ ischemic hemisphere-ribbons (3 sections per animal x 13 animals) showed two distinct patterns of their $[K^+]_{br}$ cortical profiles. One pattern (25 ribbons) had a wide ischemic plateau with low $[K^+]_{br}$ at one or both peripheral ischemic core regions. The other pattern (14 ribbons) had only a sharp minimum without a wide central ischemic plateau region so a central ICc region could not be identified. The $[K^+]_{br}$ profile of these patterns, which appeared as a "wedge", occurred because the

cortical ribbon "cut" was positioned at the caudal or rostral edges of the ischemic region. The $[K^+]_{br}$ in these cortical ribbons with "wedge" patterns were therefore excluded from analysis because K^+ loss at the peripheral parts of the ischemic core could not be compared to that in the central ischemic core.

In the remaining 25 cortical ribbons, five different within-ribbon ROIs were selected in the ischemic hemisphere-ribbon: a) two normal cortex regions of parasagittal and pyriform parts of the ischemic hemisphere (dorsal normal cortex, NCd, and ventral normal cortex, NCv, respectively); b) the ICc; and c) two ROIs at the periphery of the ischemic core, one dorsal (ICpd) and one ventral (ICpv) region identified by a consistent depression of $[K^+]_{br}$ or at those same positions in sections without such consistent depression (see Fig. 1B and 2F). Peripheral ischemic core regions (ICp) were selected based on their $[K^+]_{br}$ distribution according to our hypotheses as there is no known morphological attribute or criteria that provides such guidance.   The $[K^+]_{br}$'s of the ICpd and ICpv regions were combined into one group, ICp, for additional analysis, as they were deemed statistically similar (p>0.59, paired t-test). Five corresponding homotopic regions were chosen in the non-ischemic hemisphere ribbon.

To analyze K^+-egress from these ICp regions, their $[K^+]_{br}$, $[K^+]ICp$, was expressed as a percentage of that in the ICc region, ($[K^+]ICc$), based on the assumption that the $[K^+]_{br}$ of these peripheral ischemic core regions was originally equal to that of the central ischemic core:

$[K^+]ICp\% = 100 \times [K^+]ICp / [K^+]ICc$.

Histogram analysis was then used to explore variations in K^+-depletion within the ischemic core. The frequency distribution of $[K^+]ICp\%$ (n = 50) revealed an apparently bimodal distribution. To assess this distribution for bimodality, two models were compared; a bimodal model with two Lorentz curves and another with just one Lorentz curve (see Fig. 3). The trough of the bimodal frequency distribution, the $[K^+]ICp\%$ value at the intersection of the two Lorentz curves, was used to establish a breakpoint between peripheral ischemic core regions with (ICp-DP) and without (ICp-ND) K^+-depletion.

Finally $[K^+]_{br}$ values of the NC, ICc, ICp-DP and ICp-ND regions were plotted vs. time after stroke onset (Fig. 4). Each of these curves was assessed for linearity and differences between their

slopes and intercepts were statistically tested. Forty-four peripheral ischemic core regions were available for analysis: 6 regions from suture model animal (2 peripheral ischemic core regions * 3 sections) were excluded from the 50 peripheral ischemic core regions. This animal was excluded from the time course analysis because the suture MCA model used in this animal has a slower progression of cortical ischemic pathology (Wetterling et al., 2012) in comparison to the direct distal occlusion of the MCA (Jones et al., 2006) used in all other animals.

Immunostaining for cytoskeletal protein, microtubule-associated protein-2 (MAP2)

The decrease of IMR for the cytoskeletal protein MAP2 was used as a marker of early ischemic damage (Dawson et al., 1996; Kharlamov et al., 2001; Pettigrew et al., 1996). The MAP2 IMR was detected with anti-MAP2 primary antibody (HM2, 1:500, Sigma-Aldrich, St. Louis, MO) on adjacent sections to those histo-K^+-stained. MAP2 immunostaining in frozen sections can lead to artifacts, so only 27 immuno-stained sections (out of 39 = 3 sections per animal x 13 animals) were technically acceptable for analysis (two animals had none, six animals had 2, and the remaining 5 had 3 each that were acceptable). MAP2 IMR was normalized by expressing the ratio of the gray value (GV) in each ROI to the average GV of normal cortex for each section to correct for variations in section thickness, tissue processing, and staining intensity. Decrements of MAP2 IMR and $[K^+]_{br}$ were compared using the transposed cortical ribbon ROIs of the histo-K^+-stained sections (Fig. 2E) onto the MAP2 IMR sections (Fig. 2G) as presented in Fig. 5.

Statistics

Two-tailed paired and unpaired t-tests were used to test differences between CBF over time and between the ICc and ICp ROI values of $[K^+]_{br}$ and normalized MAP2 IMR. The distribution of ICp $[K^+]_{br}$ values was tested for bimodality by comparing the fitted Lorenz distribution of all the data to the sum of the bimodal Lorenz distributions using an F-test and a second order Akaike Information Criterion (AIC) test using OriginPro (OriginLab Corporation, Northampton, MA). Differences between slopes of the linear regression analysis of the $[K^+]_{br}$ time-courses were evaluated from the t-statistic calculated by

their difference divided by the RMS of their standard errors. These linear regression analyses were performed on pooled ROI data, which assumes independence between samples (ROIs are nested within animals). Consequently, the between-animal variance may not be negligible, and the standard errors used for slope comparisons via the t-statistic may be underestimated with the caveat that precise estimates of the two variance components would be limited with only 13 animals. Values are presented as mean ± sem. Significance was assumed for $p < 0.05$.

Results

Physiological variables (mean arterial blood pressure, blood pH, $PaCO_2$ and PaO_2) were within normal limits before ischemia and were stable after ischemia (Table 2). After ischemia, the CBF over the cortical area of the right MCA territory decreased to 16% ± 3% (mean ± sem, n = 11) of control (pre-ischemic) values (Table 2). At the end of the experiment, the mean CBF was 28% ± 7%, showing a insignificant trend higher than the immediate post-ischemic value (p = 0.063, Table 2), consistent with potentially increased collateral circulation in 3 of the animals.

Ischemic damage from reflective change (Fig. 2A) and MAP2 IMR images (Fig. 2G) was observed only in the right cortex within the MCA distribution, extending from the frontoparietal cortex to the parts of the parasagittal and pyriform cortices. The normal (non-ischemic) cortical and subcortical structures on histo-K^+-sections were uniformly stained a dark brown color with white matter less intensely stained than gray matter. The ischemic cortex demonstrated a distinct loss of histo-K^+-staining. However, the peripheral ischemic core regions often, but not always, showed more intense loss of K^+ staining, termed K^+-depletion (Fig. 1B, 2E, 2G, and 2H).

Histogram analysis for K^+-depletion

Histogram analysis of the $[K^+]ICp\%$ values was used to compare the lower $[K^+]_{br}$ values in the dorsal and ventral peripheral ischemic core regions (the K^+-depleted group, termed ICp-DP, to the non-K^+-depleted regions, termed ICp-ND. The $[K^+]ICp\%$ values conformed to a bimodal distribution (Fig. 3, p < 0.001, F-test), with an AIC weight ratio indicating that the bimodal fit is 7.5×10^7 times more likely to be

correct than a single function fit. A break-point $[K^+]ICp\%$ value of 78% separated the ICp regions into two groups: those below, K^+-depleted (ICp-DP), and those above, non-depleted (ICp-ND).   In this separation of peripheral ischemic core regions with and without K^+-depletion, the right-hand peak of Fig. 3 represents $[K^+]_{br}$'s with values similar to that of the ICc regions (median $[K^+]ICp\%$ of 90%), and the other, with a median $[K^+]ICp\%$ of 64%, represents ICp regions that have undergone K^+-depletion.

The breakpoint of 78% (Fig. 3, vertical red line) corresponds to 36 mEq/kg (calculated as 0.78 of the mean $[K^+]ICc$, 45.8 ± 3.1 mEq/kg). K^+-depletion was noted in all but 2 of the 13 animals and in 28 (or 56%) of the 50 ICp regions in the 25 cortical ribbons available from both the MCAt&biCCAo and MCA-suture models.

The comparison of $[K^+]_{br}$ in the peripheral ischemic core regions with and without K^+-depletion clearly indicates that the $[K^+]_{br}$ of the K^+-depleted ICpd and ICpv regions are approximately one half (25.9 ± 2.3 mEq/kg, n = 11 and 24.9 ± 3.4 mEq/kg, n = 11) of the ICc value (45.0 ± 3.5 mEq/kg, n = 17, p < 0.001 for both, see Figure 5B). Most importantly, the $[K^+]_{br}$ in the peripheral ischemic core regions without K^+-depletion (46.7 ± 4.8 mEq/kg, n = 11, p = 0.86 mEq/kg and 44.6 ± 4.8 mEq/kg, n = 11, p = 0.59, unpaired t-tests) are approximately equal to the ICc value (47.7 ± 3.6 mEq/kg, n = 17), bolstering our hypothesis that no accelerated K^+-efflux occurred in these particular peripheral ischemic core regions.

K^+-efflux - time course

For comparison to others' results, none of which address regional variations within the ischemic core, all the ischemic cortex $[K^+]_{br}$ values (ICpd, ICpv, and ICc) were combined into one group, $[K^+]IC$. The linear regression curve of these combined ischemic core regions over the time from stroke onset, Ta, was $[K^+]IC = 83.8 - 12.2 * Ta$, with a K^+-efflux rate of 12.2 mEq/kg/h, as presented in Table 1.

The time courses of K^+-efflux from 2.5 to 5 h presented in Fig. 4. The stability of $[K^+]_{br}$ in normal cortex and the varying efflux rates of K^+ from the other regions are clearly shown for normal cortex, non-depleted peripheral-, depleted peripheral-, and central ischemic core. The slope of the K^+-depleted

curve differed significantly (p = 0.0071) from that of the non-K^+-depleted curve and the ICc curve (p = 0.010). Quantitatively, K^+ was lost at the same linear rate in the ICc regions as in the non-K^+-depleted peripheral ischemic core, ICp-ND, regions, with a non-significant difference between their slopes (p = 0.99). When taken together, these results indicate that accelerated K^+-depletion occurred in the ICp-DP regions during the first 2.5 h after stroke onset and then became slower compared to the ICc and ICp-ND regions.

K^+-depletion - relation to MAP2 IMR

The $[K^+]_{br}$ cortical ribbons and ROIs were transposed to the aligned MAP2 images and used to assess whether the loss of normalized MAP2 IMR was associated with the loss of K^+. MAP2 IMR was decreased in the peripheral ischemic core regions compared to their corresponding NCd and NCv normal cortex regions (Fig. 5, 64% and 47%, respectively, p < 0.001 for both). When the central ischemic core MAP2 IMR was compared to the MAP2 IMR in the dorsal and ventral ICp regions, statistically significant decreases of 72% (p < 0.001) and 74% (p < 0.001) were noted. However, there was no difference between the MAP2 IMR in K^+-depleted and non-K^+-depleted dorsal or ventral peripheral ischemic core regions (ICpd-DP vs. ICpd-ND: 0.464 ± 0.051, n = 11 vs. 0.478 ± 0.037, n = 8, p = 0.83; ICpv-DP vs. ICpv-ND: 0.493 ± 0.045, n = 10 vs. 0.419 ± 0.024, n = 9, p = 0.16, unpaired t-tests) in contrast to the differences observed in $[K^+]_{br}$ between non-K^+-depleted and K^+-depleted peripheral ischemic core regions. Thus those regions with K^+-depletion did not show a similar depression of MAP2 IMR.

DISCUSSION

The main findings of this experimental study of brain potassium loss in ischemic stroke are: 1) that the loss of K^+ from ischemic brain tissue agrees with previous measurements using less regionally specific methods; 2) that there were two statistically distinct populations of $[K^+]_{br}$ and K^+-efflux in the peripheral ischemic core regions; and 3) these differences in K^+-dynamics are not related to pathological degradation of neuronal structure. These K^+ changes in the peripheral regions of the ischemic core are

at the crux of ischemic pathology with its hallmark of K^+-mediated ischemic depolarization, which - with high $[K^+]_{ex}$ - was used to initially define the ischemic threshold (Astrup et al., 1977), the penumbra (Branston et al., 1977), and ischemic cortical spreading depolarizations (SDs) (Branston et al., 1977) using K^+-sensitive electrodes.

Our measurements of total brain K^+, $[K^+]_{br}$, in ischemic tissue from histo-K^+-stain imaging can be compared to $[K^+]_{ex}$ values obtained in a similar permanent MCAo model by Sick et al. (1998). However, Sick et al.'s (1998) measurements show a stability of $[K^+]_{ex}$ in ischemic core, while our and others measurements of $[K^+]_{br}$ show a linear decrease. To explore these different time courses we used the formalism of Cserr et al. (1991) to calculate $[K^+]_{in}$ from our $[K^+]_{br}$ values and the $[K^+]_{ex}$ values of Sick et al. (1998). The equations of Cserr et al. (1991) are based on the conservation of mass, fractional water content, and fractional extracellular space volume in depolarized brain (Hansen et al., 1980; Nicholson et al., 1981). Table 3 provides the results of these calculations using our estimates of $[K^+]_{br}$ at 0, 0.5, 1.5, and 3 h and $[K^+]_{ex}$ from ischemic core Zones 2 and 3 and normal cortex Zone 8 from Sick et al. (1998). The estimates of $[K^+]_{in}$ show a decrease over time and suggest that the fall of total brain $[K^+]$ was due to the decrease in $[K^+]_{in}$ until at 3 h $[K^+]_{in}$, $[K^+]_{ex}$, and $[K^+]_{br}$ are approximately equal. This equality indicated complete depolarization as confirmed by the estimated K^+ Nernst potential of -5 mV. These estimates of $[K^+]_{in}$ are consistent with our observed decreases over time in $[K^+]_{br}$ in central ischemic cortex.

Our measurements of excessive loss of K^+ from peripheral ischemic core regions suggest several possible consequences. The first is that isoelectric SDs could become more likely during this initial 5 h period after stroke onset as without K^+, repolarization cannot occur. Alternatively even if some regions near the ischemic core can repolarize, SD propagation into adjacent ischemic regions becomes improbable because K^+ loss prevents the necessary repolarization. Even if the Na^+,K^+-ATPase could operate there is not enough K^+ left to re-supply $[K^+]_{in}$ and repolarize the cell membrane. This is consistent with Hartings et al.'s (2003) observations that SDs occur frequently (~6/h) in the initial 2 h after stroke onset but then become less frequent (~2/h) between 3-7 h. This suppression of SD

occurrence after 2-3 h from stroke onset also suggests a temporary limit to SD-mediated infarct expansion (Woitzik et al., 2013).

Perhaps the most important consequence of the loss of K^+ near the penumbra involves the reduced chance of recovery. With such low $[K^+]_{br}$ there is not enough K^+ left to reestablish $[K^+]_{in}$ even if circulatory supply is reinstituted and Na^+,K^+-ATPase functionality is re-established. Thus the probable consequence of low K^+ is that the membrane potential cannot be reestablished and neuronal/glial rescue is improbable: "Normalization of $[K^+]_e$ [after SD] is one of the critical components for restoration of neuronal activity" (Monai et al., 2021). Young et al. (1994) suggested the direct replacement of K^+ during neurosurgical procedures so that repolarization and neural transmission can occur although it is unclear how this could be accomplished. Two of these factors, limited infarct expansion and the decreased possibility of neuronal recovery, represent seemingly conflicting mechanisms that are in actuality a sequence of an immediate positive factor followed by a long-term negative. Thus K^+-depletion at the peripheral ischemic core has important clinical and therapeutic implications resulting from an enhanced understanding of the intra-ischemic core variations in the SD-mediated pathological progression of focal ischemia.

Pathophysiology of the peripheral ischemic core

Our observations of regional variations in K^+-efflux in the ischemic core compliment the changes in the peripheral ischemic core in the acute post-stroke time of up to 5 h as mentioned in the introduction, but are not often the subject of studies of focal ischemic pathology and are distinct from the micro-heterogeneity and temporal progression in the ischemic core defined by molecular and metabolic studies and observations of vascular integrity (Del Zoppo et al., 2011). However, K^+ losses from the entire ischemic region during focal ischemia has been well documented using tissue analysis by atomic absorption spectrophotometry or flame photometry (see Table 1). The estimated rate of K^+-efflux of 12.2 mEq/kg/h for the entire ischemic cortex from the pooled ischemic core ROIs (ICc, ICpd, and ICpv) in this study is consistent with an overall average of others' tissue sampling results of 10.1 mEq/kg/h

(Table 1), providing a pathophysiological validation to our quantitative K^+-imaging method. However, the very much smaller sample weights of our micro-punch methodology (0.545 mg), compared to the mean of the others from Table 1 (75 mg), made possible our intra-ischemic core analysis that disclosed a more nuanced view of K^+-egress from the ischemic core.

The bimodal analysis separating the peripheral ischemic core regions with accelerated K^+-efflux, ICp-DP, from those with K^+-efflux slopes similar to those of ICc, ICp-ND, succeeded because it was based on the comparison of each peripheral region to its own central ischemic core region within the same cortical ribbon. The finding that the bimodal distribution was statistically significant permitted the identification of a breakpoint between the depleted and non-depleted peripheral ischemic core regions as the trough of the bimodal distribution at 36 mEq/kg (see Fig. 3). This distinction in turn allowed for the observations of the significantly different K^+ efflux rates for ICp-DP and similar rates for ICp-ND and ICc as depicted in Fig. 4. Over one-half (56%) of the peripheral ischemic core regions have lower $[K^+]_{br}$ and higher K^+-efflux than the remaining and at 2.5 h these higher K^+-efflux regions lost approximately twice as much K^+ compared to those without K^+-depletion.

Our quantitative imaging measurements confirm the qualitative observation of lower gray-values in histo-K^+-stained sections in the peripheral, compared to the central, ischemic core, that was made in the gerbil model of focal ischemia (Kato et al., 1987). $[K^+]_{br}$ values from cortical brain punch samples in this study (Kato et al., 1987) did not reveal a significant decrease of $[K^+]_{br}$ in the peripheral ischemic core in comparison to the ICc when expressed as mEq/kg dry weight (p = 0.085). However, when converted to mEq/kg wet weight using their own regional brain tissue water contents, this difference between the two was significant (p = 0.0025). In another study (Mies et al., 1984) a qualitative decrease in $[K^+]_{br}$ in the peripheral ischemic core is evident in their Fig. 2B in histo-K^+-stained ischemic gerbil brain, but this K^+-depletion was not mentioned or analyzed.

Imaging vs. other methods

The quantitative imaging methodology used in this study has enabled a clear view of potassium

dynamics in the ischemic core that has not been accessed by other methods including K^+-sensitive microelectrodes, K^+-sensitive fluorochrome imaging, and micropunch tissue sampling because it combines an appropriately high spatial resolution with a width breath of view. This view allowed for the definition of the border not only between the ischemic core and normal cortex but also between the peripheral and central ischemic core.

Relation of K^+-depletion to neuronal integrity

Although we observed depressed MAP2 IMR staining in the peripheral ischemic core, there was no exaggerated MAP2 depression in those peripheral ischemic core regions that were K^+-depleted (ICp-DP) compared to peripheral core regions that were not K^+-depleted (ICp-ND). Both K^+ non-depleted (ICp-ND) and depleted regions (ICp-DP) had the same level of MAP2 IMR depression (see Fig. 5). This lack of correspondence between the decrease in MAP2 IMR and K^+-depletion is consistent with our contention that K^+-depletion is not based on calpain-mediated microtubule disassembly and that the heterogeneity of K^+-efflux dynamics lacks a neuronally specific morphological correlate.

Association of K^+-depletion with SD occurrence

A ^{42}K loading study showed that SDs cause massive amounts of K^+ to leak from the brain to the surgically exposed brain surface (Brinley, Jr. et al., 1960). This massive efflux of labeled K^+ occurred in synchrony with the DC-shift of a remotely- and electrically-elicited SD, so is consistent with the K^+-egress of our study because SDs occur with high frequency in the first hours after stroke (Hartings et al., 2003). The K^+ efflux detected in Brinley et al.'s study (1960) was detected on the exposed brain surface, so does not address the K^+-depletion we observed, although the high $[K^+]_{ex}$ from SDs is undoubtedly the first step in K^+ leaving the brain.

Possible routes of K^+-egress

Once in the subarachnoid space CSF, K^+ would have many possible efflux routes. These efflux routes include nerve sheaths, arachnoid villi/granulations, dural spaces adjacent to sinuses, and adventitial spaces of cranial foramina vessels (Rasmussen et al., 2022). One of the possible K^+-efflux

routes from the subarachnoid space is via the meningeal lymphatics, as suggested by a study of VEGF-C overexpression (Boisserand et al., 2024; Koh et al., 2024), and then via the cribriform plate into nasal lymphatics (Kida et al., 1993; Weller et al., 1992) and superficial and deep cervical lymph nodes (Boisserand et al., 2024; Cserr et al., 1992; Zhao et al., 2025). The K^+-efflux route to superficial, parotid, submandibular, and deep cervical lymph nodes would be similar to the egress pathway of lactate (Lundgaard et al., 2017).

The high $[K^+]_{ex}$ from SD occurrence in brain parenchyma would travel through the glia limitans superficialis into the subarachnoid space or would travel through the paravascular or glymphatic pathway (Iliff et al., 2012; Weller et al., 2009). Interstitial and CSF solutes, including K^+, travel via this glymphatic pathway from the arteriolar perivascular space through astrocytes and out AQP4-containing astrocytic endfeet into venule perivascular spaces eventually appearing in the subarachnoid space CSF or, alternatively, travel in the perivenular space to the nasal lymphatics. One feature of the glymphatic pathway that suggests K^+ can move through the astrocytic endfeet is the colocalization of Kir4.1 ion channels, and perhaps other Kir channels, with AQP4 in orthogonal arrays of intramembranous particles (Vella et al., 2015; Wolburg et al., 2009).

Although SD in normal brain has been shown to temporarily disable the glymphatic system flow as SD closes paravascular spaces (Schain et al., 2017) due to astrocytic endfeet swelling (Rosic et al., 2019), the situation for ischemic SDs might well be different. Indeed, during the SDs of ischemia, the vasoconstriction known as cortical spreading ischemia that occurs as a result of high $[K^+]_{ex}$ and low NO (Dreier et al., 1998) provide a situation for increases in paravascular spaces (Mestre et al., 2020). This very early increase in perivascular space after ischemic onset (Mestre et al., 2020) occurs in the same time period during which K^+-depletion occurs (before our 2.5 hour first data point in this study). However, this early influx of Na^+ and water via the glymphatic system speaks against this route also carrying the excessive increase in $[K^+]_{ex}$ out of the brain unless its bidirectional or the efflux follows at a later time, but still within the 2.5 h that we show excessive K^+ efflux in Fig 4. Even though the glymphatic pathway is eventually compromised at 3 h after ischemic stroke (Gaberel et al., 2014),

during the early acute period activation of the superficial and deep cervical lymph nodes has been demonstrated at 3 h after experimental stroke in rats (Esposito et al., 2019).

We suggest that our use of 1% isoflurane plus 70% nitrous oxide did not interfere with the glymphatic system transport (Benveniste et al., 2017; Hablitz et al., 2019) even though higher concentrations of 2.5% have been shown to effect this system (Gakuba et al., 2018).

The slower decrease in K^+-egress shown in Fig. 4 in the ICp-DP region after 2.5 h could be related to the dysfunction of the glymphatic system at this time even as other K^+-egress systems are working. After MCA occlusion via ferric chloride initiated thrombosis, Gaberel et al. (2014) showed impaired glymphatic flow at 3 h. This study (Gaberel et al., 2014) suggests that linear K^+-egress after 2.5 h Is not associated with the glymphatic pathway.

The transcellular route through endothelial cells via abluminal (Chen et al., 2005; O'Donnell et al., 2004) and luminal (Vigne et al., 1989) K^+ channels/transporters, has been shown to be a possible K^+ exit route, in addition to an Na^+ and edema entrance route, within a 3 h time period after stroke onset (O'Donnell et al., 2004). This BBB-dependent route would lead to venous clearance of K^+ from the peripheral ischemic core regions with more collateral flow potential. One other K^+ exit possibility is not active in the initial time period of 5 h after stroke: The paracellular route via disrupted tight junctions in the endothelium (Asahi et al., 2001) for edema entry takes two days to become active (Knowland et al., 2014).

Varying degrees of K^+-depletion from the peripheral ischemic core could suggest regional differences in K^+-clearance that could be due to either more SDs and more $[K^+]_{ex}$ or differential access to K^+-efflux routes. Regional differences in $[K^+]_{ex}$ are certainly possible as watershed areas of marginal collateral circulation potential would have more SDs and higher $[K^+]_{ex}$ levels compared to regions with more vibrant collateral potential (von Bornstadt et al., 2015). It might be that some of the K^+ nondepleted regions are near the regions distal to the transected MCA which might impair access to the initial glymphatic system pathway via the arterial paravascular space.

Limited study of brain waste product efflux

Although all of these are potential K^+ egress routes, K^+ has not been a subject of their many investigations extensively summarized by Rasmussen et al. (2022). "Only a limited number of endogenous brain waste solutes" (e.g., Ab (Iliff et al., 2012), tau (Patel et al., 2019), and lactate (Ball et al., 2010; Lundgaard et al., 2017)) "have been shown to clear via the glymphatic system" (Zhao et al., 2022). To our knowledge, the only specific mention of K^+ clearance by the glymphatic system in the ischemic core is made in a figure caption of a editorial comment (Hablitz et al., 2024) although clearance of "solutes" (which of course include amyloid-beta as well as K^+) is widely discussed.

Three dimensional depiction of K^+-depletion (Yushmanov et al., 2013)

The histological section ribbon analysis decrements in $[K^+]_{br}$ as shown in Fig. 2D and 2E have been previously presented three-dimensionally (Yushmanov et al., 2013). In the three-dimensional depiction of Figure 6, these lower $[K^+]_{br}$ in the peripheral ischemic core form a torus-like region of low $[K^+]_{br}$ surrounding the ischemic core. This region of K^+-depletion in the peripheral ischemic core surrounds the ischemic core with a torus-like structure.

Conclusion

Quantitative histo-K^+ analysis after acute experimental focal ischemia revealed a greater decrease of potassium in the some peripheral ischemic core regions than in the central ischemic core, possibly related to their proximity to K^+-efflux pathways. It should be clear that although we have demonstrated excessive regional variation in the well-known loss of K^+ from the brain after ischemic stroke, it is unresolved as to what route, of the many possible, this K-egress takes. In addition, differences in these potential K^+-egress pathways between rodent models and humans related to parasagittal dural lymphatics may exist (Ringstad et al., 2020). These within-ischemic-core differences are important because the relatively greater K^+-depletion from the peripheral ischemic core in comparison to the central ischemic core would temporally limit K^+ availability for the further SD depolarizations that suggests a limit to SD-mediated secondary brain injury and would make neuronal recovery less likely

even if reperfusion occurs.

Author contributions

AK developed the staining system and implemented the arterial occlusion model. AK, VEY, and SCJ performed the experimental procedures. VEY wrote the first draft and contributed additional analysis. KAE conceived of and performed the statistical analysis. SCJ, AK, and VEY edited the manuscript. SCJ conceived of the project, provided supervision, analyzed data, prepared figures, and extensively revised the manuscript,

Funding

This project was conducted at the Allegheny-Singer Research Institute, the previous work place of the authors. This work was partially supported by grants from the: US Public Health Service National Institutes of Health: NS30839, NS30839-14S1 and NS66292 to SCJ while at the Allegheny-Singer Research Institute; and Small Business Innovative Research grants 5R43NS092181 and 3R43NS092181-02S1 to SCJ for CerebroScope.

Declaration of Competing Interests

Stephen C. Jones and Alexander Kharlamov are founding partners and shareholders of CerebroScope.

Compliance with Ethical Standards

All ethical standards have been met. See section "Human study issues".

Data and Code Availability

Available upon request based on the submission and suitability of a formal project outline.

Acknowledgements

We are grateful for the technical assistance provided by Jayjayantee Dasgupta.

References

1.   Andrew RD, Hartings JA, Ayata C, Brennan KC, Dawson-Scully KD, Farkas E, Herreras O, Kirov SA, Müller M, Ollen-Bittle N, Reiffurth C, Revah O, Robertson RM, Shuttleworth CW, Ullah G, Dreier JP. The Critical Role of Spreading Depolarizations in Early Brain Injury: Consensus and Contention. *Neurocrit Care.* 2022;37:83-101. https://doi.org/10.1007/s12028-021-01431-w

2.   Asahi M, Wang X, Mori T, Sumii T, Jung JC, Moskowitz MA, Fini ME, Lo EH. Effects of matrix metalloproteinase-9 gene knock-out on the proteolysis of blood-brain barrier and white matter components after cerebral ischemia. *J Neurosci.* 2001;21:7724-7732. https://doi.org/10.1523/jneurosci.21-19-07724.2001

3.   Astrup J, Symon L, Branston NM, Lassen NA. Cortical evoked potential and extracellular K^+ and H^+ at critical levels of brain ischemia. *Stroke.* 1977;8:51-57. https://doi.org/10.1161/01.str.8.1.51

4.   Atkinson IC, Claiborne TC, Thulborn KR. Feasibility of 39-potassium MR imaging of a human brain at 9.4 Tesla. *Magn Reson Med.* 2014;71:1819-1825. https://doi.org/10.1002/mrm.24821

5.   Ball KK, Cruz NF, Mrak RE, Dienel GA. Trafficking of glucose, lactate, and amyloid-beta from the inferior colliculus through perivascular routes. *J Cereb Blood Flow Metab.* 2010;30:162-176. https://doi.org/10.1038/jcbfm.2009.206

6.   Benveniste H, Lee H, Ding F, Sun Q, Al-Bizri E, Makaryus R, Probst S, Nedergaard M, Stein EA, Lu H. Anesthesia with Dexmedetomidine and Low-dose Isoflurane Increases Solute Transport via the Glymphatic Pathway in Rat Brain When Compared with High-dose Isoflurane. *Anesthesiology.* 2017;127:976-988. https://doi.org/10.1097/aln.0000000000001888

7.   Betz AL, Coester HC. Effect of steroids on edema and sodium uptake of the brain during focal ischemia in rats. *Stroke.* 1990;21:1199-1204. https://doi.org/10.1161/01.str.21.8.1199

8.   Betz AL, Keep RF, Beer ME, Ren X. Blood-brain barrier permeability and brain concentration of sodium, potassium, and chloride during focal ischemia. *J Cereb Blood Flow Metab.* 1994;14:29-37. https://doi.org/10.1038/jcbfm.1994.5

9.   Bischof H, Rehberg M, Stryeck S, Artinger K, Eroglu E, Waldeck-Weiermair M, Gottschalk B, Rost R, Deak AT, Niedrist T et al. Novel genetically encoded fluorescent probes enable real-time detection of potassium in vitro and in vivo. *Nat Commun.* 2017;8:1422. https://doi.org/10.1038/s41467-017-01615-z

10.   Boisserand LSB, Geraldo LH, Bouchart J, El Kamouh MR, Lee S, Sanganahalli BG, Spajer M, Zhang S, Lee S, Parent M et al. VEGF-C prophylaxis favors lymphatic drainage and modulates neuroinflammation in a stroke model. *J Exp Med.* 2024;221. https://doi.org/10.1084/jem.20221983

11.   Branston NM, Strong AJ, Symon L. Extracellular potassium activity, evoked potential and tissue blood flow. Relationships during progressive ischaemia in baboon cerebral cortex. *J Neurol Sci.* 1977;32:305-321. https://doi.org/10.1016/0022-510x(77)90014-4

12.   Brinley FJ, Jr., Kandel ER, Marshall WH. Potassium outflux from rabbit cortex during spreading depression. *J Neurophysiol.* 1960;23:246-256. https://doi.org/10.1152/jn.1960.23.3.246

13.   Chen H, Sun D. The role of Na-K-Cl co-transporter in cerebral ischemia. *Neurol Res.* 2005;27:280-286. https://doi.org/10.1179/016164105x25243

14.   Chen ST, Hsu CY, Hogan EL, Maricq H, Balentine JD. A model of focal ischemic stroke in the rat: reproducible extensive cortical infarction. *Stroke.* 1986;17:738-743.

15.   Cserr HF, DePasquale M, Nicholson C, Patlak CS, Pettigrew KD, Rice ME. Extracellular volume decreases while cell volume is maintained by ion uptake in rat brain during acute hypernatremia. *J Physiol (Lond).* 1991;442:277-295.

16.   Cserr HF, Harling-Berg CJ, Knopf PM. Drainage of brain extracellular fluid into blood and deep cervical lymph and its immunological significance. *Brain Pathol.* 1992;2:269-276. https://doi.org/10.1111/j.1750-3639.1992.tb00703.x

17.   Dawson DA, Hallenbeck JM. Acute focal ischemia-induced alterations in MAP2 immunostaining: description of temporal changes and utilization as a marker for volumetric assessment of acute brain injury. *J Cereb Blood Flow Metab.* 1996;16:170-174. https://doi.org/10.1097/00004647-199601000-00020

18.   Del Zoppo GJ, Sharp FR, Heiss WD, Albers GW. Heterogeneity in the penumbra. *J Cereb Blood Flow Metab.* 2011;31:1836-1851. https://doi.org/10.1038/jcbfm.2011.93

19.   Dickinson LD, Betz AL. Attenuated development of ischemic brain edema in vasopressin-deficient rats. *J Cereb Blood Flow Metab.* 1992;12:681-690. https://doi.org/10.1038/jcbfm.1992.93

20.   Dreier JP, Korner K, Ebert N, Gorner A, Rubin I, Back T, Lindauer U, Wolf T, Villringer A, Einhaupl KM, Lauritzen M, Dirnagl U. Nitric oxide scavenging by hemoglobin or nitric oxide synthase inhibition by N-nitro-L-arginine induces cortical spreading ischemia when K+ is increased in the subarachnoid space. *J Cereb Blood Flow Metab.* 1998;18:978-990. https://doi.org/10.1097/00004647-199809000-00007

21.   Elabyad IA, Kalayciyan R, Shanbhag NC, Schad LR. First in vivo potassium-39 (³⠹K) MRI at 9.4 T using conventional copper radio frequency surface coil cooled to 77 K. *IEEE Trans Biomed Eng.* 2014;61:334-345. https://doi.org/10.1109/tbme.2013.2294277

22.   Esposito E, Ahn BJ, Shi J, Nakamura Y, Park JH, Mandeville ET, Yu Z, Chan SJ, Desai R, Hayakawa A, Ji X, Lo EH, Hayakawa K. Brain-to-cervical lymph node signaling after stroke. *Nat Commun.* 2019;10:5306. https://doi.org/10.1038/s41467-019-13324-w

23. Gaberel T, Gakuba C, Goulay R, Martinez De LS, Hanouz JL, Emery E, Touze E, Vivien D, Gauberti M. Impaired glymphatic perfusion after strokes revealed by contrast-enhanced MRI: a new target for fibrinolysis? *Stroke.* 2014;45:3092-3096. https://doi.org/10.1161/strokeaha.114.006617

24. Gakuba C, Gaberel T, Goursaud S, Bourges J, Di PC, Quenault A, Martinez De LS, Vivien D, Gauberti M. General Anesthesia Inhibits the Activity of the "Glymphatic System". *Theranostics.* 2018;8:710-722. https://doi.org/10.7150/thno.19154

25. Grochowski C, Blicharska E, Krukow P, Jonak K, Maciejewski M, Szczepanek D, Jonak K, Flieger J, Maciejewski R. Analysis of Trace Elements in Human Brain: Its Aim, Methods, and Concentration Levels. *Front Chem.* 2019;7:115. https://doi.org/10.3389/fchem.2019.00115

26. Hablitz L, Nedergaard M. Synchronized neuronal activity drives waste fluid flow. *Nature.* 2024;627:44-45. https://doi.org/10.1038/d41586-024-00422-z

27. Hablitz LM, Vinitsky HS, Sun Q, Stæger FF, Sigurdsson B, Mortensen KN, Lilius TO, Nedergaard M. Increased glymphatic influx is correlated with high EEG delta power and low heart rate in mice under anesthesia. *Sci Adv.* 2019;5:eaav5447. https://doi.org/10.1126/sciadv.aav5447

28. Hansen AJ, Olsen CE. Brain extracellular space during spreading depression and ischemia. *Acta Physiol Scand.* 1980;108:355-365. https://doi.org/10.1111/j.1748-1716.1980.tb06544.x

29. Hartings JA, Rolli ML, Lu XC, Tortella FC. Delayed secondary phase of peri-infarct depolarizations after focal cerebral ischemia: relation to infarct growth and neuroprotection. *J Neurosci.* 2003;23:11602-11610. https://doi.org/10.1523/jneurosci.23-37-11602.2003

30. Hu W, Kharlamov A, Wang Y, Perez-Trepichio AD, Jones SC. Directed sampling for electrolyte analysis and water content of micro-punch samples shows large differences between normal and ischemic rat brain cortex. *Brain Res.* 2000;868:370-375. https://doi.org/10.1016/s0006-8993(00)02360-x

31. Iliff JJ, Wang M, Liao Y, Plogg BA, Peng W, Gundersen GA, Benveniste H, Vates GE, Deane R, Goldman SA, Nagelhus EA, Nedergaard M. A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid beta. *Sci Transl Med.* 2012;4:147ra111. https://doi.org/10.1126/scitranslmed.3003748

32. Ito U, Ohno K, Nakamura R, Suganuma F, Inaba Y. Brain edema during ischemia and after restoration of blood flow: measurement of water, sodium, potassium content and plasma protein permeability. *Stroke.* 1979;10:542-547. https://doi.org/10.1161/01.str.10.5.542

33. Jones SC, Kharlamov A, Yanovski B, Kim DK, Easley KA, Yushmanov VE, Ziolko SK, Boada FE. Stroke onset time using sodium MRI in rat focal cerebral ischemia. *Stroke.* 2006;37:883-888. https://doi.org/10.1161/01.str.0000198845.79254.0f

34.   Kato H, Kogure K, Sakamoto N, Watanabe T. Greater disturbance of water and ion homeostasis in the periphery of experimental focal cerebral ischemia. *Exp Neurol.* 1987;96:118-126. https://doi.org/10.1016/0014-4886(87)90173-7

35.   Kharlamov A, Brown BR, Easley KA, Jones SC. Heterogeneous response of cerebral blood flow to hypotension demonstrated by laser speckle imaging flowmetry in rats. *Neurosci Lett.* 2004;368:151-156. https://doi.org/10.1016/j.neulet.2004.06.079

36.   Kharlamov A, Kim DK, Jones SC. Early visual changes in reflected light on non-stained brain sections after focal ischemia mirror the area of ischemic damage. *J Neurosci Methods.* 2001;111:67-73. https://doi.org/10.1016/s0165-0270(01)00444-7

37.   Kida S, Pantazis A, Weller RO. CSF drains directly from the subarachnoid space into nasal lymphatics in the rat. Anatomy, histology and immunological significance. *Neuropathol Appl Neurobiol.* 1993;19:480-488. https://doi.org/10.1111/j.1365-2990.1993.tb00476.x

38.   Knowland D, Arac A, Sekiguchi KJ, Hsu M, Lutz SE, Perrino J, Steinberg GK, Barres BA, Nimmerjahn A, Agalliu D. Stepwise recruitment of transcellular and paracellular pathways underlies blood-brain barrier breakdown in stroke. *Neuron.* 2014;82:603-617. https://doi.org/10.1016/j.neuron.2014.03.003

39.   Koh GY, McDonald DM. Meningeal lymphatics can influence stroke outcome. *J Exp Med.* 2024;221. https://doi.org/10.1084/jem.20232305

40.   Kumagai T, Walberer M, Nakamura H, Endepols H, Sue M, Vollmar S, Adib S, Mies G, Yoshimine T, Schroeter M, Graf R. Distinct spatiotemporal patterns of spreading depolarizations during early infarct evolution: evidence from real-time imaging. *J Cereb Blood Flow Metab.* 2011;31:580-592. https://doi.org/10.1038/jcbfm.2010.128

41.   Lo WD, Betz AL, Schielke GP, Hoff JT. Transport of sodium from blood to brain in ischemic brain edema. *Stroke.* 1987;18:150-157. https://doi.org/10.1161/01.str.18.1.150

42.   Lundgaard I, Lu ML, Yang E, Peng W, Mestre H, Hitomi E, Deane R, Nedergaard M. Glymphatic clearance controls state-dependent changes in brain lactate concentration. *J Cereb Blood Flow Metab.* 2017;37:2112-2124. https://doi.org/10.1177/0271678x16661202

43.   Menzies SA, Hoff JT, Betz AL. Extravasation of albumin in ischaemic brain oedema. *Acta Neurochir Suppl (Wien).* 1990;51:220-222. https://doi.org/10.1007/978-3-7091-9115-6_74

44.   Mestre H, Du T, Sweeney AM, Liu G, Samson AJ, Peng W, Mortensen KN, Staeger FF, Bork PAR, Bashford L et al. Cerebrospinal fluid influx drives acute ischemic tissue swelling. *Science.* 2020;367:eaax7171. https://doi.org/10.1126/science.aax7171

45.   Mies G, Kloiber O, Drewes LR, Hossmann K-A. Cerebral blood flow and regional potassium distribution during focal ischemia of gerbil brain. *Ann Neurol.* 1984;16:232-237. https://doi.org/10.1002/ana.410160212

46. Monai H, Koketsu S, Shinohara Y, Ueki T, Kusk P, Hauglund NL, Samson AJ, Nedergaard M, Hirase H. Adrenergic inhibition facilitates normalization of extracellular potassium after cortical spreading depolarization. *Sci Rep.* 2021;11:8150. https://doi.org/10.1038/s41598-021-87609-w

47. Nakamura H, Strong AJ, Dohmen C, Sakowitz OW, Vollmar S, Sue M, Kracht L, Hashemi P, Bhatia R, Yoshimine T, Dreier JP, Dunn AK, Graf R. Spreading depolarizations cycle around and enlarge focal ischaemic brain lesions. *Brain.* 2010;133:1994-2006. https://doi.org/10.1093/brain/awq117

48. Nicholson C, Phillips JM. Ion diffusion modified by tortuosity and volume fraction in the extracellular microenvironment of the rat cerebellum. *J Physiol.* 1981;321:225-257.

49. O'Donnell ME, Tran L, Lam TI, Liu XB, Anderson SE. Bumetanide inhibition of the blood-brain barrier Na-K-Cl cotransporter reduces edema formation in the rat middle cerebral artery occlusion model of stroke. *J Cereb Blood Flow Metab.* 2004;24:1046-1056. https://doi.org/10.1097/01.wcb.0000130867.32663.90

50. Octeau JC, Faas G, Mody I, Khakh BS. Making, Testing, and Using Potassium Ion Selective Microelectrodes in Tissue Slices of Adult Brain. *J Vis Exp.* 2018. https://doi.org/10.3791/57511

51. Pappius HM, Elliott KAC. Water distribution in incubated slices of brain and other tissues. *Can J Biochem Physiol.* 1956;34:1007-1022. https://doi.org/10.1139/y56-106

52. Patel TK, Habimana-Griffin L, Gao X, Xu B, Achilefu S, Alitalo K, McKee CA, Sheehan PW, Musiek ES, Xiong C, Coble D, Holtzman DM. Dural lymphatics regulate clearance of extracellular tau from the CNS. *Mol Neurodegener.* 2019;14:11. https://doi.org/10.1186/s13024-019-0312-x

53. Pettigrew LC, Holtz ML, Craddock SD, Minger SL, Hall N, Geddes JW. Microtubular proteolysis in focal cerebral ischemia. *J Cereb Blood Flow Metab.* 1996;16:1189-1202. https://doi.org/10.1097/00004647-199611000-00013

54. Rasmussen MK, Mestre H, Nedergaard M. Fluid transport in the brain. *Physiol Rev.* 2022;102:1025-1151. https://doi.org/10.1152/physrev.00031.2020

55. Ricks R, MacKenzie-Gray SC, Trevelyan AJ, Parrish RR. Mechanistic insight into the initiation of spreading depolarizations: Is it really all about potassium? *J Physiol.* 2025;603:3269-3273. https://doi.org/10.1113/jp288809

56. Ringstad G, Eide PK. Cerebrospinal fluid tracer efflux to parasagittal dura in humans. *Nat Commun.* 2020;11:354. https://doi.org/10.1038/s41467-019-14195-x

57. Rosic B, Dukefoss DB, Åbjørsbråten KS, Tang W, Jensen V, Ottersen OP, Enger R, Nagelhus EA. Aquaporin-4-independent volume dynamics of astroglial endfeet during cortical spreading depression. *Glia.* 2019;67:1113-1121. https://doi.org/10.1002/glia.23604

58. Schain AJ, Melo-Carrillo A, Strassman AM, Burstein R. Cortical Spreading Depression Closes Paravascular Space and Impairs Glymphatic Flow: Implications for Migraine Headache. *J Neurosci.* 2017;37:2904-2915. https://doi.org/10.1523/jneurosci.3390-16.2017

59. Schuier FJ, Hossmann K-A. Experimental brain infarcts in cats. II. Ischemic brain edema. *Stroke.* 1980;11:593-601. https://doi.org/10.1161/01.str.11.6.593

60. Shibata S, Hodge CP, Pappius HM. Effect of experimental ischemia on cerebral water and electrolytes. *J Neurosurg.* 1974;41:146-159. https://doi.org/10.3171/jns.1974.41.2.0146

61. Sick TJ, Feng ZC, Rosenthal M. Spatial stability of extracellular potassium ion and blood flow distribution in rat cerebral cortex after permanent middle cerebral artery occlusion. *J Cereb Blood Flow Metab.* 1998;18:1114-1120. https://doi.org/10.1097/00004647-199810000-00008

62. Silberstein C, Bouley R, Huang Y, Fang P, Pastor-Soler N, Brown D, Van Hoek AN. Membrane organization and function of M1 and M23 isoforms of aquaporin-4 in epithelial cells. *Am J Physiol Renal Physiol.* 2004;287:F501-F511. https://doi.org/10.1152/ajprenal.00439.2003

63. Simard JM, Woo SK, Schwartzbauer GT, Gerzanich V. Sulfonylurea receptor 1 in central nervous system injury: a focused review. *J Cereb Blood Flow Metab.* 2012;32:1699-1717. https://doi.org/10.1038/jcbfm.2012.91

64. Treadwell SD, Thanvi B. Malignant middle cerebral artery (MCA) infarction: pathophysiology, diagnosis and management. *Postgrad Med J.* 2010;86:235-242. https://doi.org/10.1136/pgmj.2009.094292

65. Vella J, Zammit C, Di GG, Muscat R, Valentino M. The central role of aquaporins in the pathophysiology of ischemic stroke. *Front Cell Neurosci.* 2015;9:108. https://doi.org/10.3389/fncel.2015.00108

66. Vigne P, Champigny G, Marsault R, Barbry P, Frelin C, Lazdunski M. A new type of amiloride-sensitive cationic channel in endothelial cells of brain microvessels. *J Biol Chem.* 1989;264:7663-7668. https://doi.org/10.1016/S0021-9258(18)83285-9

67. von Bornstadt D, Houben T, Seidel JL, Zheng Y, Dilekoz E, Qin T, Sandow N, Kura S, Eikermann-Haerter K, Endres M, Boas DA, Moskowitz MA, Lo EH, Dreier JP, Woitzik J, Sakadzic S, Ayata C. Supply-Demand Mismatch Transients in Susceptible Peri-infarct Hot Zones Explain the Origins of Spreading Injury Depolarizations. *Neuron.* 2015;85:1117-1131. https://doi.org/10.1016/j.neuron.2015.02.007

68. Walsh A. The application of atomic absorption spectra to chemical analysis. *Spectrochimica Acta.* 1955;7:108-117. https://doi.org/10.1016/0371-1951(55)80013-6

69. Weller RO, Djuanda E, Yow HY, Carare RO. Lymphatic drainage of the brain and the pathophysiology of neurological disease. *Acta Neuropathol.* 2009;117:1-14. https://doi.org/10.1007/s00401-008-0457-0

70.   Weller RO, Kida S, Zhang ET. Pathways of fluid drainage from the brain--morphological aspects and immunological significance in rat and man. *Brain Pathol.* 1992;2:277-284. https://doi.org/10.1111/j.1750-3639.1992.tb00704.x

71.   Wen X, Li Y, He X, Xu Y, Shu Z, Hu X, Chen J, Jiang H, Gong X. Prediction of Malignant Acute Middle Cerebral Artery Infarction via Computed Tomography Radiomics. *Front Neurosci.* 2020;14:708. https://doi.org/10.3389/fnins.2020.00708

72.   Wetterling F, Gallagher L, Macrae IM, Junge S, Fagan AJ. Regional and temporal variations in tissue sodium concentration during the acute stroke phase. *Magn Reson Med.* 2012;67:740-749. https://doi.org/10.1002/mrm.23031

73.   Woitzik J, Hecht N, Pinczolits A, Sandow N, Major S, Winkler MK, Weber-Carstens S, Dohmen C, Graf R, Strong AJ, Dreier JP, Vajkoczy P. Propagation of cortical spreading depolarization in the human cortex after malignant stroke. *Neurology.* 2013;80:1095-1102. https://doi.org/10.1212/wnl.0b013e3182886932

74.   Wolburg H, Noell S, Wolburg-Buchholz K, Mack A, Fallier-Becker P. Agrin, aquaporin-4, and astrocyte polarity as an important feature of the blood-brain barrier. *Neuroscientist.* 2009;15:180-193. https://doi.org/10.1177/1073858408329509

75.   Wu SY, Shen Y, Shkolnikov I, Campbell RE. Fluorescent Indicators For Biological Imaging of Monatomic Ions. *Front Cell Dev Biol.* 2022;10:885440. https://doi.org/10.3389/fcell.2022.885440

76.   Yang GY, Chen SF, Kinouchi H, Chan PH, Weinstein PR. Edema, cation content, and ATPase activity after middle cerebral artery occlusion in rats. *Stroke.* 1992;23:1331-1336. https://doi.org/10.1161/01.str.23.9.1331

77.   Young W, Constantini S. Ionic and water shifts in injured central nervous tissues. In: Salzman, S. K. and Faden, A. I., editors. *The Neurobiology of Central Nervous System Trauma,* New York: Oxford University Press; 1994. p. 123-130.

78.   Young W, Rappaport ZH, Chalif DJ, Flamm ES. Regional brain sodium, potassium, and water changes in the rat middle cerebral artery occlusion model of ischemia. *Stroke.* 1987;18:751-759. https://doi.org/10.1161/01.str.18.4.751

79.   Yushmanov VE, Kharlamov A, Boada FE, Jones SC. Monitoring of brain potassium with rubidium flame photometry and MRI. *Magn Reson Med.* 2007;57:494-500. https://doi.org/10.1002/mrm.21155

80.   Yushmanov VE, Kharlamov A, Boada FE, Jones SC. Transient blood-brain barrier disruption and changes in brain electrolytes at the edge of the ischemic core. *Proc Intl Soc Mag Reson Med.* 2009a;17:3280. https://archive.ismrm.org/2009/3281.html

81.   Yushmanov VE, Kharlamov A, Yanovski B, LaVerde G, Boada FE, Jones SC. Inhomogeneous sodium accumulation in the ischemic core in rat focal cerebral ischemia by ^{23}Na MRI. *J Magn Reson Imaging.* 2009b;30:18-24. https://doi.org/10.1002/jmri.21816

82. Yushmanov VE, Kharlamov A, Yanovski B, LaVerde G, Boada FE, Jones SC. Correlated sodium and potassium imbalances within the ischemic core in experimental stroke: A Na MRI and histochemical imaging study. *Brain Res.* 2013;1527:199-208. https://doi.org/10.1016/j.brainres.2013.06.012

83. Yushmanov VE, Yanovski B, Kharlamov A, LaVerde G, Boada FE, Jones SC. Sodium mapping in focal cerebral ischemia in the rat by quantitative [23]Na MRI. *J Magn Reson Imaging.* 2009c;29:962-966. https://doi.org/10.1002/jmri.21643

84. Zhao L, Tannenbaum A, Bakker ENTP, Benveniste H. Physiology of Glymphatic Solute Transport and Waste Clearance from the Brain. *Physiology (Bethesda).* 2022;37:349-362. https://doi.org/10.1152/physiol.00015.2022

85. Zhao MY, Ye CY, Liu YC, Wang XM, Fu JC, Liu XY, Zhu R, Li YZ, Tian Q. Role of meningeal lymphatic vessels in brain homeostasis. *Front Immunol.* 2025;16:1593630. https://doi.org/10.3389/fimmu.2025.1593630

Table 1

Table 1 Comparison with other measurements of $[K^+]_{br}$ normal cortex and ischemic cortex efflux rate in models of focal ischemia

Reference	Species	MCA Model[1]	Sample weight[2] (mg)	Directed sampling	Time Range (h)	$[K^+]_{br}$ Normal Cortex (mEq/kg ww)	$[K^+]_{br}$ Ischemic Cortex Efflux Rate (mEq/kg ww/h)
Ito et al. (1979)	gerbil	uniCCAo	H	No	0-3	105	-10.6
Schuier & Hossmann (1980)	cat	clip	100	Yes[3]	0-4	85	-8.0
Dickinson & Betz (1992)	rat	coagulation	39	No	0-4	91	-12.0
Yang et al. (1992)	rat	suture	50	No	0-4	113	-8.3
Young et al. (1987)	rat	transection	37	No	0-4	112	-8.1
Betz et al. (1994)	rat	coagulation	77	No	2-6	97	-9.0
Lo et al. (1987)	gerbil	uniCCAo	H	No	0-3	94	-9.7
Betz & Coester (1990)	rat	coagulation	77	No	0-4	103	-14.1
Shibata et al. (1974)	dog	clip & hypotension	75	Yes[4]	0-5	108	-10.3
Menzies et al. (1990)	rat	coagulation	77	No	0-3	97	-11.0
Mean ± sem			66 ± 8			101 ± 3	-10.1 ± 0.6
This study	rat	MCAt & biCCAo	0.545	Yes[5]	2.5-5	97 ± 2	-12.2 ± 1.7

[1] uni - unilateral, bi - bilateral, MCAt & biCCAo - middle cerebral artery transection and bilateral common carotid artery occlusion
[2] H - hemisphere was used
[3] India ink or Evans blue
[4] Fluorescein angiography
[5] Reflective change

Table 2

Table 2 Physiological variables at various experimental stages					
Time	MABP (mmHg)	pH (-)	$PaCO_2$ (mmHg)	PaO_2 (mmHg)	CBF (%)
Before ischemia	108±2	7.422±0.009	36±1	98±3	100
~3 min after ischemia	115±2	7.391±0.016	36±1	95±4	16±3
120 min after ischemia	104±3	7.413±0.017	35±1	97±8	25±7
End of experiment	104±2	7.400±0.017	34±1	101±9	28±7*

MABP, mean arterial blood pressure; CBF, cerebral blood flow
*, $p = 0.063$ from the immediate post occlusion value at ~3 min (n = 11)
Mean ± sem, n = 11 to 13

Table 3

Table 3 Estimates of $[K^+]_{in}$ in the central ischemic core

Parameter	Units	Time after ischemia (h)				Source
		0	0.5	1.5	3	
ECS	(-)	0.21	0.11	0.11	0.11	(Hansen et al., 1980; Nicholson et al., 1981)
$[K^+]_{br}$	(mEq/kg)	96	89	76	55	This study
$[K^+]_{ex}$	(mEq/kg)*	3.0	60	58	57	Sick et al. (1998)
$[K^+]_{in}$	(mEq/kg)	128	95	79	56	Calculated ***
K^+ Nernst potential	(mV)	-100	-17	-13	-5	

ECS, extracellular space fraction

*, assume mM = mEq/kg

**, Estimated

***, From Cserr et al. (1991)

Figures with legends

Figure 1

Figure 1   **(A)** A dorsal view of a typical rat brain with bregma and lambda showing: 1) the selection of the 2-mm-thick slices for the first and second series of micropunch samples at two levels, L1 and L2; 2) the three positions, S1, S2, and S3, for the 35-µm sections taken for K^+-staining and cortical ribbon analysis, MAP2 IMR, and reflective change; and 3) the laser Doppler flowmeter (LDF) probe position. The dashed line indicates the approximate range of the middle cerebral artery distribution that typifies the infarct area.   **(B)** Histo-K^+-stained coronal section showing the ischemic core (IC, violet), as confirmed by reflective change and MAP2 IMR in adjacent sections, and its division into regions-of-interest for dorsal peripheral (ICpd, light blue), ventral peripheral (ICpv, blue), and central ischemic core (ICc, red) regions, as identified by variations in $[K^+]_{br}$. The ischemic penumbra would be located just outside the dorsal and ventral peripheral ischemic core in the regions where $[K^+]_{br}$ is approaching normal, non-ischemic levels (see Fig. 3H).

Figure 2

Figure 2 The sequence of brain processing steps and other procedures for adjacent sections and images at the S1 level (see Figure 1). All images are aligned and processed digitally. In panels **A**, **B**, **C**, and **G** the blue dashed line indicates the infarct area. **(A)** The reflective change in a coronal brain section used to guide the micropunch locations to normal cortex, ischemic cortex, and subcortical regions. **(B)** The cut-face of the brain in the cryostat showing the micropunch hole positions used for ROI placement in panel **C**. The mm scale placed at the brain section level is at the top of the Figure. **(C)** Histo-K$^+$-section showing the micropunch ROIs (red circles) obtained from panel **B** that are used to create the calibration (see panel **D**) of the gray value (GV) by the flame photometry-determined [K$^+$]$_{br}$ in each micropunch. **(D)** The calibration curve from the linear regression of micropunch ROI GVs vs. flame photometry [K$^+$]$_{br}$ values (circles) was used to convert the GVs of the histochemical image to [K$^+$]$_{br}$. The [K$^+$]$_{br}$ values obtained from the GVs of the cortical ribbon ROIs (panels **F** and **H**) are shown along the curve as colored diamond symbols. **(E)** Pseudo-color coded histo-K$^+$-image based on the calibration curve shown in panel **D** with the color representing [K$^+$]$_{br}$ in mEq/kg as shown on the calibration bar. The red line in the ischemic right cortical region indicates the 36 mEq/kg isocontour which is the K$^+$-depletion breakpoint defined in Figure 3. **(F)** The histo-K$^+$ image shows the cortical ribbon for the normal and ischemic hemispheres (black lines) with the ROIs for dorsal normal cortex (NCd, light green), dorsal peripheral ischemic core (ICpd, light blue), central ischemic core (ICc, red),

peripheral ventral ischemic core (ICpv, dark blue), and ventral normal cortex (NCv, dark green). **(G)** Section stained for MAP2 IMR to confirm the extent of the ischemic lesion and to assess possible MAP2 IMR decrements in the peripheral ischemic core for comparison to K^+-depletion. **(H)** The $[K^+]_{br}$ profiles along the cortical ribbon for the ischemic (IH-ribbon, orange curve) and non-ischemic hemispheres (NH-ribbon, dotted-green curve). The horizontal lines indicate the position of the five ROIs (**F**, described above) along the ischemic hemisphere cortical ribbon and their $[K^+]_{br}$ values. The dorsal (ICpd) and ventral (ICpv) peripheral ischemic core regions, located at the edges of the ICc, demonstrated larger decreases of $[K^+]_{br}$ than the decrease in the ICc. In this animal, the ICpd and ICpv regions were 72% and 84% of the ICc value, respectively, so the ICpd region, but not the ICpv, was classified as K^+-depleted because it was <78%, the K^+-depletion breakpoint, as described in the text and shown in Figure 3.

Figure 3

Figure 3 Histogram analyses of [K$^+$]$_{br}$ in the combined peripheral ischemic core (ICpd and ICpv) and central ischemic core (ICc) cortical-ribbon ROIs. Frequency analysis of % changes of [K$^+$]$_{br}$ in the combined peripheral ischemic core regions (black squares and line, n = 50) revealed a bimodal distribution with medians of 64% (light blue line) and 90% (dark blue line). The minimum between the two peaks of the summed distribution (green line) was chosen as the breakpoint (78%, vertical red line) to distinguish the K$^+$-depleted peripheral ischemic core regions from those that retained [K$^+$]$_{br}$ similar to the ICc values.

Figure 4

Figure 4 Time course of $[K^+]_{br}$ (mean ± sem) from the cortical ribbon ROIs in normal cortex (NC), central ischemic core (ICc) and peripheral ischemic core (ICp) regions between 2.5 and 5 h with their linear regression curves. The ICpd and ICpv peripheral ischemic core regions have been combined, but then divided into K^+-depleted (ICp-DP) and non-K^+-depleted (ICp-ND) groups on the basis of their $[K^+]$ expressed as a percent of the ICc value. This dividing line, or breakpoint, is shown converted to mEq/kg as the horizontal dashed line at 36 mEq/kg. The time course of the combined NCd and NCv regions (NC, green line) shows stability of $[K^+]_{br}$ with a slope not significantly different from zero, whereas the ICc (red line), ICp-DP (light blue line), and ICp-ND (dark blue dotted line) time courses all show a significant decrease over time, with the slopes of the ICp-ND regions and the ICc being similar (p =0.99). The NC, ICc, and ICp-ND curves have been extrapolated to the stroke onset time, Ta = 0. The slope of the ICp-DP region is significantly different from that of the ICc (p = 0.010) and the ICp-ND (p = 0.0071) curves. This lower slope of the ICp-DP curve 2.5 h after Ta is due to the accelerated K^+-efflux from Ta = 0 to 2.5 h, as its value at Ta = 0 must be ~100 mEq/kg.

Figure 5

Figure 5 Analysis of $[K^+]_{br}$ and MAP2 IMR from ROIs in the normal and ischemic hemispheric cortical ribbons for the purpose of testing whether the ICpd and ICpv regions differ in their K^+-depletion and MAP2 IMR. The normal hemisphere values (mean ± sem) are plotted as the solid lines with filled squares. The ischemic cortical ribbon ROIs have been divided into those whose ICpd and ICpv ROIs are K^+-depleted (solid bars with n's shown inside each bar) and not K^+-depleted (stripped bars) for both panel **A**, the MAP2 normalized IMR values and panel **B**, showing the $[K^+]_{br}$ values. Thus the "K^+-depleted" normal cortex values are associated with the K^+-depleted ICpd and ICpv ROIs.

(A) The MAP2 IMR normalized to the gray value in normal cortex shows no difference between the ICpd and ICpv K^+-depleted and non-K^+-depleted values (unpaired t-test, p = 0.83, p = 0.16, respectively), but a significant decrease in ICpd and ICpv vs. ICc IMR (paired t-test, p < 0.001 for both).

(B) A significant $[K^+]_{br}$ decrease in the K^+-depleted peripheral ischemic core regions (ICpd and ICpv) in comparison to the $[K^+]_{br}$ in the central ischemic core (ICc) was observed (two-tailed paired t-test, p < 0.001 for both), and in comparison to the non-K^+-depleted regions (unpaired two-tailed t-test, p < 0.001, p = 0.0021, respectively). However, there was no significant difference between the non-K^+-depleted and ICc values.

Abbreviations: NCd, dorsal, and NCv, ventral normal cortex of ischemic hemisphere; ICc, the central ischemic core; ICpd, dorsal peripheral ischemic core; and ICpv, ventral peripheral ischemic core.

Figure 6

Figure 6 3D reconstruction of a rat brain at 4.5 h after an intraluminal suture stroke from 21 stacked histo-K$^+$-stained coronal slices (0.7-mm separation) as viewed from the front-right perspective. Regions with $[K^+]_{br}$ below 36 mEq/kg are shown in yellow color to highlight the peripheral ischemic core surrounding the central ischemic core. The rostral K$^+$-loss in the peripheral ischemic core is less intense and narrower than at the caudal, ventral, and dorsal peripheral ischemic core. Permission requested from (Yushmanov et al., 2013).

Supplemental Material

Supplement to: Kharlamov et al. "Heterogeneity of Brain Potassium Depletion in Experimental Focal Ischemia"

Non-standard Abbreviations and Acronyms

$[K^+]_{br}$ = brain tissue potassium concentration (mEq/kg)

$[K^+]_{ex}$ = extracellular potassium concentration (mM)

$[K^+]IC$ = $[K^+]_{br}$ of the ischemic cortex (mEq/kg)

$[K^+]ICc$ = $[K^+]_{br}$ of the central ischemic cortex (mEq/kg)

$[K^+]ICp$ = $[K^+]_{br}$ of the peripheral ischemic core (mEq/kg)

$[K^+]ICp\%$ = 100 x $[[K^+]ICp / [K^+]ICc$ (%)

$[Na^+]_{br}$ = brain tissue sodium concentration (mEq/kg)

BBB = Blood-Brain Barrier

CBF = Cerebral Blood Flow

CSF = CerebroSpinal Fluid

SD = cortical Spreading Depolarization

Gd-DTPA = gadolinium diethylenetriamine-pentaacetic acid

GV = gray value

Histo-K^+-stained = histochemically stained for K^+

ICc = central ischemic core

ICp = peripheral ischemic core

ICpd = dorsal peripheral ischemic core

ICpv = ventral peripheral ischemic core

ICp-DP = K^+-depleted peripheral ischemic core

ICp-ND = non-K^+-depleted peripheral ischemic core

IMR = immunoreactivity

LDF = laser Doppler flowmetry

MABP = mean arterial blood pressure

MAP2 = microtubule-associated protein 2

MCAt & biCCAo = middle cerebral artery transection and bilateral common carotid artery occlusion

NC = normal cortex

NCd = dorsal normal cortex

NCv = ventral normal cortex

PVS = perivascular spaces

ROI = regions-of-interest

Ta = time from stroke onset (h)

Used in figures only:

IH-ribbon = ischemic hemisphere cortical ribbon

NH-ribbon = non-ischemic hemisphere cortical ribbon

GLS – glia limitans superficialis

TD – terminal depolarization