Cytoglobin regulates ventricular morphogenesis and diastolic function through sGC-cGMP signaling during development.

Adam Austin Clark[1], Rasmus Hejlesen[3], Tzu-Ting, Weng[1], Muddassar Iqbal[2], Akueba Bruce[2], Paola Corti[1#]

[1]Department of Biochemistry and Molecular Biology, University of Maryland School of Medicine, Baltimore, MD 21201

[2]Department of Medicine, University of Maryland School of Medicine, Baltimore, MD 21201

[3]MRC Human Genetics Unit, Institute of Genetics and Cancer, The University of Edinburgh, Edinburgh, EH4 2XU, UK

#Correspondence: Corti, Paola. Email: PCorti@som.umaryland.edu

Abstract

Aims. Hypoplastic left heart syndrome (HLHS) is a severe congenital heart disease characterized by ventricular hypoplasia and impaired cardiac function. Clinically, inhaled nitric oxide (NO) therapy is used to reduce pulmonary vascular resistance and improve cardiopulmonary stability in HLHS patients. However, whether NO signaling contributes to HLHS pathogenesis remains unknown. Cytoglobin (CYGB) is a heme protein traditionally thought to limit NO bioavailability. Unexpectedly, our recent work shows that CYGB/Cygb enhances NO signaling through activation of the nitric oxide synthase- soluble guanylate cyclase (sGC)- cyclic guanosine monophosphate (cGMP) signaling pathway. In zebrafish embryos, Cygb-dependent NO signaling is required for normal cilia motility and for the establishment of correct cardiac laterality. Here, our aim was to determine whether Cygb-dependent NO-sGC signaling linked to cilia function regulates cardiac morphogenesis and contributes to ventricular hypoplasia in HLHS.

Methods and Results. We found that loss of *Cygb* (*cygb2*) in zebrafish disrupts NO-sGC signaling

during cardiogenesis, altering cardiac progenitor organization and migration within the anterior lateral

plate mesoderm (ALPM). Disruption of these processes impairs heart tube morphogenesis, thereby

producing a compact ventricle with increased wall thickness despite preserved cardiomyocyte number,

reduced ventricle size and decreased stroke volume, recapitulating key features of HLHS. Genetic

disruption of the sGC α-subunit (*gucy1a1)* and pharmacological NO scavenging phenocopy the *cygb2*

mutant phenotype, resulting in reduced cGMP levels, compact ventricular architecture and decreased

stroke volume (SV). Consistently, restoration of NO-sGC signaling in *cygb2* mutants rescues early

cardiac progenitor patterning, ventricular morphology and SV.

Conclusions. These findings identify Cygb-dependent NO-sGC signaling as a critical developmental

pathway for ventricular development and performance, temporally linking cardiac progenitor dynamics

to cilia-dependent signaling associated with left-right patterning. This study further suggests that

pharmacological activation of sGC may provide a therapeutic strategy for hypoplastic ventricular

disease.

Keywords: Hypoplastic left heart syndrome, soluble guanylyl cyclase, nitric oxide, cardiac

development, zebrafish.

50   **Graphical Abstract**

1. Introduction

Hypoplastic left heart syndrome (HLHS) is a complex congenital heart disease (CHD) with a multigenic etiology.[1,2] Familial clustering supports a genetic basis, yet the majority of causative genes remain unidentified. The phenotype is heterogeneous, variably affecting the left ventricle, valves, and aortic outflow tract, and no single-gene mutation has been shown to reproduce the full HLHS spectrum in animal models. Traditionally, HLHS was attributed to diminished intracardiac flow during development, leading to secondary hypoplasia. However, recent work has shifted this paradigm, suggesting that HLHS may arise from primary defects in myocardial and endocardial development, resembling a developmental cardiomyopathy in origin rather than a purely hemodynamic lesion.[3] Additionally, an emerging body of evidence links HLHS to disturbances of left–right (LR) patterning and cilia function, suggesting a new developmental mechanism contributing to disease pathogenesis.[4] Based on case reports and large registry-based analyses.[4,5] HLHS has been observed in association with heterotaxy, situs inversus, gut malrotation, biliary atresia and splenic anomalies.[6-8] These observations further support the concept that defects in laterality determination and ventricular morphogenesis may share developmental origins. According to recent interpretations, HLHS may originate from early intrinsic disruptions in cardiac progenitors arising during and prior to heart tube formation, and may occur in parallel with, or as a consequence of, abnormal left-right signaling.[9] Findings from recessive forward genetic screens in fetal mice further support this view, revealing that cilia and cilia-transduced signaling pathways play central roles in HLHS pathogenesis.[10]

Integrative genetic analysis combining human sequencing with developmental datasets and mouse models of HLHS identified CYGB as a candidate gene within a developmental gene network associated with the disease,[11] suggesting a potential role for CYGB in pathways regulating early cardiac morphogenesis. CYGB is a hexa-coordinate heme-containing globin expressed ubiquitously in vertebrate tissues.[12,13] Unlike its penta-coordinate relatives, hemoglobin and myoglobin, which primarily

bind and transport oxygen, CYGB is more prone to function as a redox-active enzyme regulating NO metabolism through reversible redox cycling of its heme iron.[14] The reduced heme iron can catalyze nitrite reduction to bioactive NO,[13,15] while the oxygenated heme iron can dioxygenate NO to inert nitrate,[16-18] thus maintaining oxygen-dependent NO homeostasis. While these reactions are possible in vitro, mouse studies indicate that the predominant physiological role of CYGB is to scavenge NO, modulating vascular tone through NO deoxygenation[19] and to protect cells against oxidative stress via superoxide dismutase-like activity.[20] Unexpectedly, we discovered a new critical role for Cygb in development, where CYGB/Cygb instead enhances NO signaling through activation of the nitric oxide synthase- sGC-cGMP signaling pathway.[21] The zebrafish *cygb2* knockout exhibits cardiac laterality defects due to impaired NO signaling, a phenotype characteristic of human disorders associated with defective cilia function. Mechanistically we demonstrated that Cygb2 localizes to cilia within the Kupffer's vesicle (KV), the zebrafish laterality organ. Loss of *cygb2* disrupts ciliary structure, motility and directional fluid flow, resulting in impaired KV activity and cardiac laterality defects.[21] Whether this function is directly linked to cardiac morphogenesis mimicking HLHS remains unknown and the extent to which Cygb dependent regulation of laterality intersects, potentially through ciliary mechanisms, has not been explored.

Zebrafish cardiac development initiates within the ALPM, where bilateral populations of mesodermal progenitors undergo progressive spatial organization prior to overt myocardial differentiation.[22,23] Between the 14–16 somite stages (ss), these progenitors undergo tightly coordinated mediolateral convergence and anterior–posterior elongation to assemble a coherent cardiac disc that subsequently transforms into the linear heart tube.[23,24] Ventricular formation is exquisitely sensitive to perturbations in progenitor number and spatial positioning as even subtle defects in coordinated cell migration or tissue architecture can give rise to small, compact ventricles, whereas atrial development may remain relatively preserved due to reliance on progenitor contributions from populations residing

in spatially segregated domains regulated by alternative signaling ques .[24,25] Importantly,

morphogenetic events critical to ventricular development occur immediately downstream of KV

mediated left–right axis establishment,[23,26] raising the possibility that laterality patterning cues

coordinating embryonic asymmetry also influence spatial patterning and migratory behavior of cardiac

progenitors, shaping ventricular growth and chamber morphogenesis. One of the earliest markers of

cardiac progenitors in zebrafish is *draculin* (*drl*), a transcription factor activated during gastrulation

that delineates bilateral cardiac progenitor fields within the ALPM.[23,27] During somitogenesis tracking

drl expression marks cardiac progenitor organization and collective migratory behavior. Within the

ALPM alterations in spatial geometry such as anterior–posterior elongation of the *drl* domain reflects

early disruption of progenitor spatial patterning,[23] defects expected to impair migratory ques and

ventricle assembly. Thus, alterations in *drl* expression may represent one of the earliest

developmental events that ultimately lead to hypoplastic ventricular growth. As development

proceeds, myocardial *drl* expression is restricted to first heart field descendants, providing a readout

of cardiac disc formation and heart tube assembly, stages that directly precede ventricular chamber

growth and functional maturation.[22]

NKX2 homeobox 5 (*Nkx2.5*) encodes a conserved cardiac homeobox transcription factor expressed

within the forming cardiac disc at the transition from migratory cardiac progenitors to differentiated

myocardium.[23,27] In both mouse models and human patients, dysregulation of NKX2.5 activity is

associated with congenital heart diseases and hypoplastic ventricular phenotypes[28]. In the zebrafish

embryo, *nkx2.5* expression is conserved and is initially induced in bilateral ALPM domains as cardiac

progenitors converge toward the midline, coinciding with progenitor stabilization and the initiation of

chamber-specific transcriptional programs.[29] By 28 hpf, *nkx2.5* expression marks specified

cardiomyocytes within the heart tube, predominantly ventricular myocardium,[24] such that its expression

at this stage reflects proper myocardial differentiation and maintenance of cardiac identity.

While Cygb-dependent regulation of embryonic laterality is now established,[21] its role in coordinating cardiac progenitor migration and myocardial differentiation remains unknown. Here, we demonstrate that *cygb2* deficiency in zebrafish disrupts early cardiac progenitor organization and ventricular chamber formation, producing a small, compact ventricle with reduced diastolic filling. These features parallel ventricular hypoplasia and impaired compliance observed in HLHS. Crucially, these defects are phenocopied in sGC mutants and fully rescued by pharmacological activation of NO-sGC signaling, identifying a developmental pathway that links ciliary NO regulation to cardiac progenitor morphogenesis and ventricular growth.

2. Methods

All animal studies were approved by the University of Maryland Institutional Animal Care and Use Committee (IACUC-1022015), Baltimore, MD, USA. The programs and facilities are USDA registered and covered under an assurance with the Office of Lab Animal Welfare (OLAW). Procedures followed guidelines outlined by the NIH Guide for the Care and Use of Laboratory Animals (8th Edition, 2011). Embryos were maintained at 28.5 °C until the desired developmental stage. Animals were euthanized according to approved protocols; embryos >3 days post fertilization (dpf) to adult (anesthetic overdose 300mg/L buffered tricane), embryos <3dpf (paraformaldehyde immersion).

2.1 Animal Studies

For all experiments, clutch-matched embryos from dome stage (~ 4hpf) to 5dpf were randomly assigned to treatment groups; sex cannot be determined at these stages. The AB (WT) genetic background was used for generation of mutant and transgenic lines. Transgenic lines in this study include Tg(*myl7:EGFP*) and Tg(*tbx5a:EGFP*;*tcf21:dsRED*). The *cygb2^801a* and the *gucy1a1* mutants were generated using CRISPR-Cas9 genome editing as previously described[21] and guide RNAs are listed in[21] and *Table S1*. Embryos were raised and screened for restriction enzyme site disruption (N1aIII)

using primers specific to exon 4 of *gucy1a1* (*Table S1*). The Tg(*myl7:EGFP*) and Tg(*tbx5a:EGFP*);Tg(*tcf21:dsRED*) lines were crossed with *cygb2[801a]* to produce *cygb2[801a]*;Tg(myl7:EGFP), *cygb2[801a]*;Tg(*tbx5a:EGFP*);Tg(*tcf21:dsRED*), and wildtype sibling lines.

2.2 Stroke Volume

Stroke volumes were calculated from live imaging. To calculate end-diastolic volume (EDV) and end-systolic volume (ESV) frames were identified from high-speed wide-field (bright-field and fluorescent) time-lapse videos of beating zebrafish hearts at the moments of maximal and minimal ventricular cavity size. The long-axis length (*L*) and short-axis diameter (*D*) of the ventricle were measured from frames using the straight line tool in FIJI imaging software. EDV and ESV were calculated using the established prolate spheroid model[30] according to the formula:

$$V = \frac{\pi}{6}LD^2$$

L represents ventricular long-axis length and D the diameter at mid-ventricle (*Figure S1*). Stroke volume (SV) was derived as SV = EDV − ESV. Ventricle wall thickness was calculated at maximal and minimal cavity size, identified from high-speed differential interference contrast (DIC) microscopy videos of beating zebrafish hearts. The measurable area and length of the ventral-facing ventricle wall was calculated using FIJI imaging software and the polygon selections and segmented line tool. Area was divided by the length of the measured wall to generate a ratio-based metric of thickness. Bright field and fluorescent images were captured using a Zeiss Axio Observer inverted microscope with attached Axiocam 807 mono camera and a Leica M165FC with attached K3C camera.

2.3 Ventricle Volume and morphology

Whole mount confocal z-series and compiled z-projections were collected on paraformaldehyde fixed zebrafish embryos from 16ss to 3 days post fertilization. Immuno-labeled embryos were mounted on glass slides and imaged on a Nikon Ti2 inverted W1 spinning disk confocal microscope with attached

Hamamatsu sCMOS camera. Collected z-series were compiled into max projections and 3D objects with surface renderings using FIJI and Imaris imaging software. Ventricle volume and wall thickness were calculated using thresholding on selected regions of interest (ROI). Z-slices covering full depth and width of ventricle were complied, images were converted to 8-bit and set to grey scale. To calculate volume of myocardium, thresholding was set to black and white, area measurements for signal-positive region were measured for each z-slice, summed, and multiplied by the distance between z-slices. Ventricular lumen volume was calculated by measuring the area of signal-negative region within the surrounding signal-positive region and summing the areas across z-slices and multiplying by the distance between z-slices. To create a ratio-based metric to measure thickness across a 3-dimensional structure, myocardium volume was divided by lumen volume. Cell density was calculated by dividing total cardiomyocyte number by myocardium volume. Total ventricle cardiomyocyte number was calculated using collected z-series and FIJI imaging software. Z-slices spanning the full depth and width of the ventricle were compiled, and analyzed frame by frame, and EGFP-positive cardiomyocytes were marked, tracked, and counted. Ventricle volume at 5 dpf was generated from z-series collected using a Zeiss LS7 Lighsheet confocal microscope. Ventricle area of z-slices was calculated by thresholding (Huang Over Under) of the EGFP positive region within a ROI defining the ventricle including the luminal space. Using FIJI and open-source macro code ventricular volume was calculated by multiplying the summed areas by the distance between z-slices. Progenitor cell fusion, cardiac disc, and heart tube areas were calculated using thresholding of EGFP and CY5 positive areas in selected ROIs of max z-projections collected using spinning disk confocal microscopy.

2.4 In situ hybridization

Whole-mount in situ hybridization was performed on paraformaldehyde fixed zebrafish embryos as previously described[21] at 10-12ss and 28 hpf using riboprobes targeting *drl*, *kinase insert domain receptor like* (*kdrl*), and *nkx2.5*. Transcripts were amplified by PCR and cDNA was obtained from pooled

embryos using transcript specific primers (*Table S1*). Images were collected using bright-field microscopy on a Zeiss Discovery V.8 SteReo stereoscope with attached Axiocam 503 color camera, brightness and contrast were unaltered, and images were formatted using FIJI imaging software. To determine expression patterning variance, raw counts of phenotype rankings were used for statistical comparisons with the Fishers exact test, percentages were plotted for graphical representation, ns - not significant ≥ 0.05, * P≤ 0.05, ** P≤ 0.01, *** P≤ 0.001, **** P≤ 0.0001.

Drug Exposure

Carboxy-PTIO potassium salt (2-4-carboxyphenyl-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide; cPTIO, Sigma–C221) and BAY 582667 (Cinaciguat, Bay58, Sigma–SML1532) were diluted in DMSO and used at final concentrations of 250µM and 120µM. (Z)-1-(N,N-diethylamino)diazen-1-ium-1,2-diolate (DETA NONOate) (Cayman-82100) and (Z)-1-[N-(2-aminoethyl)-N-(2-ammonioethyl)amino]diazen-1-ium-1,2-diolate (PAPA NONOate) (Cayman-2140) were resuspended in sodium hydroxide and diluted in phosphate buffered saline at a final concentration of 500µM and 5.73µM. Drug route of exposure was administered through incubation in embryo media (E3).

Immunohistochemistry

Whole-mount immunohistochemistry was performed as previously described[21] using primary antibodies for myosin heavy chain 6 (Developmental Studies Hybridoma Bank S46). Embryos were mounted on glass slides with Vectashield Hardset Antifade Mounting Medium (H-1500-10) cover slipped and imaged on a Nikon Ti2 inverted W1 spinning disk confocal microscope with attached Hamamatsu sCMOS camera.

cGMP ELISA

cGMP levels in zebrafish embryos were measured by competitive ELISA following manufactures protocol (Enzo-ADI-900-164). Embryos at 28 hpf were incubated in 250µM DETA NONOate for 30 minutes at 28.5 °C. Fifty embryos per sample were pooled and homogenized in 0.1M hydrochloric acid on ice. Lysates were centrifuged and the supernatant was collected for analysis. Samples were assayed in triplicate at two dilutions. cGMP concentration was calculated based on interpolation to the standard curve using a 95% confidence interval.

Cell culture

The human type II alveolar epithelial cell line A549 was kindly provided by Dr. Alan Cross (Professor, Department of Microbiology and Immunology, University of Maryland School of Medicine). A549 cells were cultured in RPMI-1640 media (Gibco-11875093) containing Penicillin/Streptomycin (Sigma-P4333) and fungizone (Amphoterecin B, Gibco-15290-026), and 10% fetal bovine serum (FBS) (Sigma-F4135) at 37^0 Celsius and 5% carbon dioxide. Cell cultures were passaged a minimum of three times before performing experiments.

Small interfering RNA

Dicer Substrate RNA duplexed small interfering RNAs targeting human cytoglobin were designed using Integrated DNA Technologies "Predesigned DsiRNA Selection Tool". DsiRNA-mediated gene silencing was performed by transfection of sub-confluent A549 monolayers using Lipofectamine RNAiMAX transfection reagent (Thermo Fisher-13778030) and non-targeting DsiRNA and targeting si-CYGB 13.1 for 48 hours. To assess cell viability, transfected cells were pelleted and resuspended in a 1:1 mixture of cell suspension and 0.4% Trypan Blue. Cells were counted using a hemocytometer in a 0.05µm section. Cells excluding trypan blue were counted as live and cells stained with trypan blue were counted as dead, viability was calculated as a percent of live cells to total cell number.

Western Blot

A549 cells were lysed on ice in RIPA buffer (Sigma-R0278) containing protease inhibitor (Halt 1X, Thermo Fisher-1860932). Lysates were centrifuged at 12,500 RPM for 20 min at 4°C. The BCA assay was used to determine total protein concentration. Samples were loaded for gel electrophoresis in 4-20% precast gels (Bio-Rad-4561094) using 100 ug total protein per sample. Following electrophoresis protein was transferred to polyvinylidene fluoride membranes. Membranes were blocked with 3% bovine serum albumin (BSA) for 2 hours at room temperature under gentle agitation. Membranes were probed with primary rabbit anti-CYGB (Sigma-HPA017757) diluted at 1:1000 in 1% BSA and 0.1% Triton X100 in phosphate buffer saline (PBST) overnight at 4^0 Celsius. Following 3 consecutive 5-minute washes at room temperature in PBST membranes were probed with anti-Rabbit IgG-HRP (Abcam-131366) at 1:10,000 in PBST with 1% BSA for 1 hour at room temperature. Following 3 consecutive 5-minute washes at room temperature in a solution of PBST protein bands were visualized using Clarity Max Western ECL substrate (Bio-Rad-1705060) following manufacturer's protocol and a Bio-Rad ChemiDoc MP imaging box. Following 3 consecutive 5-minute washes with PBST at room temperature membranes were stripped using Restore Western Blot Stripping Buffer (Thermo Scientific 21059). Blocking and probing procedure was repeated as before with primary anti-β-Actin antibody (Sigma-A2066). Band intensity was normalized to β-Actin and quantified using FIJI imaging software.

Wound healing assay

A549 cells were seeded in 24-well plates at a density of 170,000 cells per well. After 24 hours cells were transfected with 2.5pmol non-targeting DsiRNA, 1.25pmol and 2.5 pmol si-CYGB 13.1. 48 hours post transfection a scratch was introduced with a 10µl pipette tip. Following the scratch cells were washed with sterile 1x PBS and cultures were supplemented with RPMI-1640 media containing 10% (FBS). Cultures were imaged at 0 and 24 hours post scratch on a Leica DM IL LED inverted microscope using a 4X objective. Wound healing was assessed as precent migration into scratch area relative to

non-targeting DsiRNA. Image analysis was conducted using FIJI imaging software and the straight line

tool selected at lateral median.

2.5 Statistical analysis

Unless otherwise noted data are presented as mean ± SD, and statistical significance was determined

using an unpaired two-tailed *t*-test (or Mann–Whitney test for non-normal distributions), ns - not

significant ≥ 0.05, * P≤ 0.05, ** P≤ 0.01, *** P≤ 0.001, **** P≤ 0.0001.

3. Results

Cytoglobin is required for ventricular development and cardiac function

We previously defined a role for Cygb2 in cardiac laterality via regulation of cilia function. Here, we

investigated whether Cygb2 is required for cardiac morphogenesis and performance and whether these

are linked to laterality establishment. To visualize cardiac morphology we generated a *cygb2* mutant

line using the previously characterized *cygb2[801a]* mutation[21] expressed in an established transgenic

reporter background expressing EGFP under the control of the *myosin light chain 7* (*myl7)* promoter.[23]

In zebrafish, *myl7* expression begins at approximately the 16ss stage in cardiomyocyte progenitors

regardless of eventual chamber contribution to the atrium or ventricle.[23] We analyzed *myl7:EGFP*

expression at 3 days post fertilization (dpf), when cardiac morphology is established, and the ventricle

and atrium exhibit coordinated contraction and robust cardiac function. We first quantified the area of

both cardiac chambers and observed a significant decrease in the ventricle of *cygb2[801a]* embryos

compared to wildtype controls, (*Figure 1A–C, Movies S1-2*), while atrial area remained unchanged

(*Figure 1D*). To precisely assess performance, we measured EDV and ESV from in vivo high-speed

recordings and calculated SV using the prolate spheroid method *(Figure S1)*. Ventricles of *cygb2[801a]*

embryos showed a marked decrease in SV and EDV with no change in ESV (*Figure 1E–H*), consistent

with reduced diastolic filling rather than systolic impairment. As *cygb2[801a]* embryos exhibited cardiac

laterality defects, we tested whether the ventricular defect might arise secondarily from laterality disturbances. We compared SV between embryos with normal (situs solitus) and reversed (situs inversus) cardiac orientation within each genotype and detected no differences, $cygb2^{801a}$ embryos displayed reduced SV relative to wildtype independent of cardiac left-right orientation (*Figure S2A–F*). SV at 5dpf was assessed in the *myl7:EGFP* background and revealed that the phenotype persisted (*Figure S3A–D, Movies S3-4*). Additionally, ventricle volume was assessed using light sheet confocal microscopy on fixed *myl7:EGFP* samples and further supported the reduced ventricle phenotype and persistence at 5 dpf in $cygb2^{801a}$ compared to wildtype (*Figure S3E*). Collectively, these findings indicate a stable structural hypoplasia rather than a transient developmental delay. Despite reduced chamber size, ventricular cardiomyocyte number was unchanged (*Figure 1 I-J*). Instead, cells occupied a smaller volume, yielding a more compact myocardium and a thicker ventricular wall (*Figure 1K–M*). These data indicate that *cygb2* deficiency primarily affects ventricular morphogenesis and chamber remodeling, not cardiomyocyte proliferation or survival. The reduced, thick-walled ventricle observed in $cygb2^{801a}$ embryos accompanied by reduced SV mirrors key structural features of HLHS and, given the established role of Cygb2 in cilia-related laterality signaling, *cygb2* deficiency may represent a mechanistically relevant model of HLHS-like ventricular hypoplasia.

Nitric oxide restores ventricular size and function in *cygb2* mutants.

We previously showed that Cygb2 regulates cardiac laterality determination[21] through NO signaling. To test whether the reduced SV in $cygb2^{801a}$ embryos depends on NO, we first confirmed the acute physiological action of NO donors. Ten-minute exposure at 78 hpf to the pharmacological NO donor DETA-NONOate (DETA) increased SV in both wildtype and $cygb2^{801a}$ embryos *(Figure S4A-B),* consistent with the reported hemodynamic effect of NO in teleost fish larvae[31] where the increase in SV results from reduced afterload due to relaxation of the outflow vasculature. A fast-releasing NO donor, PAPA-NONOate, produced a similar acute increase, validating the assay (*Figure S4C-D*). To

determine whether the altered ventricular structure and morphology could be rescued by NO supplementation, we tested the effect of chronic DETA treatment. DETA was administered in a vehicle (sodium hydroxide), which alone did not affect SV (*Figure S4E*). We performed chronic DETA treatments from the 2ss stage to 3 dpf. Chronic exposure produced opposite outcomes depending on genotype, in *cygb2^801a* embryos ventricular size increased with consequent rescue in SV, whereas in wildtype embryos ventricular size was reduced, resulting in decreased SV (*Figure 2A-B*). Because prolonged NO exposure affects cardiac morphogenesis rather than acute mechanics, these bidirectional responses under identical dosing indicate that *cygb2* mutants possess a specific deficit in NO-dependent developmental signaling, while excess NO in wildtype embryos disrupts the finely tuned program governing ventricle morphogenesis. To provide further evidence supporting NO dependence, we treated embryos at 2ss with the NO scavenger cPTIO to phenocopy the cardiac phenotype observed in mutants (*Figure 2A,C*). As expected, ventricular volume was reduced by NO scavenging in wildtype embryos and was further exacerbated in *cygb2^801a* embryos, consistent with a requirement for NO signaling in normal ventricular development. A time course initiating DETA at stages from dome to 12ss showed maximal rescue when treatment began at dome stage to 2ss, diminishing by 12ss (*Figure 2D*), suggesting Cygb2-dependent NO signaling becomes critical for cardiac specification and morphogenesis during early somitogenesis. Together, these findings demonstrate that *cygb2* is required to maintain appropriate NO levels during early cardiac progenitor organization, thereby ensuring proper ventricular morphogenesis and SV.

Loss of Cygb2 disrupts early cardiomyocyte migration.

Given that the ventricular phenotype observed in *cygb2^801a* embryos originates during early development, we next examined cardiac morphogenesis to determine when these defects first emerge. In zebrafish, cardiac progenitors originate bilaterally within the ALPM and progressively migrate toward the midline, where they fuse to form the cardiac cone and subsequently the primitive heart tube. To

assess whether this process was altered in *cygb2* mutants, we analyzed the distribution of *myl7*-positive

cardiomyocyte precursors during heart field assembly. At 16ss when cardiomyocyte progenitors have

initiated migration but have not yet fused, *cygb2^801a* embryos displayed a significantly broader *myl7*-

positive domain compared with wildtype, indicating an expanded or delayed coalescence of cardiac

progenitors at this stage (*Figure 3A-C*). By 20ss, when the cardiac disc becomes morphologically visible

at the midline, the *myl7*-positive region remained wider and less compacted in *cygb2^801a* embryos,

occupying a larger total area relative to controls (*Figure 3D-F*). These results suggest that Cygb2 is

required for the coordinated medial convergence of cardiomyocyte progenitors during heart field

assembly. As development progressed, the cardiac tube of *cygb2^801a* embryos remained visibly smaller

than in wildtype. At 34 and 48 hpf, the ventricular area was significantly reduced indicating that the

morphogenetic defect persists through the stages of heart tube elongation and chamber ballooning,

whereas the atrium appeared morphologically normal throughout (*Figure S5A-H).* This suggests a

ventricle-specific effect consistent with the smaller, compact ventricular morphology observed at later

stages. To establish NO as a critical regulator throughout ventricular morphogenesis, embryos were

treated with the NO donor DETA beginning at 2ss. These treatments restored *myl7*-positive cardiac

field at 16 and 20ss (*Figure 3A-F)* and rescued ventricular area at 34 and 48 hpf (*Figure S5A-H*) to near

wildtype levels. These findings demonstrate that *cygb2* is essential for the proper migration and

coalescence of cardiac progenitors and for subsequent expansion of the ventricular chamber, and that

both processes depend on NO-mediated signaling.

Upon observing aberrant organization of cardiac progenitors during midline convergence and cardiac

disc formation in *cygb2^801a* embryos, we directly tested whether cell migratory behavior was disrupted

using an in vitro wound-healing assay in the human type II alveolar epithelial A549 cell line. *CYGB*

knockdown by small interfering RNA caused a dose-dependent reduction in cell migration into the

wound area (*Figure 3G-H*). CYGB expression knockdown efficiency and cell viability were confirmed

by Western Blot analysis (*Figure3I,J*) and viability assay (*Figure S6B*). These data suggest that

abnormal cardiac progenitor contribution to early heart structures may result from disruption of cellular

machinery regulating cell migration and/or cell adhesion and further indicate that Cygb-NO signaling

influences migratory behavior across cell types and species.

Soluble guanylate cyclase is required for ventricle development and heart function.

NO can signal through multiple mechanisms, including protein S-nitrosylation, nitration reactions and

formation of reactive nitrogen species. However, our data indicate that NO primarily signals through its

canonical receptor sGC. Upon ligand binding, NO-sGC catalyzes the conversion of guanosine

triphosphate (GTP) to cGMP, thereby initiating secondary downstream signaling events[32]. To test

whether the cardiac phenotype observed in *cygb2^{801a}* embryos depends on sGC, we generated an sGC

mutant (*gucy1a1^{9bp}*) using CRISPR/Cas9-mediated site-directed mutagenesis. Genomic DNA from

embryos derived from a maternal zygotic mutant line revealed a 9 bp deletion in exon 4 of the sGCα1

subunit, which was confirmed at the genomic and mRNA level by Sanger Sequencing (*Figure S7A*).

Loss of the normal transcript was validated by in situ hybridization using a 300 bp probe spanning the

edited region, which showed markedly reduced hybridization signal in *gucy1a1^{9bp}* embryos compared

to wildtype whereas a 1000 bp probe located downstream of the edited site produced comparable

staining in both mutants and wild type (*Figure S7B*). Functional loss of sGC activity in *gucy1a1^{9bp}*

embryos was confirmed by the absence of SV rescue following treatment with DETA and a

pharmacological sGC activator BAY 582667 (Bay58) (*Figure S7C-D*). Additionally, failure of DETA to

restore cGMP levels in *gucy1a1^{9bp}* compared to wildtype controls further validated the loss of function

(*Figure S7E*). In contrast, DETA treatment restored cGMP levels in *cygb2^{801a}* embryos,[21] supporting

the interpretation that Cygb2 functions upstream of sGC to regulate NO-dependent cGMP production

during cardiac development. To determine whether sGC disruption phenocopies the *cygb2^{801a}* cardiac

defect, we quantified ventricular performance in *gucy1a1^{9bp}* embryos using DIC microscopy and high-

speed imaging. SV was significantly reduced in *gucy1a1^{9bp}* compared with wildtype embryos, driven by

a decrease in EDV while ESV remained unchanged (*Figure 4A-E, Movies S5,6*). This pattern mirrors the functional phenotype of *cygb2[801a]* embryos, indicating that loss of sGCα1 function recapitulates the diastolic filling defect observed in the *cygb2* mutant. Furthermore, replicating experiments done in *cygb2* mutants we tested SV at 5dpf in *gucy1a1* mutants and found that in both reduced SV persisted at later developmental stages (*Figure S3F, Movies S7,8*). To confirm that the observed phenotype reflects disruption of a shared Cygb2-NO-sGC signaling pathway in both *cygb2[801a] and gucy1a1[9bp]* mutants, we tested whether pharmacological activation of sGC in *cygb[801a]* mutants could rescue cardiac function. Treatment of *cygb[801a]* mutants with the sGC activator Bay58 from 2ss to 34hpf fully restored SV (*Figure 4F-H),* confirming a critical role for sGC signaling in ventricular formation. Notably, neither DETA nor Bay58 rescued stroke volume in *gucy1a1[9bp]* mutants, as expected (*Figure S7C-D*). To assess a shared phenotype of increased ventricle wall thickness, measurements were performed on live embryos at 3dpf. Wall thickness at diastole and systole was increased in *gucy1a1[9bp]* (*Figure 4I-K, Movies S5,6*), similar to that observed in *cygb2[801a]* mutants (*Figure 4L-N, Movies S9,10*), indicating a conserved phenotype across mutant models and analytic approaches. To determine whether NO-sGC signaling contributes to this phenotype, *cygb2[801a]* mutants were treated from 2ss to 34hpf with DETA and Bay58. These conditions fully restored ventricular wall thickness (*Figure 4L-N),* supporting a role for Cygb-NO-sGC in ventricular morphogenesis.

NO-sGC signaling regulates early cardiac progenitor organization.

To determine whether Cygb2 and Gucy1a1 act through a shared molecular pathway during early cardiogenesis, we analyzed key markers of the ALPM at 10–12ss, when early cardiac and endothelial progenitors are established. In wildtype embryos, *drl* expression appeared as bilateral short, symmetrical stripes flanking the midline, marking anterior mesoderm that gives rise to cardiac and vascular lineages. In both *cygb2[801a]* and *gucy1a1[9bp]* embryos, *drl* domains were elongated, thinner, and extended posteriorly, indicating disrupted convergence and defective compaction of the ALPM (*Figure*

5A). Similarly, *kdrl*, which labels endothelial and endocardial progenitors, was expressed bilaterally in compact domains close to the midline in wildtype embryos but it was elongated, more widely separated, and extended posteriorly in *cygb2[801a]* and *gucy1a1[9bp]* mutants (*Figure S8*). This finding is consistent with delayed or uncoordinated migration of endothelial/endocardial precursors. Despite these spatial changes, signal intensity was not diminished, indicating that endothelial specification occurred normally, but progenitor patterning shifted from anterior (endothelial/endocardial) toward more posterior (hematopoietic) mesodermal domains. Expression of the early cardiac transcription factor nkx2.5 was also altered at 10-12ss. In wildtype embryos, *nkx2.5* was expressed as two distinct, symmetrical domains within the ALPM, whereas in *cygb2[801a]* and *gucy1a1[9bp]* mutants expression was frequently asymmetric or spatially disorganized (*Figure 5B*), suggesting loss of coordination between left and right cardiac fields. Importantly, wildtype siblings generated during establishment of the maternal-zygotic *cygb2[801a]* and *gucy1a1[9bp]* lines were outcrossed to minimize potential CRISPR off-target effects and displayed normal *drl*, *kdrl* and *nkx2.5* expression patterns at all developmental stages examined (*Figure S9*). The *drl* expression patterning abnormalities were rescued in *cygb2[801a]* embryos treated with either DETA or Bay58 (Figure 5A), confirming that early ALPM organization and synchronized cardiac specification depend on Cygb2-mediated NO–sGC signaling. Additionally, *nkx2.5* expression was also partially normalized by DETA treatment, although the strongest normalization was achieved in embryos treated with Bay58 (Figure 5B), confirming the importance of sGC signaling in this process. As *drl* marks early cardiac precursors fields and *nkx2.5* labels a broader population encompassing later myocardial progenitors, this differential rescue further supports the conclusion that NO–sGC signaling exerts its strongest influence during the earliest stages of cardiac precursor organization and initial heart structure formation.

NO–sGC signaling rescues heart tube development.

To assess whether early ALPM patterning defects persist during later stages of cardiogenesis, we examined *drl*, and *nkx2.5* expression at 28 hpf, when the linear heart tube has formed and endothelial and myocardial differentiation are underway. In wildtype embryos, *drl* expression is confined to the developing vasculature and the endocardium; in contrast, *cygb2*[801a] and *gucy1a1*[9bp] embryos exhibited reduced or absent *drl* expression in the heart tube (*Figure 6A*). This pattern is consistent with the shift observed at earlier stages from anterior (endothelial/endocardial) toward more posterior (hematopoietic) mesodermal domains. Consistently, *nkx2.5* expression in the ventricle was markedly reduced in both *cygb2*[801a] and *gucy1a1*[9bp] embryos, indicating impaired myocardial contribution to this region (*Figure 6B*). Treatment of *cygb2*[801a] embryos with DETA or Bay58 restored *drl* and *nkx2.5* expression patterns to near wildtype levels (*Figure 6A-B*). Together, these findings show that *cygb2* and *gucy1a1* mutants share a common set of early and late ALPM abnormalities that can be corrected by restoring NO-sGC signaling, linking early ALPM mis-patterning to the later emergence of a compact, hypoplastic ventricle.

4. Discussion

Nitric oxide (NO) signaling is a central regulator of cardiovascular physiology, with established roles in vascular tone, myocardial contractility, and hemodynamic homeostasis.[33-35] In contrast, the contribution of NO signaling to early cardiac morphogenesis has been far less explored. Here, we show that Cygb2-dependent NO–sGC signaling is required for ventricular formation during zebrafish heart development. Loss of *cygb2* leads to a reduced SV associated with decreased ventricle volume, increased myocardial wall thickness, elevated cellular density, and impaired diastolic filling. These defects arise from abnormal migratory behavior of cardiac precursors, which display reduced migratory efficiency, excessive cell-spreading, and subsequent hypercompaction within the forming heart tube. Importantly, genetic loss of sGC (*gucy1a1*) fully phenocopies the *cygb2* mutant phenotype, and pharmacological

activation of NO-sGC signaling in *cygb2* mutants rescues ventricular volume, cardiac progenitor

migration, myocardial organization within the forming heart tube, and diastolic function. Together, these

data identify NO–sGC signaling downstream of Cygb2 as a central pathway coordinating myocardial

progenitor dynamics, ventricular morphogenesis, and functional maturation.

Cardiac progenitor migration is a fundamental driver of heart tube assembly and ventricular

morphogenesis and depends on tightly regulated cell polarity, adhesion turnover, and cytoskeletal

dynamics.[36-39] Disruption of these processes can profoundly alter chamber size and myocardial

architecture.[40,41] In this context, our data indicate that reduced NO–sGC signaling places cardiac

progenitors into a mechanically altered migratory state. NO-sGC-cGMP signaling is known to regulate

actomyosin contractility, focal adhesion turnover, and cortical tension across multiple cellular

contexts[42-44] and cytoglobin-dependent control of NO bioavailability represents a key upstream

determinant of this pathway[13,14]. Attenuation of NO–sGC signaling during development is therefore

predicted to stabilize cell–substrate and cell–cell adhesions, resulting in excessive cell spreading,

reduced migratory efficiency, and impaired tissue-level cellular rearrangements. When such

mechanically altered cells enter the confined environment of the forming heart tube, increased adhesion

and cortical tension can bias morphogenesis toward tissue compaction and elevated cellular packing

as demonstrated in epithelial and mesenchymal morphogenetic models.[45,46] These properties provide

a mechanistic basis for the increased ventricular cell density observed in *cygb2* mutants, together with

the resulting thick-walled ventricle and impaired diastolic function.

We previously demonstrated that Cygb2 localizes to motile cilia and regulates cilia-dependent left–right

patterning through NO–sGC signaling during early zebrafish development,[21] establishing that Cygb2-

dependent NO–sGC signaling operates during early developmental windows critical for laterality

specification. The present study extends this role to ventricular formation while showing that ventricular

defects do not arise directly from altered left–right axis specification (*Figure S2*). However, ventricular

morphogenesis may still depend on cilia-associated signaling while remaining independent of final

laterality outcomes. Here, we found that pharmacological restoration of NO–sGC signaling rescues ventricular morphology only when administered during early windows encompassing laterality determination and early somitogenesis, whereas later treatment fails to restore ventricular size and stroke volume, indicating that ventricular morphogenesis is temporally linked to early cilia-associated signaling events. During this developmental window, we found that *drl* and *kdrl* expression is spatially displaced and expanded within the ALPM, phenotypes consistent with defective cardiac progenitor migration rather than primary loss of cardiogenic identity[36,38,39,41]. Notably, *nkx2.5*, normally expressed symmetrically by early somitogenesis[47] is found asymmetric in *cygb2* mutants, suggesting altered organization of subsets of cardiac progenitors. Recent single-cell studies indicate that *nkx2.5* marks transcriptionally distinct progenitor populations within the ALPM[29] supporting the idea that the disrupted signaling perturbs progenitor composition rather than uniformly impairing cardiac fate. Early migratory defects likely lead secondarily to impaired differentiation or maintenance programs later, as reflected by the reduction of *nkx2.5* and *drl* expression within ventricular territories at 28 hpf.[48,49] Importantly, dysregulation of NKX2-5 is associated with several forms of congenital heart disease in humans, including ventricular septal defects, tetralogy of Fallot, and HLHS, underscoring the relevance of these findings to human cardiac pathogenesis.[47,50,51]

Sequencing studies have identified enrichment of rare variants in cilia and ciliopathy-associated genes in subsets of HLHS patient cohorts,[4] supporting a framework in which cilia-dependent pathways contribute to disease susceptibility in a subset of patients. Within this framework, zebrafish models exhibiting reduced ventricular size with compact myocardium have emerged as informative systems for studying cardiac morphogenesis and HLHS-like phenotypes.[9,49] In our *cygb2* mutant model, HLHS-like ventricular hypoplasia originates from early events concomitant with cilia-dependent processes involved in left-right patterning. Our data indicate two possible mechanistic pathways underlying this phenotype. In one scenario, impaired NO-sGC signaling caused by loss of Cygb2 directly disrupts the coordination of cardiac progenitor migration, regional specification, and ventricle expansion, leading to

reduced ventricular growth. In a second, distinct scenario, loss of Cygb2 perturbs cilia-dependent processes that regulate early cardiac morphogenesis and myocardial growth, beyond their established role in cardiac laterality, through downstream effects on progenitor deployment and tissue remodeling,[50] independent of left–right axis determination. These mechanisms are not mutually exclusive and may interact in the multigenic context of human HLHS, where combined defects in NO signaling and cilia-associated pathways could converge on shared morphogenetic processes controlling ventricular development. The *cygb2 and gucy1a1* models characterized here converge on the hallmark ventricular phenotype of HLHS while revealing a distinct initiating axis centered on Cygb-dependent NO–sGC signaling. Together with prior work demonstrating that modulation of NO bioavailability influences myocardial plasticity and regeneration,[52] these findings establish NO–sGC signaling as a tractable developmental pathway whose disruption leads to ventricular hypoplasia and whose restoration is sufficient to rescue ventricular structure and function. Our data indicate that pharmacological activation of sGC, previously unexplored in the context of hypoplastic ventricular disease, can restore ventricular structure and function in HLHS-like phenotypes in vivo. These results position zebrafish as a valuable model for interrogating early developmental mechanisms underlying ventricular hypoplasia and for assessing sGC-targeted interventions in CHD.

Author Contributions

Adam Clark, Paola Corti designed and conducted experiments, analyzed data, and wrote the manuscript; Muddassar Iqbal and Rasmus Hejlesen reviewed the manuscript; Rasmus Hejlesen constructed a data set for transcript variance analysis; Weng Tzu-Ting conducted in situ hybridization experiments; Muddassar Iqbal designed, conducted; and analyzed data for wound healing assay experiments; Akueba Bruce conducted zebrafish husbandry and genotyping.

Acknowledgements

We are thankful to the University of Maryland School of Medicine Center for Innovative Biomedical Resources; Confocal Microscopy Core (Dr. Joseph Mauban and Dr. Shilpa Kumar) and Translational Genomics Lab, for invaluable assistance in Imaging and Sanger Sequencing.

Conflict of Interest: The authors declare no competing interests that influence the work reported in this paper.

Funding

This work was supported by National Institutes of Health under awards number 5T32AR7592-27 and NHL168775.

Data availability

The data used to produce this article will be shared at reasonable request to the corresponding author.

References

1. Iascone M, Ciccone R, Galletti L, Marchetti D, Seddio F, Lincesso AR, Pezzoli L, Vetro A, Barachetti D, Boni L, Federici D, Soto AM, Comas JV, Ferrazzi P, Zuffardi O. Identification of de novo mutations and rare variants in hypoplastic left heart syndrome. *Clin Genet* 2012;**81**:542–554.

2. Yagi H, Liu X, Gabriel GC, Wu Y, Peterson K, Murray SA, Aronow BJ, Martin LJ, Benson DW, Lo CW. The Genetic Landscape of Hypoplastic Left Heart Syndrome. *Pediatr Cardiol* 2018;**39**:1069–1081.

3. Grossfeld P, Nie S, Lin L, Wang L, Anderson RH. Hypoplastic Left Heart Syndrome: A New Paradigm for an Old Disease? *J Cardiovasc Dev Dis* 2019;**6**.

4. Gabriel GC, Lo CW. Left-right patterning in congenital heart disease beyond heterotaxy. *Am J Med Genet C Semin Med Genet* 2020;**184**:90–96.

5. Lin AE, Krikov S, Riehle-Colarusso T, Frias JL, Belmont J, Anderka M, Geva T, Getz KD, Botto LD, National Birth Defects Prevention S. Laterality defects in the national birth defects prevention study (1998-2007): birth prevalence and descriptive epidemiology. *Am J Med Genet A* 2014;**164A**:2581–2591.

6. Becker DJ, Islam S, Geiger JD. Biliary atresia associated with hypoplastic left heart syndrome: a case report and review of the literature. *J Pediatr Surg* 2004;**39**:1411–1413.

7. Stankiewicz P, Sen P, Bhatt SS, Storer M, Xia Z, Bejjani BA, Ou Z, Wiszniewska J, Driscoll DJ, Maisenbacher MK, Bolivar J, Bauer M, Zackai EH, McDonald-McGinn D, Nowaczyk MM, Murray M, Hustead V, Mascotti K, Schultz R, Hallam L, McRae D, Nicholson AG, Newbury R, Durham-O'Donnell J, Knight G, Kini U, Shaikh TH, Martin V, Tyreman M, Simonic I, Willatt L, Paterson J, Mehta S, Rajan D, Fitzgerald T, Gribble S, Prigmore E, Patel A, Shaffer LG, Carter NP, Cheung SW, Langston C, Shaw-Smith C. Genomic and genic deletions of the FOX gene

cluster on 16q24.1 and inactivating mutations of FOXF1 cause alveolar capillary dysplasia and other malformations. *Am J Hum Genet* 2009;**84**:780–791.

8. Huseynova RA, Bin Mahmoud LA, Abdelrahim A, Alroiedy MA, Huseynov O. Polysplenia syndrome with complex heart disease and jejunal atresia with malrotation in neonate: A case report. *Clin Case Rep* 2020;**8**:848–851.

9. Huang M, Akerberg AA, Zhang X, Yoon H, Joshi S, Hallinan C, Nguyen C, Pu WT, Haigis MC, Burns CG, Burns CE. Intrinsic myocardial defects underlie an Rbfox-deficient zebrafish model of hypoplastic left heart syndrome. *Nat Commun* 2022;**13**:5877.

10. Yagi H, Lo CW. Left-Sided Heart Defects and Laterality Disturbance in Hypoplastic Left Heart Syndrome. *J Cardiovasc Dev Dis* 2023;**10**.

11. Liu X, Yagi H, Saeed S, Bais AS, Gabriel GC, Chen Z, Peterson KA, Li Y, Schwartz MC, Reynolds WT, Saydmohammed M, Gibbs B, Wu Y, Devine W, Chatterjee B, Klena NT, Kostka D, de Mesy Bentley KL, Ganapathiraju MK, Dexheimer P, Leatherbury L, Khalifa O, Bhagat A, Zahid M, Pu W, Watkins S, Grossfeld P, Murray SA, Porter GA, Jr., Tsang M, Martin LJ, Benson DW, Aronow BJ, Lo CW. The complex genetics of hypoplastic left heart syndrome. *Nat Genet* 2017;**49**:1152–1159.

12. Trent JT, 3rd, Hargrove MS. A ubiquitously expressed human hexacoordinate hemoglobin. *J Biol Chem* 2002;**277**:19538–19545.

13. Corti P, Ieraci M, Tejero J. Characterization of zebrafish neuroglobin and cytoglobins 1 and 2: Zebrafish cytoglobins provide insights into the transition from six-coordinate to five-coordinate globins. *Nitric Oxide* 2016;**53**:22–34.

14. Amdahl MB, Sparacino-Watkins CE, Corti P, Gladwin MT, Tejero J. Efficient Reduction of Vertebrate Cytoglobins by the Cytochrome b(5)/Cytochrome b(5) Reductase/NADH System. *Biochemistry* 2017;**56**:3993–4004.

15. Reeder BJ, Ukeri J. Strong modulation of nitrite reductase activity of cytoglobin by disulfide bond

oxidation: Implications for nitric oxide homeostasis. *Nitric Oxide* 2018;**72**:16–23.

16. Halligan KE, Jourd'heuil FL, Jourd'heuil D. Cytoglobin is expressed in the vasculature and

regulates cell respiration and proliferation via nitric oxide dioxygenation. *J Biol Chem*

2009;**284**:8539–8547.

17. Gardner AM, Cook MR, Gardner PR. Nitric-oxide dioxygenase function of human cytoglobin with

cellular reductants and in rat hepatocytes. *J Biol Chem* 2010;**285**:23850–23857.

18. Tejero J, Shiva S, Gladwin MT. Sources of Vascular Nitric Oxide and Reactive Oxygen Species

and Their Regulation. *Physiol Rev* 2019;**99**:311–379.

19. Zweier JL, Ilangovan G. Regulation of Nitric Oxide Metabolism and Vascular Tone by Cytoglobin.

Antioxid Redox Signal 2020;**32**:1172–1187.

20. Zweier JL, Hemann C, Kundu T, Ewees MG, Khaleel SA, Samouilov A, Ilangovan G, El-Mahdy

MA. Cytoglobin has potent superoxide dismutase function. *Proc Natl Acad Sci U S A*

2021;**118**.

21. Rochon ER, Xue J, Mohammed MS, Smith C, Hay-Schmidt A, DeMartino AW, Clark A, Xu Q, Lo

CW, Tsang M, Tejero J, Gladwin MT, Corti P. Cytoglobin regulates NO-dependent cilia motility

and organ laterality during development. *Nat Commun* 2023;**14**:8333.

22. Yelon D. Cardiac patterning and morphogenesis in zebrafish. *Dev Dyn* 2001;**222**:552–563.

23. Kemmler CL, Riemslagh FW, Moran HR, Mosimann C. From Stripes to a Beating Heart: Early

Cardiac Development in Zebrafish. *J Cardiovasc Dev Dis* 2021;**8**.

24. Felker A, Prummel KD, Merks AM, Mickoleit M, Brombacher EC, Huisken J, Panakova D,

Mosimann C. Continuous addition of progenitors forms the cardiac ventricle in zebrafish. *Nat*

Commun 2018;**9**:2001.

25. Paffett-Lugassy N, Novikov N, Jeffrey S, Abrial M, Guner-Ataman B, Sakthivel S, Burns CE,

Burns CG. Unique developmental trajectories and genetic regulation of ventricular and outflow

tract progenitors in the zebrafish second heart field. *Development* 2017;**144**:4616–4624.

26. Hirokawa N, Tanaka Y, Okada Y. Left-right determination: involvement of molecular motor KIF3,

cilia, and nodal flow. *Cold Spring Harb Perspect Biol* 2009;**1**:a000802.

27. Mosimann C, Panakova D, Werdich AA, Musso G, Burger A, Lawson KL, Carr LA, Nevis KR,

Sabeh MK, Zhou Y, Davidson AJ, DiBiase A, Burns CE, Burns CG, MacRae CA, Zon LI.

Chamber identity programs drive early functional partitioning of the heart. *Nat Commun*

2015;**6**:8146.

28. Pashmforoush M, Lu JT, Chen H, Amand TS, Kondo R, Pradervand S, Evans SM, Clark B,

Feramisco JR, Giles W, Ho SY, Benson DW, Silberbach M, Shou W, Chien KR. Nkx2-5

pathways and congenital heart disease; loss of ventricular myocyte lineage specification leads

to progressive cardiomyopathy and complete heart block. *Cell* 2004;**117**:373–386.

29. Coskun H, Wang Y, Gao X, Chen K, Geoffrey Burns C, Burns CE. 3D spatial organization of

heterogeneous nkx2.5+ progenitors in the zebrafish heart field pre-patterns cardiovascular

development. *Nat Commun* 2025;**16**:11624.

30. Yalcin HC, Amindari A, Butcher JT, Althani A, Yacoub M. Heart function and hemodynamic

analysis for zebrafish embryos. *Dev Dyn* 2017;**246**:868–880.

31. Fritsche R, Schwerte T, Pelster B. Nitric oxide and vascular reactivity in developing zebrafish,

Danio rerio. *Am J Physiol Regul Integr Comp Physiol* 2000;**279**:R2200–2207.

32. Arnold WP, Mittal CK, Katsuki S, Murad F. Nitric oxide activates guanylate cyclase and increases

guanosine 3':5'-cyclic monophosphate levels in various tissue preparations. *Proc Natl Acad Sci*

U S A 1977;**74**:3203–3207.

33. Ignarro LJ. Nitric Oxide: A Unique Endogenous Signaling Molecule in Vascular Biology (Nobel

Lecture). *Angew Chem Int Ed Engl* 1999;**38**:1882–1892.

34. Gladwin MT. Deconstructing endothelial dysfunction: soluble guanylyl cyclase oxidation and the
NO resistance syndrome. *J Clin Invest* 2006;**116**:2330–2332.

35. Forstermann U, Sessa WC. Nitric oxide synthases: regulation and function. *Eur Heart J*
2012;**33**:829–837, 837a–837d.

36. Stainier DY. Zebrafish genetics and vertebrate heart formation. *Nat Rev Genet* 2001;**2**:39–48.

37. Buckingham SD, Ali DW. Computer simulations of high-pass filtering in zebrafish larval muscle
fibres. *J Exp Biol* 2005;**208**:3055–3063.

38. Abu-Issa R, Kirby ML. Heart field: from mesoderm to heart tube. *Annu Rev Cell Dev Biol*
2007;**23**:45–68.

39. Bakkers J. Zebrafish as a model to study cardiac development and human cardiac disease.
Cardiovasc Res 2011;**91**:279–288.

40. Hutson MR, Kirby ML. Model systems for the study of heart development and disease. Cardiac
neural crest and conotruncal malformations. *Semin Cell Dev Biol* 2007;**18**:101–110.

41. Vincent SD, Buckingham ME. How to make a heart: the origin and regulation of cardiac progenitor
cells. *Curr Top Dev Biol* 2010;**90**:1–41.

42. Surks HK, Mochizuki N, Kasai Y, Georgescu SP, Tang KM, Ito M, Lincoln TM, Mendelsohn ME.
Regulation of myosin phosphatase by a specific interaction with cGMP- dependent protein
kinase Ialpha. *Science* 1999;**286**:1583–1587.

43. Schlossmann J, Hofmann F. cGMP-dependent protein kinases in drug discovery. *Drug Discov*
Today 2005;**10**:627–634.

44. Prysyazhna O, Rudyk O, Eaton P. Single atom substitution in mouse protein kinase G eliminates
oxidant sensing to cause hypertension. *Nat Med* 2012;**18**:286–290.

45. Lecuit T, Lenne PF. Cell surface mechanics and the control of cell shape, tissue patterns and
morphogenesis. *Nat Rev Mol Cell Biol* 2007;**8**:633–644.

46. Heisenberg CP, Bellaiche Y. Forces in tissue morphogenesis and patterning. *Cell* 2013;**153**:948–

962.

47. Sultana N, Nag K, Hoshijima K, Laird DW, Kawakami A, Hirose S. Zebrafish early cardiac

connexin, Cx36.7/Ecx, regulates myofibril orientation and heart morphogenesis by establishing

Nkx2.5 expression. *Proc Natl Acad Sci U S A* 2008;**105**:4763–4768.

48. Moorman AF, Christoffels VM. Cardiac chamber formation: development, genes, and evolution.

Physiol Rev 2003;**83**:1223–1267.

49. Liu J, Stainier DY. Zebrafish in the study of early cardiac development. *Circ Res* 2012;**110**:870–

874.

50. Sutherland MJ, Ware SM. Disorders of left-right asymmetry: heterotaxy and situs inversus. *Am J*

Med Genet C Semin Med Genet 2009;**151C**:307–317.

51. George V, Colombo S, Targoff KL. An early requirement for nkx2.5 ensures the first and second

heart field ventricular identity and cardiac function into adulthood. *Dev Biol* 2015;**400**:10–22.

52. Rochon ER, Missinato MA, Xue J, Tejero J, Tsang M, Gladwin MT, Corti P. Nitrite Improves Heart

Regeneration in Zebrafish. *Antioxid Redox Signal* 2020;**32**:363–377.

726 **Figures**

751   **Figure 1**

752　**Figure 2**

762 **Figure 3**

771   1**Figure 5**

Figure 6

Figure Legends

Figure 1. Loss of *cygb2* leads to ventricular hypoplasia and reduced cardiac output. (A) Representative confocal projections of *cygb2wt*;Tg*(myl7*:EGFP) and *cygb2^{801a};*Tg*(myl7*:EGFP) hearts at 3 dpf, immuno-labeled for MYH6 (EGFP, green; MYH6, magenta). Scale bar, 50µm **(B)** Schematic illustrating heart chamber anatomy and cardiac orientation. A-atrium; V-ventricle; B-bulbus arteriosus **(C-D)** Comparison of ventricle and atrium area. **(E)** Representative frames from fluorescence time-lapse imaging of live *cygb2wt*;Tg*(myl7*:EGFP) and *cygb2^{801a};*Tg*(myl7*:EGFP) embryos 3 dpf showing ventricle volume at diastole and systole, dashed white line demarcates the ventricle (EGFP, green). Scale bar, 50 µm. **(F-H)** Comparison of stroke volume, diastolic volume and systolic volume. **(I)** Representative confocal projections of ventricles at 3 dpf, images are greyscale converted and inverted to highlight cardiomyocyte density. Scale bar, 20µm. **(J)** Comparison of ventricle cell number. **(K)** Representative confocal z-slices at ventricular midpoint. Images were converted to greyscale signal positive area (ventricular myocardium) shown in black. **(L)** Comparison of cardiomyocyte density. **(M)** Comparison of ventricular wall thickness. Student t-Test; *ns*, not significant; *p<0.05; **p<0.01;***p<0.001

Figure 2. NO rescues the *cygb2* mutant phenotype whereas NO scavenging phenocopies the mutant. (A) Representative frames from fluorescent time-lapse imaging of live *cygb2wt*;Tg(*myl7*:EGFP) and *cygb2^{801a};*Tg(*myl7*:EGFP) mutants at 3dpf, showing the effect of treatment with a pharmacological NO scavenger (cPTIO) and NO donor (DETA) on diastolic volume (EGFP, green). Scale bar, 50µm. **(B)** Comparison of SV at 3dpf in untreated control and DETA treated, and **(C)** cPTIO treated embryos, treatments starting at 2ss. **(D)** Comparison of stroke volumes following chronic treatments starting at different developmental stages (dome, shield, 2ss, and 30hpf). Student t-Test; *ns*, not significant; *p<0.05; **p<0.01;***p<0.001; ****p<0.0001.

Fig. 3. *cygb2* **mutants show defective cardiac progenitors migration, and CYGB knockdown inhibits cell migration in human airway epithelial cell culture. (A, D)** 3D surface rendering and surface plots from *cygb2wt*;Tg(*myl7*:EGFP) and *cygb2^{801a}*;Tg(*myl7*:EGFP) showing cardiac progenitors at midline convergence (16 ss) and cardiac disc (20 ss), renderings and surface plots generated from whole-mount confocal z-series (EGFP, green). DETA treatment starting at 2ss. Scale bar, 50μm. **(B, E)** Schematics illustrating cardiac progenitor organization and embryonic orientation. **(C, F)** Effect of chronic DETA treatment on cardiac progenitor organization, represented by EGFP positive area. Treatment starting at 2ss. **(G)** Representative bright-field images of human airway epithelial A549 cultures transfected with a non-targeting (si-Control) and *CYGB* targeting (si-Cygb) siRNAs at 0 and 24 hours post scratch. **(H)** Comparison of percent migration between si-Control and si-Cygb at 1.25pmol and 2.5pmol. **(I)** Representative images of Western blot bands for CYGB and β-Actin. **(J)** Percent CYGB protein knockdown in si-Cygb at 25pmol relative to si-Control. Student t-Test; *p<0.05; **p<0.01.

Fig. 4. Ventricle area, ventricular diastolic volume, and stroke volume are reduced, and wall thickness is increased in sGC mutants. (A) Representative frames from DIC time-lapse imaging of live *gucy1a1wt* and *gucy1a1^{9bp}* mutant embryos at 3 dpf showing volume at diastole and systole. Scale bar, 50μM. **(B)** Schematic of ventricle position and cardiac orientation. **(C-E)** Comparison of stroke volume, diastolic volume, and systolic volume between *gucy1a1wt*, and *gucy1a1^{9bp}*. **(F-H)** Comparisons of stroke volume, diastolic volume, and systolic volume at 3dpf in live untreated and Bay58 treated *cygb2^{801a}*;Tg(*myl7*:EGFP) embryos using fluorescent time-lapse imaging, treatments starting at 2ss. **(I,L)** Frames from live DIC time-lapse imaging of *gucy1a1wt*, *gucy1a1^{9bp}* , *cygb2wt* and *cygb2^{801a}* mutants at 3 dpf, depicting ventricle wall thickness at diastole and systole. Scale bar, 50μM. **(L)** DETA and Bay58 treatment starting at 2ss. **(J,K,M,N)** Comparison of wall thickness at diastole and systole. Student t-Test; *ns*, not significant; *p<0.05; **p<0.01;***p<0.001; ****p<0.0001.

Fig. 5. Early patterning of the anterior lateral plate mesoderm is disrupted in *cygb2* and *gucy1a1* mutants and rescued by NO–sGC activation. (A-B) Representative in situ hybridization images (top) and quantification of *drl* and *nkx2.5* expression patterns (bottom) in embryos at 10-12ss. **(A)** *drl* normal expression pattern is quantified in red as percentage of embryos with distinct, short, and wide patches of bilateral *drl* expression at anterior edge of ALPM. In blue, percentage of embryos with elongated incongruous thin lines of bilateral *drl* expression along ALMP. **(B)** *nkx2.5* normal expression pattern is quantified in red as percentage of embryos with symmetrical, uniform, and distinct bilateral *nkx2.5* expression along ALMP. In blue, percentage of embryos with asymmetrical, patchy, or faint bilateral *nkx2.5* expression along ALMP. Number of embryos ranked is listed within and on top of bars. Treatments starting at shield. Scale bar, 200µm. On the right, schematic of ALPM anatomy with embryonic orientation. Fishers exact test, ns - not significant ≥ 0.05, * P≤ 0.05, ** P≤ 0.01, *** P≤ 0.001, **** P≤ 0.0001.

Fig. 6. Disrupted specification of endothelial and myocardial lineages persists at 28 hpf and is normalized by restoring NO–sGC signaling. (A-B) Representative in situ hybridization images (top) and quantification of *drl* and *nkx2.5* expression patterns (bottom) in embryos at 28hpf. **(A)** *drl* normal expression pattern is quantified in red as percentage of embryos with expression in the anterior region of the heart tube. In blue, percentage of embryos with complete loss of expression of *drl* in heart tube. **(B)** *nkx2.5* normal expression pattern is quantified in red as percentage of embryos with distinct clearly defined *nkx2.5* expression along the heart tube full-length. In blue, percentage of embryos with reduced or absent *nkx2.5* expression in a shortened heart tube. Number of embryos ranked is listed within and on top of bars. Treatments starting at shield. Scale bar, 200µm. On the right, schematic of the heart tube anatomy with embryonic orientation. Fishers exact test, ns - not significant ≥ 0.05, * P≤ 0.05, ** P≤ 0.01, *** P≤ 0.001, **** P≤ 0.0001.