Cytoglobin-dependent NO-sGC-cGMP signaling regulates ventricular morphogenesis and diastolic function.

Adam Austin Clark[1], Rasmus Hejlesen[3], Tzu-Ting, Weng[1], Muddassar Iqbal[2], Akueba Bruce[2], Paola Corti[1#]

[1]Department of Biochemistry and Molecular Biology, University of Maryland School of Medicine, Baltimore, MD 21201

[2]Department of Medicine, University of Maryland School of Medicine, Baltimore, MD 21201

[3]MRC Human Genetics Unit, Institute of Genetics and Cancer, The University of Edinburgh, Edinburgh, EH4 2XU, UK

#Correspondence: Corti, Paola. Email: PCorti@som.umaryland.edu

Abstract

Aims. Hypoplastic left heart syndrome (HLHS) is a severe congenital heart disease characterized by ventricular hypoplasia and impaired cardiac function. Clinically, inhaled nitric oxide (NO) therapy is used to reduce pulmonary vascular resistance and improve cardiopulmonary stability in HLHS patients. However, whether NO signaling contributes to HLHS pathogenesis remains unknown. Cytoglobin (Cygb) is a heme-protein traditionally thought to limit NO bioavailability and downstream activation of the soluble guanylate cyclase (sGC) cyclic guanosine monophosphate (cGMP) signaling pathway. Unexpectedly, our recent work shows that Cygb enhances NO signaling through activation of NO synthase, leading to downstream activation of sGC-cGMP signaling. In zebrafish embryos, Cygb-dependent NO signaling is required for normal cilia motility and the establishment of correct cardiac laterality. Here, our aim was to determine whether Cygb-dependent NO-sGC signaling linked to cilia function regulates cardiac morphogenesis and contributes to ventricular hypoplasia in HLHS.

Methods and Results. We found that loss of *Cygb* (*cygb2*) in zebrafish disrupts NO-sGC signaling during cardiogenesis, altering cardiac progenitor organization and migration within the anterior lateral plate mesoderm. Disruption of these processes impairs heart tube morphogenesis, thereby producing a compact ventricle wall characterized by increased wall thickness (despite preserved cardiomyocyte number) reduced ventricle size and decreased stroke volume, recapitulating key features of HLHS. Genetic disruption of the sGC α-subunit (*gucy1a1)* and pharmacological NO scavenging phenocopy the *cygb2* mutant phenotype, resulting in reduced cGMP levels, compact ventricular architecture and decreased stroke volume. Consistently, restoration of NO-sGC signaling in *cygb2* mutants rescues early cardiac progenitor patterning, ventricular morphology and stroke volume.

Conclusions. These findings identify Cygb-dependent NO-sGC signaling as a critical developmental pathway for ventricular development and performance, temporally linking cardiac progenitor dynamics to cilia-dependent signaling associated with left-right patterning. This study further suggests that pharmacological activation of sGC may provide a therapeutic strategy for hypoplastic ventricular disease.

Keywords: Hypoplastic left heart syndrome, soluble guanylate cyclase, nitric oxide, cardiac development, zebrafish.

Graphical Abstract

1. Introduction

Hypoplastic left heart syndrome (HLHS) is a complex congenital heart disease with a multigenic etiology.[1,2] Familial clustering supports a genetic basis, yet the majority of causative genes remain unidentified. The phenotype is heterogeneous, variably affecting the left ventricle, valves, and aortic outflow tract, and no single-gene mutation has been shown to reproduce the full HLHS spectrum in animal models. Traditionally, HLHS was attributed to diminished intracardiac flow during development, leading to secondary hypoplasia. However, recent work has shifted this paradigm, suggesting that HLHS may arise from primary defects in myocardial and endocardial development, resembling a developmental cardiomyopathy in origin rather than a purely hemodynamic lesion.[3] Additionally, an emerging body of evidence links HLHS to disturbances of left–right patterning and cilia function, suggesting a new developmental mechanism contributing to disease pathogenesis.[4] Based on case reports and large registry-based analyses,[4,5] HLHS has been observed in association with heterotaxy, situs inversus, gut malrotation, biliary atresia and splenic anomalies.[6-8] These observations further support the concept that defects in laterality determination and ventricular morphogenesis may share developmental origins. According to recent interpretations, HLHS may originate from early intrinsic disruptions in cardiac progenitors arising during and prior to heart tube formation,[9] and may occur in parallel with, or as a consequence of, abnormal left-right signaling. Findings from recessive forward genetic screens in fetal mice further support this view, revealing that cilia and cilia-transduced signaling pathways play central roles in HLHS pathogenesis.[10]

Integrative genetic analysis combining human sequencing with developmental datasets and mouse models of HLHS, identified CYGB as a candidate gene within a developmental gene network associated with the disease,[11] suggesting a potential role for CYGB in pathways regulating early cardiac morphogenesis. CYGB is a hexa-coordinate heme-containing globin expressed ubiquitously in vertebrate tissues.[12,13] Unlike its penta-coordinate relatives, hemoglobin and myoglobin, which primarily

bind and transport oxygen, CYGB is more prone to function as a redox-active enzyme regulating NO

metabolism through reversible redox cycling of its heme iron.[14] The reduced heme iron can catalyze

nitrite reduction to bioactive NO,[13,15] while the oxygenated heme iron can dioxygenate NO to inert

nitrate,[16-18] thus maintaining oxygen-dependent NO homeostasis. While these reactions are possible

in vitro, mouse studies indicate that the predominant physiological role of CYGB is to scavenge NO,

modulating vascular tone through NO deoxygenation[19] and to protect cells against oxidative stress via

superoxide dismutase-like activity.[20] Unexpectedly, we discovered a new critical role for Cygb in

development, where Cygb instead enhances NO signaling. The zebrafish *Cygb* (*cygb2)* knockout

exhibits cardiac laterality defects due to impaired NO signaling, a phenotype characteristic of human

disorders associated with defective cilia function. Mechanistically, we demonstrated that Cygb2

localizes to cilia within the zebrafish laterality organ (Kupffer's vesicle, KV), and increases NO

bioavailability through activation of the nitric oxide synthase enzyme (Nos2b).[21] Loss of *cygb2* disrupts

ciliary structure, motility and directional fluid flow, resulting in impaired KV activity and cardiac laterality

defects.[21] Whether this function is directly linked to cardiac morphogenesis mimicking HLHS remains

unknown and the extent to which Cygb-dependent regulation of laterality intersects, potentially through

ciliary mechanisms, has not been explored.

Zebrafish cardiac development initiates within the anterior lateral plate mesoderm (ALPM), where

bilateral populations of mesodermal progenitors undergo progressive spatial organization prior to overt

myocardial differentiation.[22,23] Between the 14–16 somite stages (ss), these progenitors undergo tightly

coordinated mediolateral convergence and anterior–posterior elongation to assemble a coherent

cardiac disc that subsequently develops into the linear heart tube.[23,24] At these stages of cardiac

development, ventricular formation is exquisitely sensitive to perturbations in progenitor number and

spatial positioning as even subtle defects in coordinated cell migration or tissue architecture can give

rise to abnormalities in ventricular morphogenesis, whereas atrial development may remain relatively

preserved due to its later progenitor contribution.[24,25] Importantly, these morphogenetic events critical to ventricular development occur immediately downstream of KV mediated left–right axis establishment,[23,26] raising the possibility that laterality patterning cues coordinating embryonic asymmetry also influence spatial patterning and migratory behavior of cardiac progenitors, shaping ventricular growth and chamber morphogenesis. One of the earliest markers of cardiac progenitors in zebrafish is *draculin* (*drl*), a transcription factor activated during gastrulation that delineates bilateral cardiac progenitor fields within the ALPM.[23,27] Alterations in ALPM spatial geometry, such as anterior–posterior elongation of the *drl* domain, reflect early disruption of progenitor distribution or specification,[23] defects expected to impair migratory cues and ventricle assembly. As development proceeds, myocardial *drl* expression is restricted to first heart field-derived cardiomyocytes, providing a readout of cardiac disc formation and heart tube assembly, stages that directly precede ventricular chamber growth and functional maturation.[23]

NKX2 homeobox 5 (*Nkx2.5*) encodes a conserved cardiac homeobox transcription factor expressed within the forming cardiac disc at the transition from migratory cardiac progenitors to differentiated myocardium.[23,27] In both mouse models and human patients, dysregulation of NKX2.5 activity is associated with congenital heart diseases and hypoplastic ventricular phenotypes[28]. In the zebrafish embryo, *nkx2.5* expression is conserved and is initially induced in bilateral ALPM domains as cardiac progenitors converge toward the midline, coinciding with progenitor stabilization and the initiation of chamber-specific transcriptional programs.[29] By 28 hpf, *nkx2.5* expression marks specified cardiomyocytes within the heart tube, predominantly ventricular myocardium,[24] such that its expression at this stage reflects proper myocardial differentiation and maintenance of cardiac identity.

While Cygb-dependent regulation of embryonic laterality is now established,[21] its role in coordinating cardiac progenitor migration and myocardial differentiation remains unknown. Here, we demonstrate that *cygb2* deficiency in zebrafish disrupts early cardiac progenitor organization and ventricular

chamber formation, producing a small, compact ventricle with reduced diastolic filling. These features parallel ventricular hypoplasia and impaired diastolic function observed in HLHS. Crucially, these defects are phenocopied in sGC mutants and fully rescued by pharmacological activation of NO-sGC signaling, identifying a developmental pathway that links ciliary NO regulation to cardiac progenitor morphogenesis and ventricular growth.

2. Methods

All animal studies were approved by the University of Maryland Institutional Animal Care and Use Committee (IACUC-1022015), Baltimore, MD, USA. The programs and facilities are USDA registered and covered under an assurance with the Office of Lab Animal Welfare (OLAW). Procedures followed guidelines outlined by the NIH Guide for the Care and Use of Laboratory Animals (8th Edition, 2011). Embryos were maintained at 28.5 °C until the desired developmental stage. Animals were euthanized according to approved protocols; embryos >3 days post fertilization (dpf) to adult (anesthetic overdose 300mg/L buffered tricane), embryos <3dpf (paraformaldehyde immersion).

2.1 Animal Studies

For all experiments, clutch-matched embryos from dome stage (~ 4hpf) to 5dpf were randomly assigned to treatment groups; sex cannot be determined at these stages. The AB (WT) genetic background was used for generation of mutant and transgenic lines. Transgenic lines in this study include Tg(*myl7:EGFP*) and Tg(*tbx5a:EGFP*; *tcf21:dsRED*). The *cygb2[801a]* and the *gucy1a1[9bp]* mutants were generated using CRISPR-Cas9 genome editing as previously described[21] and guide RNAs are listed in[21] and *Table S1*. Embryos were raised and screened for restriction enzyme site disruption (N1aIII) using primers specific to exon 4 of *gucy1a1* (*Table S1*). The Tg(*myl7:EGFP*) and Tg*(tbx5a:EGFP);*Tg*(tcf21:dsRED)* lines were crossed with *cygb2[801a]* to produce *cygb2[801a]*;Tg*(myl7:EGFP)*, *cygb2[801a]*;Tg*(tbx5a:EGFP);*Tg*(tcf21:dsRED),* and wildtype sibling lines.

2.2 Stroke Volume

Stroke volumes were calculated from live imaging. To calculate end-diastolic volume (EDV) and end-systolic volume (ESV) frames were identified from high-speed wide-field (bright-field and fluorescent) time-lapse videos of beating zebrafish hearts at the moments of maximal and minimal ventricular cavity size. The long-axis length (L) and short-axis diameter (D) of the ventricle were measured from frames using FIJI imaging software. EDV and ESV were calculated using the established prolate spheroid model[30] according to the formula:

$$V = \frac{\pi}{6} LD^2$$

L represents ventricular long-axis length and D the diameter at mid-ventricle (*Figure S1*). Stroke volume (SV) was derived as SV = EDV − ESV. Ventricle wall thickness was calculated at maximal and minimal cavity size, identified from high-speed differential interference contrast (DIC) microscopy videos of beating zebrafish hearts. The measurable area and length of the ventral-facing ventricle wall was calculated using FIJI imaging software and the polygon selections and segmented line tool. The area of the selected polygon was divided by the length of the measured wall to generate a ratio-based metric of thickness. Bright field and fluorescent images were captured using a Zeiss Axio Observer inverted microscope with attached Axiocam 807 mono camera and a Leica M165FC with attached K3C camera.

2.3 Ventricle Volume and morphology

Whole mount confocal z-series and compiled z-projections were collected on paraformaldehyde fixed zebrafish embryos from 16ss to 3 days post fertilization. Immuno-labeled embryos were mounted on glass slides and imaged on a Nikon Ti2 inverted W1 spinning disk confocal microscope with attached Hamamatsu sCMOS camera. Collected z-series were compiled into max projections and 3D objects with surface renderings using FIJI and Imaris imaging software. Ventricle volume and wall thickness were calculated using thresholding on selected regions of interest (ROI). Z-slices covering full depth and width of ventricle were compiled, and images were converted to 8-bit and set to grey scale. To

calculate the volume of myocardium, thresholding was set to black and white, area measurements for signal-positive region were measured for each z-slice, summed, and multiplied by the distance between z-slices. Ventricular lumen volume was calculated by measuring the area of signal-negative region within the surrounding signal-positive region, summing the areas across z-slices and multiplying by z-slice distance. To create a ratio-based metric to measure thickness across a 3-dimensional structure, myocardium volume was divided by lumen volume. Cell density was calculated by dividing the total cardiomyocyte number by the myocardium volume. Total ventricle cardiomyocyte number was calculated using the collected z-series and FIJI imaging software. Z-slices spanning the full depth and width of the ventricle were compiled, and analyzed frame by frame, and EGFP-positive cardiomyocytes were marked, tracked, and counted. Ventricle volume at 5 dpf was generated from z-series collected using a Zeiss LS7 Lighsheet confocal microscope. Ventricle area of z-slices was calculated by thresholding (Huang Over Under) of the EGFP positive region within a ROI defining the ventricle including the luminal space. Using FIJI and open-source macro code ventricular volume was calculated by multiplying the summed areas by the distance between z-slices. Progenitor cell fusion, cardiac disc, and heart tube areas were calculated using thresholding of EGFP and CY5 positive areas in selected ROIs of max z-projections collected using spinning disk confocal microscopy.

2.4 In situ hybridization

Whole-mount in situ hybridization was performed on paraformaldehyde fixed zebrafish embryos as previously described[21] at 10-12ss and 28 hpf using riboprobes targeting *drl*, *kinase insert domain receptor like* (*kdrl*), and *nkx2.5*. Transcripts were amplified by PCR and cDNA was obtained from pooled embryos using transcript specific primers (*Table S1*). Images were collected using bright-field microscopy on a Zeiss Discovery V.8 SteReo stereoscope with attached Axiocam 503 color camera, brightness and contrast were unaltered, and images were formatted using FIJI imaging software. To determine expression patterning variance, raw counts of phenotype rankings were used for statistical

comparisons with the Fishers exact test, percentages were plotted for graphical representation, ns - not significant ≥ 0.05, * P≤ 0.05, ** P≤ 0.01, *** P≤ 0.001, **** P≤ 0.0001.

Drug Exposure

Carboxy-PTIO potassium salt (2-4-carboxyphenyl-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide; cPTIO, Sigma–C221) and BAY 582667 (Cinaciguat, Bay58, Sigma–SML1532) were diluted in DMSO and used at final concentrations of 500µM and 120µM respectively. (Z)-1-(N,N-diethylamino)diazen-1-ium-1,2-diolate (DETA NONOate) (Cayman-82100) and (Z)-1-[N-(2-aminoethyl)-N-(2-ammonioethyl)amino]diazen-1-ium-1,2-diolate (PAPA NONOate) (Cayman-2140) were resuspended in sodium hydroxide and diluted in phosphate buffered saline at a final concentration of 250µM and 5.73µM. Drug exposure was achieved by incubation in embryo medium (E3).

Immunohistochemistry

Whole-mount immunohistochemistry was performed as previously described[21] using primary antibodies for myosin heavy chain 6 (Developmental Studies Hybridoma Bank S46). Embryos were mounted on glass slides with Vectashield Hardset Antifade Mounting Medium (H-1500-10) and imaged on a Nikon Ti2 inverted W1 spinning disk confocal microscope equipped with an Hamamatsu sCMOS camera.

cGMP ELISA

cGMP levels in zebrafish embryos were measured by competitive ELISA following manufactures protocol (Enzo-ADI-900-164). Embryos at 28 hpf were incubated in 250µM DETA NONOate for 30 minutes at 28.5 °C. Fifty embryos per sample were pooled and homogenized in 0.1M hydrochloric acid on ice. Lysates were centrifuged and the supernatant was collected for analysis. Samples were

assayed in triplicate at two dilutions. cGMP concentration was calculated based on interpolation to the

standard curve using a 95% confidence interval.

Cell culture

The human type II alveolar epithelial cell line A549 was kindly provided by Dr. Alan Cross (Professor,

Department of Microbiology and Immunology, University of Maryland School of Medicine). A549 cells

were cultured in RPMI-1640 media (Gibco-11875093) containing Penicillin/Streptomycin (Sigma-

P4333) and fungizone (Amphoterecin B, Gibco-15290-026), and 10% fetal bovine serum (FBS) (Sigma-

F4135) at 37^0 Celsius and 5% carbon dioxide. Cell cultures were passaged a minimum of three times

before performing experiments.

Small interfering RNA

Dicer Substrate RNA duplexed small interfering RNAs si-CYGB 13.1 and si-CYGB 13.2 targeting

human *CYGB* were selected using Integrated DNA Technologies "Predesigned DsiRNA Selection Tool".

DsiRNA-mediated gene silencing was validated through transfection of sub-confluent A549 monolayers

using Lipofectamine RNAiMAX transfection reagent (Thermo-13778030), non-targeting and human

CYGB targeting siRNAs for 48 hours. Si-CYGB 13.1 showed higher protein knockdown in Western Blot

experiments and was chosen for migration assays. To assess cell viability, transfected cells were

pelleted and resuspended in a 1:1 mixture of cell suspension and 0.4% Trypan Blue. Cells were counted

using a hemocytometer in a 0.05μm section. Cells excluding Trypan Blue were counted as live and

cells stained with Trypan Blue were counted as dead, viability was calculated as a percent of live cells

to total cell number.

Western Blot

A549 cells were lysed on ice in RIPA buffer (Sigma-R0278) containing protease inhibitor (Halt 1X,

Thermo-1860932). Lysates were centrifuged at 12,500 RPM for 20 min at 4°C. The BCA assay was used to determine total protein concentration. Samples were loaded for gel electrophoresis in 4-20% precast gels (Bio-Rad-4561094) using 100 μg total protein per sample. Following electrophoresis protein was transferred to polyvinylidene fluoride membranes. Membranes were blocked with 3% bovine serum albumin (BSA) for 2 hours at room temperature under gentle agitation. Membranes were probed with primary rabbit anti-CYGB (Sigma-HPA017757) diluted at 1:1000 in 1% BSA and 0.1% Triton X100 in phosphate buffer saline (PBST) overnight at 4^0 Celsius. Following 3 consecutive 5-minute washes at room temperature in PBST membranes were probed with anti-Rabbit IgG-HRP (Abcam-131366) at 1:10,000 in PBST with 1% BSA for 1 hour at room temperature. Following 3 consecutive 5-minute washes at room temperature in a solution of PBST protein bands were visualized using Clarity Max Western ECL substrate (Bio-Rad-1705060) following manufacturer's protocol and a Bio-Rad ChemiDoc MP imaging box. Following 3 consecutive 5-minute washes with PBST at room temperature membranes were stripped using Restore Western Blot Stripping Buffer (Thermo-21059). Blocking and probing procedure was repeated using primary rabbit anti-β-Actin antibody (Sigma-A2066). Band intensity was normalized to β-Actin and quantified using FIJI imaging software.

Wound healing assay

A549 cells were seeded in 24-well plates at a density of 170,000 cells per well. After 24 hours cells were transfected with 2.5pmol non-targeting DsiRNA, 1.25pmol and 2.5 pmol si-CYGB 13.1. At 48 hours post transfection, a scratch was introduced with a 10μl pipette tip. Following the scratch, cells were washed with sterile 1x PBS and supplemented with RPMI-1640 media containing 10% (FBS). Cultures were imaged at 0 and 24 hours post scratch on a Leica DM IL LED inverted microscope using a 4X objective. Wound healing was assessed as percent migration into the scratch area relative to the migration of control. Image analysis was conducted using FIJI imaging software.

2.5 Statistical analysis

Unless otherwise noted data are presented as mean ± SD, and statistical significance was determined using an unpaired two-tailed *t*-test (or Mann–Whitney test for non-normal distributions), ns - not significant ≥ 0.05, * P≤ 0.05, ** P≤ 0.01, *** P≤ 0.001, **** P≤ 0.0001.

3. Results

Cytoglobin is required for ventricular development and cardiac function

We previously defined a role for Cygb2 in cardiac laterality via regulation of cilia function. Here, we investigated whether Cygb2 is required for cardiac morphogenesis and performance and whether these are linked to laterality establishment. To visualize cardiac morphology, we generated a *cygb2* mutant reporter line by crossing the previously characterized *cygb2[801a]* mutant[21] with a transgenic reporter line expressing EGFP under the control of the *myosin light chain 7* (*myl7*) promoter.[23] In zebrafish, *myl7* expression begins at approximately the 16ss stage in cardiomyocyte progenitors regardless of eventual chamber contribution to the atrium or ventricle.[23] We analyzed *myl7:EGFP* expression at 3 days post fertilization (dpf), when cardiac morphology is established, and the ventricle and atrium exhibit coordinated contraction and robust cardiac function. We first quantified the area of both cardiac chambers and observed a significant decrease in the ventricle of *cygb2[801a]* embryos compared to wildtype controls, (*Figure 1A–C, Movies S1,2*), while atrial area remained unchanged (*Figure 1D*). To precisely assess performance, we measured end diastolic volume (EDV) and end systolic volume (ESV) from in vivo high-speed recordings and calculated stroke volumes (SV) using the prolate spheroid method *(Figure S1)*. Ventricles of *cygb2[801a]* embryos showed a marked decrease in SV and EDV with no change in ESV (*Figure 1E–H*), consistent with reduced diastolic filling rather than systolic impairment. As *cygb2[801a]* embryos exhibited cardiac laterality defects, we tested whether this ventricular defect might arise secondarily from laterality disturbances. We compared embryos with normal (situs solitus) and reversed (situs inversus) cardiac orientation within each genotype and

detected no differences in SV, *cygb2[801a]* embryos displayed reduced SV relative to wildtype

independent of cardiac left-right orientation (*Figure S2A–F*). Assessment of SV at 5dpf revealed that

this phenotype persisted beyond initial ventricle morphogenesis (*Figure S3A–D, Movies S3-4*).

Additionally, ventricle volume was measured using light sheet confocal microscopy on fixed *myl7:EGFP*

samples and further supported the reduced ventricle volume observation and persistence at 5 dpf in

cygb2[801a] compared to wildtype (*Figure S3E*). Collectively, these findings indicate a stable structural

hypoplasia rather than a transient developmental delay. Despite reduced chamber size, ventricular

cardiomyocyte number was unchanged (*Figure 1 I-J*). Instead, cells occupied a smaller volume, yielding

a more compact myocardium and a thicker ventricular wall (*Figure 1K–M*). These data indicate that

cygb2 deficiency primarily affects ventricular morphogenesis and chamber remodeling, not

cardiomyocyte proliferation or survival. The reduced, thick-walled ventricle observed in *cygb2[801a]*

embryos accompanied by reduced SV mirrors key structural features of HLHS thus *cygb2* deficiency

may represent a mechanistically relevant model of HLHS-like ventricular hypoplasia.

Nitric oxide restores ventricular size and function in *cygb2* mutants.

We previously showed that Cygb2 regulates cardiac laterality determination[21] through NO signaling. To

test whether the reduced SV in *cygb2[801a]* embryos depends on NO, we first confirmed the acute

physiological action of NO donors. Ten-minute exposure at 78 hpf to the pharmacological NO donor

DETA-NONOate (DETA) increased SV in both wildtype and *cygb2[801a]* embryos *(Figure S4A,B),*

consistent with the reported hemodynamic effect of NO in teleost fish larvae[31] where the increase in

SV results from reduced afterload due to relaxation of the outflow vasculature. A fast-releasing NO

donor, PAPA-NONOate (PAPA), produced a similar acute increase, validating the assay (*Figure*

S4C,D). To determine whether the altered ventricular developmental defects could be rescued by NO

supplementation, we tested the effect of chronic DETA treatment. DETA was administered in a vehicle

(sodium hydroxide), which alone did not affect SV (*Figure S4E*). We performed DETA treatments from

the 2ss stage to 3 dpf. Chronic exposure produced opposite outcomes depending on genotype, in *cygb2*[801a] embryos ventricular size increased with consequent rescue in SV, whereas in wildtype embryos ventricular size was reduced, resulting in decreased SV (*Figure 2A,B*). Because prolonged NO exposure affects cardiac morphogenesis rather than acute mechanics, these bidirectional responses under identical dosing indicate that *cygb2* mutants possess a specific deficit in NO-dependent developmental signaling, while excess NO in wildtype embryos disrupts the finely tuned program governing ventricle morphogenesis. To provide further evidence supporting NO dependence, we treated embryos at 2ss with the NO scavenger cPTIO to phenocopy the cardiac phenotype observed in *cygb2* mutants (*Figure 2A,C*). As expected, ventricular volume was reduced by NO scavenging in wildtype embryos and was further exacerbated in *cygb2*[801a] embryos, consistent with a requirement for NO signaling in normal ventricular development. A time course initiating DETA at stages from dome to 12ss showed maximal rescue when treatment began at dome stage to 2ss, diminishing by 12ss (*Figure 2D*), suggesting Cygb2-dependent NO signaling becomes critical for cardiac specification and morphogenesis during early somitogenesis. Together, these findings demonstrate that *cygb2* is required to maintain appropriate NO levels during early cardiac progenitor organization, thereby ensuring proper ventricular morphogenesis and SV.

Loss of Cygb2 disrupts early cardiomyocyte migration.

Given that the ventricular phenotype observed in *cygb2*[801a] embryos originates during early development, we next examined cardiac morphogenesis to determine when these defects first emerge. In zebrafish, cardiac progenitors originate bilaterally within the ALPM and progressively migrate toward the midline, where they fuse to form the cardiac cone and subsequently the primitive heart tube. To assess whether this process was altered in *cygb2* mutants, we analyzed the distribution of *myl7*-positive cardiomyocyte precursors during heart field assembly. At 16ss when cardiomyocyte progenitors have initiated migration but have not yet fused, *cygb2*[801a] embryos displayed a significantly broader *myl7*-

positive domain compared with wildtype, indicating an expanded or delayed coalescence of cardiac progenitors at this stage (*Figure 3A-C*). By 20ss, when the cardiac disc becomes morphologically visible at the midline, the *myl7*-positive region remained wider and less compacted in *cygb2*[801a] embryos, occupying a larger total area relative to controls (*Figure 3D-F*). These results suggest that Cygb2 is required for the coordinated medial convergence of cardiomyocyte progenitors during heart field assembly. As development progressed, the cardiac tube of *cygb2*[801a] embryos remained visibly smaller than in wildtype. At 34 and 48 hpf, the ventricular area was significantly reduced indicating that the morphogenetic defect persists through the stages of heart tube elongation and chamber ballooning, whereas the atrium appeared morphologically normal throughout (*Figure S5A-H).* This suggests a ventricle-specific effect consistent with the smaller, compact ventricular morphology observed at later stages. To establish NO as a critical regulator throughout ventricular morphogenesis, embryos were treated with the NO donor DETA beginning at 2ss. These treatments restored *myl7*-positive cardiac field at 16 and 20ss (*Figure 3A-F)* and rescued ventricular area at 34 and 48 hpf (*Figure S5A-H*) to near wildtype levels*.* These findings demonstrate that *cygb2* is essential for the proper migration and coalescence of cardiac progenitors and for subsequent expansion of the ventricular chamber, and that both processes depend on NO-mediated signaling.

Upon observing aberrant organization of cardiac progenitors during midline convergence and cardiac disc formation in *cygb2*[801a] embryos, we directly tested whether cell migratory behavior was disrupted using an in vitro wound-healing assay in the human type II alveolar epithelial A549 cell line. *CYGB* transcript knockdown by small interfering RNA caused a dose-dependent reduction in cell migration into the wound area (*Figure 3G,H*). CYGB protein knockdown efficiency and cell viability were confirmed by Western Blot analysis (*Figure 3I,J and S6A*) and viability assay (*Figure S6B*). These data suggest that abnormal cardiac progenitor contribution to early heart structures may result from disruption of cellular machinery regulating cell migration and/or cell adhesion and further indicate that Cygb-NO signaling influences migratory behavior across cell types and species.

Soluble guanylate cyclase (sGC) is required for ventricle development and heart function.

NO can signal through multiple mechanisms, including protein S-nitrosylation, nitration reactions and formation of reactive nitrogen species. However, our data indicate that here NO primarily signals through its canonical receptor sGC. Upon ligand binding, NO-sGC catalyzes the conversion of guanosine triphosphate (GTP) to cGMP, thereby initiating secondary downstream signaling events[32]. To test whether the cardiac phenotype observed in *cygb2[801a]* embryos depends on sGC, we generated an sGC mutant (*gucy1a1[9bp]*) using CRISPR/Cas9-mediated site-directed mutagenesis. Genomic DNA from embryos derived from a maternal zygotic mutant line revealed a 9 bp deletion in exon 4 of the sGCα1 subunit, which was confirmed at the genomic and mRNA level by Sanger Sequencing (*Figure S7A*). Loss of the normal transcript was validated by in situ hybridization using a 300 bp probe spanning the edited region, which showed markedly reduced hybridization signal in *gucy1a1[9bp]* embryos compared to wildtype, whereas a 1000 bp probe located downstream of the edited site produced comparable staining in both mutants and wild type (*Figure S7B*). Functional loss of sGC activity in *gucy1a1[9bp]* embryos was confirmed by failure of DETA to restore cGMP levels in *gucy1a1[9bp]* compared to wildtype (*Figure S7E*). In contrast, DETA treatment restored cGMP levels in *cygb2[801a]* embryos,[21] supporting the interpretation that Cygb2 functions upstream of sGC to regulate NO-dependent cGMP production during cardiac development. To determine whether sGC disruption phenocopies the *cygb2[801a]* cardiac defects observed here, we quantified ventricular performance in *gucy1a1[9bp]* embryos using DIC microscopy and high-speed imaging. SV was significantly reduced in *gucy1a1[9bp]* compared with wildtype embryos, driven by a decrease in EDV while ESV remained unchanged (*Figure 4A-E, Movies S5,6*). This pattern mirrors the functional phenotype of *cygb2[801a]* embryos, indicating that loss of sGCα1 function recapitulates the diastolic filling defect observed in the *cygb2* mutant. Furthermore, we tested SV at 5dpf in *gucy1a1* mutants and found that in both, reduced SV persisted at later developmental stages (*Figure S7F-H, Movies S7,8*). To confirm that the observed phenotype reflects disruption of a shared Cygb2-NO-sGC signaling pathway in both *cygb2[801a] and gucy1a1[9bp]* mutants, we

tested whether pharmacological activation of sGC in *cygb*^{801a} mutants could rescue cardiac function.

Treatment of *cygb*^{801a} mutants with the sGC activator Bay58 from 2ss to 34hpf fully restored SV (*Figure*

4F-H), confirming a critical role for sGC signaling in ventricular formation. Notably, neither DETA nor

Bay58 rescued stroke volume in *gucy1a1*^{9bp} mutants, as expected (*Figure S7C-D*). To assess a shared

phenotype of increased ventricle wall thickness, measurements were performed on live embryos at

3dpf. Wall thickness at diastole and systole was increased in *gucy1a1*^{9bp} (*Figure 4I-K, Movies S5,6*),

similar to that observed in *cygb2*^{801a} mutants (*Figure 4L-N, Movies S9,10*), indicating a conserved

phenotype across mutant models and analytic approaches. To further determine whether NO-sGC

signaling contributes to this phenotype, *cygb2*^{801a} mutants were treated from 2ss to 34hpf with DETA

and Bay58. These conditions fully restored ventricular wall thickness (*Figure 4L-N),* supporting a role

for Cygb-NO-sGC in ventricular morphogenesis.

NO-sGC signaling regulates early cardiac progenitor organization.

To determine whether Cygb2 and Gucy1a1 act through a shared molecular pathway during early

cardiogenesis, we analyzed key markers of the ALPM at 10–12ss, when early cardiac and endothelial

progenitors first emerge. In wildtype embryos, *drl* expression appeared as bilateral short, symmetrical

stripes flanking the midline, marking anterior mesoderm that gives rise to cardiac and vascular lineages.

In both *cygb2*^{801a} and *gucy1a1*^{9bp} embryos, *drl* domains were elongated, thinner, and extended

posteriorly, indicating disrupted convergence and defective compaction of the ALPM (*Figure 5A*).

Similarly, *kdrl*, which labels endothelial and endocardial progenitors, was expressed bilaterally in

compact domains close to the midline in wildtype embryos, but it was elongated, more widely

separated, and extended posteriorly in *cygb2*^{801a} and *gucy1a1*^{9bp} mutants (*Figure S8*). This finding is

consistent with delayed or uncoordinated migration of endothelial/endocardial precursors. Despite

these spatial changes, signal intensity was not diminished, suggesting that endothelial specification

may   be   largely   preserved,   while   progenitor   patterning   appears   shifted   from   anterior

(endothelial/endocardial) toward more posterior (hematopoietic) mesodermal domains. Expression of

the early cardiac transcription factor *nkx2.5* was also altered at 10-12ss. In wildtype embryos, *nkx2.5*

was expressed as two distinct, symmetrical domains within the ALPM, whereas in *cygb2^{801a}* and

gucy1a1^{9bp} mutants, expression was frequently asymmetric or spatially disorganized (*Figure 5B*),

suggesting loss of coordination between left and right cardiac fields. Importantly, wildtype siblings

generated during establishment of the maternal-zygotic *cygb2^{801a}* and *gucy1a1^{9bp}* lines were

outcrossed to minimize potential CRISPR off-target effects and displayed normal *drl*, *kdrl* and *nkx2.5*

expression patterns at all developmental stages examined (*Figure S9*). The *drl* expression patterning

abnormalities were rescued in *cygb2^{801a}* embryos treated with either DETA or Bay58 (*Figure 5A*),

confirming that early ALPM organization and synchronized cardiac specification depend on Cygb2-

mediated NO–sGC signaling. Additionally, *nkx2.5* expression was also partially normalized by DETA

treatment, although the strongest normalization was achieved in embryos treated with Bay58 (*Figure*

5B), confirming the importance of sGC signaling in this process. As *drl* marks early cardiac precursors

fields and *nkx2.5* labels a broader population encompassing later myocardial progenitors, this

differential rescue further supports the conclusion that NO–sGC signaling exerts its strongest influence

during the earliest stages of cardiac precursor organization and initial heart structure formation.

NO–sGC signaling rescues ventricular gene expression.

To assess whether early ALPM patterning defects persist during later stages of cardiogenesis, we

examined *drl*, and *nkx2.5* expression at 28 hpf, when the heart tube has formed and endothelial and

myocardial differentiation are underway. In wildtype embryos, *drl* expression is confined to the

developing vasculature and the endocardium; in contrast, *cygb2^{801a}* and *gucy1a1^{9bp}* embryos exhibited

reduced or absent *drl* expression in the heart tube (*Figure 6A*). This pattern is consistent with the shift

observed at earlier stages from anterior (endothelial/endocardial) toward more posterior

(hematopoietic) mesodermal domains. Consistently, *nkx2.5* expression in the ventricle was markedly

reduced in both *cygb2[801a]* and *gucy1a1[9bp]* embryos, indicating impaired myocardial contribution to this

region (*Figure 6B*). Treatment of *cygb2[801a]* embryos with DETA or Bay58 restored *drl* and *nkx2.5*

expression patterns to near wildtype levels (*Figure 6A-B*). Together, these findings show that *cygb2*

and *gucy1a1* mutants share a common set of early and late ALPM abnormalities that can be corrected

by restoring NO-sGC signaling, linking early ALPM mis-patterning to the later emergence of a compact,

hypoplastic ventricle.

4. Discussion

Nitric oxide (NO) signaling is a central regulator of cardiovascular physiology, with established roles in

vascular tone, myocardial contractility, and hemodynamic homeostasis.[33-35] In contrast, the contribution

of NO signaling to early cardiac morphogenesis has been far less explored. Here, we show that Cygb2-

dependent NO–sGC signaling is required for ventricular formation during zebrafish heart development.

Loss of *cygb2* leads to a reduced SV associated with decreased ventricle volume, increased myocardial

wall thickness, elevated cellular density, and impaired diastolic filling. These defects arise from

abnormal migratory behavior of cardiac precursors, which display reduced migratory efficiency,

excessive cell-spreading, and subsequent hypercompaction within the forming heart tube. Importantly,

genetic loss of sGC (*gucy1a1*) fully phenocopies the *cygb2* mutant phenotype, and pharmacological

activation of NO-sGC signaling in *cygb2* mutants rescues ventricular volume, cardiac progenitor

migration, myocardial organization within the forming heart tube, and diastolic function. Together, these

data identify NO–sGC signaling downstream of Cygb2 as a central pathway coordinating myocardial

progenitor dynamics, ventricular morphogenesis, and functional maturation.

Cardiac progenitor migration is a fundamental driver of heart tube assembly and ventricular

morphogenesis and depends on tightly regulated cell polarity, adhesion turnover, and cytoskeletal

dynamics.[36-39] Disruption of these processes can profoundly alter chamber size and myocardial

architecture.[40,41] In this context, our data indicate that reduced NO–sGC signaling places cardiac

progenitors into a mechanically altered migratory state. NO-sGC-cGMP signaling is known to regulate actomyosin contractility, focal adhesion turnover, and cortical tension across multiple cellular contexts[42-44] and cytoglobin-dependent control of NO bioavailability represents a key upstream determinant of this pathway[13,14]. Attenuation of NO–sGC signaling during development is therefore predicted to stabilize cell–substrate and cell–cell adhesions, resulting in excessive cell spreading, reduced migratory efficiency, and impaired cellular rearrangements at the tissue level. When such mechanically altered cells enter the confined environment of the forming heart tube, increased adhesion and cortical tension can bias morphogenesis toward tissue compaction and elevated cellular packing as demonstrated in epithelial and mesenchymal morphogenetic models.[45,46] These properties provide a mechanistic basis for the increased ventricular cell density observed in *cygb2* mutants, together with the resulting thick-walled ventricle and impaired diastolic function.

We previously demonstrated that Cygb2 localizes to motile cilia and regulates cilia-dependent left–right patterning through NO–sGC signaling during early zebrafish development,[21] establishing that Cygb2-dependent NO–sGC signaling operates during early developmental windows critical for laterality specification. The present study extends this role to ventricular formation while showing that ventricular defects do not arise directly from altered left–right axis specification (*Figure S2*). However, ventricular morphogenesis may still depend on cilia-associated signaling while remaining independent of final laterality outcomes. Here, we found that pharmacological restoration of NO–sGC signaling rescues ventricular morphology only when administered during early windows encompassing laterality determination and early somitogenesis, whereas later treatment fails to restore ventricular size and stroke volume, indicating that ventricular morphogenesis is temporally linked to early cilia-associated signaling events. During this developmental window, we found that *drl* and *kdrl* expression is spatially displaced and expanded within the ALPM, phenotypes consistent with defective cardiac progenitor migration rather than primary loss of cardiogenic identity[36,38,39,41]. Notably, *nkx2.5*, normally expressed symmetrically by early somitogenesis[47] is found asymmetric in *cygb2* mutants, suggesting altered

organization of subsets of cardiac progenitors. Recent single-cell studies indicate that *nkx2.5* marks transcriptionally distinct progenitor populations within the ALPM[29] supporting the idea that the disrupted signaling perturbs progenitor composition rather than uniformly impairing cardiac fate. Early migratory defects likely lead secondarily to impaired differentiation or maintenance programs later, as reflected by the reduction of *nkx2.5* and *drl* expression within ventricular territories at 28 hpf.[48,49] Importantly, dysregulation of NKX2-5 is associated with several forms of congenital heart disease in humans, including ventricular septal defects, tetralogy of Fallot, and HLHS, underscoring the relevance of these findings to human cardiac pathogenesis.[47,50,51]

Sequencing studies have identified enrichment of rare variants in cilia and ciliopathy-associated genes in subsets of HLHS patient cohorts,[4] supporting a framework in which cilia-dependent pathways contribute to disease susceptibility in a subset of patients. Within this framework, zebrafish models exhibiting reduced ventricular size with compact myocardium have emerged as informative systems for studying cardiac morphogenesis and HLHS-like phenotypes.[9,49] In our *cygb2* mutant model, HLHS-like ventricular hypoplasia originates from early events concomitant with cilia-dependent processes involved in left-right patterning. Our data indicate two possible mechanistic pathways underlying this phenotype. In one scenario, impaired NO-sGC signaling caused by loss of Cygb2 directly disrupts the coordination of cardiac progenitor migration, regional specification, and ventricle expansion, leading to reduced ventricular growth. In a second, distinct scenario, loss of Cygb2 perturbs cilia-dependent processes that regulate early cardiac morphogenesis and myocardial growth, beyond their established role in cardiac laterality, through downstream effects on progenitor deployment and tissue remodeling,[50] independent of left–right axis determination. These mechanisms are not mutually exclusive and may interact in the multigenic context of human HLHS, where combined defects in NO signaling and cilia-associated pathways could converge on shared morphogenetic processes controlling ventricular development. The *cygb2 and gucy1a1* models characterized here converge on the hallmark ventricular phenotype of HLHS while revealing a distinct initiating axis centered on Cygb-dependent NO–sGC

signaling. Together with prior work demonstrating that modulation of NO bioavailability influences myocardial plasticity and regeneration,[52] these findings establish NO–sGC signaling as a tractable developmental pathway whose disruption leads to ventricular hypoplasia and whose restoration is sufficient to rescue ventricular structure and function. Our data indicate that pharmacological activation of sGC, previously unexplored in the context of hypoplastic ventricular disease, can restore ventricular structure and function in HLHS-like phenotypes in vivo. These results position zebrafish as a valuable model for interrogating early developmental mechanisms underlying ventricular hypoplasia and for assessing sGC-targeted interventions in CHD.

Author Contributions

Adam Clark, Paola Corti designed and conducted experiments, analyzed data, and wrote the manuscript. Paola Corti directed the research and conceptualization of experiments. Rasmus Hejlesen and Muddassar Iqbal reviewed the manuscript. Rasmus Hejlesen constructed a data set for transcript variance analysis that guided hypothesis generation. Weng Tzu-Ting conducted in situ hybridization experiments. Muddassar Iqbal designed, conducted; and analyzed data for siRNA knockdown, wound healing, and Western Blot experiments. Akueba Bruce conducted zebrafish husbandry and genotyping.

Acknowledgements

We are thankful to Dr. Mark T. Gladwin (Dean, University of Maryland School of Medicine) for expert mentorship and guidance in project design. We thank Dr. Joseph Mauban, Dr. Shilpa Kumar and the University of Maryland School of Medicine Center for Innovative Biomedical Resources, Confocal Microscopy Core for invaluable assistance in imaging, and the Translational Genomics Lab for Sanger Sequencing services.

Conflict of Interest: The authors declare no competing interests that influence the work reported in this paper.

Funding

This work was supported by National Institutes of Health under awards number 5T32AR007592 and R01HL168775.

Data availability

The data used to produce this article will be shared at reasonable request to the corresponding author.

References

1. Iascone M, Ciccone R, Galletti L, Marchetti D, Seddio F, Lincesso AR, Pezzoli L, Vetro A, Barachetti D, Boni L, Federici D, Soto AM, Comas JV, Ferrazzi P, Zuffardi O. Identification of de novo mutations and rare variants in hypoplastic left heart syndrome. *Clin Genet* 2012;**81**:542–554.

2. Yagi H, Liu X, Gabriel GC, Wu Y, Peterson K, Murray SA, Aronow BJ, Martin LJ, Benson DW, Lo CW. The Genetic Landscape of Hypoplastic Left Heart Syndrome. *Pediatr Cardiol* 2018;**39**:1069–1081.

3. Grossfeld P, Nie S, Lin L, Wang L, Anderson RH. Hypoplastic Left Heart Syndrome: A New Paradigm for an Old Disease? *J Cardiovasc Dev Dis* 2019;**6**.

4. Gabriel GC, Lo CW. Left-right patterning in congenital heart disease beyond heterotaxy. *Am J Med Genet C Semin Med Genet* 2020;**184**:90–96.

5. Lin AE, Krikov S, Riehle-Colarusso T, Frias JL, Belmont J, Anderka M, Geva T, Getz KD, Botto LD, National Birth Defects Prevention S. Laterality defects in the national birth defects prevention study (1998-2007): birth prevalence and descriptive epidemiology. *Am J Med Genet A* 2014;**164A**:2581–2591.

6. Becker DJ, Islam S, Geiger JD. Biliary atresia associated with hypoplastic left heart syndrome: a case report and review of the literature. *J Pediatr Surg* 2004;**39**:1411–1413.

7. Stankiewicz P, Sen P, Bhatt SS, Storer M, Xia Z, Bejjani BA, Ou Z, Wiszniewska J, Driscoll DJ, Maisenbacher MK, Bolivar J, Bauer M, Zackai EH, McDonald-McGinn D, Nowaczyk MM, Murray M, Hustead V, Mascotti K, Schultz R, Hallam L, McRae D, Nicholson AG, Newbury R, Durham-O'Donnell J, Knight G, Kini U, Shaikh TH, Martin V, Tyreman M, Simonic I, Willatt L, Paterson J, Mehta S, Rajan D, Fitzgerald T, Gribble S, Prigmore E, Patel A, Shaffer LG, Carter NP, Cheung SW, Langston C, Shaw-Smith C. Genomic and genic deletions of the FOX gene

cluster on 16q24.1 and inactivating mutations of FOXF1 cause alveolar capillary dysplasia and other malformations. *Am J Hum Genet* 2009;**84**:780–791.

8. Huseynova RA, Bin Mahmoud LA, Abdelrahim A, Alroiedy MA, Huseynov O. Polysplenia syndrome with complex heart disease and jejunal atresia with malrotation in neonate: A case report. *Clin Case Rep* 2020;**8**:848–851.

9. Huang M, Akerberg AA, Zhang X, Yoon H, Joshi S, Hallinan C, Nguyen C, Pu WT, Haigis MC, Burns CG, Burns CE. Intrinsic myocardial defects underlie an Rbfox-deficient zebrafish model of hypoplastic left heart syndrome. *Nat Commun* 2022;**13**:5877.

10. Yagi H, Lo CW. Left-Sided Heart Defects and Laterality Disturbance in Hypoplastic Left Heart Syndrome. *J Cardiovasc Dev Dis* 2023;**10**.

11. Liu X, Yagi H, Saeed S, Bais AS, Gabriel GC, Chen Z, Peterson KA, Li Y, Schwartz MC, Reynolds WT, Saydmohammed M, Gibbs B, Wu Y, Devine W, Chatterjee B, Klena NT, Kostka D, de Mesy Bentley KL, Ganapathiraju MK, Dexheimer P, Leatherbury L, Khalifa O, Bhagat A, Zahid M, Pu W, Watkins S, Grossfeld P, Murray SA, Porter GA, Jr., Tsang M, Martin LJ, Benson DW, Aronow BJ, Lo CW. The complex genetics of hypoplastic left heart syndrome. *Nat Genet* 2017;**49**:1152–1159.

12. Trent JT, 3rd, Hargrove MS. A ubiquitously expressed human hexacoordinate hemoglobin. *J Biol Chem* 2002;**277**:19538–19545.

13. Corti P, Ieraci M, Tejero J. Characterization of zebrafish neuroglobin and cytoglobins 1 and 2: Zebrafish cytoglobins provide insights into the transition from six-coordinate to five-coordinate globins. *Nitric Oxide* 2016;**53**:22–34.

14. Amdahl MB, Sparacino-Watkins CE, Corti P, Gladwin MT, Tejero J. Efficient Reduction of Vertebrate Cytoglobins by the Cytochrome b(5)/Cytochrome b(5) Reductase/NADH System. *Biochemistry* 2017;**56**:3993–4004.

15. Reeder BJ, Ukeri J. Strong modulation of nitrite reductase activity of cytoglobin by disulfide bond

oxidation: Implications for nitric oxide homeostasis. *Nitric Oxide* 2018;**72**:16–23.

16. Halligan KE, Jourd'heuil FL, Jourd'heuil D. Cytoglobin is expressed in the vasculature and

regulates cell respiration and proliferation via nitric oxide dioxygenation. *J Biol Chem*

2009;**284**:8539–8547.

17. Gardner AM, Cook MR, Gardner PR. Nitric-oxide dioxygenase function of human cytoglobin with

cellular reductants and in rat hepatocytes. *J Biol Chem* 2010;**285**:23850–23857.

18. Tejero J, Shiva S, Gladwin MT. Sources of Vascular Nitric Oxide and Reactive Oxygen Species

and Their Regulation. *Physiol Rev* 2019;**99**:311–379.

19. Zweier JL, Ilangovan G. Regulation of Nitric Oxide Metabolism and Vascular Tone by Cytoglobin.

Antioxid Redox Signal 2020;**32**:1172–1187.

20. Zweier JL, Hemann C, Kundu T, Ewees MG, Khaleel SA, Samouilov A, Ilangovan G, El-Mahdy

MA. Cytoglobin has potent superoxide dismutase function. *Proc Natl Acad Sci U S A*

2021;**118**.

21. Rochon ER, Xue J, Mohammed MS, Smith C, Hay-Schmidt A, DeMartino AW, Clark A, Xu Q, Lo

CW, Tsang M, Tejero J, Gladwin MT, Corti P. Cytoglobin regulates NO-dependent cilia motility

and organ laterality during development. *Nat Commun* 2023;**14**:8333.

22. Yelon D. Cardiac patterning and morphogenesis in zebrafish. *Dev Dyn* 2001;**222**:552–563.

23. Kemmler CL, Riemslagh FW, Moran HR, Mosimann C. From Stripes to a Beating Heart: Early

Cardiac Development in Zebrafish. *J Cardiovasc Dev Dis* 2021;**8**.

24. Felker A, Prummel KD, Merks AM, Mickoleit M, Brombacher EC, Huisken J, Panakova D,

Mosimann C. Continuous addition of progenitors forms the cardiac ventricle in zebrafish. *Nat*

Commun 2018;**9**:2001.

25. Paffett-Lugassy N, Novikov N, Jeffrey S, Abrial M, Guner-Ataman B, Sakthivel S, Burns CE,

Burns CG. Unique developmental trajectories and genetic regulation of ventricular and outflow

tract progenitors in the zebrafish second heart field. *Development* 2017;**144**:4616–4624.

26. Hirokawa N, Tanaka Y, Okada Y. Left-right determination: involvement of molecular motor KIF3,

cilia, and nodal flow. *Cold Spring Harb Perspect Biol* 2009;**1**:a000802.

27. Mosimann C, Panakova D, Werdich AA, Musso G, Burger A, Lawson KL, Carr LA, Nevis KR,

Sabeh MK, Zhou Y, Davidson AJ, DiBiase A, Burns CE, Burns CG, MacRae CA, Zon LI.

Chamber identity programs drive early functional partitioning of the heart. *Nat Commun*

2015;**6**:8146.

28. Pashmforoush M, Lu JT, Chen H, Amand TS, Kondo R, Pradervand S, Evans SM, Clark B,

Feramisco JR, Giles W, Ho SY, Benson DW, Silberbach M, Shou W, Chien KR. Nkx2-5

pathways and congenital heart disease; loss of ventricular myocyte lineage specification leads

to progressive cardiomyopathy and complete heart block. *Cell* 2004;**117**:373–386.

29. Coskun H, Wang Y, Gao X, Chen K, Geoffrey Burns C, Burns CE. 3D spatial organization of

heterogeneous nkx2.5+ progenitors in the zebrafish heart field pre-patterns cardiovascular

development. *Nat Commun* 2025;**16**:11624.

30. Yalcin HC, Amindari A, Butcher JT, Althani A, Yacoub M. Heart function and hemodynamic

analysis for zebrafish embryos. *Dev Dyn* 2017;**246**:868–880.

31. Fritsche R, Schwerte T, Pelster B. Nitric oxide and vascular reactivity in developing zebrafish,

Danio rerio. *Am J Physiol Regul Integr Comp Physiol* 2000;**279**:R2200–2207.

32. Arnold WP, Mittal CK, Katsuki S, Murad F. Nitric oxide activates guanylate cyclase and increases

guanosine 3':5'-cyclic monophosphate levels in various tissue preparations. *Proc Natl Acad Sci*

U S A 1977;**74**:3203–3207.

33. Ignarro LJ. Nitric Oxide: A Unique Endogenous Signaling Molecule in Vascular Biology (Nobel

Lecture). *Angew Chem Int Ed Engl* 1999;**38**:1882–1892.

34. Gladwin MT. Deconstructing endothelial dysfunction: soluble guanylyl cyclase oxidation and the

NO resistance syndrome. *J Clin Invest* 2006;**116**:2330–2332.

35. Forstermann U, Sessa WC. Nitric oxide synthases: regulation and function. *Eur Heart J*

2012;**33**:829–837, 837a–837d.

36. Stainier DY. Zebrafish genetics and vertebrate heart formation. *Nat Rev Genet* 2001;**2**:39–48.

37. Buckingham SD, Ali DW. Computer simulations of high-pass filtering in zebrafish larval muscle

fibres. *J Exp Biol* 2005;**208**:3055–3063.

38. Abu-Issa R, Kirby ML. Heart field: from mesoderm to heart tube. *Annu Rev Cell Dev Biol*

2007;**23**:45–68.

39. Bakkers J. Zebrafish as a model to study cardiac development and human cardiac disease.

Cardiovasc Res 2011;**91**:279–288.

40. Hutson MR, Kirby ML. Model systems for the study of heart development and disease. Cardiac

neural crest and conotruncal malformations. *Semin Cell Dev Biol* 2007;**18**:101–110.

41. Vincent SD, Buckingham ME. How to make a heart: the origin and regulation of cardiac progenitor

cells. *Curr Top Dev Biol* 2010;**90**:1–41.

42. Surks HK, Mochizuki N, Kasai Y, Georgescu SP, Tang KM, Ito M, Lincoln TM, Mendelsohn ME.

Regulation of myosin phosphatase by a specific interaction with cGMP- dependent protein

kinase Ialpha. *Science* 1999;**286**:1583–1587.

43. Schlossmann J, Hofmann F. cGMP-dependent protein kinases in drug discovery. *Drug Discov*

Today 2005;**10**:627–634.

44. Prysyazhna O, Rudyk O, Eaton P. Single atom substitution in mouse protein kinase G eliminates

oxidant sensing to cause hypertension. *Nat Med* 2012;**18**:286–290.

45. Lecuit T, Lenne PF. Cell surface mechanics and the control of cell shape, tissue patterns and

morphogenesis. *Nat Rev Mol Cell Biol* 2007;**8**:633–644.

46. Heisenberg CP, Bellaiche Y. Forces in tissue morphogenesis and patterning. *Cell* 2013;**153**:948–

962.

47. Sultana N, Nag K, Hoshijima K, Laird DW, Kawakami A, Hirose S. Zebrafish early cardiac

connexin, Cx36.7/Ecx, regulates myofibril orientation and heart morphogenesis by establishing

Nkx2.5 expression. *Proc Natl Acad Sci U S A* 2008;**105**:4763–4768.

48. Moorman AF, Christoffels VM. Cardiac chamber formation: development, genes, and evolution.

Physiol Rev 2003;**83**:1223–1267.

49. Liu J, Stainier DY. Zebrafish in the study of early cardiac development. *Circ Res* 2012;**110**:870–

874.

50. Sutherland MJ, Ware SM. Disorders of left-right asymmetry: heterotaxy and situs inversus. *Am J*

Med Genet C Semin Med Genet 2009;**151C**:307–317.

51. George V, Colombo S, Targoff KL. An early requirement for nkx2.5 ensures the first and second

heart field ventricular identity and cardiac function into adulthood. *Dev Biol* 2015;**400**:10–22.

52. Rochon ER, Missinato MA, Xue J, Tejero J, Tsang M, Gladwin MT, Corti P. Nitrite Improves Heart

Regeneration in Zebrafish. *Antioxid Redox Signal* 2020;**32**:363–377.

Figure 1

Figure 2

737   **Figure 3**

Figure 4

750 **Figure 5**

755 **Figure 6**

Figure Legends

Figure 1. Loss of *cygb2* leads to ventricular hypoplasia and reduced cardiac output. (A) Representative confocal projections of *cygb2wt*;Tg(*myl7*:EGFP) and *cygb2^{801a}*;Tg(*myl7*:EGFP) hearts at 3 days post fertilization (dpf), immuno-labeled for Myh6 (EGFP, green; Myh6, magenta). Scale bar, 50μm **(B)** Schematic illustrating heart chamber anatomy and cardiac orientation. A-atrium; V-ventricle; B-bulbus arteriosus **(C,D)** Comparison of ventricle and atrium area. **(E)** Representative frames from fluorescence time-lapse imaging of live *cygb2wt*;Tg(*myl7*:EGFP) and *cygb2^{801a}*;Tg(*myl7*:EGFP) embryos 3 dpf showing ventricle volume at diastole and systole, dashed white line demarcates the ventricle (EGFP, green). Scale bar, 50 μm. **(F-H)** Comparison of stroke volume, diastolic volume and systolic volume. **(I)** Representative confocal projections of ventricles at 3 dpf, images are greyscale converted and inverted to highlight cardiomyocyte density. Scale bar, 20μm. **(J)** Comparison of ventricle cell number. **(K)** Representative confocal z-slices at ventricular midpoint. Images were converted to greyscale signal positive area (ventricular myocardium) shown in black. **(L)** Comparison of cardiomyocyte density*.* **(M)** Comparison of ventricular wall thickness. Sample size (n) corresponds to independent embryos represented by individual data points on plots. Student t-Test; *ns*, not significant; *p<0.05; **p<0.01;***p<0.001

Figure 2. NO rescues the *cygb2* mutant phenotype, whereas NO scavenging phenocopies the mutant. (A) Representative frames from fluorescent time-lapse imaging of live *cygb2wt*;Tg(*myl7*:EGFP) and *cygb2^{801a}*;Tg(*myl7*:EGFP) mutants at 3 days post fertilization (dpf), showing the effect of treatment with a pharmacological NO scavenger (cPTIO) and NO donor (DETA) on diastolic volume (EGFP, green). Scale bar, 50μm. **(B)** Comparison of stroke volume at 3dpf in untreated control and NO donor (DETA) treated, and **(C)** NO scavenger (cPTIO) treated embryos,

treatments starting at 2 somite stage (ss). **(D)** Comparison of stroke volumes following chronic

treatments starting at different developmental stages (dome, shield, 2ss, and 30hpf). Sample size (n)

corresponds to independent embryos represented by individual data points on plots. Student t-Test; *ns*,

not significant; *p<0.05; **p<0.01;***p<0.001; ****p<0.0001.

Fig. 3. *cygb2* mutants show defective cardiac progenitors migration, and CYGB knockdown

inhibits cell migration in human airway epithelial cell culture. (A,D) 3D surface rendering and

surface plots from *cygb2wt*;Tg(*myl7*:EGFP) and *cygb2^{801a}*;Tg(*myl7*:EGFP) showing cardiac progenitors

at midline convergence, 16 somite stage (ss), and cardiac disc, 20 ss, renderings and surface plots

generated from whole-mount confocal z-series (EGFP, green). NO donor (DETA) treatment starting at

2ss. Scale bar, 50µm. **(B,E)** Schematics illustrating cardiac progenitor organization and embryonic

orientation. **(C,F)** Effect of chronic DETA treatment on cardiac progenitor organization, represented by

EGFP positive area. Treatment starting at 2ss. **(G)** Representative bright-field images of human airway

epithelial A549 cultures transfected with a non-targeting (si-Control) and *CYGB* targeting (si-CYGB

13.1) siRNAs at 0 and 24 hours post scratch. **(H)** Comparison of percent migration between si-Control

and si-*CYGB* at 1.25 pmol and 2.5 pmol. **(I)** Representative images of Western blot bands for CYGB

and β-Actin. **(J)** Percent CYGB protein knockdown with si-*CYGB* at 25 pmol relative to si-Control. **(A-**

F) Sample size (n) corresponds to independent embryos **(G-J)** Sample size (n) corresponds to

independent cultures, all represented by individual data points on plots. Student t-Test; *p<0.05;

**p<0.01.

Fig. 4. Ventricle area, ventricular diastolic volume, and stroke volume are reduced, and wall

thickness is increased in sGC mutants. (A) Representative frames from DIC time-lapse imaging of

live *gucy1a1wt* and *gucy1a1⁹ᵇᵖ* mutant embryos at 3 days post fertilization (dpf) showing volume at diastole and systole. Scale bar, 50µM. **(B)** Schematic of ventricle position and cardiac orientation. **(C-E)** Comparison of stroke volume, diastolic volume, and systolic volume between *gucy1a1wt*, and *gucy1a1⁹ᵇᵖ*. **(F-H)** Comparisons of stroke volume, diastolic volume, and systolic volume at 3dpf in live untreated and sGC activator (Bay58) treated *cygb2⁸⁰¹ᵃ*;Tg(*myl7*:EGFP) embryos using fluorescent time-lapse imaging, treatments starting at 2 somite stage (ss). **(I,L)** Frames from live DIC time-lapse imaging of *gucy1a1wt*, *gucy1a1⁹ᵇᵖ* , *cygb2wt* and *cygb2⁸⁰¹ᵃ* mutants at 3 dpf, depicting ventricle wall thickness at diastole and systole. Scale bar, 50µM. **(L)** NO donor (DETA) and Bay58 treatment starting at 2ss. **(J,K,M,N)** Comparison of wall thickness at diastole and systole. Sample size (n) corresponds to independent embryos represented by individual data points on plots. Student t-Test; *ns*, not significant; *p<0.05; **p<0.01;***p<0.001; ****p<0.0001.

Fig. 5. Early patterning of the anterior lateral plate mesoderm is disrupted in *cygb2* and *gucy1a1* mutants and rescued by NO–sGC activation. (A,B) Representative in situ hybridization images (top) and quantification of *drl* and *nkx2.5* expression patterns (bottom) in embryos at 10-12 somite stage. **(A)** *drl* normal expression pattern is quantified in solid-bars as percentage of embryos with distinct, short, and wide patches of bilateral *drl* expression at anterior edge of anterior lateral plate mesoderm (ALPM). In open-bars, percentage of embryos with elongated incongruous thin lines of bilateral *drl* expression along ALMP. **(B)** *nkx2.5* normal expression pattern is quantified in solid-bars as percentage of embryos with symmetrical, uniform, and distinct bilateral *nkx2.5* expression along ALMP. In open-bars, percentage of embryos with asymmetrical, patchy, or faint bilateral *nkx2.5* expression along ALMP. Number of embryos ranked is listed within and on top of bars. Treatments starting at shield. Scale bar, 200µm. On the right, schematic of ALPM anatomy with embryonic orientation. Fishers exact test, ns - not significant ≥ 0.05, * P≤ 0.05, ** P≤ 0.01, *** P≤ 0.001, **** P≤ 0.0001.

Fig. 6. Disrupted specification of endothelial and myocardial lineages persists at 28 hours post fertilization and is normalized by restoring NO–sGC signaling. (A,B) Representative in situ hybridization images (top) and quantification of *drl* and *nkx2.5* expression patterns (bottom) in embryos at 28 hours post fertilization. **(A)** *drl* normal expression pattern is quantified in solid-bars as percentage of embryos with expression in the anterior region of the heart tube. In open-bars, percentage of embryos with complete loss of expression of *drl* in heart tube. **(B)** *nkx2.5* normal expression pattern is quantified in solid-bars as percentage of embryos with distinct clearly defined *nkx2.5* expression along the heart tube full-length. In open-bars, percentage of embryos with reduced or absent *nkx2.5* expression in a shortened heart tube. Number of embryos ranked is listed within and on top of bars. Treatments starting at shield. Scale bar, 200µm. On the right, schematic of the heart tube anatomy with embryonic orientation. Fishers exact test, ns - not significant ≥ 0.05, * P≤ 0.05, ** P≤ 0.01, *** P≤ 0.001, **** P≤ 0.0001.