Variable performances of commercial eDNA inventories challenge their use for surveying stream fish communities

Roussel JM.[1,§], Quémeré E.[1], Bonnet B.[1,2], Covain R.[3], Dézerald O.[1], Lassalle G.[1], Le Bail PY.[4], Petit E.J.[1], Pottier G.[2], Quartarollo G.[2,5], Vigouroux R.[2], Lalagüe H.[2]

[1] UMR DECOD (Ecosystem Dynamics and Sustainability), INRAE, Institut Agro, IFREMER, Rennes, France

[2] HYDRECO Guyane, Kourou, French Guiana

[3] Museum of Natural History, Geneva, Switzerland

[4] LPGP (laboratoire de physiologie et génomique des poissons), INRAE, Rennes, France

[5] Guyane Wild Fish, Cayenne, French Guiana

§ Correspondence: jean-marc.roussel@inrae.fr

Highlights

- eDNA metabarcoding of water samples is widely used to detect species in streams
- Detection performances of 4 private companies were compared to an exhaustive fish inventory
- The number of undetected species varies from 32 to 92% depending on the company
- Such discrepancies challenge the use of eDNA for fish-based ecological assessments

Authors' contributions

JMR, EQ, RV and HL conceived the ideas and designed methodology; GP, RV, HL, GQ collected the data at field; PYLB, RC, GQ identified the species and standardized the names of taxa that the companies provided; BB, GL, JMR, OD and EP analysed and interpreted the data; JMR and HL led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

Acknowledgements

We are indebted to the Direction Generale des Territoires et de la Mer (DGTM Guyane) for financial support and to Nicolas Heitz for insightful discussions on the objectives and implementation of this project.

Data is available on request. No conflict of interest to declare.

Abstract

1. Environmental DNA (eDNA) metabarcoding of water samples is increasingly used to detect fish species in streams. Several studies have concluded that it can outperform traditional inventory methods and recommend using it at large scales for fish-based ecological assessments. However, there is no standard protocol that can guarantee sufficient detection rates and repeatability, despite companies offering an extensive range of analyses.

2. We compared eDNA metabarcoding performed by four companies. Following their guidelines, samples were collected in a small tropical stream in the Maroni River (French Guiana) that hosts a species-rich fish community. We compared their inventories to each other and to a list of species captured during an extensive fish inventory performed immediately after sampling eDNA, as well as to current data on the species' distributions.

3. The number of species detected by eDNA metabarcoding ranged from 5 to 48 among the companies, but these inventories contained many inaccuracies. All companies combined, 63 species were detected, of which 10 (16%) had never been reported in the Maroni River. The extensive inventory identified 50 species in the local fish community, of which 16-46 were not detected by eDNA metabarcoding (i.e. false negative detection rate of 32%-92% among the companies).

4. Reanalysis of raw sequencing data decreased differences among companies greatly, highlighting the importance of using a comprehensive and accurate DNA barcode database to assign species. Dissimilarity indices, calculated to compare the local fish community (based on presence/absence or fish catches) to eDNA detection, revealed large differences regardless of the company.

5. Summary and applications. The large percentage of species not detected by eDNA metabarcoding of water samples could strongly bias fish-diversity inventories in streams that host species-rich communities. This issue is not well documented in the literature, and we recommend that similar studies in the future focus on other stream contexts. The large differences between commercial eDNA inventories and the local fish community challenge the use of eDNA metabarcoding for fish-based ecological assessments of streams. The variable performance of eDNA companies indicates the need for a standard protocol and access to a comprehensive DNA database before beginning large-scale eDNA programmes.

Keywords: environmental DNA, mitochondrial 12S primers, metabarcoding, species-rich fish community, detection rate, ecological assessment, French Guiana

Introduction

Environmental DNA (eDNA) refers to the genetic material that organisms leave behind in the environment. It is used to detect the presence of species in environmental samples, which allows biodiversity to be studied with an unprecedented potential for action. After the pioneering study two decades ago (Ficetola et al., 2008), eDNA was soon promoted as the ultimate biomonitoring tool (Baird & Hajibabaei, 2012). Inexpensive, accurate, simple to use in the field, and ethically-based, eDNA metabarcoding was quickly set to enhance, or even replace, traditional methods based on capturing or observing organisms, thus helping policy making and future practices better prevent further biodiversity loss (Taberlet et al., 2012). This has resulted in a large amount of research and development, with nearly 1000 articles about the use of eDNA for biomonitoring published over the past decade (Blackman et al., 2024). When no previous knowledge about a community is available, eDNA metabarcoding has become the most widely used approach, as it can identify and inventory species using molecular primers that target broad taxonomic groups (Cordier et al., 2020; Deiner et al., 2017). Many studies have compared eDNA metabarcoding to traditional methods, and they often reported that the former performed as well as or better overall than the latter (*e.g.* Andres et al., 2023; Bonk et al., 2024; Czeglédi et al., 2021; Jannel et al., 2024; Keck et al., 2022; Liu et al., 2024). This has led many scientists and government agencies to recommend large-scale use of eDNA metabarcoding in monitoring programmes, especially for freshwater ecosystems (*e.g.* Keck et al., 2022; Kelly et al., 2024; Pawlowski et al., 2018; Stein et al., 2023). However, closer examination of studies that compared it to traditional methods for freshwater ecosystems revealed that its performance may be exaggerated due to several unresolved problems, and that key challenges remain to be addressed before it is routinely used for biomonitoring.

First, eDNA metabarcoding of water samples is usually compared to traditional sampling protocols, which also inventory only some of the species present. Few field studies have combined eDNA metabarcoding with exhaustive inventories of communities to explore the former's limitations (but see Blabolil et al., 2021; Di Muri et al., 2020), and the former's seemingly better performance should not mask its potential lack of completeness in the field. In general, the absence of a species is difficult to confirm unless extensive sampling is performed, and eDNA metabarcoding is no exception. Some organisms release DNA fragments into aquatic environments, depending on their taxonomy, metabolism, and ontogenetic stage (Múrria et al., 2024; Poyntz-Wright et al., 2024; Tréguier et al., 2014; Verdier et al., 2024; Yates et al., 2019; Yates et al., 2021a; Yates et al., 2021b). Thus, a species may be present, but its eDNA concentrations lie below the detection threshold. The eDNA metabarcoding protocol includes several key steps – water sampling in the field, DNA extraction and amplification in the laboratory, DNA sequencing, and bioinformatic analysis (Deiner et al., 2017) –

during which the eDNA signal of a species may be lost. The risk of misinterpreting non-detection of
eDNA as true species absence (*i.e.* false negative detection) exists and must be carefully considered
when using eDNA (Lacoursière-Roussel & Deiner, 2021).

Second, several studies have found large differences between species lists identified using eDNA
metabarcoding and traditional methods, especially at the local scale (Brantschen & Altermatt, 2024;
Cote et al., 2023; Keck et al., 2022; Kelly, 2019; Liu et al., 2024). Besides animal movement, water
currents can transport DNA fragments over long distances (Civade et al., 2016; Deiner & Altermatt,
2014; Harrison et al., 2019). Notably, rivers act as "conveyor belts" of biodiversity information (Deiner
et al., 2016; Littlefair et al., 2023) in which the area represented by an eDNA sample depends on the
persistence, shedding, decay, and downstream transport of DNA, all of which vary greatly and are
influenced by complex environmental interactions among discharge, exposure to ultraviolet light,
substrate composition, biofilm and microbial activity, and suspended-sediment characteristics
(Condachou et al., 2025; Deiner & Altermatt, 2014; Harrison et al., 2019; Hervé et al., 2023; Mauvisseau
et al., 2022; McKnight et al., 2024; Nevers et al., 2020; Scriver et al., 2023; Shogren et al., 2017; Snyder
et al., 2023; Turner et al., 2015). Animal waste can also spread eDNA in the environment (Inoue et al.,
2023; Stoeckle et al., 2016), such as when a predator excretes the remains of its prey outside of the
prey's foraging habitats (Guilfoyle & Schultz, 2017; Merkes et al., 2014). These factors imply that
detecting a species' DNA in a geolocated environmental sample does not necessarily indicate its actual
presence at that location, and that eDNA samples may represent a different spatial scale than those of
traditional methods. Unlike methods based on capturing animals, eDNA introduces a new source of
error – false positive detection – whose extent remains largely unexplored.

Third, the influence of false negatives and positives on sampling design and statistical procedures is
not sufficiently recognised. The development of eDNA metabarcoding as a tool in biodiversity studies
has been driven by technological revolutions, especially the advent of high-throughput sequencing
technologies and the initial realisation that DNA can be extracted from environmental samples
(Ficetola et al., 2008). Despite the early recommendation of Yoccoz (2012) that the eDNA community
should collaborate with quantitative ecologists, relevant sampling designs are rarely adopted, which
may hinder their ability to provide information about the state of and changes in biodiversity (Yoccoz
et al., 2001). A good indicator for eDNA inventories is the ratio of biological replicates (i.e. water
samples) to technical replicates (i.e. Polymerase Chain Reaction; PCR), which is usually low. Most eDNA
protocols prefer PCR replicates to water-sample replicates, but both are needed to assess potential
sources of errors correctly (Guillera-Arroita et al., 2017; Lahoz-Montfort et al., 2016). This disregard
for statistical issues is especially prevalent for aquatic environments, as demonstrated by recent
reviews that did not address them (*e.g.* Bruce et al., 2021; Bunholie et al., 2023; Xing et al., 2022). Most

eDNA developments have focused on increasing sample volumes rather than the number of biological
replicates (Blackman et al., 2024), and using statistical models to quantify errors remains the exception
(15 of the 431 studies reviewed by Mathieu et al. (2020)).

The increasing demand for eDNA-based tools to monitor biodiversity has spurred the worldwide
emergence in recent years of many companies that provide eDNA metabarcoding, including "turnkey"
eDNA solutions for species inventories. They provide clients (i.e. government agencies, NGOs, and
research institutions) with user-friendly sampling kits that allow the clients to filter water samples
themselves in the field. The companies process the samples in the laboratory, perform the
metabarcoding, and provide a list of the taxa detected, usually with little information about the
reliability of their results. Studies have highlighted the lack of minimum justifications for eDNA
methods and sampling design (Blackman et al., 2024; Mathieu et al., 2020), especially the volume
filtered, filter pore size, and numbers of replicates, which vary greatly among water eDNA studies
(Bunholi et al., 2023; Shu et al., 2020; Xing et al., 2022). Moreover, despite the development of large
curated barcode databases (e.g. BOLD, SILVA, MitoFish) over the past few decades, their lack of
comprehensiveness and accuracy remains a major challenge for implementing biomonitoring
programmes correctly (Weigand et al., 2019), especially for species-rich communities (Keck et al.,
2023; Marques et al., 2021). The development of private databases poses accessibility problems for
users, increases the risk of incorrect taxonomic assignment, and could result in many false positive
and/or false negative detections in eDNA inventories (Jackman et al., 2021). Companies vary greatly in
pricing, methodological transparency, and data-sharing policies, which makes it difficult for non-
academic users to select among them. It also poses challenges for comparing eDNA inventories among
companies, especially for long-term biomonitoring programmes.

Following promising scientific developments, eDNA inventories are increasingly expected to be used
to monitor biodiversity, including for regulatory surveillance of aquatic environments (Hering et al.,
2018; Pawlowski et al., 2018; Pont et al., 2021). The number of tendering processes is likely to increase
in the future, but the accuracy with which commercial eDNA inventories are able to describe fish
communities at a given location is unknown. In the present study, we compared four companies that
perform eDNA metabarcoding of water samples to assess the taxonomic composition of fish
communities. The study was performed using a species-rich stream in French Guiana to challenge the
performance of eDNA metabarcoding. After carefully following each company's guidelines for
sampling water and preserving the samples, we sent the samples to each company without disclosing
the study's objective and the exact sampling location. The species lists that each company provided
were compared to (i) an exhaustive inventory of fish species performed at the sampling site
immediately after sampling the water and (ii) current data on fish distribution in French Guiana. False

negative and false positive eDNA detection rates and fish-community composition indices were

calculated and compared among the companies. Finally, raw sequencing data supplied by three of the

companies were reanalysed using a standard bioinformatic pipeline and a recently published barcode

database of fish species in French Guiana. We then assessed how the data analysis may have

influenced differences among the companies.

Methods

Study site and eDNA sampling

The Bastien stream flows into the Maroni River in French Guiana (Supplementary Material S1). eDNA

was sampled on 18 March 2024 in a 2.5 km tributary of the Bastien stream (5.27048° N, 54.234217°

W; Supplementary Materials S1 and S2). Water temperature (26.1°C), conductivity (32.1 μS.cm^{-1}),

turbidity (2.83 NTU), pH (6.02), and dissolved oxygen (6.52 mg.L^{-1}, 79.8% saturation) were recorded

using a multi-parameter probe (Tripod Ponsel Aqualabo). Stream discharge (15 L.s^{-1}) was estimated

from water depth and velocity measured across the channel section. The four eDNA companies

selected are internationally known. Following their methodological guidelines, stream water was

pumped in running water to ensure that it was sufficiently mixed, and then filtered in the field using

sampling kits provided by each company. Differences in the number of filters used, pore size, filtration

times, and volumes filtered differed (Supplementary Material S3) revealed diverse sampling strategies

among companies. After sampling, the filters were stored according to each company's guidelines and

placed in a cool box for transport to the laboratory.

eDNA samples were sent to the companies for eDNA metabarcoding, without revealing the objective

of the study or the sampling locations, to avoid influencing the processing of samples. We asked them

only to inventory the fish species detected in the samples. As the companies provided few details about

their laboratory workflows, bioinformatic pipelines, or statistical treatments, we were not able to

compare these aspects directly. Three companies stated that they used one primer pair, either 12S-

Teleo (64 bp) (Valentini et al., 2016) (*Teleo*) or 12S-MiFish-U (163-185 bp) (Miya et al., 2015) (*MiFish*),

while the fourth stated that it used several primer pairs (targeting 12S, COI and 16S regions), but did

not specify which ones. Three companies analysed two biological replicates, while the fourth analysed

three (Supplementary Material S3). The number of technical (PCR) replicates per sample varied greatly:

from 1-12 for the three companies that provided this information (Supplementary Material S3). Finally,

three of the four companies agreed to send us their raw sequencing data (Illumina FastQ files). The

companies have been anonymised to avoid influencing their commercial activities.

Exhaustive inventory of the local fish community

Immediately after filtering the water, a barrier net (4 mm mesh) was placed across the channel
immediately on the water-pumping point (Supplementary Material S2). The bottom of the net was
buried in the substrate to prevent fish from escaping underneath. Additional barrier nets were installed
upstream to enclose a 100 m long section and prevent fish from escaping during the inventory. The
mean channel width in the section was 225 cm (range: 150-375 cm). The habitat was a mean of 16.9
cm deep (range: 2-50 cm). The stream bottom consisted mainly of pebbles, sand, and silt, with a few
macrophytes (*Thurnia sphaerocephala*). We inventoried the fish community in the enclosed section
exhaustively using electrofishing. The electrofishing device (described by Pottier et al. (2019)) was set
to deliver 1000 V (DC). The anode was a 350 mm diameter aluminium ring, and the cathode was a
double copper braid (3000 mm long × 10 mm wide × 1 mm thick). Given the ambient conductivity,
these settings ensured effective fish capture while maintaining animal welfare (Pottier et al. (2019) for
details). The fisherman who managed the anode was assisted by two people equipped with dip nets
to capture the fish in the anode's electric field. The entire water surface in the enclosed section was
explored from downstream to upstream, four times in a row, to maximise fish capture.

After capture, each fish was identified to species using the latest taxonomic knowledge about fish in
French Guiana (Le Bail et al., 2012; Le Bail et al., 2026). Fish were measured (standard length) to the
nearest mm and then gradually released downstream of the sampling section. We applied allometric
length-weight relations based on a historical database of fish captures in French Guiana
(https://eseweb.rennes.inrae.fr/RezoFleuve/) (Allard et al., 2015) to estimate the weight of each
individual, and summed the weights to calculate the biomass of each fish species in the community.
Hereafter, "local fish community" refers to the abundances and biomasses of species captured in the
100 m enclosed section.

False positive and false negative eDNA detection rates

The names of taxa that the companies provided were standardised according to Le Bail et al. (2026)
and Fricke et al. (2025). The number of taxa detected at the class, order, genus, and species levels were
counted. Companies were not penalised for incorrect spelling or if the taxonomy at the species level
was still debated by taxonomists. However, taxa reported only to genus, family, order, or class were
excluded when taxonomic knowledge could have been used to identify them to species. Each
company's list was analysed independently before being combined into a single list of all species
detected by eDNA (hereafter, the "molecular community"). We created Venn diagrams to compare
species lists among companies using the *eulerr* package (Larsson, 2024) of R software (R Core Team,
2025). We also used current knowledge about fish distribution to compare each company's species list
to those in the Maroni River catchment, other large rivers in French Guiana, and elsewhere in South
America (Le Bail et al., 2026).

As no company provided information about taxon detection across biological and technical replicates,
we used the fish captures as unambiguous detections (i.e. no false positive detections) (Guillera-
Arroita et al., 2017; Lahoz-Montfort et al., 2016; Miller et al., 2011). Species in the fish community that
were not detected by eDNA metabarcoding were classified as false negative detections, whose rates
were calculated for each company by dividing the number of undetected species by the number of
species captured in the local fish community. The same calculation was used for the molecular
community. Conversely, species detected using eDNA metabarcoding that were absent from the local
fish community were classified as false positive detections, whose rates were calculated for each
company and for the molecular community by dividing the number of species detected using eDNA
barcoding but not captured in the local fish community by the total number of species detected using
eDNA barcoding.

Reanalysis of companies' raw sequencing data

To decrease potential biases related to unequal access to DNA databases among companies and to
highlight differences caused by field sampling and/or laboratory processing, we reanalysed the raw
sequencing data provided by three of the companies using a standard bioinformatic pipeline. Forward
and reverse reads were assembled using the *illuminapairend* function of the *OBITools* v1.2.11 package
(Boyer et al., 2016), and the resulting sequences were trimmed for primers using the cutadapt v1.8.3
program (Martin, 2011). Strictly identical sequences were clustered using the *obiuniq* function
(dereplication step). Amplicon sequence variants represented by fewer than 10 reads were removed
from the dataset. Additionally, sequence lengths were filtered for each primer set: 40-100 and 140-
200 bp for *Teleo* and *MiFish*, respectively. Taxa were assigned using BLASTn (NCBI BLAST+ v2.10.0)
compared to a curated reference database (Brosse et al., 2026; NCBI Accession Number: PX415477 -
PX417033), with a minimum similarity threshold of 97% and 98%, an e-value cut-off of $1e^{-70}$ and $1e^{-20}$,
and a minimum alignment coverage of 87% and 80% for *MiFish* and *Teleo*, respectively. When several
equally good hits occurred, taxa were assigned to the lowest common ancestor. The resulting species
lists were then used to recalculate false positive and false negative detection rates using the procedure
described previously.

Statistical analysis of composition of the local fish community

Species lists provided by the four companies were used to generate presence/absence data, while read
counts from reanalysing companies' raw sequencing data were used as a proxy for fish abundance
(companies A, B, and D only). We compared eDNA (for each company individually or combined) to the
local fish community using four dissimilarity indices: Jaccard and Raup-Crick for presence/absence,
Bray-Curtis and Horn-Morisita for abundance data. Raup-Crick (Chase et al., 2011) and Horn-Morisita

(Chao et al., 2006), which considers the co-occurrence of rare and unshared species among
communities more than the Jaccard and the Bray-Curtis indices do, are more robust to differences in
species composition and sample size. All dissimilarity indices among communities were calculated
using the *vegan* package (Oksanen et al., 2024) and then visualised via principal coordinate analysis,
with Cailliez-transformation when needed, using the *ade4* package (Dray & Dufour, 2007).

Finally, we tested whether the abundance and biomass of species in the local fish community
influenced their detection by eDNA barcoding based on the reanalysis of raw sequencing data. Welch's
t-tests (unequal variance, two samples) were used to compare abundance and biomass of the species
detected, or not, by eDNA metabarcoding. Linear regressions between abundance (or biomass) of
species in the local fish community and read counts were calculated, and their significance was
assessed using F-tests. All analyses were performed in R.

Results

Exhaustive inventory of the local fish community

The 446 fish captured via electrofishing belonged to 6 orders, 22 families, 40 genera, and 50 species
(Supplementary Material S4). The local fish community was dominated by species with low abundance:
56% of the species were represented by five individuals or fewer, including nine species represented
by only one individual (Supplementary Material S4). Conversely, five species were represented by
more than 20 individuals (Supplementary Material S4 and S5). The total biomass captured was 2.037
kg for all species combined, of which 54% consisted of individuals weighing less than 10 g. The total
biomass of 11 species exceeded 50 g each, being largest (428 g) for the predator *Hoplias malabaricus*
(8 individuals).

Assessment of species lists provided by the eDNA companies

The species list reported by each company (Table 1) contained several taxonomic inaccuracies,
including 4 assignments to class (*Actinotperygii*), 1 to order, 5 to family, and 37 to genus. The
companies reported varying levels of taxonomic resolution: the percentage of taxa identified to species
was highest for company A (94%), followed by companies C (61%), B (59%), and D (24%). After
removing inaccurate detections and standardising the taxonomy, the number of species detected by
eDNA ranged from 5-48 among the four companies (Table 1). The molecular community contained 63
species. Sequencing depth varied greatly among companies, with company A producing 12 times as
many reads as company B (Supplementary Table S3). Company A detected the most species and had

the most read counts in its raw sequencing data. See Supplementary Material S6 for the complete
species list reported by each company.

The percentage of species detected by eDNA metabarcoding that were actually in the local fish
community ranged from 41%-80% among the companies (Figure 1). The molecular community
included 57% of the species actually in the local fish community, along with additional species known
to inhabit the Maroni River catchment (27%), other large rivers in French Guiana (3%), or elsewhere in
South America (13%). Company A detected 30 species that the other companies did not (Figure 1),
most of which were species known to inhabit the Maroni River catchment (Supplementary Material
S6). Company B detected 13 unique species, 7 of which had never been reported in the Maroni River
catchment or other large rivers in French Guiana. Companies A and C shared the most detections in
common: 17 species (Figure 1). Ultimately, only four species were detected by all four eDNA companies
(Figure 1).

Reanalysing the raw sequencing data greatly increased the number of species detected (43-52 per
company) and the percentage of species in the local fish community detected (60%-67%) (Figure 1). It
also decreased the differences among companies and increased the number of species detected by all
companies to 34 (Figure 1). Similarly, detections of species not present in the Maroni River catchment
or other large rivers in French Guiana decreased from 10 to 2 (Figure 1). See Supplementary Material
S6 for the complete list of species detected after reanalysing the raw sequencing data of each
company.

Among the nine species of the local fish community that were not reported by any company or
detected after reanalysis of their sequencing files, only one species (*Brachyhypopomus brevirostris*)
was absent from the Brosse et al. (2026) reference database. Another species, *Eleotris pisonis*, was
reported by one company but was not recovered in the reanalysis, due to its absence from the Brosse
et al. (2026) database.

False positive and false negative eDNA detection rates

Comparing eDNA detections to the local fish community revealed that the companies varied greatly in
their ability to detect species (Supplementary Material S7). False negative detection rates ranged from
32%-92% among the companies, whereas false positive detection rates ranged from 20.0%-59.1% (i.e.
1-14 species) (Figure 2). False negative detection rates were slightly lower for the molecular
community. Differences in detection rates among the companies decreased when raw sequencing data
were reanalysed, with false negative detection rates ranging from 30.0%-48.0% and false positive
detection rates ranging from 32.7%-39.5% (Figure 2).

Reanalysing the raw sequencing data revealed that species detected by eDNA metabarcoding were

significantly more abundant in the local fish community than undetected species for companies B and

D (Welch's t-test, p = 0.011 and 0.012, respectively) but not for company A (p = 0.489) (Supplementary

Material S8). Similarly, detected species had significantly higher biomass than undetected species,

regardless of the company (Welch's t-test, p < 0.005) (Supplementary Material S8). We observed no

significant relations between read counts of species and species abundance in the local fish community

(F-test, p > 0.065), but read counts were significantly and positively correlated with species biomass (p

< 0.0001) for all companies, with R^2 ranging from 0.31-0.59 (Supplementary Material S9).

Fish community diversity and composition

Principal coordinate analysis showed weak dissimilarity between the fish community detected by

company A and the local fish community calculated using the Jaccard index (Figure 3). The Raup-Crick

index, however, revealed strong dissimilarities between the local fish community and the fish

community detected by companies B, C, and D (first axis) and companies A and C (second axis) (Figure

3). The molecular community was also dissimilar from the local fish community, especially when using

the Raup-Crick index.

When using read counts obtained after reanalysing the raw sequencing data as a proxy for fish

abundance, we observed strong dissimilarities between eDNA metabarcoding and the local fish

community along the first axis (Horn-Morisita) and second axis (Bray-Curtis) of the principal coordinate

analysis, regardless of the company (Figure 3). As Company A provided the most read counts, it

identified a community similar to the molecular community (Bray-Curtis and Horn-Morisita) (Figure 3).

Discussion

Differences among companies require caution when using eDNA metabarcoding

The companies used different protocols, from field sampling to bioinformatic analysis of DNA

sequences. Without a standard protocol, non-specialist end-users may have no choice but to inventory

biodiversity using eDNA metabarcoding, and the company selected could ultimately influence the

quality and interpretation of the results. The eDNA inventories differed greatly in taxonomic

resolution, species richness, and community composition. Although the fish species in French Guiana

are well documented and have been the subject of several eDNA studies, we were surprised that many

of the taxa reported were identified no finer than to genus, family, order, or even class. Only company

A had satisfactory taxonomic resolution, with nearly all taxa identified to species, whereas the other

three companies identified less than half of the taxa to species. Such inaccurate inventories cannot be

used for conservation biology or biomonitoring, and they cast serious doubt on the ability of certain companies to provide taxonomically usable results.

In addition, the number of species in the eDNA inventories varied by a factor of 10 among companies, which is huge variability for biodiversity inventories. Reanalysing the raw sequencing data using the recently published database (Brosse et al. 2026) decreased differences among companies greatly, particularly for company D, for which the number of detected species increased from 5 to 43. This suggests that differences among companies were due mainly to differences in bioinformatic analyses rather than to those in sampling protocols or laboratory processing. Accessing a database of taxonomically certified barcodes helped decrease differences among the companies. Several eDNA studies have highlighted the importance of comprehensive and curated reference databases for detecting species reliably, as they directly influence detection accuracy, false positive and false negative detection rates, and the applicability of eDNA metabarcoding for monitoring biodiversity (Blackman et al., 2024; Weigand et al., 2019), especially for species-rich communities (Jackman et al., 2021; Keck et al., 2023; Marques et al., 2021).

Environmental managers who want to use eDNA inventories should request a guaranteed minimum degree of quality of the service that will be provided. Before subcontracting with a company to set up programmes to monitor biodiversity and environmental quality, company must be completely transparent about its methods, especially how it processes samples in the laboratory (e.g. DNA extraction method, number of PCR replicates, sequencing depth) and analyses the raw sequencing data (e.g. noise filtering, detection threshold, details of the barcode database used). Being able to access and use comprehensive databases is one of the main conditions for obtaining reliable results while respecting the FAIR principles (findability, accessibility, interoperability, and reusability). Companies should provide this information to customers, along with reviews of eDNA methods and best practices (*e.g.* Bruce et al., 2021; Dickie et al., 2018; Guillera-Arroita et al., 2017; Lahoz-Montfort et al., 2016; Pawlowski et al., 2020). Furthermore, it seems necessary to select a company that will provide the raw sequencing data, along with information about how these data are distributed in biological and technical replicates. Doing so would help to guarantee the quality of the service, as the raw sequencing data can be used to assess the accuracy of the inventory. It would also enable future assessment and updates based on changes in taxonomic knowledge, development of statistical models, and available barcode databases.

Finally, it is necessary to remain cautious when interpreting the results of eDNA inventories. For example, the companies listed 10 species that have never been observed in the Maroni River catchment (2 species) or other large rivers in French Guiana (8 species) (Supplementary Material S6).

A commercial eDNA inventory should not be used to inventory biodiversity without the supervision of
specialists who have taxonomic expertise in the study area.

False negative detections highlight technical limitations of eDNA inventories

This study highlights critical issues related to interpreting eDNA inventories, which illustrates some of
the limitations of this method, as in previous studies (*e.g.* Lacoursière-Roussel & Deiner, 2021; Roussel
et al., 2015). Despite extensive research on eDNA in aquatic environments over the past 10 years, it is
surprising that few studies have compared eDNA inventories of fish to the actual communities in the
field. For example, studies that drained and restocked freshwater ponds with known fish biomasses
found a moderate correlation between fish biomass and the number of reads detected in water
samples per species (Di Muri et al., 2020; Ge et al., 2025). However, this type of approach is not feasible
in running water, for which most studies compared eDNA metabarcoding to traditional sampling
protocols (*e.g.* Gehri et al., 2021; Liu et al., 2024), which are also selective of the species, body size,
and shape of fish (Cowx & Lamarque, 1990; Hamley, 1975).

The present study's assessment of the performance of eDNA metabarcoding provides an
unprecedented comparison of molecular detections to an exhaustive inventory of the local fish
community in a stream. The shallow and accessible aquatic habitats in the Bastien stream allowed for
such inventory (Supplementary Material S2). The electrofishing device was considered effective in
previous studies of similar tropical streams (Pottier et al., 2019, 2022). Moreover, the four
electrofishing passes in the study area exceeded the traditional protocol for this type of inventory
(Carle & Strub, 1978; De Lury, 1951). Thus, we believe that we obtained a comprehensive inventory of
the local fish community.

Among the 32%-92% of the species that the eDNA companies did not detect, three *Hemigrammus* spp.
were captured via electrofishing. Although species-specific DNA markers were available for these
species, only one of them was identified, and only by company B. Similarly, three species in the genera
Hyphessobrycon and *Pristella* were captured via electrofishing but were not detected by eDNA
metabarcoding. That 30%-48% of species captured via electrofishing remained undetected after
reanalysing the raw sequencing data indicates that species were missing due to limitations of eDNA
detection, likely caused by low DNA concentrations in the water and incomplete sampling of the
environment. Indeed, among the species captured via electrofishing, those detected using eDNA
metabarcoding had significantly higher biomasses than those undetected. eDNA is often considered
highly effective for detecting rare and elusive species, but this result tended to demonstrate that eDNA
was relatively ineffective at detecting fish species with low biomass, despite the extensive filtration in
the protocols of four companies.

This raises concerns about the effectiveness of using eDNA metabarcoding alone to inventory fish

biodiversity at the local scale, especially in species-rich tropical stream reaches, with rare species

represented by only one or a few small individuals in traditional inventories (Pottier et al., 2019).

Similar high false negative rates (up to 54%) were recently reported for eDNA metabarcoding

compared to traditional netting methods in Brazilian rivers (dos Anjos Santa Rosa et al., 2025), and it

is unknown whether they would be as high in temperate systems, in which eDNA is expected to persist

for longer (Lamb et al., 2022). Similar studies in contrasting ecological contexts are required to better

understand and quantify the extent to which eDNA metabarcoding provides false negatives, as well as

potential implications for inventories of river-fish biodiversity. These studies would also benefit from

using appropriate sampling designs that involve either an unambiguous inventory or relevant

calibration protocols (Guillera-Arroita et al., 2017; Lahoz-Montfort et al., 2016).

eDNA detections must be interpreted carefully

The relatively high rates of false positive detections (33%-40% among the companies) require cautious

interpretation. We inventoried the local fish community by electrofishing ca. 225 m^2 of shallow stream

habitats. Given the effectiveness of this type of inventory in tropical streams (Pottier *et al.,* 2019,

2022), we believe that few species were missed. Among the false positive eDNA detections

(Supplementary Table S6), previous inventories showed that tributaries of the Maroni River do contain

certain species (*Acestrorhynchus falcatus, Cleithracara* gr. *maronii, Hoplerythrinus vittatus,*

Electrophorus gr. *electricus, Ituglanis nebulosus, Saxatilia saxatilis*) whose presence in the study stream

cannot be ruled out. DNA fragments may have drifted upstream of the electrofishing area (*e.g.* Deiner

et al., 2016; Littlefair et al., 2023), been transported by predators, or come from contamination by

human activities (*e.g.* Guilfoyle & Schultz, 2017; Inoue *et al.*, 2023). Species that usually live in large

rivers (e.g. *Hypomasticus nijsseni* and *Ageneiosus inermis*) may occasionally enter tributaries,

especially during the wet season to feed or spawn, which could explain their eDNA detection.

However, some species detected using eDNA metabarcoding were highly improbable. For example,

Ageneiosus ucayalensis and *Pygocentrus nattereri*, reported by company B, have never been reported

in the Maroni River despite decades of inventories (Le Bail et al., 2026). These false positive detections

likely resulted from errors in bioinformatic analysis or taxonomic assignment by the companies, as they

were not replicated when we reanalysed the raw sequencing data. Indeed, if the reference database

is not comprehensive, the assignment algorithm may select a phylogenetically similar species when

using a minimum similarity threshold of 97% (Jackman et al., 2021). Another species, *Astyanax validus,*

is geographically restricted to the upper reaches of rivers in French Guiana, including the Maroni River

(Planquette et al., 1996), but its DNA was observed in the raw sequencing data of company A.

Reanalysing the companies' raw sequencing data also detected two species that had never been

reported before in the Maroni River catchment (*Gymnotus* aff. *coropinae* and *Eigenmannia* cf.
pavulagem) (Supplementary Table S6). Additional field sampling and verification of the reference
sequences are needed to ensure that these detections were not caused by DNA contamination,
assignment or identification errors. If they can be confirmed, they could be used to update the
distribution of these species in French Guiana. These examples highlight the need for expertise on fish-
species distribution at the regional and continental scales to critically interpret commercial eDNA
inventories and avoid validating those that are biased. It also shows the difficulty in assessing the
reliability of detection without transparent reporting of sequences and reference specimens.

These results support that detection at a given location does not automatically indicate that the species
occupies the sampled habitat. This spatial ambiguity is especially problematic for large-scale
biomonitoring programmes, such as those for the European Union's Water Framework Directive, in
which ecological status is expected to be based on local diversity to address the damage to biodiversity
and the functioning of entire ecosystems. The standard fish-based biotic indices are based on well-
defined habitats and quantitative sampling, which can be used to assess ecological degradation using
robust statistical indices (Pinna et al., 2023). However, high false detection rates, either positive or
negative, may strongly decrease the reliability of eDNA metabarcoding for ecological risk assessment
at the local scale (Yoccoz et al., 2001). Indeed, differences between eDNA inventories and an actual
community could influence the values of biotic indices. The present study demonstrates this concern.
The dissimilarity indices based on presence/absence data (here, Jaccard and Raup-Crick) revealed
discrepancies between eDNA-derived communities and the local community, which indicates
incomplete species coverage. The performances of abundance-based indices (Bray-Curtis and Horn-
Morisita) were not better when eDNA read counts were used as a proxy for species abundance. The
lack of consistency between the commercial eDNA inventories and the actual fish community raises
questions about using eDNA-based fish-community indices for bioindication (*e.g.* Pawlowski et al.,
2018; Pont et al., 2021), at least in species-rich contexts, such as in tropical environments. As a
precaution, future studies should focus on understanding the performances of eDNA-based
community indices before considering using them in large-scale biomonitoring programmes and
ecological risk assessments.

Conclusion and practical recommendations

Although eDNA metabarcoding has potential for inventorying fish communities, this study indicates
that its current application by eDNA companies is limited by high variability in detection rates and
taxonomic accuracy. The high rates of false negatives and positives and the large differences among
companies' results highlight current limitations of eDNA metabarcoding in capturing local biodiversity

in a species-rich context. This is particularly true in running water, in which animal movements and

passive DNA transport further complicate spatial interpretation of the results.

Advancing eDNA metabarcoding for operational monitoring requires developing standard protocols

that include field sampling, laboratory workflows, and bioinformatic analysis (Theroux et al., 2025).

Using relevant sampling designs (e.g. biological and technical replicates, calibration protocols) would

decrease uncertainty in the presence or absence of a taxon, and sharing databases would decrease

uncertainty in taxonomic assignment. The former requires balancing the number of technical and

biological replicates, as well as using blank samples for calibration to estimate false detection rates

accurately. In tropical streams, future research on species-specific eDNA shedding rates and the

environmental factors that influence DNA persistence and transport could help refine eDNA methods.

At present, eDNA users should prioritise companies that are transparent about their methods and

provide raw sequencing data based on biological and technical replicates for independent verification.

Although reanalysing the raw sequencing data with a single reference database substantially reduced

differences among eDNA companies, this alone cannot resolve all sources of error. Reliable eDNA

inventories ultimately depend on reference databases that are curated and taxonomically certified by

specialist taxonomists.

Despite the many studies that used eDNA metabarcoding and those that promoted using it for

biomonitoring, it is surprising that few of them compared eDNA inventories to existing communities.

This study does it for a fish community in a small, species-rich tropical stream, and we recommend

increasing the number of case studies in other ecological contexts. This is a direct and rapid way to

assess the performance of eDNA metabarcoding for future fish inventories and biomonitoring in

running water. Based on our findings, caution is currently warranted when using only eDNA

metabarcoding for regulatory assessments, especially in species-rich environments in which

undetected species could bias ecological interpretations. Finally, managers who use commercial eDNA

inventories to assess fish diversity should consider that interpreting them well requires expert

knowledge of the taxonomy and ecology of the species in the environment studied.

References

Allard L, Toussaint A, Vigouroux R, Brosse S (2015). Length-weight relationships of 58 fish species in French Guiana streams. J. Appl. Ichthyol., 31: 567-570. https://doi.org/10.1111/jai.12712

Andres KJ, Lambert TD, Lodge DM, Andrés J, Jackson JR. (2023). Combining sampling gear to optimally inventory species highlights the efficiency of eDNA metabarcoding. Environmental DNA 5: 146–157.

Baird D, Hajibabaei M (2012). Biomonitoring 2.0: a new paradigm in ecosystem assessment made possible by next-generation DNA sequencing. Molecular Ecology, 21, 2039–2044. https://doi.org/10.1111/j.1365-294X.2012.05519.x

Blabolil P, Jůza T, Draštík V, Knežević-Jarić J, dos Santos R, Griffiths NP, Hänfling B, Peterka J (2021). The true picture of environmental DNA, a case study on harvested fishponds. ARPHA Conference Abstracts 4: e64834. https://doi.org/10.3897/aca.4.e64834

Blackman R, Couton M, Keck F, Kirschner D, Carraro L, Cereghetti E, Perrelet K, Bossart R, Brantschen J, Zhang Y, Altermatt F (2024). Environmental DNA: The next chapter. *Molecular Ecology*, 33, e17355. https://doi.org/10.1111/mec.17355

Bonk E, Hanner R., Bartlett A., Tetreault G. (2024), Environmental DNA Metabarcoding Detects Predators at Higher Rates Than Electrofishing. Environmental DNA, 6: e70019. https://doi.org/10.1002/edn3.70019

Boyer F, Mercier C, Bonin A, Le Bras Y, Taberlet P, Coissac E. (2016). Obitools: A unix-inspired software package for DNA metabarcoding. Molecular Ecology Resources 16 (1): 176-82. https://doi.org/10.1111/1755-0998.12428.

Brantschen J, Altermatt F (2024). Contrasting strengths of eDNA and electrofishing compared to historic records for assessing fish community diversity and composition. Canadian Journal of Fisheries and Aquatic Sciences. 81(2): 178-189. https://doi.org/10.1139/cjfas-2023-0053

Brosse, S, Cuenot, Y, Condachou, C, Coutant, O, Covain, R, Quartarollo, G, Le Reun, S, Vigouroux, R, & Murienne, J (2026). Near complete 12S DNA reference library for the freshwater fish of French Guiana, northern Amazonian region. Available on the NCBI BioProject PRJNA1399107. https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1399107

Bruce, K, Blackman, R, Bourlat, S, Hellström, A, Bakker, J, Bista, I, Bohmann, K, Bouchez, A, Brys, R, Clark, K, Elbrecht, V, Fazi, S, Fonseca, V, Hänfling, B, Leese, F, Mächler, E, Mahon, A, Meissner, K, Panksep, K, & Deiner, K (2021). A practical guide to DNA-based methods for biodiversity assessment. Pensoft. https://doi.org/10.3897/ab.e68634

Bunholi IV, Foster NR, Casey JM (2023). Environmental DNA and RNA in aquatic community ecology: Toward methodological standardization. Environmental DNA, 5, 1133–1147. https://doi.org/10.1002/edn3.476

Carle FL, Strub MR (1978). A new method for estimating population size from removal data. Biometrics. 34: 621-630.

Chao A, Chazdon RL, Colwell RK, Shen T (2006). Abundance-Based Similarity Indices and Their Estimation When There are Unseen Species in Samples, Biometrics, Volume 62, Issue 2, Pages 361–371, https://doi.org/10.1111/j.1541-0420.2005.00489.x

Chase JM, Kraft NJB, Smith KG, Vellend M, Inouye BD (2011). Using null models to disentangle variation in community dissimilarity from variation in α-diversity. Ecosphere 2(2):art24. doi:10.1890/ES10-00117.1

Civade R, Dejean T, Valentini A, Roset N, Raymond J-C, Bonin A, et al. (2016). Spatial Representativeness of Environmental DNA Metabarcoding Signal for Fish Biodiversity Assessment in a Natural Freshwater System. PLoS ONE 11(6): e0157366. doi:10.1371

Condachou C, Coutant O, Fouquet A, Valentini A, Brosse S, Murienne J (2025). Tropical Rainfall and
eDNA Washout Impact Estimations of Amazonian Biodiversity Patterns From Environmental Samples.
Environmental DNA ; 7:e70141

Cordier T., Alonso-Sáez L., Apothéloz-Perret-Gentil L., Aylagas E., Bohan D.A., Bouchez A., Chariton A.,
Creer S., Frühe L., Keck F., Keeley N., Laroche O., Leese F., Pochon X., Stoeck T., Pawlowski J., Lanzén
A. (2020). Ecosystems monitoring powered by environmental genomics: a review of current strategies
with an implementation roadmap. Molecular Ecology, 00, 1–22. https://doi. org/10.1111/mec.15472

Cote D, McClenaghan B, Desforges J, Fahner NA, Hajibabaei M, Chawarski J, Roul S, Singer G, Aubry C,
Geoffroy M. (2023). Comparing eDNA metabarcoding and conventional pelagic netting to inform
biodiversity monitoring in deep ocean environments. Miller-Saunders K, ed. ICES Journal of Marine
Science 80: 2545–2562.

Cowx I.G., Lamarque P. (1990) – Fishing with electricity. Applications in Freshwater Fisheries
Management, Fishing News Books, Oxford, England, Blackwell Scientific Publications, 248 p.

Czeglédi I, Sály P, Specziár A, Preiszner B, Szalóky Z, Maroda Á, Pont D, Meulenbroek P, Valentini A,
Erős T. (2021). Congruency between two traditional and eDNA-based sampling methods in
characterising taxonomic and trait-based structure of fish communities and community-environment
relationships in lentic environment. *Ecological Indicators* 129: 107952.

De Lury DB (1951). On the planning of experiments for estimation of fish populations. Journal of the
Fisheries Research Board of Canada, 8: 281-307.

Deiner K, Altermatt F (2014). Transport distance of invertebrate environmental DNA in a natural river.
PLoS One, 9(2), e88786

Deiner K, Bik HM, Mächler E, et al.(2017). Environmental DNA metabarcoding: Transforming how we
survey animal and plant communities. Mol Ecol. 26: 5872–5895. https://doi.org/10.1111/mec.14350

Deiner K., Fronhofer E., Mächler E., Walser JC., Altermatt F. (2016). Environmental DNA reveals that
rivers are conveyer belts of biodiversity information. Nat Commun 7, 12544.
https://doi.org/10.1038/ncomms12544

Di Muri C, Lawson Handley L, Bean CW, Li J, Peirson G, Sellers GS, Walsh K, Watson HV, Winfield IJ,
Hänfling B (2020). Read counts from environmental DNA (eDNA) metabarcoding reflect fish abundance
and biomass in drained ponds. Metabarcoding and Metagenomics 4: e56959.
https://doi.org/10.3897/mbmg.4.56959

Dickie IA, Boyer S, Buckley HL, et al. (2018). Towards robust and repeatable sampling methods in eDNA-
based studies. *Mol Ecol Resour*. 18: 940–952. https://doi.org/10.1111/1755-0998.12907

Dos Anjos Santa Rosa F., Carvalho CO, Queiroz AKOD, Silva DJF, Barreto SB, Thorbek BLG, Sales JBDL,
Mauvisseau Q, De Boer H, Ready JS. (2025). From nets to molecules: a comparative study of stream
fish diversity recovery using different sampling methods in Eastern Amazonia. BioRxiv preprint,
https://doi.org/10.1101/2025.01.02.631091

Dray S, Dufour A (2007). The ade4 Package: Implementing the Duality Diagram for Ecologists. Journal
of Statistical Software, 22(4), 1-20. https://doi.org/10.18637/jss.v022.i04

Ficetola GF, Miaud C, Pompanon F, Taberlet P (2008). Species detection using environmental DNA from
water samples. Biology letters, 4(4), 423-425.

Fricke R., Eschmeyer W. N., Fong J. D. (2025). Eschmeyer's catalog of fishes: genera/species by
family/subfamily.
http://researcharchive.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp

Ge Q, Piao Y, Li Z, Yang Y, Pan M, Bai Y. (2025). Environmental DNA integrity index is sensitive for species biomass estimation in freshwater. Science of The Total Environment, Volume 966. 178734. https://doi.org/10.1016/j.scitotenv.2025.178734

Gehri R.R., Larson W.A., Gruenthal K., Sard N.M. and Shi Y. (2021), eDNA metabarcoding outperforms traditional fisheries sampling and reveals fine-scale heterogeneity in a temperate freshwater lake. Environmental DNA, 3: 912-929. https://doi.org/10.1002/edn3.197

Guilfoyle MP, Schultz MT (2017). The contribution of double-crested cormorants (Phalacrocorax auritus) to silver carp (*Hypophthalmichthys molitrix*) DNA loads in the Chicago Area Waterway System. Journal of Great Lakes Research, Volume 43, Issue 6, Pages 1181-1185, ISSN 0380-1330, https://doi.org/10.1016/j.jglr.2017.09.008.

Guillera-Arroita G., Lahoz-Monfort J.J., van Rooyen A.R., Weeks A.R., Tingley R. (2017). Dealing with false positive and false negative errors about species occurrence at multiple levels. Methods Ecol Evol 8, 1081–1091. https://doi.org/10.1111/2041-210X.12743.

Hamley JM (1975). Review of gillnet selectivity. Journal of the Fisheries Board of Canada. Vol. 32. https://doi.org/10.1139/f75-233

Harrison JB, Sunday JM, Rogers SM (2019). Predicting the fate of eDNA in the environment and implications for studying biodiversity. Proc R Soc B 286: 20191409.

Hering D, Borja A, Jones JI, Pont D, Boets P, Bouchez A, et al. (2018). Implementation options for DNA-based identification into ecological status assessment under the European Water Framework Directive. Water Res., 138., pp. 192-205.

Hervé A, Domaizon I, Baudoin J-M, Dejean T, Gibert P, Valentini A, Logez M. (2023). Do temperature and population size structure influence the quantity of fish eDNA in water? Front Environ Sci 11: 1193393. doi: 10.3389/fenvs.2023.1193393

Inoue Y, Miyata K, Yamane M, Honda H (2023). Environmental Nucleic Acid Pollution: Characterization of Wastewater Generating False Positives in Molecular Ecological Surveys. ACS ESandT Water 2023 3 (3), 756-764. DOI: 10.1021/acsestwater.2c00542

Jackman JM, Benvenuto C, Coscia I, Oliveira Carvalho C, Ready JS, Boubli JP, Guimarães Sales N (2021). eDNA in a bottleneck: Obstacles to fish metabarcoding studies in megadiverse freshwater systems. Environmental DNA, 3(4), 837-849.

Jannel L-A, Guilhaumon F, Valade P, Chabanet P, Borie G, Grondin H, Jourand P (2024). eDNA Metabarcoding, a Promising Tool for Monitoring Aquatic Biodiversity in the Estuaries of Reunion Island (South-West Indian Ocean). Environmental DNA, 6: e70044. https://doi.org/10.1002/edn3.70044

Keck F, Blackman RC, Bossart R, Brantschen J, Couton M, Hürlemann S, Kirschner D, Locher N, Zhang H, Altermatt F (2022). Meta-analysis shows both congruence and complementarity of DNA and eDNA metabarcoding to traditional methods for biological community assessment. *Molecular Ecology*, 31, 1820–1835. https://doi.org/10.1111/mec.16364

Keck F, Couton M, Altermatt F (2023). Navigating the seven challenges of taxonomic reference databases in metabarcoding analyses. Molecular Ecology Resources, 23(4), 742-755.

Kelly M (2019). Adapting the (fast-moving) world of molecular ecology to the (slow-moving) world of environmental regulation: lessons from the UK diatom metabarcoding exercise. MBMG 3: e39041.

Kelly RP, Lodge DM, Lee KN, Theroux S, Sepulveda AJ, Scholin C A, Weisberg SB (2024). Toward a national eDNA strategy for the United States. Environmental DNA, 6(1), e432.

Lacoursière-Roussel A, Deiner K. (2021). Environmental DNA is not the tool by itself. *J Fish Biol*. 98: 383–386. https://doi.org/10.1111/jfb.14177

Lahoz-Monfort J.J., Guillera-Arroita G., Tingley R. (2016). Statistical approaches to account for false-
positive errors in environmental DNA samples. Mol Ecol Resour 16, 673–685.
https://doi.org/10.1111/1755-0998.12486 .

Lamb, PD, Fonseca, VG, Maxwell, DL, & Nnanatu, CC (2022). Systematic review and meta-analysis:
Water type and temperature affect environmental DNA decay. Molecular Ecology Resources, 22(7),
2494-2505.

Larsson J (2024). eulerr: Area-Proportional Euler and Venn Diagrams with Ellipses. R package version
7.0.2, https://CRAN.R-project.org/package=eulerr.

Le Bail P-Y, Covain R, Jégu M, Fisch-Muller S, Vigouroux R, Keith P (2012). Updated checklist of the
freshwater and estuarine fishes of French Guiana. Cybium, 36(1), 293-319.

Le Bail P-Y, Covain R, Melki F (2026). Guide expert des Poissons d'eau douce de Guyane. Manuel
d'identification. Edition Biotope. 600 p. ISBN 978-2-36662-296-6.

Littlefair JE, Hleap JS, Palace V, Rennie MD, Paterson MJ, Cristescu ME (2023). Freshwater connectivity
transforms spatially integrated signals of biodiversity. Proc. R. Soc. B 290: 20230841.
https://doi.org/10.1098/rspb.2023.0841

Liu Z, H, C, You W, Li S, Wu Y, Liang Y, Chu L, Yan Y, Zhang C (2024). Comparison Between Environmental
DNA Metabarcoding and Traditional Survey Method to Identify Community Composition and Assembly
of Stream Fish. Ecol Evol, 14: e70627. https://doi.org/10.1002/ece3.70627

Marques V, Milhau T, Albouy C, Dejean T, Manel S, Mouillot D, Juhel J B (2021). GAPeDNA: Assessing
and mapping global species gaps in genetic databases for eDNA metabarcoding. Diversity and
Distributions, 27(10), 1880-1892.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet j. Vol. 17(1):10. DOI: 10.14806/ej.17.1.200 http://www-huber.embl.de/users/an-.

Mathieu C, Hermans SM, Lear G, Buckley TR, Lee KC and Buckley HL (2020). A Systematic Review of
Sources of Variability and Uncertainty in eDNA Data for Environmental Monitoring. Front. Ecol. Evol.
8:135.

Mauvisseau Q, Harper LR, Sander M, Hanner RH, Kleyer H, Deiner K (2022). The Multiple States of
Environmental DNA and What Is Known about Their Persistence in Aquatic Environments.
Environmental Science and Technology 2022 56 (9), 5322-5333. DOI: 10.1021/acs.est.1c07638

McKnight EGW, Shafer ABA, Frost PC (2024). Quantifying the effect of water quality on eDNA
degradation using microcosm and bioassay experiments. *Environmental DNA*, 6, e530.
https://doi.org/10.1002/edn3.530

Merkes CM, McCalla SG, Jensen NR, Gaikowski MP, Amberg JJ (2014). Persistence of DNA in Carcasses,
Slime and Avian Feces May Affect Interpretation of Environmental DNA Data. PLoS ONE 9(11):
e113346. doi:10.1371/journal.pone.0113346

Miller D.A., Nichols J.D., McClintock B.T., Grant E.H.C., Bailey L.L., Weir L.A. (2011). Improving
occupancy estimation when two types of observational error occur: non-detection and species
misidentification. Ecology, 92, 1422–1428. https://doi.org/10.1890/10-1396.1

Miya M, Sato Y, Fukunaga T, Sado T, Poulsen J.Y, Sato K, Minamoto T, Yamamoto S, Yamanaka H, Araki
H, Kondoh M, Iwasaki W (2015). MiFish, a set of universal PCR primers for metabarcoding
environmental DNA from fishes: detection of more than 230 subtropical marine species. R. Soc. Open
Sci.2150088http://doi.org/10.1098/rsos.150088

Múrria C, Wangensteen OS, Somma S, Väisänen L, Fortuno P, Arnedo MA, Prat N. (2024). Taxonomic
accuracy and complementarity between bulk and eDNA metabarcoding provides an alternative to

morphology for biological assessment of freshwater macroinvertebrates. Science of the Total
Environment 935. 173243. https://doi.org/10.1016/j.scitotenv.2024.173243

Nevers MB, Przybyla-Kelly K, Shively D, Morris CC, Dickey J, Byappanahalli MN. (2020). Influence of
sediment and stream transport on detecting a source of environmental DNA. Doi H, ed. PLoS ONE 15:
e0244086.

Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P, Stevens M,
Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho G, Chirico M, De Caceres
M, Durand S, Evangelista H, FitzJohn R, Friendly M, Furneaux B, Hannigan G, Hill M, Lahti L, McGlinn D,
Ouellette M, Ribeiro Cunha E, Smith T, Stier A, Ter Braak C, Weedon J (2024). Vegan: Community
Ecology Package. R package version 2.6-8, <https://CRAN.R-project.org/package=vegan>.

Pawlowski J, Apothéloz-Perret-Gentil L, Altermatt F (2020). Environmental DNA: What's behind the
term? Clarifying the terminology and recommendations for its future use in biomonitoring. *Mol Ecol*
2020; 29: 4258–4264. https://doi.org/10.1111/mec.15643

Pawlowski, J., Kelly-Quinn, M., Altermatt, F., & et al. (2018). The future of biotic indices in the
ecogenomic era: Integrating (e)DNA metabarcoding in biological assessment of aquatic ecosystems.
Science of the Total Environment, 637–638, 1295–1310.
https://doi.org/10.1016/j.scitotenv.2018.05.002

Pinna M, Zangaro F, Saccomanno B, Scalone C, Bozzeda F, Fanini L, Specchia V. 2023. An Overview of
Ecological Indicators of Fish to Evaluate the Anthropogenic Pressures in Aquatic Ecosystems: From
Traditional to Innovative DNA-Based Approaches. Water 15: 949.

Planquette P., Keith P., Le Bail P.Y., (1996). Atlas des poissons d'eau douce de Guyane (tome 1) :
Collection du Patrimoine Naturel. IEGB/Muséum national d'Histoire naturelle, Paris, INRA, CSP, Min.
Env., Paris, vol 22, 429 pages.

Pont D, Valentini A, Rocle M, Maire A, Delaigue O, Jean P, Dejean T (2021). The future of fish-based
ecological assessment of European rivers: from traditional EU Water Framework Directive compliant
methods to eDNA metabarcoding-based approaches. Journal of fish biology, 98(2), 354-366.
https://doi.org/10.1111/jfb.14176

Pottier G., Bargier N., Bennevault Y., Vigouroux R., Azam D., Marchand F., Nevoux M., Roussel J.-M.
(2022). Optimising electrofishing settings for shrimp and fish in shallow tropical streams. Fisheries
Research, 256, 106457, https://dx.doi.org/10.1016/j.fishres.2022.106457

Pottier G., Beaumont W.R., Marchand F., Le Bail P.Y., Azam D., Rives J., Vigouroux R., Roussel J.M.
(2019). Electrofishing in streams of low water conductivity but high biodiversity value: challenges,
limits and perspectives. Fisheries Management and Ecology 27: 52-63.
https://doi.org/10.1111/fme.12384

Poyntz-Wright IP, Harrison XA, Pedersen S, Tyler CR. (2024). Effectiveness of eDNA for monitoring
riverine macroinvertebrates, Science of The Total Environment, Vol. 941,
173621. https://doi.org/10.1016/j.scitotenv.2024.173621

Roussel J.-M., Paillisson J.-M., Treguier A., Petit E. (2015). The downside of eDNA as a survey tool in
water bodies. Journal of Applied Ecology, 52 (4), 823-826, https://dx.doi.org/10.1111/1365-
2664.12428

Scriver M, ZAiko A, Pochon X, Von Ammon U. (2023) Harnessing decay rates for coastal marine
biosecurity applications: A review ofenvironmental DNA and RNA fate. Environmental DNA. 5:960–
972. DOI: 10.1002/edn3.405

Shogren AJ, Tank JL, Andruszkiewicz E et al. (2017). Controls on eDNA movement in streams: Transport,
Retention, and Resuspension. *Sci Rep* **7**, 5065. https://doi.org/10.1038/s41598-017-05223-1

Shu L, Ludwig A, Peng Z. (2020). Standards for Methods Utilizing Environmental DNA for Detection of
Fish Species. Genes (Basel). 11;11(3):296. doi: 10.3390/genes11030296. PMID: 32168762; PMCID:
PMC7140814.

Snyder ED, Tank JL, Brandao-Dias PFP, Bibly K, Shogren AJ, Biins AW, Peters B, Curtis EM, Bolster D,
Egan, SP, Lamberti GA. (2023). Environmental DNA (eDNA) removal rates in streams differ by particle
size under varying substrate and light conditions. Science of the Total Environment 903
166469. https://doi.org/10.1016/j.scitotenv.2023.166469

Stein, E. D., Jerde, C. L., Allan, E. A.,Sepulveda, A. J., Abbott, C. L., Baerwald, M. R., Darling, J.,Goodwin,
K. D., Meyer, R. S., Timmers, M. A., & Thielen, P. M.(2023). Critical considerations for communicating
environmental DNA science. Environmental DNA, 00, 1–12.https://doi.org/10.1002/edn3.472

Stoeckle BC, Kuehn R, Geist J (2016). Environmental DNA as a monitoring tool for the endangered
freshwater pearl mussel (*Margaritifera margaritifera* L.): a substitute for classical monitoring
approaches?, Aquatic Conserv: Mar. Freshw. Ecosyst., 26: 1120–1129. doi: 10.1002/aqc.2611.

Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH. (2012). Environmental DNA. Mol Ecol. 21(8):1789-
93. doi: 10.1111/j.1365-294X.2012.05542.x. PMID: 22486819.

Theroux S, Sepulveda A, Abbott CL, Gold Z, Watts AW, Hunter ME, Klymus KE, Hirsch SL, Craine JM,
Jones DN, Brown RJ, Steele JA, Takahashi M, Noble RT, Darling JA (2025) What is eDNA method
standardisation and why do we need it? Metabarcoding and Metagenomics 9: e132076.
https://doi.org/10.3897/mbmg.9.132076

Tréguier A., Paillisson J.-M., Dejean T., Valentini A., Schlaepfer M. A., Roussel J.-M. (2014).
Environmental DNA surveillance for invertebrate species: advantages and technical limitations to
detect invasive crayfish Procambarus clarkii infreshwater ponds. Journal of Applied Ecology, 51 (4),
871-879, https://dx.doi.org/10.1111/1365-2664.12262

Turner C.R., Uy K.L., Everhart R.C. (2015). Fish environmental DNA is more concentrated in aquatic
sediments than surface water. Biological Conservation, 183: 93-102.
http://dx.doi.org/10.1016/j.biocon.2014.11.017

Valentini A, Taberlet P, Miaud C, Civade R, Herder J, Thomsen PF, Bellemain E, Besnard A, Coissac E,
Boyer F, Gaboriaud C, Jean P, Poulet N, Roset N, Copp G, Geniez P, Pont D, Argillier C, Baudoin J, Peroux
T, Crivelli A, Olivier A, Acqueberge M, Le Brun M, Møller P, Willerslev E, Dejean T (2016) Next-
generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. Molecular
Ecology 25 (4): 929-942. https:// doi.org/10.1111/mec.13428

Verdier H, Datry T, Logez M, Konecny-Duprè L, Gauthier M, Lefébure T (2024). Environmental DNA
Particle Size Distribution and Quantity Differ Across Taxa and Organelles. Environmental DNA, 6: e598.
https://doi.org/10.1002/edn3.598

Weigand H, Beermann AJ, Čiampor F, Costa FO, Csabai Z, Duarte S, Geiger MF, Grabowski M, Rimet F,
Rulik B, Strand M, Szucsich N, Weigand AM, Willassen E, Wyler SA, Bouchez A, Borja A, Čiamporová-
Zaťovičová Z, Ferreira S, … Ekrem T (2019). DNA barcode reference libraries for the monitoring of
aquatic biota in Europe: Gap-analysis and recommendations for future work. *Science of The Total
Environment*, 678, 499–524. https://doi.org/10.1016/j.scitotenv.2019.04.247

Xing Y, Gao W, Shen Z, Zhang Y, Bai J, Cai X, Ouyang J and Zhao Y (2022). A Review of Environmental
DNA Field and Laboratory Protocols Applied in Fish Ecology and Environmental Health. Front. Environ.
Sci.10:725360. doi: 10.3389/fenvs.2022.725360

Yates MC, Cristescu ME, Derry AM (2021a). Integrating physiology and environmental dynamics to
operationalize environmental DNA (eDNA) as a means to monitor freshwater macro-organism
abundance. *Molecular Ecology*, 30, 6531–6550. https://doi.org/10.1111/mec.16202

Yates MC, Fraser DJ, Derry AM. (2019). Meta-analysis supports further refinement of eDNA for
monitoring aquatic species-specific abundance in nature. Environmental DNA 1: 5–13.

Yates MC, Glaser D, Post J, Cristescu ME, Fraser DJ, Derry AM. (2021b). The relationship between eDNA
particle concentration and organism abundance in nature is strengthened by allometric scaling. *Mol*
Ecol. 2021; 30: 3068–3082. https://doi.org/10.1111/mec.15543

Yoccoz N.G. (2012). The future of environmental DNA in ecology. Molecular Ecology 21, 2031–2038.
https://doi.org/10.1111/j.1365-294X.2012.05505.x.

Yoccoz N.G., Nichols J.D., Boulinier T. (2001). Monitoring of biological diversity in space and time.
Trends in Ecology and Evolution 16, 446–453. https://doi.org/10.1016/S0169-5347(01)02205-4

Table 1. Number of taxa reported by eDNA companies (A, B, C, D and all combined) at the class, order,
family, genus, and species levels.
807

Level	A	B	C	D	All
Class	0	3	0	1	4
Order	0	0	1	0	1
Family	0	2	2	1	4
Genus	3	11	9	14	19
Species [1]	48	22	19	5	63

808 [1] *Including species for whose taxonomy is currently under debate in the scientific community*
Figure 1. Number of fish species detected in water samples using eDNA metabarcoding based on lists
provided by eDNA companies (top) and after reanalysing their raw sequencing data (bottom).
Histograms show the geographic origin of the species detected: local community (*i.e.* captured on site
immediately after sampling eDNA), in the Maroni River catchment, other large rivers in French Guiana,
or elsewhere in South America. Venn diagrams show the number of species detections shared among
companies.
821

822
823
824
825

Figure 2. False negative detection rate (i.e. percentage of species captured in the local community not
detected by eDNA barcoding) (left) and false positive detection rate (i.e. percentage of species
detected by eDNA barcoding not captured in the local fish community) (right) for the inventories
provided by each eDNA company and all companies combined before (white bars) and after (grey bars)
reanalysing their raw sequencing data.

Figure 3. Community indices for fish captured in the local community (green circles), eDNA
metabarcoding of water samples for each eDNA company (A, B, C, and D), and all companies combined
(molecular community) (blue circles) on the first two axes of principal coordinate analysis (PCoA).
Indices based on the presence/absence of species (Jaccard and Raup-Crick) were calculated using the
commercial eDNA inventories (top). Indices based on abundance per species (Bray-Curtis and Horn-
Morisita) were calculated using read counts as a proxy for abundance after reanalysing raw sequencing
data (bottom).

841