1
2

# Variable performances of commercial eDNA inventories challenge their use for surveying stream fish communities

Roussel JM.[1,§], Quéméré E.[1], Bonnet B.[1-2], Covain R.[3], Dézerald O.[1], Lassalle G.[1], Le Bail PY.[4], Petit E.J.[1],
Pottier G.[2], Quartarollo G.[2], Vigouroux R.[2], Lalagüe H.[2]

[1] UMR DECOD (Ecosystem Dynamics and Sustainability), INRAE, Institut Agro, IFREMER,
Rennes, France
[2] HYDRECO Guyane, Kourou, French Guiana
[3] Museum of Natural History, Geneva, Switzerland.
[4] LPGP (laboratoire de physiologie et génomique des poissons), INRAE, Rennes, France

§ Correspondence: jean-marc.roussel@inrae.fr

**Highlights**

18  -    eDNA metabarcoding of water samples is widely used to detect species in streams
19  -    Detection performances of 4 private companies were compared to an exhaustive fish
20       inventory
21  -    The number of undetected species varies from 32 to 92% depending on the company
22  -    Such discrepancies challenge the use of eDNA for fish-based ecological assessments

**Authors' contributions**
25  JMR, EQ, RV and HL conceived the ideas and designed methodology; GP, RV, HL, GQ collected the
26  data at field; PYLB, RC, GQ identified the species and standardized the names of taxa that the
27  companies provided; BB, GL, JMR, OD and EP analysed and interpreted the data; JMR and HL led the
28  writing of the manuscript. All authors contributed critically to the drafts and gave final approval for
29  publication.

**Acknowledgements**
32  We are indebted to the Direction Generale des Territoires et de la Mer (DGTM Guyane) for financial
33  support and to Nicolas Heitz for insightful discussions on the objectives and implementation of this
34  project.

36  **Data is available on request. No conflict of interest to declare.**

**Abstract**

1. Environmental DNA (eDNA) metabarcoding of water samples is increasingly used to detect fish species in streams. Several studies have concluded that it can outperform traditional inventory methods and recommend using it at large scales for fish-based ecological assessments. However, there is no standard protocol that can guarantee sufficient detection rates and repeatability, despite companies offering an extensive range of analyses.

2. We compared eDNA metabarcoding performed by four companies. Following their guidelines, samples were collected in a small tropical stream in the Maroni River (French Guiana) that hosts a species-rich fish community. We compared their inventories to each other and to a list of species captured during an extensive fish inventory performed immediately after sampling eDNA, as well as to current data on the species' distributions.

3. The number of species detected by eDNA metabarcoding ranged from 5 to 48 among the companies, but these inventories contained many inaccuracies. All companies combined, 63 species were detected, of which 10 (16%) had never been reported in the Maroni River. The extensive inventory identified 50 species in the local fish community, of which 16-46 were not detected by eDNA metabarcoding (i.e. false negative detection rate of 32%-92% among the companies).

4. Reanalysis of raw sequencing data decreased differences among companies greatly, highlighting the importance of using a comprehensive and accurate DNA barcode database to assign species. Dissimilarity indices, calculated to compare the local fish community (based on presence/absence or fish catches) to eDNA detection, revealed large differences regardless of the company.

5. Summary and applications. The large percentage of species not detected by eDNA metabarcoding of water samples could strongly bias fish-diversity inventories in streams that host species-rich communities. This issue is not well documented in the literature, and we recommend that similar studies in the future focus on other stream contexts. The large differences between commercial eDNA inventories and the local fish community challenge the use of eDNA metabarcoding for fish-based ecological assessments of streams. The variable performance of eDNA companies indicates the need for a standard protocol and access to a comprehensive DNA database before beginning large-scale eDNA programmes.

**Keywords**: environmental DNA, mitochondrial 12S primers, metabarcoding, species-rich fish community, detection rate, ecological assessment, French Guiana

**Introduction**

Environmental DNA (eDNA) refers to the genetic material that organisms leave behind in the
environment. It is used to detect the presence of species in environmental samples, which allows
biodiversity to be studied with an unprecedented potential for action. After the pioneering study two
decades ago (Ficetola et al., 2008), eDNA was soon promoted as the ultimate biomonitoring tool
(Baird & Hajibabaei, 2012). Inexpensive, accurate, simple to use in the field, and ethically-based,
eDNA metabarcoding was quickly set to enhance, or even replace, traditional methods based on
capturing or observing organisms, thus helping policy making and future practices better prevent
further biodiversity loss (Taberlet et al., 2012). This has resulted in a large amount of research and
development, with nearly 1000 articles about the use of eDNA for biomonitoring published over the
past decade (Blackman et al., 2024). When no previous knowledge about a community is available,
eDNA metabarcoding has become the most widely used approach, as it can identify and inventory
species using molecular primers that target broad taxonomic groups (Cordier et al., 2020; Deiner et
al., 2017). Many studies have compared eDNA metabarcoding to traditional methods, and they often
reported that the former performed as well as or better overall than the latter (*e.g.* Andres et al.,
2023; Bonk et al., 2024; Czeglédi et al., 2021; Jannel et al., 2024; Keck et al., 2022; Liu et al., 2024).
This has led many scientists and government agencies to recommend large-scale use of eDNA
metabarcoding in monitoring programmes, especially for freshwater ecosystems (*e.g.* Keck et al.,
2022; Kelly et al., 2024; Pawlowski et al., 2018; Stein et al., 2023). However, closer examination of
studies that compared it to traditional methods for freshwater ecosystems revealed that its
performance may be exaggerated due to several unresolved problems, and that key challenges
remain to be addressed before it is routinely used for biomonitoring.

First, eDNA metabarcoding of water samples is usually compared to traditional sampling protocols,
which also inventory only some of the species present. Few field studies have combined eDNA
metabarcoding with exhaustive inventories of communities to explore the former's limitations (but
see Blabolil et al., 2021; Di Muri et al., 2020), and the former's seemingly better performance should
not mask its potential lack of completeness in the field. In general, the absence of a species is difficult
to confirm unless extensive sampling is performed, and eDNA metabarcoding is no exception. Some
organisms release DNA fragments into aquatic environments, depending on their taxonomy,
metabolism, and ontogenetic stage (Múrria et al., 2024; Poyntz-Wright et al., 2024; Tréguier et al.,
2014; Verdier et al., 2024; Yates et al., 2019; Yates et al., 2021a; Yates et al., 2021b). Thus, a species
may be present, but its eDNA concentrations lie below the detection threshold. The eDNA
metabarcoding protocol includes several key steps – water sampling in the field, DNA extraction and
amplification in the laboratory, DNA sequencing, and bioinformatic analysis (Deiner et al., 2017) –

during which the eDNA signal of a species may be lost. The risk of misinterpreting non-detection of
eDNA as true species absence (*i.e.* false negative detection) exists and must be carefully considered
when using eDNA (Lacoursière-Roussel & Deiner, 2021).

Second, several studies have found large differences between species lists identified using eDNA
metabarcoding and traditional methods, especially at the local scale (Brantschen & Altermatt, 2024;
Cote et al., 2023; Keck et al., 2022; Kelly, 2019; Liu et al., 2024). Besides animal movement, water
currents can transport DNA fragments over long distances (Civade et al., 2016; Deiner & Altermatt,
2014; Harrison et al., 2019). Notably, rivers act as "conveyor belts" of biodiversity information
(Deiner et al., 2016; Littlefair et al., 2023) in which the area represented by an eDNA sample depends
on the persistence, shedding, decay, and downstream transport of DNA, all of which vary greatly and
are influenced by complex environmental interactions among discharge, exposure to ultraviolet light,
substrate composition, biofilm and microbial activity, and suspended-sediment characteristics
(Condachou et al., 2025; Deiner & Altermatt, 2014; Harrison et al., 2019; Hervé et al., 2023;
Mauvisseau et al., 2022; McKnight et al., 2024; Nevers et al., 2020; Scriver et al., 2023; Shogren et al.,
2017; Snyder et al., 2023; Turner et al., 2015). Animal waste can also spread eDNA in the
environment (Inoue et al., 2023; Stoeckle et al., 2016), such as when a predator excretes the remains
of its prey outside of the prey's foraging habitats (Guilfoyle & Schultz, 2017; Merkes et al., 2014).
These factors imply that detecting a species' DNA in a geolocated environmental sample does not
necessarily indicate its actual presence at that location, and that eDNA samples may represent a
different spatial scale than those of traditional methods. Unlike methods based on capturing animals,
eDNA introduces a new source of error – false positive detection – whose extent remains largely
unexplored.

Third, the influence of false negatives and positives on sampling design and statistical procedures is
not sufficiently recognised. The development of eDNA metabarcoding as a tool in biodiversity studies
has been driven by technological revolutions, especially the advent of high-throughput sequencing
technologies and the initial realisation that DNA can be extracted from environmental samples
(Ficetola et al., 2008). Despite the early recommendation of Yoccoz (2012) that the eDNA community
should collaborate with quantitative ecologists, relevant sampling designs are rarely adopted, which
may hinder their ability to provide information about the state of and changes in biodiversity (Yoccoz
et al., 2001). A good indicator for eDNA inventories is the ratio of biological replicates (i.e. water
samples) to technical replicates (i.e. Polymerase Chain Reaction; PCR), which is usually low. Most
eDNA protocols prefer PCR replicates to water-sample replicates, but both are needed to assess
potential sources of errors correctly (Guillera-Arroita et al., 2017; Lahoz-Montfort et al., 2016). This
disregard for statistical issues is especially prevalent for aquatic environments, as demonstrated by

recent reviews that did not address them (*e.g.* Bruce et al., 2021; Bunholie et al., 2023; Xing et al.,
2022). Most eDNA developments have focused on increasing sample volumes rather than the
number of biological replicates (Blackman et al., 2024), and using statistical models to quantify errors
remains the exception (15 of the 431 studies reviewed by Mathieu et al. (2020)).

The increasing demand for eDNA-based tools to monitor biodiversity has spurred the worldwide
emergence in recent years of many companies that provide eDNA metabarcoding,
including "turnkey" eDNA solutions for species inventories. They provide clients (i.e. government
agencies, NGOs, and research institutions) with user-friendly sampling kits that allow the clients to
filter water samples themselves in the field. The companies process the samples in the laboratory,
perform the metabarcoding, and provide a list of the taxa detected, usually with little information
about the reliability of their results. Studies have highlighted the lack of minimum justifications for
eDNA methods and sampling design (Blackman et al., 2024; Mathieu et al., 2020), especially the
volume filtered, filter pore size, and numbers of replicates, which vary greatly among water eDNA
studies (Bunholi et al., 2023; Shu et al., 2020; Xing et al., 2022). Moreover, despite the development
of large curated barcode databases (e.g. BOLD, SILVA, MitoFish) over the past few decades, their lack
of comprehensiveness and accuracy remains a major challenge for implementing biomonitoring
programmes correctly (Weigand et al., 2019), especially for species-rich communities (Keck et al.,
2023; Marques et al., 2021). The development of private databases poses accessibility problems for
users, increases the risk of incorrect taxonomic assignment, and could result in many false positive
and/or false negative detections in eDNA inventories (Jackman et al., 2021). Companies vary greatly
in pricing, methodological transparency, and data-sharing policies, which makes it difficult for non-
academic users to select among them. It also poses challenges for comparing eDNA inventories
among companies, especially for long-term biomonitoring programmes.

Following promising scientific developments, eDNA inventories are increasingly expected to be used
to monitor biodiversity, including for regulatory surveillance of aquatic environments (Hering et al.,
2018; Pawlowski et al., 2018; Pont et al., 2021). The number of tendering processes is likely to
increase in the future, but the accuracy with which commercial eDNA inventories are able to describe
fish communities at a given location is unknown. In the present study, we compared four companies
that perform eDNA metabarcoding of water samples to assess the taxonomic composition of fish
communities. The study was performed using a species-rich stream in French Guiana to challenge the
performance of eDNA metabarcoding. After carefully following each company's guidelines for
sampling water and preserving the samples, we sent the samples to each company without disclosing
the study's objective and the exact sampling location. The species lists that each company provided
were compared to (i) an exhaustive inventory of fish species performed at the sampling site

immediately after sampling the water and (ii) current data on fish distribution in French Guiana. False

negative and false positive eDNA detection rates and fish-community composition indices were

calculated and compared among the companies. Finally, raw sequencing data supplied by three of

the companies were reanalysed using a standard bioinformatic pipeline and a recently published

barcode database of fish species in French Guiana. We then assessed how the data analysis may have

influenced differences among the companies.

**Methods**

***Study site and eDNA sampling***

The Bastien stream flows into the Maroni River in French Guiana (Supplementary Material S1). eDNA

was sampled on 18 March 2024 in a 2.5 km tributary of the Bastien stream (5.27048° N, 54.234217°

W; Supplementary Materials S1 and S2). Water temperature (26.1°C), conductivity (32.1 $\mu$S.cm$^{-1}$),

turbidity (2.83 NTU), pH (6.02), and dissolved oxygen (6.52 mg.L$^{-1}$, 79.8% saturation) were recorded

using a multi-parameter probe (Tripod Ponsel Aqualabo). Stream discharge (15 L.s$^{-1}$) was estimated

from water depth and velocity measured across the channel section. The four eDNA companies

selected are internationally known. Following their methodological guidelines, stream water was

pumped in running water to ensure that it was sufficiently mixed, and then filtered in the field using

sampling kits provided by each company. Differences in the number of filters used, pore size,

filtration times, and volumes filtered differed (Supplementary Material S3) revealed diverse sampling

strategies among companies. After sampling, the filters were stored according to each company's

guidelines and placed in a cool box for transport to the laboratory.

eDNA samples were sent to the companies for eDNA metabarcoding, without revealing the objective

of the study or the sampling locations, to avoid influencing the processing of samples. We asked

them only to inventory the fish species detected in the samples. As the companies provided few

details about their laboratory workflows, bioinformatic pipelines, or statistical treatments, we were

not able to compare these aspects directly. Three companies stated that they used one primer pair,

either 12S-Teleo (64 bp) (Valentini et al., 2016) (*Teleo*) or 12S-MiFish-U (163-185 bp) (Miya et al.,

2015) (*MiFish*), while the fourth stated that it used several primer pairs (targeting 12S, COI and 16S

regions), but did not specify which ones. Three companies analysed two biological replicates, while

the fourth analysed three (Supplementary Material S3). The number of technical (PCR) replicates per

sample varied greatly: from 1-12 for the three companies that provided this information

(Supplementary Material S3). Finally, three of the four companies agreed to send us their raw

sequencing data (Illumina FastQ files). The companies have been anonymised to avoid influencing
their commercial activities.

### *Exhaustive inventory of the local fish community*

Immediately after filtering the water, a barrier net (4 mm mesh) was placed across the channel
immediately on the water-pumping point (Supplementary Material S2). The bottom of the net was
buried in the substrate to prevent fish from escaping underneath. Additional barrier nets were
installed upstream to enclose a 100 m long section and prevent fish from escaping during the
inventory. The mean channel width in the section was 225 cm (range: 150-375 cm). The habitat was a
mean of 16.9 cm deep (range: 2-50 cm). The stream bottom consisted mainly of pebbles, sand, and
silt, with a few macrophytes (*Thurnia sphaerocephala*). We inventoried the fish community in the
enclosed section exhaustively using electrofishing. The electrofishing device (described by Pottier et
al. (2019)) was set to deliver 1000 V (DC). The anode was a 350 mm diameter aluminium ring, and
the cathode was a double copper braid (3000 mm long × 10 mm wide × 1 mm thick). Given the
ambient conductivity, these settings ensured effective fish capture while maintaining animal welfare
(Pottier et al. (2019) for details). The fisherman who managed the anode was assisted by two people
equipped with dip nets to capture the fish in the anode's electric field. The entire water surface in
the enclosed section was explored from downstream to upstream, four times in a row, to maximise
fish capture.

After capture, each fish was identified to species using the latest taxonomic knowledge about fish in
French Guiana (Le Bail et al., 2012; Le Bail et al., 2026). Fish were measured (standard length) to the
nearest mm and then gradually released downstream of the sampling section. We applied allometric
length-weight relations based on a historical database of fish captures in French Guiana
(https://eseweb.rennes.inrae.fr/RezoFleuve/) (Allard et al., 2015) to estimate the weight of each
individual, and summed the weights to calculate the biomass of each fish species in the community.
Hereafter, "local fish community" refers to the abundances and biomasses of species captured in the
100 m enclosed section.

### *False positive and false negative eDNA detection rates*

The names of taxa that the companies provided were standardised according to Le Bail et al. (2026)
and Fricke et al. (2025). The number of taxa detected at the class, order, genus, and species levels
were counted. Companies were not penalised for incorrect spelling or if the taxonomy at the species
level was still debated by taxonomists. However, taxa reported only to genus, family, order, or class
were excluded when taxonomic knowledge could have been used to identify them to species. Each
company's list was analysed independently before being combined into a single list of all species

detected by eDNA (hereafter, the "molecular community"). We created Venn diagrams to compare

species lists among companies using the *eulerr* package (Larsson, 2024) of R software (R Core Team,

2025). We also used current knowledge about fish distribution to compare each company's species

list to those in the Maroni River catchment, other large rivers in French Guiana, and elsewhere in

South America (Le Bail et al., 2026).

As no company provided information about taxon detection across biological and technical

replicates, we used the fish captures as unambiguous detections (i.e. no false positive detections)

(Guillera-Arroita et al., 2017; Lahoz-Montfort et al., 2016; Miller et al., 2011). Species in the fish

community that were not detected by eDNA metabarcoding were classified as false negative

detections, whose rates were calculated for each company by dividing the number of undetected

species by the number of species captured in the local fish community. The same calculation was

used for the molecular community. Conversely, species detected using eDNA metabarcoding that

were absent from the local fish community were classified as false positive detections, whose rates

were calculated for each company and for the molecular community by dividing the number of

species detected using eDNA barcoding but not captured in the local fish community by the total

number of species detected using eDNA barcoding.

*Reanalysis of companies' raw sequencing data*

To decrease potential biases related to unequal access to DNA databases among companies and to

highlight differences caused by field sampling and/or laboratory processing, we reanalysed the raw

sequencing data provided by three of the companies using a standard bioinformatic pipeline.

Forward and reverse reads were assembled using the *illuminapairend* function of the *OBITools*

v1.2.11 package (Boyer et al., 2016), and the resulting sequences were trimmed for primers using the

cutadapt v1.8.3 program (Martin, 2011). Strictly identical sequences were clustered using the *obiuniq*

function (dereplication step). Amplicon sequence variants represented by fewer than 10 reads were

removed from the dataset. Additionally, sequence lengths were filtered for each primer set: 40-100

and 140-200 bp for *Teleo* and *MiFish*, respectively. Taxa were assigned using BLASTn (NCBI BLAST+

v2.10.0) compared to a curated reference database (Brosse et al., 2026; NCBI Accession Number:

PX415477 - PX417033), with a minimum similarity threshold of 97% and 98%, an e-value cut-off of

$1e^{-70}$ and $1e^{-20}$, and a minimum alignment coverage of 87% and 80% for *MiFish* and *Teleo*,

respectively. When several equally good hits occurred, taxa were assigned to the lowest common

ancestor. The resulting species lists were then used to recalculate false positive and false negative

detection rates using the procedure described previously.

*Statistical analysis of composition of the local fish community*

Species lists provided by the four companies were used to generate presence/absence data, while
read counts from reanalysing companies' raw sequencing data were used as a proxy for fish
abundance (companies A, B, and D only). We compared eDNA (for each company individually or
combined) to the local fish community using four dissimilarity indices: Jaccard and Raup-Crick for
presence/absence, Bray-Curtis and Horn-Morisita for abundance data. Raup-Crick (Chase et al., 2011)
and Horn-Morisita (Chao et al., 2006), which considers the co-occurrence of rare and unshared
species among communities more than the Jaccard and the Bray-Curtis indices do, are more robust
to differences in species composition and sample size. All dissimilarity indices among communities
were calculated using the *vegan* package (Oksanen et al., 2024) and then visualised via principal
coordinate analysis, with Cailliez-transformation when needed, using the *ade4* package (Dray &
Dufour, 2007).

Finally, we tested whether the abundance and biomass of species in the local fish community
influenced their detection by eDNA barcoding based on the reanalysis of raw sequencing data.
Welch's t-tests (unequal variance, two samples) were used to compare abundance and biomass of
the species detected, or not, by eDNA metabarcoding. Linear regressions between abundance (or
biomass) of species in the local fish community and read counts were calculated, and their
significance was assessed using F-tests. All analyses were performed in R.

**Results**

***Exhaustive inventory of the local fish community***

The 446 fish captured via electrofishing belonged to 6 orders, 22 families, 40 genera, and 50 species
(Supplementary Material S4). The local fish community was dominated by species with low
abundance: 56% of the species were represented by five individuals or fewer, including nine species
represented by only one individual (Supplementary Material S4). Conversely, five species were
represented by more than 20 individuals (Supplementary Material S4 and S5). The total biomass
captured was 2.037 kg for all species combined, of which 54% consisted of individuals weighing less
than 10 g. The total biomass of 11 species exceeded 50 g each, being largest (428 g) for the predator
*Hoplias malabaricus* (8 individuals).

***Assessment of species lists provided by the eDNA companies***

The species list reported by each company (Table 1) contained several taxonomic inaccuracies,
including 4 assignments to class (*Actinotperygii*), 1 to order, 5 to family, and 37 to genus. The
companies reported varying levels of taxonomic resolution: the percentage of taxa identified to

species was highest for company A (94%), followed by companies C (61%), B (59%), and D (24%).

After removing inaccurate detections and standardising the taxonomy, the number of species

detected by eDNA ranged from 5-48 among the four companies (Table 1). The molecular community

contained 63 species. Sequencing depth varied greatly among companies, with company A producing

12 times as many reads as company B (Supplementary Table S3). Company A detected the most

species and had the most read counts in its raw sequencing data. See Supplementary Material S6 for

the complete species list reported by each company.

The percentage of species detected by eDNA metabarcoding that were actually in the local fish

community ranged from 41%-80% among the companies (Figure 1). The molecular community

included 57% of the species actually in the local fish community, along with additional species known

to inhabit the Maroni River catchment (27%), other large rivers in French Guiana (3%), or elsewhere

in South America (13%). Company A detected 30 species that the other companies did not (Figure 1),

most of which were species known to inhabit the Maroni River catchment (Supplementary Material

S6). Company B detected 13 unique species, 7 of which had never been reported in the Maroni River

catchment or other large rivers in French Guiana. Companies A and C shared the most detections in

common: 17 species (Figure 1). Ultimately, only four species were detected by all four eDNA

companies (Figure 1).

Reanalysing the raw sequencing data greatly increased the number of species detected (43-52 per

company) and the percentage of species in the local fish community detected (60%-67%) (Figure 1).

It also decreased the differences among companies and increased the number of species detected by

all companies to 34 (Figure 1). Similarly, detections of species not present in the Maroni River

catchment or other large rivers in French Guiana decreased from 10 to 2 (Figure 1). See

Supplementary Material S6 for the complete list of species detected after reanalysing the raw

sequencing data of each company.

Among the nine species of the local fish community that were not reported by any company or

detected after reanalysis of their sequencing files, only one species (*Brachyhypopomus brevirostris*)

was absent from the Brosse et al. (2026) reference database. Another species, *Eleotris pisonis*, was

reported by one company but was not recovered in the reanalysis, due to its absence from the

Brosse et al. (2026) database.

*False positive and false negative eDNA detection rates*

Comparing eDNA detections to the local fish community revealed that the companies varied greatly

in their ability to detect species (Supplementary Material S7). False negative detection rates ranged

from 32%-92% among the companies, whereas false positive detection rates ranged from 20.0%-

59.1% (i.e. 1-14 species) (Figure 2). False negative detection rates were slightly lower for the
molecular community. Differences in detection rates among the companies decreased when raw
sequencing data were reanalysed, with false negative detection rates ranging from 30.0%-48.0% and
false positive detection rates ranging from 32.7%-39.5% (Figure 2).

Reanalysing the raw sequencing data revealed that species detected by eDNA metabarcoding were
significantly more abundant in the local fish community than undetected species for companies B
and D (Welch's t-test, p = 0.011 and 0.012, respectively) but not for company A (p = 0.489)
(Supplementary Material S8). Similarly, detected species had significantly higher biomass than
undetected species, regardless of the company (Welch's t-test, p < 0.005) (Supplementary Material
S8). We observed no significant relations between read counts of species and species abundance in
the local fish community (F-test, p > 0.065), but read counts were significantly and positively
correlated with species biomass (p < 0.0001) for all companies, with $R^2$ ranging from 0.31-0.59
(Supplementary Material S9).

*Fish community diversity and composition*

Principal coordinate analysis showed weak dissimilarity between the fish community detected by
company A and the local fish community calculated using the Jaccard index (Figure 3). The Raup-Crick
index, however, revealed strong dissimilarities between the local fish community and the fish
community detected by companies B, C, and D (first axis) and companies A and C (second axis)
(Figure 3). The molecular community was also dissimilar from the local fish community, especially
when using the Raup-Crick index.

When using read counts obtained after reanalysing the raw sequencing data as a proxy for fish
abundance, we observed strong dissimilarities between eDNA metabarcoding and the local fish
community along the first axis (Horn-Morisita) and second axis (Bray-Curtis) of the principal
coordinate analysis, regardless of the company (Figure 3). As Company A provided the most read
counts, it identified a community similar to the molecular community (Bray-Curtis and Horn-Morisita)
(Figure 3).

**Discussion**

*Differences among companies require caution when using eDNA metabarcoding*

The companies used different protocols, from field sampling to bioinformatic analysis of DNA
sequences. Without a standard protocol, non-specialist end-users may have no choice but to
inventory biodiversity using eDNA metabarcoding, and the company selected could ultimately

influence the quality and interpretation of the results. The eDNA inventories differed greatly in
taxonomic resolution, species richness, and community composition. Although the fish species in
French Guiana are well documented and have been the subject of several eDNA studies, we were
surprised that many of the taxa reported were identified no finer than to genus, family, order, or
even class. Only company A had satisfactory taxonomic resolution, with nearly all taxa identified to
species, whereas the other three companies identified less than half of the taxa to species. Such
inaccurate inventories cannot be used for conservation biology or biomonitoring, and they cast
serious doubt on the ability of certain companies to provide taxonomically usable results.

In addition, the number of species in the eDNA inventories varied by a factor of 10 among
companies, which is huge variability for biodiversity inventories. Reanalysing the raw sequencing
data using the recently published database (Brosse et al. 2026) decreased differences among
companies greatly, particularly for company D, for which the number of detected species increased
from 5 to 43. This suggests that differences among companies were due mainly to differences in
bioinformatic analyses rather than to those in sampling protocols or laboratory processing. Accessing
a database of taxonomically certified barcodes helped decrease differences among the companies.
Several eDNA studies have highlighted the importance of comprehensive and curated reference
databases for detecting species reliably, as they directly influence detection accuracy, false positive
and false negative detection rates, and the applicability of eDNA metabarcoding for monitoring
biodiversity (Blackman et al., 2024; Weigand et al., 2019), especially for species-rich communities
(Jackman et al., 2021; Keck et al., 2023; Marques et al., 2021).

Environmental managers who want to use eDNA inventories should request a guaranteed minimum
degree of quality of the service that will be provided. Before subcontracting with a company to set
up programmes to monitor biodiversity and environmental quality, company must be completely
transparent about its methods, especially how it processes samples in the laboratory (e.g. DNA
extraction method, number of PCR replicates, sequencing depth) and analyses the raw sequencing
data (e.g. noise filtering, detection threshold, details of the barcode database used). Being able to
access and use comprehensive databases is one of the main conditions for obtaining reliable results
while respecting the FAIR principles (findability, accessibility, interoperability, and reusability).
Companies should provide this information to customers, along with reviews of eDNA methods and
best practices (e.g. Bruce et al., 2021; Dickie et al., 2018; Guillera-Arroita et al., 2017; Lahoz-Montfort
et al., 2016; Pawlowski et al., 2020). Furthermore, it seems necessary to select a company that will
provide the raw sequencing data, along with information about how these data are distributed in
biological and technical replicates. Doing so would help to guarantee the quality of the service, as the
raw sequencing data can be used to assess the accuracy of the inventory. It would also enable future

assessment and updates based on changes in taxonomic knowledge, development of statistical
models, and available barcode databases.

Finally, it is necessary to remain cautious when interpreting the results of eDNA inventories. For
example, the companies listed 10 species that have never been observed in the Maroni River
catchment (2 species) or other large rivers in French Guiana (8 species) (Supplementary Material S6).
A commercial eDNA inventory should not be used to inventory biodiversity without the supervision
of specialists who have taxonomic expertise in the study area.

**_False negative detections highlight technical limitations of eDNA inventories_**

This study highlights critical issues related to interpreting eDNA inventories, which illustrates some of
the limitations of this method, as in previous studies (_e.g._ Lacoursière-Roussel & Deiner, 2021;
Roussel et al., 2015). Despite extensive research on eDNA in aquatic environments over the past 10
years, it is surprising that few studies have compared eDNA inventories of fish to the actual
communities in the field. For example, studies that drained and restocked freshwater ponds with
known fish biomasses found a moderate correlation between fish biomass and the number of reads
detected in water samples per species (Di Muri et al., 2020; Ge et al., 2025). However, this type of
approach is not feasible in running water, for which most studies compared eDNA metabarcoding to
traditional sampling protocols (_e.g._ Gehri et al., 2021; Liu et al., 2024), which are also selective of the
species, body size, and shape of fish (Cowx & Lamarque, 1990; Hamley, 1975).

The present study's assessment of the performance of eDNA metabarcoding provides an
unprecedented comparison of molecular detections to an exhaustive inventory of the local fish
community in a stream. The shallow and accessible aquatic habitats in the Bastien stream allowed for
such inventory (Supplementary Material S2). The electrofishing device was considered effective in
previous studies of similar tropical streams (Pottier et al., 2019, 2022). Moreover, the four
electrofishing passes in the study area exceeded the traditional protocol for this type of inventory
(Carle & Strub, 1978; De Lury, 1951). Thus, we believe that we obtained a comprehensive inventory
of the local fish community.

Among the 32%-92% of the species that the eDNA companies did not detect, three _Hemigrammus_
spp. were captured via electrofishing. Although species-specific DNA markers were available for
these species, only one of them was identified, and only by company B. Similarly, three species in the
genera _Hyphessobrycon_ and _Pristella_ were captured via electrofishing but were not detected by
eDNA metabarcoding. That 30%-48% of species captured via electrofishing remained undetected
after reanalysing the raw sequencing data indicates that species were missing due to limitations of
eDNA detection, likely caused by low DNA concentrations in the water and incomplete sampling of

the environment. Indeed, among the species captured via electrofishing, those detected using eDNA

metabarcoding had significantly higher biomasses than those undetected. eDNA is often considered

highly effective for detecting rare and elusive species, but this result tended to demonstrate that

eDNA was relatively ineffective at detecting fish species with low biomass, despite the extensive

filtration in the protocols of four companies.

This raises concerns about the effectiveness of using eDNA metabarcoding alone to inventory fish

biodiversity at the local scale, especially in species-rich tropical stream reaches, with rare species

represented by only one or a few small individuals in traditional inventories (Pottier et al., 2019).

Similar high false negative rates (up to 54%) were recently reported for eDNA metabarcoding

compared to traditional netting methods in Brazilian rivers (dos Anjos Santa Rosa et al., 2025), and it

is unknown whether they would be as high in temperate systems, in which eDNA is expected to

persist for longer (Lamb et al., 2022). Similar studies in contrasting ecological contexts are required

to better understand and quantify the extent to which eDNA metabarcoding provides false negatives,

as well as potential implications for inventories of river-fish biodiversity. These studies would also

benefit from using appropriate sampling designs that involve either an unambiguous inventory or

relevant calibration protocols (Guillera-Arroita et al., 2017; Lahoz-Montfort et al., 2016).

*eDNA detections must be interpreted carefully*

The relatively high rates of false positive detections (33%-40% among the companies) require

cautious interpretation. We inventoried the local fish community by electrofishing ca. 225 $m^2$ of

shallow stream habitats. Given the effectiveness of this type of inventory in tropical streams (Pottier

*et al.,* 2019, 2022), we believe that few species were missed. Among the false positive eDNA

detections (Supplementary Table S6), previous inventories showed that tributaries of the Maroni

River do contain certain species (*Acestrorhynchus falcatus, Cleithracara* gr. *maronii, Hoplerythrinus*

*vittatus, Electrophorus* gr. *electricus, Ituglanis nebulosus, Saxatilia saxatilis*) whose presence in the

study stream cannot be ruled out. DNA fragments may have drifted upstream of the electrofishing

area (*e.g.* Deiner et al., 2016; Littlefair et al., 2023), been transported by predators, or come from

contamination by human activities (*e.g.* Guilfoyle & Schultz, 2017; Inoue *et al.*, 2023). Species that

usually live in large rivers (e.g. *Hypomasticus nijsseni* and *Ageneiosus inermis*) may occasionally enter

tributaries, especially during the wet season to feed or spawn, which could explain their eDNA

detection.

However, some species detected using eDNA metabarcoding were highly improbable. For example,

*Ageneiosus ucayalensis* and *Pygocentrus nattereri*, reported by company B, have never been

reported in the Maroni River despite decades of inventories (Le Bail et al., 2026). These false positive

detections likely resulted from errors in bioinformatic analysis or taxonomic assignment by the
companies, as they were not replicated when we reanalysed the raw sequencing data. Indeed, if the
reference database is not comprehensive, the assignment algorithm may select a phylogenetically
similar species when using a minimum similarity threshold of 97% (Jackman et al., 2021). Another
species, *Astyanax validus,* is geographically restricted to the upper reaches of rivers in French Guiana,
including the Maroni River (Planquette et al., 1996), but its DNA was observed in the raw sequencing
data of company A. Reanalysing the companies' raw sequencing data also detected two species that
had never been reported before in the Maroni River catchment (*Gymnotus* aff. *coropinae* and
*Eigenmannia* cf. *pavulagem*) (Supplementary Table S6). Additional field sampling and verification of
the reference sequences are needed to ensure that these detections were not caused by DNA
contamination, assignment or identification errors. If they can be confirmed, they could be used to
update the distribution of these species in French Guiana. These examples highlight the need for
expertise on fish-species distribution at the regional and continental scales to critically interpret
commercial eDNA inventories and avoid validating those that are biased. It also shows the difficulty
in assessing the reliability of detection without transparent reporting of sequences and reference
specimens.

These results support that detection at a given location does not automatically indicate that the
species occupies the sampled habitat. This spatial ambiguity is especially problematic for large-scale
biomonitoring programmes, such as those for the European Union's Water Framework Directive, in
which ecological status is expected to be based on local diversity to address the damage to
biodiversity and the functioning of entire ecosystems. The standard fish-based biotic indices are
based on well-defined habitats and quantitative sampling, which can be used to assess ecological
degradation using robust statistical indices (Pinna et al., 2023). However, high false detection rates,
either positive or negative, may strongly decrease the reliability of eDNA metabarcoding for
ecological risk assessment at the local scale (Yoccoz et al., 2001). Indeed, differences between eDNA
inventories and an actual community could influence the values of biotic indices. The present study
demonstrates this concern. The dissimilarity indices based on presence/absence data (here, Jaccard
and Raup-Crick) revealed discrepancies between eDNA-derived communities and the local
community, which indicates incomplete species coverage. The performances of abundance-based
indices (Bray-Curtis and Horn-Morisita) were not better when eDNA read counts were used as a
proxy for species abundance. The lack of consistency between the commercial eDNA inventories and
the actual fish community raises questions about using eDNA-based fish-community indices for
bioindication (*e.g.* Pawlowski et al., 2018; Pont et al., 2021), at least in species-rich contexts, such as
in tropical environments. As a precaution, future studies should focus on understanding the

performances of eDNA-based community indices before considering using them in large-scale
biomonitoring programmes and ecological risk assessments.

*Conclusion and practical recommendations*

Although eDNA metabarcoding has potential for inventorying fish communities, this study indicates
that its current application by eDNA companies is limited by high variability in detection rates and
taxonomic accuracy. The high rates of false negatives and positives and the large differences among
companies' results highlight current limitations of eDNA metabarcoding in capturing local
biodiversity in a species-rich context. This is particularly true in running water, in which animal
movements and passive DNA transport further complicate spatial interpretation of the results.

Advancing eDNA metabarcoding for operational monitoring requires developing standard protocols
that include field sampling, laboratory workflows, and bioinformatic analysis (Theroux et al., 2025).
Using relevant sampling designs (e.g. biological and technical replicates, calibration protocols) would
decrease uncertainty in the presence or absence of a taxon, and sharing databases would decrease
uncertainty in taxonomic assignment. The former requires balancing the number of technical and
biological replicates, as well as using blank samples for calibration to estimate false detection rates
accurately. In tropical streams, future research on species-specific eDNA shedding rates and the
environmental factors that influence DNA persistence and transport could help refine eDNA
methods. At present, eDNA users should prioritise companies that are transparent about their
methods and provide raw sequencing data based on biological and technical replicates for
independent verification. Although reanalysing the raw sequencing data with a single reference
database substantially reduced differences among eDNA companies, this alone cannot resolve all
sources of error. Reliable eDNA inventories ultimately depend on reference databases that are
curated and taxonomically certified by specialist taxonomists.

Despite the many studies that used eDNA metabarcoding and those that promoted using it for
biomonitoring, it is surprising that few of them compared eDNA inventories to existing communities.
This study does it for a fish community in a small, species-rich tropical stream, and we recommend
increasing the number of case studies in other ecological contexts. This is a direct and rapid way to
assess the performance of eDNA metabarcoding for future fish inventories and biomonitoring in
running water. Based on our findings, caution is currently warranted when using only eDNA
metabarcoding for regulatory assessments, especially in species-rich environments in which
undetected species could bias ecological interpretations. Finally, managers who use commercial
eDNA inventories to assess fish diversity should consider that interpreting them well requires expert
knowledge of the taxonomy and ecology of the species in the environment studied.

## References

Allard L, Toussaint A, Vigouroux R, Brosse S (2015). Length-weight relationships of 58 fish species in French Guiana streams. J. Appl. Ichthyol., 31: 567-570. https://doi.org/10.1111/jai.12712

Andres KJ, Lambert TD, Lodge DM, Andrés J, Jackson JR. (2023). Combining sampling gear to optimally inventory species highlights the efficiency of eDNA metabarcoding. Environmental DNA 5: 146–157.

Baird D, Hajibabaei M (2012). Biomonitoring 2.0: a new paradigm in ecosystem assessment made possible by next-generation DNA sequencing. Molecular Ecology, 21, 2039–2044. https://doi.org/10.1111/j.1365-294X.2012.05519.x

Blabolil P, Jůza T, Draštík V, Knežević-Jarić J, dos Santos R, Griffiths NP, Hänfling B, Peterka J (2021). The true picture of environmental DNA, a case study on harvested fishponds. ARPHA Conference Abstracts 4: e64834. https://doi.org/10.3897/aca.4.e64834

Blackman R, Couton M, Keck F, Kirschner D, Carraro L, Cereghetti E, Perrelet K, Bossart R, Brantschen J, Zhang Y, Altermatt F (2024). Environmental DNA: The next chapter. *Molecular Ecology*, 33, e17355. https://doi.org/10.1111/mec.17355

Bonk E, Hanner R., Bartlett A., Tetreault G. (2024), Environmental DNA Metabarcoding Detects Predators at Higher Rates Than Electrofishing. Environmental DNA, 6: e70019. https://doi.org/10.1002/edn3.70019

Boyer F, Mercier C, Bonin A, Le Bras Y, Taberlet P, Coissac E. (2016). Obitools: A unix-inspired software package for DNA metabarcoding. Molecular Ecology Resources 16 (1): 176⊠82. https://doi.org/10.1111/1755-0998.12428.

Brantschen J, Altermatt F (2024). Contrasting strengths of eDNA and electrofishing compared to historic records for assessing fish community diversity and composition. Canadian Journal of Fisheries and Aquatic Sciences. 81(2): 178-189. https://doi.org/10.1139/cjfas-2023-0053

Brosse, S, Cuenot, Y, Condachou, C, Coutant, O, Covain, R, Quartarollo, G, Le Reun, S, Vigouroux, R, & Murienne, J (2026). Near complete 12S DNA reference library for the freshwater fish of French Guiana, northern Amazonian region. Available on the NCBI BioProject PRJNA1399107. https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1399107

Bruce, K, Blackman, R, Bourlat, S, Hellström, A, Bakker, J, Bista, I, Bohmann, K, Bouchez, A, Brys, R, Clark, K, Elbrecht, V, Fazi, S, Fonseca, V, Hänfling, B, Leese, F, Mächler, E, Mahon, A, Meissner, K, Panksep, K, & Deiner, K (2021). A practical guide to DNA-based methods for biodiversity assessment. Pensoft. https://doi.org/10.3897/ab.e68634

Bunholi IV, Foster NR, Casey JM (2023). Environmental DNA and RNA in aquatic community ecology: Toward methodological standardization. Environmental DNA, 5, 1133–1147. https://doi.org/10.1002/edn3.476

Carle FL, Strub MR (1978). A new method for estimating population size from removal data. Biometrics. 34: 621-630.

Chao A, Chazdon RL, Colwell RK, Shen T (2006). Abundance-Based Similarity Indices and Their Estimation When There are Unseen Species in Samples, Biometrics, Volume 62, Issue 2, Pages 361–371, https://doi.org/10.1111/j.1541-0420.2005.00489.x

Chase JM, Kraft NJB, Smith KG, Vellend M, Inouye BD (2011). Using null models to disentangle variation in community dissimilarity from variation in α-diversity. Ecosphere 2(2):art24. doi:10.1890/ES10-00117.1

Civade R, Dejean T, Valentini A, Roset N, Raymond J-C, Bonin A, et al. (2016). Spatial
Representativeness of Environmental DNA Metabarcoding Signal for Fish Biodiversity Assessment in
a Natural Freshwater System. PLoS ONE 11(6): e0157366. doi:10.1371

Condachou C, Coutant O, Fouquet A, Valentini A, Brosse S, Murienne J (2025). Tropical Rainfall and
eDNA Washout Impact Estimations of Amazonian Biodiversity Patterns From Environmental Samples.
Environmental DNA ; 7:e70141

Cordier T., Alonso-Sáez L., Apothéloz-Perret-Gentil L., Aylagas E., Bohan D.A., Bouchez A., Chariton A.,
Creer S., Frühe L., Keck F., Keeley N., Laroche O., Leese F., Pochon X., Stoeck T., Pawlowski J., Lanzén
A. (2020). Ecosystems monitoring powered by environmental genomics: a review of current
strategies with an implementation roadmap. Molecular Ecology, 00, 1–22. https://doi.
org/10.1111/mec.15472

Cote D, McClenaghan B, Desforges J, Fahner NA, Hajibabaei M, Chawarski J, Roul S, Singer G, Aubry C,
Geoffroy M. (2023). Comparing eDNA metabarcoding and conventional pelagic netting to inform
biodiversity monitoring in deep ocean environments. Miller-Saunders K, ed. ICES Journal of Marine
Science 80: 2545–2562.

Cowx I.G., Lamarque P. (1990) – Fishing with electricity. Applications in Freshwater Fisheries
Management, Fishing News Books, Oxford, England, Blackwell Scientific Publications, 248 p.

Czeglédi I, Sály P, Specziár A, Preiszner B, Szalóky Z, Maroda Á, Pont D, Meulenbroek P, Valentini A,
Erős T. (2021). Congruency between two traditional and eDNA-based sampling methods in
characterising taxonomic and trait-based structure of fish communities and community-environment
relationships in lentic environment. *Ecological Indicators* 129: 107952.

De Lury DB (1951). On the planning of experiments for estimation of fish populations. Journal of the
Fisheries Research Board of Canada, 8: 281-307.

Deiner K, Altermatt F (2014). Transport distance of invertebrate environmental DNA in a natural
river. PLoS One, 9(2), e88786

Deiner K, Bik HM, Mächler E, et al.(2017). Environmental DNA metabarcoding: Transforming how we
survey animal and plant communities. Mol Ecol. 26: 5872–5895. https://doi.org/10.1111/mec.14350

Deiner K., Fronhofer E., Mächler E., Walser JC., Altermatt F. (2016). Environmental DNA reveals that
rivers are conveyer belts of biodiversity information. Nat Commun 7, 12544.
https://doi.org/10.1038/ncomms12544

Di Muri C, Lawson Handley L, Bean CW, Li J, Peirson G, Sellers GS, Walsh K, Watson HV, Winfield IJ,
Hänfling B (2020). Read counts from environmental DNA (eDNA) metabarcoding reflect fish
abundance and biomass in drained ponds. Metabarcoding and Metagenomics 4: e56959.
https://doi.org/10.3897/mbmg.4.56959

Dickie IA, Boyer S, Buckley HL, et al. (2018). Towards robust and repeatable sampling methods in
eDNA-based studies. *Mol Ecol Resour*. 18: 940–952. https://doi.org/10.1111/1755-0998.12907

Dos Anjos Santa Rosa F., Carvalho CO, Queiroz AKOD, Silva DJF, Barreto SB, Thorbek BLG, Sales JBDL,
Mauvisseau Q, De Boer H, Ready JS. (2025). From nets to molecules: a comparative study of stream
fish diversity recovery using different sampling methods in Eastern Amazonia. BioRxiv preprint,
https://doi.org/10.1101/2025.01.02.631091

Dray S, Dufour A (2007). The ade4 Package: Implementing the Duality Diagram for Ecologists. Journal
of Statistical Software, 22(4), 1-20. https://doi.org/10.18637/jss.v022.i04

Ficetola GF, Miaud C, Pompanon F, Taberlet P (2008). Species detection using environmental DNA
from water samples. Biology letters, 4(4), 423-425.

Fricke R., Eschmeyer W. N., Fong J. D. (2025). Eschmeyer's catalog of fishes: genera/species by family/subfamily. http://researcharchive.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp

Ge Q, Piao Y, Li Z, Yang Y, Pan M, Bai Y. (2025). Environmental DNA integrity index is sensitive for species biomass estimation in freshwater. Science of The Total Environment, Volume 966. 178734. https://doi.org/10.1016/j.scitotenv.2025.178734

Gehri R.R., Larson W.A., Gruenthal K., Sard N.M. and Shi Y. (2021), eDNA metabarcoding outperforms traditional fisheries sampling and reveals fine-scale heterogeneity in a temperate freshwater lake. Environmental DNA, 3: 912-929. https://doi.org/10.1002/edn3.197

Guilfoyle MP, Schultz MT (2017). The contribution of double-crested cormorants (Phalacrocorax auritus) to silver carp (*Hypophthalmichthys molitrix*) DNA loads in the Chicago Area Waterway System. Journal of Great Lakes Research, Volume 43, Issue 6, Pages 1181-1185, ISSN 0380-1330, https://doi.org/10.1016/j.jglr.2017.09.008.

Guillera-Arroita G., Lahoz-Monfort J.J., van Rooyen A.R., Weeks A.R., Tingley R. (2017). Dealing with false positive and false negative errors about species occurrence at multiple levels. Methods Ecol Evol 8, 1081–1091. https://doi.org/10.1111/2041-210X.12743.

Hamley JM (1975). Review of gillnet selectivity. Journal of the Fisheries Board of Canada. Vol. 32. https://doi.org/10.1139/f75-233

Harrison JB, Sunday JM, Rogers SM (2019). Predicting the fate of eDNA in the environment and implications for studying biodiversity. Proc R Soc B 286: 20191409.

Hering D, Borja A, Jones JI, Pont D, Boets P, Bouchez A, et al. (2018). Implementation options for DNA-based identification into ecological status assessment under the European Water Framework Directive. Water Res., 138., pp. 192-205.

Hervé A, Domaizon I, Baudoin J-M, Dejean T, Gibert P, Valentini A, Logez M. (2023). Do temperature and population size structure influence the quantity of fish eDNA in water? Front Environ Sci 11: 1193393. doi: 10.3389/fenvs.2023.1193393

Inoue Y, Miyata K, Yamane M, Honda H (2023). Environmental Nucleic Acid Pollution: Characterization of Wastewater Generating False Positives in Molecular Ecological Surveys. ACS ESandT Water 2023 3 (3), 756-764. DOI: 10.1021/acsestwater.2c00542

Jackman JM, Benvenuto C, Coscia I, Oliveira Carvalho C, Ready JS, Boubli JP, Guimarães Sales N (2021). eDNA in a bottleneck: Obstacles to fish metabarcoding studies in megadiverse freshwater systems. Environmental DNA, 3(4), 837-849.

Jannel L-A, Guilhaumon F, Valade P, Chabanet P, Borie G, Grondin H, Jourand P (2024). eDNA Metabarcoding, a Promising Tool for Monitoring Aquatic Biodiversity in the Estuaries of Reunion Island (South-West Indian Ocean). Environmental DNA, 6: e70044. https://doi.org/10.1002/edn3.70044

Keck F, Blackman RC, Bossart R, Brantschen J, Couton M, Hürlemann S, Kirschner D, Locher N, Zhang H, Altermatt F (2022). Meta-analysis shows both congruence and complementarity of DNA and eDNA metabarcoding to traditional methods for biological community assessment. *Molecular Ecology*, 31, 1820–1835. https://doi.org/10.1111/mec.16364

Keck F, Couton M, Altermatt F (2023). Navigating the seven challenges of taxonomic reference databases in metabarcoding analyses. Molecular Ecology Resources, 23(4), 742-755.

Kelly M (2019). Adapting the (fast-moving) world of molecular ecology to the (slow-moving) world of environmental regulation: lessons from the UK diatom metabarcoding exercise. MBMG 3: e39041.

Kelly RP, Lodge DM, Lee KN, Theroux S, Sepulveda AJ, Scholin C A, Weisberg SB (2024). Toward a national eDNA strategy for the United States. Environmental DNA, 6(1), e432.

Lacoursière-Roussel A, Deiner K. (2021). Environmental DNA is not the tool by itself. *J Fish Biol*. 98: 383–386. https://doi.org/10.1111/jfb.14177

Lahoz-Monfort J.J., Guillera-Arroita G., Tingley R. (2016). Statistical approaches to account for false-positive errors in environmental DNA samples. Mol Ecol Resour 16, 673–685. https://doi.org/10.1111/1755-0998.12486 .

Lamb, PD, Fonseca, VG, Maxwell, DL, & Nnanatu, CC (2022). Systematic review and meta-analysis: Water type and temperature affect environmental DNA decay. Molecular Ecology Resources, 22(7), 2494-2505.

Larsson J (2024). eulerr: Area-Proportional Euler and Venn Diagrams with Ellipses. R package version 7.0.2, https://CRAN.R-project.org/package=eulerr.

Le Bail P-Y, Covain R, Jégu M, Fisch-Muller S, Vigouroux R, Keith P (2012). Updated checklist of the freshwater and estuarine fishes of French Guiana. Cybium, 36(1), 293-319.

Le Bail P-Y, Covain R, Melki F (2026). Guide expert des Poissons d'eau douce de Guyane. Manuel d'identification. Edition Biotope. 600 p. ISBN 978-2-36662-296-6.

Littlefair JE, Hleap JS, Palace V, Rennie MD, Paterson MJ, Cristescu ME (2023). Freshwater connectivity transforms spatially integrated signals of biodiversity. Proc. R. Soc. B 290: 20230841. https://doi.org/10.1098/rspb.2023.0841

Liu Z, H, C, You W, Li S, Wu Y, Liang Y, Chu L, Yan Y, Zhang C (2024). Comparison Between Environmental DNA Metabarcoding and Traditional Survey Method to Identify Community Composition and Assembly of Stream Fish. Ecol Evol, 14: e70627. https://doi.org/10.1002/ece3.70627

Marques V, Milhau T, Albouy C, Dejean T, Manel S, Mouillot D, Juhel J B (2021). GAPeDNA: Assessing and mapping global species gaps in genetic databases for eDNA metabarcoding. Diversity and Distributions, 27(10), 1880-1892.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnet j. Vol. 17(1):10. DOI: 10.14806/ej.17.1.200 http://www-huber.embl.de/users/an-.

Mathieu C, Hermans SM, Lear G, Buckley TR, Lee KC and Buckley HL (2020). A Systematic Review of Sources of Variability and Uncertainty in eDNA Data for Environmental Monitoring. Front. Ecol. Evol. 8:135.

Mauvisseau Q, Harper LR, Sander M, Hanner RH, Kleyer H, Deiner K (2022). The Multiple States of Environmental DNA and What Is Known about Their Persistence in Aquatic Environments. Environmental Science and Technology 2022 56 (9), 5322-5333. DOI: 10.1021/acs.est.1c07638

McKnight EGW, Shafer ABA, Frost PC (2024). Quantifying the effect of water quality on eDNA degradation using microcosm and bioassay experiments. *Environmental DNA*, 6, e530. https://doi.org/10.1002/edn3.530

Merkes CM, McCalla SG, Jensen NR, Gaikowski MP, Amberg JJ (2014). Persistence of DNA in Carcasses, Slime and Avian Feces May Affect Interpretation of Environmental DNA Data. PLoS ONE 9(11): e113346. doi:10.1371/journal.pone.0113346

Miller D.A., Nichols J.D., McClintock B.T., Grant E.H.C., Bailey L.L., Weir L.A. (2011). Improving occupancy estimation when two types of observational error occur: non-detection and species misidentification. Ecology, 92, 1422–1428. https://doi.org/10.1890/10-1396.1

Miya M, Sato Y, Fukunaga T, Sado T, Poulsen J.Y, Sato K, Minamoto T, Yamamoto S, Yamanaka H,
Araki H, Kondoh M, Iwasaki W (2015). MiFish, a set of universal PCR primers for metabarcoding
environmental DNA from fishes: detection of more than 230 subtropical marine species. R. Soc. Open
Sci.2150088http://doi.org/10.1098/rsos.150088

Múrria C, Wangensteen OS, Somma S, Väisänen L, Fortuno P, Arnedo MA, Prat N. (2024). Taxonomic
accuracy and complementarity between bulk and eDNA metabarcoding provides an alternative to
morphology for biological assessment of freshwater macroinvertebrates. Science of the Total
Environment 935. 173243. https://doi.org/10.1016/j.scitotenv.2024.173243

Nevers MB, Przybyla-Kelly K, Shively D, Morris CC, Dickey J, Byappanahalli MN. (2020). Influence of
sediment and stream transport on detecting a source of environmental DNA. Doi H, ed. PLoS ONE 15:
e0244086.

Oksanen J, Simpson G, Blanchet F, Kindt R, Legendre P, Minchin P, O'Hara R, Solymos P, Stevens M,
Szoecs E, Wagner H, Barbour M, Bedward M, Bolker B, Borcard D, Carvalho G, Chirico M, De Caceres
M, Durand S, Evangelista H, FitzJohn R, Friendly M, Furneaux B, Hannigan G, Hill M, Lahti L, McGlinn
D, Ouellette M, Ribeiro Cunha E, Smith T, Stier A, Ter Braak C, Weedon J (2024). Vegan: Community
Ecology Package. R package version 2.6-8, <https://CRAN.R-project.org/package=vegan>.

Pawlowski J, Apothéloz-Perret-Gentil L, Altermatt F (2020). Environmental DNA: What's behind the
term? Clarifying the terminology and recommendations for its future use in biomonitoring. *Mol Ecol*
2020; 29: 4258–4264. https://doi.org/10.1111/mec.15643

Pawlowski, J., Kelly-Quinn, M., Altermatt, F., & et al. (2018). The future of biotic indices in the
ecogenomic era: Integrating (e)DNA metabarcoding in biological assessment of aquatic ecosystems.
Science      of      the      Total      Environment,      637–638,      1295–1310.
https://doi.org/10.1016/j.scitotenv.2018.05.002

Pinna M, Zangaro F, Saccomanno B, Scalone C, Bozzeda F, Fanini L, Specchia V. 2023. An Overview of
Ecological Indicators of Fish to Evaluate the Anthropogenic Pressures in Aquatic Ecosystems: From
Traditional to Innovative DNA-Based Approaches. Water 15: 949.

Planquette P., Keith P., Le Bail P.Y., (1996). Atlas des poissons d'eau douce de Guyane (tome 1) :
Collection du Patrimoine Naturel. IEGB/Muséum national d'Histoire naturelle, Paris, INRA, CSP, Min.
Env., Paris, vol 22, 429 pages.

Pont D, Valentini A, Rocle M, Maire A, Delaigue O, Jean P, Dejean T (2021). The future of fish-based
ecological assessment of European rivers: from traditional EU Water Framework Directive compliant
methods to eDNA metabarcoding-based approaches. Journal of fish biology, 98(2), 354-366.
https://doi.org/10.1111/jfb.14176

Pottier G., Bargier N., Bennevault Y., Vigouroux R., Azam D., Marchand F., Nevoux M., Roussel J.-M.
(2022). Optimising electrofishing settings for shrimp and fish in shallow tropical streams. Fisheries
Research, 256, 106457, https://dx.doi.org/10.1016/j.fishres.2022.106457

Pottier G., Beaumont W.R., Marchand F., Le Bail P.Y., Azam D., Rives J., Vigouroux R., Roussel J.M.
(2019). Electrofishing in streams of low water conductivity but high biodiversity value: challenges,
limits     and     perspectives.     Fisheries     Management     and     Ecology     27:     52-63.
https://doi.org/10.1111/fme.12384

Poyntz-Wright IP, Harrison XA, Pedersen S, Tyler CR. (2024). Effectiveness of eDNA for monitoring
riverine     macroinvertebrates,     Science     of     The     Total     Environment,     Vol.     941,
173621. https://doi.org/10.1016/j.scitotenv.2024.173621

Roussel J.-M., Paillisson J.-M., Treguier A., Petit E. (2015). The downside of eDNA as a survey tool in
water bodies. Journal of Applied Ecology, 52 (4), 823-826, https://dx.doi.org/10.1111/1365-
2664.12428

Scriver M, ZAiko A, Pochon X, Von Ammon U. (2023) Harnessing decay rates for coastal marine
biosecurity applications: A review ofenvironmental DNA and RNA fate. Environmental DNA. 5:960–
972. DOI: 10.1002/edn3.405

Shogren AJ, Tank JL, Andruszkiewicz E et al. (2017). Controls on eDNA movement in streams:
Transport, Retention, and Resuspension. *Sci Rep* **7**, 5065. https://doi.org/10.1038/s41598-017-
05223-1

Shu L, Ludwig A, Peng Z. (2020). Standards for Methods Utilizing Environmental DNA for Detection of
Fish Species. Genes (Basel). 11;11(3):296. doi: 10.3390/genes11030296. PMID: 32168762; PMCID:
PMC7140814.

Snyder ED, Tank JL, Brandao-Dias PFP, Bibly K, Shogren AJ, Biins AW, Peters B, Curtis EM, Bolster D,
Egan, SP, Lamberti GA. (2023). Environmental DNA (eDNA) removal rates in streams differ by particle
size under varying substrate and light conditions. Science of the Total Environment 903
166469. https://doi.org/10.1016/j.scitotenv.2023.166469

Stein, E. D., Jerde, C. L., Allan, E. A.,Sepulveda, A. J., Abbott, C. L., Baerwald, M. R., Darling,
J.,Goodwin, K. D., Meyer, R. S., Timmers, M. A., & Thielen, P. M.(2023). Critical considerations for
communicating environmental DNA science. Environmental DNA, 00, 1–
12.https://doi.org/10.1002/edn3.472

Stoeckle BC, Kuehn R, Geist J (2016). Environmental DNA as a monitoring tool for the endangered
freshwater pearl mussel (*Margaritifera margaritifera* L.): a substitute for classical monitoring
approaches?, Aquatic Conserv: Mar. Freshw. Ecosyst., 26: 1120–1129. doi: 10.1002/aqc.2611.

Taberlet P, Coissac E, Hajibabaei M, Rieseberg LH. (2012). Environmental DNA. Mol Ecol. 21(8):1789-
93. doi: 10.1111/j.1365-294X.2012.05542.x. PMID: 22486819.

Theroux S, Sepulveda A, Abbott CL, Gold Z, Watts AW, Hunter ME, Klymus KE, Hirsch SL, Craine JM,
Jones DN, Brown RJ, Steele JA, Takahashi M, Noble RT, Darling JA (2025) What is eDNA method
standardisation and why do we need it? Metabarcoding and Metagenomics 9: e132076.
https://doi.org/10.3897/mbmg.9.132076

Tréguier A., Paillisson J.-M., Dejean T., Valentini A., Schlaepfer M. A., Roussel J.-M. (2014).
Environmental DNA surveillance for invertebrate species: advantages and technical limitations to
detect invasive crayfish Procambarus clarkii infreshwater ponds. Journal of Applied Ecology, 51 (4),
871-879, https://dx.doi.org/10.1111/1365-2664.12262

Turner C.R., Uy K.L., Everhart R.C. (2015). Fish environmental DNA is more concentrated in aquatic
sediments than surface water. Biological Conservation, 183: 93-102.
http://dx.doi.org/10.1016/j.biocon.2014.11.017

Valentini A, Taberlet P, Miaud C, Civade R, Herder J, Thomsen PF, Bellemain E, Besnard A, Coissac E,
Boyer F, Gaboriaud C, Jean P, Poulet N, Roset N, Copp G, Geniez P, Pont D, Argillier C, Baudoin J,
Peroux T, Crivelli A, Olivier A, Acqueberge M, Le Brun M, Møller P, Willerslev E, Dejean T (2016) Next-
generation monitoring of aquatic biodiversity using environmental DNA metabarcoding. Molecular
Ecology 25 (4): 929□942. https:// doi.org/10.1111/mec.13428

Verdier H, Datry T, Logez M, Konecny-Duprè L, Gauthier M, Lefébure T (2024). Environmental DNA
Particle Size Distribution and Quantity Differ Across Taxa and Organelles. Environmental DNA, 6:
e598. https://doi.org/10.1002/edn3.598

Weigand H, Beermann AJ, Čiampor F, Costa FO, Csabai Z, Duarte S, Geiger MF, Grabowski M, Rimet F,
Rulik B, Strand M, Szucsich N, Weigand AM, Willassen E, Wyler SA, Bouchez A, Borja A, Čiamporová-
Zaťovičová Z, Ferreira S, … Ekrem T (2019). DNA barcode reference libraries for the monitoring of
aquatic biota in Europe: Gap-analysis and recommendations for future work. *Science of The Total
Environment*, 678, 499–524. https://doi.org/10.1016/j.scitotenv.2019.04.247

Xing Y, Gao W, Shen Z, Zhang Y, Bai J, Cai X, Ouyang J and Zhao Y (2022). A Review of Environmental
DNA Field and Laboratory Protocols Applied in Fish Ecology and Environmental Health. Front.
Environ. Sci.10:725360. doi: 10.3389/fenvs.2022.725360

Yates MC, Cristescu ME, Derry AM (2021a). Integrating physiology and environmental dynamics to
operationalize environmental DNA (eDNA) as a means to monitor freshwater macro-organism
abundance. *Molecular Ecology*, 30, 6531–6550. https://doi.org/10.1111/mec.16202

Yates MC, Fraser DJ, Derry AM. (2019). Meta-analysis supports further refinement of eDNA for
monitoring aquatic species-specific abundance in nature. Environmental DNA 1: 5–13.

Yates MC, Glaser D, Post J, Cristescu ME, Fraser DJ, Derry AM. (2021b). The relationship between
eDNA particle concentration and organism abundance in nature is strengthened by allometric
scaling. *Mol Ecol*. 2021; 30: 3068–3082. https://doi.org/10.1111/mec.15543

Yoccoz N.G. (2012). The future of environmental DNA in ecology. Molecular Ecology 21, 2031–2038.
https://doi.org/10.1111/j.1365-294X.2012.05505.x.

Yoccoz N.G., Nichols J.D., Boulinier T. (2001). Monitoring of biological diversity in space and time.
Trends in Ecology and Evolution 16, 446–453. https://doi.org/10.1016/S0169-5347(01)02205-4

817
818

819 **Table 1.** Number of taxa reported by eDNA companies (A, B, C, D and all combined) at the class,
820 order, family, genus, and species levels.
821

| Level | A | B | C | D | All |
|---|---|---|---|---|---|
| Class | 0 | 3 | 0 | 1 | 4 |
| Order | 0 | 0 | 1 | 0 | 1 |
| Family | 0 | 2 | 2 | 1 | 4 |
| Genus | 3 | 11 | 9 | 14 | 19 |
| Species [1] | 48 | 22 | 19 | 5 | 63 |

822 [1] *Including species for whose taxonomy is currently under debate in the scientific community*
**Figure 1.** Number of fish species detected in water samples using eDNA metabarcoding based on lists
provided by eDNA companies (top) and after reanalysing their raw sequencing data (bottom).
Histograms show the geographic origin of the species detected: local community (*i.e.* captured on
site immediately after sampling eDNA), in the Maroni River catchment, other large rivers in French
Guiana, or elsewhere in South America. Venn diagrams show the number of species detections
shared among companies.
835

**Figure 2.** False negative detection rate (i.e. percentage of species captured in the local community
not detected by eDNA barcoding) (left) and false positive detection rate (i.e. percentage of species
detected by eDNA barcoding not captured in the local fish community) (right) for the inventories
provided by each eDNA company and all companies combined before (white bars) and after (grey
bars) reanalysing their raw sequencing data.

**Figure 3.** Community indices for fish captured in the local community (green circles), eDNA
metabarcoding of water samples for each eDNA company (A, B, C, and D), and all companies
combined (molecular community) (blue circles) on the first two axes of principal coordinate analysis
(PCoA). Indices based on the presence/absence of species (Jaccard and Raup-Crick) were calculated
using the commercial eDNA inventories (top). Indices based on abundance per species (Bray-Curtis
and Horn-Morisita) were calculated using read counts as a proxy for abundance after reanalysing raw
sequencing data (bottom).

856