Title:

Exposure to (*Z*)-3-hexenol primes tobacco plants for faster and stronger defense without

negatively affecting their ability to grow and reproduce

Running Title: (*Z*)-3-hexenol primes tobacco defense without growth or reproductive costs

Authors:

Bipana Paudel Timilsena[1*], Irmgard Seidl-Adams[1], Sarah Refi Hind[2], Gary Felton[1], James H.

Tumlinson[†1]

[†] Deceased 9 February 2022

[1]Department of Entomology, The Pennsylvania State University, PA, USA

[2]Department of Crop Science, University of Illinois, IL, USA

[*]**Author for correspondence:** Bipana Paudel Timilsena, Email: bpp5121@psu.edu

Abstract

Plants exposed to volatile signals from herbivore-infested neighbors can activate faster and

stronger defenses against subsequent herbivore attack, a phenomenon called defense priming.

However, the specific volatile components responsible for activating defense priming remain

unclear. Here, we examined the role of green leaf volatiles (GLV) by silencing their biosynthesis

using virus-induced gene silencing technique. Exposure to full blend of herbivore-induced plant

volatiles (HIPV) primed receiver plants for enhanced production of all 5 groups of HIPV (GLV,

monoterpenes, sesquiterpenes, aldoximes, and indole). When GLV production was silenced in

emitter plants, receiver plants were no longer primed for terpene production. However, exposure

to (*Z*)-3-hexenol (Z3HOL) alone primed receiver plants for terpene production. These results

suggest that GLV are necessary, and Z3HOL alone is sufficient, to prime terpene production in

receiver plants. Consistent with enhanced resistance, *Manduca sexta* larvae feeding on Z3HOL-

or HIPV-primed plants consumed less leaf tissue and exhibited reduced growth compared with

controls. Importantly, priming did not impose fitness costs, as Z3HOL-exposed plants showed

normal growth but produced more seed capsules and seeds than control plants. Together, these

findings suggest that Z3HOL alone is sufficient to prime plants for better defense without

compromising their ability to grow and reproduce.

Key words: Plant defense priming, herbivore-induced plant volatiles, green leaf volatiles (GLV),

Manduca sexta, plant-insect interactions, fitness cost, virus-induced gene silencing (VIGS)

1. Introduction:

Numerous studies have demonstrated that when plants are exposed to volatile signals from herbivore-infested neighbors, they tailor their defense reactions to incoming attack. After perceiving these volatile signals, some plants immediately exhibit direct induction of defense responses, such as upregulation of defense related genes and/or production of defense metabolites, even in the absence of herbivore attack (Farag & Paré 2002; Kost & Heil 2006). While others do not show immediate changes in defense status but respond more rapidly and more strongly following subsequent attack by herbivores, a phenomenon known as defense priming (Engelberth *et al*., 2004; Ton *et al*. 2007; Frost *et al*., 2008). The first evidence that plants can respond to volatile signals emitted by damaged neighbors was reported in the 1980s (Rhoades 1983; Baldwin & Schultz 1983). Since then, numerous studies have reported similar observations in both natural and laboratory settings (Heil & Karban 2010; Karban *et al*., 2014). However, after decades of research, it is still unknown what the distinguishing characteristics of the active compounds responsible for defense priming. Moreover, while most of the priming studies have focused on upregulation of defense genes and metabolites in primed plants, relatively little is known about how defense priming affects the performance of attacking herbivores and fitness of plants themselves.

Several different volatile compounds have been reported to induce defense priming in plants. The phytohormone ethylene was identified as the first gaseous signaling molecule in the early 1900s (Sasidharan 2022). Other volatile phytohormones, such as methyl jasmonate (meJA) and methyl salicylate (meSA), were also reported to induce defense responses in non-stressed neighboring plants (Farmer & Ryan 1990; Karban *et al*., 2000; Zavala & Baldwin 2004; Baldwin *et al*., 2006; Kessler *et al*., 2006; Tamogami *et al*., 2008). Yao *et al.* (2011) proposed

involvement of additional signaling compounds beside volatile hormones (MeJA and MeSA) in

plant defense priming. Accordingly, other candidate compounds for defense signaling include

Green Leaf Volatiles (GLV), terpenes and aromatic compounds.

GLV (carbon-6 aldehydes, alcohols, and their esters) are among the most studied

signaling molecules involved in defense priming (Cofer *et al*., 2018). Plants release GLV

immediately after wounding or herbivory. Exposure to GLV activates several defense related

genes and downstream metabolites (Zeringue 1992; Bate & Rothstein 1998; Engelberth *et al*.,

2013). In addition to GLV, homoterpenes (*(E)*-3,8-dimethyl-1,4,7-nonatriene (DMNT) and

(*E,E*)-4,8,12-trimethyl tridecane-1,3,7,11-tetraene (TMTT)) and monoterpenes (myrcene alone

or blend of ocimenes consisting *(E)*-β-ocimene, *(Z)*-β-ocimene and allo-ocimene) have also been

shown to induce defense related genes and meJA accumulation in exposed plants (Arimura *et al*.

2000; Godard *et al*., 2008; Meents *et al*. 2019). Similarly, exposure to physiologically relevant

concentrations of indole (50 ng hr^{-1}) also primes neighboring plants for enhanced production of

JA, JA isoleucine (JA-Ile), abscisic acid (ABA) and plant volatiles following subsequent

herbivore damage (Erb *et al.* 2015).

Yet most studies on priming, to date, have been performed using high doses of synthetic

volatiles in airtight containers (Bate & Rothstein 1998; Engelberth *et al*. 2004). This high dose

volatile exposure does not realistically represent what occurs in nature (von Mérey *et al*. 2011)

and it also damages the plant tissue. The airtight container itself imposes physiological stress on

plants. To address these limitations, a few studies attempted to mimic natural exposure by using

genetically manipulated plants or slow-release dispensers but reported negative or unexpected

results. For example, under field conditions, GLV-exposed plants experienced increased

herbivore attraction and damage compared to unexposed control plants (von Mérey *et al*. 2011).

Similarly, *N. attenuata* plants genetically impaired to emit GLV or terpenes failed to prime

neighboring plants for defense gene expression and defense metabolite production following a

single wounding event (Paschold *et al.*, 2006). In contrast, volatiles from detached leaves of

Artemisia tridentata, containing both GLV and terpenes, primed *N. attenuata* for enhanced

proteinase activity only under continuous damage by *M. sexta* larvae (Kessler *et al.* 2006). These

contrasting results suggest that, beyond volatile composition, the frequency and intensity of

herbivore damage play a critical role in shaping plant defense and priming responses. In tobacco,

continuous damage on plants built up JA production which is necessary to induce herbivore

resistance mediated by both HIPV and defensive metabolites, such as nicotine and trypsin

proteinase inhibitor (TPI) (Ziegler *et al.*, 2001; Zavala *et al.*, 2004; Zavala & Baldwin 2004;

Halitschke *et al.*, 2004). By contrast, a single wounding event elicits a quick burst of JA

accumulation that rapidly returns to basal levels and is insufficient to confer herbivore resistance

(Ziegler *et al.*, 2001).

In a previous study, we found that the complete blend of HIPV primes neighboring *N.*

benthamiana plants for enhanced JA and volatile production following subsequent herbivore

challenge (Paudel Timilsena *et al.*, 2020). Here we examined the role of GLV by silencing their

biosynthetic pathway using virus-induced gene silencing (VIGS). We also exposed plants to

individual GLV at physiologically relevant concentrations by using slow-release dispensers. In

addition to quantifying defense metabolites, we also evaluated whether defense priming affects

herbivore performance on primed plants, as well as the growth and reproduction of the plants

themselves.

2. Materials and Methods:

2.1 Plant materials and growing condition:

Wild type (WT) tobacco plants, *N. benthamiana* (Solanaceae), were grown from seeds in individual pots (7.5 cm height, 9 cm diameter) filled with commercial potting soil (Pro-mix® PGX, Quakertown, PA) supplemented with Osmocote slow-release fertilizer (8-45-14 N-P-K, Scotts Company, Marysville, OH, USA). WT plants were grown in a greenhouse at 26 ± 2 °C under a 16 h light/ 8 h dark photoperiod. Plants used for experiments were 3-4 weeks old and had 4 fully developed leaves.

All VIGS plants were grown in a growth chamber (20 ± 2 °C and 16 h light/ 8 h dark) post VIGS inoculation. VIGS technique was used to silence the NbLOX2 gene as described in section below. Two days before the experiment, VIGS plants were transferred to and kept under the greenhouse conditions described above (26 ± 2°C, 16 h light/ 8 h dark). Transferring VIGS plants to the greenhouse condition with high temperature (26 ± 2°C) and high light intensity did not change the silenced phenotype (Nethra *et al*., 2006).

2.2 Insect and regurgitant:

The Schilder lab at Penn State University, PA, generously provided *M. sexta* eggs. *M. sexta* neonates were reared on artificial diet under a 16 h light/ 8 h dark photoperiod at 25 ± 2°C. Oral secretions were collected from fourth- to fifth-instar *M. sexta* larvae feeding on *N. benthamiana* leaves for at least 24 h using previously described methods (Paudel Timilsena *et al*. 2020). Briefly, regurgitant was collected by squeezing larvae's head and placing a pipette on their mouthpart. The collected regurgitant was immediately diluted with deionized water (1:1 v/v) and stored at -80 °C until use.

2.3 Chemicals

(*Z*)-3-hexenal (50%, SAFC, USA), (*Z*)-3-hexenol (>98%, Aldrich), (*Z*)-3-hexenyl acetate (>98%, Aldrich) were used to make slow-releasing dispensers. Hexane (>98.5%, J.T.Baker, USA) and dichloromethane (99.9%, OmniSolv, Germany) were used to elute volatiles from Super-Q (Alltech, USA) filters, and to wash Super-Q filters and glassware. Nonyl acetate (>97%, Aldrich, USA) was used as an internal standard to quantify volatiles. (*E*)-2-hexenal (98%, Bedoukian Research Inc., USA), linalool (97%, Aldrich), limonene (97%, Aldrich), (1, 8-) cineol (99%, Sigma-Aldrich), myrcene (90%, Aldrich), α-pinene (99%, Aldrich), β-pinene (99%, Aldrich), (*E*)-β-farnesene (99%, mixtures of isomers, Sigma-Aldrich) were used as reference standards for compounds identification based on mass spectra and retention time comparison using GC-MS and GC-FID, respectively.

2.4 Manipulation of HIPV release by emitter plants:

We previously showed that exposure to the whole blend of HIPV primed neighboring plants for enhanced defense responses after subsequent challenge with herbivore regurgitant. To investigate the role of GLVs in priming, we generated emitter plants whose HIPV profile was lacking GLVs by silencing the lipoxygenase (LOX) gene in the hydroperoxide lyase pathway (HPL) branch of the oxylipin pathway (Bate & Rothstein 1998). We used virus-induced gene silencing (VIGS) to generate these GLV-compromised plants. These plants were exclusively used as emitter plants to produce defined volatile profiles at physiologically relevant concentrations. Therefore, any possible side effects of the VIGS would not affect their experimental role.

GLV are carbon-6 aldehyde, alcohol and their esters produced by the HPL branch of the

oxylipin pathway. We silenced the homolog of the *N. attenuata* NaLOX2 gene in *N.*

benthamiana plants (accession no. AY254348; Halitschke & Baldwin, 2003). Following

herbivore damage, LOX2-silenced plants produced greatly reduced amounts of GLV,

comparable to basal levels (**Fig. 2B**). Therefore, we used herbivore damaged LOX2 silenced

plants as emitter plants and wild type (WT) as receiver plants. After 48 hours of exposure to

volatiles, we measured the defense metabolites; volatiles and JA, in the WT receiver plants.

2.5 Plasmid construction for VIGS*:*

Constructs used for this study include TRV::LOX2 (LOX2, lipoxygenase gene from *N.*

benthamiana), TRV::GFP (GFP, green fluorescent protein, a non-plant gene obtained from

jellyfish *Aequorea victoria* (Haseloff *et al*., 1997)) and TRV::PDS (PDS, phytoene desaturase

from *N. benthamiana*). TRV::GFP was used as a VIGS control and TRV::PDS as a visual marker

to confirm VIGS efficiency in all our VIGS experiments (Supplementary Fig. **S1**).

For cloning individual genes into pTRV2-LIC, the gene of interest (GOI) was amplified

using the primers: 5' CGACGACAAGACCCT (LIC vector adaptor) – GOI specific sequence-3'

and 5'-GAGGAGAAGAGCCCT (LIC vector adaptor) – GOI specific sequence - 3' (Table **S1,**

Dong *et al*., 2007). The insert size of each target gene was 300 bp in the TRV2-LIC vector. The

PCR product of GOI was purified with a PCR clean-up kit. The 1000 ng of TRV2-LIC (pYY13)

vector plasmid was digested by PstI-HF restriction enzyme in a 50 μL reaction, incubated for 1 h

at 37 ˚C and purified with PCR-cleanup kit. A total of 50 ng of purified PCR product was treated

with T4 DNA Polymerase in 1X reaction buffer containing 5 mM dATP and 5 mM Dithiothreitol

(DTT) for 30 min at 22 ˚C followed by 20 min of inactivation of T4 DNA polymerase at 70 ˚C.

A total of 500 ng of digested and purified TRV2-LIC (pYY13) plasmid was treated with T4

DNA polymerase and dTTP instead of dATP. Then the PCR product and TRV2-LIC (pYY13)

vector were mixed in 1:1 ratio and incubated at 65 ˚C for 2 min and then 22 ˚C for 10 min. The

mixture was then transformed into *Escherichia coli* strains TOP10 according to One Shot®

Top10 Competent Cells protocol (Thermo Fisher Scientific). A total of 100 μL transformation

reaction was spread onto LB plates containing 50 μg mL^{-1} of Kanamycin and incubated at 37 ˚C

for up to 48 h. Transformants were selected by PCR and confirmed by running agarose gel

electrophoresis. The plasmids from the confirmed transformants were isolated using Mini-Prep

kit and were introduced into competent *Agrobacterium tumefaciens* strains GV2260 using the

electroporation method by using a MicroPulser Electroporation according to the manufacturer's

(Bio-Rad) instructions.

2.6 Virus inoculation:

Agrobacterium-mediated inoculation was conducted as described previously (Senthil-

Kumar & Mysore, 2014). *Agrobacterium tumefaciens* strains GV2260 cells transformed with

individual constructs were grown in LB (Luria-Bertani) broth, harvested and resuspended in

MES (2-(N-morpholino) ethane-sulfonic acid), and incubation at room temperature for 3 h.

Suspensions of Agrobacterium carrying pTRV1 and pTRV2 carrying GOI were mixed in 1:1

(v/v) ratio and infiltrated into leaves using a needleless syringe. Inoculated plants were grown in

a growth chamber at 20 ˚C for 4 weeks.

2.7 Volatile dispenser:

Plants were exposed to individual GLV for 48 hours using a slow-release dispenser

designed to release physiologically relevant amounts for 5-7 days. Briefly, synthetic compound

of interest was mixed with lanolin and 25 μL of the mixture was placed into a GC-insert. The

GC-insert was immediately placed into the GC-vial and capped. A 1 μL, 64 mm micropipette

was then inserted into the GC-vial. The emission rates were adjusted to physiologically relevant

levels. For example, Z3HOL was released at 500 ng h^{-1} (mean, n = 6), which closely mimicked

the emission rate observed in *M. sexta* damaged plant at 48 h post infestation (Supplementary

Fig. **S2**). These calibrated dispensers were used as a source of volatile throughout the priming

experiments.

2.8 Treatments on receiver plants, plant volatile collection and analysis:

To examine the role of GLV in plant defense priming, WT receiver plants were exposed

to volatiles emitted from *M. sexta* damaged WT or LOX2-silenced or GFP-silenced (VIGS

control) plants as described in Paudel Timilsena *et al.* (2020). Briefly, two *N. benthamiana* plants

were placed inside a bell jar (10 L); one was treated as an emitter plant and the other as a

receiver plant (**Fig. 1**). Emitter plants were enclosed inside the wire case to prevent herbivore

escape. In volatile exposure treatments, emitter plants were continuously damaged by *M. sexta*

feeding, whereas control treatment, emitter plants remained undamaged. Volatiles emissions

from undamaged plants were below the detection limit of GC-FID (Supplementary **Fig. S3**).

To examine the role of individual GLV, a slow-release dispenser loaded with the

compound of interest was used as the volatile source and placed inside the bell jar with a receiver

plant. Control plants were exposed to undamaged WT plant together with a dispenser containing

lanolin only. To expose receiver plants to whole blend of volatiles, emitter plants were damaged

by *M. sexta* as described above. During the priming phase, there was continuous airflow inside

the bell jar (push air: 1.2 L min^{-1}, pull air: 1 L min^{-1}).

After 48 h of exposure to volatiles, receiver plants were transferred to individual clean

bell jars (8 L) and repeatedly wounded every 4 h during the light period for two consecutive days

(Paudel Timilsena *et al.* 2020). During each wounding event, two leaves were damaged once on

each side of the mid rib using a fine cheese grater (3 cm $\times$ 2 cm) that made ~180 holes per

wounding site. Thus made wounds were immediately treated with 10 µL of *M. sexta* regurgitant

(diluted with water, 1:1 v/v), which was identified previously to mimic insect herbivory (Heil *et*

al. 2001, 2012).

2.9 Headspace volatiles collection and analysis:

Following simulated herbivory, headspace volatiles from individual receiver plants were

collected at 4, 8, 12, 16, 20, 24, 28, 32 and 36 h using a dynamic headspace sampling system as

previously described by Paudel Timilsena *et al.* (2020). Briefly, clean, filtered air was pushed

into the bell jar at 1 L min^{-1}, and outgoing air was pulled through glass filter containing 30 mg of

Super-Q adsorbent (Alltech, Deerfield, IL, USA) at 0.8 L min^{-1}. Trapped volatiles on the filters

were eluded using 120 µL of a hexane: dichloromethane (1:1 v/v) containing nonyl acetate (4 ng

µL^{-1}) as an internal standard. Volatile compounds were quantified using gas chromatograph

coupled with a flame ionization detector (Agilent 6890 GC-FID, Agilent, CA, USA) fitted with a

capillary column (HP-5, 30 m $\times$ 0.25 mm ID $\times$ 0.25 µm film thickness; Supelco, PA, USA).

Samples (1 µL) were injected using an automated injector. The GC oven was programmed from

40 °C (2 min hold) to 150 °C at 4 °C min^{-1}, then to 290 °C at 40 °C min^{-1}, with a final 3 min

hold. Helium was used as the carrier gas at 14.79 psi. Peak areas were integrated using GC

ChemStation software, and volatile quantities were calculated relative to an internal standard.

Compound identities were verified by GC-MS (Agilent 6890N GC coupled to a 5973N

MS) using the same column and temperature program. (Z)-3-hexenal, (Z)-3-hexenol, (Z)-3-

hexenyl acetate, (E)-2-hexenal, linalool, limonene, (1, 8-) cineol, myrcene, α-pinene, β-pinene,

sabinene, and (E)-β-farnesene were verified by comparing retention times and mass spectra with

synthetic standards. Remaining compounds were identified by matching spectra to the NIST

library (GCMS Solution, Shimadzu, MD, USA).

2.10 Phytohormone JA extraction and quantification:

We measured JA accumulation at the damaged site of receiver plants as an indicator of

primed defense response. After 48 h of exposure to volatiles, receiver plants were damaged with

simulated herbivory. The third leaf from the base was selected and mechanically wounded on

both sides of the mid rib and immediately treated with *M. sexta* regurgitant. The damaged

portion of the leaf was harvested 30 min after challenge. The tissue was flash frozen into liquid

nitrogen and stored at -80 ˚C until analyzed. All the samples were ground into fine powder with

pre-cooled mortar and pestle under liquid nitrogen. The powdered sample (100 mg) was

transferred into a 2-mL screw-cap Fast Prep Tubes (Qbiogene, CA, USA) previously loaded with

1 g of Zirmil beads (1.1 mm; SEPR Ceramic Beads and Powders, NJ, USA), 400 μL extraction

solvent (isopropanol: H2O, 2:1 v/v, adjusted to pH 3 with hydrochloric acid) and 20 μL dihydro-

JA (10 ng μL^{-1}) an internal standard and vortexed thoroughly. Then 1 mL of dichloromethane

was added to each sample vortexed vigorously for 20 sec using a FastPrep FP 120 tissue

homogenizer (Qbiogene) and centrifuged at 10,000 rpm for 2 min. The bottom organic phase

(dichloromethane and isopropanol layer) was transferred to a 4-mL screw cap glass vial, dried

under nitrogen gas. The dried extract was re-suspended in 200 μL diethyl ether: methanol (9:1

v/v) and then esterified by adding 3 μL trimethylsilyldiazomethane solution (2 M in hexanes,

Andrich). The vial was then capped, vortexed, and incubated at room temperature for 30 min.

The excess of trimethylsilyldiazomethane was inactivated by adding 3 μL of 2 M acetic acid.

The vial was again capped, vortexed, and incubated at room temperature for 30 min. To collect

methylated and volatilized JA, a SuperQ filter attached to the vacuum source was inserted into

the vial through a cut in the septum. The vial was heated in a heating block to 180 °C for 2 min.

The methylated and volatilized JA was then trapped on the SuperQ filter for 2 min at 400 mL

min^{-1}. Trapped JA was eluded with 150 μL of dichloromethane and analyzed using a gas

chromatograph coupled to a mass spectrometer as previously described (Paudel Timilsena *et al.*

2020). The methylated product of endogenous JA was quantified by correlating the peak area of

the compound with that of the dihydro-JA (internal standard) using GC Chemstation software.

2.11 Herbivore performance bioassay:

First, we investigated the mortality rate of freshly hatched *M. sexta* on the plants that

were exposed to Z3HOL and the volatiles from undamaged emitter plants (here after air-

exposed). After 48 h of exposure, all the receiver plants were treated with 20 neonates per plant.

The mortality rate of the neonates was recorded at 24 h after the insect treatment.

To analyze the rate of feeding consumption, two third-instar larvae of *M. sexta* were

placed in an oblong container (30 cm × 15 cm) with four leaves of HIPV-exposed, Z3HOL-

exposed or air-exposed plants. The containers were previously coated with a thin layer of 1%

agar to prevent leaf wilting and lids had nylon mesh-covered holes for ventilation. Twenty-four h

after the start of the experiment, the area of leaf tissue removed was analyzed using the software

ImageJ (Schneider *et al*., 2012).

In a separate experiment, growth of third-instar larvae on Z3HOL-exposed or air-exposed

plants was analyzed. After 48 h of exposure to volatiles, receiver plants were treated with third-

instar larvae of *M. sexta* (4 plant^{-1}) for 7 days and the insect weight gain was recorded at the

beginning (day 0) and end of the experiment (day 7).

2.12 Herbivore behavioral response and dual choice bioassay:

We characterized behavioral responses of *M. sexta* larvae to HIPV-exposed, Z3HOL-

exposed and air-exposed control plants using a dual-choice assay in an oblong container (30 cm

$\times$ 15 cm). The 3rd and 4th leaves from HIPV- or Z3HOL-exposed or unexposed tobacco plants (a

combination of two treatments) were collected and placed on opposite sides of the container. The

container previously was coated with a thin layer of 1% agar to prevent leaf wilting or drying

out. Four 3rd instar *M. sexta* larvae were released at the center of the container and covered with

the lid containing nylon mesh for ventilation. Larvae were free to move inside the container and

choose between two leaf sources. The number of *M. sexta* on each side of the container was

recorded at 1, 6, 12 and 24 h.

2.13 Plant growth and development experiment:

To assess the effect of volatile exposure on plant growth and development, growth

parameters (such as plant height, number of secondary branches, secondary branch length) and

reproduction parameters (such as number of seed capsules, seed mass and seed germination

percentage) were measured. Two weeks after transplanting, *N. benthamiana* plants were exposed

to Z3HOL using the slow release volatile dispenser. Dispensers were calibrated to release

Z3HOL at the rate of 500 ng hr^{-1} for 1 week and were replaced weekly. Once the plants started to

flower, volatile dispensers were removed. Control treatments were exposed to identical

dispensers containing only lanolin. All the plants were kept in a greenhouse at $26 \pm 2\,^{\circ}C$ under

16L: 8D h photoperiod.

2.14 Statistical analysis:

The volatile data from the time course experiments were analyzed using the mixed-model

for repeated measures, SAS PROC MIXED (Saxton 1998), procedure in SAS statistical software

(SAS Institute Inc. 2013; SAS/STAT 9.4 User's Guide, NC, USA) as described previously

(Seidl-Adams *et al.* 2015; Paudel Timilsena *et al.* 2020). For repeated measures analysis with

Wald F-statistics, the best covariance structure was selected based on the Akaike Information

Criterion (AIC) and Bayesian Information Criterion (BIC), and the degrees of freedom were

adjusted using the Kenward-Roger method (Gomez *et al.* 2005). Since volatile emission follows

diurnal rhythm (Rim *et al.* 2019), volatile data were divided into three subsets (Day-1, Night-1,

and Day-2) and analyzed separately. For each dataset, two separate hypotheses were tested: 1)

mean volatile emissions at particular time points were same for all treatments, 2) mean volatile

emission for particular treatments were same at all time points. Then, the comparisons of interest

were extracted and evaluated for significance using the Bonferroni adjustment for multiple

comparisons.

All the other statistical analyses were conducted in R (R Core Team 2025). Data from

insect feeding assays were analyzed using one-way ANOVA, after confirming normally

distributed residuals (Shapiro-Wilk test), and homogeneity of variances (Levene's F-test). When

significant differences (P < 0.05) were found, pairwise post hoc comparisons of least squares

means were conducted using Least Significant Difference (LSD) at P□≤□0.05. Plant growth and

development data were analyzed by Student's *t*-tests or Welch two sample *t*-tests, depending on

whether samples violated the assumption of equal variance or not. The nature of dual-choice

assay with *M. sexta* larvae per case generated discrete data. Therefore, we used the

nonparametric statistical procedure, Wilcoxon matched-pair test, to determine the difference in

the number of larvae settled on either treatment.

3. Results

3.1 LOX2-silenced plant produces all the other volatiles except GLVs

Approximately 25 d after inoculation of early rosette-stage *N. benthamiana* with

Agrobacterium tumefaciens GV3101 harboring pTRV::LOX2 (gene of interest), pTRV::GFP

(vector control) and pTRV::PDS plasmid, new growth of PDS-silenced *N. benthamiana* plants

became completely bleached, demonstrating successful gene silencing (Supplementary Figure **1**).

Silencing of NbLOX2 did not markedly affect plant growth compared to the vector control

(NbGFP-silenced) or WT plants grown under the same environment conditions. To determine

the silencing efficiency of NaLOX2, we analyzed volatiles emitted from *M. sexta* feeding

damaged NaLOX2-silenced plants and compared them with similarly treated NaGFP-silenced

plants (vector control) and WT plants. Headspace collection followed by GC-FID analysis shows

no significance difference in mono- and sesquiterpene production by *M. sexta* fed WT, GFP-

silenced and LOX2-silenced plants (Fig. **2A**). However, LOX2-silenced plants emitted only

basal amounts of GLVs after *M. sexta* feeding, whereas WT and GFP-silenced plants showed

robust GLV induction (Fig. **2A**).

355   The main purpose of silencing GLV production was to investigate the role of GLV in

356 defense priming. In previous priming experiments, receiver plants were exposed to HIPVs for 48

357 h (Paudel Timilsena *et al*. 2020). Therefore, we used the same time window of 48 h and

358 confirmed that NbLOX2-silenced plants produce basal level of GLV after *M. sexta* feeding for

359 48 h (Fig. **2B**).

360 **3.2 GLVs are necessary for terpene priming**

361   WT receiver plants were exposed for 48 h to volatiles emitted from *M. sexta* damaged

362 WT, LOX-2 silenced, or GFP-silenced plants or from undamaged WT plants. Following volatile

363 exposure, all receiver plants were challenged (mechanical damage followed by *M. sexta*

364 regurgitant application) and emitted volatiles were collected for 36 h at 4-h intervals. When GLV

365 production was silenced in emitter plants, receiver plants produced significantly lower amount of

366 mono- and sesquiterpenes compared with plants exposed to the whole blend of volatile (i.e,

367 volatiles from *M. sexta* fed WT and GFP-silenced emitters). Receiver plants exposed to LOX2-

368 silenced plants and the undamaged WT plants produced similar amounts of mono- and

369 sesquiterpenes after challenge (Fig. **3A-B**).

370   Similarly, JA accumulation at the damaged site of the receiver plants was only strongly

371 primed after exposure to the whole blend of HIPV. Receiver plants exposed to *M. sexta* damaged

372 WT and GFP-silenced emitters produced significantly higher amounts of JA than those exposed

373 to undamaged emitter plants (Fig. **3C**). Although not statistically significant, receiver plants

374 exposed to *M. sexta* damaged LOX2-silenced emitters produced 82.39 % and 53.11 % less JA

375 than plants exposed to similarly treated WT and GFP-silenced emitters, respectively. Thus,

removing GLV from the HIPV blend diminished priming but did not completely abolish JA

induction, likely because other HIPV components still contribute to the response.

Together, these results suggest that GLV is necessary for full priming of mono- and

sesquiterpene production and contribute subsequently to JA accumulation in neighboring *N.*

benthamiana plants (Fig. **3**).

3.3 Exposure to (*Z*)-3-hexenol alone primes terpene emission

M. sexta damaged *N. benthamiana* plants emit four major GLV: (*Z*)-3-hexenal, (*E*)-2-

hexenal, (*Z*)-3-hexenol and (*Z*)-3-hexenyl acetate (Supplementary Fig. **S2D**). To determine

whether any individual GLV is sufficient for priming, plants were exposed to the physiologically

relevant concentration of each GLV using slow-release dispensers.

To test whether (*Z*)-3-hexenol (here after Z3HOL) acts as a priming agent between

plants, healthy *N. benthamiana* plants were exposed to synthetic Z3HOL at the release rate of

500 ng hr^{-1} for 48 h. Following exposure, receiver plants were repeatedly challenged (mechanical

damage followed by *M. sexta* regurgitant application) and volatiles were collected for 36 h. On

the second day of challenge, emission of sesquiterpene was significantly enhanced in Z3HOL-

exposed plants, demonstrating that exposure to synthetic Z3HOL alone is sufficient to prime

mono- and sesquiterpene emission in *N. benthamiana* plants (Fig. **4A-B**). However, exposure to

the full blend of volatiles resulted in stronger terpene than exposure to Z3HOL alone across all

sampling periods. These results suggest that while Z3HOL can independently induce priming,

primed responses are amplified when the plants are exposed to the complete blend of volatiles.

Although (*Z*)-3-hexenal and (*Z*)-3-hexenyl acetate were also tested in separate experiments,

Z3HOL was the only compound tested that primed *N. benthamiana* plants for induced volatile

emissions and JA production (**Fig. S4-5**).

3.4 *M. sexta* larvae performed poorly on volatile exposed plants

To further examine whether Z3HOL alone is sufficient to prime defense responses

against herbivory, *N. benthamiana* plants were exposed for 48 h to this volatile at comparable

amounts that were emitted from *M. sexta* damaged plants, i.e 500 ng hr^{-1}. After exposure, plants

were infested with freshly hatched *M. sexta* neonates (15 per plant), and neonate mortality was

recorded 48 h after release. Neonate mortality rate did not differ significantly among treatments

(F = 3.374, p = 0.054, n = 8; **Fig. 5A**). However, the overall neonate mortality rate was very

high, above 50%, across all treatments. We therefore evaluated *M. sexta* performance using early

third-instar larva. Larvae were fed on tobacco leaves for at least for 24 h prior to the experiment,

and only healthy, actively feeding larvae were selected. The mortality rate of these larvae was

lower than that of freshly hatched neonates (Supplementary Fig. **S6**).

N. benthamiana plants exposed to volatiles (either Z3HOL or HIPV) showed enhanced

defense against freshly molted 3rd-instar *M. sexta* larvae, as indicated by significantly reduced

leaf consumption and larval weight gain. *M. sexta* larvae feeding on Z3HOL- and HIPV-exposed

plants gained significantly less weight than those feeding on air-exposed control plants, with no

significant difference between the Z3HOL and HIPV treatments (F = 11.27, $p < 0.001$, Fig. **5b**).

Similarly, during a 24 h no-choice feeding assay, *M. sexta* consumed significantly less leaf

materials when they were placed on detached leaves of Z3HOL- and HIPV-exposed plants

compared to leaves from air-exposed plants (F = 7.84, p = 0.004; Fig. **5C**). Leaf consumption

was decreased by 46.65% and 32.78% on HIPV- and Z3HOL-exposed plants, respectively,

compared to air-exposed controls.

3.5 *M. sexta* larvae prefer air-exposed plants over volatile-exposed plants:

We characterized behavioral responses of *M. sexta* larvae to air-exposed, Z3HOL-

exposed or HIPV-exposed *N. benthamiana* plants using dual-choice assays with all the possible

combinations. At the 1 and 6 h after assay initiation, the 3rd-instar *M. sexta* larvae did not

discriminate between leaves from differently treated plants in the dual choice assay. However, at

12 and 24 h, significantly higher numbers of larvae settled on leaves from air-exposed plants,

suggesting reduced preference to Z3HOL- or HIPV-exposed plants **(Fig. 6A)**. Similarly, when

larvae were given a choice between Z3HOL- and HIPV-exposed plants, significantly more

larvae settled on Z3HOL-exposed plants than HIPV-exposed plant at 6, 12 and 24 h **(Fig. 6A),**

suggesting that HIPV-exposure might have primed additional changes or stronger responses in

the receiver plants that further reduced larval preference.

In addition to larval settlement, volatile exposure also significantly influenced the amount

of leaf consumption. During a 24 h feeding period, *M. sexta* larvae consumed significantly lower

amounts of leaf materials on HIPV-exposed plants when given a choice between air-exposed vs

HIPV-exposed plants, as well as between Z3HOL-exposed vs HIPV-exposed plants **(Fig. 6B-C)**.

When given a choice between air-exposed and Z3HOL-exposed plants, *M. sexta* larvae

consumed 12% less leaf materials from Z3HOL-exposed plants, although this difference was not

statistically significant ($p = 0.057$). Together, these results indicate that both HIPV- and Z3HOL-

exposed plants are less preferred by *M. sexta* larvae compared to air-exposed plants, with HIPV-

exposure eliciting the strongest deterrent effect.

3.6 *N. benthamiana* plants exposed to Z3HOL or HIPV exhibit enhanced defense responses

To understand the reduced performance and preference of *M. sexta* larvae on Z3HOL-
and HIPV-exposed plants, we quantified the levels of the key defense phytohormone, jasmonic
acid (JA), in exposed and air-exposed plants before and after herbivore damage. Herbivore
damage induced accumulation of JA in all plants at the wound site but the magnitude of
induction was significantly greater in Z3HOL- and HIPV-exposed plants. At 30 minutes after
challenge (mechanical damage + *M. sexta* regurgitant application), JA levels were 2.57 and 2.63
folds higher in Z3HOL- and HIPV-exposed plants, respectively, compared with air-exposed
plants (Fig. **7**).

3.7 (*Z*)-3-hexenol exposure enhanced the reproductive fitness of the plant:

To explore whether the exposure to Z3HOL affects plant growth and development,
tobacco plants were exposed to Z3HOL from two-week after transplanting until the onset of
flowering under insect-free conditions. The Z3HOL dispensers were replaced weekly to maintain
a consistent supply of Z3HOL. Z3HOL exposure did not affect vegetative growth, as exposed
plants showed no significant differences in plant height, number of secondary branches per plant,
and average length of secondary branch compared with air-exposed control plants (Figure **8A-
C**). In contrast, reproductive performance was significantly enhanced by Z3HOL exposure. At
harvest, Z3HOL-exposed plants produced significantly more seed capsules and greater seed mass
compared to unexposed control plants. Specifically, Z3HOL-exposed plants had about 13% more
seed capsules and a 24% increase in seed yield compared to control plants (Figure **8E-F**).

4. Discussion:

Herbivore induced plant volatiles (HIPV) play an important role in plant defense priming against insect herbivores. To identify the key volatile component required for plant defense priming, most studies used synthetic compounds at high concentrations, which led to controversies for many years (Dicke *et al.* 2003). Using transgenic plants impaired in their ability to emit specific volatile groups provides a robust approach for dissecting the role of individual volatiles in defense priming. In this study, we used *N. benthamiana* plants with silenced Green leaf Volatile (GLV) biosynthetic pathway (i.e, LOX3 gene silenced plants) to study the role of GLV as airborne priming signals. We assessed priming outcomes using insects' behavioral and physiological responses as indicators of defense activation, and plants growth and reproductive traits to evaluate potential fitness costs or benefits associated with priming.

Our results demonstrate that when GLV production was silenced in emitter plants, receiver plants no longer exhibited enhanced terpene priming and also showed reduced JA accumulation following simulated herbivory. However, exposure to a physiologically realistic concentration of (*Z*)-3-hexenol (Z3HOL) was sufficient to prime receiver plants for increased JA and terpene production following herbivore challenge. Together, these results indicate that GLV are necessary for full terpene priming and contribute substantially to JA amplification, while Z3HOL alone is sufficient to prime these defense responses. We further demonstrated that this priming has clear ecological consequences. *M. sexta* larvae consumed less leaf tissue and exhibited reduced growth on tobacco plants exposed to Z3HOL or the full HIPV blend, confirming that volatile-mediated priming translated to enhanced resistance. Importantly, we observed no evidence of fitness cost associated with priming. Z3HOL-exposed plants showed no differences in vegetative growth compared to air-exposed controls but produced more seed

capsules and greater seed mass. This suggests a positive or neutral trade-off between defense

priming and reproduction. Consistent with the definition of priming, elevated production of

terpenes and JA was observed only after simulated herbivory, rather than immediately following

volatile exposure, indicating primed rather than directly induced defense response (van Hulten *et*

al. 2006).

N. benthamiana plants emit four main components of GLV, such as (*Z*)-3-hexenal, (*E*)-2-

hexenal, (*Z*)-3-hexenol, and (*Z*)-3-hexenyl acetate, after *M. sexta* feeding damage. Among these,

only exposure to Z3HOL at physiologically relevant concentration (500 ng h^{-1}) primed *N.*

benthamianana for enhanced defense against herbivory. However, exposure to (*Z*)-3-hexenal,

(*E*)-2-hexenal, and (*Z*)-3-hexenyl acetate neither directly induced nor primed plant defenses

against herbivory. These results differ from studies using higher concentration of GLV, which

reported direct induction of defense gene expression and increased resistance to herbivores

(Farag *et al.* 2005; Yamauchi *et al.* 2018). Moreover, moderate concentrations of (*Z*)-3-hexenyl

acetate have been shown to prime *Zea mays* (maize) and *Populus deltoides × nigra* (hybrid

poplar) for enhanced JA production and JA-responsive defense genes expression after herbivory

(Engelberth *et al.* 2007; Frost *et al.* 2008). This suggests that plants may have evolved species-

species mechanisms for responding to volatile cues.

Although Z3HOL alone was sufficient to prime *N. benthamiana*, exposure to the full

blend of HIPV consistently primed for stronger defense responses than the exposure to Z3HOL

alone. This pattern also aligns with LOX2-silencing results, where removal of the GLV fraction

from the complex HIPV blend reduced the magnitude of JA priming but did not abolish it,

indicating that additional volatile may have partially compensate for the loss of GLV. Previous

studies observed the similar pattern, where exposure to indole and (*Z*)-3-hexenyl acetate induced

stronger defense in *Z. mays* than single compounds (Hu *et al.* 2019). These findings support the

concept that the full blend of HILV provides a more reliable indicator of herbivore presence than

individual volatiles, as proposed by Douma *et al.* (2019).

515   Behavioral assays further revealed that *M. sexta* preferred to feed on air-exposed plants

over Z3HOL- or HIPV-exposed plants. While *M. sexta* larvae initially showed no discrimination

among treatments, after several hours, they settled on the leaves from the control plant over

Z3HOL- or HIPV-exposed plants. We expected that, after herbivore attack, Z3HOL- or HIPV-

exposed plants emitted volatiles in larger amounts than the control; and therefore, be less

preferred by *M. sexta* larvae. HIPV-exposed plants do not only accelerate the production of

volatiles after herbivore damage but also absorb and reemit significantly higher amounts of

volatiles than air-exposed plants for a certain amount of time without herbivore damage (Paudel

Timilsena *et al.* 2020). Yet, the higher emission of the volatiles from HIPV-exposed plants did

not affect *M. sexta* settlement at the beginning of the experiment. This suggests that feeding

preference of *M. sexta* is based on their gustatory response rather than olfactory response alone.

The *M. sexta* larval olfactory response to volatiles in our study is in line with previous studies

showing that HIPV induced by lepidopteral herbivore do not repel conspecific larvae (Carroll *et*

al. 2006, 2008; von Mérey *et al.* 2013). Although not repelled by volatiles, *M. sexta* larvae

consumed significantly less leaf materials from Z3HOL- or HIPV-exposed plants than from

unexposed control plants, suggesting induction of feeding deterrents or toxic compounds.

Reduced herbivore feeding on volatile-exposed plants has been widely reported under both

laboratory and field conditions. Under laboratory conditions, Arimura *et al.* (2000) reported less

feeding damage by *Teranychus urticae* on *Phaseolus lunatus* plants that were previously exposed

to herbivore-damaged conspecific plants. Under field conditions, *N. attenuata* plants and

sagebrush suffered less damage from naturally occurring herbivores when exposed to volatiles

from artificially damaged sagebrush plants (Karban *et al.* 2000; Karban *et al.* 2006). Similarly,

Salix exigua and *S. lemmonii* exposed to volatiles from mechanically damaged conspecific plants

also suffered less herbivore damage in the fields (Pearse *et al.* 2013). The reduced feeding

preference by herbivores may be mediated by direct toxicity of terpenes or presence of

antifeedant compound/toxin or both, but this would require additional data to demonstrate.

Consistent with reduced feeding, *M. sexta* larvae grew poorly on plants exposed to

Z3HOL or HIPV. Reduced herbivore performance on the primed/volatile-exposed plants has

been reported across multiple systems, including reduced growth of *Spodoptera littoralis* on

maize (von Mérey *et al.* 2013) and reduced consumption and weight gain of *Plagiodera*

versicolora on willow (Yoneya *et al.* 2014). Interestingly, such reductions in herbivore

performance can be achieved by exposing herbivores to mixture of GLV (von Mérey *et al.* 2013)

and treating plants with individual GLV, such as Z3HOL (in this study) and (Z)-3-hexenyl

acetate (Hu *et al.* 2019).

One likely explanation for the reduced weight gain of *M. sexta* on volatiles-exposed

plants is enhanced activation of plant defenses. Since JA plays a central role in regulating direct

defenses against herbivores (Yang *et al.* 2019), we quantified JA levels in volatile-exposed

plants. Our results show that the Z3HOL- and HIPV-exposed plants accumulated significantly

higher amounts of JA at the wounding sites than the air-exposed control plants. Consistent with

our findings, Z3HOL-exposure has been shown to increase the expression of JA-biosynthesis

related genes and JA accumulation in several plants, including *Z. mays* (Farag *et al.* 2005;

Engelberth *et al.* 2013), *Camellia sinensis* (Xin *et al.* 2016; Chen *et al.* 2020), and *Solanum*

lycopersicum (Su *et al.* 2020; Yang *et al.* 2020). Although, we did not directly test how JA

regulated specific defense traits in this study, numerous studies showed that JA activates genes

encoding antiherbivore proteins and defense metabolites (Engelberth *et al.* 2004; Farag *et al.*

2005; Ton *et al.* 2007; Frost *et al.* 2008). For instance, exogenous Z3HOL treatment primes JA

accumulation and induces JA-responsive genes encoding digestion inhibitors such as proteinase

inhibitors, polyphenol oxidases and xylanase inhibitors, and enhance resistance against chewing

herbivores (Engelberth *et al.* 2013; Xin *et al.* 2016). Altogether, these findings suggest that

enhanced JA accumulation in the Z3HOL- and HIPV-exposed plants likely contributed to

increased production of anti-herbivore defense metabolites and reduced larval performance.

In addition to enhanced direct defense, Z3HOL- and HIPV-exposed plants emitted higher

levels of defensive volatiles than control plants after herbivory, potentially increasing attraction

of natural enemies. (*E*)-α-bergamotene and (*E*)-β-farnesene, two major sesquiterpenes emitted by

herbivore damaged *N. benthamiana* plants (Paudel Timilsena *et al.* 2020), were strongly induced

in HIPV- and Z3HOL-exposed plants after herbivore damages. Previous studies have shown that

(*E*)-α-bergamotene reduces herbivore population on *N. attenuate* by attracting a generalist

predator and deterring ovipositing *M. sexta* moths (Kessler & Baldwin 2001; Halitschke *et al.*

2008). Similarly, transgenic *A. thaliana* expressing maize terpene synthase (ZmTPS10), gene

that produce (*E*)-α-bergamotene and (*E*)-β-farnesene, attracted the parasitoid *Cotesia*

marginiventris and reduced performance of *S. littoralis* (Schnee *et al.* 2006). Thus, volatile

exposure (Z3HOL and HIPV) likely enhances both direct and indirect defense against

herbivores.

Despite enhanced defense activation, Z3HOL exposure did not negatively affect plant

growth and development. Instead, Z3HOL-exposed plants produced more seed capsules, which

significantly increased seed weight. Similar fitness benefits have been reported in higher

seedling survivorship, branch growth and inflorescence production in sagebrush and *N. attenuata*

plants primed by volatiles from clipped neighbors, which showed increased branch growth,

numbers of flower and seed production (Karban & Maron 2002; Karban *et al.* 2012). In addition,

recent evidence shows that GLV exposure can enhance plant growth and yield through

jasmonate-dependent plant-soil feedback that alter rhizosphere microbial communities (Hu et al.,

2025). These results suggest that defense priming does not necessarily impose growth or

reproductive penalties. If priming were inherently costly, primed plants would be expected to

show reduced reproductive fitness due to the cost of secondary metabolite production (reviewed

in Guo *et al.* 2018). However, primed plants enhance defense metabolite production only after

herbivore-attack. Under herbivore-free conditions, as in this study, Z3HOL exposure did not

directly induce defense, allowing plants to allocate resources to growth and reproduction.

Although priming cost may become apparent under stressful conditions, such as soil nutrient

deficiency, low light, drought, frost or herbivory (Karban & Maron 2002; Engelberth &

Engelberth 2019; Yip *et al*. 2019), priming with GLV (Z3HAC) can temporarily reduce leaf

growth and shift plant physiology toward defense (Engelberth & Engelberth 2019), no such costs

were detected under the controlled conditions in this study.

Exposure to Z3HOL enhanced reproductive fitness in *N. benthamiana*. Although the

reason behind the enhanced reproductive fitness is unclear, we speculated that JA, which

calibrates growth-defense balance (Guo *et al.* 2018), might have shifted resource allocation

towards reproduction in the absence of herbivory. JA signaling is known to regulate flower

development, especially flower abscission and male sterility (Acosta *et al.* 2009; Song *et al.*

2011; Oh *et al.* 2013; Niwa *et al.* 2018). For example, *N. attenuate* plants with impaired JA

signaling (NaJAZd-silenced) increased flower abscission in later stages of flower development

which resulted in reduced number of seed capsules and seed mass than wild type plant (Oh *et al.*

2013). Similarly, the constitutive level of JA and its conjugate JA-Ile were much higher in flower

buds than those in wounded leaves (Hause *et al.* 2000; Suza *et al.* 2010), suggesting importance

of JA signaling in plant reproductive development. Here, our data showed that Z3HOL-exposure

increased JA accumulation in leaves after herbivore attack. However, when a primed plant is not

under attack, plants could shift resources for reproduction. Whether Z3HOL-exposure also

promoted JA biosynthesis in flowers and directly affected flower and seed capsule development,

remains to be elucidated.

In summary, silencing GLV production in emitter plants abolished priming of JA

accumulation and terpenes emission in receiver plants after simulated herbivory. This

demonstrates that GLVs are necessary for defense priming in *N. benthamiana* plants under the

experimental condition tested. Moreover, exposure to physiologically relevant concentration of

Z3HOL, a common GLV, alone was sufficient to enhance herbivore resistance and reproductive

fitness in *N. benthamiana.* Since defense was activated only upon herbivore attack, primed plants

did not incur detectable fitness cost under herbivore free conditions. Further long-term field

studies are needed to determine whether Z3HOL-mediated priming consistently confers benefits

under variable environmental biotic stress, and whether it can be effectively exploited for

sustainable pest management.

Acknowledgments:

The authors would like to thank the Department of Entomology and the Center for Chemical

Ecology at The Pennsylvania State University for providing support to this study. We dedicate

this paper to the memory of late Prof. James H. Tumlinson, who supervised this project with

dedication until the end.

Author Contribution:

Conceptualization; BPT, ISA, and JHT. Methodology; BPT, ISA, and SRH. Formal Analysis;

BPT and ISA. Visualization; BPT. Writing - original draft: BPT. Writing – Review & Editing;

BPT, ISA, GF, SRH, and JHT. Supervision; GF and JHT.

Conflict of Interest:

The authors have no conflict of interest to declare.

Funding Statement:

This research received no specific grant from any funding agency in the public, commercial or

not-for-profit sectors.

Data Availability:

The datasets generated and analyzed during the current study are available from the

corresponding author upon reasonable reques.

References:

Acosta IF, Laparra H, Romero SP, Schmelz E, Hamberg M, Mottinger JP, Moreno MA, Dellaporta SL. 2009. tasselseed1 is a lipoxygenase affecting jasmonic acid signaling in sex determination of maize. Science (New York, N.Y.) **323**, 262–265.

Arimura G, Ozawa R, Shimoda T, Nishioka T, Boland W, Takabayashi J. 2000. Herbivory-induced volatiles elicit defence genes in lima bean leaves. Nature **406**, 512–515.

Baldwin IT, Halitschke R, Paschold A, von Dahl CC, Preston CA. 2006. Volatile signaling in plant-plant interactions: 'talking trees' in the genomics era. Science (New York, N.Y.) **311**, 812–815.

Baldwin IT, Schultz JC. 1983. Rapid changes in tree leaf chemistry induced by damage: evidence for communication between plants. Science (New York, N.Y.) **221**, 277–279.

Bate NJ, Rothstein SJ. 1998. C6-volatiles derived from the lipoxygenase pathway induce a subset of defense-related genes. The Plant Journal: For Cell and Molecular Biology **16**, 561–569.

Carroll MJ, Schmelz EA, Meagher RL, Teal PEA. 2006. Attraction of Spodoptera frugiperda Larvae to Volatiles from Herbivore-Damaged Maize Seedlings. Journal of Chemical Ecology **32**, 1911–1924.

Carroll MJ, Schmelz EA, Teal PEA. 2008. The attraction of Spodoptera frugiperda neonates to cowpea seedlings is mediated by volatiles induced by conspecific herbivory and the elicitor inceptin. Journal of Chemical Ecology **34**, 291–300.

Chen SL, Zhang LP, Cai XM, Bian L, Luo ZX, Li ZQ, Ge LG, Chen ZM, Xin ZJ. 2020. (Z)-3-Hexenol Induces Tea Defense against Ectropis grisescens in the Field. Russian Journal of Plant Physiology **67**, 572–580.

Cofer TM, Seidl-Adams I, Tumlinson JH. 2018. From Acetoin to (Z)-3-Hexen-1-ol: The Diversity of Volatile Organic Compounds that Induce Plant Responses. Journal of Agricultural and Food Chemistry **66**, 11197–11208.

Dicke M, Agrawal AA, Bruin J. 2003. Plants talk, but are they deaf? Trends in Plant Science **8**, 403–405.

Douma JC, Ganzeveld LN, Unsicker SB, Boeckler GA, Dicke M. 2019. What makes a volatile organic compound a reliable indicator of insect herbivory? Plant, Cell & Environment **42**, 3308–3325.

Engelberth J, Alborn HT, Schmelz EA, Tumlinson JH. 2004. Airborne signals prime plants against insect herbivore attack. Proceedings of the National Academy of Sciences **101**, 1781–1785.

Engelberth J, Contreras CF, Dalvi C, Li T, Engelberth M. 2013. Early Transcriptome Analyses of Z-3-Hexenol-Treated Zea mays Revealed Distinct Transcriptional Networks and Anti-Herbivore Defense Potential of Green Leaf Volatiles. PLOS ONE **8**, e77465.

Engelberth J, Engelberth M. 2019. The Costs of Green Leaf Volatile-Induced Defense Priming: Temporal Diversity in Growth Responses to Mechanical Wounding and Insect Herbivory | MDPI. Plants **8**, 1–13.

Engelberth J, Seidl-Adams I, Schultz JC, Tumlinson JH. 2007. Insect elicitors and exposure to green leafy volatiles differentially upregulate major octadecanoids and transcripts of 12-oxo phytodienoic acid reductases in Zea mays. Molecular plant-microbe interactions: MPMI **20**, 707–716.

Erb M, Veyrat N, Robert CAM, Xu H, Frey M, Ton J, Turlings TCJ. 2015. Indole is an essential herbivore-induced volatile priming signal in maize. Nature Communications **6**, 6273.

Farag MA, Fokar M, Abd H, Zhang H, Allen RD, Paré PW. 2005. (Z)-3-Hexenol induces defense genes and downstream metabolites in maize. Planta **220**, 900–909.

Farag MA, Paré PW. 2002. C6-Green leaf volatiles trigger local and systemic VOC emissions in tomato. Phytochemistry **61**, 545–554.

Farmer EE, Ryan CA. 1990. Interplant communication: airborne methyl jasmonate induces synthesis of proteinase inhibitors in plant leaves. Proceedings of the National Academy of Sciences **87**, 7713–7716.

Frost CJ, Mescher MC, Carlson JE, De Moraes CM. 2008. Plant Defense Priming against Herbivores: Getting Ready for a Different Battle. Plant Physiology **146**, 818–824.

Godard K-A, White R, Bohlmann J. 2008. Monoterpene-induced molecular responses in Arabidopsis thaliana. Phytochemistry **69**, 1838–1849.

Gomez EV, Schaalje GB, Fellingham GW. 2005. Performance of the Kenward–Roger Method when the Covariance Structure is Selected Using AIC and BIC. Communications in Statistics - Simulation and Computation **34**, 377–392.

Guo Q, Major IT, Howe GA. 2018. Resolution of growth-defense conflict: mechanistic insights from jasmonate signaling. Current Opinion in Plant Biology **44**, 72–81.

Halitschke R, Baldwin IT. 2003. Antisense LOX expression increases herbivore performance by decreasing defense responses and inhibiting growth-related transcriptional reorganization in Nicotiana attenuata. The Plant Journal **36**, 794–807.

Halitschke R, Stenberg JA, Kessler D, Kessler A, Baldwin IT. 2008. Shared signals -'alarm calls' from plants increase apparency to herbivores and their enemies in nature. Ecology Letters **11**, 24–34.

Halitschke R, Ziegler J, Keinänen M, Baldwin IT. 2004. Silencing of hydroperoxide lyase and allene oxide synthase reveals substrate and defense signaling crosstalk in Nicotiana attenuata. The Plant Journal: For Cell and Molecular Biology **40**, 35–46.

Hause B, Stenzel I, Miersch O, Maucher H, Kramell R, Ziegler J, Wasternack C. 2000. Tissue-specific oxylipin signature of tomato flowers: allene oxide cyclase is highly expressed in distinct flower organs and vascular bundles. The Plant Journal: For Cell and Molecular Biology **24**, 113–126.

Heil M, Ibarra-Laclette E, Adame-Álvarez RM, Martínez O, Ramirez-Chávez E, Molina-Torres J, Herrera-Estrella L. 2012. How Plants Sense Wounds: Damaged-Self Recognition Is Based on Plant-Derived Elicitors and Induces Octadecanoid Signaling. PLOS ONE **7**, e30537.

Heil M, Karban R. 2010. Explaining evolution of plant communication by airborne signals. Trends in Ecology & Evolution **25**, 137–144.

Heil M, Koch T, Hilpert A, Fiala B, Boland W, Linsenmair KE. 2001. Extrafloral nectar production of the ant-associated plant, Macaranga tanarius, is an induced, indirect, defensive response elicited by jasmonic acid. Proceedings of the National Academy of Sciences **98**, 1083–1088.

Hu L, Ye M, Erb M. 2019. Integration of two herbivore-induced plant volatiles results in synergistic effects on plant defence and resistance. Plant, Cell & Environment **42**, 959–971.

Hu L, Zhang K, Xu Y, *et al.* 2025. Herbivory-induced green leaf volatiles increase plant performance through jasmonate-dependent plant–soil feedbacks. Nature Plants **11**, 1001–1017.

van Hulten M, Pelser M, van Loon LC, Pieterse CMJ, Ton J. 2006. Costs and benefits of priming for defense in Arabidopsis. Proceedings of the National Academy of Sciences **103**, 5602–5607.

Karban R, Baldwin IT, Baxter KJ, Laue G, Felton GW. 2000. Communication between plants: induced resistance in wild tobacco plants following clipping of neighboring sagebrush. Oecologia **125**, 66–71.

Karban R, Ishizaki S, Shiojiri K. 2012. Long-term demographic consequences of eavesdropping for sagebrush. Journal of Ecology **100**, 932–938.

Karban R, Maron J. 2002. The Fitness Consequences of Interspecific Eavesdropping between Plants. Ecology **83**, 1209–1213.

Karban R, Shiojiri K, Huntzinger M, McCall AC. 2006. Damage-induced resistance in sagebrush: volatiles are key to intra- and interplant communication. Ecology **87**, 922–930.

Karban R, Yang LH, Edwards KF. 2014. Volatile communication between plants that affects herbivory: a meta-analysis. Ecology Letters **17**, 44–52.

Kessler A, Baldwin IT. 2001. Defensive function of herbivore-induced plant volatile emissions in nature. Science (New York, N.Y.) **291**, 2141–2144.

Kessler A, Halitschke R, Diezel C, Baldwin IT. 2006. Priming of plant defense responses in nature by airborne signaling between Artemisia tridentata and Nicotiana attenuata. Oecologia **148**, 280–292.

Kost C, Heil M. 2006. Herbivore-induced plant volatiles induce an indirect defence in neighbouring plants. Journal of Ecology **94**, 619–628.

Meents AK, Chen S-P, Reichelt M, Lu H-H, Bartram S, Yeh K-W, Mithöfer A. 2019. Volatile DMNT systemically induces jasmonate-independent direct anti-herbivore defense in leaves of sweet potato (Ipomoea batatas) plants. Scientific Reports **9**, 17431.

von Mérey GE, Veyrat N, D'Alessandro M, Turlings T. 2013. Herbivore-induced maize leaf volatiles affect attraction and feeding behavior of Spodoptera littoralis caterpillars. Frontiers in Plant Science **4**.

von Mérey G, Veyrat N, Mahuku G, Valdez RL, Turlings TCJ, D'Alessandro M. 2011. Dispensing synthetic green leaf volatiles in maize fields increases the release of sesquiterpenes by the plants, but has little effect on the attraction of pest and beneficial insects. Phytochemistry **72**, 1838–1847.

Nethra P, Nataraja KN, Rama N, Udayakumar M. 2006. Standardization of environmental conditions for induction and retention of post-transcriptional gene silencing using tobacco rattle virus vector. Current Science **90**, 431–435.

Niwa T, Suzuki T, Takebayashi Y, Ishiguro R, Higashiyama T, Sakakibara H, Ishiguro S. 2018. Jasmonic acid facilitates flower opening and floral organ development through the upregulated expression of SlMYB21 transcription factor in tomato. Bioscience, Biotechnology, and Biochemistry **82**, 292–303.

Oh Y, Baldwin IT, Galis I. 2013. A Jasmonate ZIM-Domain Protein NaJAZd Regulates Floral Jasmonic Acid Levels and Counteracts Flower Abscission in Nicotiana attenuata Plants. (GK Pandey, Ed.). PLoS ONE **8**, e57868.

Paschold A, Halitschke R, Baldwin IT. 2006. Using 'mute' plants to translate volatile signals. The Plant Journal: For Cell and Molecular Biology **45**, 275–291.

Paudel Timilsena B, Seidl-Adams I, Tumlinson JH. 2020. Herbivore-specific plant volatiles prime neighboring plants for nonspecific defense responses. Plant, Cell & Environment **43**, 787–800.

Pearse IS, Hughes K, Shiojiri K, Ishizaki S, Karban R. 2013. Interplant volatile signaling in willows: revisiting the original talking trees. Oecologia **172**, 869–875.

R Core Team. 2025. R: The R Project for Statistical Computing. https://www.r-project.org/. Accessed January 2026.

Rhoades DF. 1983. Responses of Alder and Willow to Attack by Tent Caterpillars and Webworms: Evidence for Pheromonal Sensitivity of Willows. ACS Symposium Series. Plant Resistance to Insects. American Chemical Society, 55–68.

Rim H, Sekiguchi S, Ozawa R, Kainoh Y, Arimura G. 2019. Diurnal rhythm of volatile emissions from damaged Brachypodium distachyon affects the temporal preferences of tritrophic interactions. Journal of Plant Interactions **14**, 143–150.

Sasidharan R. 2022. Ethylene — the lifeguard hormone. Nature Reviews Molecular Cell Biology **23**, 643–643.

Saxton AM. 1998. A Macro for Converting Mean Separation Output to Letter Groupings in PROC MIXED. 23rd SAS User Group Intl. SAS Institute, Cary, NC, 1243–1246.

Schnee C, Köllner TG, Held M, Turlings TCJ, Gershenzon J, Degenhardt J. 2006. The products of a single maize sesquiterpene synthase form a volatile defense signal that attracts natural enemies of maize herbivores. Proceedings of the National Academy of Sciences **103**, 1129–1134.

Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis. Nature Methods **9**, 671–675.

Seidl-Adams I, Richter A, Boomer KB, Yoshinaga N, Degenhardt J, Tumlinson JH. 2015. Emission of herbivore elicitor-induced sesquiterpenes is regulated by stomatal aperture in maize (Zea mays) seedlings. Plant, Cell & Environment **38**, 23–34.

Song S, Qi T, Huang H, Ren Q, Wu D, Chang C, Peng W, Liu Y, Peng J, Xie D. 2011. The Jasmonate-ZIM domain proteins interact with the R2R3-MYB transcription factors MYB21 and MYB24 to affect Jasmonate-regulated stamen development in Arabidopsis. The Plant Cell **23**, 1000–1013.

Su Q, Yang F, Zhang Q, Tong H, Hu Y, Zhang X, Xie W, Wang S, Wu Q, Zhang Y. 2020. Defence priming in tomato by the green leaf volatile (Z)-3-hexenol reduces whitefly transmission of a plant virus. Plant, Cell & Environment **43**, 2797–2811.

Suza WP, Rowe ML, Hamberg M, Staswick PE. 2010. A tomato enzyme synthesizes (+)-7-iso-jasmonoyl-L-isoleucine in wounded leaves. Planta **231**, 717–728.

Tamogami S, Rakwal R, Agrawal GK. 2008. Interplant communication: Airborne methyl jasmonate is essentially converted into JA and JA-Ile activating jasmonate signaling pathway and VOCs emission. Biochemical and Biophysical Research Communications **376**, 723–727.

Ton J, D'Alessandro M, Jourdie V, Jakab G, Karlen D, Held M, Mauch-Mani B, Turlings TCJ. 2007. Priming by airborne signals boosts direct and indirect resistance in maize. The Plant Journal: For Cell and Molecular Biology **49**, 16–26.

Xin Z, Li X, Li J, Chen Z, Sun X. 2016. Application of chemical elicitor (Z)-3-hexenol enhances direct and indirect plant defenses against tea geometrid Ectropis obliqua. BioControl **61**, 1–12.

Yamauchi Y, Matsuda A, Matsuura N, Mizutani M, Sugimoto Y. 2018. Transcriptome analysis of Arabidopsis thaliana treated with green leaf volatiles: possible role of green leaf volatiles as self-made damage-associated molecular patterns. Journal of Pesticide Science **43**, 207–213.

Yang J, Duan G, Li C, Liu L, Han G, Zhang Y, Wang C. 2019. The Crosstalks Between Jasmonic Acid and Other Plant Hormone Signaling Highlight the Involvement of Jasmonic Acid as a Core Component in Plant Response to Biotic and Abiotic Stresses. Frontiers in Plant Science **10**.

Yang F, Zhang Q, Yao Q, Chen G, Tong H, Zhang J, Li C, Su Q, Zhang Y. 2020. Direct and indirect plant defenses induced by (Z)-3-hexenol in tomato against whitefly attack. Journal of Pest Science **93**, 1243–1254.

Yao Y, Danna CH, Zemp FJ, Titov V, Ciftci ON, Przybylski R, Ausubel FM, Kovalchuk I. 2011. UV-C–Irradiated Arabidopsis and Tobacco Emit Volatiles That Trigger Genomic Instability in Neighboring Plants[W]. The Plant Cell **23**, 3842–3852.

Yip EC, Tooker JF, Mescher MC, De Moraes CM. 2019. Costs of plant defense priming: exposure to volatile cues from a specialist herbivore increases short-term growth but reduces rhizome production in tall goldenrod (Solidago altissima). BMC plant biology **19**, 209.

Yoneya K, Kugimiya S, Takabayashi J. 2014. Leaf beetle larvae, Plagiodera versicolora (Coleoptera: Chrysomelidae), show decreased performance on uninfested host plants

exposed to airborne factors from plants infested by conspecific larvae. Applied
Entomology and Zoology **49**, 249–253.

Zavala JA, Baldwin IT. 2004. Fitness benefits of trypsin proteinase inhibitor expression in
Nicotiana attenuata are greater than their costs when plants are attacked. BMC ecology **4**,
11.

Zavala JA, Patankar AG, Gase K, Hui D, Baldwin IT. 2004. Manipulation of endogenous
trypsin proteinase inhibitor production in Nicotiana attenuata demonstrates their function
as antiherbivore defenses. Plant Physiology **134**, 1181–1190.

Zeringue HJ. 1992. Effects of C6C10 alkenals and alkanals on eliciting a defence response in
the developing cotton boll. Phytochemistry 3**1,** 2305–2308.

Ziegler J, Keinänen M, Baldwin IT. 2001. Herbivore-induced allene oxide synthase transcripts
and jasmonic acid in Nicotiana attenuata. Phytochemistry 5**8,** 729–738.

Figure Legends:

Fig. 1 Experimental design for GLV-mediated defense priming. (A) Priming phase (0-48 h): In the first experiment, receiver plants (R) were exposed to volatiles from *M. sexta*-damaged WT, LOX2-silenced, or GFP-silenced (VIGS-control) emitter plants (E). Receiver plants exposed to undamaged WT plants served as the negative control. In separate experiments, receiver plants were exposed to synthetic GLV [(*Z*)-3-hexenal (Z3HAL), (*Z*)-3hexenol (Z3HOL), or (*Z*)-3-hexenyl acetate (Z3HAC)] via slow-release dispensers. Receiver plants exposed to lanolin dispensers and undamaged WT plants as negative control, while those exposed to *M. sexta*-damaged WT served as positive control. (**B) Post-exposure challenge (0-36 h):** After priming, receiver plants (R) were transferred to clean bell jars and subjected to repetitive wounding every 4 h for 36 h using a cheese grater. Wounds were immediately treated with 10 μL of *M. sexta* regurgitant (1:1 v/v) to simulate herbivory. Volatiles were collected via SuperQ filters.

Fig. 2 HIPVs emitted from *M. sexta* damaged *N. benthamiana* plants. **A)** Chromatographic profiles of HIPVs emitted by *M. sexta* damaged wild type (top), GFP-silenced (middle), and LOX2-silenced (bottom) plants. *N. benthamiana* plants were continuously damaged by fourth-instar *M. sexta* feeding for 48 h. The chromatogram represents HIPV profile collected from 24 to 28 h after feeding initiation. Five major families of HIPVs were induced by *M. sexta* feeding on *N. benthamiana*: **green leaf volatiles (GLV):** (*Z*)-3-hexenal [3], (*E*)-2-hexenal [4], (*Z*)-3-hexenol [5], (*Z*)-3-hexenyl acetate [11], **monoterpenes:** α-pinene [8], sabinene [9], β-pinene [10], limonene [12], 1,8-cineole [13], (*E*)-β-ocimene [14], linalool [15], nonanal [16], **sesquiterpenes:** (*E*)-α-bergamotene [21], unknown sesquiterpene#1 [22], (*E*)-β-farnesene

[23], aristolochene (4,4-di epi) [24], β-bisabolene [25], β-sesquiphellandrene [26], **aldoximes:** propyl aldoxime, 2 methyl-,syn-[1], propyl aldoxime, 2 methyl-,anti-[2], butyl aldoxime, 2 methyl-,syn- [6], butyl aldoxime, 2 methyl-,anti-[7], and **aromatic compound**: indole [17]. **B)** Levels of GLVs emitted by M. sexta damaged plants. Early fourth-instar M. sexta (one larva per plant) was feeding continuously on the plant for 48 hours. HIPVs collection started on 2nd day, 16 h after feeding initiation and continued up to 48 h in 4-h intervals. Values represent mean ± SE (n = 4). Shaded areas indicate nighttime volatile collections.

Fig. 3 Receiver *N. benthamiana* plants did not get primed for terpene emissions when GLV emission was silenced in emitter plants. *N. benthamiana* plants were exposed to volatiles from *M. sexta* damaged WT (green), LOX2-silenced (dark red), GFP-silenced (blue) or undamaged (gray) plants for 48 h. After 48 h of exposure to volatiles, the receiver plants were transferred to individual clean bell jars and challenged repeatedly by mechanical wounding followed by *M. sexta* regurgitant application. The graph shows the total amount of **A)** monoterpenes and **B)** sesquiterpenes emitted from the challenged receiver plants at different time intervals after initial challenge. Values represent mean ± SE (n = 3). Data were analyzed with a mixed model for repeated measures. Different letters indicate significant ($p <$ 0.05) differences between treatments within time points, with Bonferroni's correction for multiple comparisons. Shaded areas indicate nighttime volatile collections. **C)** For JA analysis, after 48 h of exposure to volatiles, receiver plants were challenged with mechanical damage followed by *M. sexta* regurgitant application. Wounded portions of the leaf were harvested 30 min after challenge and JA accumulation in the harvested tissue was analyzed (ANOVA, Tukey HSD, n = 3, $p < 0.01$).

Fig. 4 Exposure to Z3HOL alone primed *N. benthamiana* plants for enhanced production of mono and sesquiterpenes. *N. benthamiana* plants were exposed to volatiles from *M. sexta* damaged WT (green), undamaged WT plants (gray) or synthetic Z3HOL (blue) for 48 h. After 48 h of exposure to volatiles, the receiver plants were transferred to individual clean bell jars and challenged repeatedly by mechanical wounding followed by *M. sexta* regurgitant application. The graph shows the total amount of **A)** monoterpenes and **B)** sesquiterpenes emitted from the challenged receiver plants at different time intervals after initial challenge. Values represent mean ± SE (n = 4). Data were analyzed with a mixed model for repeated measures. Different letters indicate significant ($p < 0.05$) differences between treatments within time points, with Bonferroni's correction for multiple comparisons. Shaded areas indicate nighttime volatile collections.

Fig. 5 Volatile (Z3HOL and HIPV) exposure primed defenses against *M. sexta* larvae. *N. benthamiana* plants were exposed for 48 h to volatile from undamaged plant (air-exposed, gray), 500ng of Z3HOl (Z3HOL-exposed, blue) or HIPVs from *M. sexta* damaged plants (HIPV-exposed, green). **A)** Mortality rate of freshly hatched *M. sexta* neonates 48 h after the neonate release on receiver plants (ANOVA, F = 3.374, n = 8, $p = 0.054$), **B)** Amount of leaf area consumed by two early-3rd-instar *M. sexta* larvae during 24 h feeding period (ANOVA, Tukey HSD, F = 11.27, n = 6, $p < 0.001$), **C)** Larval weight gain at 0 and 7 days after feeding on receiver plants (ANOVA, Tukey HSD, F = 7.84, n = 6, $p < 0.01$). The bar represents the mean ± SE. Letters indicate significant differences among different treatments.

Fig. 6 Feeding preference of *M. sexta* larvae is influenced by volatile exposure treatments. **A)** The preference of *M. sexta* larvae on detached leaves of air-exposed, Z3HOL-exposed or HIPV-exposed plants. Four early-3rd-instar larvae were used for each replication, with 8 replications in total. The bar represents mean $\pm$ SE (n = 8). * and *** indicate significant difference between treatments at $p < 0.05$, $p < 0.001$, respectively, according to Wilcoxon rank sum test. **B)** Representative differences in amount of leaf material consumed by four 3rd-instar *M. sexta* larvae. Pictures were taken at 6, 12 and 24 h after the release of caterpillars in the tub. **c)** Leaf area consumed by four 3rd-instar *M. sexta* larvae during 24 h feeding period (dual-choice bioassay). The bar represents mean $\pm$ SE (n = 8). * indicates significant differences between treatments at $p < 0.05$ according to Student's *t* test.

Fig. 7 Volatile (Z3HOL or HIPV) exposed plants respond more strongly to simulated herbivory than air-exposed plants. *N. benthamiana* plants were exposed for 48 h to volatile from undamaged plant (air-exposed, gray), 500 ng of Z3HOL (Z3HOL-exposed, blue) or HIPVs from *M. sexta* damaged plants (HIPV-exposed, green). After 48 h of exposure to volatiles, receiver plants were challenged with mechanical damage followed by *M. sexta* regurgitant application. Wounded portions of the leaf were harvested 30 min after challenge for JA analysis (ANOVA, Tukey HSD, n = 4, $p < 0.05$). The bar represents the mean $\pm$ SE. Letters indicate significant differences among different treatments.

Fig. 8 Effect of Z3HOL-exposure on *N. benthamiana* growth and development. A) Plant height (cm), **B)** Average number of secondary branches per plant, **C)** Average length of secondary branch (cm), **D)** Number of seed capsules per plant, e) Seed weight per plant, and f)

seed germination percentage. The bar represents mean ± SE (n = 12). * and ** indicate significant difference between control and Z3HOL-exposure at $p < 0.05$, $p < 0.01$, respectively, according to Student's *t* test.

Supporting Information

Fig. S1 Silencing of the phytoene desaturase gene (PDS) using VIGS in *N. benthamiana*. *N. benthamiana* plants were syringe infiltrated with TRV2 with PDS fragments (TRV::PDS) and photographed 25 d post-infiltration. TRV::PDS inoculated plants had newly emerged leaves with the photo bleached phenotype characteristic of PDS silencing.

Fig. S2 Slow-releasing dispenser designed to release physiologically relevant amount of GLV. A) A mixture of (*Z*)-3-hexenol and lanolin in 1:20 ratio (v/v) inside a GC-insert (25 μL of a mixture per insert), **B)** Slow releasing dispenser, **C)** Volatile exposure set up, **D)** Amount of GLVs (mean ± SE, n = 6) released by *M. sexta* damaged *N. benthamiana* plants at 48 h after feeding initiation, **E)** Amount of individual GLV released from the slow-releasing dispensers (mean ± SE, n = 6).

Fig. S3 Comparison of constitutive volatiles emissions from undamaged wild-type (WT) plants and *M. sexta*-damaged WT plant. Volatiles emitted by undamaged plants were below the detection limit of GC-FID.

Fig. S4 Exposure to (*Z*)-3-hexenal (Z3HAL) alone did not primed *N. benthamiana* plants for enhanced production of monoterpene, sesquiterpene and JA. *N. benthamiana* plants were exposed to volatiles from *M. sexta* damaged WT (green), undamaged WT plants (gray) or synthetic Z3HAL (dark purple) for 48 h. After 48 h of exposure to volatiles, the receiver plants were transferred to individual clean bell jars and challenged repeatedly by mechanical wounding followed by *M. sexta* regurgitant application. The graph shows the total amount of **A)** monoterpenes and **B)** sesquiterpenes emitted from the challenged receiver plants at different time intervals after initial challenge. Values represent mean ± SE (n = 4). Data were analyzed with a mixed model for repeated measures. Different letters indicate significant ($p < 0.05$) differences between treatments within time points, with Bonferroni's correction for multiple comparisons. Shaded areas indicate nighttime volatile collections. **C)** For JA analysis, after 48 h of exposure, receiver plants were challenged with mechanical damage followed by *M. sexta* regurgitant application. Wounded portions of the leaf were harvested 30 min after challenge and JA accumulation in the harvested tissue was analyzed (ANOVA, Tukey HSD, n = 4, $p < 0.05$).

Fig. S5 Exposure to (*Z*)-3-hexenyl acetate (Z3HAC) alone did not primed *N. benthamiana* plants for enhanced production of monoterpene, sesquiterpene and JA. *N. benthamiana* plants were exposed to volatiles from *M. sexta* damaged WT (green), undamaged WT plants (gray) or synthetic Z3HAL (purple) for 48 h. After 48 h of exposure to volatiles, the receiver plants were transferred to individual clean bell jars and challenged repeatedly by mechanical wounding followed by *M. sexta* regurgitant application. The graph shows the total amount of **A)** monoterpenes and **B)** sesquiterpenes emitted from the challenged receiver plants at different time intervals after initial challenge. Values represent mean ± SE (n = 4). Data were analyzed with a

mixed model for repeated measures. Different letters indicate significant ($p < 0.05$) differences between treatments within time points, with Bonferroni's correction for multiple comparisons. Shaded areas indicate nighttime volatile collections. **C)** For JA analysis, after 48 h of exposure, receiver plants were challenged with mechanical damage followed by *M. sexta* regurgitant application. Wounded portions of the leaf were harvested 30 min after challenge and JA accumulation in the harvested tissue was analyzed (ANOVA, Tukey HSD, n = 4, $p < 0.05$).

Fig. S6 Volatile (Z3HOL and HIPV) exposure primed defenses against *M. sexta* larvae. *N. benthamiana* plants were exposed for 48 h to volatile from undamaged plant (air-exposed, gray), 500ng of Z3HOl (Z3HOL-exposed, red) or HIPVs from *M. sexta* damaged plants (HIPV-exposed, green). Values represent mean mortality rate of 3rd-instar *M. sexta* larvae 48 h after releasing them on receiver plants (ANOVA, F = 0.389, n = 6, $p = 0.689$).

Table S1 List of primer sequences used in this study.

Title: Exposure to (*Z*)-3-hexenol primes tobacco plants for faster and stronger defense without negatively affecting their ability to grow and reproduce

Figures

Fig. 1 Experimental design for GLV-mediated defense priming. (A) Priming phase (0-48 h): In the first experiment, receiver plants (R) were exposed to volatiles from *M. sexta*-damaged WT, LOX2-silenced, or GFP-silenced (VIGS-control) emitter plants (E). Receiver plants exposed to undamaged WT plants served as the negative control. In separate experiments, receiver plants were exposed to synthetic GLV [(*Z*)-3-hexenal (Z3HAL), (*Z*)-3hexenol (Z3HOL), or (*Z*)-3-hexenyl acetate (Z3HAC)] via slow-release dispensers. Receiver plants exposed to lanolin dispensers and undamaged WT plants as negative control, while those exposed to *M. sexta*-damaged WT served as positive control. **(B) Post-exposure challenge (0-36 h):** After priming, receiver plants (R) were transferred to clean bell jars and subjected to repetitive wounding every 4 h for 36 h using a cheese grater. Wounds were immediately treated with 10 μL of *M. sexta* regurgitant (1:1 v/v) to simulate herbivory. Volatiles were collected via SuperQ filters.

Fig. 2 HIPVs emitted from *M. sexta* damaged *N. benthamiana* plants. **A)** Chromatographic profiles of HIPVs emitted by *M. sexta* damaged wild type (top), GFP-silenced (middle), and LOX2-silenced (bottom) plants. *N. benthamiana* plants were continuously damaged by fourth-instar *M. sexta* feeding for 48 h. The chromatogram represents HIPV profile collected from 24 to 28 h after feeding initiation. Five major families of HIPVs were induced by *M. sexta* feeding on

N. benthamiana: **green leafy volatiles (GLVs):** (*Z*)-3-hexenal [3], (*E*)-2-hexenal [4], (*Z*)-3-hexenol [5], (*Z*)-3-hexenyl acetate [11], **monoterpenes:** α-pinene [8], sabinene [9], β-pinene [10], limonene [12], 1,8-cineole [13], (*E*)-β-ocimene [14], linalool [15], nonanal [16], **sesquiterpenes:** (*E*)-α-bergamotene [21], unknown sesquiterpene#1 [22], (*E*)-β-farnesene [23], aristolochene (4,4-di epi) [24], β-bisabolene [25], β-sesquiphellandrene [26], **aldoximes:** propyl aldoxime, 2 methyl-,syn-[1], propyl aldoxime, 2 methyl-,anti-[2], butyl aldoxime, 2 methyl-,syn- [6], butyl aldoxime, 2 methyl-,anti-[7], and **aromatic compound**: indole [17].

B) Levels of GLVs emitted by *M. sexta* damaged plants. Early fourth-instar *M. sexta* (one larva per plant) was feeding continuously on the plant for 48 hours. HIPVs collection started on 2nd day, 16 h after feeding initiation and continued up to 48 h in 4-h intervals. Values represent mean ± SE (n = 4). Shaded areas indicate nighttime volatile collections.

Fig. 3 Receiver *N. benthamiana* plants did not get primed for terpene emissions when GLV emission was silenced in emitter plants. *N. benthamiana* plants were exposed to volatiles from *M. sexta* damaged WT (green), LOX2-silenced (dark red), GFP-silenced (blue) or undamaged (gray) plants for 48 h. After 48 h of exposure to volatiles, the receiver plants were transferred to individual clean bell jars and challenged repeatedly by mechanical wounding followed by *M. sexta* regurgitant application. The graph shows the total amount of **A)** monoterpenes and **B)** sesquiterpenes emitted from the challenged receiver plants at different time intervals after initial challenge. Values represent mean $\pm$ SE (n = 3). Data were analyzed with a mixed model for repeated measures. Different letters indicate significant ($p < 0.05$) differences between treatments within time points, with Bonferroni's correction for multiple comparisons. Shaded areas indicate nighttime volatile collections. **C)** For JA analysis, after 48 h of exposure to volatiles, receiver plants were challenged with mechanical damage followed by *M. sexta* regurgitant application. Wounded portions of the leaf were harvested 30 min after challenge and JA accumulation in the harvested tissue was analyzed (ANOVA, Tukey HSD, n = 3, $p < 0.01$).

Fig. 4 Exposure to Z3HOL alone primed *N. benthamiana* plants for enhanced production of mono and sesquiterpenes. *N. benthamiana* plants were exposed to volatiles from *M. sexta* damaged WT (green), undamaged WT plants (gray) or synthetic Z3HOL (red) for 48 h. After 48 h of exposure to volatiles, the receiver plants were transferred to individual clean bell jars and challenged repeatedly by mechanical wounding followed by *M. sexta* regurgitant application. The graph shows the total amount of **A)** monoterpenes and **B)** sesquiterpenes emitted from the challenged receiver plants at different time intervals after initial challenge. Values represent mean ± SE (n = 4). Data were analyzed with a mixed model for repeated measures. Different letters indicate significant ($p < 0.05$) differences between treatments within time points, with Bonferroni's correction for multiple comparisons. Shaded areas indicate nighttime volatile collections.

Fig. 5 Volatile (Z3HOL and HIPV) exposure primed defenses against *M. sexta* larvae. *N. benthamiana* plants were exposed for 48 h to volatile from undamaged plant (air-exposed, gray), 500ng of Z3HOl (Z3HOL-exposed, blue) or HIPVs from *M. sexta* damaged plants (HIPV-exposed, green). **A)** Mortality rate of freshly hatched *M. sexta* neonates 48 h after the neonate release on receiver plants (ANOVA, F = 3.374, n = 8, $p = 0.054$), **B)** Amount of leaf area consumed by two early-3rd-instar *M. sexta* larvae during 24 h feeding period (ANOVA, Tukey HSD, F = 11.27, n = 6, $p < 0.001$), **C)** Larval weight gain at 0 and 7 days after feeding on receiver plants (ANOVA, Tukey HSD, F = 7.84, n = 6, $p < 0.01$). The bar represents the mean ± SE. Letters indicate significant differences among different treatments.

Fig. 6 Feeding preference of *M. sexta* larvae is influenced by volatile exposure treatments.
A) The preference of *M. sexta* larvae on detached leaves of air-exposed, Z3HOL-exposed or HIPV-exposed plants. Four early-3[rd]-instar larvae were used for each replication, with 8 replications in total. The bar represents mean ± SE (n = 8). * and *** indicate significant difference between treatments at $p < 0.05$, $p < 0.001$, respectively, according to Wilcoxon rank sum test. **B)** Representative differences in amount of leaf material consumed by four 3[rd]-instar *M. sexta* larvae. Pictures were taken at 6, 12 and 24 h after the release of caterpillars in the tub. **C)** Leaf area consumed by four 3[rd]-instar *M. sexta* larvae during 24 h feeding period (dual-choice bioassay). The bar represents mean ± SE (n = 8). * Indicates significant differences between treatments at $p < 0.05$ according to Student's *t* test.

Fig. 7 Volatile (Z3HOL or HIPV) exposed plants respond more strongly to simulated herbivory than air-exposed plants. *N. benthamiana* plants were exposed for 48 h to volatile from undamaged plant (air-exposed, gray), 500 ng of Z3HOL (Z3HOL-exposed, blue) or HIPVs from *M. sexta* damaged plants (HIPV-exposed, green). After 48 h of exposure to volatiles, receiver plants were challenged with mechanical damage followed by *M. sexta* regurgitant application. Wounded portions of the leaf were harvested 30 min after challenge for JA analysis (ANOVA, Tukey HSD, n = 4, $p < 0.05$). The bar represents the mean ± SE. Letters indicate significant differences among different treatments.

Fig. 8 Effect of Z3HOL-exposure on *N. benthamiana* growth and development.

A) Plant height (cm), **B)** Average number of secondary branches per plant, **C)** Average length of secondary branch (cm), **D)** Number of seed capsules per plant, **E)** Seed weight per plant, and **F)** seed germination percentage. The bar represents mean ± SE (n = 12). * and ** indicate significant difference between control and Z3HOL-exposure at $p < 0.05$, $p < 0.01$, respectively, according to Student's *t* test.