Hierarchical TBX6-FOXC Regulatory Logic Governs Human Trunk Mesoderm Diversification

John-Poul Ng-Blichfeldt[1,*], Rosalind Drummond[1], David Lando[1], Samina Kausar[1], Melania Barile[1], and Anna Philpott[1,2,*,‡]

[1] Cambridge Stem Cell Institute, University of Cambridge, Cambridge CB2 0AW, UK

[2] Department of Oncology, University of Cambridge, Cambridge CB2 0AH, UK

* Correspondence to:

Dr John-Poul Ng-Blichfeldt jpn36@cam.ac.uk

Professor Anna Philpott ap113@cam.ac.uk

‡ Lead Contact

Highlights

- Human trunk-like structures model mesodermal diversification during early post-gastrulation human trunk development
- TBX6 duration acts as a temporal gate controlling mesodermal fate potential
- FOXC1/2 function downstream of TBX6 to lock in somitic identity and restrict alternative fates
- DegTrace enables lineage tracing following transient transcription factor activation

Summary

Mesoderm forms during gastrulation and diversifies to generate many embryonic
tissues, yet the regulatory logic governing early human mesodermal lineage decisions
remains poorly defined. We use human iPSC-derived 3D trunk-like structures (hTLS) to
model lineage specification during early post-gastrulation development, a stage
experimentally inaccessible in humans. Developing hTLS generate neural and
mesodermal lineages and recapitulate progressive mesodermal diversification into
paraxial (somitic) and intermediate (renal) identities. Using genetic perturbations and
temporally controlled transcription factor activation, we identify duration-dependent
TBX6 activity as a critical determinant of mesodermal diversification. Downstream, the
forkhead box transcription factors FOXC1 and FOXC2 stabilise somitic identity and are
required for sclerotome differentiation. These findings reveal a duration-dependent
hierarchical regulatory logic in early human mesoderm, in which competence for
diversification is established before downstream lineage identities are stabilised,
positioning hTLS as a powerful platform to experimentally dissect human developmental
programs.

Introduction

How transient developmental states are converted into stable lineage identities remains a fundamental problem in developmental biology. Embryonic cell identity begins to diversify prior to and during gastrulation as epiblast cells progressively generate the three embryonic germ layers, with cells that ingress through the primitive streak forming mesoderm and endoderm while non-ingressing cells adopt ectodermal identity [1,2]. Mesoderm further diversifies into distinct lineages that form the primordia of a wide range of tissues, with specific mesodermal identities arising from discrete positions along the rostral-caudal axis of the streak [3]. Emerging from the prospective trunk region of the embryo, paraxial mesoderm undergoes segmentation into somites, transient epithelial structures that further subdivide into sclerotome that forms vertebrae, and dermomyotome that produces skeletal muscle [4]. Adjacent to paraxial mesoderm, intermediate mesoderm gives rise to the urogenital system, including the tubular epithelial components of the kidneys and gonads [5,6]. Despite its fundamental importance, the regulatory logic by which early mesoderm diversifies into distinct lineages, including paraxial and intermediate mesoderm and their derivatives, remains incompletely understood.

Classic fate-mapping and grafting experiments in avian and mouse embryos demonstrated that cells of the primitive streak represent a transient multipotent state characterised by broad equivalence in lineage competence, with positional cues within the streak directing specific mesodermal identities [7-13]. These positional cues are established by signalling gradients that both pattern the embryonic axes and bias lineage specification: Wnt activity posteriorises streak identity and promotes mesodermal differentiation, whereas retinoic acid (RA) antagonises Wnt/FGF signalling to permit differentiation of trunk progenitors while also contributing to posterior neural patterning [14,15]. However, downstream of local signalling environments, how cell-intrinsic transcription factor dynamics convert transient multipotent states into stable lineage-specific mesodermal identities remains poorly defined.

Recent comparative studies have highlighted divergence in developmental

mechanisms between mouse and human mesodermal derivatives, including differences

in morphogenesis and patterning of the kidney [16-19] and in the tempo and signalling

mechanisms underlying somitogenesis [20,21]. However, direct investigation of these early

developmental events in humans is constrained by technical and ethical limitations [22].

Advances in single-cell transcriptomics have provided valuable insights into

transcriptional states associated with early lineage diversification in human embryos, yet

these stages remain experimentally inaccessible *in vivo* [23-25]. Recently, the development

of human pluripotent stem cell–based 3D differentiation systems has begun to enable

functional investigations of early human developmental programs by modelling key

aspects of embryonic development *in vitro* [26-34].

Here, we employ human induced pluripotent stem cell (iPSC)-derived trunk-like

structures (hTLS) to investigate the molecular mechanisms governing early

diversification of paraxial (somitic) and intermediate (renal) mesoderm lineages during

early post-gastrulation human trunk development. By combining temporal control of

transcription factor activity with genetic perturbations, we uncover a previously unknown

regulatory program governing mesodermal diversification. We find that the duration of

upstream TBX6 activity establishes a transient window of multipotent competence,

which is subsequently stabilised by the downstream transcription factors FOXC1 and

FOXC2 to drive lineage commitment. Together, these findings reveal a duration-

dependent hierarchical regulatory logic governing human mesoderm diversification and

demonstrate the utility of hTLS as a tractable experimental system to dissect early

human developmental programs.

Results

Human trunk-like structures (hTLS) model progressive mesodermal diversification

To investigate how trunk mesodermal lineages emerge and diversify during early post-gastrulation human development, we adapted published protocols to generate 3D trunk-like structures (hTLS) from human iPSCs. In this system, iPSCs are exposed to a transient pulse of high Wnt signalling, followed by reduced Wnt signalling combined with RA, prior to aggregation [27]. Under these signalling conditions, neural and mesodermal identities emerge concomitantly, recapitulating lineage diversification characteristic of posterior (post-cranial) embryonic development (Figure 1A) [27]. Differentiating aggregates embedded in Matrigel elongated and generated SOX2$^+$ neural and SOX2$^-$ non-neural tissues in parallel (Figures 1B and S1A,A').

At day 5, SOX2$^+$ structures exhibited lumens and expressed PAX6, consistent with neural tube-like identity (Figure S1B) [35]. Surrounding SOX2$^-$ tissue expressed the paraxial mesoderm marker forkhead box C2 (FOXC2), and the nephron progenitor marker PAX8, indicating the concurrent emergence of both paraxial (somitic) and intermediate (renal) mesodermal identities (Figures 1C and S1B). By day 8, hTLS contained somitic (DACH2$^+$) and renal (PAX8$^+$) epithelial structures with apical-basal polarity, indicated by the apical localisation of ZO1 (Figure 1D). Renal epithelial structures expressed CDH1 (E-Cadherin/ECAD), hepatocyte nuclear factor beta (HNF1B) and PAX8, consistent with nephron tubular identity (Figure S1C). Similar neural, somitic, and renal lineages were observed in hTLS generated from an independent iPSC line (IMR-90-4), demonstrating protocol robustness across different genetic backgrounds (Figure S1D).

To define the transcriptional programs underlying lineage emergence in hTLS, we performed single nucleus RNA-sequencing (snRNA-seq) across a time course from day 0 to day 7, including undifferentiated iPSCs as a control (Figures 1E,E' and S1E-I). Unsupervised clustering revealed *MEOX1*$^+$ somitic mesoderm, *PAX2*$^+$ intermediate mesoderm-derived kidney progenitors and their derivatives (*HNF1B*$^+$ nephron tube and

NPHS1[+] podocytes), as well as neural tube-derived *MAP2*[+] neurons and *SOX10*[+] neural

crest-like populations (Figures S1G-I). By immunofluorescence, cells adjacent to neural

tube-like structures exhibited migratory morphology and were SOX9[+], consistent with a

neural crest-like identity (Figure S1J). *PECAM-1*[+] endothelial cells were also identified

by both snRNA-seq and immunofluorescence (Figures S1G-I and S1K).

To assess the developmental correspondence of hTLS to human embryogenesis

in vivo, we compared the hTLS snRNA-seq dataset to reference atlases spanning

zygote to post-conception week 2 (PCW2/CS7) [25], and to an independent PCW3/CS10

human embryo dataset [24]. hTLS cells from days 0-4 mapped predominantly to CS7-

stage embryonic populations (Figures S1G and SL-L"), whereas cells from days 5-7

showed strong correspondence to PCW3 neural tube, somite, intermediate mesoderm

and endothelial lineages, while lacking notochord, endoderm, cardiac and erythroid

populations (Figures 1F and S1G,H). These analyses support the use of hTLS as an

experimentally tractable model of early post-gastrulation human trunk development,

capturing neural tube, and paraxial (somitic) and intermediate (renal) mesodermal

diversification.

To examine mesodermal diversification in greater detail, we isolated mesodermal

cells in the hTLS snRNA-seq dataset and performed secondary unsupervised clustering

(Figures 1G and S1M-P). Early time points were dominated by a single mesodermal

population enriched for *TBX6*, *FGF10* and *PDGFRA* (Early mesoderm; Figures 1H and

S1P). From day 3 onward, additional clusters emerged corresponding to nephron

progenitors (*PAX8*[+] *CDH6*[+]) and somite progenitors (*MEOX1*[+] *PAX3*[+]), followed by

diversification into nephron derivatives, alongside dorsal somite (*PAX3*[+] *DACH2*[+])-

derived muscle (*NEB*[+]) and mesenchymal (*FAP*[+]) populations at later time points

(Figures 1G,H and S1P). These data indicate that hTLS recapitulate progressive

mesodermal diversification into paraxial (somitic) and intermediate (renal) mesoderm

derivatives, and therefore enable detailed investigation of mechanisms controlling their

divergence.

To reconstruct developmental trajectories underlying mesodermal diversification,

we performed pseudotime inference using Slingshot [36], identifying four major

trajectories corresponding to nephron tube, podocytes, somite-muscle and somite-
mesenchymal lineages (Figure 1J). While cluster identities were used to define the
overall trajectory structure, gene expression dynamics were analysed along pseudotime
at single-cell resolution. Using tradeSeq [37], we identified lineage-specific transcriptional
programs associated with terminal states along each trajectory (Figures 1K,L). Nephron
tube and podocyte lineages exhibited distinct renal epithelial (*PAX2*, *PATJ*) and
glomerular (*WT1*, *MAGI2*) gene signatures, whereas somite-muscle and somite-
mesenchymal lineages were marked by sarcomeric (*TTN*, *NEB*) or extracellular matrix-
associated (*COL23A1*, *SULF1*) gene expression. Together, these analyses establish
hTLS as a system in which trunk mesodermal lineages progressively diversify along
transcriptionally distinct trajectories, providing a foundation for mechanistic interrogation
of human trunk mesoderm specification and diversification.

166 TBX6 is required for establishment of a transient multipotent mesodermal
167 competence state in hTLS

To identify factors regulating the initial establishment of mesodermal multipotency
in hTLS, we compared the earliest emerging mesodermal and neural populations within
the hTLS snRNA-seq dataset. Differential expression analysis identified 27 transcription
factors enriched in early mesoderm, with the T-box transcription factor *TBX6* as one of
the most highly enriched (Figures 2A,A'). Although TBX6 is classically associated with
somitic mesoderm specification, recent work has suggested that its expression
precedes overt lineage diversification [38,39], raising the possibility that TBX6 functions
more broadly to establish mesodermal competence. We therefore examined the
temporal dynamics of TBX6 expression and its functional requirement during early hTLS
differentiation.

We observed by immunofluorescence that TBX6 was expressed at low levels
and heterogeneously in iPSCs after 24 hours of differentiation (day 0), whereas the
primitive streak marker TBXT/Brachyury was broadly expressed (Figures S2A,B).
Notably, TBX6 expression was restricted to a subset of TBXT[+] cells, indicating that
TBX6 marks a subpopulation within the broader TBXT[+] mesodermal compartment. A
subset of TBXT[+] cells also co-expressed SOX2 at early stages (Figures 2B and S2B),

consistent with neuromesodermal progenitor (NMP)-like states that may contribute to
both neural and mesodermal lineages in this system. By day 2 in aggregated hTLS,
TBX6 and SOX2 marked distinct, non-overlapping cell populations, with TBX6
expression declining at day 3 and largely depleted by day 4 (Figure 2B). Time course
RT-qPCR analyses revealed that *TBX6* induction preceded markers of both somitic
mesoderm (*FOXC2*, *MEOX1*, and *PAX3)* and intermediate (renal) mesoderm (*OSR1*,
PAX2, *PAX8*; Figures S2C,D). Thus, consistent with observations in mouse embryos *in*
vivo [38], TBX6 is transiently expressed in mesoderm prior to lineage diversification in
hTLS, suggesting a broader role in establishing mesodermal competence.

We next investigated the role of TBX6 in mesodermal emergence. We
engineered iPSCs with a CRISPR interference (CRISPRi) system to enable inducible
suppression of TBX6 expression (Figures 2C,D) [40]. In this system, a catalytically dead
Cas9 (dCas9) fused to the Krüppel-associated box (KRAB) repressor domain is
targeted to the *TBX6* locus under the control of Doxycycline (Dox), with cytoplasmic
GFP as a readout for dCas9 expression (TBX6-dCas9-KRAB). Addition of Dox at the
onset of differentiation (day -1) resulted in ~82% knockdown of *TBX6* expression at day
0 compared to untreated cells (p = 0.0042; Figure 2E). *TBX6* knockdown was
associated with decreased expression of the early mesodermal markers *TBXT* (p =
0.011) and *MSGN1* (p = 0.0031; Figure S2E). Depletion of TBX6 protein after Dox
treatment was confirmed by immunofluorescence (Figure S2F). Next, TBX6-dCas9-
KRAB iPSCs were differentiated to hTLS and harvested at day 5 of the protocol (Figure
2D). Control hTLS did not express GFP, indicating lack of knockdown cassette
expression, and exhibited somitic (FOXC2$^+$), renal (PAX8$^+$) and neural (SOX2$^+$) tissue
as observed previously (Figure 2F). By contrast, Dox-treated TBX6-dCas9-KRAB hTLS
showed heterogeneous GFP expression with a reduction in both FOXC2$^+$ and PAX8$^+$
tissue and an expanded SOX2$^+$ domain. Remaining FOXC2$^+$PAX8$^+$SOX2$^-$ cells were
GFP$^-$ and thus did not express the knockdown cassette, indicating that renal and somitic
mesodermal differentiation was mutually exclusive with TBX6 depletion (Figure 2F).
Quantification revealed a 78% decrease in the number of FOXC2$^+$ cells per confocal
image (p < 0.0001; Figure 2G), and an 85% decrease in PAX8$^+$ cells (p < 0.0001; Figure
2H), with a corresponding 84% increase in SOX2$^+$ cells (p < 0.0001; Figure 2I). Similar

results were obtained with an independent iPSC line with a distinct gRNA to target
TBX6 (Figure S2G). These data indicate that TBX6 is necessary for establishment of a
multipotent mesodermal population in hTLS, from which both somitic and renal lineages
emerge.

219 Acute TBX6 expression promotes mesodermal transcriptional progression
220 without commitment to a specific lineage

TBX6 is transiently and heterogeneously expressed early in hTLS differentiation
(Figures 2B and S2B). This raised the question of whether TBX6 is permissive for
mesodermal diversification or actively instructs mesodermal progression. To address
this, we engineered a piggyBac-insertable, Dox-inducible TBX6 overexpression system
with cytoplasmic mCherry as a fluorescent readout of transgene induction, to more
uniformly elevate TBX6 expression in the earliest stages of hTLS differentiation (PB-
Dox-TBX6; Figures 3A-C).

After 24 hours of differentiation (day 0 of the protocol), control cells lacked
mCherry expression and formed compact colonies, whereas cells differentiated in the
presence of Dox showed robust mCherry expression and adopted a more dispersed,
mesenchymal-like morphology (Figure 3D). RT-qPCR confirmed strong induction of
TBX6 expression (p<0.0001) and a concomitant reduction in *SOX2* levels (p = 0.0093;
Figure S3A). TBX6 overexpression significantly upregulated markers associated with
both paraxial mesoderm (*FOXC1*, *FOXC2*), and intermediate mesoderm (*OSR1*),
suggesting activation of diverse mesodermal lineage programs (Figure S3A), consistent
with features of an early, multipotent mesodermal state.

To define global transcriptional changes induced by acute TBX6 overexpression,
we performed bulk RNA-seq after 24 hours of Dox treatment (Figures 3B,C and S3B,C).
TBX6 induction resulted in widespread transcriptional rewiring, including downregulation
of genes characteristic of pluripotency (*NANOG*), early mesendoderm (*EOMES*), and
alternative lineage programs (endoderm regulator *FOXA2*), alongside upregulation of
genes associated with differentiated mesodermal states (Figure 3E). These included
somitic and paraxial mesoderm (*FOXC1, DLL1, ALDH1A2*), intermediate mesoderm
(*OSR1, LHX1*) and cardiac/muscle lineages (*MYL7, NRAP*) (Figure 3E). Consistent with

this, gene ontology (GO) analysis revealed enrichment of biological processes related to
mesodermal differentiation and morphogenesis, including somite, renal nephron, and
cardiac lineages (Figure S3D).

Gene set enrichment analysis (GSEA) using a reference human PCW2
embryonic atlas [23] demonstrated selective enrichment of differentiated mesodermal
transcriptional programs following TBX6 induction, with highest enrichment for
emergent, advanced and axial mesoderm signatures, alongside depletion of epiblast
and primitive streak signatures (Figure 3E). In a PCW3 reference atlas [24], TBX6-
induced programs aligned with multiple distinct mesodermal lineages, including paraxial
and intermediate mesoderm, while neural programs were depleted (Figure S3E).
Reciprocal analysis, in which a TBX6-induced gene module was mapped onto the
PCW2 and PCW3 human embryonic atlases, further supported these findings (Figures
S3F-M). Together, these data indicate that acute TBX6 overexpression is sufficient to
promote transcriptional progression beyond early mesoderm toward differentiated
mesodermal states, but does not enforce commitment to a single mesodermal lineage.

Duration of TBX6 activity determines commitment to somitic or alternative
mesodermal fates

Previous studies in mouse embryonic stem cells suggested that the duration of
TBX6 expression can influence mesodermal differentiation [39]. Since acute TBX6
induction activates transcriptional programs associated with multiple mesodermal
lineages, we next tested whether altering the duration of TBX6 activity would influence
mesodermal lineage commitment in hTLS. We first investigated the consequences of
sustained TBX6 overexpression by differentiating PB-Dox-TBX6 cells to hTLS in the
continuous presence of Dox (Figures 4A,B). As a control, iPSCs harbouring Dox-
inducible mCherry alone were differentiated to hTLS (PB-Dox-mCherry; Figures 4A,B).
At day 5, hTLS generated from both PB-Dox-TBX6 and PB-Dox-mCherry cells exhibited
numerous mCherry$^+$ cells, indicating sustained transgene expression. This suggests
that the piggyBac-based Dox-inducible system remained unsilenced in these cells, likely
due to the presence of multiple insulator sequences (Figures 4B-D).

At day 5 in hTLS generated from PB-Dox-TBX6 cells, mCherry[+] cells

predominantly expressed the somitic marker FOXC2, lacked expression of PAX8 and

SOX2, and adopted a mesenchymal morphology, indicating restriction toward a somitic

mesoderm fate (Figure 4C). By contrast, hTLS generated from PB-Dox-mCherry cells

exhibited mCherry expression within epithelial structures across somitic, renal, and

neural compartments (Figure 4D), confirming that sustained TBX6 activity specifically,

and not Dox-inducible transgene expression per se, drives a specific somitic lineage

bias.

Since TBX6 activity is transient both *in vivo* and during hTLS development

(Figures 2B and S2B) [38], we next tested whether restricting the duration of ectopic TBX6

activity would enable mesodermal diversification. To precisely control TBX6 protein

stability while retaining a record of prior TBX6 expression, we utilised a FK506 binding

protein (FKBP) domain fused to TBX6 to develop a conditional degradation and lineage-

tracing system that we termed DegTrace (Figures 4E,E') [41]. In this system, Dox induces

transcription of the TBX6-FKBP-T2A-mCherry transgene, whereas subsequent addition

of dTAG selectively depletes TBX6 protein by engaging the fused FKBP domain with E3

ubiquitin ligases for proteasomal degradation [41], without affecting the co-translated

mCherry reporter, thus enabling lineage tracing of cells that have transiently expressed

TBX6 (Figures 4E,E'). We validated DegTrace in iPSCs, confirming that dTAG-

mediated degradation selectively depleted TBX6 protein while preserving mCherry

expression (Figures S4A,B).

We next applied DegTrace to 3D trunk differentiation. TBX6-FKBP-T2A-mCherry

transgene overexpression was induced with Dox from day -1, and TBX6 protein

degradation was triggered by dTAG addition at day 0 (after 24 h), day 1 (48 h), or day 2

(72 h) (Figures 4E,E'). RT-qPCR for the mCherry transcript showed that Dox alone

induced transgene expression irrespective of subsequent dTAG addition ($p < 0.001$

versus control; Figure 4F). Sustained TBX6 overexpression (Dox alone) robustly

induced the somitic mesoderm markers *FOXC1* and *MEOX1* (both $p < 0.01$ versus

control; Figure 4F). By contrast, restricting TBX6 activity to the first 24 h led to

diminished expression of these somitic markers, whereas inducing degradation after 48

or 72 h of TBX6 activity resulted in partial marker retention (Figure 4F).

Consistent with these transient transcriptional effects, immunofluorescence

analysis revealed that cells exposed to transient TBX6 activity (24 h) failed to maintain

mesodermal identity and instead adopted a SOX2$^+$PAX3$^+$ neural epithelial fate (Figures

4G,H and S4C,D), whereas cells exposed to an intermediate duration of TBX6 activity

(72 h) differentiated into mesodermal epithelial structures expressing both the somitic

mesoderm marker FOXC2 and the renal marker PAX8 (Figure 4H). By contrast,

sustained TBX6 overexpression promoted mesenchymal differentiation and restriction

to a FOXC2$^+$PAX8$^-$ somitic fate (Figures 4C,G).

Together, these findings indicate that a minimum duration of TBX6 activity is

needed to suppress neural identity and promote mesodermal identity. An intermediate

duration enables acquisition of a multipotent epithelial trunk mesoderm state with

competence for intermediate (renal) mesoderm differentiation. By contrast, prolonged

TBX6 activity biases cells toward a somitic mesoderm fate and reduces competence for

alternative mesodermal lineages. This is accompanied by loss of epithelial

characteristics, suggesting progression beyond an epithelial somite-like state toward a

more mesenchymal somite derivative. Notably, the intermediate duration of exposure

closely mirrors the endogenous dynamics of TBX6 expression observed in hTLS

(Figures 2B and S2B,D).

FOXC1 and FOXC2 establish and safeguard somitic mesoderm identity

Having established that TBX6 duration governs mesodermal fate output, we next

sought to identify downstream mechanisms that stabilise specific mesodermal identities.

TBX6 induction alone biased differentiation toward somite-like fates in hTLS, so we

sought to investigate downstream transcriptional regulators of the somitic lineage.

Analysis of the hTLS snRNA-seq mesodermal subset (Figure 1G) identified a somitic

progenitor population enriched for canonical somitic transcription factors including

FOXC1, *FOXC2*, *PAX3*, *TCF15* (Paraxis), *SIX1*, *MEOX1* and the Hedgehog pathway
mediator *GLI2* (Figures 5A,B).

We focussed on investigating the function of FOXC1 and FOXC2 in human
embryoids based on prior genetic evidence that combined loss of these factors in mice
disrupts somite formation and leads to expansion of intermediate (renal) mesoderm [42].
As prior mouse studies indicated partial functional redundancy between FOXC1 and
FOXC2 in early development [42-44], we engineered a dual knockdown system using
dCas9-KRAB, hereafter referred to as FOXC1/2-dCas9-KRAB (Figures 5C and S5C).

To examine the role of FOXC1/2 specifically in somitic mesoderm differentiation,
we adapted a published 2D human somitic mesoderm differentiation protocol [45] that
recapitulates progression from primitive streak through paraxial mesoderm to an early
somite-like state (Figures 5D and S5A,B). Dual knockdown of *FOXC1* and *FOXC2* was
induced at the primitive streak stage, and cells were harvested at the early somite stage
for bulk RNA-seq (Figures 5D and S5D,E).

Dual FOXC1/2 knockdown efficiently suppressed expression of both factors and
was accompanied by a robust decrease in canonical somitic markers, including
RIPPLY1, *PAX3*, *TCF15, HAND1* and *TWIST1* (Figure 5E). Concomitantly, dual
FOXC1/2 knockdown induced genes associated with alternative mesodermal lineages.
These included intermediate (renal) mesoderm (e.g. *NPHS1, EYA2, JAG1),* lateral plate
mesoderm (*FOXF1, BMP2, BMP7*) and cardiomyocyte/smooth muscle-associated
genes (e.g. *TNNT2*, *CNN1*, *MYH11, MYH7B*; Figures 5E and S5F). Primitive streak
(*TBXT*, *CDX2*) and paraxial mesodermal genes (*TBX6, MSGN1*) were unchanged or
upregulated, indicating that dual FOXC1/2 knockdown did not prevent establishment of
paraxial mesoderm but instead impaired progression from paraxial mesoderm to an
early somite state (Figure S5G). GO analysis confirmed depletion of patterning and
skeletal development programs, with concomitant enrichment of contractile, heart, and
renal system processes (Figure S5H).

To better understand how FOXC1/2-dependent genes correspond to human

embryonic programs *in vivo*, we performed GSEA using PCW3-5 human embryonic

reference atlases [24]. This analysis revealed that FOXC1/2 knockdown led to depletion of

signatures associated with somitic mesoderm derivatives and enrichment of alternative

lineage programs, including intermediate (renal) mesoderm and cardiomyocyte

signatures (Figure S5I). Moreover, GSEA using independent human embryonic heart [46]

and kidney [47] reference atlases further demonstrated that genes upregulated upon

FOXC1/2 knockdown were enriched for cardiomyocyte (Figure S5J) and renal nephron

tube and podocyte signatures (Figure S5K). Together, these results indicate that

FOXC1 and FOXC2 are required to establish somite transcriptional identity and to

restrict alternative mesodermal differentiation programs.

To investigate mechanisms regulating FOXC1/2-dependent gene expression, we

used published ATAC-seq profiles from the equivalent early somite population

generated with the same protocol [45]. Motif enrichment analysis was performed on

accessible chromatin regions adjacent to genes we found to be downregulated following

dual FOXC1/2 knockdown, using accessible regions associated with expressed genes

that were not FOXC1/2-dependent as background. This analysis revealed enrichment of

forkhead, homeobox, and basic helix-loop-helix (bHLH) motifs within FOXC1/2-

dependent regulatory regions, suggesting that FOXC1 and FOXC2 act within a broader

regulatory landscape of combinations of transcriptional regulators rather than as solitary

lineage-determining factors (Figure 5F).

To test whether FOXC1/2 regulate somite identity in hTLS, we induced dual

knockdown during hTLS differentiation (Figure 5G). While neural and renal tissues were

preserved, PAX3$^+$SOX2$^-$ somitic tissue was markedly reduced (68% reduction

compared to control; p < 0.0001), demonstrating a specific requirement for FOXC1/2 in

establishing somitic mesoderm identity in hTLS (Figures 5H-K and S5L).

Finally, to determine whether TBX6 function relies on FOXC1/2 and thereby

define their epistatic relationship, we generated iPSCs harbouring both a Dox-inducible

TBX6 overexpression cassette and the FOXC1/2-dCas9-KRAB knockdown system

(Figure S5M). TBX6 overexpression alone induced multiple somitic genes, including
TWIST1, *TWIST2*, *MEOX1*, *MEOX2* and *TBX18*, however, this induction was abolished
upon simultaneous FOXC1/2 knockdown, indicating that TBX6-driven somitic
differentiation depends on FOXC1/2 activity (Figure S5N). By contrast, *MESP2,*
previously described as a TBX6-responsive gene [39] was unaffected by FOXC1/2
knockdown, suggesting a parallel FOXC-independent TBX6 pathway (Figure S5N).

Together, these data indicate that mesodermal diversification is regulated by the
duration of TBX6 activity, while FOXC transcription factors function downstream of
TBX6 to constrain mesodermal plasticity and establish somitic identity.

396 FOXC1 and FOXC2 stabilise somitic identity and preserve competence for further
397 diversification

Having found that FOXC1/2 are required to establish somite mesoderm identity,
we next asked whether they are additionally needed to maintain somite identity. To
investigate this, we differentiated FOXC1/2-dCas9-KRAB cells to early somite stage and
then induced dual knockdown (Figure 6A). Dual FOXC1/2 knockdown alone led to a
reduction in somite markers *TCF15*, *TBX18* and *MEOX2,* whereas *MEOX1* and *SIX1*
were unchanged, suggesting that FOXC1/2 loss partially destabilised the early somite
program (Figure 6B).

We next tested whether dual knockdown affected the ability of early somite
cultures to differentiate into sclerotome, a transient embryonic structure that gives rise to
the vertebral skeleton. Studies of avian and mouse development have established that
sclerotome differentiation *in vivo* is triggered in the ventral somite by Hedgehog ligands
derived from the notochord and floorplate [48,49]. This process is characterised by
repression of PAX3, upregulation of FOXC1/2, and induction of downstream markers
including TWIST1, PAX1, PAX9, and SOX9 together with an epithelial-to-mesenchymal
transition (EMT) (Figure S6A).

We adapted a 2D human somite-to-sclerotome differentiation protocol in which early somite cultures were exposed to smoothened agonist (SAG) to activate Hedgehog signalling and trigger sclerotome differentiation (Figure 6A) [45]. SAG treatment resulted in efficient generation of sclerotome characterised by suppression of *PAX3* and upregulation of *FOXC1*, *FOXC2*, *TWIST1*, *PAX9*, *PAX1* and *SOX9* compared to untreated cells (Figures 6A,C,D).

Dual FOXC1/2 knockdown induced at the early somite stage, in combination with SAG treatment, strongly attenuated induction of sclerotome markers (Figures 6C,D), whereas induction of canonical Hedgehog pathway targets *PTCH1* and *GLI1* by SAG was unaffected by FOXC1/2 knockdown (Figure 6D). These data indicate that FOXC1/2 are not necessary for Hedgehog signalling per se, but instead support maintenance of somite identity and thereby preserve competence for sclerotome differentiation, or indeed may play a more direct role in sclerotome differentiation.

As FOXC1/2 knockdown inhibited the establishment and maintenance of somite identity, we next asked whether they are sufficient to promote somite differentiation when overexpressed in hTLS. Given the functional redundancy of FOXC1 and FOXC2 *in vivo* and in hTLS, we focussed on FOXC1 and generated a Dox-inducible FOXC1 overexpression line (PB-Dox-FOXC1; Figure 6E). FOXC1 overexpression was induced at day 2 of differentiation to match the timeframe of the previous FOXC1/2 knockdown experiments in hTLS (Figures 5G-K and 6F).

RT-qPCR analysis of 2D-cultured hTLS generated with PB-Dox-FOXC1 cells revealed that FOXC1 overexpression robustly upregulated *TWIST1* (~2-fold; $p < 0.01$), a bHLH transcription factor associated with ventral somite identity, sclerotome differentiation and EMT (Figure 6G). FOXC1 overexpression also upregulated the ventral somite marker *MEOX1* (~1.4-fold; $p < 0.05$), but only modestly induced *PAX9* and *TWIST2* and failed to induce *SOX9* (Figure 6G).

Overexpression of a FOXC1 variant lacking the forkhead DNA-binding domain (FOXC1ΔFKH) failed to upregulate *TWIST1,* even at the highest Dox concentration (1 µM) despite robust transgene induction, indicating that *TWIST1* induction is specific to transcriptionally active FOXC1 (Figure 6H).

To investigate the effects of FOXC1 overexpression in a context more closely
resembling native embryonic tissue, PB-Dox-FOXC1 cells were differentiated to 3D-
hTLS, and FOXC1 overexpression was induced from day 2 of hTLS differentiation
(Figure 6I). Robust, heterogeneous mCherry induction was observed in elongating
hTLS and maintained through days 3-5 (Figure 6J). FOXC1-overexpressing cells were
largely excluded from the epithelial cysts that formed and instead occupied a sub-
epithelial position, consistent with acquisition of a mesenchymal-like state (Figure 6K).
By contrast, FOXC1ΔFKH-overexpressing cells were observed within epithelial cysts,
indicating that FOXC1 transcriptional activity is needed to drive the mesenchymal
phenotype (Figure 6N).

Cells overexpressing wild type FOXC1 (FOXC1::mCherry[+]) within hTLS did not
express SOX2, PAX3, or PAX8, indicating that FOXC1 overexpression does not
promote neural, dorsal somite, or renal fates (Figures 6L,M). Moreover, FOXC1-
overexpressing cells robustly expressed TWIST1 (Figure 6M) but rarely co-expressed
the sclerotome marker SOX9 (Figure S6H), suggesting that FOXC1 is insufficient to fully
execute the sclerotome differentiation program. Overexpression of FOXC2 yielded a
similar phenotype, with sub-epithelial localisation of overexpressing cells and absence
of SOX2 and PAX8 expression (Figure S6I).

Consistent with this, snRNA-seq analysis of untreated hTLS did not reveal a
sclerotome population (Figures S1G-J), despite expression of the Hedgehog pathway
mediators *GLI2* and *GLI3* (Figures S6A,B). This likely reflects the absence of notochord-
and floorplate-derived Hedgehog ligands in this system (Figures S6A,B). To test
whether hTLS retain competence for sclerotome differentiation, we activated Hedgehog
signalling with SAG from day 5 to day 8 of differentiation (Figure S6C). SAG treatment
led to expansion of FOXC2[+] tissue and depletion of PAX3[+] dorsal somitic domains,
consistent with somite ventralisation and sclerotome differentiation (Figures S6D-G).

Together, these results indicate that FOXC1 and FOXC2 are required to establish
and stabilise somitic mesoderm identity and to enable somite-to-sclerotome
differentiation. Moreover, while FOXC1 overexpression is sufficient for *TWIST1*

induction and EMT-like behaviour, full sclerotome differentiation requires additional cues

such as Hedgehog signalling.

TWIST1 is required for Hedgehog-driven sclerotome differentiation

Because our ATAC-seq analysis suggested that FOXC factors co-operate with

bHLH factors to establish the somitic transcriptional landscape, we hypothesised that a

bHLH factor might act as the direct effector of FOXC1/2. FOXC1 overexpression

upregulated the bHLH factor TWIST1 and drove EMT in hTLS, and TWIST1 has known

roles in inducing EMT [50,51], positioning TWIST1 as a plausible candidate bHLH

functioning downstream of FOXC1. We edited iPSCs to harbour a Dox-inducible

TWIST1 overexpression cassette and differentiated them to hTLS (Figures S7A,B).

Inducible TWIST1 overexpression led to sub-epithelial localisation of TWIST1::mCherry$^+$

cells and exclusion from epithelial cysts, closely resembling the FOXC1 overexpression

phenotype (Figure S7C), suggesting that TWIST1 may function downstream of FOXC1

to drive EMT in this system.

Finally, as FOXC1 and FOXC2 are necessary for sclerotome differentiation, we

asked whether TWIST factors are also required. Dual knockdown of TWIST1 and

TWIST2 during SAG-induced differentiation (Figure S7D) strongly reduced SAG-

induced expression of sclerotome markers, but did not affect induction of *FOXC1* or

FOXC2 by SAG (Figure S7E). Moreover, SAG-induced expression of *PTCH1* and *GLI1*

was unaffected by TWIST1/2 knockdown (Figure S7E). Together, these data suggest

that FOXC factors may promote EMT via TWIST factors, and support a model in which

TWIST factors do not affect Hedgehog signalling per se, but instead act as necessary

effectors of Hedgehog-driven sclerotome differentiation in human embryonic mesoderm.

Discussion

How lineage identities first emerge and diversify during early human embryogenesis remains a fundamental unresolved question, largely owing to the experimental inaccessibility of post-gastrulation human development *in vivo*. While recent single-cell atlases have mapped transcriptomic states during early human development, experimental dissection of the mechanisms governing lineage diversification during this window has remained limited. To address this knowledge gap, we use an iPSC-derived hTLS embryoid model, combined with genetic tools that enable precise temporal control of transcription factor activity with dual gene knockdown, to uncover a hierarchical regulatory logic underlying early human mesodermal diversification. Our data support a model in which TBX6 functions as a "temporal gate" governing mesodermal diversification, with the duration of TBX6 activity determining intermediate (renal) versus somitic mesodermal fates (Figure 7). Downstream of TBX6, FOXC1/2 act as a "lock" to establish and stabilise somitic mesoderm identity, preserving the ability of somite cells to undergo sclerotome differentiation in response to Hedgehog signalling, in part through induction of TWIST factors. Given that many developmental transcription factors are transiently expressed, our findings suggest that the duration of transcription factor activity may act as a temporal parameter that helps define competence windows to hierarchically restrict lineage plasticity during embryogenesis.

We benchmarked our embryoid model against human *in vivo* embryonic reference atlases, validating hTLS as an experimentally tractable model that recapitulates key aspects of post-gastrulation human embryonic development. The presence of paraxial (somitic) and intermediate (renal) mesoderm in our hTLS model enabled investigation of the logic underpinning the emergence of these distinct mesodermal subtypes. Fate-mapping of peri-gastrulation avian and mouse embryos showed that mesodermal subtypes including paraxial and intermediate mesoderm arise from discrete positions along the rostral-caudal axis of the primitive streak [7-10,13]. Moreover, rostrally-located streak cells fated to become head mesenchyme and anterior somites would adopt lateral fates including intermediate and lateral plate mesoderm when grafted caudally [7], indicating developmental equivalence within a multipotent

cellular pool. While morphogen gradients including BMP and FGF pattern the
mediolateral axis of the mesoderm [3,52], specific transcription factors that stabilise
lineage identity remain poorly defined. Our finding that FOXC1/2 promotes commitment
to somitic mesoderm and restricts alternative fates including intermediate mesoderm
provides a molecular basis for mesodermal fate commitment likely operating
downstream of local microenvironmental cues. Our epistasis experiments, and our
finding that *FOXC1* and *FOXC2* are induced by acute TBX6 overexpression, position
FOXC1/2 functionally downstream of TBX6, either through direct chromatin-level
regulation or via intermediate factors.

Our DegTrace system enabled precise control over the duration of transcription
factor activity while simultaneously tracing the fate of modified cells, allowing us to
experimentally interrogate how transcription factor dynamics influence lineage
outcomes. Using this system, we uncovered a duration-dependent TBX6 regulatory
logic underlying mesodermal diversification. TBX6 is classically described as a critical
regulator of somitic mesoderm specification. TBX6 null mice lack somitic mesoderm and
instead exhibit three neural tubes [53,54], and TBX6 has been proposed to repress the
SOX2 locus to permit somitic mesoderm differentiation in NMPs [54]. Accordingly, recent
human population genetics studies have revealed *TBX6* variants to be associated with
congenital vertebral malformations, reflecting the somitic origin of vertebrae [55-57]. Our
finding that FOXC1/2 act downstream of TBX6 to establish and stabilise somitic
mesoderm identity provides a mechanistic explanation for these defects. Consistent
with this, *FOXC1* mutations in humans can cause Axenfeld-Rieger syndrome, which
includes skeletal abnormalities such as scoliosis in addition to ocular defects [58,59].

Our observation that *TBX6* knockdown depletes both kidney and somite
mesoderm is consistent with a model in which a multipotent mesodermal population
expressing *TBX6* gives rise to both intermediate and paraxial mesoderm during post-
gastrulation development. Studies of early mouse development mapped initial *Tbx6*
expression broadly throughout the primitive streak, becoming progressively more
posteriorly restricted as development proceeds [38]. A multipotent TBX6[+] progenitor of
intermediate and somitic mesoderm is supported by lineage tracing studies of *Tbx6-*

expressing cells in mice, which showed labelling of somitic and intermediate mesoderm
when traced at early stages, and labelling restricted to somites if traced later [60]. These
observations support a model in which longer durations of TBX6 activity bias cells
toward somitic fate, whereas shorter exposure permits alternative mesodermal
outcomes. Supporting this broader role, *TBX6* variants have been associated with
defects in intermediate mesoderm derivatives including the kidneys, in both human
genetic studies and mouse models [61-66]. Together with these studies, our findings
position TBX6 as a potential regulator of intermediate mesoderm development in
addition to its known role as a somitic mesoderm specifier.

The duration-dependent effects of TBX6 activity revealed by DegTrace are
consistent with a recent report showing that a defined pulse of TBX6 induction
enhances cardiomyocyte differentiation efficiency in pluripotent stem cell protocols [39].
These observations raise the question of how the activity duration of a single
transcription factor can encode competence for divergent lineage outcomes. One
possibility is that prolonged TBX6 exposure permits accumulation of downstream
cofactors that modify transcriptional output, whereas transient exposure fails to stabilise
these networks. Additionally, mutually repressive interactions may occur between
downstream targets as has recently been suggested for *Osr1* and *Foxc2* in mouse [66].
Dissecting the chromatin and network dynamics underlying this temporal encoding will
be an important direction for future work. The relative resistance of our piggyBac system
to transgene silencing in many cells enabled inducible, sustained TBX6 activity,
highlighting its utility for interrogating temporal transcription factor dynamics in
differentiating systems, and offers advantages over recently described non-inducible
transgene expression systems [67].

Our findings support a model in which FOXC1 and FOXC2 establish and stabilise
somite identity in part by restricting alternative lineage programs. Comparison with
human embryonic reference atlases revealed that dual FOXC1/2 knockdown caused
loss of canonical somite markers alongside induction of intermediate mesoderm and
cardiomyocyte/smooth muscle-associated gene programs. This is consistent with a shift
toward alternative mesodermal fates when somite development is impaired, suggesting

that FOXC1/2 help restrict, and may actively repress, intermediate and lateral plate
mesoderm identities during development. In support of this, dual FOXC1/2 knockdown
upregulated *FOXF1,* a lateral plate mesoderm marker *in vivo* [68], and *BMP2* and *BMP7,*
which regulate lateral plate mesoderm development [69]. Our findings contrast with prior
studies of *Foxc1/2* double knockout mice which exhibited loss of both somitic and
cardiac lineages [44], suggesting potential divergence in FOXC1/2 function between
human and mice and highlighting the importance of using human-specific models,
including hTLS, to study early human developmental mechanisms.

Notably, FOXC1 overexpression alone was not sufficient to drive somite
differentiation. Our analysis of published ATAC data of early somite cultures revealed
that FOXC motifs were co-enriched with bHLH and homeobox factors specifically at
FOXC1/2-dependent genes in open chromatin compared to non-dependent genes. It is
therefore likely that FOXC1 acts in a broader combinatorial regulatory framework to
establish somite identity that may require cooperativity with bHLH transcription factors.
We found that potential FOXC1/2 targets included bHLH factors such as *TCF15* and
HAND1 and homeobox factors including *MEOX1*, which could function as cooperative
partners with FOXC1/2 in somite differentiation. Moreover, FOXC1 overexpression
drove induction of the bHLH factor *TWIST1* and EMT-like behaviour. Thus, FOXC1-
TWIST1 cooperativity may have a specific role in regulating the acquisition of a
migratory, mesenchymal phenotype during sclerotome differentiation. However, *FOXC1*
expression is not sufficient to drive full sclerotome differentiation, and additional cues
such as Hedgehog signalling are needed. The established role of FOXC1 in cancer-
associated EMT [70,71] suggests that this developmental function in promoting
mesenchymal transition may be co-opted in pathological contexts.

In summary, our study reveals a duration-dependent hierarchical regulatory logic
governing human mesodermal diversification and highlights the utility of iPSC-derived
embryoid models, combined with transgene overexpression and genetic perturbations,
for gaining insights into otherwise inaccessible stages of early human embryogenesis.

Figure Legends

Figure 1 - Human trunk-like structures recapitulate progressive mesodermal diversification

(A) Schematic of the protocol used to generate human trunk-like structures (hTLS) from induced pluripotent stem cells (iPSCs), adapted from [27].

(B) Brightfield and corresponding green fluorescence images of hTLS generated from SOX2:mNeon iPSCs from day 1 to day 5 of the protocol, showing expansion of mNeon⁺ and mNeon⁻ domains. Scale bars = 300 µm.

(C and D) Immunofluorescence of hTLS showing (C) FOXC2 (magenta), SOX2 (yellow), PAX8 (cyan) expression at day 5, with DAPI as nuclear counterstain (grey), or (D) DACH2 (magenta), SOX2 (yellow), PAX8 (cyan), and ZO1 (grey) expression at day 8. hTLS are outlined with dotted white lines. Scale bars: main, 100 µm; insets, 20 µm.

(E) Schematic of the hTLS single-nucleus RNA-sequencing (snRNA-seq) strategy. (E') UMAP representation of 13,618 hTLS cells coloured by time point, with major lineages annotated (see Supplemental Figure 1G,H). PS, pluripotent stem cells.

(F) Heatmap showing transcriptional similarity between clusters identified from hTLS snRNA-seq (days 5-7) and clusters from post-conception week 3 (PCW3) human embryo scRNA-seq data [24].

(G) UMAP representation of 8,088 cells of hTLS mesodermal cells identified from the snRNA-seq dataset (see Supplemental Figure S1M, N), coloured by unsupervised clustering and annotated based on marker expression.

(H) Alluvial plot showing the proportion of hTLS mesodermal cells at each time point assigned to clusters defined in (G).

(I) Expression of representative marker genes for clusters in (G) overlaid on UMAP plots.

(J) UMAP representation of hTLS mesodermal cells (as in G) with pseudotime trajectories inferred using Slingshot [36].

(K) Within-lineage standardized expression dynamics of the top 20 genes defining each
terminal state along the pseudotime trajectories shown in (J), identified using tradeSeq
endpoint differential expression [37]. Genes are ranked by the strength and consistency of
their differential expression at lineage endpoints. Heatmaps show z-scored fitted
expression along pseudotime within each lineage (red, higher; blue, lower).

(L) Expression of representative genes identified in (K) (*MECOM, WT1, SULF1* and
TTN) plotted along pseudotime for each lineage. Individual cells are shown as dots, with
fitted expression curves shown as lines and coloured by lineage.

See also Figure S1.

Figure 2 - TBX6 defines early mesoderm and is critical for its diversification

(A) Schematic of bifurcation of caudal epiblast cells into mesodermal and neural-fated
cells (coloured in UMAP representation of 13,618 hTLS cells underneath).
(A') Genes differentially expressed between mesoderm and neural clusters (as in A).
Genes with adjusted p-value < 0.05 and |log2 fold change| > 0.58 (±1.5-fold) are
highlighted (red, upregulated; blue, downregulated). Genes encoding the top 10 up- and
downregulated transcription factors, ranked by log2 fold change, are shown in darker
colours and annotated.

(B) Immunofluorescence showing TBX6 (green), TBXT/Brachyury (yellow), and SOX2
(magenta) expression in hTLS at days 2, 3 and 4, with nuclei labelled with DAPI (blue).
Putative neuromesodermal progenitors (NMP)-like cells (SOX2$^+$, TBXT$^+$, TBX6$^-$) are
outlined with a dotted line. Arrowhead indicates TBX6$^+$ mesodermal domain showing
progressive loss of TBX6 expression. Scale bars, 100 μm.

(C) Schematic of the dCas9-KRAB CRISPR interference (CRISPRi) gene knockdown
system [40]. Created with BioRender.com.

(D) Schematic of experimental protocol to suppress *TBX6* expression in hTLS.

(E) RT-qPCR for *TBX6* in TBX6-dCas9-KRAB iPSCs differentiated in the hTLS protocol
and harvested after 24 hours, either untreated (control) or treated with Dox (1 µM) from
day -1. Statistical significance was assessed using two-way ANOVA on log-transformed
values, with treatment as the column factor and differentiation batch as the row factor; p
value corresponds to the main effect of treatment (** $p < 0.01$). Data are plotted as $2^{-\Delta\Delta Ct}$
for visualisation; bars represent mean ± SEM; dots represent n = 4 independent
differentiation batches per group.

(F) Immunofluorescence of GFP (green), SOX2 (yellow), FOXC2 (magenta), or PAX8
(grey) expression in day 5 hTLS generated from TBX6-dCas9-KRAB iPSCs, either
untreated (control, top) or treated with Dox from day -1 (1 µM, bottom). White brackets
indicate SOX2$^-$, FOXC2$^+$, PAX8$^+$ cells confined to a GFP$^-$ region. Scale bars, 100 µm.

(G-I) Quantification of number of nuclei per confocal image positive for (G) FOXC2, (H)
PAX8, or (I) SOX2, in day 5 TBX6-dCas9-KRAB hTLS either untreated (control, white
bars; n=20 hTLS across 2 independent batches) or treated with Dox from day -1 (1 µM,
coloured bars; n=22 hTLS across 2 independent batches). Dots represent individual
hTLS; bars represent median ± IQR; Mann-Whitney test.

See also Figure S2.

Figure 3 - TBX6 drives transcriptional progression toward advanced mesodermal
fates

(A) Schematic of a piggyBac-insertable construct for doxycycline (Dox)-inducible
overexpression of TBX6-FKBP-T2A-mCherry.

(B) Inducible TBX6 transgene overexpression strategy. A piggyBac donor plasmid
encoding a Dox-inducible TBX6-FKBP-T2A-mCherry cassette with puromycin
resistance was integrated into iPSCs using piggyBac transposase. Following selection,
edited cells were differentiated to hTLS with or without Dox to induce transgene
expression.

(C) Experimental design for TBX6 overexpression during the first 24 hours of the hTLS protocol in piggyBac-edited iPSCs (PB-Dox-TBX6), followed by bulk RNA-seq.

(D) Brightfield and corresponding red fluorescence images of PB-Dox-TBX6 cells 24 hours after initiation of the hTLS protocol, either without Dox (control) or with Dox (0.1 µM). Scale bars, 300 µm.

(E) Volcano plot of differential gene expression following 24 h of TBX6 overexpression versus control (n = 4 independent differentiation batches per condition). Genes with adjusted p-value < 0.05 and |log2 fold change| > 0.58 (±1.5-fold) are highlighted (red, upregulated; blue, downregulated). Selected genes associated with alternative lineages are annotated and shown in darker colour. (Somitic: *FOXC1, DLL1, ALDH1A2*; intermediate (renal) mesoderm: *OSR1, LHX1*; cardiac/muscle lineages: *MYL7, NRAP*; pluripotency: *NANOG*; mesendoderm/endoderm: *EOMES, FOXA2*).

(F) Ridgeplots of gene set enrichment analysis (GSEA) of genes following TBX6 overexpression (Dox versus control), ranked by the DESeq2 Wald statistic [72], and tested against the top 50 marker genes for each annotated PCW2 human embryo scRNA-seq population from Tyser et al. [23]. Ridgeplots show the distribution of DESeq2 Wald statistics for genes within each population marker set along the ranked gene list; populations are ordered by normalised enrichment score (NES), and colour denotes FDR-adjusted p-values.

ITR, inverted terminal repeat; TRE3G, third-generation tetracycline-responsive promoter; cHS4, chicken hypersensitive site 4 insulator; pA, polyadenylation signal; PuroR, puromycin resistance gene; GSEA, gene set enrichment analysis; PCW2, post-conception week 2.

See also Figure S3.

Figure 4 - Duration of TBX6 activity determines commitment to a somitic mesoderm fate

(A) Experimental design to overexpress *TBX6* during the hTLS protocol in PB-Dox-TBX6 iPSCs.

(B) Schematics of piggyBac-insertable constructs for Dox-inducible overexpression of
TBX6-FKBP-V-T2A-mCherry (top), or mCherry alone (bottom).

(C, D) Immunofluorescence of hTLS made with (C) PB-Dox-TBX6 iPSCs, or (D) PB-
Dox-mCherry iPSCs, following sustained transgene induction (1 µM Dox from day -1),
showing expression of SOX2 (grey), PAX8 (cyan), FOXC2 (yellow), and the mCherry
transgene (magenta). Dotted white boxes indicate regions magnified in subsequent
panels. Scale bars: main, 100 µm; insets, 20 µm.

(E) Schematic of the DegTrace system. Dox induces TBX6-FKBP-T2A-mCherry
expression, while subsequent dTAG treatment selectively depletes FKBP-fused TBX6
protein. mCherry fluorescence is retained, enabling lineage tracing of cells that
transiently overexpressed TBX6.

(E') Experimental design to test various lengths of TBX6 protein activity in 2D- or 3D-
cultured hTLS followed by RT-qPCR or IF analysis.

(F) RT-qPCR for mCherry to assess transgene induction, or somitic mesoderm markers
FOXC1 and *MEOX1* in 2D-cultured hTLS generated from PB-Dox-TBX6 iPSCs either
untreated (Ctl [control], white bars), treated with Dox only from day -1 (red bars), or
treated with Dox from day -1 plus dTAG added either at day 0, day 1 or day 2 (blue
bars). . Statistical significance was assessed using two-way ANOVA on log-transformed
values, with treatment as the column factor and differentiation batch as the row factor; p
value corresponds to the main effect of treatment (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$,
ns = not significant). Data are plotted as $2^{-\Delta\Delta Ct}$ for visualisation; bars represent mean ±
SEM; dots represent mean of technical replicates from n = 5 independent differentiation
batches per group.

(G) Immunofluorescence of mCherry (magenta), FOXC2 (yellow), PAX3 (cyan) and
SOX2 (blue) in representative day 5 PB-Dox-TBX6 hTLS. Top row: hTLS treated with
Dox only from day -1 (6 days of TBX6 activity). Bottom row: hTLS treated with Dox from
day -1 followed by dTAG from day 0 (24 h TBX6 activity). Dotted white boxes indicate
regions of mCherry-expressing cells shown at higher magnification; an mCherry[+]

FOXC2⁻, PAX3⁺, SOX2⁺ epithelial structure is outlined in the bottom row. Scale bars:
main, 100 μm; insets, 20 μm.

(H) Immunofluorescence of mCherry (magenta), FOXC2 (yellow) and PAX8 (blue)
expression in representative day 5 PB-Dox-TBX6 hTLS. Top row: hTLS treated with Dox
from day -1 followed by dTAG from day 0 (24 h of TBX6 activity). Bottom row: hTLS
treated with Dox from day -1 followed by dTAG from day 2 (72 h of TBX6 activity).
Dotted white boxes indicate mCherry-expressing structures shown at higher
magnification; mCherry⁺ epithelial structures are outlined with dotted lines in the insets.
Scale bars: main, 100 μm; insets, 20 μm.

See also Figure S4.

Figure 5 – FOXC1 and FOXC2 establish and safeguard somitic mesoderm identity

(A) UMAP of 8,088 mesodermal subset cells, with the somite progenitor cluster (as in
Figure 1G) highlighted.

(B) Volcano plot of genes differentially expressed in the somite progenitor cluster
compared with the remaining mesodermal subset. Genes with adjusted p-value < 0.05
and |log2 fold change| > 0.58 (±1.5-fold) are highlighted (red, upregulated; blue,
downregulated). The top 20 upregulated transcription factors according to log2 fold
change are shown in darker colour. Box highlights FOXC1 and FOXC2.

(C) Inducible dual FOXC1 and FOXC2 CRISPRi strategy. Two piggyBac donor
constructs each encoding a Dox-inducible dCas9-KRAB-IRES-GFP cassette and a
single gRNA targeting either FOXC1 (with puromycin resistance; PuroR), or FOXC2
(with neomycin resistance; NeoR) were co-integrated into iPSCs using piggyBac
transposase. Dual antibiotic selection yields a single cell line harbouring both cassettes.
Dox treatment induces dCas9-KRAB expression, enabling simultaneous knockdown of
FOXC1 and *FOXC2* during somitic mesoderm differentiation.

(D) Experimental design to test dual knockdown of *FOXC1* and *FOXC2* in a 2D somitic
mesoderm differentiation protocol adapted from [45], followed by bulk RNA-seq.

(E) Volcano plot of differential gene expression following dual knockdown of *FOXC1* and
FOXC2 in 2D somitic mesoderm differentiation (n = 3 independent differentiation
batches per condition). Genes with adjusted p-value < 0.05 and |log2 fold change| >
0.58 (±1.5-fold) are highlighted (red, upregulated; blue, downregulated). Selected genes
associated with somitic, kidney, and lateral plate/cardiomyocyte/smooth muscle
identities are highlighted in darker colours. (Somitic: *FOXC1, FOXC2, MEOX1, HAND1,*
TCF15, TWIST1, RIPPLY1, SIX1, PAX3; kidney: *OSR1, NPHS1, EYA2, EPCAM, CDH1,*
JAG1, HEYL, HES5; lateral plate/cardiomyocyte/smooth muscle: *FOXF1*, *BMP2*, *BMP7*,
TNNT2, *CNN1*, *MYH7B*, *MYH11*).

(F) Dot plot showing enrichment of transcription factor motif families identified in
regulatory regions of FOXC1/2-dependent genes versus expressed non-dependent
genes (defined in (E)), using early somite ATAC-seq data [45]. Dot position indicates the
number of significantly enriched motifs per family (BH-adjusted q < 0.01); dot size
reflects total motifs representation in HOMER; colour denotes the significance ($-\log_{10}$ q)
of the most enriched motif in each family.

(G) Experimental design to test dual knockdown of *FOXC1* and *FOXC2* in hTLS.

(H-J) Quantification of the number of nuclei per confocal image that are (G) SOX2$^+$, (H)
PAX8$^+$, or (I) PAX3$^+$SOX2$^-$, in FOXC1/2-dCas9-KRAB hTLS either untreated (control,
white bars) or treated with 1µM Dox from day -1 (Dox, coloured bars). Dots represent
n=16 individual hTLS across 2 independent batches per group; bars represent median $\pm$
IQR; Mann-Whitney test (*** p < 0.001).

(K) Immunofluorescence of hTLS generated from FOXC1/2-dCas9-KRAB iPSCs and
harvested at day 5, either untreated (control, top) or treated with Dox from day -1 (1 µM,
bottom), showing GFP (green), SOX2 (yellow), PAX3 (magenta) and PAX8 (grey).
Dotted white lines outline PAX3$^+$SOX2$^+$ neural structures, white arrowheads indicate
PAX3$^+$SOX2$^-$ somitic tissue depleted upon Dox treatment. Scale bars, 100 µm.

See also Figure S5.

Figure 6 - FOXC1 drives a partial sclerotome differentiation program

(A) Experimental design to test dual FOXC1 and FOXC2 knockdown in a 2D sclerotome
differentiation protocol adapted from [45] followed by RT-qPCR.

(B-D) RT-qPCR for B) early somite markers, C) *FOXC1* and *FOXC2*, and D)
dermomyotome and sclerotome markers and canonical Hedgehog target genes, in
FOXC1/FOXC2-dCas9-KRAB iPSCs differentiated to early somite and then either
untreated (control) or treated from day 3 with doxycycline (Dox, B-D), or also with
smoothened agonist (SAG), or combined Dox and SAG treatment (C,D). Statistical
significance was assessed using two-way ANOVA on log-transformed values, with
treatment as the column factor and differentiation batch as the row factor; p-values
correspond to the main effect of treatment (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p <$
0.0001, ns = not significant). Data are plotted as $2^{-\Delta\Delta Ct}$; bars represent mean ± SEM;
dots represent mean of technical replicates from n = 5 independent differentiation
batches per group.

(E) Schematic of the FOXC1 overexpression strategy

(F) Schematic of experimental design to overexpress *FOXC1* in 2D-cultured hTLS for
subsequent RT-qPCR.

(G) Schematic of FOXC1 protein showing domain architecture, together with RT-qPCR
analysis of *FOXC1* transgene, *TWIST1*, *MEOX1*, *PAX9, TWIST2* and *SOX9* expression
in hTLS derived from PB-Dox-FOXC1 iPSCs at day 5, either untreated (control) or
treated with Dox (0.1 µM) from day 2. Statistical significance was assessed using two-
way ANOVA on log-transformed values, with treatment as the column factor and
differentiation batch as the row factor; p-values shown correspond to the main effect of
treatment (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$). Data are plotted as $2^{-\Delta\Delta Ct}$
for visualisation; bars represent mean ± SEM; dots represent mean of technical
replicates from n = 4 independent differentiation batches per group. N-TAD, N-terminal
activation domain; FKH, forkhead domain; C-TAD, C-terminal activation domain; grey
box, putative inhibitory domain [73].

(H) Schematic of FOXC1[ΔFKH] protein showing domain architecture with FKH DNA-binding domain replaced with a flexible linker; plots show RT-qPCR analysis for transgene and TWIST1 in day 5 2D-hTLS generated from iPSCs harbouring Dox-inducible FOXC1[ΔFKH] and treated with varying concentrations of Dox (0, 0.01, 0.1, 1 µM) from day 2. Statistical significance was assessed using two-way ANOVA on log-transformed values (ΔCt), with treatment as the column factor and differentiation batch as the row factor. The p values shown correspond to the main effect of treatment (** $p < 0.01$, **** $p < 0.0001$). Data are plotted as $2^{-\Delta\Delta Ct}$; bars represent mean ± SEM; dots represent mean of technical replicates from n=2-3 independent differentiation batches per group.N-TAD, N-terminal activation domain; FKH, forkhead domain; C-TAD, C-terminal activation domain; grey box, putative inhibitory domain [73].

(I) Experimental design to overexpress FOXC1 in hTLS

(J) Brightfield and corresponding red fluorescence images of hTLS generated from PB-Dox-FOXC1 iPSCs at days 3, 4, and 5. Scale bars, 300 µm.

(K) Immunofluorescence of hTLS generated from PB-Dox-FOXC1 iPSCs following FOXC1 overexpression from day 2, stained for mCherry (magenta) and F-actin (phalloidin, grey). White dotted box indicates region shown at higher magnification, in which individual epithelial cysts are outlined with dotted lines. Arrows indicate FOXC1-overexpressing cells occupying a subepithelial position. Scale bars: main, 100 µm; insets, 20 µm.

(L) Immunofluorescence of hTLS generated from PB-Dox-FOXC1 iPSCs following FOXC1 overexpression from day 2, stained for mCherry (magenta), PAX3 (cyan), and SOX2 (yellow). Arrows indicate FOXC1-overexpressing cells lacking both PAX3 and SOX2. Scale bars: main, 100 µm; insets, 20 µm.

(M) Immunofluorescence of hTLS generated from PB-Dox-FOXC1 iPSCs following FOXC1 overexpression from day 2, stained for mCherry (magenta), PAX8 (grey), and TWIST1 (green). Arrows indicate FOXC1-overexpressing cells co-expressing TWIST1. Scale bars: main, 100 µm; insets, 20 µm.

(N) Immunofluorescence of hTLS generated from PB-Dox-FOXC1[ΔFKH] iPSCs following overexpression from day 2, stained for mCherry (magenta), PAX3 (cyan), and F-actin (phalloidin, grey). White dotted box indicates region shown at higher magnification, in which individual epithelial cysts are outlined with dotted lines. Arrows indicate FOXC1[ΔFKH]-overexpressing cells within epithelial cysts. Scale bars: main, 100 µm; insets, 20 µm.

See also Figures S6 and S7.

Figure 7 – Proposed temporal gate and lock mechanism for human mesodermal diversification

In this model, TBX6 activity duration functions as a temporal gate that determines lineage competence during early human mesoderm specification. In the absence or at low levels of TBX6, cells adopt a neural fate, whereas intermediate TBX6 activity permits differentiation into intermediate (renal) mesoderm. Prolonged TBX6 activity biases cells toward a somitic mesodermal fate. Downstream, FOXC1/2 act as a stabilising "lock" that consolidates somitic identity, restricting alternative lineage potential and preserving competence for subsequent differentiation. Together, these findings define a hierarchical regulatory logic in which TBX6 activity duration regulates competence for mesodermal diversification, followed by downstream stabilisation of committed cell identities.

Supplementary Figure 1. snRNA-seq quality control and additional immunofluorescence analyses (related to Figure 1)

(A) Immunofluorescence of day 5 hTLS generated from SOX2::mNeon reporter iPSCs showing endogenous mNeon signal (green), and staining for SOX2 (grey), FOXC2 (magenta) and PAX8 (cyan) expression. A SOX2$^+$ neural tube-like structure is outlined with a dotted line. mNeon expression is specific to SOX2$^+$ cells within the neural tube-like structure but does not label all SOX2$^+$ cells. Scale bars, 100 µm.

(A') Representative hTLS from two independent differentiation batches generated using SOX2::mNeon iPSCs, shown in brightfield and corresponding green fluorescence to illustrate intra- and inter-batch variability.

(B) Immunofluorescence of FOXC2 (red), PAX6 (white) and SOX2 (green) expression in hTLS at day 5, with nuclei counterstained with DAPI (blue). Arrows indicate SOX2$^+$ PAX6$^+$ neural tube-like structures; arrowheads indicate FOXC2$^+$ epithelial somite-like structures. Scale bars, 100 µm.

(C) Immunofluorescence of the nephron markers PAX8 (green), ECAD/CDH1 (yellow), and HNF1B (magenta) in a day 8 hTLS. hTLS is outlined with dotted white line. Scale bar, 100 µm.

(D) Immunofluorescence of PAX8 (cyan), SOX2 (yellow), and FOXC2 (magenta) in a day 5 hTLS derived from an independent iPSC line, IMR-90-4. Nuclei are counterstained with DAPI, grey. hTLS is outlined with dotted line. Scale bars, 100 µm.

(E) Violin plots of hTLS snRNA-seq data showing the number of detected genes per cell at each time point.

(F) Violin plots of hTLS snRNA-seq data showing the percentage of mitochondrial reads per cell at each time point.

(G) UMAP representation of 13,618 hTLS cells coloured by unsupervised clustering with annotations determined by marker gene expression.

(H) Dot plot showing expression of selected marker genes for each cluster in (G).

(I) UMAP plots of 13,618 hTLS cells as in (G), with expression of selected marker genes
overlaid.

(J) Immunofluorescence of PAX3 (cyan), SOX2 (yellow), and the neural crest marker
SOX9 (magenta) in hTLS at day 8. White box indicates magnified region. Dotted white
line in the insets outline a PAX3$^+$SOX2$^+$ neural tube-like structure; arrowheads indicate
delaminating SOX9$^+$ neural crest cells. Scale bars: main image, 100 µm; insets, 20 µm.

(K) Immunofluorescence of the neural/somite marker PAX3 (cyan), the neural marker
SOX2 (yellow) and the endothelial marker PECAM-1/CD31 (magenta), in hTLS at day
5. White box indicates magnified region. Arrowheads indicate a PECAM-1$^+$ vessel-like
structure nestled between SOX2$^+$ and PAX3$^+$ neural/somitic tissue. Scale bars: main
image, 100 µm; insets, 20 µm.

(L, L', L") Transcriptional similarity between hTLS snRNA-seq dataset and human
reference embryogenesis dataset from Zhao et al., [25]. UMAP representations of human
reference embryogenesis dataset coloured by (L) time point and (L') cell type using
annotations from the reference. (L") projection of hTLS snRNA-seq cells (iPSC and
days 0-4) onto reference UMAP, coloured by time point, with reference cells in grey.

(M) UMAP of hTLS cells as in (G), coloured by unsupervised clustering at resolution =
0.03 to identify mesodermal cells (cluster 0), which were subsequently sub-clustered to
isolate the mesodermal subset.

(N) Heatmap of top 10 marker genes defining clusters in (M). Enriched genes were first
determined with Wilcoxon rank sum test with thresholds log2 fold change > 1 and
adjusted p-value < 1e-8.

(O) UMAP of mesodermal subset (as in Figure 1G) coloured by time point.

(P) Heatmap showing top 50 genes defining each cluster of the mesodermal subset
annotated as in Figure 1G, with selected marker genes presented. Enriched genes were
first determined with Wilcoxon rank sum test with thresholds log2 fold change > 1 and
adjusted p-value < 0.005.

Supplementary Figure 2. TBX6 is expressed transiently throughout the mesoderm before mesoderm diversification (related to Figure 2)

(A) Experimental design to differentiate iPSCs in the hTLS protocol on glass-bottomed dishes without aggregation for subsequent immunofluorescence analysis.

(B) Immunofluorescence of differentiated iPSCs (as outlined in A) showing expression of TBX6 (green), TBXT/Brachyury (yellow) and SOX2 (magenta) expression, with nuclei counterstained with DAPI (blue). Scale bars, 100 μm.

(C) Strategy to differentiate iPSCs using the hTLS protocol followed by dissociation then reseeding at day 0 to generate hTLS pools maintained in 2D culture for subsequent RT-qPCR analysis.

(D) Time-course RT-qPCR analysis of selected of early mesoderm, somite, kidney and neural markers during the hTLS protocol. Dots and bars represent mean ± SEM; n = 4 wells of hTLS per time point across 2 independent differentiation batches.

(E) RT-qPCR analysis of *TBXT* and *MSGN1* expression in TBX6-dCas9-KRAB iPSCs differentiated using the hTLS protocol and harvested after 24 hours (as in Figure 2E), either untreated (control) or treated with Dox (1 μM) from day -1. Dots represent n = 4 independent differentiation batches. Two-way ANOVA on log-transformed values (ΔCt), with treatment as the column factor and differentiation batch as the row factor. P values correspond to treatment effect (** $p < 0.01$). Data are plotted as $2^{-\Delta\Delta Ct}$; bars represent mean ± SEM.

(F) Immunofluorescence of TBX6-dCas9-KRAB iPSC cells differentiated as in (A) either without or with Dox (1 μM) treatment from day -1 to induce TBX6 knockdown, showing expression of TBX6 (magenta), and F-actin (grey) to outline cells, with GFP (green) as a readout for expression of the knockdown cassette, which is expressed heterogeneously after Dox treatment. Dotted box indicates magnified region. Arrowheads indicate expression of TBX6 in cells that lack GFP expression and thus did not express the knockdown cassette. Scale bars main images 100 μm; insets, 20 μm.

(G) Immunofluorescence of day 5 hTLS generated from a second independent TBX6-
dCas9-KRAB iPSC line with a distinct gRNA to target TBX6, either with no treatment
(control, top) or following Dox treatment from day -1 (1µM Dox, bottom), showing GFP
(green), SOX2 (yellow), FOXC2 (magenta), and PAX8 (grey). White brackets indicate
FOXC2$^+$PAX8$^+$ cells confined to a GFP$^-$ region. Scale bars, 100µm.

Supplementary Figure 3. TBX6 RNA-seq quality control and additional analysis
(related to Figure 3)

(A) RT-qPCR analysis of *TBX6, SOX2,* somitic mesoderm markers *FOXC1* and *FOXC2,*
and the intermediate mesoderm marker *OSR1,* in PB-Dox-TBX6 iPSCs differentiated
using the hTLS protocol and harvested after 24 h, either untreated (control) or treated
with Dox (0.1 µM) from day -1. Dots represent n = 5 independent differentiation batches.
Two-way ANOVA on log-transformed values (ΔCt), with treatment as the column factor
and differentiation batch as the row factor. P values correspond to treatment effect (** p
< 0.01, **** p < 0.0001). Data are plotted as $2^{-\Delta\Delta Ct}$; bars represent mean ± SEM.

(B) Principal component plots of the TBX6 overexpression bulk RNA-seq dataset.

(C) Sample-to-sample correlation heatmap of the TBX6 overexpression bulk RNA-seq
dataset.

(D) Dot plot showing gene ontology (GO) biological processes (BP) analysis,
highlighting selected processes upregulated following TBX6 overexpression. Dot size
indicates gene count; colour denotes FDR-adjusted p-value.

(E) Ridge plots of gene set enrichment analysis (GSEA) of genes following TBX6
overexpression (Dox versus control), ranked by the DESeq2 Wald statistic [72], and tested
against the top 50 marker genes for each annotated PCW3 human embryo scRNA-seq
population from Zeng et al. [24]. Ridgeplots show the distribution of DESeq2 Wald
statistics for genes within each population marker set along the ranked gene list;
populations are ordered by normalised enrichment score (NES), and colour denotes
FDR-adjusted p-values.

(F) UMAP of scRNA-seq data from PCW2 (CS7) human embryo from Tyser et al. [23],
coloured by cell type.

(G) UMAP of TBX6 expression in the Tyser et al. dataset.

(H) Projection of TBX6-upregulated gene module onto the Tyser et al. UMAP. The
module was defined as the top 500 genes upregulated upon TBX6 overexpression in
hTLS (adjusted p-value < 0.05, log2 fold change > 2) and filtered for overlap between
the Tyser and Zeng datasets (221 genes total remaining).

(I) Violin plot showing enrichment of the TBX6-upregulated gene module across cell
types in the Tyser et al. dataset, using cluster annotations as in (E).

(J) UMAP of scRNA-seq data from PCW3 (CS10) human embryo dataset from Zeng et
al. [24], coloured by cell type.

(K) UMAP of TBX6 expression in the Zeng et al. dataset.

(L) Projection of the TBX6-upregulated gene module (as in H) onto the Zeng et al.
dataset.

(M) Violin plot showing enrichment of the TBX6-upregulated gene module across cell
types in the Zeng et al. dataset, using cluster annotations as in (J).

1009

Supplementary Figure 4. DegTrace (TBX6-FKBP-V-T2A-mCherry lineage tracing)
system (related to Figure 4).

(A) Experimental design to test depletion of the TBX6-FKBP fusion protein following
dTAG treatment in PB-Dox-TBX6 iPSCs cultured in iPSC maintenance medium
(Stemflex).

(B) Immunofluorescence of F-Actin (phalloidin, grey), mCherry (magenta) and TBX6
(green), with nuclei counterstained with DAPI (blue), in representative PB-Dox-TBX6
iPSCs cultured as in (A) and either untreated (no Dox, top row), treated with Dox for 48
h (0.1 µM, middle row), or treated with Dox for 48 h followed by dTAG for 24 h (2 µM,
bottom row). Scale bars, 100 µm.

(C) Experimental design to test depletion of the TBX6-FKBP fusion protein in hTLS
generated from PB-Dox-TBX6 iPSCs, followed by immunofluorescence (IF) analysis.

(D) Immunofluorescence of mCherry (magenta), TBX6 (green), PAX8 (cyan), and PAX3
(grey) in representative day 5 PB-Dox-TBX6 hTLS treated as in (C), with Dox (0.1 μM)
from day -1 followed by dTAG (2 μM) from day 0. White boxes indicate mCherry[+]
epithelial structures shown at higher magnification; dotted lines in the insets outline the
mCherry[+] structures. The top row shows an hTLS retaining low levels of TBX6 within the
mCherry[+] structure, whereas the bottom row shows an mCherry[+] structure with TBX6
fully depleted. Scale bars: main, 100 μm; insets, 20 μm.

1029

Supplementary Figure 5. Dual FOXC1 and FOXC2 knockdown quality control and
additional analyses (related to Figure 5)

(A) Schematic of the 2D somitic mesoderm differentiation protocol, adapted from Loh et
al. [45].

(B) Time-course RT-qPCR analysis of selected marker genes during differentiation. Dots
represent n = 2 independent differentiation batches per group, bars represent mean ±
SEM.

(C) Schematic of piggyBac-based constructs for Dox-inducible dual CRISPRi-mediated
knockdown of *FOXC1* and *FOXC2*.

(D) Principal component analysis of the dual FOXC1 and FOXC2 knockdown bulk RNA-
seq dataset.

(E) Sample-to-sample correlation heatmap of the dual FOXC1 and FOXC2 knockdown
bulk RNA-seq dataset.

(F, G) Gene-level boxplots showing DESeq2 [72] normalised counts (log10 scale) for (F)
representative genes differentially expressed during somite differentiation including
downregulated canonical somite genes (blue), upregulated kidney-associated genes
(orange), and upregulated genes associated with lateral plate (LP), cardiomyocyte, or
smooth muscle (SM) lineages (red), and (G) early and paraxial mesodermal markers,

following dual FOXC1 and FOXC2 knockdown. FOXC1 and FOXC2 were among the

most strongly downregulated genes (log2FC ≈ –3; FDR < $10^{-1\square\square}$), confirming efficient

repression. Dots represent biological replicates; boxplots indicate the IQR across

replicates; grey indicates control and coloured indicates Dox treatment. Asterisks

denote DESeq2 adjusted p-values (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

(H) Dot plot showing selected enriched gene ontology (GO) biological process (BP)

terms among genes upregulated (UP) or downregulated (DOWN) following dual

FOXC1/FOXC2 knockdown. Dot size indicates gene ratio; colour denotes FDR-adjusted

p-value.

(I-K) Ridge plots of gene set enrichment analysis (GSEA) of genes following dual

FOXC1 and FOXC2 knockdown (Dox versus control), ranked by the DESeq2 Wald

statistic [72], tested against marker gene sets for (I) PCW3-5 human embryo scRNA-seq

populations from Zeng et al. [24], (J) fetal human nephron cell types from Stewart et al. [47],

and (K) PCW6.5-7 human heart cell types from Asp et al. [46]. Ridgeplots show the

distribution of DESeq2 Wald statistics for genes within each population marker set along

the ranked gene list; populations are ordered by normalised enrichment score (NES),

and colour denotes FDR-adjusted p-values.

(L) Immunofluorescence of hTLS generated from FOXC1/FOXC2-dCas9-KRAB iPSCs

and harvested at day 5, either untreated (control, top) or treated with Dox from day -1 (1

µM, bottom), showing GFP (green), SOX2 (yellow), FOXC2 (magenta) and PAX8

(cyan), illustrating depletion of FOXC2 upon dual FOXC1 and FOXC2 knockdown.

Scale bars, 100 µm.

(M) Schematic of experimental design to test epistatic relationship between TBX6 and

FOXC1/2 in 2D-cultured hTLS.

(N) RT-qPCR analysis of somite markers in 2D-cultured hTLS derived from PB-Dox-

TBX6 or Dox-TBX6-dCas9-FOXC1/2 iPSCs at day 5, either untreated (control) or

treated with Dox (0.1 µM) from day -1. Dots represent mean of technical replicates from

n = 4 independent differentiation batches per condition. Two-way ANOVA on log-

transformed values (ΔCt), with treatment as the column factor and differentiation batch

as the row factor, followed by post hoc comparisons of Dox-treated versus control
conditions within each cell line. P values correspond to main treatment effect (* p <
0.05, ** p < 0.01). Data are plotted as $2^{-\Delta\Delta Ct}$ for visualisation, normalised to the
untreated control condition within each iPSC line; bars represent mean ± SEM.

ITR, inverted terminal repeat; TRE3G, third-generation tetracycline-responsive
promoter; cHS4, chicken hypersensitive site 4 insulator; pA, polyadenylation signal;
PuroR, puromycin resistance gene.

1084 Supplementary Figure 6. hTLS retain sclerotome differentiation potential (related
1085 to Figure 6)

(A) Schematic of sclerotome differentiation *in vivo*. Hedgehog ligands secreted by the
notochord signal to the ventral domain of epithelial somites to activate sclerotome-
associated genes including FOXC1, FOXC2, TWIST1, PAX1, PAX9, and SOX9, to
downregulate PAX3, and to initiate epithelial-to-mesenchymal transition (EMT), while
dorsal somites remain PAX3^{+}.

(B) UMAP plots of 13,618 hTLS snRNA-seq cells (as in Figure 1E) showing minimal
expression of Hedgehog ligands (*SHH*, *IHH*, *DHH*), and expression of downstream
Hedgehog pathway mediators (*GLI1*, *GLI2, GLI3*).

(C) Experimental design to test the effect of Hedgehog pathway activation using
smoothened agonist (SAG; 500 nM) in hTLS from day 5 to day 8 of the protocol.

(D) Immunofluorescence of SOX2 (yellow), FOXC1 (grey), and PAX9 (cyan) expression
in day 8 hTLS cultured as in (C), either untreated (control, top row) or treated with SAG
from day 5 (500 nM, bottom row). Scale bars, 100 µm.

(E-G) Quantification of the number of nuclei per confocal image positive for (E) SOX2,
(F) FOXC2, or (G) PAX3 (whilst also SOX2^{-}) in day 8 hTLS either untreated (control,
white bars) or treated with SAG from day 5 (500 nM, coloured bars). Dots represent
individual hTLS (n = 12 hTLS across 2 independent differentiation batches); bars
represent median ± IQR; Mann-Whitney test (** p < 0.01, *** p < 0.001).

(H) Immunofluorescence of mCherry (magenta), FOXC2 (grey), and SOX9 (green)
expression in day 5 hTLS generated from PB-Dox-FOXC1 iPSCs treated with Dox from
day 2 (0.1 μM). White box indicates magnified region; white arrows indicate mCherry[+]
FOXC2[+] cells expressing SOX9; arrowheads indicate a rare mCherry[-] FOXC2[+] SOX9[+]
cell. Scale bars: main, 100 μm; insets, 20 μm.

(I) Schematic of the Dox-inducible FOXC2-overexpression strategy and corresponding
immunofluorescence of SOX2 (blue), mCherry (magenta), PAX8 (grey), and PAX3
(green) expression in a day 5 hTLS generated from PB-Dox-FOXC2 iPSCs and treated
with Dox from day 2 (0.1 μM). White box indicates magnified region; the dotted lines
outline an mCherry[+], PAX8[-], PAX3[-] cell cluster positioned between PAX8[+] and PAX3[+]
tissue. Scale bars: main, 100 μm; insets, 20 μm.

1115

Supplementary Figure 7. Dual TWIST1 and TWIST2 knockdown in sclerotome
differentiation (related to Figure 7)

(A) Schematic of a piggyBac-insertable construct for Dox-inducible overexpression of
TWIST1.

(B) Experimental design to test TWIST1 overexpression in hTLS.

(C) Immunofluorescence of hTLS generated from PB-Dox-TWIST1 iPSCs showing
mCherry (magenta), FOXC2 (cyan), F-actin (phalloidin, yellow) and SOX2 (blue)
expression at day 5 following Dox treatment from day 2. The white box indicates
magnified region. Dotted white lines in magnified panels indicate FOXC2[+] epithelial
structures, and white arrows indicate mCherry[+] TWIST1-overexpressing cells occupying
a subepithelial position. Scale bars: main, 100 μm; insets, 20 μm.

(D) Experimental design to test dual TWIST1 and TWIST2 knockdown in a 2D
sclerotome differentiation protocol (adapted from Loh et al., [45]), followed by RT-qPCR.

(E) RT-qPCR analysis of *FOXC1, FOXC2, TWIST1, TWIST2, PAX9, PAX1, SOX9,*
PTCH1 and *GLI1* expression in TWIST1/2-dCas9-KRAB iPSCs differentiated to

sclerotome and either untreated (control), or treated from day 3 with Dox (1 μM), SAG,

or combined Dox and SAG treatment. Dots represent mean of technical replicates from

n = 3 independent differentiation batches per condition. Two-way ANOVA on log-

transformed values (ΔCt), with treatment as the column factor and differentiation batch

as the row factor. P values correspond to main treatment effect (* $p < 0.05$, ** $p < 0.01$,

*** $p < 0.001$, **** $p < 0.0001$). Data are plotted as $2^{-\Delta\Delta Ct}$ for visualisation, normalised to

the untreated control condition within each iPSC line; bars represent mean ± SEM.

1138

 STAR METHODS

 EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Cell Lines

Human iPSCs were cultured as previously described with modifications [74]. Human
iPSCs used in this study were kindly gifted by Austin Smith [75]. iPS(IMR90)-4 cells were
obtained from WiCell. All iPSCs were maintained in StemFlex (ThermoFisher,
A3349401) containing Pen/Strep on Geltrex (Gibco, A14133-01)-coated plates. Cells
were passaged every 3-4 days using PBS with 0.5µM EDTA and re-seeded in StemFlex
containing Pen/Strep and 10 µM ROCK inhibitor Y27623 (ROCKi; Tocris, 1254).
Transgenic iPSC lines (TBX6-dCas9-KRAB, FOXC1/2-dCas9-KRAB, TWIST1/2-dCas9-
KRAB, PB-Dox-FOXC1, PB-Dox-FOXC1[ΔFKH], PB-Dox-TBX6, PB-Dox-FOXC2, PB-
Dox-TWIST1 and PB-Dox-mCherry), generated as described below, were maintained in
StemFlex containing Pen/Strep, 0.5 µg/mL puromycin and/or 200 µg/mL G418 as
appropriate. All cell cultures were incubated at 37^{o}C with 7% CO_2 and 5% O_2.

Ethical Statement about use of human embryoid models

The human trunk-like structure system used in this study is a human pluripotent stem
cell–derived model that although reproduces features of early post-gastrulation human
developmental stages, does not generate extraembryonic tissues, and is therefore
classified as a *non-integrated embryo model* under the terminology established in the
2021 International Society for Stem Cell Research (ISSCR) guidelines [76]. In addition,
the model lacks multiple embryonic lineages including, but not limited to, all anterior
neural tissues, lateral mesoderm, notochord, endoderm, and non-neural ectoderm. No
human embryos were used in this research. This research was conducted in
accordance with the 2021 ISSCR Guidelines.

1163

METHOD DETAILS

Plasmid constructs

Target gene knockdown in iPSCs was performed as previously described with

modifications [74]. pPB-Ins-U6p-sgRNAentry-EF1Ap-TetOn3G-IRES-Neo (Addgene

plasmid # 183411; RRID:Addgene 183411) and pPB-Ins-TRE3Gp-KRAB-dCas9-

ecDHFR-IRES-GFP-EF1Ap-Puro (Addgene plasmid # 183410; RRID:Addgene 183410)

were kind gifts from Azim Surani. The SOX2:mNeon iPSC line was generated with the

following plasmids: pCAS9-mCherry-Frame +0 [77] (gift from Veit Hornung, Addgene

plasmid # 66939 ; http://n2t.net/addgene:66939 ; RRID:Addgene_66939), pCRISPaint-

mNeon-PuroR [M1G] [78] (gift from Michael Hoelzel, Addgene plasmid # 171800 ;

http://n2t.net/addgene:171800 ; RRID:Addgene_171800), and pSPgRNA [79] (gift from

Charles Gersbach, Addgene plasmid # 47108 ; http://n2t.net/addgene:47108 ;

RRID:Addgene_47108) containing gRNA targeting the human SOX2 locus

(TGCCCCTCTCACACATGTGA). All gRNA oligos used were ordered using the custom

DNA oligo synthesis service from Merck. Specific gRNA sequences for dCas9-KRAB

mediated gene repression were designed by identifying putative transcription start sites

(TSSs) for target genes according to FANTOM CAGE data publicly available on ZENBU

[80], then the Wellcome Sanger Institute Genome Editing (WGE) CRISPR Finder [81] was

used to identify CRISPR sequences 50-100 bp downstream of a putative TSS for

optimal gene knockdown [82]. Two gRNA sequences were tested per TSS. For generation

of gRNA expression vectors against TBX6, complementary gRNA oligos were annealed

and ligated into pPB-Ins-U6p-sgRNAentry-EF1Ap-TetOn3G- IRES-Neo (Addgene

plasmid, 183411) by Esp3I digestion (NEB, R0734) followed by ligation with T4 DNA

ligase (New England Biolabs, M0202S). Piggybac-based Dox-inducible transgene

overexpression plasmids were created by first PCR amplifying (with Cloneamp Takara,

639298) the Dox-response element from pPB-Ins-TRE3Gp-KRAB-dCas9-ecDHFR-

IRES-GFP-EF1Ap-Puro (Addgene plasmid # 183410) and the gRNA-rtTA expression

cassette from pPB-Ins-U6p-sgRNAentry-EF1Ap-TetOn3G-IRES-Neo (Addgene plasmid,

183411), and cloning into VB240515-1632gka backbone (Vectorbuilder) by digesting

with EcoRV (NEB, R3195) and PacI (NEB, R0547) and inserting using In Fusion Snap

Assembly master mix (Takara, 638948). This yielded Dox-inducible dCas9-KRAB

vectors that were used for knockdown of FOXC1/2 and TWIST1/2 factors; versions with

Puro and Neo resistance were generated to combine 2 distinct gRNA-expressing

plasmids into the same cells. For Dox-inducible expression of transgenes of interest, the
dCas9-KRAB-U6-gRNA cassette was excised with NotI (NEB, R3189) and AsiSI (NEB,
R0630) and transgenes were cloned in by In Fusion. FOXC1 and TWIST1 transgenes
were synthesised by Twist Biosciences. mCherry was synthesised by Vectorbuilder
(VB240515-1632gka). TBX6 was PCR amplified from TFORF1979 (Addgene 142303,
kind gift from Feng Zhang). FKBP-V was PCR amplified from
pBS_GGSG_Fkbp_2xHA_BB2 (Addgene 235503, kind gift from Brian Hendrich). PCR
amplified fragments were gel extracted using Monarch gel extraction kit (NEB, T1120).
Stellar competent E. coli (Takara, 636763) were used for transformation. Successful
transformants were purified using QIAprep Spin Miniprep Kit (Qiagen, 27104) or
QIAprep Plasmid Plus Midiprep Kit (Qiagen, 12943), then validated with Nanopore
sequencing (Oxford Nanopore Technologies) by Plasmidsaurus.

Quantitative reverse transcriptase PCR (RT-qPCR)

To isolate RNA for RT-qPCR analysis, cells were washed twice with PBS and then lysed
in 350 µl RLT lysis buffer prior to RNA extraction using the RNeasy mini kit (Qiagen,
74106) with on-column DNase digestion (Qiagen, 79254) and reverse transcribed to
cDNA using the QuantiTect Reverse Transcription kit (Qiagen, 205314). Quantitative
PCRs were performed using PowerUp SYBR Green Master Mix (Applied Biosystems,
A25742) with forward and reverse primers diluted to final concentration of 1 µM per
reaction and run either on the Quantstudio 5 384-well qPCR machine (Applied
Biosystems) or the StepOne Plus with 96-well block (Applied Biosystems). Statistical
analyses were performed on ΔCt values (Ct_target – Ct_reference). Data were
converted to $2^{-\Delta\Delta Ct}$ values for visualisation [83]. Details of statistical tests used are
described in the figure legends. RT-qPCR data were analysed using Prism (Graphpad,
v11.0.0).

1222

Transgenic cell lines

TBX6-dCas9-KRAB, FOXC1/2-dCas9-KRAB, TWIST1/2-dCas9-KRAB, PB-Dox-
FOXC1, PB-Dox-FOXC1[ΔFKH], PB-Dox-TBX6, PB-Dox-FOXC2, PB-Dox-TWIST1 and

PB-Dox-mCherry transgenic lines were established as previously described with
modifications [74]. 6.5x10^5 iPSCs were seeded into Geltrex-coated 6-well plates in
StemFlex with ROCKi, and the following day were washed with PBS and incubated in
1ml Optimem media (Gibco, 31985070) with Revitacell supplement (Fisher, 15317447).
For each well of a 6-well plate, cells were transfected with Super piggyBac Transposase
expression vector (Stratech, PB210PA-1-SBI) (500 ng), together with the donor
plasmid(s) to a total of 2 µg plasmid DNA mixed in 3.5 µl Lipofectamine Stem
transfection reagent (Life Technologies, STEM00003) in 400 µl Opti-MEM with
Revitacell. DNA-lipofectamine mixtures were incubated at room temperature for 20
minutes and then added dropwise to cells, which were then incubated at 37^{o}C for 4
hours. Media were then replaced with 3 ml StemFlex with Revitacell. The following day,
media were refreshed with StemFlex with Revitacell. 48-72 hours later, media were
refreshed with fresh StemFlex with Revitacell containing antibiotics 0.5 µg/mL
puromycin and/or 200 µg/mL G418 for selection. Media were refreshed with selection
media every 2-3 days thereafter. A week later, surviving colonies were pooled, and
knockdown efficiency was assessed by addition of 1µM Dox (a concentration we found
in previous studies to be optimal [74]) followed by RT-qPCR with primers specific to the
target gene.

To make the SOX2:mNeon CRISPR-edited iPSC line, iPSCs were transfected as
described above, using the following plasmids: pCAS9-mCherry-Frame +0 [77],
pCRISPaint-mNeon-PuroR [M1G] [78], and pSPgRNA [79] containing gRNA targeting the
human SOX2 locus (TGCCCCTCTCACACATGTGA). 72 hours after transfection, cells
were selected with 0.5 µg/mL puromycin, and successful colonies were isolated, with
mNeon expression confirmed by fluorescence microscopy.

1250

Human trunk-like structure generation

Human trunk-like structures (hTLS) were generated using a previously described
protocol with modifications [27]. Human iPSCs were grown to 70% confluency, washed
with PBS containing 0.5µM EDTA and dissociated with Accutase (Day -2), counted
using Countess II (Life Technologies, AMQAX1000), then seeded evenly onto Geltrex-

coated 6-well plates at 400,000 cells per well in 2ml Nutristem hPSC XF medium
(Sartorius, 05-100-1A) with ROCKi. The following day (Day -1), cells were incubated in
Nutristem with 6µM CHIR (ABCR, AB253776). The next day (Day 0), cells were washed
with PBS with EDTA, then dissociated with Accutase, centrifuged at 300g and
resuspended in Essential 6 media (E6; Thermo, A1516401) containing 3µM CHIR and
500nM retinoic acid (RA), and counted. For 3D hTLS, 400 cells were seeded in 50µl of
E6 with 3µM CHIR and 500nM RA per well of a round-bottom low-adhesion 96 well
plate (Corning, CLS7007); the plate was then centrifuged at 200g for 2 minutes and
incubated at 37^{o}C to promote aggregation. By the following day (day 1) cell aggregates
were visible by light microscopy at 4x magnification, at which point 150 µl E6 was added
to each well gently to avoid disrupting the aggregates. On day 2, 150 µl media was
carefully removed from each well, with care taken to avoid aspirating the aggregates,
and 150 µl E6 containing 10% Matrigel (Corning, 356231) and 10 nM RA was gently
added to each well, with Dox added as indicated. Plates were incubated at 37^{o}C until
Day 5, by which point the aggregates had elongated into trunk-like structures. To
harvest for immunofluorescence analysis, structures were gently aspirated using a glass
Pasteur pipette and fixed in 3% PFA for 30 minutes as described below. Alternatively, if
the experiment was continued, 50 µl E6 was added to each well together with any
indicated treatment, and hTLS were cultured further until harvest. For 2D hTLS, on day
0 after dissociation and counting, 4.8×10^4 cells were seeded into Geltrex-coated 24-well
plates in 500 µl of E6 with 3µM CHIR and 500nM RA per well. The following day (day 1)
375 µl medium was removed from each well and replaced with 375 µl E6 only. On day
2, 375 µl medium was removed and replaced with 375 µl E6 containing 10% Matrigel
and 10 nM RA. Plates were incubated at 37^{o}C, then on day 5 wells were washed twice
with PBS then lysed for RNA extraction. The 2D hTLS protocol mirrors the 3D protocol
in terms of treatment conditions with the exception that instead of growing as individual
3D structures, the 2D hTLS adhere to the 24 well plate and interconnect. Both protocols
show similar marker gene expression dynamics, with the emergence of neural (SOX2$^+$)
cells, somitic mesoderm (FOXC2$^+$) and renal mesoderm (PAX8$^+$) identities.

Immunofluorescence analysis

To harvest for immunofluorescence (IF) analysis, hTLS were transferred using a glass

Pasteur pipette into a 30 ml sterilin, were centrifuged at 200g for 1 minute, washed once

with PBS, then fixed in 3% PFA for 30 minutes at room temperature with gentle rotation,

and then washed in PBS (3x10 min) and stored in PBS with 0.01% sodium azide at 4^oC.

Prior to antibody staining, samples were incubated in blocking buffer overnight at 4^oC

(PBS containing 0.25% Triton X-100, 5% BSA and 0.01% sodium azide). Samples were

incubated in primary antibodies diluted in wash buffer (PBS containing 0.1% Triton X-

100, 0.1% BSA and 0.01% sodium azide) at the indicated concentrations overnight at

4^oC. Next, samples were washed in wash buffer (3x15 min at room temperature) and

incubated in secondary antibodies, with DAPI (abcam, ab228549) where indicated,

diluted in wash buffer at the indicated concentrations overnight at 4^oC. The following

day, samples were washed once in wash buffer and mounted on glass slides in

Vectorshield (Vector Labs, H1000) in 120 µm spacers (Invitrogen, S24735) with

coverslips overlaid.

Imaging and image analysis

Brightfield images were taken using an EVOS M5000 microscope (Life Technologies,

AMF5000SV). Confocal images were taken using a Zeiss LSM 710 or 880. Image files

were analysed using Zeiss Zen (3.12) or Fiji (Image J). All immunofluorescence images

presented are representative confocal slices from at least 2 independent hTLS

differentiation batches. To quantify number of cells expressing a given marker, single

confocal images were analysed in CellProfiler using a custom analysis pipeline [84].

Briefly, positively stained nuclei were identified using a thresholding and declumping

module (IdentifyPrimaryObject module with Otsu two-class adaptive thresholding).

Smoothing factors were determined empirically. Objects (nuclei) were filtered by size,

and the number of objects within that channel was measured.

1311

2D somitic mesoderm differentiation

Human iPSCs were differentiated to somitic mesoderm using a previously described

protocol with modifications [45]. Human iPSCs were grown to 70% confluency, washed

with PBS containing 0.5µM EDTA and dissociated with Accutase (day -1), counted using

Countess II (Life Technologies, AMQAX1000), then seeded evenly onto Geltrex-coated
24-well plates at 50,000 cells per well in 500 µl Nutristem hPSC XF medium (Sartorius,
05-100-1A) with ROCKi. The following day (day 0), cells were incubated in Nutristem
with 6µM CHIR (ABCR, AB253776) and 100 nM PIK90 (Tocris, 7902). Recombinant
FGF2 and Activin A were omitted as Nutristem contains FGF2 and TGFß activity. The
next day (day 1, primitive streak stage) media were refreshed with E6 containing 3 µM
CHIR, 20 ng/ml FGF2, 1 µM A-83-01 (Tocris 2939) and 250 nM LDN 193189 (Tocris
6053). On day 2 (paraxial mesoderm stage), media were refreshed with E6 containing 1
µM A-83-01, 250 nM LDN 193189, 1 µM XAV 939 (Tocris 3748) and 500 nM PD0325.
On day 3 cells were early somite stage, at which point cells were harvested for RNA
extraction of bulk RNA-seq. Alternatively, for sclerotome differentiation cells media were
refreshed with E6 containing 500 nM SAG (Tocris 4366) and cells were cultured for a
further 72 hours. For dCas9-KRAB knockdown experiments, Dox (1 µM) was added
either at day 1 or day 3 as indicated.

1330

Antibodies

Primary antibodies used for protein detection, with their corresponding dilutions for
immunofluorescence (IF) were as follows: rabbit anti-PAX8 [EPR13511] (Abcam,
ab189249, 1:200), sheep anti-FOXC2 (R&D #AF6989, 1:200), rat anti-SOX2 (Invitrogen
#14-9811-82, 1:200), goat anti-DACH2 (R&D #AF5230, 1:100), rabbit anti-ZO1 (Cell
Signaling #13663, 1:200), rabbit anti-PAX6 (abcam #195045, 1:200), goat anti-TCF-2/
HNF-1 beta (R&D Systems, AF3330, 1:200), mouse anti-E-Cadherin (BD Bioscience,
610181, 1:200), mouse anti-PAX3 (R&D #MAB2457, 1:100), mouse anti-PAX3 (DSHB
#PAX3-c, 1:100), rabbit anti-SOX9 (Sigma #AB5535, 1:200), goat anti-PECAM-1 (R&D
#AF3628, 1:200), goat anti-TBX6 (R&D #AF4744, 1:200), rabbit anti-TBXT (Cell
Signaling #81694, 1:200), Phalloidin-647 (Thermo #A22287, 1:400), RFP-568 booster
(Proteintech #rb2AF568-50, 1:100), and mouse anti-Twist1 (abcam #ab50887, 1:100).
Alexa Fluor 405-, 488-, 568- and 647-conjugated secondary antibodies (Invitrogen)
were used for detection of primary antibodies.

1345

Bulk RNA-sequencing (RNA-seq)

Bulk RNA-seq of TBX6 overexpression (Figures 3 and S3) was performed on n = 4 independent differentiation batches per condition. Bulk RNA-seq of dual FOXC1/2 knockdown (Figures 5 and S5) was performed on n = 3 independent differentiation batches per condition. For each sample, at least 100,000 cells per condition were collected at the indicated time points. Cells were washed with PBS containing EDTA, dissociated using Accutase, pelleted by centrifugation, and lysed in DNA/RNA Shield (Zymo, R1100-50).

Bulk RNA-seq libraries were prepared and sequenced by Plasmidsaurus using an Illumina-based 3′ end counting approach. Total RNA was extracted using a bead-based method, and mRNA was reverse transcribed using poly(dT) priming, followed by second-strand synthesis, tagmentation, and PCR amplification. Libraries incorporated unique molecular identifiers (UMIs) to enable deduplication of PCR duplicates and were sequenced as single-end reads (~90 bp) enriched at the 3′ end of transcripts.

Sequencing reads were processed using a standardized pipeline including quality filtering, alignment to the reference genome using STAR, UMI-based deduplication, and gene-level quantification using featureCounts to generate count matrices for downstream analysis.

Bulk RNA-seq analysis

Count matrices were imported into RStudio (version 2024.09.1+394) running R version 4.4.2, and differential gene expression analysis between control and Dox-treated conditions was analysed using DESeq2 on raw count data [72]. The significance threshold for differential expression was set as log2 fold change > 0.58 (corresponding to fold change > 1.5) and adjusted p-value < 0.05. Transformed, variance-stabilised expression values were used for principal component analysis and sample-to-sample correlation heatmaps. Expression plots for individual genes used DESeq2 normalised counts.

1373   Gene ontology (GO) analysis was performed with the enrichGO function from
1374   clusterProfiler [85].

1375

Single-nucleus RNA-seq library preparation

For single-nucleus RNA-sequencing experiments, cell pellets or hTLS containing 0.5-1
M cells harvested at the indicated time points were snap frozen. Pellets were thawed
and resuspended in 1 ml of ice-cold nuclei permeabilization buffer (10 mM Tris-HCl
pH7.4, 10 mM NaCl, 3 mM MgCl$_2$, 0.2% IGEPAL CA300, 0.1 mg/ml BSA, 1x protease
inhibitor cocktail (Roche cat.11873580001), 0.5 U/µl ea Superase RNAse Inhibitor and
RNAaseOUT). Cells were then permeabilized for 5 min and centrifuged at 600g for 5
min at 4°C. Cells were lightly fixed by resuspending in 475 µl fixation buffer (1xPBS,
0.2% formaldehyde, 0.5 U/µl ea Superase RNAse Inhibitor and RNAaseOUT) and
incubated on ice for 5 min. Fixation was stopped by adding 2 µl 1M Tris-HCl pH7.4
containing 0.1 mg/ml BSA. Nuclei were pelleted at 600g for 5 min at 4°C and
resuspended in 1xPBS containing 1x protease inhibitor cocktail and 0.5 U/µl Superase
RNAse Inhibitor and 0.5 U/µl RNAaseOUT.

For in situ reverse transcription (RT) reaction 50,000 cells for each sample were
aliquoted in 0.2 ml PCR tubes containing 2.5 µM sample specific barcoded primers as
previously described [86]. To each tube RT mix (1x Maxima H buffer, 0.5 mM dNTPs, 0.1
mg/ml BSA, 400U Maxima Reverse H Minus Reverse Transcriptase (Thermo Scientific)
and 0.5 U/ul ea Superase RNAse Inhibitor and RNAaseOUT) was added to a final
volume of 20 µl. The reverse transcription was performed in a thermocycler with the
following program: Step 1: 50°C × 10 min; Step 2 (3 cycles): 8°C × 12 s, 15°C × 45 s,
20°C × 45 s, 30°C × 30 s, 42°C × 2 min, 50°C × 5 min; Step 3: 50°C × 10 min. After the
reaction, the nuclei were transferred and pooled on ice into a 1.5-ml tube prewashed
with 5% BSA in 1x PBS. Then Triton X-100 was added to a final percent of 0.075 and
nuclei were pelleted at 800g for 5 min at 4°C.

For ligation-based combinatorial barcoding the pooled nuclei were resuspended in 1 ml
of 1x NEBuffer3.1 (NEB) and transferred to 2,910 µl of ligation buffer (1x T4 ligase

buffer (NEB), 0.35 mg/ml BSA, 25 mM NaCl, 40,000 U T4 ligase (NEB), and 0.25 U/μl

Superase RNAse Inhibitor and 0.25 U/μl RNAaseOUT). Using a multichannel pipette,

40 μl of the ligation mix was distributed to the 96 wells of the first R02 barcode plate [86].

The plate was sealed and placed on a Thermomixer (Eppendorf) and mixed for 30 sec

at 37°C at 1200 rpm. Then incubated for a further 30 min at 37°C at 300 rpm. After

incubation the plate was removed and 10 μl of R02 blocking solution (26 μM R02

blocking oligo 5' ATCCACGTGCTTGAGAGGCCAGAGCATTCG in 1x T4 ligase buffer)

was added to each well. The plate was re-sealed and mixed again for 30 sec at 1200

rpm before incubating for an additional 30 min at 37°C. The nuclei from each well were

then pooled into 1.5 ml tubes and pelleted at 800g for 5 min at 4°C. The second round

of ligation was then carried out using R03 biotin tagged barcode plate [86] similarly to the

first round, except that after the 30 min ligation reaction, blocking solution (26 μM

quencher oligo 5' GTGGCCGATGTTTCGGTGCGAACTCAGACC in 0.25 M EDTA) was

added to quench the reaction. After 2 rounds of ligation-based barcoding, the pooled

nuclei were resuspended in 1x PBS with 0.5 U/μl Superase RNAse Inhibitor and 0.5

U/μl RNAaseOUT, counted and aliquoted into 10 μl sublibraries containing 3,000 nuclei.

For each sublibrary nuclei were lysed and cross-links reversed by adding 10 μl of lysis

buffer (10 mM Tris-pH 7.5, 1 M NaCl, 0.5 mM EDTA, 2% SDS) and incubating at 65°C

for 5 hrs. Biotin barcoded cDNA molecules were purified by adding 80 μl of dynabead

solution (10 ul MyOne Streptavidin C1 dynabeads (Invitrogen) in 10 mM Tris-pH 7.5, 1

M NaCl, 0.5 mM EDTA) and incubating for 1 h at 24°C at 1000 rpm. Bound cDNAs were

washed twice with 200 μl BW buffer (10 mM Tris-pH 7.5, 1 M NaCl, 0.5 mM EDTA)

followed by 2 additional washes with 200 μl Tris-T solution (10 mM Tris-HCl pH7.5,

0.1% Tween 20). For incorporating a 5' adaptor onto the streptavidin bound cDNA a 50

μl template switching mix (1x Maxima H buffer, 4% Ficoll, 1 mM dNTPs, 2.5 μM TS oligo

(AAGCAGTGGTATCAACGCAGAGTGAATrGrG+G where rG is RNA base), 500 U

Maxima H minus RT, 15U Superase RNAse Inhibitor and 20 U RNAaseOUT) was

added to each sublibrary. The bead solution was then incubated at 25°C for 30 minutes

and then at 42°C for 90 minutes with mixing at 1000 rpm. After the incubation beads

were washed twice with 200 μl Tris-T solution.

Washed beads were resuspended in 50 µl of PCR reaction mix (1x Q5 reaction buffer,
0.1 mM dNTPs, 0.5 uM ea Rev (AAGCAGTGGTATCAACGCAGAGT) and For
(CAGACGTGTGCTCTTCCGATCT) primer, 1U Q5 High-Fidelity DNA polymerase
(NEB)) and amplified with the following program: Step 1: 95°C × 1 min; Step 2 (6
cycles): 98°C × 30 s, 70°C × 15 s, 72°C × 3 min; Step 3: 72°C × 1 min. PCR reactions
were purified using a 0.85X ratio of SPRI beads (Beckman). To determine the optimal
PCR cycle to give 1/3rd of a saturated signal we repeated the PCR using 1/10th of the
purified PCR reaction and EvaGreen (Biotium) in a qPCR machine. Using the optimal
cycle we PCR amplified the remaining cDNA and purified the reactions using 0.85X ratio
of SPRI beads. To normalize the amplified cDNA fragment lengths for sequencing we
trimmed the libraries using Tn5 transposase (Diagenode) loaded with (5'-[Phos]
CTGTCTCTTATACACATCT, 5'-TCGTCGGCAGCGTCAGATGTGTATAAGAGACAG) to
a size range of 200-1000bp. Each trimmed sublibrary was then PCR amplified using a
unique   forward   index   P7Trueseq   barcode   oligo   (5'
CAAGCAGAAGACGGCATACGAGATXXXXXXGTGACTGGAGTTCAGACGTGTGCTCT
TCCGATC)   and   reverse   oligo
(AATGATACGGCGACCACCGAGATCTACACTAGATCGCTCGTCGGCAGCGTCAG).
Purified libraries were then sequenced on a NovaSeq 6000 using 100 bp paired end
reads and a read depth of 25-30,000 reads per cell.

Single-nucleus RNA-seq data preprocessing

The Paired-Tag pipeline as previously described [87] (https://github.com/cxzhu/Paired-
Tag) was used to process raw sequencing reads. Briefly this set of scripts takes raw
reads and extracts cellular barcodes from read2 and then uses read1 sequence to map
reads to human genome reference hg38. Finally demultiplexed and mapped data from
different sublibraries are merged to generate the final cell barcode gene count matrix.

Single-nucleus RNA-seq analysis

Single nucleus transcriptomes were analysed from hTLS at days 2-7 across 2
independent differentiation batches, with additional day 0 and day 7 samples from
independent batches, and undifferentiated SOX2:mNeon iPSC cells, included. Filtered
feature-barcode matrices of snRNA-seq data were converted into Seurat objects using

the Seurat R package (version 5.3.1) [88], merged into a single object, and quality control

filtered for cells containing > 500 and < 5000 genes and < 10% mitochondrial reads. An

average of ~1,500 genes and ~1% mitochondrial reads were detected per cell,

consistent across time points (Figures S1E,F). Gene expression counts were

normalised using SCTransform [89], followed by PCA. Visualisation and clustering were

performed using the functions RunUMAP and FindNeighbors, followed by FindClusters

using clustering resolution as stated in the figure legends. Clustering identified a

minority of clusters composed of cells with consistently lower gene detection relative to

the dataset average. These clusters lacked detectable expression of established

lineage marker genes and were excluded from downstream analyses due to insufficient

information for reliable cell type annotation. Mesodermal cells were isolated by

clustering the full filtered dataset at low resolution (0.03; Figure S1M), and re-clustered

using the top 41 principal components at a resolution of 0.5. Marker genes for each

cluster were identified with Wilcoxon rank sum test in Seurat (FindMarkers). For

differential expression analyses, thresholds of log2 fold change > 1 and adjusted p-

value < 0.005 (or 1e-8 for selected analyses) were applied, while full results were used

for visualisation in volcano plots. Cell identities were manually annotated by expression

of established canonical marker genes.

1480

Pseudotime analysis

To identify distinct mesodermal lineages arising during hTLS development, the

mesoderm subset was clustered with resolution 0.5, and Slingshot [36] trajectories were

inferred using cluster 0 as the root and clusters 2, 1, 6 and 4 as terminal states.

tradeSeq [37] was used to model gene expression dynamics along each trajectory and for

statistical analyses. To identify genes defining terminal cell states, the tradeSeq

endpoint differential expression test (diffEndTest) [37] was used, which tests for differential

expression between lineage endpoints. For each lineage, genes were ranked using a

combined endpoint significance score defined as the sum of -log10-transformed p-

values from the three pairwise endpoint contrasts involving that lineage. The top 20

ranked genes for each lineage were selected for visualisation. For heatmap

visualisation (Figure 1K), expression dynamics along pseudotime were obtained from
the fitted tradeSeq GAM as predicted expression values and transformed using log1p.
Gene expression trajectories were standardised by computing z-scores relative to the
gene's mean fitted expression across all pseudotime lineages. Heatmap values thus
represent global standardised expression, with positive values (red) indicating
expression above the gene's global mean and negative values (blue) indicating
expression below the mean. To reduce the visual influence of genes with low absolute
expression levels, standardised values were scaled by an expression-dependent weight
derived from the fitted log1p expression. Specifically, z-scores were multiplied by an
expression-dependent logistic weighting function derived from the fitted log1p
expression, thereby downweighting pseudotime regions with low fitted expression and
therefore less likely to reflect biologically meaningful expression. Weighted z-scores
were clipped to ±2.5 to limit the visual influence of extreme values.

1505

1506 Motif enrichment

Motif enrichment analysis was performed using HOMER (v4.11) [90] on ATAC-seq peaks
from early somite populations generated using the same differentiation protocol as
described by Loh et al [45]. Public ATAC-seq peaks corresponding to early somite cells
(GEO accession GSM2257295) were used as the accessible chromatin reference set.
Peaks located within ±50 kb of transcription start sites of genes significantly
downregulated following FOXC1/2 knockdown were identified using BEDTools. These
regions were tested for transcription factor motif enrichment relative to a background set
consisting of ATAC-seq peaks located within ±50 kb of transcription start sites of genes
expressed but not significantly altered following FOXC1/2 knockdown. Motif enrichment
was calculated using HOMER (findMotifsGenome.pl) with sequences extracted from the
hg19 human genome assembly. Significantly enriched motifs were ranked by
hypergeometric p-value and FDR. Enriched motifs were interpreted as candidate
transcriptional regulators acting at FOXC-dependent regulatory elements during early
somite specification.

1521

Mapping scRNA-seq to reference human embryogenesis datasets

We projected our data (query dataset) on two reference datasets: Zhao et al. for the earlier time points (days 0-4) [25], and Zeng et al. for the later ones (days 5-7) [24]. For the Zhao et al. reference, we used the projection pipeline provided by the authors themselves for projecting any user's query dataset onto their reference (R function: projecting_query_dataset.R). We modified only one parameter of the standard pipeline: para.list$cor.cutoff <- 0.2 instead of 0.5. We projected our datasets one sample at the time. The output of this pipeline matches each cell of the target to its closest location in the reference's umap, which allows to represent the reference landscape as published by the authors with the query dataset projected on top. For the Zeng et al dataset, we built our own projection pipeline, which consists of two parts: integration and projection. For the integration, we used the scVI method from the scVI-tools (10.1038/s41592-018-0229-2) to correct the raw counts of the reference and target datasets. Correcting the counts was necessary due to the different chemistries of the reference and the query datasets. Once the counts were corrected, we computed 50 pcs, that were then used for projecting one sample of the query datasets at the time, following these steps: we computed the distance matrix among the pcs of the reference and those of the query dataset; for each cell in a query sample, we gave a score 1 to its 15 nearest neighbours (NN) of the reference based on the distance matrix, and 0 otherwise; we computed the adjacency (or similarity) score of each cell of the reference to each cell of the query dataset as the average of the scores 1 (cell being a NN) or 0 (cell not being a NN). As a result, we can visualize the originally published umap of the reference with our own data projected on top in terms of similarity scores.

QUANTIFICATION AND STATISTICAL ANALYSIS

Statistical analyses of bulk RNA-seq data were performed with DESeq2. Analyses of snRNA-seq data were performed with Seurat. Marker genes were analysed with Wilcoxon rank sum test. Thresholds for volcano plots were set as |log2FoldChange| > 0.58 (±1.5-fold) and adjusted p-value < 0.05, and for heatmaps were set as

log2FoldChange >1 (>2-fold) and adjusted p-value < 1e-8 (Fig S1N) and adjusted p-

value < 0.005 (Fig S1P). Statistical analyses of confocal microscopy quantifications and

RT-qPCR data were performed within Prism. For image quantification, data were

analysed with Mann-Whitney test and visualised in plots with dots representing

individual hTLS and bars represent median ± IQR. For RT-qPCR analyses, individual

data points represent means of up to 4 technical replicates from a single independent

differentiation batch, which were used as 'n's for statistical analyses. Biological replicate

data were assumed to have a normal distribution, while to account for batch effects,

data were analysed using two-way ANOVA with batch as row factor and treatment as

column factor. RT-qPCR data are visualised in plots with bars representing mean ±

SEM. Asterisks in plots represent p values as follows: * ($p < 0.05$), ** ($p < 0.01$), *** ($p <$

0.001), **** ($p < 0.0001$). All the details of quantifications and statistical analyses are

fully described in the main text, figure legends, and method details section.

1564

References

1565

1566   1.   Ghimire, S., Mantziou, V., Moris, N., and Martinez Arias, A. (2021). Human
1567   gastrulation: The embryo and its models. Dev Biol *474*, 100–108.
1568   10.1016/j.ydbio.2021.01.006.

1569   2.   Sheng, G., Martinez Arias, A., and Sutherland, A. (2021). The primitive streak
1570   and cellular principles of building an amniote body through gastrulation. Science
1571   *374*, abg1727. 10.1126/science.abg1727.

1572   3.   Ferretti, E., and Hadjantonakis, A.K. (2019). Mesoderm specification and
1573   diversification: from single cells to emergent tissues. Curr Opin Cell Biol *61*, 110–
1574   116. 10.1016/j.ceb.2019.07.012.

1575   4.   Miao, Y., and Pourquie, O. (2024). Cellular and molecular control of vertebrate
1576   somitogenesis. Nat Rev Mol Cell Biol *25*, 517–533. 10.1038/s41580-024-00709-
1577   z.

1578   5.   Costantini, F., and Kopan, R. (2010). Patterning a complex organ: branching
1579   morphogenesis and nephron segmentation in kidney development. Dev Cell *18*,
1580   698–712. 10.1016/j.devcel.2010.04.008.

1581   6.   McMahon, A.P. (2016). Development of the Mammalian Kidney. Curr Top Dev
1582   Biol *117*, 31–64. 10.1016/bs.ctdb.2015.10.010.

1583   7.   Garcia-Martinez, V., and Schoenwolf, G.C. (1992). Positional control of
1584   mesoderm movement and fate during avian gastrulation and neurulation. Dev
1585   Dyn *193*, 249–256. 10.1002/aja.1001930305.

1586   8.   Kinder, S.J., Tsang, T.E., Quinlan, G.A., Hadjantonakis, A.K., Nagy, A., and Tam,
1587   P.P. (1999). The orderly allocation of mesodermal cells to the extraembryonic
1588   structures and the anteroposterior axis during gastrulation of the mouse embryo.
1589   Development *126*, 4691–4701. 10.1242/dev.126.21.4691.

1590   9.   Parameswaran, M., and Tam, P.P. (1995). Regionalisation of cell fate and
1591   morphogenetic movement of the mesoderm during mouse gastrulation. Dev
1592   Genet *17*, 16–28. 10.1002/dvg.1020170104.

1593   10.   Tam, P.P., and Beddington, R.S. (1987). The formation of mesodermal tissues in
1594   the mouse embryo during gastrulation and early organogenesis. Development
1595   *99*, 109–126. 10.1242/dev.99.1.109.

11.   Tam, P.P., Parameswaran, M., Kinder, S.J., and Weinberger, R.P. (1997). The
allocation of epiblast cells to the embryonic heart and other mesodermal
lineages: the role of ingression and tissue movement during gastrulation.
Development *124*, 1631–1642. 10.1242/dev.124.9.1631.

12.   Wilson, V., and Beddington, R.S. (1996). Cell fate and morphogenetic movement
in the late mouse primitive streak. Mech Dev *55*, 79–89. 10.1016/0925-
4773(95)00493-9.

13.   James, R.G., and Schultheiss, T.M. (2003). Patterning of the avian intermediate
mesoderm by lateral plate and axial tissues. Dev Biol *253*, 109–124.
10.1006/dbio.2002.0863.

14.   Duester, G. (2008). Retinoic acid synthesis and signaling during early
organogenesis. Cell *134*, 921–931. 10.1016/j.cell.2008.09.002.

15.   Yamaguchi, T.P. (2001). Heads or tails: Wnts and anterior-posterior patterning.
Curr Biol *11*, R713–724. 10.1016/s0960-9822(01)00417-1.

16.   Kim, S., Koppitch, K., Parvez, R.K., Guo, J., Achieng, M., Schnell, J., Lindstrom,
N.O., and McMahon, A.P. (2024). Comparative single-cell analyses identify
shared and divergent features of human and mouse kidney development. Dev
Cell *59*, 2912–2930 e2917. 10.1016/j.devcel.2024.07.013.

17.   Lindstrom, N.O., Tran, T., Guo, J., Rutledge, E., Parvez, R.K., Thornton, M.E.,
Grubbs, B., McMahon, J.A., and McMahon, A.P. (2018). Conserved and
Divergent Molecular and Anatomic Features of Human and Mouse Nephron
Patterning. J Am Soc Nephrol *29*, 825–840. 10.1681/ASN.2017091036.

18.   Lindstrom, N.O., McMahon, J.A., Guo, J., Tran, T., Guo, Q., Rutledge, E., Parvez,
R.K., Saribekyan, G., Schuler, R.E., Liao, C., et al. (2018). Conserved and
Divergent Features of Human and Mouse Kidney Organogenesis. J Am Soc
Nephrol *29*, 785–805. 10.1681/ASN.2017080887.

19.   Lindstrom, N.O., Guo, J., Kim, A.D., Tran, T., Guo, Q., De Sena Brandine, G.,
Ransick, A., Parvez, R.K., Thornton, M.E., Baskin, L., et al. (2018). Conserved
and Divergent Features of Mesenchymal Progenitor Cell Types within the Cortical
Nephrogenic Niche of the Human and Mouse Kidney. J Am Soc Nephrol *29*,
806–824. 10.1681/ASN.2017080890.

20.   Matsuda, M., Hayashi, H., Garcia-Ojalvo, J., Yoshioka-Kobayashi, K., Kageyama,
1628   R., Yamanaka, Y., Ikeya, M., Toguchida, J., Alev, C., and Ebisuya, M. (2020).
1629   Species-specific segmentation clock periods are due to differential biochemical
1630   reaction speeds. Science *369*, 1450–1455. 10.1126/science.aba7668.
1631   21.   Xi, H., Fujiwara, W., Gonzalez, K., Jan, M., Liebscher, S., Van Handel, B.,
1632   Schenke-Layland, K., and Pyle, A.D. (2017). In Vivo Human Somitogenesis
1633   Guides Somite Development from hPSCs. Cell Rep *18*, 1573–1585.
1634   10.1016/j.celrep.2017.01.040.
1635   22.   Rivron, N.C., Martinez Arias, A., Pera, M.F., Moris, N., and M'Hamdi H, I. (2023).
1636   An ethical framework for human embryology with embryo models. Cell *186*,
1637   3548–3557. 10.1016/j.cell.2023.07.028.
1638   23.   Tyser, R.C.V., Mahammadov, E., Nakanoh, S., Vallier, L., Scialdone, A., and
1639   Srinivas, S. (2021). Single-cell transcriptomic characterization of a gastrulating
1640   human embryo. Nature *600*, 285–289. 10.1038/s41586-021-04158-y.
1641   24.   Zeng, B., Liu, Z., Lu, Y., Zhong, S., Qin, S., Huang, L., Zeng, Y., Li, Z., Dong, H.,
1642   Shi, Y., et al. (2023). The single-cell and spatial transcriptional landscape of
1643   human gastrulation and early brain development. Cell Stem Cell *30*, 851–866
1644   e857. 10.1016/j.stem.2023.04.016.
1645   25.   Zhao, C., Plaza Reyes, A., Schell, J.P., Weltner, J., Ortega, N.M., Zheng, Y.,
1646   Bjorklund, A.K., Baque-Vidal, L., Sokka, J., Trokovic, R., et al. (2025). A
1647   comprehensive human embryo reference tool using single-cell RNA-sequencing
1648   data. Nat Methods *22*, 193–206. 10.1038/s41592-024-02493-2.
1649   26.   Gribaudo, S., Robert, R., van Sambeek, B., Mirdass, C., Lyubimova, A., Bouhali,
1650   K., Ferent, J., Morin, X., van Oudenaarden, A., and Nedelec, S. (2024). Self-
1651   organizing models of human trunk organogenesis recapitulate spinal cord and
1652   spine co-morphogenesis. Nat Biotechnol *42*, 1243–1253. 10.1038/s41587-023-
1653   01956-9.
1654   27.   Hamazaki, N., Yang, W., Kubo, C.A., Qiu, C., Martin, B.K., Garge, R.K.,
1655   Regalado, S.G., Nichols, E.K., Pendyala, S., Bradley, N., et al. (2024). Retinoic
1656   acid induces human gastruloids with posterior embryo-like structures. Nat Cell
1657   Biol *26*, 1790–1803. 10.1038/s41556-024-01487-8.

28.   Shahbazi, M.N., and Pasque, V. (2024). Early human development and stem cell-
based human embryo models. Cell Stem Cell *31*, 1398–1418.
10.1016/j.stem.2024.09.002.

29.   Makwana, K., Tilley, L., Chakravarty, P., Thompson, J., Baillie-Benson, P.,
Rodriguez-Polo, I., and Moris, N. (2025). Modelling co-development between the
somites and neural tube in human trunk-like structures. Nat Cell Biol *27*, 2049–
2062. 10.1038/s41556-025-01813-8.

30.   Wu, H., and Wang, H. (2026). Progress in stem cell-based embryo models and
their applications in developmental biology and biomedicine. Nat Rev Mol Cell
Biol *27*, 178–193. 10.1038/s41580-025-00942-0.

31.   Yamanaka, Y., Hamidi, S., Yoshioka-Kobayashi, K., Munira, S., Sunadome, K.,
Zhang, Y., Kurokawa, Y., Ericsson, R., Mieda, A., Thompson, J.L., et al. (2023).
Reconstituting human somitogenesis in vitro. Nature *614*, 509–520.
10.1038/s41586-022-05649-2.

32.   Miao, Y., Djeffal, Y., De Simone, A., Zhu, K., Lee, J.G., Lu, Z., Silberfeld, A., Rao,
J., Tarazona, O.A., Mongera, A., et al. (2023). Reconstruction and deconstruction
of human somitogenesis in vitro. Nature *614*, 500–508. 10.1038/s41586-022-
05655-4.

33.   Sanaki-Matsumiya, M., Matsuda, M., Gritti, N., Nakaki, F., Sharpe, J., Trivedi, V.,
and Ebisuya, M. (2022). Periodic formation of epithelial somites from human
pluripotent stem cells. Nat Commun *13*, 2325. 10.1038/s41467-022-29967-1.

34.   Moris, N., Anlas, K., van den Brink, S.C., Alemany, A., Schroder, J., Ghimire, S.,
Balayo, T., van Oudenaarden, A., and Martinez Arias, A. (2020). An in vitro model
of early anteroposterior organization during human development. Nature *582*,
410–415. 10.1038/s41586-020-2383-9.

35.   Zhang, X., Huang, C.T., Chen, J., Pankratz, M.T., Xi, J., Li, J., Yang, Y., Lavaute,
T.M., Li, X.J., Ayala, M., et al. (2010). Pax6 is a human neuroectoderm cell fate
determinant. Cell Stem Cell *7*, 90–100. 10.1016/j.stem.2010.04.017.

36.   Street, K., Risso, D., Fletcher, R.B., Das, D., Ngai, J., Yosef, N., Purdom, E., and
Dudoit, S. (2018). Slingshot: cell lineage and pseudotime inference for single-cell
transcriptomics. BMC Genomics *19*, 477. 10.1186/s12864-018-4772-0.

37.   Van den Berge, K., Roux de Bezieux, H., Street, K., Saelens, W., Cannoodt, R.,
1690   Saeys, Y., Dudoit, S., and Clement, L. (2020). Trajectory-based differential
1691   expression analysis for single-cell sequencing data. Nat Commun *11*, 1201.
1692   10.1038/s41467-020-14766-3.

1693   38.   Chapman, D.L., Agulnik, I., Hancock, S., Silver, L.M., and Papaioannou, V.E.
1694   (1996). Tbx6, a mouse T-Box gene implicated in paraxial mesoderm formation at
1695   gastrulation. Dev Biol *180*, 534–542. 10.1006/dbio.1996.0326.

1696   39.   Sadahiro, T., Isomi, M., Muraoka, N., Kojima, H., Haginiwa, S., Kurotsu, S.,
1697   Tamura, F., Tani, H., Tohyama, S., Fujita, J., et al. (2018). Tbx6 Induces Nascent
1698   Mesoderm from Pluripotent Stem Cells and Temporally Controls Cardiac versus
1699   Somite Lineage Diversification. Cell Stem Cell *23*, 382–395 e385.
1700   10.1016/j.stem.2018.07.001.

1701   40.   Tang, W.W.C., Castillo-Venzor, A., Gruhn, W.H., Kobayashi, T., Penfold, C.A.,
1702   Morgan, M.D., Sun, D., Irie, N., and Surani, M.A. (2022). Sequential enhancer
1703   state remodelling defines human germline competence and specification. Nat
1704   Cell Biol *24*, 448–460. 10.1038/s41556-022-00878-z.

1705   41.   Nabet, B., Roberts, J.M., Buckley, D.L., Paulk, J., Dastjerdi, S., Yang, A., Leggett,
1706   A.L., Erb, M.A., Lawlor, M.A., Souza, A., et al. (2018). The dTAG system for
1707   immediate and target-specific protein degradation. Nat Chem Biol *14*, 431–441.
1708   10.1038/s41589-018-0021-8.

1709   42.   Wilm, B., James, R.G., Schultheiss, T.M., and Hogan, B.L. (2004). The forkhead
1710   genes, Foxc1 and Foxc2, regulate paraxial versus intermediate mesoderm cell
1711   fate. Dev Biol *271*, 176–189. 10.1016/j.ydbio.2004.03.034.

1712   43.   Kume, T., Deng, K., and Hogan, B.L. (2000). Murine forkhead/winged helix genes
1713   Foxc1 (Mf1) and Foxc2 (Mfh1) are required for the early organogenesis of the
1714   kidney and urinary tract. Development *127*, 1387–1395. 10.1242/dev.127.7.1387.

1715   44.   Kume, T., Jiang, H., Topczewska, J.M., and Hogan, B.L. (2001). The murine
1716   winged helix transcription factors, Foxc1 and Foxc2, are both required for
1717   cardiovascular development and somitogenesis. Genes Dev *15*, 2470–2482.
1718   10.1101/gad.907301.

1719   45.   Loh, K.M., Chen, A., Koh, P.W., Deng, T.Z., Sinha, R., Tsai, J.M., Barkal, A.A.,

1720   Shen, K.Y., Jain, R., Morganti, R.M., et al. (2016). Mapping the Pairwise Choices

1721   Leading from Pluripotency to Human Bone, Heart, and Other Mesoderm Cell

1722   Types. Cell *166*, 451–467. 10.1016/j.cell.2016.06.011.

1723   46.   Asp, M., Giacomello, S., Larsson, L., Wu, C., Furth, D., Qian, X., Wardell, E.,

1724   Custodio, J., Reimegard, J., Salmen, F., et al. (2019). A Spatiotemporal Organ-

1725   Wide Gene Expression and Cell Atlas of the Developing Human Heart. Cell *179*,

1726   1647–1660 e1619. 10.1016/j.cell.2019.11.025.

1727   47.   Stewart, B.J., Ferdinand, J.R., Young, M.D., Mitchell, T.J., Loudon, K.W., Riding,

1728   A.M., Richoz, N., Frazer, G.L., Staniforth, J.U.L., Vieira Braga, F.A., et al. (2019).

1729   Spatiotemporal immune zonation of the human kidney. Science *365*, 1461–1466.

1730   10.1126/science.aat5031.

1731   48.   Fan, C.M., Porter, J.A., Chiang, C., Chang, D.T., Beachy, P.A., and Tessier-

1732   Lavigne, M. (1995). Long-range sclerotome induction by sonic hedgehog: direct

1733   role of the amino-terminal cleavage product and modulation by the cyclic AMP

1734   signaling pathway. Cell *81*, 457–465. 10.1016/0092-8674(95)90398-4.

1735   49.   Johnson, R.L., Laufer, E., Riddle, R.D., and Tabin, C. (1994). Ectopic expression

1736   of Sonic hedgehog alters dorsal-ventral patterning of somites. Cell *79*, 1165–

1737   1173. 10.1016/0092-8674(94)90008-6.

1738   50.   Yang, J., Mani, S.A., Donaher, J.L., Ramaswamy, S., Itzykson, R.A., Come, C.,

1739   Savagner, P., Gitelman, I., Richardson, A., and Weinberg, R.A. (2004). Twist, a

1740   master regulator of morphogenesis, plays an essential role in tumor metastasis.

1741   Cell *117*, 927–939. 10.1016/j.cell.2004.06.006.

1742   51.   Thiery, J.P., Acloque, H., Huang, R.Y., and Nieto, M.A. (2009). Epithelial-

1743   mesenchymal transitions in development and disease. Cell *139*, 871–890.

1744   10.1016/j.cell.2009.11.007.

1745   52.   Row, R.H., Pegg, A., Kinney, B.A., Farr, G.H., 3rd, Maves, L., Lowell, S., Wilson,

1746   V., and Martin, B.L. (2018). BMP and FGF signaling interact to pattern mesoderm

1747   by controlling basic helix-loop-helix transcription factor activity. Elife *7*.

1748   10.7554/eLife.31018.

1749   53.   Chapman, D.L., and Papaioannou, V.E. (1998). Three neural tubes in mouse
1750   embryos with mutations in the T-box gene Tbx6. Nature *391*, 695–697.
1751   10.1038/35624.

1752   54.   Takemoto, T., Uchikawa, M., Yoshida, M., Bell, D.M., Lovell-Badge, R.,
1753   Papaioannou, V.E., and Kondoh, H. (2011). Tbx6-dependent Sox2 regulation
1754   determines neural or mesodermal fate in axial stem cells. Nature *470*, 394–398.
1755   10.1038/nature09729.

1756   55.   Liu, J., Wu, N., Deciphering Disorders Involving, S., study, C.O., Yang, N.,
1757   Takeda, K., Chen, W., Li, W., Du, R., Liu, S., et al. (2019). TBX6-associated
1758   congenital scoliosis (TACS) as a clinically distinguishable subtype of congenital
1759   scoliosis: further evidence supporting the compound inheritance and TBX6 gene
1760   dosage model. Genet Med *21*, 1548–1558. 10.1038/s41436-018-0377-x.

1761   56.   Yang, N., Wu, N., Zhang, L., Zhao, Y., Liu, J., Liang, X., Ren, X., Li, W., Chen, W.,
1762   Dong, S., et al. (2019). TBX6 compound inheritance leads to congenital vertebral
1763   malformations in humans and mice. Hum Mol Genet *28*, 539–547.
1764   10.1093/hmg/ddy358.

1765   57.   Wu, N., Ming, X., Xiao, J., Wu, Z., Chen, X., Shinawi, M., Shen, Y., Yu, G., Liu, J.,
1766   Xie, H., et al. (2015). TBX6 null variants and a common hypomorphic allele in
1767   congenital scoliosis. N Engl J Med *372*, 341–350. 10.1056/NEJMoa1406829.

1768   58.   Garza Flores, A., Nordgren, I., Pettersson, M., Dias-Santagata, D., Nilsson, D.,
1769   Hammarsjo, A., Lindstrand, A., Batkovskyte, D., Wiggs, J., Walton, D.S., et al.
1770   (2023). Case report: Extending the spectrum of clinical and molecular findings in
1771   FOXC1 haploinsufficiency syndrome. Front Genet *14*, 1174046.
1772   10.3389/fgene.2023.1174046.

1773   59.   Reis, L.M., Maheshwari, M., Capasso, J., Atilla, H., Dudakova, L., Thompson, S.,
1774   Zitano, L., Lay-Son, G., Lowry, R.B., Black, J., et al. (2023). Axenfeld-Rieger
1775   syndrome: more than meets the eye. J Med Genet *60*, 368–379. 10.1136/jmg-
1776   2022-108646.

1777   60.   Concepcion, D., Washkowitz, A.J., DeSantis, A., Ogea, P., Yang, J.I., Douglas,
1778   N.C., and Papaioannou, V.E. (2017). Cell lineage of timed cohorts of Tbx6-

expressing cells in wild-type and Tbx6 mutant embryos. Biol Open *6*, 1065–1073.

10.1242/bio.026203.

61. Sandbacka, M., Laivuori, H., Freitas, E., Halttunen, M., Jokimaa, V., Morin-

Papunen, L., Rosenberg, C., and Aittomaki, K. (2013). TBX6, LHX1 and copy

number variations in the complex genetics of Mullerian aplasia. Orphanet J Rare

Dis *8*, 125. 10.1186/1750-1172-8-125.

62. Dong, S., Wang, C., Li, X., Shen, Q., Fu, X., Wu, M., Song, C., Yang, N., Wu, Y.,

Wang, H., et al. (2019). Noncoding rare variants of TBX6 in congenital anomalies

of the kidney and urinary tract. Mol Genet Genomics *294*, 493–500.

10.1007/s00438-018-1522-6.

63. Ma, C., Chen, N., Jolly, A., Zhao, S., Coban-Akdemir, Z., Tian, W., Kang, J., Ye,

Y., Wang, Y., Koch, A., et al. (2022). Functional characteristics of a broad

spectrum of TBX6 variants in Mayer-Rokitansky-Kuster-Hauser syndrome. Genet

Med *24*, 2262–2273. 10.1016/j.gim.2022.08.012.

64. Li, G., Strong, A., Wang, H., Kim, J.S., Watson, D., Zhao, S., Vaccaro, C.,

Hartung, E., Hakonarson, H., Zhang, T.J., et al. (2022). TBX6 as a cause of a

combined skeletal-kidney dysplasia syndrome. Am J Med Genet A *188*, 3469–

3481. 10.1002/ajmg.a.62972.

65. Chapman, D.L. (2019). Impaired intermediate formation in mouse embryos

expressing reduced levels of Tbx6. Genesis *57*, e23270. 10.1002/dvg.23270.

66. Hayashi, S., Suzuki, H., and Takemoto, T. (2021). The nephric mesenchyme

lineage of intermediate mesoderm is derived from Tbx6-expressing derivatives of

neuro-mesodermal progenitors via BMP-dependent Osr1 function. Dev Biol *478*,

155–162. 10.1016/j.ydbio.2021.07.006.

67. Uenaka, T., Napole, A., Saha, A.D., Sun, D., Singavarapu, A., Calzada, E., Chen,

J., Erlebach, L., McQuade, A., Ramos, D.M., et al. (2026). Prevention of

transgene silencing during human pluripotent stem cell differentiation. Cell Stem

Cell. 10.1016/j.stem.2026.01.007.

68. Ormestad, M., Astorga, J., and Carlsson, P. (2004). Differences in the embryonic

expression patterns of mouse Foxf1 and -2 match their distinct mutant

phenotypes. Dev Dyn *229*, 328–333. 10.1002/dvdy.10426.

69. Funayama, N., Sato, Y., Matsumoto, K., Ogura, T., and Takahashi, Y. (1999).
Coelom formation: binary decision of the lateral plate mesoderm is controlled by
the ectoderm. Development *126*, 4129–4138. 10.1242/dev.126.18.4129.

70. Huang, L., Huang, Z., Fan, Y., He, L., Ye, M., Shi, K., Ji, B., Huang, J., Wang, Y.,
and Li, Q. (2017). FOXC1 promotes proliferation and epithelial-mesenchymal
transition in cervical carcinoma through the PI3K-AKT signal pathway. Am J
Transl Res *9*, 1297–1306.

71. Ou-Yang, L., Xiao, S.J., Liu, P., Yi, S.J., Zhang, X.L., Ou-Yang, S., Tan, S.K., and
Lei, X. (2015). Forkhead box C1 induces epithelial◻mesenchymal transition and
is a potential therapeutic target in nasopharyngeal carcinoma. Mol Med Rep *12*,
8003–8009. 10.3892/mmr.2015.4427.

72. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold
change and dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.
10.1186/s13059-014-0550-8.

73. Berry, F.B., Saleem, R.A., and Walter, M.A. (2002). FOXC1 transcriptional
regulation is mediated by N- and C-terminal activation domains and contains a
phosphorylated transcriptional inhibitory domain. J Biol Chem *277*, 10292–
10297. 10.1074/jbc.M110266200.

74. Ng-Blichfeldt, J.P., Stewart, B.J., Clatworthy, M.R., Williams, J.M., and Roper, K.
(2024). Identification of a core transcriptional program driving the human renal
mesenchymal-to-epithelial transition. Dev Cell *59*, 595–612 e598.
10.1016/j.devcel.2024.01.011.

75. Takashima, Y., Guo, G., Loos, R., Nichols, J., Ficz, G., Krueger, F., Oxley, D.,
Santos, F., Clarke, J., Mansfield, W., et al. (2014). Resetting transcription factor
control circuitry toward ground-state pluripotency in human. Cell *158*, 1254–
1269. 10.1016/j.cell.2014.08.029.

76. Lovell-Badge, R., Anthony, E., Barker, R.A., Bubela, T., Brivanlou, A.H.,
Carpenter, M., Charo, R.A., Clark, A., Clayton, E., Cong, Y., et al. (2021). ISSCR
Guidelines for Stem Cell Research and Clinical Translation: The 2021 update.
Stem Cell Reports *16*, 1398–1408. 10.1016/j.stemcr.2021.05.012.

77.   Schmid-Burgk, J.L., Honing, K., Ebert, T.S., and Hornung, V. (2016). CRISPaint
1841   allows modular base-specific gene tagging using a ligase-4-dependent
1842   mechanism. Nat Commun *7*, 12338. 10.1038/ncomms12338.

1843   78.   Effern, M., Glodde, N., Braun, M., Liebing, J., Boll, H.N., Yong, M., Bawden, E.,
1844   Hinze, D., van den Boorn-Konijnenberg, D., Daoud, M., et al. (2020). Adoptive T
1845   Cell Therapy Targeting Different Gene Products Reveals Diverse and Context-
1846   Dependent Immune Evasion in Melanoma. Immunity *53*, 564–580 e569.
1847   10.1016/j.immuni.2020.07.007.

1848   79.   Perez-Pinera, P., Kocak, D.D., Vockley, C.M., Adler, A.F., Kabadi, A.M., Polstein,
1849   L.R., Thakore, P.I., Glass, K.A., Ousterout, D.G., Leong, K.W., et al. (2013). RNA-
1850   guided gene activation by CRISPR-Cas9-based transcription factors. Nat
1851   Methods *10*, 973–976. 10.1038/nmeth.2600.

1852   80.   Severin, J., Lizio, M., Harshbarger, J., Kawaji, H., Daub, C.O., Hayashizaki, Y.,
1853   Consortium, F., Bertin, N., and Forrest, A.R. (2014). Interactive visualization and
1854   analysis of large-scale sequencing datasets using ZENBU. Nat Biotechnol *32*,
1855   217–219. 10.1038/nbt.2840.

1856   81.   Hodgkins, A., Farne, A., Perera, S., Grego, T., Parry-Smith, D.J., Skarnes, W.C.,
1857   and Iyer, V. (2015). WGE: a CRISPR database for genome engineering.
1858   Bioinformatics *31*, 3078–3080. 10.1093/bioinformatics/btv308.

1859   82.   Gilbert, L.A., Horlbeck, M.A., Adamson, B., Villalta, J.E., Chen, Y., Whitehead,
1860   E.H., Guimaraes, C., Panning, B., Ploegh, H.L., Bassik, M.C., et al. (2014).
1861   Genome-Scale CRISPR-Mediated Control of Gene Repression and Activation.
1862   Cell *159*, 647–661. 10.1016/j.cell.2014.09.029.

1863   83.   Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression
1864   data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method.
1865   Methods *25*, 402–408. 10.1006/meth.2001.1262.

1866   84.   Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A.,
1867   and Goodman, A. (2021). CellProfiler 4: improvements in speed, utility and
1868   usability. BMC Bioinformatics *22*, 433. 10.1186/s12859-021-04344-9.

1869   85.   Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W.,
1870   Zhan, L., et al. (2021). clusterProfiler 4.0: A universal enrichment tool for

interpreting omics data. Innovation (Camb) *2*, 100141.

10.1016/j.xinn.2021.100141.

86. Rosenberg, A.B., Roco, C.M., Muscat, R.A., Kuchina, A., Sample, P., Yao, Z.,

Graybuck, L.T., Peeler, D.J., Mukherjee, S., Chen, W., et al. (2018). Single-cell

profiling of the developing mouse brain and spinal cord with split-pool barcoding.

Science *360*, 176–182. 10.1126/science.aam8999.

87. Zhu, C., Zhang, Y., Li, Y.E., Lucero, J., Behrens, M.M., and Ren, B. (2021). Joint

profiling of histone modifications and transcriptome in single cells from mouse

brain. Nat Methods *18*, 283–292. 10.1038/s41592-021-01060-3.

88. Hao, Y., Stuart, T., Kowalski, M.H., Choudhary, S., Hoffman, P., Hartman, A.,

Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., and Satija, R.

(2024). Dictionary learning for integrative, multimodal and scalable single-cell

analysis. Nat Biotechnol *42*, 293–304. 10.1038/s41587-023-01767-y.

89. Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of

single-cell RNA-seq data using regularized negative binomial regression.

Genome Biol *20*, 296. 10.1186/s13059-019-1874-1.

90. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X.,

Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-

determining transcription factors prime cis-regulatory elements required for

macrophage and B cell identities. Mol Cell *38*, 576–589.

10.1016/j.molcel.2010.05.004.

1892

Figure 1

A Human iPSC-derived 3D hTLS differentiation

B (Day 1–Day 5, Brightfield and SOX2:mNeon)

C Day 5 — FOXC2, SOX2, PAX8, DAPI

D Day 8 — ZO1, PAX8, DACH2, SOX2

E Time-course snRNA-seq

E' UMAP — Mesoderm, Endothelium, Neural, PS

F Transcriptional similarity of hTLS (Days 5-7) to human PCW3 embryo

G Mesoderm subset

H Mesoderm subset

I Paraxial mesoderm derivatives / Intermediate mesoderm derivatives

J Pseudotime trajectories

K Expression dynamics of lineage-defining endpoint genes

L MECOM, WT1, SULF1, TTN

A

B Inducible TBX6-overexpression strategy

C *TBX6* overexpression, first 24h of hTLS protocol

D Day 0 (24 hours of differentiation +/- Dox)

Control 0.1μM Dox

E Differential gene expression following TBX6 overexpression

F GSEA of human PCW2 embryo lineage signatures following TBX6 overexpression

A Mesoderm subset

B Somite progenitor vs rest of cells

C Inducible dual-gene CRISPRi strategy for FOXC1 and FOXC2

D *2D somitic mesoderm differentiation*

E Dual FOXC1 and FOXC2 knockdown

F Motif families enriched at FOXC1/2-dependent regulatory regions

G *3D hTLS differentiation*

H SOX2⁺ nuclei

I PAX8⁺ nuclei

J PAX3⁺ SOX2⁻ nuclei

K Dual FOXC1 and FOXC2 knockdown

A 2D somitic mesoderm differentiation

B RT-qPCR time courses

F Somite genes downregulated · Kidney genes upregulated · LPM/cardiomyocyte/SM genes upregulated

C

D

E Pearson correlation

G Early mesodermal genes

H GO biological processes after dual FOXC1/2 knockdown

I GSEA of human PCW3–5 lineage signatures following FOXC1/2 knockdown

J GSEA of human PCW6.5-7 heart signatures following FOXC1/2 knockdown

K GSEA of human fetal nephron signatures following FOXC1/2 knockdown

L Day 5 hTLS, dual FOXC1 and FOXC2 knockdown (Dox from day 2)

M Testing TBX6-FOXC1/2 epistasis

N

A *Somite diversification* in vivo:

Neural tube

Somite

Noto-chord

HH

EMT

Pre-skin/muscle (dermomyotome)
PAX3+

Pre-cartilage/bone (sclerotome)
FOXC1+ FOXC2+ TWIST1+
PAX1+ PAX9+ SOX9+

B *SHH* *IHH* *DHH* *GLI1* *GLI2* *GLI3*

C
3D hTLS differentiation
+/- SAG (500nM)
Day -2 -1 0 2 5 8
2D culture *3D culture*

E SOX2+ nuclei
SOX2+ nuclei per image
Control 500nM SAG
***

F FOXC2+ nuclei
FOXC2+ nuclei per image
Control 500nM SAG
**

G PAX3+ SOX2- nuclei
PAX3+ SOX2- nuclei per image
Control 500nM SAG
**

D Day 8
Merge | SOX2 (neural) | FOXC2 (somitic mesoderm) | PAX3 (somite / dorsal neural)
Control
+SAG (500nM)

H Day 5
SOX2
SOX9
FOXC2
[FOXC1]:mCherry
[FOXC1]::mCherry | FOXC2 (somite / sclerotome) | SOX9 (sclerotome)

I FOXC2 overexpression
ITR cHS4 TRE3G FOXC2 mCherry T2A pA cHS4 pEF1α rtTA T2A PuroR pA cHS4 ITR

Day 5
SOX2
PAX8
PAX3
[FOXC2]::mCherry
[FOXC2]::mCherry | PAX8 | PAX3

A TWIST1 overexpression

B *3D hTLS differentiation*

C [TWIST1]::mCherry / F-actin / FOXC2 / SOX2 · [TWIST1]::mCherry · FOXC2 · F-actin

D *2D somite-to-sclerotome differentiation*

E *FOXC1* *FOXC2* *TWIST1* *TWIST2* *PAX1* *PAX9* *SOX9* *PTCH1* *GLI1*