A 'capture-handling-restraint protocol' elicits short-term stress responses in

wild great tits (*Parus major*) but has little impact on reproductive success

Franziska Fröhlich[1*], Lucia Mentesana[1,2*¥], Caroline Deimel[1,3] and Michaela Hau[1,4†]

[1] Max-Planck-Institute for Biological Intelligence, Germany.

[2] Biology Department, Facultad de Ciencias, Universidad de la República, Uruguay.

[3] Department of Behavioral and Cognitive Biology, University of Vienna, Austria.

[4] Department of Biology, University of Konstanz, Germany.

Franziska Fröhlich: franziska.froehlich.pr@gmail.com (https://orcid.org/0009-0006-8440-
7992)

Dr. Lucia Mentesana: lucia.mentesana@fcien.edu.uy (https://orcid.org/0000-0003-4090-2570)

Caroline Deimel: caroline.deimel@univie.ac.at (https://orcid.org/0000-0001-9450-7088)

Prof. Dr. Michaela Hau (https://orcid.org/0000-0002-3836-4083)

[*] Franziska Fröhlich and Lucia Mentesana share first authorship.

[†]Deceased; contributions made before 09/2025.

[¥]Author for correspondence: lucia.mentesana@fcien.edu.uy

Abstract

Capturing and handling wild animals is essential for ecological and evolutionary research, yet their effects on physiology, behaviour, and reproductive success remain poorly understood. We investigated short- and longer-term consequences of a 'capture-handling-restraint protocol' in wild great tits (*Parus major*) over three breeding seasons. To assess short-term responses, we measured circulating corticosterone, a metabolic hormone that responds to unpredictable challenges, and automatically recorded provisioning behaviour. We also explored whether environmental and individual traits were related to provisioning latency (i.e., time to resume provisioning after capture). To evaluate longer-term effects, we monitored provisioning in the days following capture and related it to reproductive success (fledgling number and body condition). We predicted that longer handling would increase stress-induced corticosterone and provisioning latency, that these variables would be positively correlated, and that higher corticosterone and longer latencies would be associated with lower reproductive success. After capture, great tits showed elevated corticosterone and delayed provisioning. Contrary to our predictions, handling duration was negatively associated with stress-induced corticosterone in males (but not females) and did not affect provisioning latency. Provisioning latency was unrelated to corticosterone, environmental, or individual variables. Following capture, parents resumed provisioning, and short-term responses had little influence on reproductive success. We show that parental behaviour and physiology are affected by capture restraint protocols on the short term, but offspring condition and survival are not. However, these results should be interpreted cautiously, as our study lacks an uncaptured control group. Our findings highlight that evaluating welfare impacts requires rigorous study design incorporating both immediate and longer-term behavioural and fitness effects.

Keywords: Wild Animal Welfare, Corticosterone, Provisioning, Reproductive Success, RFID

1. Introduction

As scientists, we bear responsibility not only towards research integrity but also towards the welfare of the animals we study. Ethical evaluations of research procedures are guided by the principles of the 'Three Rs' (Replacement, Refinement and Reduction; Russell & Burch, 1959), which were initially established for laboratory animals and later expanded into the 'Nine Rs' for studies involving wildlife (Curzer et al., 2013). The primary goal of these principles is to minimise harm experienced by animals used in research, as physiological and behavioural indicators suggest that animals perceive these procedures as negative affective states (Broom & Johnson, 2019; reviewed by Kaplan, 2022; Soulsbury et al., 2020). Negative welfare states can affect the validity, generalisation, and interpretation of research results (Poole, 1997; Prescott and Lidster, 2017). Moreover, when animals are subjected to research experiments during reproductive life-history stages, the effects of research activities might extend beyond the animal used for research and potentially also affect the well-being of their offspring.

Capturing, sampling, handling and restraining wild animals is a common practice in biological research (Boitani and Fuller, 2000; Duarte, 2013; Soulsbury et al., 2020). These human-animal interactions are conceptualised as eliciting negative experiences in the affected animals, resulting in physiological and behavioural changes that are mounted in response to this event that can be used as welfare indicators (Mellor et al., 2020). Yet our understanding of how these procedures affect the short-term behavioural responses of wild research animals remains incomplete (Gillies et al., 2020; Schlicht and Kempenaers, 2015; Soulsbury et al., 2020), particularly in relation to how these effects align with well-documented physiological responses. This is important because the application of the 'Three Rs', or rather 'Nine Rs' principles, relies on accurately documenting the adverse effects of procedures that animals involved in research experience.

The short-term physiological effects of 'capture-handling-restraint protocols' (Wingfield et al., 1982) are well documented in vertebrates. Upon encountering an unexpected environmental challenge (e.g., being captured by researchers), catecholamines are rapidly released from the adrenal cortex to mediate the Fight-or-Flight response (Romero and Butler, 2007). Even short periods of high adrenal activity can weaken the immune response, potentially increasing mortality (Broom and Johnson, 2019). As a second, slower response to a challenge, glucocorticoid levels begin to rise within three to five minutes (Wingfield et al., 1982). They typically peak between 30 and 60 minutes, reaching what is commonly referred to as 'stress-induced glucocorticoid levels' (reviewed by Romero, 2004; Romero & Reed, 2005). Since glucocorticoids can be more easily measured than the rapidly changing catecholamines,

glucocorticoids are widely used as indicators of the physiological stress response (reviewed by
De Kloet et al., 2008; reviewed by Dickens & Romero, 2013; Huber et al., 2021; Romero &
Beattie, 2022; reviewed by Sapolsky et al., 2000; reviewed by Wingfield & Sapolsky, 2003).
Elevated glucocorticoid levels (i.e., allostatic state) enhance the action of catecholamines and
modulate physiological and behavioural responses, mediating allostasis, a process to maintain
stability through change, helping the animal to cope with the stressor (McEwen and Wingfield,
2003). The cumulative cost of coping adds to the required energy (i.e., allostatic load). If the
energy demand exceeds the energy supply, leading to an allostatic overload and resulting in an
emergency life history stage, immediate survival can be promoted at the expense of functions
such as reproduction (McEwen & Wingfield, 2003; Romero & Beattie, 2022; Romero &
Wingfield, 2015; Wingfield et al., 1998; reviewed by Wingfield & Sapolsky, 2003). On the
contrary, the short-term behavioural effects of 'capture-handling-restraint protocols' are poorly
documented in vertebrates. For example, blue tits (*Cyanistes caeruleus*) exposed to this
protocol during the breeding period stayed away much longer from the nest after being released
(on average 4.2 hours) compared to their normal intervals between nest visits (Schlicht and
Kempenaers, 2015). Although this study did not examine the relationship between short-term
behavioural changes and physiological parameters, these behavioural responses seem to be
contingent on physiological responses (i.e., the activation of the hypothalamic-pituitary-adrenal
(HPA) axis and subsequent increases in glucocorticoids).

During the breeding season, short-term effects of 'capture-handling-restraint protocols'
may also lead to longer-term consequences by altering parental behaviour, thereby indirectly
affecting offspring development and survival. Experimentally induced short-term physiological
stress responses have been associated with reduced reproductive success. In pied flycatchers
(*Ficedula hypoleuca*), experimentally increased glucocorticoid levels to stress-induced levels
during the nestling period led to reduced feeding and fewer surviving fledglings (Silverin,
1986). In black-legged kittiwakes (*Rissa tridactyla*), an experimentally induced short-term
increase in corticosterone, a type of glucocorticoid, during the chick-rearing period led to
decreased prolactin levels, which were associated with reduced nest attendance and increased
latency to return to the nest, which in turn significantly reduced breeding success (Angelier et
al., 2009). Similarly, in mammals, elevated cortisol, another type of glucocorticoid, reduced
maternal care in postpartum marmosets (*Callithrix jacchus*), which carried their infants less
than control females (Saltzman and Abbott, 2009). However, there seems to be little evidence
that intrinsic stress-induced increases in glucocorticoids following 'capture-handling-restraint
protocols' negatively affect breeding success. For example, eastern bluebird (*Sialia sialis*)

females exhibited increased corticosterone levels following a 'capture-handling-restraint protocol', but these did not affect fledging success (Lynn et al., 2010). Likewise, Lane et al. (2025) showed that incubating female song sparrows (*Melospiza melodia*) exhibited elevated glucocorticoid levels and longer intervals between nest visits following capture. Still, glucocorticoid levels did not explain the latency to return to the nest after the capture event. On the other hand, parental short-term behavioural stress responses to 'capture-handling-restraint protocols' may also lead to longer-term consequences affecting offspring condition and survival. Nestlings may have to wait unusually long for parents to return after capture and during this time they are exposed to adverse conditions that cause negative affective states, such as hunger and cold. This short-term exposure to adverse conditions can potentially negatively affect proper development and survival on the long term. These consequences may persist if parents return to the nest after the capture event but provide reduced care compared to their behaviour before capture. However, empirical evidence from wild populations addressing these effects remains limited. One notable example is the study by Schlicht and Kempenaers (2015), which found that although blue tits stayed away from their nests for extended periods after release, parents resumed normal feeding rates upon returning and no negative association with breeding success was detected.

Since a 'capture-handling-restraint protocol' can affect multiple welfare indicators over different time scales, studies assessing their short- and longer-term physiological and behavioural relationships and their links to reproductive success are greatly needed. For birds, despite their common use as model species in field studies, such research remains limited.

In this study, we therefore investigated short- and longer-term relationships of a 'capture-handling-restraint protocol' with physiological and behavioural traits in a wild population of great tits (*Parus major;* Figure 1A). We conducted this study during the nestling provisioning period across three consecutive breeding seasons (2020-2022). To do so, we captured adult great tits on days seven to twelve after their eggs had hatched. We subjected them to a standardised 'capture-handling-restraint protocol', which included blood sampling and is commonly used in wild bird research (Figure 1B). After catching an individual, we took a first blood sample within three minutes to measure baseline corticosterone levels. Then, we marked and tagged the bird to allow for the unequivocal identification and restrained it in a cloth bag for 20-30 minutes. After this, we took the bird's morphometric measures and a second blood sample to assess stress-induced corticosterone levels. Finally, we released the bird back into its nestbox. We automatically recorded feeding behaviour and nest abandonment by equipping each adult great tit with RFID leg transponders (Figure 1C) and using RFID antenna

devices located at the nest-box entrance (125 kHz, NatureCounters, UK; Figure 1D). To assess reproductive success, we measured nestling condition on day 15 or 16 post-hatching and determined the final number of fledglings at fledging.

First, we predicted that increases in handling duration would be correlated with higher stress-induced corticosterone levels, a greater likelihood of nest abandonment and increased time to resume provisioning (i.e., provisioning latency) after the catching event (Canoine et al., 2002; Monaghan & Spencer, 2014; reviewed by Romero, 2004). Because stress-induced corticosterone levels are suggested to support behavioural responses to cope with a stressor (reviewed by Andrews, 2022; Veen & Sivars, 2000; Wingfield et al., 1998), we predicted provisioning latency to increase with higher stress-induced corticosterone levels.

Secondly, an individual's capacity to respond to the stress of capture and handling is contingent on its own body condition, health, life history traits and perceived trade-offs, as well as environmental mediators. Therefore, we examined whether provisioning latency was explained by ambient temperature at capture, parental condition or brood quality. For parental condition, we considered variables such as age, parasite infestation and mass and tarsus length, as these factors influence an individual's allostatic load and therefore its capacity to sustain physiological and behavioural resources that are required for a response (McEwen & Wingfield, 2003; reviewed by Wu et al., 2025).

Thirdly, we examined if the 'capture-handling-restraint protocol' had longer-term relationships with behaviour (i.e., average time between two consecutive nest visits by a parent) and offspring condition (i.e., nest success, fledgling number, fledgling mass and tarsus length) and survival. Following (Schlicht and Kempenaers, 2015), we predicted that birds that returned to their nests after being captured would resume normal provisioning activity on the days following the capture. We also predicted that higher stress-induced corticosterone levels and longer provisioning latencies would be related to poorer nestling condition and decreased survival (Keller and Noordwijk, 1993; Monaghan and Spencer, 2014).

Lastly, because both male and female great tits care for the nestlings and nest success depends on contributions from both parents (Hinde, 2006; Norris, 1990), we examined whether nest abandonment by one parent would reduce fledging success. We predicted that the abandonment by one parent leads to a reduced fledging success, but not necessarily to complete nest failure.

Figure 1. (**A**) Schematic representation of our research question that aimed at studying the combined short- and longer-term relationship of a 'capture-handling-restraint protocol' for breeding great tits. (**B**) Upon capture, we measured birds and extracted a blood sample to quantify the levels of baseline (not shown in the figure) and stress-induced corticosterone levels (green clock). Upon release (blue clock), we recorded provisioning latency (orange clock) and following nest visits (grey clock) using RFID transponders. (**C**) Great tit ringed with a numbered metal ring and an RFID transponder (blue ring). (**D**) Nest box with an RFID logger at the entrance hole.

2. Materials and Methods

2.1 Study site

We collected data for both first and second clutches during three consecutive breeding seasons (May-July of 2020-2022) in a nest-box population of wild great tits, located in the Ettenhofer

Holz (48°03'26.7"N 11°15'11.0"E) in Upper Bavaria, Germany. Our study area contained 124

nest boxes and is a mosaic deciduous-coniferous forest of predominantly European beech

(*Fagus sylvatica*) and Norway spruce (*Picea abies*), typical of managed forests in this area.

This population has been the subject of a long-term study since 2013 and many individuals are

fitted with a uniquely numbered metal leg ring (Hau et al., 2022).

2.2 Field Procedures

At the beginning of the breeding season, we visited the nest boxes every second day to

determine when the first egg was laid, the final clutch size and when the first chick hatched

(hatching date = nestling day 0, for more details regarding the study site and the standard nest

box monitoring protocol (see Hau et al., 2022; Mentesana et al., 2024). When nestlings were

between seven and twelve days old (mean ± SD = 8.56 ± 0.88 days), we caught the parents in

the nest box between 07:00 and 14:00 hours. We took a blood sample (50-75 µl) from the

brachial vein using a cannula needle (gauge size 26) and heparinised microhematocrit capillary

for determining baseline corticosterone levels within three minutes after capture (mean ± SD =

118 ± 34 sec). We recorded the ID of the individual from the metal ring and RFID transponders,

if present. Unmarked birds were fitted with metal rings and with a silicone leg ring containing

a uniquely identifiable PIT tag (PIT Bird tag, EM4102 based EccelTechnology). Phenotypic

sex was determined based on plumage characteristics and age (1st year breeding vs. older) was

estimated based on the colouration of the primary coverts. The age of birds that were ringed as

nestlings was based on the year of birth. We proceeded with the 'capture-handling-restraint

protocol' by placing the bird inside an opaque cloth bag for approximately 20 minutes

(Wingfield et al., 1992, 1982). Then, the bird was taken out of the bag, morphometric

measurements were taken (body mass to the nearest 0.25 g, tarsus and wing length to the nearest

0.1 mm) and the bird was checked for the presence of parasites (i.e., ticks, louse flies, feather

mites). Next, we took a second blood sample (max. 25-30 µl) to determine stress-induced

corticosterone levels and recorded the sampling time (i.e., the time between capture and

finishing the second blood sample; mean ± SD = 28.70 ± 3.74 min, min. = 21.00 min, max. =

40.67 min). After the bleeding had stopped, we released the bird into its nest box through the

entrance hole and again recorded the time to determine the total handling time (i.e., the time

between capture and release; mean ± SD = 32.19 ± 5.07 min, min. = 8.00 min, max. = 45.00

min). Both blood samples were kept on ice until centrifugation in our laboratory. The plasma

fraction was stored at -20°C until analysis (as described in Hau et al., 2022). Ambient

temperature at capture was recorded automatically to the nearest 0.01°C every 30 min by a

HOBO weather station (equipped with S-THB-M002 12 Bit Temperature sensors) located within the study area. We used the temperature recordings that were closest to our time of capture.

As proxies for nestling condition, we measured fledgling mass and tarsus length of nestlings shortly before fledging (97.7% on nestling day 15 and 2.3% on day 16). We checked nests daily for fledging and recorded if any nestlings had died in the nest since day 15 to determine the number of fledglings.

2.3 Corticosterone analysis

We quantified corticosterone levels as described in detail in Hau et al. (2022) using the Arbor Assays DetectX® Corticosterone Enzyme Immunoassay Kit with the high sensitivity protocol (Cat. No. K014). Before quantification, we conducted a double liquid-liquid extraction with diethyl ether of the plasma samples (Baugh et al., 2012; Hau et al., 2022; Ouyang et al., 2011). After evaporation of the diethyl ether, samples were reconstituted over night at 4°C in the assay buffer provided with the kit and assayed in duplicate on the following day. We included several control samples in each assay batch, which were taken through the entire extraction and analysis procedure. Two positive controls were included, containing stripped chicken plasma to which a known amount of a corticosterone stock solution was added, as well as two negative controls containing only ultrapure water. In total, we extracted and analysed 26 batches and plates (for further details on assay procedures, as well as intra- and inter-assay variation, see Supplementary Information 1). We measured two quality control (QC) samples on each plate, one at around 60% (QC 1) and one at around 30% (QC 2) percent binding. We measured both QC samples twice in duplicate on each plate and used these values to calculate intra- and inter-assay variation to evaluate extraction and assay consistency. Intra-assay variation was calculated by averaging the coefficient of variation between duplicate measurements of the control samples across all plates. Intra-assay variation was $13 \pm 10\%$ for QC 1, and $8.1 \pm 4.5\%$ for QC 2. Inter-assay variation was calculated by calculating the coefficient of variation across the corticosterone concentrations of the two control samples across plates. Inter-assay variation was 15.9% for QC1, and 9.1% for QC. Note that these values include not just variation on assay performance across plates, but also variation in extraction efficiency.

For nine individuals (three females, nine males), we measured unusually high corticosterone levels (range: 134.50-313.62 ng/ml) as compared to our long-term data from this population (Hau et al., 2022). Since these samples were processed together on the same plate,

the values are likely not true biological measurements but rather the result of human error and
were excluded from the analyses.

2.4 Automatic nest visit recordings
We used an RFID system (Radio Frequency IDentification) to automatically record the date and
time of each bird's visit to a nest box. Each nest box was equipped with a reading antenna,
positioned around the entrance hole.

The reliability and consistency of RFID devices can vary significantly between studies
(Hughes et al., 2021; Iserbyt et al., 2018; Schlicht and Kempenaers, 2015), making it essential
to perform validations for each study. Therefore, we performed validation tests to assess the
reliability of our setup. In 2020, we validated the RFID system by comparing its nest visit data
with video recordings prior to the capture (for details see Supplementary Information 2). This
allowed us to determine how accurately entries and exits were recorded and to estimate the time
spent inside and outside the nest box, which is important for identifying separate nest visits.
Entry registration was reliable, with 94.3% of entries correctly registered. Exit registration was
much lower, with only 22.4% of exits properly registered because birds exited the nest boxes
too quickly for the system to respond and record. Ninety per cent of the nest visits lasted less
than one minute and 52 seconds. Based on these results, we excluded any consecutive RFID
recordings by the same bird that occurred within a two-minute interval, as they likely belonged
to the same visit (see Supplementary Information 2).

We conducted a second video validation in 2022 to test the performance of the RFID
system following the release of birds into the nest box after capture. When we placed birds at
the entrance hole, the entry was properly recorded in only 53.6% of cases. The registration of
exits was 42.9%, likely due to birds exiting more slowly and cautiously. Birds also stayed much
longer inside the nest box after the release, up to 40 minutes. Therefore, to calculate
provisioning latency (i.e., the time between release and the bird's first return to resume
provisioning) and to avoid confusing the exit after release with the subsequent re-entry, we
filtered the data to include only the first registered movement that occurred at least 51 minutes
after release. This cutoff was based on the latest exit (after 40 minutes) and the earliest return
(after 62 minutes) observed in the video analysis in 2022. Thus, 51 minutes was a conservative
midpoint between these values. To filter the data, we created a timestamp variable using the
'as.POSIXct' function (R Core Team, 2024). Birds that abandoned their nests were excluded
from the final analyses. For the birds that returned on the next day, the duration of the night was
included in the calculation of provisioning latency.

One common error with these automatic recording systems is the registration of non-
parental individuals, either due to incorrect PIT tag readings or because birds other than the
social parents inspected the nest box. Non-parental IDs (e.g., IDs of other tagged birds, misread
IDs) were recorded in ~60% of the nests and were excluded from the analyses. Other errors
were time-shifted RFID recordings (in ~5% of the nests), due to failing batteries and automatic
resetting of the logger clock. To detect such instances, we visually inspected the data for entries
occurring outside the birds' usual activity times (normal activity time: ~04:30 to 21:30h),
compared them with notes on battery changes and discrepancies between the logger clock and
field-recorded notes and adjusted the data accordingly. If a time shift could not be resolved due
to missing information on the correct time and it occurred around the time of the capture of a
bird, the data were excluded from the analyses (this occurred for only one nest).

Since this study did not include a control group of uncaptured birds, we calculated the
average provisioning interval (i.e., regular intervals between the nest visits during the two days
before the capture) to better interpret provisioning latency. For this, we used data from birds
that had already been equipped with a PIT tag during a previous capture (N = 83, females: 43,
males: 40).

2.5 Statistical Analyses
General Information
All statistical analyses were performed with the statistical software R (version 4.4.0, R Core
Team, 2024) using the '*lm*', '*lmer*' or the '*glmer*' function of the R-package 'lme4' (Bates et al.,
2015). To ensure that model assumptions were met, we conducted a series of diagnostic checks
by visually inspecting the residuals. All covariates were mean-centred to allow for comparisons
of effect sizes across variables. We fitted statistical models using a Bayesian approach using the
'*sim*' function of the R-package '*arm*' (Gelman and Su, 2021), to obtain 10,000 simulations for
the posterior modes, from which we estimated the mean and the 95% Bayesian credible
intervals (CrI). Following recommendations by Amrhein et al. (2019), we present our results
by first describing their direction and then their statistical significance (defined as variables for
which the 95% CI did not overlap with zero). To plot the results, we used the '*ggplot*' function
of the library '*ggplot2*' (Wickham, 2016).

2.5.1 Short-term relationship of the 'capture-handling-restraint protocol' with
physiology and behaviour
To test for a relationship of our 'capture-handling-restraint protocol' with either physiology or
behaviour, we ran three univariate linear mixed-effect models with a Gaussian distribution. Bird

ID and nest ID were included as random factors. To test for the influence of our handling protocol on the birds' physiology, we fitted stress-induced corticosterone levels as the response variable. Sampling time, date and time of capture were included as covariates. Sex (levels: females vs. males) was added as a fixed factor, as well as its interaction with sampling time. As random factors, we additionally included year and assay number. To test whether corticosterone levels increased in response to the protocol, we compared baseline (< 3 min) to stress-induced corticosterone levels using a two-sided paired t-test.

To test for a relationship between handling and the birds' behaviour, we used log-transformed provisioning latency as the response variable. Handling time, date and time of capture were added as covariates. Sex was added as a fixed effect. The interaction between handling time and sex was also included. Additionally, bird ID, nest ID and year were included as random factors, but since year explained no variance, we removed it from the final model. For the interpretation of provisioning latency, we first compared the average provisioning interval one and two days before the capture. We performed two-sided paired t-tests to compare the average provisioning intervals between the two days preceding capture, analysing females and males separately to account for potential sex differences.

To test if our handling protocol led birds to abandon their nests (i.e., whether birds did not return to the nest after being captured), we ran a univariate generalised linear mixed-effect model with a binomial distribution fitting the return of a bird (levels: no vs. yes) as the response variable, handling time as a covariate, and sex as a fixed effect. We set the covariate handling time in interaction with sex. Nest ID and year were included as random factors.

To test for the influence of stress-induced corticosterone levels on the birds' behaviour, we fitted the log-transformed provisioning latency as the response variable. Stress-induced corticosterone levels, date and time of capture were added as covariates. Sex was included as a fixed factor together with its interaction with stress-induced corticosterone levels. Additionally, we included bird ID and nest ID as random factors. We also included year as a random factor, but since it explained no variance, we removed it from the final model.

2.5.2 *Relationship of ambient temperature, parental condition and brood quality with behaviour*

To test for the potential influence of other factors besides handling on provisioning latency, we performed five additional models with a Gaussian distribution and log-transformed provisioning latency as the response variable. We first investigated the influence of ambient temperature at capture by fitting this variable, date and time of capture as covariates, sex as a

fixed factor and the interaction between temperature and sex. Bird ID and nest ID were included as random factors. Elevated temperatures may facilitate quicker energy replenishment for parents following a predation event due to increased foraging profitability (Avery and Krebs, 1984). According to this, higher temperatures could be expected to be associated with a lower provisioning latency.

Second, we examined whether the condition of the parents at capture influenced the time it took them to return to their nest box. For this, we ran two separate models for females and males, respectively. We included mass and tarsus length as covariates; age (levels: one year vs. older) and parasite infection (levels: no vs. yes) as fixed factors. For females, we included bird ID and year as random factors. However, for males, these variables did not explain any variance; thus, we ran a linear model. Parents in better condition might cope with stress more effectively than those in poorer condition. Conversely, individuals in poor condition might resume provisioning sooner, as their reduced prospects for future reproduction could favour greater investment in the current brood. Parental age might also influence provisioning latency. Older birds might have greater stress tolerance and respond less sensitively to capture, resulting in shorter provisioning delays (Monaghan and Spencer, 2014). Limited future mating opportunities might also drive them to suppress the behavioural short-term stress response of avoiding the nest after being captured at the nest and to return sooner to feed their offspring (Kitaysky et al., 1999).

Third, we tested for the association that brood quality (i.e., clutch size, offspring age and offspring body condition) might have with the parents' behaviour. We ran two separate models for females and males. In each model, we added offspring age (median = 8 days, range: 7-12 days) and number of chicks (median = 6, range: 2-8) as continuous variables. We added clutch number (levels: 1 vs. 2) as a fixed factor. In both models, we included bird ID as a random factor. Solely for females, we also included year (because for males, it did not explain any variance). Brood quality may encourage parents to return to their nests more quickly, as offspring that are more numerous, older and in better body condition are likely to beg more loudly for food than offspring that are fewer, smaller and weaker (Bowers et al., 2019).

2.5.4 Longer-term relationship of the 'capture-handling-restraint protocol' with proxies for reproductive success

To examine whether the protocol led to prolonged behavioural changes following capture, we analysed average provisioning intervals in birds that returned to their nests, separately for females and males. We first compared the average provisioning interval one day before and one

day after the capture using two-sided paired t-tests to determine whether the birds returned to regular feeding behaviour after the initial capture. To determine if the average provisioning interval remained constant or if missed feeding time due to post-capture nest absence was later compensated for, we compared the average provisioning interval one and two days after the capture, again using a two-sided paired t-test for females and males separately.

We studied the longer-term implications of the protocol on fitness using first fledgling number and then fledgling mass and tarsus length as proxies. We first ran two generalised linear mixed-effect models with Poisson distribution fitting fledgling number (median = 5, range: 0-12) as the response variable and stress-induced corticosterone levels or provisioning latency as covariates in separate models, together with hatching date and clutch size. Nest ID and year were included as random factors. Next, we conducted four linear mixed-effect models with a Gaussian distribution fitting fledgling mass and tarsus length as response variables, respectively, and stress-induced corticosterone levels or provisioning latency as covariates. In each of these four models, we also included hatching date and clutch size as covariates. Since body mass can change throughout the day, we also added time of the day as a covariate for the two models looking at this variable. Sex and its interaction with either stress-induced corticosterone levels or provisioning latency were also included in the models. The random effects included in these models were parental bird ID, person ID (i.e., the person who took morphological data) and year.

Lastly, we examined whether the abandonment of a parent would lead to nest failure or reduce fledging success. For this, we fitted nest success (levels: no vs. yes) and fledgling number as response variables of two separate generalised linear mixed-effect models with Poisson distribution and a binomial distribution, respectively. For both models, we set the fixed factor return (levels: no vs. yes) in interaction with sex and added nest ID and year as random factors. In the model testing for fledgling number, we additionally included the hatch date of the offspring as a covariate to control for within-season differences.

Ethical Note

All field procedures were conducted in compliance with the legal requirements of Germany and the EU and were approved by the cognizant regional governmental authority, the Regierungspräsidium von Oberbayern, Germany (license no. ROB-55.2-2532-Vet_02-13-204 & ROB-55.2-2532-Vet_02-15-25).

3. Results

Over the course of three breeding seasons, we captured and collected morphological data for 185 individual adult great tits. Of these 185 adults, 86 were captured repeatedly; 48 were recorded twice, 32 were recorded three times, five were recorded four times and one bird was even recorded five times. We also collected morphological data for 722 fledglings from 136 nests (1st clutch: 83%, 2nd clutch: 17%; N nests: 2020: 406, 2021: 185, 2022: 331).

We collected blood samples to quantify stress-induced corticosterone levels from 183 adult individuals, resulting in 299 samples. Nine samples had to be excluded due to their exceptionally high values (see 2.3 Corticosterone analysis), leaving us with 290 samples (2020: 81, 2021: 100, 2022: 109) from 181 individuals with 160 nests (1st clutch: 83%, 2nd clutch: 17%). Out of the 181 individuals, 48 were sampled twice, 26 were sampled three times and three were sampled four times. Average stress-induced sampling time was 28.70 ± 3.74 minutes (range: 21.00 - 40.67 min).

We collected behavioural data for 169 individuals. Out of these 169 individuals, 39 were recorded for two different breeding attempts, 33 were recorded for three attempts and eight were recorded for four attempts, resulting in 298 recordings (N females: 2020: 42, 2021: 55, 2022: 57; N males: 2020: 37, 2021: 55, 2022: 52) from 165 nests (1st clutch: 82%, 2nd clutch: 18%; N nests: 2020: 45, 2021: 61, 2022: 59). For 82 out of the 169 individuals, we had average provisioning interval data the two days before and after the capture. Out of these 82 individuals, 23 were recorded twice (N females: 12, N males: 11) and four individuals were recorded three times (N females: 2, N males: 2). This resulted in 113 paired recordings (N females: 59, N males: 54) of the average provisioning interval the two days before and the two days after the capture.

3.1 Short-term relationship of the 'capture-handling-restraint protocol' with physiology and behaviour

Baseline corticosterone levels (mean $\pm$ SD = 4.196 ± 2.633 ng/ml, range: 0.061-17.691 ng/ml) were lower compared to stress-induced corticosterone levels (mean $\pm$ SD = 28.687 ± 10.672 ng/ml, range: 3.950-66.479 ng/ml) upon capture ($t(275) = -38.847$, p-value < 0.000). These results hold after looking at each sex separately (females: $t(138) = -27.125$, p-value < 0.000; males: $t(136) = -28.002$, p-value < 0.000).

Out of 240 capture events, 217 captured birds returned to their nests (209 on the same day and eight on the following day) and 23 birds abandoned the nest (11 female individuals and 11 male individuals, with one male which abandoned twice, both times within one year). Parents

resumed feeding on average after 2.53 ± 3.20 hours. Females and males had similar provisioning latencies of 2.47 ± 2.91 and 2.59 ± 3.52 hours, respectively. The provisioning latency differed significantly from the average provisioning interval the day before the capture for both sexes (females: t(58) = -6.158, p-value < 0.000; males: t(53) = -5.168, p-value < 0.000), which was 8.72 ± 3.57 minutes for females and 6.87 ± 1.97 minutes for males (females: min. = 4.20 min, max. = 24.45 min; males: min. = 4.08 min, max. = 14.23 min; for more details see Supplementary Information 5).

While sampling time explained differences in physiology observed among male great tits, this was not the case for females. Handling time did not explain differences in observed behaviour for either sex. Sampling time tended to be negatively related to stress-induced corticosterone levels (Table 1A; Figure 2A) and handling time appeared to be negatively related to both abandonment rate (Table 2A) and provisioning latency (Table 1B; Figure 2B). However, of these relationships, only the relationship of sampling time and stress-induced corticosterone levels of male birds was significant (posterior probabilities for females and males: stress-induced corticosterone levels: > 74% and > 99%; abandonment: > 62% and > 52%; provisioning latency: > 72% and > 27%). Stress-induced corticosterone levels were positively correlated with provisioning latency, although this correlation too was not statistically significant (posterior probabilities: females > 13% and males > 56%; Figure 2; Table 1C).

Table 1. Results from mixed models testing the short-term relationships of sampling and handling protocol with physiology and behaviour. Statistical significance was considered when the CI did not overlap zero.

A) Relationship between sampling time and stress-induced corticosterone levels		
Fixed effects		
	β	**95% CI**
Intercept	29.984	[25.396, 34.660]
Sampling time[a] (scaled)	-0.617	[-2.843, 1.232]
Sex [male]	-1.882	[-4.408, 0.658]
Date of capture (scaled)	-0.004	[-1.212, 1.193]
Time of day at capture (scaled)	-2.054	[-3.343, -0.774]
Sampling time (scaled) : Sex [male]	-1.117	[-3.425, 1.183]
Random effects		
	σ^2	95% CI
Bird ID (N = 179)	34.276	[27.228, 43.011]
Nest ID (N = 159)	11.167	[8.611, 14.361]
Assay batch ID[b] (N = 26)	3.094	[1.724, 5.198]
Year (N = 3)	13.793	[2.049, 55.818]
Residual	58.957	[50.213, 69.842]
B) Relationship between handling time and provisioning latency[c]		
Fixed effects		
	β	**95% CI**
Intercept	0.747	[0.603, 0.885]
Handling time[d] (scaled)	-0.033	[-0.143, 0.079]
Sex [male]	0.036	[-0.120, 0.194]
Date of capture (scaled)	0.023	[-0.092, 0.135]
Time of day at capture (scaled)	-0.111	[-0.223, -0.001]
Handling time (scaled) : Sex [male]	0.057	[-0.124, 0.234]
Random effects		
	σ^2	95% CI
Bird ID (N = 136)	0.040	[0.030, 0.054]
Nest ID (N = 128)	0.378	[0.296, 0.483]
Residual	0.197	[0.163, 0.240]
C) Relationship between stress-induced corticosterone levels and provisioning latency		
Fixed effects		
	β	**95% CI**
Intercept	0.731	[0.584, 0.875]
Stress-induced levels[e] (scaled)	0.074	[-0.054, 0.200]
Sex [male]	0.061	[-0.090, 0.212]
Date of capture (scaled)	0.029	[-0.094, 0.152]
Time of day at capture (scaled)	-0.114	[-0.228, 0.001]
Stress-induced levels (scaled) : Sex [male]	-0.151	[-0.327, 0.019]
Random effects		
	σ^2	95% CI
Bird ID (N = 130)	0.020	[0.015, 0.028]
Nest ID (N = 122)	0.411	[0.340, 0.567]
Residual	0.208	[0.171, 0.255]

[a] Time span between capture and sampling of the stress-induced corticosterone levels
[b] Batch identity of samples assayed together for their stress-induced corticosterone levels
[c] Time it took a bird to resume provisioning activity after being captured and handled

[d] Time span between capture and release of the bird
[e] Corticosterone levels around 30 minutes after the capture

Figure 2. Relationship of sampling and handling time with physiology and behaviour for
breeding female (circles) and male (triangles) wild great tits. Association of **A**) sampling time
with stress-induced corticosterone levels, **B**) handling time with the provisioning latency (i.e.,
time between release and first return to the nest box) and **C**) stress-induced corticosterone levels
with provisioning latency. Figures show mean estimates (coloured lines), 95% credible intervals
(shaded grey) and raw data (grey symbols). The asterisk (*) indicates statistical significance for
the relationship between sampling time and stress-induced corticosterone levels in males,
meaning males that experienced longer sampling times had lower stress-induced corticosterone
levels.

Table 2. Results from mixed models testing the relationship between abandonment and
indicators of fitness. Statistical significance was considered when the CI did not overlap zero.

A) **Relationship between handling time[a] and abandonment**		
Fixed effects		
	β	95% CI
Intercept	28.648	[-67.162, 122.636]
Handling time	-27.543	[-215.820, 136.776]
Sex [male]	-27.434	[-149.576, 91.168]
Handling time: Sex [male]	50.091	[-149.559, 255.413]
Random effects		
	σ^2	95% CI
Nest ID (N = 142)	1247.662	[967.369, 1577.736]
Year (N = 3)	0.011	[0.000, 0.057]
B) **Relationship between abandonment and nest success[b]**		
Fixed effects		
	β	95% CI
Intercept	10.586	[-31.104, 53.190]
Return [yes]	4.195	[-43.832, 51.905]
Sex [male]	0.976	[-47.817, 48.473]
Return [yes] : Sex [male]	-0.971	[-51.857, 50.658]
Random effects		
	σ^2	95% CI
Nest ID (N = 142)	2311.270	[1766.692, 2929.550]
Year (N = 3)	25.025	[0.909, 128.759]
C) **Relationship between abandonment and fledgling number**		
Fixed effects		
	β	95% CI
Intercept	2.128	[0.611, 3.640]
Return [yes]	0.706	[0.234, 1.182]
Sex [male]	0.306	[-0.240, 0.859]
Hatch date (numeric)	-0.261	[-0.534, 0.013]
Return [yes] : Sex [male]	-0.305	[-0.867, 0.263]
Random effects		
	σ^2	95% CI
Nest ID (N = 142)	0.140	[0.112, 0.171]
Year (N = 3)	0.079	[0.007, 0.247]

495 [a] Time span between capture and release of the bird
496 [b] Binomial variable; 'nest success' (any number of fledged offspring) or 'nest failure' (no fledged offspring)
497

      3.2 Relationship of ambient temperature, parental condition and brood quality with

499   *behaviour*

Ambient temperature at capture (Table 3A), parental condition (Tables 3B & 3C) and brood

quality (Tables 3D & 3E) did not explain variation in provisioning latency. Great tits captured

and handled during warmer times of the day seemed to return to their nests later than those

captured during cooler periods. However, these relationships were not statistically significant

(posterior probabilities for females and males: > 18% and > 42%). While females appeared to

take longer to return to their nests when caring for a second clutch or for more chicks, males

returned more quickly. Both sexes seemed to return faster when nestlings were older. However,

once again, none of these relationships were statistically significant (posterior probabilities for

females and males: clutch number: > 4% and > 91%; number of nestlings: > 40% and > 93%;

nestling age: > 59% and > 67%).

Table 3. Results from mixed models testing the relationship between non-handling related
variables with provisioning latency. Statistical significance was considered when the CI did not
overlap zero.

A) Relationship between temperature at capture and provisioning latency

Fixed effects

	β	95% CI
Intercept	0.755	[0.613, 0.899]
Temperature at capture (°C)	0.067	[-0.076, 0.211]
Sex [male]	0.041	[-0.115, 0.202]
Capture date (scaled)	0.007	[-0.128, 0.139]
Capture time (scaled)	-0.125	[-0.241, -0.005]
Temperature at capture (scaled) : Sex [male]	-0.084	[-0.228, 0.057]

Random effects

	σ^2	95% CI
Bird ID (N = 135)	0.051	[0.038, 0.069]
Nest ID (N = 126)	0.377	[0.296, 0.481]
Residual	0.188	[0.155, 0.230]

B) Relationship between parental condition and provisioning latency (females)

Fixed effects

	β	95% CI
Intercept	-0.247	[-5.439, 4.885]
Mass (g)	-0.114	[-0.281, 0.052]
Tarsus length (mm)	0.154	[-0.100, 0.412]
Age [adult]	-0.091	[-0.361, 0.168]
Parasites [yes]	0.103	[0.145, -0.346]

Random effects

	σ^2	95% CI
Bird ID (N = 75)	0.273	[0.206, 0.358]
Year (N = 3)	0.004	[0.000, 0.021]
Residual	0.227	[0.176, 0.300]

C) Relationship between parental condition and provisioning latency (males)

Fixed effects

	β	95% CI
Intercept	0.257	[-3.828, 4.372]
Mass (g)	-0.094	[-0.300, 0.110]
Tarsus length (mm)	0.121	[-0.138, 0.375]
Age [adult]	-0.348	[-0.725, 0.025]
Parasites [yes]	0.207	[-0.113, 0.532]

D) Relationship between brood quality and provisioning latency (females)

Fixed effects

	β	95% CI
Intercept	0.780	[-0.557, 2.129]
Clutch number [2]	0.275	[-0.028, 0.578]
Age nestlings (days)	-0.015	[-0.148, 0.118]
Clutch size	0.011	[-0.078, 0.101]

Random effects

	σ^2	95% CI
Parental bird ID (N = 74)	0.279	[0.210, 0.368]
Year (N = 3)	0.001	[0.000, 0.006]
Residual	0.251	[0.195, 0.332]

E) Relationship between brood quality and provisioning latency (males)

Fixed effects

	β	95% CI
Intercept	1.730	[0.018, 3.414]
Clutch number [2]	-0.315	[-0.780, 0.147]
Age nestlings (days)	-0.037	[-0.204, 0.132]
Clutch size	-0.096	[-0.225, 0.031]

Random effects		
	σ^2	95% CI
Parental bird ID (N = 61)	0.073	[0.047, 0.111]
Residual	0.561	[0.429, 0.754]

3.3 Longer-term relationship of the 'capture-handling-restraint protocol' with proxies for reproductive success

Parents that returned to their nests after capture and handling resumed provisioning with intervals on the following day comparable to the day preceding capture (females and males: N = 59, 54; average provisioning interval on the following day after capture = 8.09 ± 5.43 min, 6.91 ± 2.47 min; p-values = 0.229, 0.856; Supplementary Table 1). Stress-induced corticosterone levels appeared positively associated with fledgling number and negatively associated with fledgling mass and tarsus length. But these relationships were not statistically significant (posterior probabilities for females and males: fledgling number: > 44% and > 54%; mass: > 89% and > 12%; tarsus length only for males: > 23%). However, females with lower stress-induced corticosterone levels raised nestlings with significantly longer tarsi (posterior probability > 99%; Figure 3; Table 4C). Provisioning latency seemed positively related to all three fitness proxies, but none of these relationships were statistically significant (posterior probabilities for females and males: fledgling number: > 26% and > 50% (Table 5A); fledgling mass: > 23% and > 57% (Table 5B); fledgling tarsus length: > 41% and 47% (Table 5C); Figure 3). The return of a parent did not explain whether a nest was successful or not (posterior probabilities for both sexes > 43%; Table 2B). However, individuals that returned to the nest raised more fledglings than individuals that did not (posterior probabilities for both sexes > 0.1%; Table 2C).

Table 4. Results from mixed models testing the relationship between the physiology of breeding great tits with indicators of fitness. Statistical significance was considered when the CI did not overlap zero.

A) Relationship between stress-induced corticosterone levels[a] and fledgling number		
Fixed effects		
	β	**95% CI**
Intercept	1.310	[0.998, 1.624]
Stress-induced levels (scaled)	0.007	[-0.089, 0.106]
Sex [male]	-0.006	[-0.122, 0.112]
Hatch date (scaled)	-0.076	[-0.188, 0.034]
Clutch size	0.075	[-0.035, 0.183]
Stress-induced levels (scaled) : Sex [male]	-0.010	[-0.144, 0.126]
Random effects		
	σ^2	95% CI
Nest ID (N = 157)	0.309	[0.257, 0.366]
Year (N = 3)	0.092	[0.012, 0.261]
B) Relationship between stress-induced corticosterone levels and fledgling mass		
Fixed effects		
	β	**95% CI**
Intercept	16.875	[15.388, 18.335]
Stress-induced levels (scaled)	-0.127	[-0.327, 0.079]
Sex [male]	0.127	[-0.281, 0.540]
Hatch date (scaled)	0.102	[-0.022, 0.223]
Clutch size	-0.164	[-0.265, -0.058]
Capture time of the chicks (scaled)	0.138	[-0.009, 0.285]
Stress-induced levels (scaled) : Sex [male]	0.289	[-0.002, 0.580]
Random effects		
	σ^2	95% CI
Parental bird ID (N = 161)	1.405	[1.212, 1.621]
Observer ID (N = 5)	0.240	[0.080, 0.498]
Year (N = 3)	0.984	[0.616, 1.453]
Residual	2.007	[1.855, 2.173]
C) Relationship between stress-induced corticosterone levels and fledgling tarsus length		
Fixed effects		
	β	**95% CI**
Intercept	18.846	[18.326, 19.361]
Stress-induced levels (scaled)	-0.103	[-0.187, -0.018]
Sex [male]	0.066	[-0.084, 0.217]
Hatch date (scaled)	-0.023	[-0.079, 0.035]
Clutch size	0.046	[0.001, 0.091]
Stress-induced levels (scaled) : Sex [male]	0.107	[-0.022, 0.235]
Random effects		
	σ^2	95% CI
Parental bird ID (N = 161)	0.143	[0.118, 0.171]
Observer ID (N = 5)	0.066	[0.023, 0.140]
Year (N = 3)	0.067	[0.029, 0.124]
Residual	0.564	[0.522, 0.610]

[a] Corticosterone levels around 30 minutes after capture

Table 5. Results from mixed models testing the relationship between behaviour of breeding great tits and indicators of their fitness. Statistical significance was considered when the CI did not overlap zero.

A) Relationship between provisioning latency[a] and fledgling number		
Fixed effects		
	β	**95% CI**
Intercept	1.491	[1.271, 1.704]
Provisioning latency (scaled)	0.036	[-0.073, 0.144]
Sex [male]	0.008	[-0.117, 0.135]
Hatch date (scaled)	-0.073	[-0.161, 0.014]
Clutch size	0.097	[0.013, 0.179]
Provisioning latency (scaled) : Sex [male]	-0.045	[-0.179, 0.089]
Random effects		
	σ^2	95% CI
Nest ID (N = 127)	0.082	[0.064, 0.102]
Year (N = 3)	0.043	[0.003, 0.149]
B) Relationship between provisioning latency and fledgling mass		
Fixed effects		
	β	**95% CI**
Intercept	15.785	[14.609, 16.948]
Provisioning latency (scaled)	0.101	[-0.168, 0.365]
Sex [male]	0.032	[-0.402, 0.472]
Hatch date (scaled)	0.054	[-0.077, 0.185]
Clutch size	-0.188	[-0.333, -0.045]
Capture time of the chicks (scaled)	-0.075	[-0.233, 0.082]
Provisioning latency (scaled) : Sex [male]	-0.175	[-0.495, 0.139]
Random effects		
	σ^2	95% CI
Parental bird ID (N = 131)	1.302	[1.100, 1.536]
Observer ID (N = 5)	0.810	[0.258, 1.781]
Year (N = 3)	0.499	[0.105, 1.461]
Residual	2.125	[1.951 2.323]
C) Relationship between provisioning latency and fledgling tarsus length		
Fixed effects		
	β	**95% CI**
Intercept	19.235	[18.796, 19.671]
Provisioning latency (scaled)	0.013	[-0.098, 0.124]
Sex [male]	0.037	[-0.136, 0.216]
Hatch date (scaled)	-0.030	[-0.093, 0.033]
Clutch size	0.086	[0.018, 0.153]
Provisioning latency (scaled) : Sex [male]	-0.044	[-0.180, 0.095]
Random effects		
	σ^2	95% CI
Parental bird ID (N = 131)	0.167	[0.136, 0.201]
Observer ID (N = 5)	0.156	[0.041, 0.362]
Year (N = 3)	0.040	[0.004, 0.127]
Residual	0.606	[0.556, 0.662]

[a] Time it took a bird to resume provisioning activity after being captured and handled

Figure 3. Relationship of physiology and behaviour with fitness indicators for breeding female (circles) and male (triangles) wild great tits. Association of stress-induced corticosterone levels with **A)** number of fledglings, **B)** mass of fledglings and **C)** tarsus length of fledglings. Association of provisioning latency (i.e., time between release and first return to the nest box) with **D)** number of fledglings, **E)** mass of fledglings and **F)** tarsus length of fledglings. Figures show mean estimates (coloured lines: yellow: stress-induced corticosterone levels, orange: provisioning latency), 95% credible intervals (shaded grey) and raw data (grey symbols).

4. Discussion

We examined the short- and longer-term effects of a standardised 'capture-handling-restraint protocol' by analysing its relationship behavioural and physiological responses of adult great tits during the nestling provisioning phase and the relationship of these responses with reproductive success. Overall, in 90.4% of capture events, birds returned to their nests. The protocol strongly affected adult physiological and behavioural traits on the short term, but had no longer-term consequences on behaviour. Moreover, except for abandonment, these short-term physiological and behavioural changes did not explain variation in fledgling condition and number, likely because, once parents returned to their nests, they resumed normal average provisioning intervals. While abandonment reduced the number of fledglings, it did not affect overall nest success or failure.

On a short-time scale, we found that fewer than 10% of capturing events (23 out of 240) resulted in parents abandoning their nests after being captured and this was not related to the duration of handling. Nonetheless, it is worth noting that the abandonment by one parent did not automatically lead to a complete nest failure, likely due to continued provisioning by the other parent. This suggests that, if a parent abandons the nest when the nestlings are between seven and twelve days old, the other parent may be capable of caring for the nestlings until fledging, albeit less efficiently, as indicated by the decreased number of fledglings in nests where one parent abandoned We also found significant increases in corticosterone levels following capture, confirming the expected physiological activation of the HPA axis. However, the lack of a relationship between sampling time and stress-induced corticosterone levels in females and the negative correlation for males did not align with our expectations. The females' result might be explained by the fact that most bird species reach their peak corticosterone levels in response to the initial encounter of an unpredictable stimulus at around 30 min (Baugh et al., 2012; Monaghan and Spencer, 2014; Romero et al., 2000; Wilson and McMahon, 2006;

Wingfield et al., 1998), which is approximately when we sampled most great tits (i.e., 28.70

min). Consistently, Huber et al. (2021) found a difference in stress-induced concentrations

between 15 and 30 min, irrespective of the three different handling protocols used. One possible

explanation for our findings in males is that there are sex differences in HPA axis regulation

during the breeding season, with the negative feedback being induced earlier in males than in

females. However, to our knowledge, there are no studies in birds supporting this hypothesis.

We only found results indicating the opposite: breeding males being insensitive to negative

glucocorticoid feedback (Astheimer et al., 1994). Furthermore, free-living animals' stress

responses are context-dependent and modulated by various other factors (e.g., season and

weather, development, previous experiences or remaining reproductive opportunities), making

it difficult to connect the consequences to a single immediate stressor (reviewed by Bonier et

al., 2009; Crespi et al., 2013; Kitaysky et al., 1999; Romero et al., 2000; Wingfield et al., 1998).

For example, Lynn et al. (2010) studied whether one capture experience can influence the stress

response to the same protocol later on. They subjected free-living eastern female bluebirds to a

capture-handling-restraint protocol, distinguishing between three test groups: birds that were

captured twice (once during their first and once during their second brood, i.e., repeaters) or

birds that were subjected to the protocol only once (either during their first or second brood).

They found a decline in corticosterone secretion in repeaters compared to birds experiencing

the protocol for the first time, indicating that they acclimated or habituated to the protocol after

only one experience. This suggests that a singular previous stressful experience can

significantly influence HPA responsiveness.

Likewise, to the significant increases in corticosterone levels following capture,

provisioning latency was also significantly delayed, suggesting a behavioural stress response.

But its magnitude was not related to the duration of handling. Our result aligns with the results

of a study on blue tits, where researchers retained birds on average for 30 minutes (range 7-84

min; Schlicht & Kempenaers, 2015). It is possible that the birds displayed a behavioural

response immediately after capture and/or that the response was related to other factors not

measured in our study.

Stress-induced corticosterone levels also did not explain the duration birds stayed away

from their nests. This result is in line with Lane et al. (2025), who reported that neither baseline

nor stress-induced corticosterone concentrations were correlated with return latency in female

song sparrows, arguing that corticosterone levels are not the only mechanism that mediates

behavioural stress responses. To test if an increase in stress-induced corticosterone levels affects

provisioning latency following capture and handling, one could conduct a follow-up study and

experimentally manipulate corticosterone's actions (see review by Hau et al., 2016). In general, experimental studies have linked glucocorticoids to parental provisioning in birds. For instance, administration of corticosterone implants to pied flycatcher parents reduced provisioning intervals and, at high concentrations, induced brood abandonment (Silverin, 1986). Likewise, corticosterone implants in male, white-crowned sparrows (*Zonotrichia albicollis*) from the high-provisioning (tan-striped) morph reduced offspring feeding rates, while the glucocorticoid receptor antagonist RU486 induced an increase in provisioning in the low-provisioning, white-striped morph (Almasi et al., 2008; Horton and Holberton, 2009). Also, daily pulses of high corticosterone exposure in incubating female tree swallows (*Tachycineta bicolor*) reduced later provisioning rates, especially when caring for large broods (Vitousek et al., 2018). Yet, these results concern provisioning rates and not provisioning latency after an acute stressor and therefore might represent different underlying mechanisms. Taken together, it is likely that the increase in corticosterone helped great tit parents in our study cope with the unexpected experience of being captured and supported changes in their provisioning behaviour (e.g., replenishing energy before returning to the nest). However, differences in provisioning latency are likely driven by factors other than stress-induced corticosterone levels (and also not by ambient temperature, parental condition or brood quality). For example, variation in the timing of behavioural resumption may be associated with the efficiency of HPA axis downregulation (Hau et al., 2016; Lattin and Kelly, 2020; Romero, 2004; Taff et al., 2018). Experimental work in eastern bluebirds has shown that more effective negative feedback mechanisms might predict an individual's resilience in response to transient stressors (Taff et al., 2018). A promising avenue for future research would therefore be to relate provisioning latency to glucocorticoid negative feedback. Likewise, to investigate physiological mechanisms mediating such behaviour, one could focus on both corticosterone and prolactin, the latter known as the 'parental hormone', since these hormones influence parental behaviour and may affect different components of the stress response (reviewed by Angelier & Chastel, 2009; Chastel et al., 2005; Ouyang et al., 2013). Additionally, one could explore the social dynamics within breeding pairs, as in great tits, the behaviour of one parent might influence the other (Székely et al., 1996), as well as genetic differences in the behavioural stress response (Mutzel et al., 2013; Van Oers et al., 2004).

On a longer-time scale, great tit parents that returned to their nests resumed normal provisioning behaviour. This result is in line with Schlicht & Kempenaers (2015), who found similar time spans between nest blue tit visits preceding capture and after the initial return. Contrary to our predictions, neither stress-induced corticosterone levels nor provisioning

latency explained variation in fitness proxies, except for fledglings of parents with higher corticosterone levels having shorter tarsus lengths. Given that tarsus length is a morphological trait largely determined by genetics (Husby et al., 2011; Noordwijk et al., 1988) and that birds are exposed to multiple challenging events during the breeding season that may trigger a stress response (Huber et al., 2021; Lack, 1968; Soulsbury et al., 2020), it seems unlikely that a single capture event caused nestlings to develop shorter tarsi. This raises the question: why does an interruption of several hours in a usually frequent parental care behaviour have no major impact on reproductive success? Was the missing food delivery subsequently compensated for? Curiously, we did not find a significant increase in provisioning that would support the idea of compensation after a parent's return to the nest. Despite clear physiological and behavioural responses to the 'capture-handling-restraint protocol', the absence of measurable consequences for offspring number or condition, aside from a weak relationship with tarsus length, suggests a large degree of resilience in this (and many other) species during the provisioning period. These findings, therefore, contribute to a growing body of literature that questions the longer-term impacts of acute, short-term stressors on reproductive success in wild bird populations (Lynn et al., 2010; Nicolaus et al., 2008; Ouyang et al., 2013; Schlicht and Kempenaers, 2015). However, this should not be interpreted as evidence that such disturbances are inconsequential. Our study naturally lacked a true control group of uncaptured individuals (because assessing corticosterone levels requires capture and handling), limiting our ability to detect subtle differences in longer-term relationships with reproductive output.

5. Conclusion

Capture, handling and sampling are integral to research on wild animals in fields of biology as diverse as ecology, evolution, physiology, genetics and conservation. Yet our understanding of the short- and long-term implications of these procedures has remained limited. In this study, we assessed immediate physiological and behavioural responses to a standardised 'capture-handling-restraint protocol', as well as longer-term behaviour and reproductive success over three consecutive breeding seasons. While the protocol triggered clear acute stress responses, our findings suggest that great tits are behaviourally resilient, with no detectable long-term implications on reproductive success. However, these conclusions must be interpreted with caution. The absence of an uncaptured control group limits our ability to fully assess potential subtle or cumulative relationships. It also remains possible that such brief disruptions may have longer-term consequences not captured by our metrics or that become evident only across

successive breeding attempts. To advance both scientific understanding and ethical standards, future research should prioritise the inclusion of true control groups, consider the role of social dynamics within breeding pairs and monitor reproductive success over multiple breeding seasons. Additionally, comparative studies would be valuable for determining species' sensitivity to capture-handling protocols. These studies should include species with divergent life-history strategies, varying reproductive investment per breeding bout and different survival rates. Such efforts are essential for refining best practices in field research, ensuring that the burden on the welfare of the wild animals studied does not outweigh the benefits of scientific data collection. This approach will enable researchers and regulatory agencies responsible for evaluating experimental studies on wild populations to conduct and assess scientific research responsibly and ethically.

Acknowledgements

We would like to dedicate this paper to the memory of Michaela (Ela) Hau, whose insight, dedication and passion for research greatly shaped this study. Her contributions and enthusiasm continue to inspire us.

We highly appreciate the help from Tanguy Deville, Carlotta Bonaldi, Michi Stiegler and Natalia Perez with field and laboratory work. Many thanks to the Erzdiozöse München und Freising, especially K. Meindl, B. Vollmar and M. Laußer for allowing us to work in their forest. The Hau-Goymann laboratory group provided input that greatly improved data collection and manuscript preparation.

Authors' contributions

Conceptualisation: all authors. Data curation: FF, LM and CD. Formal analysis: FF and LM. Funding acquisition: MH. Investigation: all authors. Methodology: all authors. Project administration: LM and MH. Resources: MH and CD. Software: FF, LM and CD. Supervision: LM and MH. Validation: FF, LM and CD. Visualisation: FF and LM. Writing - original draft: FF and LM. Writing - review and editing: all authors.

Conflict of interest declaration

The authors declare that no competing interests exist.

Data availability statement

All data and code are available at Zenodo:
https://zenodo.org/records/19094415?token=eyJhbGciOiJIUzUxMiJ9.eyJpZCI6ImMxNDlhY

Funding

This work was funded by the Max Planck Society.

Declaration of Generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used ChatGPT in order to improve readability and language. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

References

Almasi, B., Roulin, A., Jenni-Eiermann, S., Jenni, L., 2008. Parental investment and its sensitivity to corticosterone is linked to melanin-based coloration in barn owls. Horm. Behav. 54, 217–223. https://doi.org/10.1016/j.yhbeh.2008.02.021

Amrhein, V., Trafimow, D., Greenland, S., 2019. Inferential Statistics as Descriptive Statistics: There Is No Replication Crisis if We Don't Expect Replication. Am. Stat. 73, 262–270. https://doi.org/10.1080/00031305.2018.1543137

Andrews, C., 2022. On the use of body mass measures in severity assessment in laboratory passerine birds. Anim. Welf. 31, 387–401. https://doi.org/10.7120/09627286.31.1.013

Angelier, F., Chastel, O., 2009. Stress, prolactin and parental investment in birds: A review. Gen. Comp. Endocrinol. 163, 142–148. https://doi.org/10.1016/j.ygcen.2009.03.028

Angelier, F., Clément-Chastel, C., Welcker, J., Gabrielsen, G.W., Chastel, O., 2009. How does corticosterone affect parental behaviour and reproductive success? A study of prolactin in black-legged kittiwakes. Funct. Ecol. 23, 784–793. https://doi.org/10.1111/j.1365-2435.2009.01545.x

Astheimer, L.B., Buttemer, W.A., Wingfield, J.C., 1994. Gender and seasonal differences in the adrenocortical response to ACTH challenge in an Arctic passerine, Zonotrichia leucophrys gambelii. Gen. Comp. Endocrinol. 94, 33–43.

Avery, M.I., Krebs, J.R., 1984. Temperature and foraging success of great tits *Parus major* hunting for spiders. Ibis 126, 33–38.

Bates, D., Maechler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models Using lme4. J. Stat. Softw. 67, 1–48. https://doi.org/doi:10.18637/jss.v067.i01

Baugh, A.T., Schaper, S.V., Hau, M., Cockrem, J.F., de Goede, P., van Oers, K., 2012. Corticosterone responses differ between lines of great tits (*Parus major*) selected for divergent personalities. Gen. Comp. Endocrinol. 175, 488–494.

Boitani, L., Fuller, T. (Eds.), 2000. Research techniques in animal ecology: controversies and consequences. Columbia University Press.

Bonier, F., Martin, P.R., Moore, I.T., Wingfield, J.C., 2009. Do baseline glucocorticoids predict fitness? Trends Ecol. Evol. 24, 634–642. https://doi.org/10.1016/j.tree.2009.04.013

Bowers, E.K., Jenkins, J.B., Mueller, A.J., Miller, K.D., Thompson, C.F., Sakaluk, S.K., 2019. Condition-Dependent Begging Elicits Increased Parental Investment in a Wild Bird Population. Am. Nat. 193, 725–737. https://doi.org/10.1086/702848

Broom, D.M., Johnson, K.G., 2019. Stress and Animal Welfare: Key Issues in the Biology of Humans and Other Animals, Animal Welfare. Springer International Publishing, Cham. https://doi.org/10.1007/978-3-030-32153-6

Canoine, V., Hayden, T., Rowe, K., Goymann, W., 2002. The stress response of European stonechats depends on the type of stressor. Behaviour 139, 1303–1311. https://doi.org/10.1163/156853902321104172

Chastel, O., Lacroix, A., Weimerskirch, H., Gabrielsen, G., 2005. Modulation of prolactin but not corticosterone responses to stress in relation to parental effort in a long-lived bird. Horm. Behav. 47, 459–466. https://doi.org/10.1016/j.yhbeh.2004.10.009

Crespi, E.J., Williams, T.D., Jessop, T.S., Delehanty, B., 2013. Life history and the ecology of stress: how do glucocorticoid hormones influence life-history variation in animals? Funct. Ecol. 27, 93–106. https://doi.org/10.1111/1365-2435.12009

Curzer, H.J., Wallace, M.C., Perry, G., Muhlberger, P.J., Perry, D., 2013. The Ethics of Wildlife Research: A Nine R Theory. ILAR J. 54, 52–57. https://doi.org/10.1093/ilar/ilt012

De Kloet, E.R., Karst, H., Joëls, M., 2008. Corticosteroid hormones in the central stress response: Quick-and-slow. Front. Neuroendocrinol. 29, 268–272. https://doi.org/10.1016/j.yfrne.2007.10.002

Dickens, M.J., Romero, L.M., 2013. A consensus endocrine profile for chronically stressed wild
animals does not exist. Gen. Comp. Endocrinol. 191, 177–189.
https://doi.org/10.1016/j.ygcen.2013.06.014
Duarte, L.M.G., 2013. Impacts of capture and handling on wild birds (Doctoral thesis). Cardiff
UK Cardiff Univ.
Gelman, A., Su, Y.-S., 2021. arm: Data Analysis Using Regression and Multilevel/Hierarchical
Models. R Package Version 112-2.
Gillies, N., Fayet, A.L., Padget, O., Syposz, M., Wynn, J., Bond, S., Evry, J., Kirk, H., Shoji,
A., Dean, B., Freeman, R., Guilford, T., 2020. Short-term behavioural impact contrasts
with long-term fitness consequences of biologging in a long-lived seabird. Sci. Rep. 10,
15056. https://doi.org/10.1038/s41598-020-72199-w
Hau, M., Casagrande, S., Ouyang, J.Q., Baugh, A.T., 2016. Glucocorticoid-Mediated
Phenotypes in Vertebrates, in: Advances in the Study of Behavior. Elsevier, pp. 41–115.
https://doi.org/10.1016/bs.asb.2016.01.002
Hau, M., Deimel, C., Moiron, M., 2022. Great tits differ in glucocorticoid plasticity in response
to spring temperature. Proc. R. Soc. B 289, 20221235.
Hinde, C.A., 2006. Negotiation over offspring care?—a positive response to partner-
provisioning rate in great tits. Behav. Ecol. 17, 6–12.
https://doi.org/10.1093/beheco/ari092
Horton, B.M., Holberton, R.L., 2009. Corticosterone manipulations alter morph-specific
nestling provisioning behavior in male white-throated sparrows, *Zonotrichia albicollis*.
Horm. Behav. 56, 510–518. https://doi.org/10.1016/j.yhbeh.2009.09.001
Huber, N., Mahr, K., Tóth, Z., Szarka, E.Z., Çınar, Y.U., Salmón, P., Lendvai, Á.Z., 2021. The
stressed bird in the hand: Influence of sampling design on the physiological stress
response in a free-living songbird. Physiol. Behav. 238, 113488.
https://doi.org/10.1016/j.physbeh.2021.113488
Hughes, E.J., Mady, R.P., Bonter, D.N., 2021. Evaluating the accuracy and biological meaning
of visits to RFID-enabled bird feeders using video. Ecol. Evol. 17132–17141.
https://doi.org/10.1002/ece3.8352
Husby, A., Hille, S.M., Visser, M.E., 2011. Testing Mechanisms of Bergmann's Rule:
Phenotypic Decline but No Genetic Change in Body Size in Three Passerine Bird
Populations. Am. Nat. 178, 202–213. https://doi.org/10.1086/660834
Iserbyt, A., Griffioen, M., Borremans, B., Eens, M., Müller, W., 2018. How to quantify animal
activity from radio-frequency identification (RFID) recordings. Ecol. Evol. 10166–
10174. https://doi.org/10.1002/ece3.4491
Kaplan, G., 2022. Casting the Net Widely for Change in Animal Welfare: The Plight of Birds
in Zoos, Ex Situ Conservation, and Conservation Fieldwork. Animals 12, 31.
https://doi.org/10.3390/ani12010031
Keller, L.F., Noordwijk, A.J.V., 1993. A Method to Isolate Environmental Effects on Nestling
Growth, Illustrated with Examples from the Great Tit (*Parus major*). Funct. Ecol. 7,
493. https://doi.org/10.2307/2390037
Kitaysky, A.S., Wingfield, J.C., Piatt, J.F., 1999. Dynamics of food availability, body condition
and physiological stress response in breeding Black-legged Kittiwakes: Food
availability and stress response in Kittiwakes. Funct. Ecol. 13, 577–584.
https://doi.org/10.1046/j.1365-2435.1999.00352.x
Lack, D., 1968. Lack, D. 1968. Ecological adaptations for breeding in Birds. Methuen, London.
Methuen Press, London.
Lane, S.J., Fossett, T.E., VanDiest, I.J., Sewall, K.B., 2025. Recovery through resistance?
Nesting urban female song sparrows (*Melospiza melodia*) have a lower glucocorticoid
response to disturbance and return to parental care as quickly as rural females. Front.
Physiol. 16, 1520208. https://doi.org/10.3389/fphys.2025.1520208

Lattin, C.R., Kelly, T.R., 2020. Glucocorticoid negative feedback as a potential mediator of
trade-offs between reproduction and survival. Gen. Comp. Endocrinol. 286, 113301.
https://doi.org/10.1016/j.ygcen.2019.113301

Lynn, S.E., Prince, L.E., Phillips, M.M., 2010. A single exposure to an acute stressor has lasting
consequences for the hypothalamo–pituitary–adrenal response to stress in free-living
birds. Gen. Comp. Endocrinol. 165, 337–344.
https://doi.org/10.1016/j.ygcen.2009.07.018

Marks, C., Allen, L., Gigliotti, F., Busana, F., Gonzalez, T., Lindeman, M., Fisher, P., 2004.
Evaluation of the tranquilliser trap device (TTD) for improving the humaneness of
dingo trapping. Anim. Welf. 13, 393–399. https://doi.org/10.1017/S0962728600028645

McEwen, B.S., Wingfield, J.C., 2003. The concept of allostasis in biology and biomedicine.
Horm. Behav. 43, 2–15. https://doi.org/10.1016/S0018-506X(02)00024-7

Mellor, D.J., Beausoleil, N.J., Littlewood, K.E., McLean, A.N., McGreevy, P.D., Jones, B.,
Wilkins, C., 2020. The 2020 Five Domains Model: Including Human–Animal
Interactions in Assessments of Animal Welfare. Animals 10, 1870.
https://doi.org/10.3390/ani10101870

Mentesana, L., Casagrande, S., Hau, M., 2024. Baseline glucocorticoids alone do not predict
reproductive success across years, but in interaction with enzymatic antioxidants. Ecol.
Evol. 14, e11193. https://doi.org/10.1002/ece3.11193

Monaghan, P., Spencer, K.A., 2014. Stress and life history. Curr. Biol. 24, R408–R412.
https://doi.org/10.1016/j.cub.2014.04.017

Mutzel, A., Dingemanse, N.J., Araya-Ajoy, Y.G., Kempenaers, B., 2013. Parental provisioning
behaviour plays a key role in linking personality with reproductive success. Proc. R.
Soc. B Biol. Sci. 280, 20131019. https://doi.org/10.1098/rspb.2013.1019

Nicolaus, M., Bouwman, K.M., Dingemanse, N.J., 2008. Effect of PIT Tags on the Survival
and Recruitment of Great Tits *Parus major*. Ardea 96, 286–292.
https://doi.org/10.5253/078.096.0215

Noordwijk, A.J.V., Balen, J.H.V., Scharloo, W., 1988. Heritability of body size in a natural
population of the Great Tit (*Parus major*) and its relation to age and environmental
conditions during growth. Genet. Res. 51, 149–162.
https://doi.org/10.1017/s0016672300024162

Norris, K.J., 1990. Female choice and the quality of parental care in the great tit *Parus major*.
Behav. Ecol. Sociobiol. 27. https://doi.org/10.1007/BF00164900

Ouyang, J.Q., Hau, M., Bonier, F., 2011. Within seasons and among years: when are
corticosterone levels repeatable? Horm. Behav. 60, 559–564.

Ouyang, J.Q., Sharp, P., Quetting, M., Hau, M., 2013. Endocrine phenotype, reproductive
success and survival in the great tit, Parus major. J. Evol. Biol. 26, 1988–1998.
https://doi.org/10.1111/jeb.12202

Poole, T., 1997. Happy animals make good science. Lab. Anim. 31, 116–124.
https://doi.org/10.1258/002367797780600198

Prescott, M.J., Lidster, K., 2017. Improving quality of science through better animal welfare:

the NC3Rs strategy. Lab Anim. 46, 152–156. https://doi.org/10.1038/laban.1217R Core Team,

2024. R: A Language and Environment for Statistical Computing.

Romero, L.M., 2004. Physiological stress in ecology: lessons from biomedical research. Trends
Ecol. Evol. 19, 249–255. https://doi.org/10.1016/j.tree.2004.03.008

Romero, L.M., Beattie, U.K., 2022. Common myths of glucocorticoid function in ecology and
conservation. J. Exp. Zool. Part Ecol. Integr. Physiol. 337, 7–14.
https://doi.org/10.1002/jez.2459

Romero, L.M., Butler, L.K., 2007. Endocrinology of Stress. Int. J. Comp. Psychol. 20. https://doi.org/10.46867/IJCP.2007.20.02.15

Romero, L.M., Reed, J.M., 2005. Collecting baseline corticosterone samples in the field: is under 3 min good enough? Comp. Biochem. Physiol. A. Mol. Integr. Physiol. 140, 73–79. https://doi.org/10.1016/j.cbpb.2004.11.004

Romero, L.M., Reed, J.M., Wingfield, J.C., 2000. Effects of Weather on Corticosterone Responses in Wild Free-Living Passerine Birds. Gen. Comp. Endocrinol. 118, 113–122. https://doi.org/10.1006/gcen.1999.7446

Romero, L.M., Wingfield, J.C., 2015. Tempests, Poxes, Predators, and People: Stress in Wild Animals and How They Cope, Oxford Series in Behavioral Neuroendocrinology, Oxford Series in Behavioral Neuroendocrinology. Oxford University Press, New York.

Russell, W.M.S., Burch, R.L., 1959. The principles of humane experimental technique. Methuen Press, London.

Saltzman, W., Abbott, D.H., 2009. Effects of elevated circulating cortisol concentrations on maternal behavior in common marmoset monkeys (*Callithrix jacchus*). Psychoneuroendocrinology 34, 1222–1234. https://doi.org/10.1016/j.psyneuen.2009.03.012

Sapolsky, R.M., Romero, L.M., Munck, A.U., 2000. How Do Glucocorticoids Influence Stress Responses? Integrating Permissive, Suppressive, Stimulatory, and Preparative Actions 21.

Schlicht, E., Kempenaers, B., 2015. Immediate effects of capture on nest visits of breeding blue tits, *Cyanistes caeruleus*, are substantial. Anim. Behav. 105, 63–78. https://doi.org/10.1016/j.anbehav.2015.04.010

Silverin, B., 1986. Corticosterone-binding proteins and behavioral effects of high plasma levels of corticosterone during the breeding period in the pied flycatcher. Gen. Comp. Endocrinol. 67–74. https://doi.org/10.1016/0016-6480(86)90029-8

Soulsbury, C.D., Gray, H.E., Smith, L.M., Braithwaite, V., Cotter, S.C., Elwood, R.W., Wilkinson, A., Collins, L.M., 2020. The welfare and ethics of research involving wild animals: A primer. Methods Ecol. Evol. 11, 1164–1181. https://doi.org/10.1111/2041-210X.13435

Székely, T., Webb, J.N., Houston, A.I., McNamara, J.M., 1996. An Evolutionary Approach to Offspring Desertion in Birds, Current Ornithology. Springer, Boston, MA.

Taff, C.C., Zimmer, C., Vitousek, M.N., 2018. Efficacy of negative feedback in the HPA axis predicts recovery from acute challenges. Biol. Lett. 14, 20180131. https://doi.org/10.1098/rsbl.2018.0131

Van Oers, K., Drent, P.J., De Jong, G., Van Noordwijk, A.J., 2004. Additive and nonadditive genetic variation in avian personality traits. Heredity 93, 496–503. https://doi.org/10.1038/sj.hdy.6800530

Veen, I.T.V.D., Sivars, L.E., 2000. Causes and consequences of mass loss upon predator encounter: feeding interruption, stress or fit-for-flight? Funct. Ecol. 14, 638–644. https://doi.org/10.1046/j.1365-2435.2000.t01-1-00465.x

Vitousek, M.N., Taff, C.C., Hallinger, K.K., Zimmer, C., Winkler, D.W., 2018. Hormones and Fitness: Evidence for Trade-Offs in Glucocorticoid Regulation Across Contexts. Front. Ecol. Evol. 6, 42. https://doi.org/10.3389/fevo.2018.00042

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verl. N. Y.

Wilson, R.P., McMahon, C.R., 2006. Measuring devices on wild animals: what constitutes acceptable practice? Front. Ecol. Environ. 4, 147–154. https://doi.org/10.1890/1540-9295(2006)004%255B0147:MDOWAW%255D2.0.CO;2

Wingfield, J.C., Maney, D.L., Breuner, C.W., Jacobs, J.D., Lynn, S., Ramenofsky, M., Richardson, R.D., 1998. Ecological Bases of Hormone—Behavior Interactions: The

"Emergency Life History Stage." Am. Zool. 38, 191–206.
https://doi.org/10.1093/icb/38.1.191
Wingfield, J.C., Sapolsky, R.M., 2003. Reproduction and Resistance to Stress: When and How:
Reproduction and resistance to stress. J. Neuroendocrinol. 15, 711–724.
https://doi.org/10.1046/j.1365-2826.2003.01033.x
Wingfield, J.C., Smith, J.P., Farner, D.S., 1982. Endocrine Responses of White-Crowned
Sparrows to Environmental Stress. The Condor 84, 399.
https://doi.org/10.2307/1367443
Wingfield, J.C., Vleck, C.M., Moore, M.C., 1992. Seasonal changes of the adrenocortical
response to stress in birds of the Sonoran desert. J. Exp. Zool. 264, 419–428.
https://doi.org/10.1002/jez.1402640407
Wu, F., Tilbrook, A., Maloney, S.K., Blache, D., 2025. The intricate relationship between stress
and animal welfare: from historical perspective to new avenues. Biol. Rev. brv.70057.
https://doi.org/10.1111/brv.70057
Zimmer, C., Boogert, N.J., Spencer, K.A., 2013. Developmental programming: Cumulative
effects of increased pre-hatching corticosterone levels and post-hatching unpredictable
food availability on physiology and behaviour in adulthood. Horm. Behav. 64, 494–500.
https://doi.org/10.1016/j.yhbeh.2013.07.002
942