Title

Maintenance cost of photosynthesis sets key ecological constraints on zooxanthellate corals

Authors
Kelly Gomez-Campo[1,2,3]*, M. Isabel Martinez-Rugerio[1,2,4], Miguel Ángel Gómez Reali[5], Raúl A. González-Pech[3,6], Michel Geovanni Santiago-Martínez[7], Kira E. Turnham[3], Todd C. LaJeunesse[3], Iliana B. Baums[1,2,3,4], Susana Enríquez[5], Roberto Iglesias-Prieto[1]*

Affiliations
[1]Helmholtz-Institute for Functional Marine Biodiversity at the University of Oldenburg (HIFMB), Im Technologiepark 5, 26129, Germany.
[2]Alfred Wegener Institute, Helmholtz-Centre for Polar and Marine Research (AWI), Bremerhaven, Germany.
[3]Department of Biology, The Pennsylvania State University, University Park, PA 16802, USA.
[4]Institute for Chemistry and Biology of the Marine Environment (ICBM), School of Mathematics and Science, Carl von Ossietzky Universität Oldenburg, Oldenburg, Germany,
[5]Unidad Académica de Sistemas Arrecifales Puerto Morelos, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, 77580, México.
[6]Department of Biology, Texas State University, San Marcos, TX 78666, USA.
[7]Department of Molecular and Cell Biology, The University of Connecticut, Storrs, Connecticut, 06269, USA.

Corresponding authors: KGC kelly.gomezcampo@hifmb.de and RIP rzi3@psu.edu

Key words: Symbiodiniaceae, corals, symbiosis, light acclimation, D1 protein, photosystem II, photodamage, PSII repair.

Abstract

Ecological models using light limitation to explain coral depth distribution have largely disregarded the energetic cost of sustaining photosynthetic activity. Here, we quantified photosystem II (PSII) turnover across a depth-simulated light gradient in a zooxanthellate coral, measuring PSII half-life, D1 protein abundance, and PSII-complex gene expression. Maximum photosynthetic capacity remained stable across irradiance levels while respiration rose and PSII turnover accelerated as a power law, imposing increasing ATP demand at the shallowest depths. Declining D1 protein abundance alongside stable transcript levels demonstrated that this escalating maintenance cost operates through post-transcriptional regulation. Consequently, a decreasing fraction of photosynthetic usable energy is available for translocation to the coral host at high irradiance, as the energy required for PSII repair increases. Integrating these physiological constraints into a bio-optical model revealed that the balance between photosynthetic capacity and its maintenance cost defines an optimal depth, the Photosynthetic Usable Energy Supply (*PUES*) maximum, where host energetic returns are maximized. This framework provides a mechanistic basis for understanding depth distributions in symbiotic corals and extends as a predictive tool for any photosynthetic organism operating under variable irradiance, including forecasting how environmental degradation contracts viable depth ranges.

MAIN TEXT

Introduction

Light supplies the energy that sustains coral reef ecosystems, but its role in shaping coral
distribution cannot be explained by light availability alone. Variation in light quantity and
spectral composition set physiological limits that structure community composition,
species richness, and colony morphology across depth gradients (*1–6*). These ecological
patterns emerge from fundamental biophysical constraints on how efficiently light energy
is captured and converted into metabolic output. Corals are modular, sessile, mixotrophic
organisms that host intracellular dinoflagellates in the family Symbiodiniaceae (*7*).
Endosymbionts translocate a large fraction of their photosynthetically fixed carbon to the
host, fueling basal metabolism and calcification (*8–10*). Light harvesting by the coral
holobiont is determined by the emergent optical properties of symbionts, host tissue, and
skeleton (*11*). Consequently, efficient photosynthesis requires coordinated adjustment
across multiple levels of biological organization, from symbiont cell pigmentation and
density, to host tissue optics, skeletal architecture, and whole colony morphology (*1, 12–*
26). These traits are integrated through photoacclimatory processes that optimize energy
conversion under different light environments (*24, 27–30*). Yet, the physiological limits
through which photosynthesis constrains coral ecological niches across depth remain
unresolved (*5, 9, 31, 32*).

Depth gradients impose continuous selective pressures on coral–symbiont associations as
sunlight simultaneously determines energy acquisition and physiological demands. If coral
distribution were determined by solar energy alone, productivity and species richness
would be expected to peak in shallow reefs and decline monotonically with depth as light
attenuates. This pattern would reflect reduced growth rates and increased extinction risk
under lower energy supply (*33*). However, for symbiotic corals, light availability by itself
often fails to predict vertical patterns of biodiversity and performance (*3*). Instead, the
amount of light that can be converted into biologically usable energy for the host better
explains species richness and depth zonation in zooxanthellate corals (*5*). This distinction
has relevant ecological implications. At intermediate depths, many corals maximize their
net usable energy, expanding niche space and promoting greater species coexistence
relative to both shallow and deep extremes. Under this framework, depth zonation cannot
be simply considered a response to light attenuation with depth, but the outcome of
selection on traits that determine how efficiently corals convert depth-specific available
light into net energetic return. This is achieved through coordinated optimization of coral
light capture, physiological adjustments of symbionts and host, and colony morphology.
While lower depth limits are attributed to light limitation, upper depth limits are
hypothesized to be constrained by the increasing energetic demands required to maintain
performance under high irradiance.

Among the physiological processes that impose these energetic demands, protein damage
and repair rank among the most energetically expensive cellular processes across living
systems (*34–37*). However, their contribution to whole-organism energy budgets remains
difficult to quantify. In photosynthetic organisms, this challenge is exemplified by the
photosystem II (PSII) reaction center D1 protein, which undergoes rapid, light-dependent
cycles of damage, degradation, and resynthesis (*38–43*). Replacement of photodamaged
PSII reaction centers requires substantial ATP investment to support D1 synthesis,
assembly, and regulation (*44–47*). In coral-associated Symbiodiniaceae, accelerated D1

turnover has been documented under both light- and thermal stress, consistent with a central role in sustaining photosynthetic function under stress conditions and maintaining the stability of the mutualism (*48–50*). Thus, empirical data on the energetic cost of sustaining this repair activity, and its consequences for energy partitioning within the symbiosis, are critical to determine energy utilization and delineate energetic boundaries in depth gradients.

We hypothesize that changes in PSII damage and repair across a light gradient affect the net energy that can be translocated from the symbiont to the coral host. Enhanced repair in high light environments sustains photosynthesis but requires more energy for ATP production and protein synthesis, reducing the fraction of photosynthetic energy available in support of host metabolism. Conversely, low light conditions limit energy production regardless of maintenance demands. These mechanisms predict a non-linear relationship between light availability and net autotrophic benefit to the coral host, providing an explanation for why abundant light in shallow reefs does not necessarily translate into enhanced coral growth or physiological performance (*5*).

To measure the effect of metabolic cost across high and low light conditions, we conducted a series of experiments to characterize coral photoacclimatory responses, quantify energetic demands of PSII damage-repair activity, and its impact on energy translocation to the coral host (**Fig. S1**). First, (1) we engineered a custom light system capable of simulating natural diurnal light cycles under controlled irradiance and spectral quality. This system minimized the confounding effects of ultra violet (UV) light (**Fig. S2**) and stabilized daily light integrals (DLI in mol quanta m^{-2} day^{-1}) against the variability caused by cloud cover (*51*, *52*). Second, (2) coral fragments from the depth generalist *Orbicella faveolata* were acclimated to DLI treatments for four weeks, followed by a detailed characterization of its optical, structural, and physiological light phenotypes, including PSII half-life and turnover dynamics (**Table 1**). Third, (3) coral fragments were exposed to short-term shifts in DLI to evaluate repair capacity under light conditions that induce light limitation and light stress. Lastly, (4), we developed a bio-optical model integrating empirical results and theoretical considerations to predict shifts in photosynthetic usable energy available to the coral host across a depth gradient. This mechanistic framework provides a tool to predict the natural habitat and depth distribution of various coral taxa as well as quantify the impact of environmental changes and increase turbidity (diminished water quality) on the energetic cost to reef-building corals.

Results

Rapid photoacclimation across a realistic depth-dependent light gradient

Colonies in replicates were exposed to daily light integrals of 4 (DLI-4), 12 (DLI-12), 22 (DLI-22), and 32 (DLI-32) mol quanta m^{-2} day^{-1} (**Fig. 1A**, **Fig. S3A**), which simulates a depth gradient of 5–34 m (assuming a diffuse attenuation coefficient for downwelling irradiance, K_d, of 0.07 m^{-1}; typical for clear-water tropical reefs). PSII maximum photochemical efficiency (F_v/F_m) for experimental replicates (sample fragments) reached a steady state by day 10 (**Fig. S3B**), indicating that samples had acclimated to each of their respective daily doses of irradiance.

Diurnal photoprotection scaled with light exposure

Traits were treated as continuous functions of DLI rather than as discrete treatment means, characterizing phenotypic adjustments across the light gradient (**Fig. S4**). F_v/F_m was highest under DLI-4 and declined linearly with increasing light intensity (**Fig. 1B**). To quantify diurnal variation in PSII excitation pressure, we measured maximum pressure over PSII (Qm), which increased linearly corresponding to the transition from low to high irradiance (**Fig. S3C**). Qm was positively correlated with non-photochemical quenching (NPQ; $r^2 = 0.96$), consistent with increases in photoprotective energy dissipation capacity as excitation pressure increased.

PSII photochemical efficiencies, maximum (F_v/F_m) and effective ($\Delta F/F_m'$), measured every hour captured diurnal variation in PSII performance over the diurnal cycle. Differences in photochemical efficiency during the day were driven by light exposure, with a characteristic decline during the morning, a minimum at midday, and a reciprocal recovery in the afternoon (**Fig. 1C**). These diurnal dynamics reflected the balance among photoprotective mechanisms: induced dissipation as heat of the energy absorbed in excess (NPQ), induced PSII inactivation, and recovery (including NPQ relaxation) over the course of the day. The amplitude of diurnal oscillations increased proportionally with light exposure, reaching a maximum under the highest light treatment (DLI-32), which resulted in progressive increases in NPQ (**Fig. 1D**).

Optical reorganization reduces bulk light absorption under high irradiance

Coral tissue pigmentation darkened or lightened in response to the corresponding light treatment, resulting in systematic changes in tissue optical properties (**Fig. 2A–B, Fig. S3D, Table S1**). From minimum to maximum exposure, Chlorophyll a (Chl$a$) content declined by $\approx$ 36% (112 $\pm$ 13 to 41 $\pm$ 4 mg Chl a m^{-2}), while intact tissue absorbance peak at 675 nm (De_{675}) decreased by $\approx$ 11%. This corresponded to a decline in average absorptance in the PAR range (A_{PAR}) of 6% from low to high light conditions. The reduction in Chl$a$ pigment and slight reduction in bulk absorption capacity resulted in a non-linear increase in holobiont efficiency to absorb light (chlorophyll-specific absorption coefficient a*$_{Chla}$) (**Fig. 2B**).

Photosynthetic capacity is maintained despite increasing respiratory demands

Metabolic rates were quantified using photosynthesis–irradiance (P/E) curves fitted with a
hyperbolic tangent model, from which four photosynthetic parameters were extracted:
photosynthetic efficiency (α), maximum photosynthetic rate (P_{max}), saturation irradiance
(E_k), and compensation irradiance (E_c) (53). Photosynthetic efficiency (α) increased as
light exposure decreased, indicating enhanced efficiency under low irradiance conditions
(**Fig. 2C**). In contrast, maximum photosynthetic rate (P_{max}) remained stable across all light
treatments at approximately 12 ± 0.33 μmol O_2 m^{-2} s^{-1} (**Fig. 2D**), indicating that
photoacclimation preserved maximal photosynthetic capacity despite large differences in
light exposure and pigmentation.

P/E curves allowed estimation of dark respiration rates (Rd), which increased linearly with
light exposure (**Fig. 2E**), indicating a higher ATP demand in corals acclimated to higher
light conditions. The phototrophic contribution to host energy requirements also increased
with light exposure, as reflected by the higher daily integrated photosynthesis-to-
respiration ratios (54) (**Fig. S3E**). The minimum quantum requirements of oxygenic
photosynthesis (Φ^{-1}_{O2}) ranged from 10 to 17 quanta O_2^{-1} across treatments (**Fig. 2F**) and
the quantum efficiency of the photosynthetic process (Φ_{O2}) from 0.1 to 0.059 mol O_2
released per mol quanta absorbed. Both parameters revealed values close to the theoretical
minimum of 8 quanta O_2^{-1} for Φ^{-1}_{O2} and the theoretical maximum of 0.125 for Φ_{O2} (55,
56) for functional oxygenic photosynthesis.

PSII functional half-life shortens sharply with increasing light exposure

Having quantified how absorbed light energy is dissipated as excitation pressure increases,
and the resulting metabolic adjustments, we next examined processes regulating
maintenance of functional PSII reaction centers under contrasting light conditions. First, to
separate damage accumulation (before noon) from repair (before dusk), we inhibited *de
novo* protein synthesis using chloramphenicol (CAP) (**Fig. 3A**). In control corals (−CAP),
PSII activity declined through midday and recovered by the end of the light cycle as
expected, whereas CAP-treated corals (+CAP) showed sustained loss of PSII activity
consistent with inhibited repair.

This approach enabled estimation of PSII functional half-life ($t_{1/2}$, h), defined as the
number of daylight hours required to inactivate 50% of functional PSII reaction centers
(**Fig. 3B**). PSII half-life declined exponentially from ~13 h under low light to ~3 h under
high light exposure (**Fig. 4A**). This resulted in PSII turnover rates (τ_{PSII} day^{-1}) increasing
significantly from low to high light exposure (**Fig. 4B**), revealing accelerated replacement
of photodamaged reaction centers under high irradiance.

To assess molecular regulation underlying these dynamics, we quantified transcriptional
and translational responses of major PSII reaction center proteins. Quantitative
immunoblotting revealed significant differences in D1 protein abundance across the light
conditions (**Fig. 4D–F**). D1 content declined sharply with increasing light exposure,
mirroring the exponential decrease in PSII half-life and supporting regulation at the
translational or post-translational level (**Fig. 4E**). Normalization of D1 content to
chlorophyll *a* content yielded the same pattern (**Fig. 4F)**. In contrast, transcript abundance
of the core components of the reaction center *psbA* (D1) and *psbD* (D2), and inner antenna
proteins (that flank the D1/D2 heterodimer) *psbB* (CP47) and *psbC* (CP43), showed no
systematic relationship with light treatment (**Fig. 4G**; **Fig. S5**).

PSII half-life rapidly adjusts to acute changes in light exposure

To quantify short-term adjustments in PSII half-life under acute light stress, corals acclimated to a given DLI were transferred to the three alternative light conditions for a day, following the same CAP protein synthesis inhibition protocol (**Fig. S6**). PSII half-life responded rapidly to the new light environment and was independent of the prior acclimatory state, indicating that short-term regulation was driven by the destination light condition rather than the acclimation history (**Fig. 4H**).

To place these responses in the appropriate environmental context, we compared changes in half-life of PSII activity under our experimental PAR, and under natural sunlight, which also includes ultraviolet radiation (UV). In both cases, PSII half-life exhibited comparable exponential declines as a function of the destination light condition (**Fig. 4H**). Corals transferred from lower light treatments (DLI-4, DLI-12, DLI-22) to the highest light level (DLI-32) converged on a PSII half-life of approximately 3 h, matching that of colonies acclimated to DLI-32. These results demonstrate that PSII half-life can be rapidly adjusted within a single day across a wide range of light environments.

Despite this rapid adjustment, short-term regulation was insufficient to prevent the accumulation of PSII photodamage under high excitation pressure conditions. The magnitude of light stress, conveyed as the balance between PSII damage and repair, and quantified as *relative-F_v/F_m* changes (see methods) (*57*), declined with increasing excess excitation energy (EEE) (**Fig. 4I**). This indicated an increase in the magnitude of physiological light stress due to the rapid accumulation of damaged PSII.

Energetic cost of photoacclimation constrains usable energy across depth

The bio-optical model developed here integrates theoretical considerations, empirical measurements, and equations describing phenotypic adjustments to predict the photosynthetic energy available for the coral host in depth-dependent light gradients (see methods). In a previous model implementation (*5*), community-averaged photophysiological responses were represented using a lumped-parameter approach, with photosynthetic and maintenance parameters optimized to reproduce observed biodiversity patterns across depth. Here, we incorporated light-dependent photoacclimatory profiles to explicitly predict photosynthetic energy allocation.

First, we computed DLI as a function of depth across a range of diffuse attenuation coefficients (K_d = 0.04–0.40 m^{-1}), spanning from clear to turbid water conditions, to account for variation in underwater light environments, (**Eq. 1–2; Fig. S7**). Then, for each optical scenario, daily gross photosynthetic output ($DPg_{(z)}$) was estimated from photosynthesis–irradiance curve parameters (**Eq. 3–4**), while depth-specific maintenance cost ($ECMpt_{(z)}$) was derived from empirically constrained PSII turnover rates (**Eq. 5**). These components were computed to predict the photosynthetic usable energy supply ($PUES$) available to the coral host after accounting for symbiont metabolic demands (**Eq. 6**).

Across all optical scenarios, $PUES$ exhibited a unimodal relationship with depth (**Fig. 5A**). In shallow waters, high irradiance required elevated maintenance cost to sustain photosynthetic rates. Consequently, an increasing fraction of photosynthetic energy was allocated to symbiont maintenance rather than host translocation. At greater depths, light

limitation reduced carbon fixation, constraining energy availability despite lower PSII
repair activity. Importantly, the depth at which *PUES* was maximized shifted
systematically with K_d (**Fig. 5B**). Consequently, clearer waters supported deeper energetic
optima, whereas increased turbidity compressed the energetic window for coral
distribution toward shallower depths. These results define an optimal bio-optical state for
coral performance that emerges from the balance between photosynthetic output and
maintenance cost, and whose position and extent depend on water-column optical
properties.

Discussion

This investigation provides a mechanistic basis for disentangling the apparent paradox
between light availability, coral physiological performance, and vertical distribution of
coral species by demonstrating that energetic constraints arise at both ends of the depth
gradient. Clearly, carbon acquisition is limited by low irradiance. However, in shallow
waters, high light intensity increases the metabolic cost of maintaining photosynthetic
rates. Across a typical light gradient, maximum photosynthetic capacity remained stable
(**Fig. 2D**) despite rising excitation pressure (**Fig. 1B, Fig. S3C**), enhanced photoprotective
capacity (**Fig. 1C-D**), and elevated respiration (**Fig. 2E**). This homeostasis under high
irradiance was sustained by accelerated PSII turnover and shortened reaction center half-
lives (**Fig. 4**), imposing increasing ATP demand. Consequently, greater light availability
(*i.e.* energy acquisition) did not directly translate into greater net energetic benefit to the
host. Instead, a growing fraction of photosynthetic usable energy was retained by the
symbiont for its cellular maintenance.

A bio-optical model based on these physiological dynamics predicted a unimodal
relationship between light and usable energy supply (**Fig. 5A**), defining an optimal depth
range bounded by the maintenance cost of living in shallow waters and carbon limitation
at depth (**Fig. 5B**). Photosynthetic Usable Energy Supply *(PUES)* was previously
conceptualized in the context of Species–Energy Theory (*33*) to explain patterns of coral
biodiversity (*5*); here we provide the physiological basis for these distributions by
incorporating the metabolic cost of sustaining photosynthesis under a wide range of light
conditions. Classic carbon-budget studies show that zooxanthellae translocate a large
proportion of their net daily fixed carbon to the coral host, with the fraction of host
respiratory demand varying substantially with species, depth, and light regime (*8, 58*).
Translocated products such as glycerol, glucose, and other metabolites that vary among
Symbiodiniaceae lineages (*7, 59–63*) support host metabolism, growth, and calcification,
whereas a smaller fraction is retained for symbiont cell growth (*58, 64*). Our results
demonstrate that this translocation is predictable. As irradiance increases, ATP-intensive
photoprotective processes, particularly rapid PSII repair via D1 turnover, reduce the
fraction of organic carbon available for host use.

The cost of maintaining cellular function is rarely included in photosynthesis or primary
productivity models, which traditionally rely on photosynthesis:respiration ratios and/or
Chl*a* to identify conditions of positive carbon balance (*5, 65–67*). By coupling depth-
dependent photosynthetic output with associated maintenance cost, *PUES* reframes coral
energetics by emphasizing photosynthetic usable energy for the host. The resulting bio-
optical model provides a tool to translate physiological limits to the ecological distribution
of photosynthetic organisms, which in the case of zooxanthellate corals, involves colony
growth and morphology as well as gamete production and the timing of reproduction.

Additionally, it offers a framework for predicting how environmental stressors contract
depth distributions by necessitating greater investment in metabolic maintenance.

The stability of P_{max} across the range of light exposure reveals the central energetic
problem: photoacclimation in the coral holobiont preserves photosynthetic capacity
despite large changes in light availability. Disentangling the mechanisms requires
examining each component of the acclimatory response – photoprotection, optical
adjustment, metabolic adjustments, and reaction center maintenance – and what each
reveals about the limits of metabolic flexibility.

The first and energetically cheapest line of defense against excess irradiance is thermal
dissipation via NPQ. The tight coupling between Q_m and NPQ ($r^2 = 0.96$) reflects the
dynamic photoprotective equilibrium, driven by xanthophyll cycling, that scales
continuously with excitation pressure across diel cycles and light conditions (*1, 16, 68–*
71), consistent with prior field and laboratory observations (*16, 70, 72–75*). Critically,
however, NPQ alone cannot prevent photodamage when excitation pressure exceeds the
capacity for thermal energy dissipation. The declining F_v/F_m with increasing irradiance
marks precisely this transition, which is the point where the NPQ buffer is insufficient and
repair via D1 turnover becomes the dominant maintenance strategy, with an escalating
ATP cost (*47, 88, 89*).

The optical organization of the coral host and its zooxanthellae operate under strict
acclimatory limits at both ends of the depth gradient. Coral skeleton and tissue optics have
evolved for light amplification rather than attenuation (*11, 24*). Despite a nearly threefold
decline in Chl*a* content across light conditions, overall absorptance remained effectively
constant, with total energy absorbed preserved regardless of pigment reduction (**Table
S1**). Rather than reducing total light absorption, pigment loss increased chlorophyll-
specific absorption efficiency (a*$_{Chla}$) as the holobiont approached its high-light
acclimatory limit (*11, 76, 77*). At this boundary, the capacity for further pigment reduction
is exhausted, leaving the symbiosis more vulnerable to additional stressors such as thermal
stress (*78*), which is consistent with the higher occurrence of bleaching in high-irradiance
environments. The same optical architecture that enables colonization of light-limited
deep reefs (*27, 29*) simultaneously constrains the upper irradiance tolerance of the
symbiosis, as it prevents meaningful reduction of absorbed energy through pigment
adjustment alone. Thus, in shallow reef environments, when this optical limit is reached
and repair capacity is overwhelmed, selection favors colony morphologies with structural
self-shading, *i.e.* branching, foliose, and massive forms with strong within-colony light
attenuation (*27, 30, 79*).

The stability of P_{max} across the range of light exposure marks a physiological constraint on
carbon fixation, not merely a correlate of photoacclimation. Additional irradiance above
saturation does not generate additional fixed carbon, hence it can only be dissipated as
heat or directed toward repair. Evidence for this limit emerges from both ends of the light
gradient. Low light phenotypes approached the theoretical maximum for PSII
photochemical efficiency in dinoflagellates *in hospite* ($F_v/F_m \approx 0.7$) and the theoretical
minimum quantum requirement for oxygen evolution (~10 quanta O_2^{-1}; **Fig. 2F**) (*55, 56*),
marking the lower boundary of photoacclimatory capacity (*1, 54*). On the other hand, high
light phenotypes, while more efficient at absorbing light (higher a*$_{Chla}$), were
proportionally less efficient at converting it into fixed carbon (lower α and Φ_{max}). The
symbiosis thus operates across a bounded photoacclimatory range between light-limited in

deep water and light-saturated in shallow water, with P_{max} as the stable outcome. This reveals that coral photoacclimation operates through modulation of light absorption, heat dissipation, and protein turnover rather than by regulating photosynthetic capacity. Such decoupling suggests that photosynthetic rates are constrained by downstream processes, such as inorganic carbon diffusion or alternative electron sinks within the symbiosome (*80–83*) and, more critically, demonstrates that maintaining photosynthetic performance under high light requires progressively greater metabolic support without proportional energetic return. The primary source of this increasing cost is PSII turnover and D1 protein dynamics.

Protein turnover is among the most energetically expensive cellular processes (*34–37*). In higher plants, thylakoid protein turnover accounts for 13–38% of total ATP consumption, with D1 replacement representing a major fraction of this budget (*37*). Energy for this repair cycle is supplied through respiration, linking PSII maintenance directly to metabolic demand. By preventing protein synthesis in the symbiont, we decoupled PSII damage from repair and quantified changes in PSII half-life across experimental light conditions. PSII half-life declined significantly with increasing irradiance and closely paralleled reductions in D1 protein abundance. Accordingly, corals acclimated to high light conditions maintained maximum photosynthetic capacity via faster replacement of photodamaged reaction centers and lower D1 pools. Similar relationships between D1 abundance, PSII half-life, and light history have been documented in higher plants, including how rapid D1 turnover accompanies elevated metabolic demand under high light exposure (*37, 84–86*). The elevated dark respiration (R_d) found under high light conditions is therefore consistent with increased ATP requirements for D1 synthesis and PSII reassembly (*36, 44, 87*). ATP availability is known to constrain *de novo* D1 synthesis and full PSII repair (*47, 88, 89*), suggesting that ATP supply becomes the limiting currency sustaining photosynthetic function under light stress.

In this context, D1 turnover operates as a regulated sink for excess excitation, preventing over-reduction of the electron transport chain and protecting PSI under acceptor-side limitations (*45, 86*). This hierarchy reflects structural cost optimization within the photosynthetic apparatus. Because antenna complexes are far more abundant ($\sim 7.6 \times 10^6$ cell^{-1}) and energetically more expensive than D1 protein ($\sim 1.0 \times 10^5$ cell^{-1}) (*90*), it makes selective turnover of PSII reaction centers a comparatively economical strategy. In fact, regulation of reactive oxygen species within the photosynthetic apparatus shows that while chlorophyll triplet states in antenna complexes are efficiently quenched by carotenoids, spatial constraints within the PSII reaction center prevent such protection, allowing singlet oxygen to target D1 directly and requiring rapid turnover as a controlled safeguard (*40, 91, 92*). D1 protein, then, becomes the unavoidable target of oxidative damage, making its rapid turnover a necessary mechanism to preserve photosynthetic integrity.

This feature of D1 damage and its costly turnover has direct ecological implications. Earlier work has primarily focused on how PSII repair impairment under thermal stress leads to coral bleaching. When repair fails to match damage rates, D1 protein abundance drops below the threshold needed to sustain photosynthetic activity, resulting in pigment loss, symbiont degradation, and ultimately, expulsion or symbiont digestion by the host (*48, 93–96*). In our experiments, even in the absence of heat stress, high light and PAR alone shortened PSII half-life from ~13 h to ~3 h (**Fig. 4B**) and reduced D1 abundance by 80% (**Fig. 4E**), imposing substantial ATP demand. Thus, the upper tolerance limits of the coral holobiont are defined not solely by photodamage, but by the energetic capacity to

sustain PSII repair. This constraint is always active in high light conditions, revealing the
high susceptibility of shallow water corals to any additional environmental stress. We
confirmed the environmental regulation of this mechanism by comparing PSII half-life
under experimental PAR (400–700 nm) and natural sunlight (PAR + UV) (**Fig. 4H**). No
significant differences in PSII half-life were found, consistent with PAR being the primary
driver of photoinactivation and daily damage accumulation. This result challenges the
assumption that UV is the dominant driver of PSII photodamage in shallow-water corals
(*83, 97*).

Ultimately, PSII repair is a fundamental energetic 'bottleneck' that constrains coral
holobiont performance at high irradiance. Sustaining photosynthesis under light stress
requires continuous ATP investment in protein repair and replacement, thereby diverting
resources from growth, reproduction, and calcification. In this way, energetic maintenance
cost of photosynthesis sets the upper limits of energy acquisition by reef corals and their
stress tolerance to episodes of acute environmental change.

A bio-optical model incorporating Photosynthetic Usable Energy Supply provides a
mechanistic scaling of coral bioenergetics to explain their ecological patterns of depth and
latitude distribution (**Fig. 5A**). The model predicts the fraction of photosynthetically
derived energy that is available for host metabolic requirements, including maintenance,
growth, reproduction, and ecological interactions, and thus, defines fundamental
ecological niches of the coral holobiont. In the bioenergetic framework, the *PUES* bio-
optical model, specifically *PUES* optimum (*Depth$_{opt}$*), defines the depth at which
photosynthetic supply and maintenance demand are optimally balanced, maximizing
usable energy translocation while minimizing the cost of sustaining photosynthetic
integrity (**Fig. 5B**). The *Depth$_{opt}$* position and extent of coral colonization depend on water
column optical properties. When exploring a range of light attenuation coefficients (K_d),
spanning clear to turbid conditions, our analysis revealed that increased light attenuation
shifts *Depth$_{opt}$* toward shallower waters and compresses the energetic envelope. Therefore,
environmental degradation, including progressive ocean darkening (*i.e*., increased
turbidity), drives a compression of the viable depth range for coral colonization and
growth.

The development of adaptations to better meet the energetic needs for living at various
depths and water conditions carries significant evolutionary consequences. Intermediate
depths, where energy use is maximized without excessive physiological costs, support
multiple photophysiological strategies and species coexistence. In contrast, the higher
energetic costs of protein maintenance in shallow water or extremes of carbon limitation at
greater depth, ultimately restrict possible trait combinations channeling ecological success
through specialization under strong selection pressure. Such patterns align with general
Species-Energy Theory and with emerging genomic evidence that closely related coral
lineages repeatedly diverge across narrow depth gradients, with adaptation to distinct light
conditions linked to environmental sensing and reproductive timing (*98–100*).

438
439 **Materials and Methods**
440
PAR LED Light System
We developed an experimental light system by engineering custom-made lamps and
software able to simulate diurnal light cycles. Natural light in the tanks was absent
throughout the experiments. The illumination system relied on 100W light-emitting diodes
(LED)-lamps that enabled automation of the diurnal cycles (**Fig. S3A**). The four light
treatments (Daily Light Integral, DLI) were: 4 mol quanta m^{-2} day^{-1} (DLI-4, peak of
irradiance 200 µmol quanta m^{-2} s^{-1}); 12 mol quanta m^{-2} day^{-1} (DLI-12; peak of irradiance
450 µmol quanta m^{-2} s^{-1}); 22 mol quanta m^{-2} day^{-1} (DLI-22; peak of irradiance 750 µmol
quanta m^{-2} s^{-1}); and 32 mol quanta m^{-2} day^{-1} (DLI-32; peak of irradiance 1100 µmol
quanta m^{-2} s^{-1}).
451
Three parameters were prioritized for the control of each light condition: light intensity,
light quality and time. We built each LED-lamp with a BC Series High CRI COB-LED-
400L-100W- Unit (YujiLEDS®) which best covered the sunlight spectrum in the range of
the Photosynthetic Active Radiation (PAR). LEDs were operated in continuous mode with
a multifunction I/O card (USB-6001, National Instruments Corp., USA) to avoid potential
artifacts on photosynthesis and fluorescence responses related to the "flashing effect" of
commercial LEDs at different pulsating frequencies (*101*). Light intensity was controlled
with a custom-made software in LabVIEW (Laboratory Virtual Instrument Engineering
Workbench, National Instruments Corp., USA). To eliminate natural sunlight inside
experimental tanks, we placed LED-lamps inside black boxes that fully covered each tank
(**Fig. S1**).
463
Diurnal light exposure (mol quanta m^{-2} day^{-1}) was described by integrating the per second
change in irradiance (µmol quanta m^{-2} s^{-1}) over a day as Daily Light Integral (DLI). Light
exposure as an integral has proven to be a useful tool for assessing irradiance accumulated
when the response of interest is growth or acclimation (*102*). DLIs conditions captured a
wide range of coral vertical distribution in clear waters (5 to 35 m depth for a
downwelling light attenuation coefficient, K_d, of 0.07 m^{-1}). Irradiance at depth depends on
the optical properties of the water column given by the Kd at any reef location. Irradiance
was calibrated and continuously monitored with the quantum sensor LI-1400 (LI-COR,
USA).
Light spectra also play an important role in photosynthesis performance. The effective
activation of the repair cycle of photodamaged PSII has its own action spectrum. Studies
have shown that UV-radiation damages PSII more effectively than visible light (*97*), and
that the reported action spectra are rather dissimilar in the range between 400 and 700 nm.
We used a sun-like spectrum and UV was excluded by design (**Fig. S2**). We were able to
effectively induce PSII damage repair cycles in the simulated DLIs treatments.
Acclimation to Daily Light Integrals
We sampled 64 fragments from 4 *Orbicella faveolata* colonies (each a different genet)
between 4-5 m depth under similar light conditions in La Bocana, Puerto Morelos
National Park. Experimental coral fragments (~5-9 cm^2) were set for a 2-week healing
period after sampling by placing them on PVC-deployed structures located at 5 m depth
near the sampling area. Corals were then transported to the National Autonomous

485 University of Mexico (UNAM) mesocosm facility, where each colony was equally
486 distributed in four outdoor tanks (shaded natural sunlight to 16 ± 3 mol quanta m^2 day^{-1})
487 equipped with running natural seawater (n = 16 fragments per tank). Water temperature
488 and flow were maintained at $28 \pm 1°C$ and 1100 L h^{-1} using commercial aquaria chillers
489 and heaters. Water temperature in the tanks was continuously monitored with Hobo data
490 loggers (Onset Computer Corporation, MA, USA). Corals were acclimated to the
491 mesocosm for two weeks before LED installation, followed by 4+ weeks under
492 experimental DLI conditions before phenotyping.

493 We monitored daily PSII maximum photochemical efficiency (F_v/F_m) at dusk and
494 effective photochemical efficiency ($\Delta F/F_m{'}$) at noon, with a pulse-amplitude modulated
495 fluorometer (Diving-PAM; Heinz Walz, Effeltrich). Maximum excitation pressure over
496 PSII ($Q_m = 1 - [(\Delta F/F_m{'}$ at noon$) / (F_v/F_m$ at dusk$)]$) was also estimated to monitor the
497 moderate light stress induced during natural and daily photo-acclimation, and the
498 adjustment to a stable photo-acclimatory condition (*1*).

499 **Phenotyping in the range distribution of zooxanthellate corals**

500 Following the 4+ weeks of acclimation to the experimental light gradient, we measured a
501 suite of traits to characterize coral phenotypes (**Table 1**). Chl*a* fluorescence measurements
502 were recorded as photochemical efficiency (F_v/F_m and $\Delta F/F_m{'}$), and non-photochemical
503 quenching was calculated as $NPQ = [(F_m-F_m{'})/F_m]$.

504 To describe changes in coral pigmentation, we estimated Chl*a* and Chl*c* contents. Briefly,
505 an air gun and filtered sea water (FSW) were used to obtain tissue slurries subsequently
506 homogenized at low temperature (Tissue-Tearor Homogenizer BioSpec Inc, USA) and
507 centrifuged. The resulting pellet was re-suspended in filtered seawater and preserved for
508 pigment concentration (extracted with acetone/dimethyl sulfoxide 95:5 vol/vol) and
509 Symbiodiniaceae cell density (counted in a haemocytometer after the addition of 200 μL
510 of iodine preservation solution). Chl*a* and Chl*c* concentrations were estimated
511 spectrophotometrically (3 reads per sample) with a modular spectrometer (Flame-T-UV-
512 VIS, Ocean Optics Inc., USA) using the equations described by Jeffrey and Humphrey
513 (*103*).

514 Light absorption spectra were determined on intact *in vivo* coral tissues from reflectance
515 (R) measurements. R was used to estimate Absorbance (De) $[De = \log (1/R)]$ and
516 Absorptance (A), assuming that Transmittance through the coral skeleton was negligible
517 (*104*). A was estimated for the PAR range (400 and 700 nm; $A_{PAR} = 1 - R_{PAR}$); and at the
518 675 peak of Chl*a* ($A_{675} = 1 - R_{675}$).

519 Specific-absorption coefficients (a*) were normalized to Chl*a* content (a*$_{Chla}$; m^{-2} mg
520 Pigm^{-1}) as $[$a$^{*}_{Chla}= (De_{675}/\rho)\cdot\ln(10)$, where $De_{675} = \log(1/R_{675})]$ (*11, 77, 105*). Likewise, a*
521 was normalized to symbiont cell density (a*$_{Sym}$; m^{-2} sym^{-1}) as $[$a$^{*}_{sym} = (De_{675}/$sym $\rho) \cdot$
522 $\ln(10)]$. These descriptors estimate the holobiont efficiency to absorb light (a*$_{Chla}$) and
523 symbiont specific light absorption efficiency (a^{*}_{sym}) (*78*).

524 To describe changes in coral metabolic rates, we used photosynthesis / irradiance curves
525 (P/E) fitted to a hyperbolic tangent model (**Fig. S4A**). Briefly, coral fragments were
526 incubated using a laboratory-made water-jacketed respirometer (*76*) with filtered sea
527 water at a constant temperature (28 °C) and constant water flow generated by continuous

agitation from magnetic stirrers, which ensure that the boundary layer around coral
surfaces is broken. One chamber was measured as a blank (no coral fragment) to correct
for background activity in the water. A LED-system was designed to enable automation of
light increments every 10 min. Oxygen evolution was measured continuously with a fiber-
optic oxygen meter system (FireSting, Pyroscience) to estimate the rate in the 10 min
range of: dark (pre-illumination), light (10 light increments), dark (post-illumination) for a
total period of the 2 h incubation. Photosynthetic efficiency (α), compensation irradiance
(E_c), saturation irradiance (E_k), respiration rates in the dark (R_D), light-enhanced
respiration (R_L), and maximum photosynthetic rates (P_{max}), were estimated from the PE
curves (*106, 107*). Each curve reflects the photosynthetic potential of the light phenotype.
To estimate the energy absorbed in excess, we calculated the hours of light exposure
above E_k (number of hours in P_{max}; H_{sat} (*108*)) and the magnitude of energy absorbed
under these conditions (mol quanta m^{-2} day^{-1}).

Using absorptance values determined above [$A = 1- R$], we calculated the maximum
quantum yield of photosynthesis (Φ_{O2}; mol oxygen evolved per mol quanta absorbed^{-1})
and its inverse, the minimum quantum requirements of photosynthesis [(Φ^{-1}_{O2}; $= 1/(\alpha *$
A_{PAR}); quanta required to release an O_2 molecule^{-1})] (*106, 107*). Φ_{O2} was estimated from
α, the slope of the linear regression of photosynthesis in the sub-saturating region of the
P/E curve; corrected for the light absorbed (A) in our experimental setting [(photosynthetic
utilizable radiation (PUR) absorbed (A_{PUR})] (*76, 106, 107*). To estimate the integrative
index of the holobiont's metabolism, we calculated daily integrated photosynthesis-to-
respiration ratios [Pg *(E)* day^{-1}: R *(E)* day^{-1}].

Algal symbiont identification

Orbicella faveolata hosts distinct species of algal symbionts in this region (*e.g.*,
Symbiodinium sp. A3, *Breviolum faviinorum*, *Breviolum meandrinum*, *Cladocopium* sp.
C3, *Cladocopium* sp. C7, and *Durusdinium trenchii*), with the possibility of multiple
species simultaneously inhabiting a same colony (*109–111*). To account for this functional
diversity, symbiont identity was determined using two complementary approaches.

(1) One cm^2 coral fragments were used to extract genomic DNA using a modified
Promega Wizard protocol (Promega Madison, WI) (*112, 113*). The large ribosomal
subunit (*LSU* rDNA) was amplified using *28S* forward (5'-
CCCGCTGAATTTAAGCATATAAGTAAGCGG-3') and *28S* reverse (5'-
GTTAGACTCCTTGGTCCGTGTTTCAAGA-3') (*114*) and an annealing temperature of
66 ºC. Amplifications were Sanger sequenced using the Big- Dye Terminator 3.1 Cycle
Sequencing Kit (ThermoFisher Scientific, Waltham, MA) and analyzed on the Applied
Biosciences Sequencer (Applied Biosciences, Foster City, CA) at the Penn State
University Genomics Core Facility. Sequences were checked and edited in Geneious
v11.0.5 (Biomatters Ltd, Auckland, NZ) and were queried against the NCBI nucleotide
database using BLASTn (National Center for Biotechnology Information).

(2) We separately aligned raw RNA-Seq reads (see below) from each sample against
reference transcriptomes using bwa v0.7.18 (*115*); mapping rates were determined using
SAMStat v2.2.3. The reference transcriptomes encompassed eight datasets from the most
closely related Symbiodiniaceae that *Orbicella* associates with: *Symbiodinium* sp. A3
(*116*), *Breviolum faviinorum* (*116*), *Breviolum* sp. B5 (*116*), *Cladocopium proliferum*
(*117*), *Cladocopium* sp. C1 MI (*118*), *Cladocopium* sp. C1 SM (*118*), and *Durusdinium*

trenchii (*117*, *118*). Symbionts present in each sample were determined by comparing
LSU rDNA and mapping results (**Fig. S5A**).

PSII half-life and turnover rates under acclimatory conditions

After the 4-week acclimation, the PSII damage and repair cycle was determined through
monitoring the variation of F_v/F_m and $\Delta F/Fm'$ in 12 h diurnal cycles. We tracked the
accumulation of photo-inactivated PSII and the accumulation of photo-repaired PSII. The
damage accumulation phase runs from dawn to midday, and repair from midday to dusk.

To quantify the effect of light acclimation on the photodamage and repair of the PSII
active pool, we inhibited the PSII repair process. In each acclimatory condition, fragments
were separated into two subgroups: control (n = 8, no inhibitor -CAP) and the inhibitory
treatment (n = 8, treated with D1 protein synthesis inhibitor chloramphenicol; +CAP).
Subgroups were placed separately in 1 L plastic containers with internal seawater
circulation (FSW) and positioned within tanks (water baths). Water was exchanged in the
containers every 2 h for both treatments (FSW+CAP) and controls (FSW-CAP). PSII
photochemical efficiency was measured every hour, and samples were snap frozen after
dusk for further analyses. This protocol was repeated for every light condition after 4+
weeks of acclimation. To estimate PSII half-life ($t_{1/2}$, h) photochemical efficiency
treatment-to-control ratios [+CAP / -CAP] were curve fitted using a logistic sigmoidal
function and expressed for each light condition as the number of hours required to
inactivate 50% of PSII ($t_{1/2}$, h).

Abundance of PSII D1 protein

Algal pellet assigned for protein analyses was flash frozen in liquid nitrogen and stored at
-80 °C. We quantified D1 protein content in algal symbionts isolated from three coral
fragments for each light regime. Each immunoblotting assay was performed multiple
times to load: (lane 1) a negative control, (lanes 2–6) 0.10-2.00 pmol *psbA*/D1 standard,
(lanes 7–10) one sample from each light regime to be normalized by total protein (µg
protein^{-1}), and (lanes 11–14) one sample from each light regime to be normalized by Chl *a*
(mg Chl a^{-1}). We performed quantitative immunoblotting following established protocols
(*93*, *94*, *96*, *119*, *120*). Briefly, proteins were extracted using protein extraction buffer PEB
(Agrisera AS08 300) supplemented with Pefabloc® SC (Sigma-Aldrich 11429868001) to
inhibit proteolysis. Samples were subjected to three cycles of sonication-to-liquid N_2 to
break cell wall and avoid over-heating. Protein concentrations were determined using The
Better Bradford assay with a standard curve. Equal amounts of total protein were mixed
with Bolt™ LDS sample buffer (Invitrogen B0007) and Bolt™ Sample Reducing Agent
(ThermoFisher B0004), denatured, and separated on NuPAGE™ 4–12% Bis-Tris gels
(Invitrogen NP0329BOX) alongside PageRuler™ Plus pre-stained protein ladder
(ThermoScientific 26619). Following electrophoresis, proteins were transferred to PVDF
membranes (Invitrogen LC2005) using Bolt™ transfer buffer (Invitrogen BT0006) and
blocked with ECL™ Advance Blocking Reagent (GE Healthcare GERPN418).
Membranes were probed with a rabbit polyclonal anti-PsbA (D1) antibody (Agrisera
AS05 084), followed by HRP-conjugated chicken anti-rabbit secondary antibody
(Agrisera AS10 833). Membranes were stained post-transfer with SimplyBlue™ SafeStain
(Invitrogen LC6060). D1 protein quantification was achieved by co-loading known
amounts of PsbA/D1 standard (Agrisera AS01 016S), imaging on a chemiluminescent
imager (ChemiDoc BIO-RAD), and pixel-quantifying in ImageJ.

Expression of PSII-complex genes

Expression of PSII genes from algal symbionts was assessed following an RNA-Seq approach. Total RNA was extracted from 16 fragments, 4 untreated fragments (-CAP) per light exposure. Each sample was homogenized in TRIzol reagent (Ambion, Life Technology) before centrifugation with chloroform for 15 min at 12,000 g at 4 °C. The aqueous phase was then isolated and cleaned using Qiagen RNeasy Mini kit (Qiagen), as per the manufacturer's protocol with an additional on-column DNase treatment using RNase-Free DNase Set (Qiagen). To maximize concentration of eluted RNA, the same 35 μL of RNAse-free water was twice passed through the spin column for the final isolation step. Concentration and purity were pre-screened via Qubit spectrometer and quality assessed (RNA integrity number; RIN > 7) with Agilent 2100 Bioanalyzer (Agilent Technologies). Purified RNA was sent to Biomarker Technologies (BMK) GmbH where concentration and purity were re-analyzed (Qubit® dsDNA BR Assay Kit on a Qubit® 2.0. Fluorometer). RNA libraries for 2×150 bp paired-end sequencing were non-stranded, prepared using Hieff NGS Ultima Dual-mode mRNA Library Prep Kit for Illumina (Yeasen) model:13533ES96. Samples were run in platform Illumina NovaSeq X.

Raw reads were assessed for quality using FastQC v0.12.1 (github.com/s-andrews/FastQC) and results visualized using MultiQC v1.28 (*121*), followed by filtering and trimming using Trimmomatic v0.39 (*122*). Pseudoalignments of the processed read pairs and quantification of expression were done using the quant function of kallisto v0.51.1 (*123*) in two manners: against all reference transcriptomes combined into a single database and against separate transcriptomes according to the symbiont type identified per sample. For the combined database, all reference transcriptomes were concatenated removing redundant sequences using cd-hit-est v4.8.1 (*124*) at a sequence similarity threshold of 0.99. The transcripts per million (TPM) output by kallisto were used as measure of gene expression level. Genes of the core complex of PSII were identified by a BLASTn v2.16.0 (*125*) search of Symbiodiniaceae GenBank records (JN557844 for *psbA*, HG515018 for *psbB*, HG515019 for *psbC*, and HG515020 for *psbD*) as queries against the reference transcriptomes with an e-value $\leq 10^{-5}$. The BLAST hit with the highest mean expression across samples per genus was selected as the representative transcript for each gene.

PSII half-life under light-stress conditions

To quantify the effect of light stress on the PSII photodamage and repair cycle, and the variation of PSII half-life ($t_{1/2}$, h), we replicated the experimental protocol to inhibit the PSII repair process. However, here the 4-weeks acclimated corals were switched (new coral fragments for each experiment) to an alternative light exposure (*i.e.*, DLI-4$_{acclimated}$ switched to DLI-12$_{destination}$, DLI-22$_{destination}$ and DLI-32 $_{destination}$) for one diurnal cycle. The experimental protocol was repeated for each light exposure for a total of 12 experiments: DLI-12$_{acclimated}$ switched to DLI-4$_{destination}$, DLI-22$_{destination}$ and DLI-32$_{destination}$; DLI-22$_{acclimated}$ switched to DLI-4$_{destination}$, DLI-12$_{destination}$ and DLI-32$_{destination}$; and DLI-32$_{acclimated}$ switched to DLI-4$_{destination}$, DLI-12$_{destination}$ and DLI-22$_{destination}$.

The severity of light stress was quantified with the metric Excess Excitation Energy *EEE* (mol quanta m^{-2} day^{-1}) as follows (*57*): (1) Conditions where *EEE* = 0 (*EEE$_0$*) indicate that corals are well acclimated (F_v/F_m is steady day-to-day), no net light stress is induced, hence no photodamage is accumulated. (2) Conditions where *EEE* ≠ 0 (*EEE$_{\neq 0}$)* indicate

corals were switched to a different light condition and may experience light stress. *EEE*
was quantified as the difference between DLI in the switched condition (DLI$_{destination}$) and
DLI acclimated (DLI$_{acclimated}$). In the case that the switched condition resulted in lower
light exposure, *EEE* shows negative values. The magnitude of the physiological stress was
measured as the ratio between F_v/F_m in the switched condition and F_v/F_m acclimated,
which results in the parameter *relative-F_v/F_m*.

PSII half-life (t$_{1/2}$, h) was estimated following the same protocol described above for PSII
half-life under acclimatory conditions as the number of hours required to inactivate 50%
of the active PSII pool.

Modeling the energetic cost of photo-acclimation

To estimate how the light environment shapes the energetic balance of the coral-algal
symbiosis, we developed a bio-optical model that integrates theoretical light attenuation in
the water column, gross photosynthetic output, and the energetic costs of PSII turnover.
The analysis focused on the estimation of the potentially maximum photosynthetic output
of the symbionts after subtracting the costs of maintenance of photosynthetic activity
through PSII repair, which can be translocatable to the host. Other metabolic costs
required by the maintenance and/or growth of algae and host were not considered into the
model due to the difficulties of parameterization of the large diverse of biological
processes potentially involved. Light availability at depth was expressed as daily light
integrals (DLI) and computed from surface irradiance. Different light attenuation
coefficients (K_d) were used. Depth-dependent photosynthetic output was then estimated
using a photosynthesis-irradiance relationship parameterized with function-valued
phenotyping derived from experimental profiles. To capture the costs of maintaining
photosynthetic function under different light regimes, we incorporated PSII turnover
dynamics by linking PSII lifetime (t$_{1/2}$, h) and its cost, maximum photochemical efficiency
(F_v/F_m), and repair potential to DLI. The resulting framework yields the Photosynthetic
Usable Energy Supply (*PUES*) by the symbionts to the host, a metric that represents the
potentially maximum energy available for the coral host after subtracting the light-
dependent costs of photosynthesis maintenance. Considerations and computations
underlying the model proceeded as follows:

Theoretical and empirical considerations

As light is the primary factor that regulates primary production and the photoacclimatory
response of photosynthetic organisms, we first estimated the amount of light available at a
given depth (*126*):

1. Solar irradiance as a function of depth

The intensity of a parallel-beam radiation of a particular wavelength decreases along its
path in pure water, obeying the Lambert Law. The general equation of light attenuation in
seawater is (*126*):

$$E_{(z)} = E_0 * e^{-Kd*z}$$ (Eq. 1)

Where $E_{(z)}$ is light intensity at depth z (here irradiance value for Photosynthetic Active
Radiation, PAR, as UVR was not considered), E_0 is the PAR just beneath the water

surface (when $z = 0$) and Kd (m^{-1}) is the vertical attenuation coefficient of downwelling
PAR irradiance.

2. Changes in daily light integral (DLI) with depth: Light exposure

Irradiance at any point on the Earth's surface is determined by solar elevation. This rises
during the day from zero at dawn to its maximum value at noon, and then diminishes in a
precisely symmetrical manner to zero at dusk under no cloud cover. Light exposure
integrates the variation in irradiance per second and m^2 over time. The integrated quantum
unit for light exposure at a certain depth and day is mol quanta m^{-2} day^{-1}, which describes
the amount of quanta received by m^{-2} during a 24 h period (*126*):

$$DLI_{(z)} = \int_0^N E_0 * Sin\left(\frac{\pi t}{N}\right) * e^{(-Kdz)}\, dt \tag{Eq. 2}$$

The parameter $DLI_{(z)}$ is then the daily light integral at depth, E_0 is the superficial irradiance
at solar noon, N is the daylength (in seconds) and t is time along a day (seconds).

3. Photosynthesis as a function of DLI

The response of photosynthetic organisms to available light (from the light saturation
curve), depends on two fundamental parameters in our model: the slope α at very low light
availability (carbon fixed/oxygen evolved per unit of available light), and the asymptotic
maximum production P_{max} (the rate of carbon fixation/oxygen evolved under saturating
light) (*127*):

$$GPg_{(DLI)} = Pmax * \tanh\left(\alpha * \frac{DLI}{Pmax}\right) + R(DLI) \tag{Eq. 3}$$

Where $GPg_{(DLI)}$ is the rate of gross photosynthesis measured at a given quantum
irradiance, DLI; P_{max} is the maximal photosynthetic capacity; α is the is the photosynthetic
efficiency; and $R_{(DLI)}$ is the light dependent variation of net coral respiration. As indicated
previously, we used here the direct values of R_L and R_D measured for each coral
phenotype, instead a single and general function for *Orbicella faveolata*, maintaining the
same criteria: R_L was used for the period of the day at photosynthesis saturation (P_{max}), R_D
in darkness or very low light levels, and the average for the sub-saturating region,
according to each acclimatory condition.

4. Photosynthetic output as a function of depth

To estimate the photosynthetic output $\Delta Pg(z)$ at a given z, we needed to integrate daily
photosynthesis at z (modified from (*127*)):

$$GPg_{(z)} = GPmax \int_0^N \tanh \frac{\alpha_{(z)} DLI_{(t)(z)}}{GPmax}\, dt \tag{Eq. 4}$$

5. Photosynthetic usable energy for coral maintenance

We tracked the relationship between photosynthetic and maintenance at depth and
incorporated the energy cost of photoacclimation *PUES(z)* as the cost of protein turnover

rate at depth. Energetic cost of 1 turnover (0 if no energy cost is involved in protein
turnover) (37, 54):

740
$$ECMpt_{(z)} = Protein\ turnover_{(z)} * Energetic\ cost \qquad \text{(Eq. 5)}$$

741 Photosynthetic usable energy supply for coral at depth z was expressed by integrating
742 equations 4 and 5:

743
$$PUES_{(z)} = DPg_{(z)} * (1 - ECMpt_{(z)}) \qquad \text{(Eq. 6)}$$

744 <u>Function-valued phenotyping included in the model, where *z* is depth and DLI daily light</u>
745 <u>integral:</u>

746 *1. Photosynthetic parameters as a function of DLI(z):*
747   a. Gross maximum photosynthetic rate *GPmax* = 15 µmol O_2 m^{-2} s^{-1}
748   b. Saturation irradiance *Ek* as a function of *DLI(z)* (mol quanta m^{-2} day^{-1})

749
$$Ek_{(z,DLI)} = 9.50DLI_{(z)} + 300.32$$

750   c. Photosynthetic efficiency α as a function of *DLI(z)*

751
$$\alpha_{(z,DLI)} = -0.0005DLI_{(z)} + 0.0374$$

752 *2. Protein turnover rate as a function of DLI(z):*

753   a. PSII-half-time (hours) as a function of *DLI* (mol quanta m^{-2} day^{-1}):

754
$$PSII1/2_{(z,DLI)} = \left[27.8374DLI_{(z)}^{-0.54723}\right]$$

755   b. Relative maximum photochemical efficiency (*Fv/Fm*):

756
$$FvFm_{DLI} = -0.0044DLI + 0.698$$

757
$$Rel.FvFm_{(DLI)} = \left[\frac{FvFm_{(DLI)}}{maxFvFm}\right]$$

758   c. Repair potential (dimensionless):

759
$$Repair\ potential_{(DLI)} = \left[\frac{12\ hours}{PSII1/2_{(DLI)}}\right]$$

760   d. Protein turnover rate:

762
$$Protein\ turnover_{(z)} = Repair\ potential_{(z)} * Rel.FvFm_{(z)}$$

References

1. R. Iglesias-Prieto, V. H. Beltrán, T. C. LaJeunesse, H. Reyes-Bonilla, P. E. Thomé, Different algal symbionts explain the vertical distribution of dominant reef corals in the eastern Pacific. *Proceedings. Biological sciences / The Royal Society* **271**, 1757–1763 (2004).

2. G. S. E. Chow, Y. K. S. Chan, S. S. Jain, D. Huang, Light limitation selects for depth generalists in urbanised reef coral communities. *Marine Environmental Research* **147**, 101–112 (2019).

3. T. E. Roberts, S. A. Keith, C. Rahbek, T. C. L. Bridge, M. J. Caley, A. H. Baird, Testing biodiversity theory using species richness of reef-building corals across a depth gradient. *Biol Lett* **15**, 20190493 (2019).

4. R. Tamir, G. Eyal, N. Kramer, J. H. Laverick, Y. Loya, Light environment drives the shallow-to-mesophotic coral community transition. *Ecosphere* **10**, e02839 (2019).

5. T. López-Londoño, K. Gómez-Campo, X. Hernández-Pech, S. Enríquez, R. Iglesias-Prieto, Photosynthetic usable energy explains vertical patterns of biodiversity in zooxanthellate corals. *Sci Rep* **12**, 20821 (2022).

6. G. Pérez-Rosales, M. Pichon, H. Rouzé, S. Villéger, G. Torda, P. Bongaerts, J. Carlot, U. T. P. Consortium, V. Parravicini, L. Hédouin, Mesophotic coral ecosystems of French Polynesia are hotspots of alpha and beta generic diversity for scleractinian assemblages. *Diversity and Distributions* **28**, 1391–1403 (2022).

7. T. C. LaJeunesse, J. E. Parkinson, P. W. Gabrielson, H. J. Jeong, J. D. Reimer, C. R. Voolstra, S. R. Santos, Systematic Revision of Symbiodiniaceae Highlights the Antiquity and Diversity of Coral Endosymbionts. *Current Biology* **28**, 2570-2580.e6 (2018).

8. L. Muscatine, L. R. Mccloskey, R. E. Marian, Estimating the daily contribution of carbon from zooxanthellae to coral animal respiration. *Limnology and Oceanography* **26**, 601–611 (1981).

9. P. G. Falkowski, Z. Dubinsky, L. Muscatine, J. W. Porter, Light and the Bioenergetics of a Symbiotic Coral. *BioScience* **34**, 705–709 (1984).

10. J. P. Gattuso, D. Allemand, M. Frankignoulle, Photosynthesis and calcification at cellular, organismal and community levels in coral reefs: A review on interactions and control by carbonate chemistry. *American Zoologist* **39**, 160–183 (1999).

11. S. Enríquez, E. R. Méndez, R. Iglesias-Prieto, S. Enrı, E. R. Me, Multiple scattering on coral skeletons enhances light absorption by symbiotic algae. *Limnology and Oceanography* **50**, 1025–1032 (2005).

12. J. W. Porter, L. Muscatine, Z. Dubinsky, P. G. Falkowski, Primary Production and Photoadaptation in Light- and Shade-Adapted Colonies of the Symbiotic Coral, Stylophora pistillata. *Proceedings of the Royal Society of London. Series B, Biological Sciences* **222**, 161–180 (1984).

803   13.   R. K. Trench, Microalgal-Invertebrate symbioses : a Review. *Endocytobiosis & Cell*
804   *Research* **9**, 135–175 (1993).

805   14.   R. Iglesias-Prieto, R. K. Trench, Acclimation and adaptation to irradiance in symbiotic
806   dinoflagellates. II. Response of chlorophyll-protein complexes to different photon-flux
807   densities. *Marine Biology* **130**, 23–33 (1997).

808   15.   W. K. Fitt, F. K. McFarland, M. E. Warner, G. C. Chilcoat, Seasonal patterns of tissue
809   biomass and densities of symbiotic dinoflagellates in reef corals and relation to coral
810   bleaching. *Limnology and Oceanography* **45**, 677–685 (2000).

811   16.   M. Y. Gorbunov, Z. S. Kolber, M. P. Lesser, P. G. Falkowski, Photosynthesis and
812   photoprotection in symbiotic corals. *Association for the Sciences of Limnology and*
813   *Oceanography* **46**, 75–85 (2001).

814   17.   E. A. Titlyanov, T. V. Titlyanova, K. Yamazato, R. van Woesik, Photo-acclimation
815   dynamics of the coral *Stylophora pistillata* to low and extremely low light. *Journal of*
816   *Experimental Marine Biology and Ecology* **263**, 211–225 (2001).

817   18.   K. R. N. Anthony, O. Hoegh-Guldberg, Kinetics of Photoacclimation in Corals. *Oecologia*
818   **134**, 23–31 (2003).

819   19.   P. R. Frade, P. Bongaerts, A. J. S. Winkelhagen, L. Tonk, R. P. M. Bak, In situ
820   photobiology of corals over large depth ranges: A multivariate analysis on the roles of
821   environment, host, and algal symbiont. *Limnology and Oceanography* **53**, 2711–2723
822   (2008).

823   20.   S. J. Hennige, D. J. Smith, R. Perkins, M. Consalvey, D. M. Paterson, D. J. Suggett,
824   Photoacclimation, growth and distribution of massive coral species in clear and turbid
825   waters. *Marine Ecology Progress Series* **369**, 77–88 (2008).

826   21.   P. A. Todd, Morphological plasticity in scleractinian corals. *Biological Reviews* **83**, 315–
827   337 (2008).

828   22.   M. P. Lesser, S. Marc, S. Michael, O. Michiko, R. D. Gates, G. Andrea,
829   Photoacclimatization by the coral Montastraea cavernosa in the mesophotic zone: Light,
830   food, and genetics. *Ecology* **91**, 990–1003 (2010).

831   23.   M. S. Roth, The engine of the reef: Photobiology of the coral-algal symbiosis. *Frontiers in*
832   *Microbiology* **5** (2014).

833   24.   S. Enríquez, E. R. Méndez, O. Hoegh-Guldberg, R. Iglesias-Prieto, Key functional role of
834   the optical properties of coral skeletons in coral ecology and evolution. *Proceedings of the*
835   *Royal Society B: Biological Sciences* **284**, 20161667 (2017).

836   25.   K. E. Lohr, E. F. Camp, U. Kuzhiumparambil, A. Lutz, W. Leggat, J. T. Patterson, D. J.
837   Suggett, Resolving coral photoacclimation dynamics through coupled photophysiological
838   and metabolomic profiling. *J Exp Biol* **222**, jeb195982 (2019).

839   26.   A. Malik, S. Einbinder, S. Martinez, D. Tchernov, S. Haviv, R. Almuly, P. Zaslansky, I.
840   Polishchuk, B. Pokroy, J. Stolarski, T. Mass, Molecular and skeletal fingerprints of

841 scleractinian coral biomineralization: From the sea surface to mesophotic depths. *Acta*
842 *Biomaterialia* **120**, 263–276 (2021).

843 27. M. O. Hoogenboom, S. R. Connolly, K. R. N. Anthony, Interactions Between
844 Morphological and Physiological Plasticity Optimize Energy Acquisition in Corals.
845 *Ecology* **89**, 1144–1154 (2008).

846 28. Y. X. Ow, P. A. Todd, Light-induced morphological plasticity in the scleractinian coral
847 Goniastrea pectinata and its functional significance. *Coral Reefs* **29**, 797–808 (2010).

848 29. N. Kramer, R. Tamir, O. Ben-Zvi, S. L. Jacques, Y. Loya, D. Wangpraseurt, Efficient light-
849 harvesting of mesophotic corals is facilitated by coral optical traits. *Functional Ecology* **36**,
850 406–418 (2022).

851 30. K. Gomez-Campo, R. Sanchez, I. Martinez-Rugerio, X. Yang, T. Maher, C. C. Osborne, S.
852 Enriquez, I. B. Baums, S. A. Mackenzie, R. Iglesias-Prieto, Phenotypic plasticity for
853 improved light harvesting, in tandem with methylome repatterning in reef-building corals.
854 *Molecular Ecology* **33**, e17246 (2024).

855 31. P. Hallock, Algal symbiosis: A mathematical analysis. *Mar. Biol.* **62**, 249–255 (1981).

856 32. M. O. Hoogenboom, K. R. N. Anthony, S. R. Connolly, Energetic cost of photoinhibition
857 in corals. *Marine Ecology Progress Series* **313**, 1–12 (2006).

858 33. D. H. Wright, Species-Energy Theory: An Extension of Species-Area Theory. *Oikos* **41**,
859 496–506 (1983).

860 34. R. De Visser, C. J. T. Spitters, T. J. Bouma, "Energy cost of protein turnover: theorectical
861 calculation and experimental estimation from regression of respiration on protein
862 concentration of full-grown leaves" in *Molecular, Biochemical and Physiological Aspects*
863 *of Plant Respiration* (1992), pp. 493–508.

864 35. T. J. Bouma, R. De Visser, J. H. J. A. Janssen, M. J. De Kock, P. H. Van Leeuwen, H.
865 Lambers, Respiratory energy requirements and rate of protein turnover in vivo determined
866 by the use of an inhibitor of protein synthesis and a probe to assess its effect. *Physiologia*
867 *Plantarum* **92**, 585–594 (1994).

868 36. A. Quigg, J. Beardall, Protein turnover in relation to maintenance metabolism at low
869 photon flux in two marine microalgae. *Plant, Cell and Environment* **26**, 693–703 (2003).

870 37. L. Li, C. J. Nelson, J. Trösch, I. Castleden, S. Huang, A. H. Millar, Protein degradation rate
871 in Arabidopsis thaliana leaf growth and development. *Plant Cell* **29**, 207–228 (2017).

872 38. G. Schuster, R. Timberg, I. Ohad, Turnover of thylakoid photosystem II proteins during
873 photoinhibition of Chlamydomonas reinhardtii. *European Journal of Biochemistry* **177**,
874 403–410 (1988).

875 39. E. M. Aro, S. McCaffery, J. M. Anderson, Photoinhibition and D1 Protein Degradation in
876 Peas Acclimated to Different Growth Irradiances. *Plant physiology* **103**, 835–843 (1993).

40. E.-M. Aro, I. Virgin, B. Andersson, Photoinhibition of Photosystem II. Inactivation, protein
damage and turnover. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1143**, 113–134
(1993).

41. S. Järvi, M. Suorsa, E. M. Aro, Photosystem II repair in plant chloroplasts--Regulation,
assisting proteins and shared components with photosystem II biogenesis. *Biochimica et*
biophysica acta **1847**, 900–909 (2015).

42. P. Chotewutmontri, A. Barkan, Dynamics of Chloroplast Translation during Chloroplast
Differentiation in Maize. *PLOS Genetics* **12**, e1006106 (2016).

43. L. Li, E. M. Aro, A. H. Millar, Mechanisms of Photodamage and Protein Turnover in
Photoinhibition. *Trends in Plant Science* **23**, 667–676 (2018).

44. K. Miyata, K. Noguchi, I. Terashima, Cost and benefit of the repair of photodamaged
photosystem II in spinach leaves: Roles of acclimation to growth light. *Photosynthesis*
Research **113**, 165–180 (2012).

45. M. Tikkanen, N. R. Mekala, E. M. Aro, Photosystem II photoinhibition-repair cycle
protects Photosystem i from irreversible damage. *Biochimica et Biophysica Acta -*
Bioenergetics **1837**, 210–215 (2014).

46. S. Järvi, J. Isojärvi, S. Kangasjärvi, J. Salojärvi, F. Mamedov, M. Suorsa, E.-M. Aro,
Photosystem II Repair and Plant Immunity: Lessons Learned from Arabidopsis Mutant
Lacking the THYLAKOID LUMEN PROTEIN 18.3. *Front. Plant Sci.* **7** (2016).

47. N. Murata, Y. Nishiyama, ATP is a driving force in the repair of photosystem II during
photoinhibition. *Plant, Cell & Environment* **41**, 285–299 (2018).

48. S. Takahashi, T. Nakamura, M. Sakamizu, R. Van Woesik, H. Yamasaki, Repair nachinery
of symbiotic photosynthesis as the primary target of heat stress for reef-building corals.
Plant and Cell Physiology **45**, 251–255 (2004).

49. R. Hill, S. Takahashi, Photosystem II recovery in the presence and absence of chloroplast
protein repair in the symbionts of corals exposed to bleaching conditions. *Coral Reefs* **33**,
1101–1111 (2014).

50. S. Roberty, D. Fransolet, P. Cardol, J.-C. Plumier, F. Franck, Imbalance between oxygen
photoreduction and antioxidant capacities in Symbiodinium cells exposed to combined heat
and high light stress. *Coral Reefs* **34**, 1063–1073 (2015).

51. K. R. N. Anthony, P. V. Ridd, A. R. Orpin, P. Larcombe, J. Lough, Temporal variation of
light availability in coastal benthic habitats: Effects of clouds, turbidity, and tides.
Limnology and Oceanography **49**, 2201–2211 (2004).

52. S. DiPerna, M. Hoogenboom, S. Noonan, K. Fabricius, Effects of variability in daily light
integrals on the photophysiology of the corals Pachyseris speciosa and Acropora millepora.
PLoS ONE **13**, e0203882 (2018).

53. P. G. Falkowski, J. A. Raven, *Aquatic Photosynthesis* (Princeton University Press, 2014).

54.   T. López-Londoño, C. T. Galindo-Martínez, K. Gómez-Campo, L. A. González-Guerrero,
915   S. Roitman, F. J. Pollock, V. Pizarro, M. López-Victoria, M. Medina, R. Iglesias-Prieto,
916   Physiological and ecological consequences of the water optical properties degradation on
917   reef corals. *Coral Reefs* **40**, 1243–1256 (2021).

918   55.   R. Hill, F. Bendall, Function of the Two Cytochrome Components in Chloroplasts: A
919   Working Hypothesis. *Nature* **186**, 136–137 (1960).

920   56.   J. F. Hill, Govindjee, The controversy over the minimum quantum requirement for oxygen
921   evolution. *Photosynth Res* **122**, 97–112 (2014).

922   57.   W. Skirving, S. Enríquez, J. D. Hedley, S. Dove, C. M. Eakin, R. A. B. Mason, J. L. D. L.
923   Cour, G. Liu, O. Hoegh-Guldberg, A. E. Strong, P. J. Mumby, R. Iglesias-Prieto, Remote
924   sensing of coral bleaching using temperature and light: Progress towards an operational
925   algorithm. *Remote Sensing* **10**, 18 (2018).

926   58.   L. Muscatine, P. G. ; Falkowski, J. W. ; Porter, Z. Dubinsky, Fate of photosynthetic fixed
927   carbon in light- and shade-adapted colonies of the symbiotic coral Stylophora pistillata.
928   *Proceedings of the Royal Society of London. Series B. Biological Sciences* **222**, 181–202
929   (1984).

930   59.   L. Muscatine, Symbiosis of hydra and algae—III. Extracellular products of the algae.
931   *Comparative Biochemistry and Physiology* **16**, 77–92 (1965).

932   60.   L. Muscatine, Glycerol excretion by symbiotic algae from corals and Tridacna and its
933   control by the host. *Science* **156**, 516–519 (1967).

934   61.   M. S. Burriesci, T. K. Raab, J. R. Pringle, Evidence that glucose is the major transferred
935   metabolite in dinoflagellate–cnidarian symbiosis. *Journal of Experimental Biology* **215**,
936   3467–3477 (2012).

937   62.   R. K. Trench, The Physiology and Biochemistry of Zooxanthellae Symbiotic with Marine
938   Coelenterates. II. Liberation of Fixed 14C by Zooxanthellae in vitro. *Proceedings of the
939   Royal Society of London. Series B, Biological Sciences* **177**, 237–250 (1971).

940   63.   R. K. Trench, The Physiology and Biochemistry of Zooxanthellae Symbiotic with Marine
941   Coelenterates. I. The Assimilation of Photosynthetic Products of Zooxanthellae by Two
942   Marine Coelenterates. *Proceedings of the Royal Society of London. Series B, Biological
943   Sciences* **177**, 225–235 (1971).

944   64.   P. Tremblay, R. Grover, J. F. Maguer, M. Hoogenboom, C. Ferrier-Pagès, Carbon
945   translocation from symbiont to host depends on irradiance and food availability in the
946   tropical coral Stylophora pistillata. *Coral Reefs* **33**, 1–13 (2014).

947   65.   S. Sathyendranath, T. Platt, Ž. Kovač, J. Dingle, T. Jackson, R. J. W. Brewin, P. Franks, E.
948   Marañón, G. Kulk, H. A. Bouman, Reconciling models of primary production and
949   photoacclimation [Invited]. *Applied Optics* **59**, C100 (2020).

950   66.   R. D. Gates, P. J. Edmunds, The physiological mechanisms of acclimatization in tropical
951   reef corals. *American Zoologist* **39**, 30–43 (1999).

952   67.   M. O. Hoogenboom, S. R. Connolly, K. R. N. Anthony, Effects of photoacclimation on the
953   light niche of corals: A process-based approach. *Marine Biology* **156**, 2493–2503 (2009).

954   68.   B. Demmig-Adams, W. W. Adams, The role of xanthophyll cycle carotenoids in the
955   protection of photosynthesis. *Trends in Plant Science* **1**, 21–26 (1996).

956   69.   B. E. Brown, I. Ambarsari, M. E. Warner, W. K. Fitt, R. P. Dunne, S. W. Gibb, D. G.
957   Cummings, Diurnal changes in photochemical efficiency and xanthophyll concentrations in
958   shallow water reef corals: Evidence for photoinhibition and photoprotection. *Coral Reefs*
959   **18**, 99–105 (1999).

960   70.   P. J. Ralph, R. Gademann, A. W. D. Larkum, U. Schreiber, In situ underwater
961   measurements of photosynthetic activity of coral zooxanthellae and other reef-dwelling
962   dinoflagellate endosymbionts. *Marine Ecology Progress Series* **180**, 139–147 (1999).

963   71.   E. Erickson, S. Wakao, K. K. Niyogi, Light stress and photoprotection in Chlamydomonas
964   reinhardtii. *Plant Journal*, 449–465 (2015).

965   72.   R. J. Jones, O. Hoegh-Guldberg, Diurnal changes in the photochemical efficiency of the
966   symbiotic dinoflagellates (Dinophyceae) of corals: photoprotection, photoinactivation and
967   the relationship to coral bleaching. *Plant, Cell & Environment* **24**, 89–99 (2001).

968   73.   M. P. Lesser, M. Y. Gorbunov, Diurnal and bathymetric changes in chlorophyll
969   fluorescence yields of reef corals measured in situ with a fast repetition rate fluorometer.
970   **212**, 69–77 (2001).

971   74.   G. Winters, Y. Loya, R. Röttgers, S. Beer, Photoinhibition in shallow-water colonies of the
972   coral Stylophora pistillata as measured in situ. *Limnology and Oceanography* **48**, 1388–
973   1393 (2003).

974   75.   O. Levy, Z. Dubinsky, K. Schneider, Y. Achituv, D. Zakai, M. Y. Gorbunov, Diurnal
975   hysteresis in coral photosynthesis. *Marine Ecology Progress Series* **268**, 105–117 (2004).

976   76.   A. Rodríguez-Román, X. Hernández-Pech, P. E. Thomé, S. Enríquez, R. Iglesias-Prieto, X.
977   Herna, P. E. Thome, S. Enrı, R. Iglesias-Prieto, Photosynthesis and light utilization in the
978   Caribbean coral Montastraea faveolata recovering from a bleaching event. *Limnology and*
979   *Oceanography* **51**, 2702–2710 (2006).

980   77.   T. Scheufen, R. Iglesias-Prieto, S. Enríquez, Changes in the number of symbionts and
981   Symbiodinium cell pigmentation modulate differentially coral light absorption and
982   photosynthetic performance. *Frontiers in Marine Science* **4**, 309 (2017).

983   78.   T. Scheufen, W. E. Krämer, R. Iglesias-Prieto, S. Enríquez, Seasonal variation modulates
984   coral sensibility to heat-stress and explains annual changes in coral productivity. *Scientific*
985   *Reports* **7**, 4937 (2017).

986   79.   N. Kramer, R. Tamir, G. Eyal, Y. Loya, Coral Morphology Portrays the Spatial
987   Distribution and Population Size-Structure Along a 5–100 m Depth Gradient. *Front. Mar.*
988   *Sci.* **7** (2020).

989   80.   J. P. Krall, G. E. Edwards, Relationship between photosystem II activity and CO2 fixation
990   in leaves. *Physiologia Plantarum* **86**, 180–187 (1992).

81.   M. J. Fryer, J. R. Andrews, K. Oxborough, D. A. Blowers, N. R. Baker, Relationship
between CO2 Assimilation, Photosynthetic Electron Transport, and Active O2 Metabolism
in Leaves of Maize in the Field during Periods of Low Temperature1. *Plant Physiology*
116, 571–580 (1998).

82.   O. Hoegh-Guldberg, R. Jones, Photoinhibition and photoprotection in symbiotic
dinoflagellates from reef-building corals. *Mar. Ecol. Prog. Ser.* **183**, 73–86 (1999).

83.   M. Lesser, Depth-dependent photoacclimatization to solar ultraviolet radiation in the
Caribbean coral Montastraea faveolata. *Mar. Ecol. Prog. Ser.* **192**, 137–151 (2000).

84.   J. A. Raven, The cost of photoinhibition. *Physiologia Plantarum* **142**, 87–104 (2011).

85.   X.-P. Yi, H.-S. Yao, D.-Y. Fan, X.-G. Zhu, P. Losciale, Y.-L. Zhang, W.-F. Zhang, W. S.
Chow, The energy cost of repairing photoinactivated photosystem II: an experimental
determination in cotton leaf discs. *New Phytologist* **235**, 446–456 (2022).

86.   S. Gunell, T. Lempiäinen, E. Rintamäki, E.-M. Aro, M. Tikkanen, Enhanced function of
non-photoinhibited photosystem II complexes upon PSII photoinhibition. *Biochimica et*
Biophysica Acta (BBA) - Bioenergetics **1864**, 148978 (2023).

87.   C. Langdon, On the causes of interspecific differences in the growth-irradiance relationship
for phytoplankton. II. A general review. *Journal of Plankton Research* **10**, 1291–1312
(1988).

88.   N. Murata, S. Takahashi, Y. Nishiyama, S. I. Allakhverdiev, Photoinhibition of
photosystem II under environmental stress. *Biochimica et Biophysica Acta - Bioenergetics*
1767, 414–421 (2007).

89.   S. I. Allakhverdiev, Y. Nishiyama, S. Takahashi, S. Miyairi, I. Suzuki, N. Murata,
Systematic Analysis of the Relation of Electron Transport and ATP Synthesis to the
Photodamage and Repair of Photosystem II in Synechocystis. *Plant Physiology* **137**, 263–
273 (2005).

90.   R. Iglesias-prieto, R. K. Trench, Acclimation and adaptation to irradiance in symbiotic
dinoflagellates. 1.Responses of the photosynthetic unit to changes in photon flux density.
Marine Ecology Progress Series **113**, 163–176 (1994).

91.   A. Trebst, Function of β-Carotene and Tocopherol in Photosystem II. *Zeitschrift für*
Naturforschung C **58**, 609–620 (2003).

92.   A. Krieger-Liszkay, Singlet oxygen production in photosynthesis. *J Exp Bot* **56**, 337–346
(2005).

93.   M. E. Warner, W. K. Fitt, G. W. Schmidt, Damage to photosystem II in symbiotic
dinoflagellates: a determinant of coral bleaching. *Proceedings of the National Academy of*
Sciences of the United States of America **96**, 8007–12 (1999).

94.   S. Takahashi, S. Whitney, S. Itoh, T. Maruyama, M. Badger, Heat stress causes inhibition
of the de novo synthesis of antenna proteins and photobleaching in cultured Symbiodinium.
Proceedings of the National Academy of Sciences of the United States of America **105**,
4203–8 (2008).

95.   S. Takahashi, M. Yoshioka-Nishimura, D. Nanba, M. R. Badger, Thermal Acclimation of
the Symbiotic Alga Symbiodinium spp. Alleviates Photobleaching under Heat Stress.
PLANT PHYSIOLOGY **161**, 477–485 (2013).

96.   R. Hill, C. M. Brown, K. DeZeeuw, D. a Campbell, P. J. Ralph, Increased rate of D1 repair
in coral symbionts during bleaching is insufficient to counter accelerated photo-
inactivation. *Limnology and Oceanography* **56**, 139–146 (2011).

97.   S. Takahashi, S. E. Milward, W. Yamori, J. R. Evans, W. Hillier, M. R. Badger, The solar
action spectrum of photosystem II damage. *Plant Physiology* **153**, 988–993 (2010).

98.   J. M. Pandolfi, C. E. Lovelock, A. F. Budd, Character Release Following Extinction in a
Caribbean Reef Coral Species Complex. *Evolution* **56**, 479–501 (2002).

99.   C. Prada, M. Gómez-Corrales, A. González, B. Kamel, M. Medina, N. Knowlton, D.
Levitan, Speciation across depth gradients in reef corals. In Review [Preprint] (2024).
https://doi.org/10.21203/rs.3.rs-4202515/v1.

100. H. E. Aichelman, B. E. Benson, K. Gomez-Campo, M. I. Martinez-Rugerio, J. E. Fifer, L.
Tsang, A. M. Hughes, C. B. Bove, O. C. Nieves, A. M. Pereslete, D. Stanizzi, N. G.
Kriefall, J. H. Baumann, J. P. Rippe, P. Gondola, K. D. Castillo, S. W. Davies, Cryptic
coral diversity is associated with symbioses, physiology, and response to thermal
challenge. *Science Advances* **11**, eadr5237 (2025).

101. D. Iluz, I. Alexandrovich, Z. Dubinsky, The enhancement of photosynthesis by fluctuating
light. *Artifical Photosynthesis* (2012).

102. J. E. Faust, V. Holcombe, N. C. Rajapakse, D. R. Layne, The effect of daily light integral
on bedding plant growth and flowering. *HortScience* **40**, 645–649 (2005).

103. S. W. Jeffrey, G. F. Humphrey, New spectrophotometric equations for determining
chlorophylls a, b, c1 and c2 in higher plants, algae and natural phytoplankton. *Biochemie
und Physiologie der Pflanzen* **167**, 191–194 (1975).

104. R. M. Vásquez-Elizondo, L. Legaria-Moreno, M. Á. Pérez-Castro, W. E. Krämer, T.
Scheufen, R. Iglesias-Prieto, S. Enríquez, Absorptance determinations on multicellular
tissues. *Photosynthesis Research* **132**, 311–324 (2017).

105. S. Enríquez, K. Sand-Jensen, Variation in Light Absorption Properties of *Mentha aquatica*
L. as a Function of Leaf Form: Implications for Plant Growth. *International Journal of
Plant Sciences* **164**, 125–136 (2003).

106. N. M. Cayabyab, S. Enríquez, Leaf photoacclimatory responses of the tropical seagrass
Thalassia testudinum under mesocosm conditions: a mechanistic scaling-up study. *New
Phytologist* **176**, 108–123 (2007).

107. L. A. González-Guerrero, R. M. Vásquez-Elizondo, T. López-Londoño, G. Hernán, R.
Iglesias-Prieto, S. Enríquez, Validation of parameters and protocols derived from
chlorophyll a fluorescence commonly utilised in marine ecophysiological studies.
Functional Plant Biol., doi: 10.1071/FP21101 (2021).

108. W. C. Dennison, R. S. Alberte, Photosynthetic Responses of Zostera marina L. (Eelgrass)
to in situ Manipulations of Light Intensity. *Oecologia* **55**, 137–144 (1982).

109. M. K. DeSalvo, S. Sunagawa, P. L. Fisher, C. R. Voolstra, R. Iglesias-Prieto, M. Medina,
Coral host transcriptomic states are correlated with Symbiodinium genotypes. *Molecular*
Ecology **19**, 1174–1186 (2010).

110. D. W. Kemp, D. J. Thornhill, R. D. Rotjan, R. Iglesias-Prieto, W. K. Fitt, G. W. Schmidt,
Spatially distinct and regionally endemic Symbiodinium assemblages in the threatened
Caribbean reef-building coral Orbicella faveolata. *Coral Reefs* **34**, 535–547 (2015).

111. A. M. Lewis, A. N. Chan, T. C. LaJeunesse, New Species of Closely Related
Endosymbiotic Dinoflagellates in the Greater Caribbean have Niches Corresponding to
Host Coral Phylogeny. *Journal of Eukaryotic Microbiology* **66**, 469–482 (2019).

112. T. C. LaJeunesse, W. K. W. Loh, R. Van Woesik, O. Hoegh-Guldberg, G. W. Schmidt, W.
K. Fitt, Low symbiont diversity in southern Great Barrier Reef corals, relative to those of
the Caribbean. *Limnology and Oceanography* **48**, 2046–2054 (2003).

113. T. C. Lajeunesse, R. K. Trench, Biogeography of two species of Symbiodinium
(Freudenthal) inhabiting the intertidal sea anemone Anthopleura elegantissima (Brandt).
https://doi.org/10.2307/1542872 **199**, 126–134 (2016).

114. R. Zardoya, E. Costas, V. López-Rodas, A. Garrido-Pertierra, J. M. Bautista, Revised
dinoflagellate phylogeny inferred from molecular analysis of large-subunit ribosomal RNA
gene sequences. *Journal of Molecular Evolution* **41**, 637–645 (1995).

115. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv arXiv:1303.3997 [Preprint] (2013). https://doi.org/10.48550/arXiv.1303.3997.

116. V. Avila-Magaña, B. Kamel, M. DeSalvo, K. Gómez-Campo, S. Enríquez, H. Kitano, R. V.
Rohlfs, R. Iglesias-Prieto, M. Medina, Elucidating gene expression adaptation of
phylogenetically divergent coral holobionts under heat stress. *Nat Commun* **12**, 5731
(2021).

117. E. F. Camp, T. Kahlke, B. Signal, C. A. Oakley, A. Lutz, S. K. Davy, D. J. Suggett, W. P.
Leggat, Proteome metabolome and transcriptome data for three Symbiodiniaceae under
ambient and heat stress conditions. *Sci Data* **9**, 153 (2022).

118. R. A. Levin, V. H. Beltran, R. Hill, S. Kjelleberg, D. McDougald, P. D. Steinberg, M. J. H.
van Oppen, Sex, Scavengers, and Chaperones: Transcriptome Secrets of Divergent
Symbiodinium Thermal Tolerances. *Mol Biol Evol* **33**, 2201–2215 (2016).

119. C. M. Brown, D. A. Campbell, J. E. Lawrence, Resource dynamics during infection of
Micromonas pusilla by virus MpV-Sp1. *Environmental Microbiology* **9**, 2720–2727
(2007).

120. C. Six, Z. V. Finkel, A. J. Irwin, D. A. Campbell, Light Variability Illuminates Niche-
Partitioning among Marine Picocyanobacteria. *PLoS ONE* **2**, e1341 (2007).

121. P. Ewels, M. Magnusson, S. Lundin, M. Käller, MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

122. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

123. N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol* **34**, 525–527 (2016).

124. L. Fu, B. Niu, Z. Zhu, S. Wu, W. Li, CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

125. C. Camacho, G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, T. L. Madden,
BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

126. J. Kirk, *Light and Photosynthesis* (Cambridge University Press, New York, NY, 2011;
https://www.cambridge.org/core/books/light-and-photosynthesis-in-aquatic-
ecosystems/C19B28AE07B1CDEBDA5593194DE4E304).

127. A. D. Jassby, T. Platt, Mathematical formulation of the relationship between photosynthesis
and light for phytoplankton. *Limnology and Oceanography* **21**, 540–547 (1976).

1120

1121
Acknowledgments
We thank UNAM and ICMyL for staff support, facility assistance, SAMMO for data, S
Enriquez for laboratory space and mesocosm facility. V Avila-Magana, CT Galindo-
Martinez, NS Locatelli, CC Osborne for assistance in laboratory work. T López-Londono
for assistance in modeling. VA Gómez-Chavez for Figure S1. Collection permit No.
PPF/DGOPA-020/2018 and CITES permit No. MX96989, MX102782 to RIP and SE.
1128
Funding:
This project has received funding from The Pennsylvania State University Start-up (RIP).
Authors acknowledge support by the HIFMB, a collaboration between the Alfred-
Wegener-Institute, Helmholtz-Center for Polar and Marine Research, and the Carl-von-
Ossietzky University Oldenburg, initially funded by the Ministry of Science and Culture
of Lower Saxony (MWK).
1135
Author contributions:
Conceptualization: RIP, KGC.
Data curation: KGC.
Formal analysis: KGC and RIP.
Funding acquisition: RIP, SE, TCL, IBB.
Investigation and methodology: KGC performed the experiments analyze and integrated
data. IMR contributed to fieldwork, phenotypic descriptions and light systems. MAGR
engineered the light system. RGP to transcriptomic computational work and
interpretations. MGS to translational laboratory work and interpretations. KET to algal
identification laboratory work and interpretations. TLC and IBB to analysis and
interpretations.
Project administration: RIP.
Resources: RIP, SE, TCL, IBB, MGSM.
Software: MAGR.
Supervision: RIP.
Validation: KGC, RIP.
Visualization: KGC, RIP.
Writing - original draft: KGC, RIP.
Writing - review & editing: KGC, IMR, MAGR, RGP, MGSM, KET, TCL, IBB, SE, RIP.
1155
1156   **Competing interests:** Authors declare that they have no competing interests.
1157
1158   **Data and materials availability:** Data is available in the main text or the supplementary
1159   materials. All detailed data, custom codes used in the analyses will be available for
1160   purposes of reproducing or extending the analyses in GitHub. All data generated by this
1161   study will be deposited to the National Center for Biotechnology Information (NCBI).
1162

1163
1164   **Figures and Tables**
1165
1166

1167
Fig. 1. Phenotyping after acclimation to light conditions. (**A**) The general equation of
light attenuation in seawater ($DLI_{(z)} = DLI_{(o)} \times e^{(-Kd \times z)}$; Eq. 1) maps the four
experimental DLI conditions to their corresponding depths (5–35 m, assuming K_d
= 0.07 m^{-1}), illustrated alongside a depth gradient. (**B**) Maximum photochemical
efficiency (F_v/F_m at dusk) as a function of DLI after 4+ weeks of acclimation
illustrating a linear trend. Adjustments in photoprotection shown as (**C**) diurnal
changes in photochemical efficiency (maximum F_v/F_m at dawn/dusk, and effective
$\Delta F/F_m{}'$ during daylight) and (**D**) non-photochemical quenching (*NPQ*) measured
hourly across light conditions. Symbols represent mean ± SE (n = 16). Shaded
areas show bootstrapped 95% confidence intervals.

Fig. 2. Optical reorganization and metabolic rates for light harvesting across light conditions. (**A**) Estimated absorbance (*De*) from reflectance measurements indicating reduced absorption under high light (n = 16) and (photo) qualitative changes in tissue pigmentation. (**B**) Pigment-specific absorption coefficient (a*$_{Chla}$) showing the non-linear increase in absorption efficiency in the coral tissue as pigment content declines, consistent with the package effect. (**C**) Photosynthetic efficiency (α), (**D**) maximum photosynthetic rate (P_{max} normalized to host area, and P_{sym} normalized to symbiont cell) showing no significant changes in P_{max} across light regimes (n = 8). (**E**) Dark respiration rates (R_D) (n = 8), and (**F**) maximum photochemical efficiency of PSII (F_v/F_m) as a function of the minimum quantum requirement for photosynthesis (MQR, Φ^{-1}; mol photons mol^{-1} O$_2$). Lower MQR values indicate higher photosynthetic efficiency or fewer photons required per molecule of O$_2$ evolved. DLI-4 corals achieved MQR values approaching the theoretical minimum (~8 quanta O$_2^{-1}$), indicating near-maximal photosynthetic efficiency and delineating the lower boundary of photoacclimatory capacity for the symbiosis. With increasing irradiance (toward DLI-32), both F_v/F_m and quantum efficiency declined, reflecting the increasing energetic cost of sustaining photosynthesis under high light. Shaded areas show bootstrapped 95% confidence intervals; symbols are means ± SE.

Fig. 3. Diurnal PSII activity and PSII half-life. (A) PSII photochemical efficiency measured hourly across a 12 h photoperiod in the presence (+CAP; empty circles) and absence (−CAP; filled circles) of the protein synthesis inhibitor chloramphenicol. Blocking *de novo* protein synthesis revealed net PSII damage accumulation, while control samples showed oscillations with characteristic daily photodamage and repair cycles. **(B)** Relative photochemical efficiency (+CAP/-CAP) decay showing proportional rate of photodamage with increased light intensity, reflecting the rate of PSII inactivation in the absence of repair. Dashed lines illustrate the time of day when 50% of PSII reaction centers are photo-inactivated. In DLI-4 PSII $t_{1/2}$ exceeded the 12 h photoperiod. Curves show four-parameter logistic fits (mean ± 95% CI).

Fig. 4. PSII regulation in zooxanthellate corals across light conditions. (A) PSII large pigment–protein complex with its core reaction center D1 (encoded by *psbA* gene). D1 provides key ligands to the oxygen-evolving complex, harbors the QB plastoquinone binding site, and is the primary target of photodamage. (**B**) Power law decay of PSII half-life ($t_{1/2}$, h) with increasing DLI. (**C**) PSII turnover rate (day^{-1}) accelerated significantly under higher DLI. (**D**) Quantitative immunoblot of the D1 protein, showing patterns across light conditions. Lanes include a negative control (lane1, 0.00 pmol), quantification standards (lanes2–6, 0.10-2.00 pmol), and samples loaded by total protein (lanes7–10, total protein^{-1}) and Chl*a* content (11–14, mg Chl*a*$^{-1}$). Quantification of D1 protein normalized to total protein (**E**; pmol D1 protein μg total protein^{-1}) and Chl*a* (**F**; pmol D1 protein mg Chl*a*$^{-1}$), showing a similar power law decay with light exposure (n = 3 immunoblots). (**G**) Transcriptional abundance (transcripts per million; TPM) of PSII complex genes (*psbA, psbB, psbC, psbD*). Transcript levels showed no systematic relationship with DLI, in contrast to the exponential decline in D1 protein abundance shown in panels E and F, indicating post-transcriptional regulation. (**H**) PSII half-life ($t_{1/2}$, h) when acclimated corals (DLI) were changed to an alternative light condition (DLI-destination). Lines show a power law fit to experimental PAR (black line and colored circles; this study) and natural sunlight measurements (blue line and grey circles; data from (*5*)). Shaded areas show bootstrapped 95% confidence intervals; symbols are means ± SE. (**I**) Photodamage accumulation after the change in light condition. Results are shown as the relative change in F_v/F_m (*relative-F_v/F_m* = F_v/F_m-destination / F_v/F_m-acclimation) when coral fragments were exposed to *EEE* (*EEE* = DLI-destination - DLI-acclimation); the linear prediction is consistent with prior findings (*57*).

1239
1240

1241
Fig. 5. Bio-optical model of *photosynthetic usable energy supply* (*PUES*). (**A**) Modeled
profiles of *PUES* across depth for a range of vertical attenuation coefficients of
downwelling irradiance (K_d = 0.04–0.40 m⁻¹), representing varying water column
optical properties. *PUES* represents the photosynthetic energy available to the
coral host after subtracting the depth-dependent energetic cost of PSII repair. Each
curve represents a distinct K_d value, color-coded from clear to turbid conditions.
(**B**) Optimal depth (*Depth_opt*) at which *PUES* is maximized, as a function of Kd.
This bio-optical state represents the depth at which the balance between
photosynthetic output and PSII repair cost is optimized, maximizing the energy
potentially available for host metabolic demands.
1252

Table 1. Table of terms. Descriptors used for phenotyping coral holobionts in the different levels of organization.

Units	Parameters and notations	Units
PSII	Maximum photochemical efficiency (F_v/F_m)	Dimensionless
	Effective photochemical efficiency ($\Delta F/F_m'$)	Dimensionless
	Maximum excitation pressure over PSII (Q_m)	Dimensionless
	PSII half-life ($PSII_{1/2}$)	Hours
	PSII D1 protein abundance per protein	pmol D1 µg total protein^{-1}
	PSII D1 protein abundance per Chl$a$	pmol D1 µg Chla^{-1}
Structural and optical traits	Chl$a$ density per area of host tissue	mg Chl$a$ m^{-2}
	Zoox. per area of host tissue	Cells cm^2
	Chl$a$ per zoox. cell (Ci)	Pg Chl$a$ sym^{-1}
	Absorptance PAR (A_{PAR})	Dimensionless
	Absorptance 675 (A_{675})	Dimensionless
	Absorbance PAR (De_{PAR})	Dimensionless
	Absorbance 675 (De_{675})	Dimensionless
	Chl$a$ specific absorption coefficient (a^*_{Chla})	m^2 mg Chla^{-1}
	Symbiont specific absorption coefficient (a^*_{Sym})	m^2 sym^{-1}
Metabolic rates	Photosynthetic efficiency (α)	mol O$_2$ mol incident quanta^{-1}
	Quantum yield of photosynthesis (Φ_{O2})	mol O$_2$ mol absorbed quanta^{-1}
	Minimum Quantum Requirement ($1/\Phi_{O2}$)	photons absorbed per O$_2$ evolved
	Maximum photosynthetic rate (P_{max})	µmol O$_2$ m^{-2} s^{-1}
	Saturation irradiance (E_k)	µmol quanta m^{-2} s^{-1}
	Compensation irradiance (E_c)	µmol quanta m^{-2} s^{-1}
	Dark respiration (R_D)	µmol O$_2$ m^{-2} s^{-1}
	Light-enhanced respiration (R_l)	µmol O$_2$ m^{-2} s^{-1}
	Photosynthetic rate per zoox. cell ($Psym$)	µmol O$_2$ m^{-2} s^{-1}