Title

Maintenance cost of photosynthesis sets key ecological constraints on zooxanthellate corals

Authors
Kelly Gomez-Campo[1]*, M. Isabel Martinez-Rugerio[1], Miguel Ángel Gómez Reali[2], Raúl A. González-Pech[1,3], Michel Geovanni Santiago-Martínez[4], Kira E. Turnham[1], Todd C. LaJeunesse[1], Iliana B. Baums[1], Susana Enríquez[2], Roberto Iglesias-Prieto[1]*

Affiliations
[1]Department of Biology, The Pennsylvania State University, University Park, PA 16802, USA.
[2]Unidad Académica de Sistemas Arrecifales Puerto Morelos, Instituto de Ciencias del Mar y Limnología, Universidad Nacional Autónoma de México, 77580, México.
[3]Department of Biology, Texas State University, San Marcos, TX 78666, USA.
[4]Department of Molecular and Cell Biology, The University of Connecticut, Storrs, Connecticut, 06269, USA.

Corresponding authors: KGC kelly.gomezcampo@hifmb.de and RIP rzi3@psu.edu

Abstract

Ecological models using light limitation to explain coral depth distribution have largely disregarded the energetic cost of sustaining photosynthetic activity. Here, we quantified photosystem II (PSII) turnover across a depth-simulated light gradient in a zooxanthellate coral, measuring PSII half-life, D1 protein abundance, and PSII-complex gene expression. Maximum photosynthetic capacity remained stable across irradiance levels while respiration rose and PSII turnover accelerated as a power law, imposing increasing ATP demand at the shallowest depths. Declining D1 protein abundance alongside stable transcript levels demonstrated that this escalating maintenance cost operates through post-transcriptional regulation. Consequently, a decreasing fraction of photosynthetic usable energy is available for translocation to the coral host at high irradiance, as the energy required for PSII repair increases. Integrating these physiological constraints into a bio-optical model revealed that the balance between photosynthetic capacity and its maintenance cost defines an optimal depth, the Photosynthetic Usable Energy Supply (*PUES*) maximum, where host energetic returns are maximized. This framework provides a mechanistic basis for understanding depth distributions in symbiotic corals and extends as a predictive tool for any photosynthetic organism operating under variable irradiance, including forecasting how environmental degradation contracts viable depth ranges.

MAIN TEXT

Introduction

Light supplies the energy that sustains coral reef ecosystems, but its role in shaping coral distribution cannot be explained by light availability alone. Variation in light quantity and spectral composition set physiological limits that structure community composition, species richness, and colony morphology across depth gradients (*1–6*). These ecological patterns emerge from fundamental biophysical constraints on how efficiently light energy is captured and converted into metabolic output. Corals are modular, sessile, mixotrophic organisms that host intracellular dinoflagellates in the family Symbiodiniaceae (*7*). Endosymbionts translocate a large fraction of their photosynthetically fixed carbon to the host, fueling basal metabolism and calcification (*8–10*). Light harvesting by the coral holobiont is determined by the emergent optical properties of symbionts, host tissue, and skeleton (*11*). Consequently, efficient photosynthesis requires coordinated adjustment across multiple levels of biological organization, from symbiont cell pigmentation and density, to host tissue optics, skeletal architecture, and whole colony morphology (*1, 12–26*). These traits are integrated through photoacclimatory processes that optimize energy conversion under different light environments (*24, 27–30*). Yet, the physiological limits through which photosynthesis constrains coral ecological niches across depth remain unresolved (*5, 9, 31, 32*).

Depth gradients impose continuous selective pressures on coral–symbiont associations as sunlight simultaneously determines energy acquisition and physiological demands. If coral distribution were determined by solar energy alone, productivity and species richness would be expected to peak in shallow reefs and decline monotonically with depth as light attenuates. This pattern would reflect reduced growth rates and increased extinction risk under lower energy supply (*33*). However, for symbiotic corals, light availability by itself often fails to predict vertical patterns of biodiversity and performance (*3*). Instead, the amount of light that can be converted into biologically usable energy for the host better explains species richness and depth zonation in zooxanthellate corals (*5*). This distinction has relevant ecological implications. At intermediate depths, many corals maximize their net usable energy, expanding niche space and promoting greater species coexistence relative to both shallow and deep extremes. Under this framework, depth zonation cannot be simply considered a response to light attenuation with depth, but the outcome of selection on traits that determine how efficiently corals convert depth-specific available light into net energetic return. This is achieved through coordinated optimization of coral light capture, physiological adjustments of symbionts and host, and colony morphology. While lower depth limits are attributed to light limitation, upper depth limits are hypothesized to be constrained by the increasing energetic demands required to maintain performance under high irradiance.

Among the physiological processes that impose these energetic demands, protein damage and repair rank among the most energetically expensive cellular processes across living systems (*34–37*). However, their contribution to whole-organism energy budgets remains difficult to quantify. In photosynthetic organisms, this challenge is exemplified by the photosystem II (PSII) reaction center D1 protein, which undergoes rapid, light-dependent cycles of damage, degradation, and resynthesis (*38–43*). Replacement of photodamaged PSII reaction centers requires substantial ATP investment to support D1 synthesis, assembly, and regulation (*44–47*). In coral-associated Symbiodiniaceae, accelerated D1

turnover has been documented under both light- and thermal stress, consistent with a
central role in sustaining photosynthetic function under stress conditions and maintaining
the stability of the mutualism (*48–50*). Thus, empirical data on the energetic cost of
sustaining this repair activity, and its consequences for energy partitioning within the
symbiosis, are critical to determine energy utilization and delineate energetic boundaries
in depth gradients.

We hypothesize that changes in PSII damage and repair across a light gradient affect the
net energy that can be translocated from the symbiont to the coral host. Enhanced repair in
high light environments sustains photosynthesis but requires more energy for ATP
production and protein synthesis, reducing the fraction of photosynthetic energy available
in support of host metabolism. Conversely, low light conditions limit energy production
regardless of maintenance demands. These mechanisms predict a non-linear relationship
between light availability and net autotrophic benefit to the coral host, providing an
explanation for why abundant light in shallow reefs does not necessarily translate into
enhanced coral growth or physiological performance (*5*).

To measure the effect of metabolic cost across high and low light conditions, we
conducted a series of experiments to characterize coral photoacclimatory responses,
quantify energetic demands of PSII damage-repair activity, and its impact on energy
translocation to the coral host (**Fig. S1**). First, (1) we engineered a custom light system
capable of simulating natural diurnal light cycles under controlled irradiance and spectral
quality. This system minimized the confounding effects of ultra violet (UV) light (**Fig. S2**)
and stabilized daily light integrals (DLI in mol quanta m^{-2} day^{-1}) against the variability
caused by cloud cover (*51*, *52*). Second, (2) coral fragments from the depth generalist
Orbicella faveolata were acclimated to DLI treatments for four weeks, followed by a
detailed characterization of its optical, structural, and physiological light phenotypes,
including PSII half-life and turnover dynamics. Third, (3) coral fragments were exposed to
short-term shifts in DLI to evaluate repair capacity under light conditions that induce light
limitation and light stress. Lastly, (4), we developed a bio-optical model integrating
empirical results and theoretical considerations to predict shifts in photosynthetic usable
energy available to the coral host across a depth gradient. This mechanistic framework
provides a tool to predict the natural habitat and depth distribution of various coral taxa as
well as quantify the impact of environmental changes and increase turbidity (diminished
water quality) on the energetic cost to reef-building corals.

Results

Rapid photoacclimation across a realistic depth-dependent light gradient

Colonies in replicates were exposed to daily light integrals of 4 (DLI-4), 12 (DLI-12), 22 (DLI-22), and 32 (DLI-32) mol quanta m^{-2} day^{-1} (**Fig. 1A**, **Fig. S3A**), which simulates a depth gradient of 5–34 m (assuming a diffuse attenuation coefficient for downwelling irradiance, K_d, of 0.07 m^{-1}; typical for clear-water tropical reefs). PSII maximum photochemical efficiency (F_v/F_m) for experimental replicates (sample fragments) reached a steady state by day 10 (**Fig. S3B**), indicating that samples had acclimated to each of their respective daily doses of irradiance.

Diurnal photoprotection scaled with light exposure

Traits were treated as continuous functions of DLI rather than as discrete treatment means, characterizing phenotypic adjustments across the light gradient (**Fig. S4**). F_v/F_m was highest under DLI-4 and declined linearly with increasing light intensity (**Fig. 1B**). To quantify diurnal variation in PSII excitation pressure, we measured maximum pressure over PSII (Qm), which increased linearly corresponding to the transition from low to high irradiance (**Fig. S3C**). Qm was positively correlated with non-photochemical quenching (NPQ; $r^2 = 0.96$), consistent with increases in photoprotective energy dissipation capacity as excitation pressure increased.

PSII photochemical efficiencies, maximum (F_v/F_m) and effective ($\Delta F/F_m^{'}$), measured every hour captured diurnal variation in PSII performance over the diurnal cycle. Differences in photochemical efficiency during the day were driven by light exposure, with a characteristic decline during the morning, a minimum at midday, and a reciprocal recovery in the afternoon (**Fig. 1C**). These diurnal dynamics reflected the balance among photoprotective mechanisms: induced dissipation as heat of the energy absorbed in excess (NPQ), induced PSII inactivation, and recovery (including NPQ relaxation) over the course of the day. The amplitude of diurnal oscillations increased proportionally with light exposure, reaching a maximum under the highest light treatment (DLI-32), which resulted in progressive increases in NPQ (**Fig. 1D**).

Optical reorganization reduces bulk light absorption under high irradiance

Coral tissue pigmentation darkened or lightened in response to the corresponding light treatment (**Fig. 2A**), resulting in systematic changes in tissue optical properties (**Fig. 2A–B, Fig. S3D**). From minimum to maximum exposure, Chlorophyll a (Chl$a$) content declined by $\approx 36\%$ (113 ± 13 to 41 ± 7 mg Chl a m^{-2}), while intact tissue absorbance peak at 675 nm (De_{675}) decreased by $\approx 10\%$. This corresponded to a decline in average absorptance in the PAR range (A_{PAR}) of 6% from low to high light conditions. The reduction in bulk absorption resulted in a non-linear increase in chlorophyll-specific absorption (a^*_{Chla}) as Chl$a$ declined with increasing light exposure (**Fig. 2B**).

Photosynthetic capacity is maintained despite increasing respiratory demands

Metabolic rates were quantified using photosynthesis–irradiance (P/E) curves fitted with a hyperbolic tangent model, from which four photosynthetic parameters were extracted: photosynthetic efficiency (α), maximum photosynthetic rate (P_{max}), saturation irradiance (E_k), and compensation irradiance (E_c) (53). Photosynthetic efficiency (α) increased as light exposure decreased, indicating enhanced efficiency under low irradiance conditions (**Fig. 2C**). In contrast, maximum photosynthetic rate (P_{max}) remained stable across all light treatments at approximately 12 ± 0.33 μmol O_2 m^{-2} s^{-1} (**Fig. 2D**), indicating that photoacclimation preserved maximal photosynthetic capacity despite large differences in light exposure and pigmentation.

P/E curves allowed estimation of dark respiration rates (Rd), which increased linearly with light exposure (**Fig. 2E**), indicating a higher ATP demand in corals acclimated to higher light conditions. The phototrophic contribution to host energy requirements also increased with light exposure, as reflected by the higher daily integrated photosynthesis-to-respiration ratios (54) (**Fig. S3E**). The minimum quantum requirements of oxygenic photosynthesis (Φ^{-1}_{O2}) ranged from 10 to 17 quanta O_2^{-1} across treatments (**Fig. 2F**) and the quantum efficiency of the photosynthetic process (Φ_{O2}) from 0.1 to 0.059 mol O_2 released per mol quanta absorbed. Both parameters revealed values close to the theoretical minimum of 8 quanta O_2^{-1} for Φ^{-1}_{O2} and the theoretical maximum of 0.125 for Φ_{O2} (55, 56) for functional oxygenic photosynthesis.

PSII functional half-life shortens sharply with increasing light exposure

Having quantified how absorbed light energy is dissipated as excitation pressure increases, and the resulting metabolic adjustments, we next examined processes regulating maintenance of functional PSII reaction centers under contrasting light conditions. First, to separate damage accumulation (before noon) from repair (before dusk), we inhibited *de novo* protein synthesis using chloramphenicol (CAP) (**Fig. 3A**). In control corals (−CAP), PSII activity declined through midday and recovered by the end of the light cycle as expected, whereas CAP-treated corals (+CAP) showed sustained loss of PSII activity consistent with inhibited repair.

This approach enabled estimation of PSII functional half-life ($t_{1/2}$, h), defined as the number of daylight hours required to inactivate 50% of functional PSII reaction centers (**Fig. 3B**). PSII half-life declined exponentially from ~13 h under low light to ~3 h under high light exposure (**Fig. 4A**). This resulted in PSII turnover rates (τ_{PSII} day^{-1}) increasing significantly from low to high light exposure (**Fig. 4B**), revealing accelerated replacement of photodamaged reaction centers under high irradiance.

To assess molecular regulation underlying these dynamics, we quantified transcriptional and translational responses of major PSII reaction center proteins. Quantitative immunoblotting revealed significant differences in D1 protein abundance across the light conditions (**Fig. 4D–F**). D1 content declined sharply with increasing light exposure, mirroring the exponential decrease in PSII half-life and supporting regulation at the translational or post-translational level (**Fig. 4E**). Normalization of D1 content to chlorophyll *a* content yielded the same pattern (**Fig. 4F**). In contrast, transcript abundance of the core components of the reaction center *psbA* (D1) and *psbD* (D2), and inner antenna proteins (that flank the D1/D2 heterodimer) *psbB* (CP47) and *psbC* (CP43), showed no systematic relationship with light treatment (**Fig. 4G**; **Fig. S5**).

PSII half-life rapidly adjusts to acute changes in light exposure

To quantify short-term adjustments in PSII half-life under acute light stress, corals acclimated to a given DLI were transferred to the three alternative light conditions for a day, following the same CAP protein synthesis inhibition protocol (**Fig. S6**). PSII half-life responded rapidly to the new light environment and was independent of the prior acclimatory state, indicating that short-term regulation was driven by the destination light condition rather than the acclimation history (**Fig. 4H**).

To place these responses in the appropriate environmental context, we compared changes in half-life of PSII activity under our experimental PAR, and under natural sunlight, which also includes ultraviolet radiation (UV). In both cases, PSII half-life exhibited comparable exponential declines as a function of the destination light condition (**Fig. 4H**). Corals transferred from lower light treatments (DLI-4, DLI-12, DLI-22) to the highest light level (DLI-32) converged on a PSII half-life of approximately 3 h, matching that of colonies acclimated to DLI-32. These results demonstrate that PSII half-life can be rapidly adjusted within a single day across a wide range of light environments.

Despite this rapid adjustment, short-term regulation was insufficient to prevent the accumulation of PSII photodamage under high excitation pressure conditions. The magnitude of light stress, conveyed as the balance between PSII damage and repair, and quantified as *relative-F_v/F_m* changes (see methods) (*57*), declined with increasing excess excitation energy (EEE) (**Fig. 4I**). This indicated an increase in the magnitude of physiological light stress due to the rapid accumulation of damaged PSII.

Energetic cost of photoacclimation constrains usable energy across depth

The bio-optical model developed here integrates theoretical considerations, empirical measurements, and equations describing phenotypic adjustments to predict the photosynthetic energy available for the coral host in depth-dependent light gradients (see methods). In a previous model implementation (*5*), community-averaged photophysiological responses were represented using a lumped-parameter approach, with photosynthetic and maintenance parameters optimized to reproduce observed biodiversity patterns across depth. Here, we incorporated light-dependent photoacclimatory profiles to explicitly predict photosynthetic energy allocation.

First, we computed DLI as a function of depth across a range of diffuse attenuation coefficients (K_d = 0.04–0.40 m^{-1}), spanning from clear to turbid water conditions, to account for variation in underwater light environments, (**Eq. 1–2; Fig. S7**). Then, for each optical scenario, daily gross photosynthetic output ($DPg_{(z)}$) was estimated from photosynthesis–irradiance curve parameters (**Eq. 3–4**), while depth-specific maintenance cost ($ECMpt_{(z)}$) was derived from empirically constrained PSII turnover rates (**Eq. 5**). These components were computed to predict the photosynthetic usable energy supply (*PUES*) available to the coral host after accounting for symbiont metabolic demands (**Eq. 6**).

Across all optical scenarios, *PUES* exhibited a unimodal relationship with depth (**Fig. 5A**). In shallow waters, high irradiance required elevated maintenance cost to sustain photosynthetic rates. Consequently, an increasing fraction of photosynthetic energy was allocated to symbiont maintenance rather than host translocation. At greater depths, light

limitation reduced carbon fixation, constraining energy availability despite lower PSII
repair activity. Importantly, the depth at which *PUES* was maximized shifted
systematically with K_d (**Fig. 5B**). Consequently, clearer waters supported deeper energetic
optima, whereas increased turbidity compressed the energetic window for coral
distribution toward shallower depths. These results define an optimal bio-optical state for
coral performance that emerges from the balance between photosynthetic output and
maintenance cost, and whose position and extent depend on water-column optical
properties.

Discussion
This investigation provides a mechanistic basis for disentangling the apparent paradox
between light availability, coral physiological performance, and vertical distribution of
coral species by demonstrating that energetic constraints arise at both ends of the depth
gradient. Clearly, carbon acquisition is limited by low irradiance. In shallow waters,
however, abundant light increases the metabolic cost of maintaining photosynthetic rates.
Across a realistic light gradient, maximum photosynthetic capacity remained stable
despite rising excitation pressure, enhanced photoprotective capacity, and elevated
respiration. This stability was sustained by accelerated PSII turnover and shortened
reaction center half-lives, imposing increasing ATP demand under high irradiance levels.
Consequently, greater light availability did not directly translate into greater net energetic
benefit to the host. Instead, a growing fraction of photosynthetic usable energy was
withheld by the symbiont for its cellular maintenance.

When these physiological dynamics were integrated into a depth-resolved bio-optical
model, they predicted a unimodal relationship between light and usable energy supply,
defining an optimal depth range bounded by shallow maintenance costs and deep carbon
limitation. Photosynthetic Usable Energy Supply *(PUES)* was previously framed
conceptually within species–energy theory (*33*) to explain patterns of coral biodiversity
(*5*); here we provide the physiological basis for these distributions patterns by explicitly
incorporating the metabolic cost of sustaining photosynthesis under a range of high to low
light conditions. Classic carbon-budget studies show that zooxanthellae translocate the
vast majority (more than 95%) of their net daily fixed carbon to the coral host, with the
fraction of host respiratory demand this covers varying substantially with species, depth,
and light regime, (*8, 58*). Translocated products such as glycerol, glucose, and other
metabolites that vary among Symbiodiniaceae lineages (*7, 59–63*) support host
metabolism, growth, and calcification, whereas only a small fraction is retained for algal
growth (*58, 64*). Our results demonstrated that this translocation is predictable. As
irradiance increases, ATP-intensive photoprotective processes, particularly rapid PSII
repair via D1 turnover, reduce the fraction of organic carbon available for host use.

The cost of maintaining cellular function is rarely included in photosynthesis or primary
productivity models, which traditionally rely on photosynthesis:respiration ratios and/or
chl*a* to identify conditions of positive carbon balance (*5, 65–67*). By coupling depth-
dependent photosynthetic output with associated maintenance cost, *PUES* reframes coral
energetics as it centers in photosynthetic usable energy for the host. This bio-optical
model emerges as a tool to scale physiological limits to the ecological potential of
photosynthetic organisms, which in the case of zooxanthellate corals, involves colony
growth and morphology as well as gamete production and the timing of reproduction.
Additionally, it offers a framework for predicting how environmental stressors contract
energetic boundaries by necessitating greater investment in metabolic maintenance.

The stability of P_{max} across the range of light exposure reveals the central energetic
problem: photoacclimation in the coral holobiont preserves photosynthetic capacity
without increasing it. Understanding why requires examining each component of the
acclimatory response — photoprotection, optical adjustment, and reaction center
maintenance — and what each reveal about the limits of metabolic flexibility.

The first and energetically cheapest line of defense against excess irradiance is thermal
dissipation via NPQ. The tight coupling between Q_m and NPQ ($r^2 = 0.96$) reflects the
dynamic photoprotective equilibrium, driven by xanthophyll cycling, that scales
continuously with excitation pressure across diel cycles and light conditions (1, 16, 68–
71), consistent with prior field and laboratory observations (16, 70, 72–75). Critically,
however, NPQ alone cannot prevent photodamage when excitation pressure exceeds the
capacity for thermal energy dissipation. The declining F_v/F_m with increasing irradiance
marks precisely this transition, the point at which the NPQ buffer is insufficient and repair
via D1 turnover becomes the dominant maintenance strategy, at escalating ATP cost.

The optical organization of the coral holobiont reflect similar acclimatory limits. The coral
skeleton and tissue optics are engineered for light amplification (11, 24). Despite a nearly
threefold decline in Chl$a$ content, overall absorptance remained effectively constant, with
total energy absorbed by the holobiont preserved regardless of pigment reduction. Rather
than lowering light absorption, pigment loss increased chlorophyll-specific (a^*_{Chla}) and
symbiont-specific (a^*_{sym}) light absorption efficiency, driving the symbiosis toward the
upper boundary of holobiont acclimatory capacity (11, 76, 77). This proximity to the
optical maximum, constrains further pigment reduction under additional stressors such as
thermal stress (78), explaining why bleaching thresholds are usually reached more rapidly
in high light environments. Ecological implications are significant, as the same optical
trait that enables colonization of light-limited deep reefs (27, 29) simultaneously prevents
meaningful reduction of absorbed energy through pigment adjustment at high irradiance.
In shallow reef environments, when this optical limit is reached and repair capacity is
overwhelmed, selection favors colony morphologies with structural self-shading, $i.e.$
branching, foliose, and massive forms with strong within-colony light attenuation (27, 30,
79).

The stability of P_{max} across light treatments reflects a key constraint itself. It means that no
additional irradiance above saturation serves to enhance carbon fixation rates, and thus,
the absorbed energy above E_k can only be dissipated as heat, channeled into NPQ, or
directed into PSII damage:repair. In our investigation, low light phenotypes approached
the theoretical maxima for PSII photochemical efficiency in dinoflagellates in $hospite$
($F_v/F_m \approx 0.7$) and theoretical minimum quantum requirements for oxygen evolution (~10
quanta O_2^{-1}) (55, 56), marking the lower boundary of photoacclimatory capacity (1, 54).
At the other extreme, high light phenotypes with higher light absorption efficiencies
(a^*_{Chla}) were proportionally less efficient at converting absorbed energy into carbon
fixation (lower α and Φ_{max}). The symbiosis therefore operates near the boundaries of its
photosynthetic range at both ends of the depth gradient, saturated above and limited
below, with maximum photosynthetic capacity as stable outcome.

Notably, the stability of maximum photosynthetic rates across treatments indicates that
coral photoacclimation primarily operates through modulation of light absorption, energy
dissipation and protein turnover, and not by regulating photosynthetic capacity. This
decoupling suggests that photosynthetic rates in $hospite$ are constrained by downstream

processes, such as inorganic carbon diffusion or alternative electron sinks within the symbiosome (*80–83*). Crucially, it demonstrates that maintaining photosynthetic performance under high light requires progressively greater metabolic support without proportional energetic return. The source of this increasing cost is PSII turnover and D1 protein dynamics.

By arresting protein synthesis, we decoupled PSII damage from repair and quantified changes in PSII half-life across experimental light conditions. PSII half-life declined exponentially with increasing irradiance and closely paralleled reductions in D1 protein abundance. Accordingly, corals acclimated to high light conditions maintained maximum photosynthetic capacity via faster replacement of photodamaged reaction centers and lower D1 pools. Similar relationships between D1 abundance, PSII half-life, and light history have been documented in higher plants, including how rapid D1 turnover accompanies elevated metabolic demand under high light exposure (*37, 84–86*).

Protein turnover is among the most energetically expensive cellular processes (*34–37*). In higher plants, thylakoid protein turnover accounts for 13–38% of total ATP consumption, with D1 replacement representing a major fraction of this budget (*37*). Energy for this repair cycle is supplied through respiration, linking PSII maintenance directly to metabolic demand. The elevated dark respiration (R_d) found under high light conditions is therefore consistent with increased ATP requirements for D1 synthesis and PSII reassembly (*36, 44, 87*). ATP availability is known to constrain *de novo* D1 synthesis and full PSII repair (*47, 88, 89*), suggesting that ATP supply becomes the limiting currency sustaining photosynthetic function under light stress.

Under high irradiance, absorbed energy that exceeds photosynthetic capacity cannot be converted into additional carbon fixation and is instead partitioned between dissipation as heat through *NPQ* and investment in accelerated PSII repair. In this context, D1 turnover operates as a regulated sink for excess excitation, preventing over-reduction of the electron transport chain and protecting PSI under acceptor-side limitations (*45, 86*). This hierarchy reflects structural cost optimization within the photosynthetic apparatus. Because antenna complexes are far more abundant ($\sim 7.6 \times 10^6$ cell^{-1}) and energetically more expensive than D1 protein ($\sim 1.0 \times 10^5$ cell^{-1}) (*90*), it makes selective turnover of PSII reaction centers a comparatively economical strategy. In fact, regulation of reactive oxygen species within the photosynthetic apparatus shows that while chlorophyll triplet states in antenna complexes are efficiently quenched by carotenoids, spatial constraints within the PSII reaction center prevent such protection, allowing singlet oxygen to target D1 directly and requiring rapid turnover as a controlled safeguard (*40, 91, 92*). D1 protein, then, becomes the unavoidable target of oxidative damage, making its rapid turnover a necessary mechanism to preserve photosynthetic integrity.

This bottleneck has direct ecological implications. Earlier work has primarily focused on how PSII repair impairment under thermal stress leads to coral bleaching. When repair fails to match damage rates, D1 protein abundance drops below the threshold needed to sustain photosynthetic activity, resulting in pigment loss, symbiont degradation, and ultimately, expulsion or symbiont digestion by the host (*48, 93–96*). Even in the absence of heat stress, high light and even PAR alone shortened PSII half-life to approximately three hours and reduced D1 abundance, imposing substantial ATP demand. Thus, the upper tolerance limits of the coral holobiont are defined not solely by photodamage, but by the energetic capacity to sustain PSII repair. This constraint is always active in high

light conditions, revealing the high susceptibility of shallow water corals to any additional
environmental stress. We confirmed the environmental regulation of this mechanism
comparing PSII half-life under experimental PAR (400–700 nm) and natural sunlight
(PAR + UV). No significant differences in PSII half-life were found, consistent with PAR
being the primary driver of photoinactivation and daily damage accumulation. This result
challenges the assumption that UV is the dominant driver of PSII photodamage in
shallow-water corals (*83*, *97*).

Eco-evolutionary significance

We identified PSII repair as a fundamental energetic bottleneck that constrains coral
holobiont performance at high irradiance. Sustaining photosynthesis under light stress
requires continuous ATP investment in protein turnover, and thus diverting resources from
growth, reproduction, and calcification. In this way, energetic maintenance cost of
photosynthesis sets the upper boundaries of coral environmental tolerance.

In the bioenergetic framework, the *PUES* bio-optical model, specifically *PUES* optimum
($Depth_{opt}$) defines the depth at which photosynthetic supply and maintenance demand are
optimally balanced, maximizing usable energy translocation while minimizing the cost of
sustaining photosynthetic integrity. The $Depth_{opt}$ position and extent of coral colonization
depend on water column optical properties. When exploring a range of light attenuation
coefficients (K_d), spanning clear to turbid conditions, our analysis revealed that increased
light attenuation shifts $Depth_{opt}$ toward shallower waters and compresses the energetic
envelope. Therefore, environmental degradation that alters underwater light conditions,
including progressive ocean darkening, reshapes the energetic landscape rather than
simply reducing light availability.

This energetic landscape has further evolutionary consequences. Intermediate depths,
where usable energy is maximized, represent zones of expanded energetic niche space
capable of supporting multiple photophysiological strategies, species coexistence and
diversification. In contrast, energetic extremes due to carbon limitation at depth or higher
energetic demands of maintenance cost in shallow water, restrict viable trait combinations
and narrow ecological opportunities through exclusion or specialization. Such patterns
align with general species–energy theory and with emerging genomic evidence that
closely related coral lineages repeatedly diverge across narrow depth gradients, with
adaptation to distinct light conditions linked to environmental sensing and reproductive
timing (*98–100*).

The bio-optical model developed here and focused on *PUES* provides a mechanistic
scaling of coral bioenergetics to ecological patterns, with the power to delineate energetic
boundaries of coral performance in the depth distribution of zooxanthellate corals. This
allows predicting the fraction of photosynthetically derived energy that is available for
supporting host metabolic requirements, including maintenance, growth, reproduction, and
ecological interactions, hence, contributing to define fundamental ecological niches of the
coral holobiont.

430
431 **Materials and Methods**
432
PAR LED Light System
We developed an experimental light system by engineering custom-made lamps and
software able to simulate diurnal light cycles. Natural light in the tanks was absent
throughout the experiments. The illumination system relied on 100W light-emitting diodes
(LED)-lamps that enabled automation of the diurnal cycles (**Fig. S3A**). The four light
treatments (Daily Light Integral, DLI) were: 4 mol quanta m^{-2} day^{-1} (DLI-4, peak of
irradiance 200 µmol quanta m^{-2} s^{-1}); 12 mol quanta m^{-2} day^{-1} (DLI-12; peak of irradiance
450 µmol quanta m^{-2} s^{-1}); 22 mol quanta m^{-2} day^{-1} (DLI-22; peak of irradiance 750 µmol
quanta m^{-2} s^{-1}); and 32 mol quanta m^{-2} day^{-1} (DLI-32; peak of irradiance 1100 µmol
quanta m^{-2} s^{-1}).
443
Three parameters were prioritized for the control of each light condition: light intensity,
light quality and time. We built each LED-lamp with a BC Series High CRI COB-LED-
400L-100W- Unit (YujiLEDS®) which best covered the sunlight spectrum in the range of
the Photosynthetic Active Radiation (PAR). LEDs were operated in continuous mode with
a multifunction I/O card (USB-6001, National Instruments Corp., USA) to avoid potential
artifacts on photosynthesis and fluorescence responses related to the "flashing effect" of
commercial LEDs at different pulsating frequencies (*101*). Light intensity was controlled
with a custom-made software in LabVIEW (Laboratory Virtual Instrument Engineering
Workbench, National Instruments Corp., USA). To eliminate natural sunlight inside
experimental tanks, we placed LED-lamps inside black boxes that fully covered each tank
(**Fig. S1**).
455
Diurnal light exposure (mol quanta m^{-2} day^{-1}) was described by integrating the per second
change in irradiance (µmol quanta m^{-2} s^{-1}) over a day as Daily Light Integral (DLI). Light
exposure as an integral has proven to be a useful tool for assessing irradiance accumulated
when the response of interest is growth or acclimation (*102*). DLIs conditions captured a
wide range of coral vertical distribution in clear waters (5 to 35 m depth for a
downwelling light attenuation coefficient, K_d, of 0.07 m^{-1}). Irradiance at depth depends on
the optical properties of the water column given by the Kd at any reef location. Irradiance
was calibrated and continuously monitored with the quantum sensor LI-1400 (LI-COR,
USA).
465
Light spectra also play an important role in photosynthesis performance. The effective
activation of the repair cycle of photodamaged PSII has its own action spectrum. Studies
have shown that UV-radiation damages PSII more effectively than visible light (*97*), and
that the reported action spectra are rather dissimilar in the range between 400 and 700 nm.
We used a sun-like spectrum and UV was excluded by design (**Fig. S2**). We were able to
effectively induce PSII damage repair cycles in the simulated DLIs treatments.

Acclimation to Daily Light Integrals

We sampled 64 fragments from 4 *Orbicella faveolata* colonies (each a different genet)
between 4-5 m depth under similar light conditions in La Bocana, Puerto Morelos
National Park. Experimental coral fragments (~5-9 cm^2) were set for a 2-week healing
period after sampling by placing them on PVC-deployed structures located at 5 m depth
near the sampling area. Corals were then transported to the National Autonomous

University of Mexico (UNAM) mesocosm facility, where each colony was equally
distributed in four outdoor tanks (shaded natural sunlight to 16 ± 3 mol quanta m^2 day^{-1})
equipped with running natural seawater (n = 16 fragments per tank). Water temperature
and flow were maintained at $28 \pm 1°C$ and 1100 L h^{-1} using commercial aquaria chillers
and heaters. Water temperature in the tanks was continuously monitored with Hobo data
loggers (Onset Computer Corporation, MA, USA). Corals were acclimated to the
mesocosm for two weeks before LED installation, followed by 4+ weeks under
experimental DLI conditions before phenotyping.

We monitored daily PSII maximum photochemical efficiency (F_v/F_m) at dusk and
effective photochemical efficiency ($\Delta F/F_m'$) at noon, with a pulse-amplitude modulated
fluorometer (Diving-PAM; Heinz Walz, Effeltrich). Maximum excitation pressure over
PSII ($Q_m = 1 - [(\Delta F/F_m'$ at noon) / (F_v/F_m at dusk)]) was also estimated to monitor the
moderate light stress induced during natural and daily photo-acclimation, and the
adjustment to a stable photo-acclimatory condition (1).

Phenotyping in the range distribution of zooxanthellate corals

Following the 4+ weeks of acclimation to the experimental light gradient, we measured a
suite of traits to characterize coral phenotypes (**Table 1**). Chl$a$ fluorescence measurements
were recorded as photochemical efficiency (F_v/F_m and $\Delta F/F_m'$), and non-photochemical
quenching was calculated as $NPQ = [(F_m-F_m')/F_m]$.

To describe changes in coral pigmentation, we estimated Chl$a$ and Chl$c$ contents. Briefly,
an air gun and filtered sea water (FSW) were used to obtain tissue slurries subsequently
homogenized at low temperature (Tissue-Tearor Homogenizer BioSpec Inc, USA) and
centrifuged. The resulting pellet was re-suspended in filtered seawater and preserved for
pigment concentration (extracted with acetone/dimethyl sulfoxide 95:5 vol/vol) and
Symbiodiniaceae cell density (counted in a haemocytometer after the addition of 200 μL
of iodine preservation solution). Chl$a$ and Chl$c$ concentrations were estimated
spectrophotometrically (3 reads per sample) with a modular spectrometer (Flame-T-UV-
VIS, Ocean Optics Inc., USA) using the equations described by Jeffrey and Humphrey
(103).

Light absorption spectra were determined on intact *in vivo* coral tissues from reflectance
(R) measurements. R was used to estimate Absorbance (De) [$De = \log(1/R)$] and
Absorptance (A), assuming that Transmittance through the coral skeleton was negligible
(104). A was estimated for the PAR range (400 and 700 nm; $A_{PAR} = 1 - R_{PAR}$); and at the
675 peak of Chl$a$ ($A_{675} = 1 - R_{675}$).

Specific-absorption coefficients (a*) were normalized to Chl$a$ content (a*$_{Chla}$; m^{-2} mg
Pigm^{-1}) as [a*$_{Chla}$= (De_{675}/ρ)·ln(10), where $De_{675} = \log(1/R_{675})$] (11, 77, 105). Likewise, a*
was normalized to symbiont cell density (a*$_{Sym}$; m^{-2} sym^{-1}) as [a*$_{sym}$ = (De_{675}/sym ρ) ·
ln(10)]. These descriptors estimate the holobiont efficiency to absorb light (a*$_{Chla}$) and
symbiont specific light absorption efficiency ($a*_{sym}$) (78).

To describe changes in coral metabolic rates, we used photosynthesis / irradiance curves
(P/E) fitted to a hyperbolic tangent model (**Fig. S4A**). Briefly, coral fragments were
incubated using a laboratory-made water-jacketed respirometer (76) with filtered sea
water at a constant temperature (28 °C) and constant water flow generated by continuous

| 520 | agitation from magnetic stirrers, which ensure that the boundary layer around coral |

agitation from magnetic stirrers, which ensure that the boundary layer around coral
surfaces is broken. One chamber was measured as a blank (no coral fragment) to correct
for background activity in the water. A LED-system was designed to enable automation of
light increments every 10 min. Oxygen evolution was measured continuously with a fiber-
optic oxygen meter system (FireSting, Pyroscience) to estimate the rate in the 10 min
range of: dark (pre-illumination), light (10 light increments), dark (post-illumination) for a
total period of the 2 h incubation. Photosynthetic efficiency (α), compensation irradiance
(E_c), saturation irradiance (E_k), respiration rates in the dark (R_D), light-enhanced
respiration (R_L), and maximum photosynthetic rates (P_{max}), were estimated from the PE
curves (*106*, *107*). Each curve reflects the photosynthetic potential of the light phenotype.
To estimate the energy absorbed in excess, we calculated the hours of light exposure
above E_k (number of hours in P_{max}; H_{sat} (*108*)) and the magnitude of energy absorbed
under these conditions (mol quanta m^{-2} day^{-1}).

Using absorptance values determined above [$A = 1- R$], we calculated the maximum
quantum yield of photosynthesis (Φ_{O2}; mol oxygen evolved per mol quanta absorbed^{-1})
and its inverse, the minimum quantum requirements of photosynthesis [(Φ^{-1}_{O2}; $= 1/(\alpha *$
A_{PAR}); quanta required to release an O_2 molecule^{-1})] (*106*, *107*). Φ_{O2} was estimated from
α, the slope of the linear regression of photosynthesis in the sub-saturating region of the
P/E curve; corrected for the light absorbed (A) in our experimental setting [(photosynthetic
utilizable radiation (PUR) absorbed (A_{PUR})] (*76*, *106*, *107*). To estimate the integrative
index of the holobiont's metabolism, we calculated daily integrated photosynthesis-to-
respiration ratios [Pg (E) day^{-1}: R (E) day^{-1}].

Algal symbiont identification

Orbicella faveolata hosts distinct species of algal symbionts in this region (*e.g.*,
Symbiodinium sp. A3, *Breviolum faviinorum*, *Breviolum meandrinum*, *Cladocopium* sp.
C3, *Cladocopium* sp. C7, and *Durusdinium trenchii*), with the possibility of multiple
species simultaneously inhabiting a same colony (*109–111*). To account for this functional
diversity, symbiont identity was determined using two complementary approaches.

(1) One cm^2 coral fragments were used to extract genomic DNA using a modified
Promega Wizard protocol (Promega Madison, WI) (*112*, *113*). The large ribosomal
subunit (*LSU* rDNA) was amplified using *28S* forward (5'-
CCCGCTGAATTTAAGCATATAAGTAAGCGG-3') and *28S* reverse (5'-
GTTAGACTCCTTGGTCCGTGTTTCAAGA-3') (*114*) and an annealing temperature of
66 ºC. Amplifications were Sanger sequenced using the Big- Dye Terminator 3.1 Cycle
Sequencing Kit (ThermoFisher Scientific, Waltham, MA) and analyzed on the Applied
Biosciences Sequencer (Applied Biosciences, Foster City, CA) at the Penn State
University Genomics Core Facility. Sequences were checked and edited in Geneious
v11.0.5 (Biomatters Ltd, Auckland, NZ) and were queried against the NCBI nucleotide
database using BLASTn (National Center for Biotechnology Information).

(2) We separately aligned raw RNA-Seq reads (see below) from each sample against
reference transcriptomes using bwa v0.7.18 (*115*); mapping rates were determined using
SAMStat v2.2.3. The reference transcriptomes encompassed eight datasets from the most
closely related Symbiodiniaceae that *Orbicella* associates with: *Symbiodinium* sp. A3
(*116*), *Breviolum faviinorum* (*116*), *Breviolum* sp. B5 (*116*), *Cladocopium proliferum*
(*117*), *Cladocopium* sp. C1 MI (*118*), *Cladocopium* sp. C1 SM (*118*), and *Durusdinium*

trenchii (*117*, *118*). Symbionts present in each sample were determined by comparing
LSU rDNA and mapping results (**Fig. S5A**).

PSII half-life and turnover rates under acclimatory conditions

After the 4-week acclimation, the PSII damage and repair cycle was determined through
monitoring the variation of F_v/F_m and $\Delta F/Fm'$ in 12 h diurnal cycles. We tracked the
accumulation of photo-inactivated PSII and the accumulation of photo-repaired PSII. The
damage accumulation phase runs from dawn to midday, and repair from midday to dusk.

To quantify the effect of light acclimation on the photodamage and repair of the PSII
active pool, we inhibited the PSII repair process. In each acclimatory condition, fragments
were separated into two subgroups: control (n = 8, no inhibitor -CAP) and the inhibitory
treatment (n = 8, treated with D1 protein synthesis inhibitor chloramphenicol; +CAP).
Subgroups were placed separately in 1 L plastic containers with internal seawater
circulation (FSW) and positioned within tanks (water baths). Water was exchanged in the
containers every 2 h for both treatments (FSW+CAP) and controls (FSW-CAP). PSII
photochemical efficiency was measured every hour, and samples were snap frozen after
dusk for further analyses. This protocol was repeated for every light condition after 4+
weeks of acclimation. To estimate PSII half-life ($t_{1/2}$, h) photochemical efficiency
treatment-to-control ratios [+CAP / -CAP] were curve fitted using a logistic sigmoidal
function and expressed for each light condition as the number of hours required to
inactivate 50% of PSII ($t_{1/2}$, h).

Abundance of PSII D1 protein

Algal pellet assigned for protein analyses was flash frozen in liquid nitrogen and stored at
-80 °C. We quantified D1 protein content in algal symbionts isolated from three coral
fragments for each light regime. Each immunoblotting assay was performed multiple
times to load: (lane 1) a negative control, (lanes 2–6) 0.10-2.00 pmol *psbA*/D1 standard,
(lanes 7–10) one sample from each light regime to be normalized by total protein (μg
protein^{-1}), and (lanes 11–14) one sample from each light regime to be normalized by Chl *a*
(mg Chl a^{-1}). We performed quantitative immunoblotting following established protocols
(*93*, *94*, *96*, *119*, *120*). Briefly, proteins were extracted using protein extraction buffer PEB
(Agrisera AS08 300) supplemented with Pefabloc® SC (Sigma-Aldrich 11429868001) to
inhibit proteolysis. Samples were subjected to three cycles of sonication-to-liquid N_2 to
break cell wall and avoid over-heating. Protein concentrations were determined using The
Better Bradford assay with a standard curve. Equal amounts of total protein were mixed
with Bolt™ LDS sample buffer (Invitrogen B0007) and Bolt™ Sample Reducing Agent
(ThermoFisher B0004), denatured, and separated on NuPAGE™ 4–12% Bis-Tris gels
(Invitrogen NP0329BOX) alongside PageRuler™ Plus pre-stained protein ladder
(ThermoScientific 26619). Following electrophoresis, proteins were transferred to PVDF
membranes (Invitrogen LC2005) using Bolt™ transfer buffer (Invitrogen BT0006) and
blocked with ECL™ Advance Blocking Reagent (GE Healthcare GERPN418).
Membranes were probed with a rabbit polyclonal anti-PsbA (D1) antibody (Agrisera
AS05 084), followed by HRP-conjugated chicken anti-rabbit secondary antibody
(Agrisera AS10 833). Membranes were stained post-transfer with SimplyBlue™ SafeStain
(Invitrogen LC6060). D1 protein quantification was achieved by co-loading known
amounts of PsbA/D1 standard (Agrisera AS01 016S), imaging on a chemiluminescent
imager (ChemiDoc BIO-RAD), and pixel-quantifying in ImageJ.

Expression of PSII-complex genes

Expression of PSII genes from algal symbionts was assessed following an RNA-Seq
approach. Total RNA was extracted from 16 fragments, 4 untreated fragments (-CAP) per
light exposure. Each sample was homogenized in TRIzol reagent (Ambion, Life
Technology) before centrifugation with chloroform for 15 min at 12,000 g at 4 °C. The
aqueous phase was then isolated and cleaned using Qiagen RNeasy Mini kit (Qiagen), as
per the manufacturer's protocol with an additional on-column DNase treatment using
RNase-Free DNase Set (Qiagen). To maximize concentration of eluted RNA, the same
35 μL of RNAse-free water was twice passed through the spin column for the final
isolation step. Concentration and purity were pre-screened via Qubit spectrometer and
quality assessed (RNA integrity number; RIN > 7) with Agilent 2100 Bioanalyzer (Agilent
Technologies). Purified RNA was sent to Biomarker Technologies (BMK) GmbH where
concentration and purity were re-analyzed (Qubit® dsDNA BR Assay Kit on a Qubit®
2.0. Fluorometer). RNA libraries for 2 × 150 bp paired-end sequencing were non-stranded,
prepared using Hieff NGS Ultima Dual-mode mRNA Library Prep Kit for Illumina
(Yeasen) model:13533ES96. Samples were run in platform Illumina NovaSeq X.

Raw reads were assessed for quality using FastQC v0.12.1 (github.com/s-
andrews/FastQC) and results visualized using MultiQC v1.28 (*121*), followed by filtering
and trimming using Trimmomatic v0.39 (*122*). Pseudoalignments of the processed read
pairs and quantification of expression were done using the quant function of kallisto
v0.51.1 (*123*) in two manners: against all reference transcriptomes combined into a single
database and against separate transcriptomes according to the symbiont type identified per
sample. For the combined database, all reference transcriptomes were concatenated
removing redundant sequences using cd-hit-est v4.8.1 (*124*) at a sequence similarity
threshold of 0.99. The transcripts per million (TPM) output by kallisto were used as
measure of gene expression level. Genes of the core complex of PSII were identified by a
BLASTn v2.16.0 (*125*) search of Symbiodiniaceae GenBank records (JN557844 for *psbA*,
HG515018 for *psbB*, HG515019 for *psbC*, and HG515020 for *psbD*) as queries against the
reference transcriptomes with an e-value ≤ 10^{-5}. The BLAST hit with the highest mean
expression across samples per genus was selected as the representative transcript for each
gene.

PSII half-life under light-stress conditions

To quantify the effect of light stress on the PSII photodamage and repair cycle, and the
variation of PSII half-life ($t_{1/2}$, h), we replicated the experimental protocol to inhibit the
PSII repair process. However, here the 4-weeks acclimated corals were switched (new
coral fragments for each experiment) to an alternative light exposure (*i.e.*, DLI-4$_{acclimated}$
switched to DLI-12$_{destination}$, DLI-22$_{destination}$ and DLI-32 $_{destination}$) for one diurnal cycle. The
experimental protocol was repeated for each light exposure for a total of 12 experiments:
DLI-12$_{acclimated}$ switched to DLI-4$_{destination}$, DLI-22$_{destination}$ and DLI-32$_{destination}$; DLI-
22$_{acclimated}$ switched to DLI-4$_{destination}$, DLI-12$_{destination}$ and DLI-32$_{destination}$; and DLI-
32$_{acclimated}$ switched to DLI-4$_{destination}$, DLI-12$_{destination}$ and DLI-22$_{destination}$.

The severity of light stress was quantified with the metric Excess Excitation Energy *EEE*
(mol quanta m^{-2} day^{-1}) as follows (*57*): (1) Conditions where $EEE = 0$ (EEE_0) indicate that
corals are well acclimated (F_v/F_m is steady day-to-day), no net light stress is induced,
hence no photodamage is accumulated. (2) Conditions where $EEE \neq 0$ ($EEE_{\neq 0}$) indicate

655	corals were switched to a different light condition and may experience light stress. *EEE*
656	was quantified as the difference between DLI in the switched condition ($DLI_{destination}$) and
657	DLI acclimated ($DLI_{acclimated}$). In the case that the switched condition resulted in lower
658	light exposure, *EEE* shows negative values. The magnitude of the physiological stress was
659	measured as the ratio between F_v/F_m in the switched condition and F_v/F_m acclimated,
660	which results in the parameter *relative-F_v/F_m*.

PSII half-life ($t_{1/2}$, h) was estimated following the same protocol described above for PSII
half-life under acclimatory conditions as the number of hours required to inactivate 50%
of the active PSII pool.

Modeling the energetic cost of photo-acclimation

To estimate how the light environment shapes the energetic balance of the coral-algal
symbiosis, we developed a bio-optical model that integrates theoretical light attenuation in
the water column, gross photosynthetic output, and the energetic costs of PSII turnover.
The analysis focused on the estimation of the potentially maximum photosynthetic output
of the symbionts after subtracting the costs of maintenance of photosynthetic activity
through PSII repair, which can be translocatable to the host. Other metabolic costs
required by the maintenance and/or growth of algae and host were not considered into the
model due to the difficulties of parameterization of the large diverse of biological
processes potentially involved. Light availability at depth was expressed as daily light
integrals (DLI) and computed from surface irradiance. Different light attenuation
coefficients (K_d) were used. Depth-dependent photosynthetic output was then estimated
using a photosynthesis-irradiance relationship parameterized with function-valued
phenotyping derived from experimental profiles. To capture the costs of maintaining
photosynthetic function under different light regimes, we incorporated PSII turnover
dynamics by linking PSII lifetime ($t_{1/2}$, h) and its cost, maximum photochemical efficiency
(F_v/F_m), and repair potential to DLI. The resulting framework yields the Photosynthetic
Usable Energy Supply (*PUES*) by the symbionts to the host, a metric that represents the
potentially maximum energy available for the coral host after subtracting the light-
dependent costs of photosynthesis maintenance. Considerations and computations
underlying the model proceeded as follows:

<u>Theoretical and empirical considerations</u>

As light is the primary factor that regulates primary production and the photoacclimatory
response of photosynthetic organisms, we first estimated the amount of light available at a
given depth (*126*):

1. Solar irradiance as a function of depth

The intensity of a parallel-beam radiation of a particular wavelength decreases along its
path in pure water, obeying the Lambert Law. The general equation of light attenuation in
seawater is (*126*):

$$E_{(z)} = E_0 * e^{-Kd*z} \qquad \text{(Eq. 1)}$$

694 Where $E_{(z)}$ is light intensity at depth z (here irradiance value for Photosynthetic Active
695 Radiation, PAR, as UVR was not considered), E_0 is the PAR just beneath the water

696 surface (when $z = 0$) and Kd (m^{-1}) is the vertical attenuation coefficient of downwelling
697 PAR irradiance.

2. Changes in daily light integral (DLI) with depth: Light exposure

Irradiance at any point on the Earth's surface is determined by solar elevation. This rises
during the day from zero at dawn to its maximum value at noon, and then diminishes in a
precisely symmetrical manner to zero at dusk under no cloud cover. Light exposure
integrates the variation in irradiance per second and m^2 over time. The integrated quantum
unit for light exposure at a certain depth and day is mol quanta m^{-2} day^{-1}, which describes
the amount of quanta received by m^{-2} during a 24 h period (*126*):

$$DLI_{(z)} = \int_0^N E_0 * Sin\left(\frac{\pi t}{N}\right) * e^{(-Kdz)} \, dt \qquad \text{(Eq. 2)}$$

706 The parameter $DLI_{(z)}$ is then the daily light integral at depth, E_0 is the superficial irradiance
707 at solar noon, N is the daylength (in seconds) and t is time along a day (seconds).

3. Photosynthesis as a function of DLI

709 The response of photosynthetic organisms to available light (from the light saturation
710 curve), depends on two fundamental parameters in our model: the slope α at very low light
711 availability (carbon fixed/oxygen evolved per unit of available light), and the asymptotic
712 maximum production P_{max} (the rate of carbon fixation/oxygen evolved under saturating
713 light) (*127*):

$$GPg_{(DLI)} = Pmax * \tanh\left(\alpha * \frac{DLI}{Pmax}\right) + R(DLI) \qquad \text{(Eq. 3)}$$

Where $GPg_{(DLI)}$ is the rate of gross photosynthesis measured at a given quantum
irradiance, DLI; P_{max} is the maximal photosynthetic capacity; α is the is the photosynthetic
efficiency; and $R_{(DLI)}$ is the light dependent variation of net coral respiration. As indicated
previously, we used here the direct values of R_L and R_D measured for each coral
phenotype, instead a single and general function for *Orbicella faveolata*, maintaining the
same criteria: R_L was used for the period of the day at photosynthesis saturation (P_{max}), R_D
in darkness or very low light levels, and the average for the sub-saturating region,
according to each acclimatory condition.

4. Photosynthetic output as a function of depth

724 To estimate the photosynthetic output $\Delta Pg(z)$ at a given z, we needed to integrate daily
725 photosynthesis at z (modified from (*127*)):

$$GPg_{(z)} = GPmax \int_0^N \tanh\frac{\alpha_{(z)} DLI_{(t)(z)}}{GPmax} \, dt \qquad \text{(Eq. 4)}$$

5. Photosynthetic usable energy for coral maintenance

728 We tracked the relationship between photosynthetic and maintenance at depth and
729 incorporated the energy cost of photoacclimation *PUES(z)* as the cost of protein turnover

730 rate at depth. Energetic cost of 1 turnover (0 if no energy cost is involved in protein
731 turnover) (37, 54):

$$ECMpt_{(z)} = Protein\ turnover_{(z)} * Energetic\ cost \qquad \text{(Eq. 5)}$$

733 Photosynthetic usable energy supply for coral at depth z was expressed by integrating
734 equations 4 and 5:

$$PUES_{(z)} = DPg_{(z)} * (1 - ECMpt_{(z)}) \qquad \text{(Eq. 6)}$$

736 <u>Function-valued phenotyping included in the model, where *z* is depth and DLI daily light</u>
737 <u>integral:</u>

738 *1. Photosynthetic parameters as a function of DLI_{(z)}:*
739   a. Gross maximum photosynthetic rate *GPmax* = 15 µmol O_2 m^{-2} s^{-1}
740   b. Saturation irradiance *Ek* as a function of *DLI_{(z)}* (mol quanta m^{-2} day^{-1})

$$Ek_{(z,DLI)} = 9.50DLI_{(z)} + 300.32$$

742   c. Photosynthetic efficiency α as a function of *DLI_{(z)}*

$$\alpha_{(z,DLI)} = -0.0005DLI_{(z)} + 0.0374$$

744 *2. Protein turnover rate as a function of DLI_{(z)}:*

745   a. PSII-half-time (hours) as a function of *DLI* (mol quanta m^{-2} day^{-1}):

$$PSII1/2_{(z,DLI)} = \left[27.8374DLI_{(z)}^{-0.54723}\right]$$

747   b. Relative maximum photochemical efficiency (*Fv/Fm*):

$$FvFm_{DLI} = -0.0044DLI + 0.698$$

$$Rel.FvFm_{(DLI)} = \left[\frac{FvFm_{(DLI)}}{maxFvFm}\right]$$

750   c. Repair potential (dimensionless):

$$Repair\ potential_{(DLI)} = \left[\frac{12\ hours}{PSII1/2_{(DLI)}}\right]$$

752   d. Protein turnover rate:

$$Protein\ turnover_{(z)} = Repair\ potential_{(z)} * Rel.FvFm_{(z)}$$

References

1. R. Iglesias-Prieto, V. H. Beltrán, T. C. LaJeunesse, H. Reyes-Bonilla, P. E. Thomé, Different algal symbionts explain the vertical distribution of dominant reef corals in the eastern Pacific. *Proceedings. Biological sciences / The Royal Society* **271**, 1757–1763 (2004).

2. G. S. E. Chow, Y. K. S. Chan, S. S. Jain, D. Huang, Light limitation selects for depth generalists in urbanised reef coral communities. *Marine Environmental Research* **147**, 101–112 (2019).

3. T. E. Roberts, S. A. Keith, C. Rahbek, T. C. L. Bridge, M. J. Caley, A. H. Baird, Testing biodiversity theory using species richness of reef-building corals across a depth gradient. *Biol Lett* **15**, 20190493 (2019).

4. R. Tamir, G. Eyal, N. Kramer, J. H. Laverick, Y. Loya, Light environment drives the shallow-to-mesophotic coral community transition. *Ecosphere* **10**, e02839 (2019).

5. T. López-Londoño, K. Gómez-Campo, X. Hernández-Pech, S. Enríquez, R. Iglesias-Prieto, Photosynthetic usable energy explains vertical patterns of biodiversity in zooxanthellate corals. *Sci Rep* **12**, 20821 (2022).

6. G. Pérez-Rosales, M. Pichon, H. Rouzé, S. Villéger, G. Torda, P. Bongaerts, J. Carlot, U. T. P. Consortium, V. Parravicini, L. Hédouin, Mesophotic coral ecosystems of French Polynesia are hotspots of alpha and beta generic diversity for scleractinian assemblages. *Diversity and Distributions* **28**, 1391–1403 (2022).

7. T. C. LaJeunesse, J. E. Parkinson, P. W. Gabrielson, H. J. Jeong, J. D. Reimer, C. R. Voolstra, S. R. Santos, Systematic Revision of Symbiodiniaceae Highlights the Antiquity and Diversity of Coral Endosymbionts. *Current Biology* **28**, 2570-2580.e6 (2018).

8. L. Muscatine, L. R. Mccloskey, R. E. Marian, Estimating the daily contribution of carbon from zooxanthellae to coral animal respiration. *Limnology and Oceanography* **26**, 601–611 (1981).

9. P. G. Falkowski, Z. Dubinsky, L. Muscatine, J. W. Porter, Light and the Bioenergetics of a Symbiotic Coral. *BioScience* **34**, 705–709 (1984).

10. J. P. Gattuso, D. Allemand, M. Frankignoulle, Photosynthesis and calcification at cellular, organismal and community levels in coral reefs: A review on interactions and control by carbonate chemistry. *American Zoologist* **39**, 160–183 (1999).

11. S. Enríquez, E. R. Méndez, R. Iglesias-Prieto, S. Enrı, E. R. Me, Multiple scattering on coral skeletons enhances light absorption by symbiotic algae. *Limnology and Oceanography* **50**, 1025–1032 (2005).

12. J. W. Porter, L. Muscatine, Z. Dubinsky, P. G. Falkowski, Primary Production and Photoadaptation in Light- and Shade-Adapted Colonies of the Symbiotic Coral, Stylophora pistillata. *Proceedings of the Royal Society of London. Series B, Biological Sciences* **222**, 161–180 (1984).

13.   R. K. Trench, Microalgal-Invertebrate symbioses : a Review. *Endocytobiosis & Cell*
Research **9**, 135–175 (1993).

14.   R. Iglesias-Prieto, R. K. Trench, Acclimation and adaptation to irradiance in symbiotic
dinoflagellates. II. Response of chlorophyll-protein complexes to different photon-flux
densities. *Marine Biology* **130**, 23–33 (1997).

15.   W. K. Fitt, F. K. McFarland, M. E. Warner, G. C. Chilcoat, Seasonal patterns of tissue
biomass and densities of symbiotic dinoflagellates in reef corals and relation to coral
bleaching. *Limnology and Oceanography* **45**, 677–685 (2000).

16.   M. Y. Gorbunov, Z. S. Kolber, M. P. Lesser, P. G. Falkowski, Photosynthesis and
photoprotection in symbiotic corals. *Association for the Sciences of Limnology and*
Oceanography **46**, 75–85 (2001).

17.   E. A. Titlyanov, T. V. Titlyanova, K. Yamazato, R. van Woesik, Photo-acclimation
dynamics of the coral *Stylophora pistillata* to low and extremely low light. *Journal of*
Experimental Marine Biology and Ecology **263**, 211–225 (2001).

18.   K. R. N. Anthony, O. Hoegh-Guldberg, Kinetics of Photoacclimation in Corals. *Oecologia*
134, 23–31 (2003).

19.   P. R. Frade, P. Bongaerts, A. J. S. Winkelhagen, L. Tonk, R. P. M. Bak, In situ
photobiology of corals over large depth ranges: A multivariate analysis on the roles of
environment, host, and algal symbiont. *Limnology and Oceanography* **53**, 2711–2723
(2008).

20.   S. J. Hennige, D. J. Smith, R. Perkins, M. Consalvey, D. M. Paterson, D. J. Suggett,
Photoacclimation, growth and distribution of massive coral species in clear and turbid
waters. *Marine Ecology Progress Series* **369**, 77–88 (2008).

21.   P. A. Todd, Morphological plasticity in scleractinian corals. *Biological Reviews* **83**, 315–
337 (2008).

22.   M. P. Lesser, S. Marc, S. Michael, O. Michiko, R. D. Gates, G. Andrea,
Photoacclimatization by the coral Montastraea cavernosa in the mesophotic zone: Light,
food, and genetics. *Ecology* **91**, 990–1003 (2010).

23.   M. S. Roth, The engine of the reef: Photobiology of the coral-algal symbiosis. *Frontiers in*
Microbiology **5** (2014).

24.   S. Enríquez, E. R. Méndez, O. Hoegh-Guldberg, R. Iglesias-Prieto, Key functional role of
the optical properties of coral skeletons in coral ecology and evolution. *Proceedings of the*
Royal Society B: Biological Sciences **284**, 20161667 (2017).

25.   K. E. Lohr, E. F. Camp, U. Kuzhiumparambil, A. Lutz, W. Leggat, J. T. Patterson, D. J.
Suggett, Resolving coral photoacclimation dynamics through coupled photophysiological
and metabolomic profiling. *J Exp Biol* **222**, jeb195982 (2019).

26.   A. Malik, S. Einbinder, S. Martinez, D. Tchernov, S. Haviv, R. Almuly, P. Zaslansky, I.
Polishchuk, B. Pokroy, J. Stolarski, T. Mass, Molecular and skeletal fingerprints of

scleractinian coral biomineralization: From the sea surface to mesophotic depths. *Acta*
Biomaterialia **120**, 263–276 (2021).

27. M. O. Hoogenboom, S. R. Connolly, K. R. N. Anthony, Interactions Between
Morphological and Physiological Plasticity Optimize Energy Acquisition in Corals.
Ecology **89**, 1144–1154 (2008).

28. Y. X. Ow, P. A. Todd, Light-induced morphological plasticity in the scleractinian coral
Goniastrea pectinata and its functional significance. *Coral Reefs* **29**, 797–808 (2010).

29. N. Kramer, R. Tamir, O. Ben-Zvi, S. L. Jacques, Y. Loya, D. Wangpraseurt, Efficient light-
harvesting of mesophotic corals is facilitated by coral optical traits. *Functional Ecology* **36**,
406–418 (2022).

30. K. Gomez-Campo, R. Sanchez, I. Martinez-Rugerio, X. Yang, T. Maher, C. C. Osborne, S.
Enriquez, I. B. Baums, S. A. Mackenzie, R. Iglesias-Prieto, Phenotypic plasticity for
improved light harvesting, in tandem with methylome repatterning in reef-building corals.
Molecular Ecology **33**, e17246 (2024).

31. P. Hallock, Algal symbiosis: A mathematical analysis. *Mar. Biol.* **62**, 249–255 (1981).

32. M. O. Hoogenboom, K. R. N. Anthony, S. R. Connolly, Energetic cost of photoinhibition
in corals. *Marine Ecology Progress Series* **313**, 1–12 (2006).

33. D. H. Wright, Species-Energy Theory: An Extension of Species-Area Theory. *Oikos* **41**,
496–506 (1983).

34. R. De Visser, C. J. T. Spitters, T. J. Bouma, "Energy cost of protein turnover: theorectical
calculation and experimental estimation from regression of respiration on protein
concentration of full-grown leaves" in *Molecular, Biochemical and Physiological Aspects*
of Plant Respiration (1992), pp. 493–508.

35. T. J. Bouma, R. De Visser, J. H. J. A. Janssen, M. J. De Kock, P. H. Van Leeuwen, H.
Lambers, Respiratory energy requirements and rate of protein turnover in vivo determined
by the use of an inhibitor of protein synthesis and a probe to assess its effect. *Physiologia*
Plantarum **92**, 585–594 (1994).

36. A. Quigg, J. Beardall, Protein turnover in relation to maintenance metabolism at low
photon flux in two marine microalgae. *Plant, Cell and Environment* **26**, 693–703 (2003).

37. L. Li, C. J. Nelson, J. Trösch, I. Castleden, S. Huang, A. H. Millar, Protein degradation rate
in Arabidopsis thaliana leaf growth and development. *Plant Cell* **29**, 207–228 (2017).

38. G. Schuster, R. Timberg, I. Ohad, Turnover of thylakoid photosystem II proteins during
photoinhibition of Chlamydomonas reinhardtii. *European Journal of Biochemistry* **177**,
403–410 (1988).

39. E. M. Aro, S. McCaffery, J. M. Anderson, Photoinhibition and D1 Protein Degradation in
Peas Acclimated to Different Growth Irradiances. *Plant physiology* **103**, 835–843 (1993).

869 40. E.-M. Aro, I. Virgin, B. Andersson, Photoinhibition of Photosystem II. Inactivation, protein
870   damage and turnover. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1143**, 113–134
871   (1993).

872 41. S. Järvi, M. Suorsa, E. M. Aro, Photosystem II repair in plant chloroplasts--Regulation,
873   assisting proteins and shared components with photosystem II biogenesis. *Biochimica et*
874   *biophysica acta* **1847**, 900–909 (2015).

875 42. P. Chotewutmontri, A. Barkan, Dynamics of Chloroplast Translation during Chloroplast
876   Differentiation in Maize. *PLOS Genetics* **12**, e1006106 (2016).

877 43. L. Li, E. M. Aro, A. H. Millar, Mechanisms of Photodamage and Protein Turnover in
878   Photoinhibition. *Trends in Plant Science* **23**, 667–676 (2018).

879 44. K. Miyata, K. Noguchi, I. Terashima, Cost and benefit of the repair of photodamaged
880   photosystem II in spinach leaves: Roles of acclimation to growth light. *Photosynthesis*
881   *Research* **113**, 165–180 (2012).

882 45. M. Tikkanen, N. R. Mekala, E. M. Aro, Photosystem II photoinhibition-repair cycle
883   protects Photosystem i from irreversible damage. *Biochimica et Biophysica Acta -*
884   *Bioenergetics* **1837**, 210–215 (2014).

885 46. S. Järvi, J. Isojärvi, S. Kangasjärvi, J. Salojärvi, F. Mamedov, M. Suorsa, E.-M. Aro,
886   Photosystem II Repair and Plant Immunity: Lessons Learned from Arabidopsis Mutant
887   Lacking the THYLAKOID LUMEN PROTEIN 18.3. *Front. Plant Sci.* **7** (2016).

888 47. N. Murata, Y. Nishiyama, ATP is a driving force in the repair of photosystem II during
889   photoinhibition. *Plant, Cell & Environment* **41**, 285–299 (2018).

890 48. S. Takahashi, T. Nakamura, M. Sakamizu, R. Van Woesik, H. Yamasaki, Repair nachinery
891   of symbiotic photosynthesis as the primary target of heat stress for reef-building corals.
892   *Plant and Cell Physiology* **45**, 251–255 (2004).

893 49. R. Hill, S. Takahashi, Photosystem II recovery in the presence and absence of chloroplast
894   protein repair in the symbionts of corals exposed to bleaching conditions. *Coral Reefs* **33**,
895   1101–1111 (2014).

896 50. S. Roberty, D. Fransolet, P. Cardol, J.-C. Plumier, F. Franck, Imbalance between oxygen
897   photoreduction and antioxidant capacities in Symbiodinium cells exposed to combined heat
898   and high light stress. *Coral Reefs* **34**, 1063–1073 (2015).

899 51. K. R. N. Anthony, P. V. Ridd, A. R. Orpin, P. Larcombe, J. Lough, Temporal variation of
900   light availability in coastal benthic habitats: Effects of clouds, turbidity, and tides.
901   *Limnology and Oceanography* **49**, 2201–2211 (2004).

902 52. S. DiPerna, M. Hoogenboom, S. Noonan, K. Fabricius, Effects of variability in daily light
903   integrals on the photophysiology of the corals Pachyseris speciosa and Acropora millepora.
904   *PLoS ONE* **13**, e0203882 (2018).

905 53. P. G. Falkowski, J. A. Raven, *Aquatic Photosynthesis* (Princeton University Press, 2014).

54. T. López-Londoño, C. T. Galindo-Martínez, K. Gómez-Campo, L. A. González-Guerrero,
S. Roitman, F. J. Pollock, V. Pizarro, M. López-Victoria, M. Medina, R. Iglesias-Prieto,
Physiological and ecological consequences of the water optical properties degradation on
reef corals. *Coral Reefs* **40**, 1243–1256 (2021).

55. R. Hill, F. Bendall, Function of the Two Cytochrome Components in Chloroplasts: A
Working Hypothesis. *Nature* **186**, 136–137 (1960).

56. J. F. Hill, Govindjee, The controversy over the minimum quantum requirement for oxygen
evolution. *Photosynth Res* **122**, 97–112 (2014).

57. W. Skirving, S. Enríquez, J. D. Hedley, S. Dove, C. M. Eakin, R. A. B. Mason, J. L. D. L.
Cour, G. Liu, O. Hoegh-Guldberg, A. E. Strong, P. J. Mumby, R. Iglesias-Prieto, Remote
sensing of coral bleaching using temperature and light: Progress towards an operational
algorithm. *Remote Sensing* **10**, 18 (2018).

58. L. Muscatine, P. G. ; Falkowski, J. W. ; Porter, Z. Dubinsky, Fate of photosynthetic fixed
carbon in light- and shade-adapted colonies of the symbiotic coral Stylophora pistillata.
Proceedings of the Royal Society of London. Series B. Biological Sciences **222**, 181–202
(1984).

59. L. Muscatine, Symbiosis of hydra and algae—III. Extracellular products of the algae.
Comparative Biochemistry and Physiology **16**, 77–92 (1965).

60. L. Muscatine, Glycerol excretion by symbiotic algae from corals and Tridacna and its
control by the host. *Science* **156**, 516–519 (1967).

61. M. S. Burriesci, T. K. Raab, J. R. Pringle, Evidence that glucose is the major transferred
metabolite in dinoflagellate–cnidarian symbiosis. *Journal of Experimental Biology* **215**,
3467–3477 (2012).

62. R. K. Trench, The Physiology and Biochemistry of Zooxanthellae Symbiotic with Marine
Coelenterates. II. Liberation of Fixed 14C by Zooxanthellae in vitro. *Proceedings of the
Royal Society of London. Series B, Biological Sciences* **177**, 237–250 (1971).

63. R. K. Trench, The Physiology and Biochemistry of Zooxanthellae Symbiotic with Marine
Coelenterates. I. The Assimilation of Photosynthetic Products of Zooxanthellae by Two
Marine Coelenterates. *Proceedings of the Royal Society of London. Series B, Biological
Sciences* **177**, 225–235 (1971).

64. P. Tremblay, R. Grover, J. F. Maguer, M. Hoogenboom, C. Ferrier-Pagès, Carbon
translocation from symbiont to host depends on irradiance and food availability in the
tropical coral Stylophora pistillata. *Coral Reefs* **33**, 1–13 (2014).

65. S. Sathyendranath, T. Platt, Ž. Kovač, J. Dingle, T. Jackson, R. J. W. Brewin, P. Franks, E.
Marañón, G. Kulk, H. A. Bouman, Reconciling models of primary production and
photoacclimation [Invited]. *Applied Optics* **59**, C100 (2020).

66. R. D. Gates, P. J. Edmunds, The physiological mechanisms of acclimatization in tropical
reef corals. *American Zoologist* **39**, 30–43 (1999).

67. M. O. Hoogenboom, S. R. Connolly, K. R. N. Anthony, Effects of photoacclimation on the
light niche of corals: A process-based approach. *Marine Biology* **156**, 2493–2503 (2009).

68. B. Demmig-Adams, W. W. Adams, The role of xanthophyll cycle carotenoids in the
protection of photosynthesis. *Trends in Plant Science* **1**, 21–26 (1996).

69. B. E. Brown, I. Ambarsari, M. E. Warner, W. K. Fitt, R. P. Dunne, S. W. Gibb, D. G.
Cummings, Diurnal changes in photochemical efficiency and xanthophyll concentrations in
shallow water reef corals: Evidence for photoinhibition and photoprotection. *Coral Reefs*
18, 99–105 (1999).

70. P. J. Ralph, R. Gademann, A. W. D. Larkum, U. Schreiber, In situ underwater
measurements of photosynthetic activity of coral zooxanthellae and other reef-dwelling
dinoflagellate endosymbionts. *Marine Ecology Progress Series* **180**, 139–147 (1999).

71. E. Erickson, S. Wakao, K. K. Niyogi, Light stress and photoprotection in Chlamydomonas
reinhardtii. *Plant Journal*, 449–465 (2015).

72. R. J. Jones, O. Hoegh-Guldberg, Diurnal changes in the photochemical efficiency of the
symbiotic dinoflagellates (Dinophyceae) of corals: photoprotection, photoinactivation and
the relationship to coral bleaching. *Plant, Cell & Environment* **24**, 89–99 (2001).

73. M. P. Lesser, M. Y. Gorbunov, Diurnal and bathymetric changes in chlorophyll
fluorescence yields of reef corals measured in situ with a fast repetition rate fluorometer.
212, 69–77 (2001).

74. G. Winters, Y. Loya, R. Röttgers, S. Beer, Photoinhibition in shallow-water colonies of the
coral Stylophora pistillata as measured in situ. *Limnology and Oceanography* **48**, 1388–
1393 (2003).

75. O. Levy, Z. Dubinsky, K. Schneider, Y. Achituv, D. Zakai, M. Y. Gorbunov, Diurnal
hysteresis in coral photosynthesis. *Marine Ecology Progress Series* **268**, 105–117 (2004).

76. A. Rodríguez-Román, X. Hernández-Pech, P. E. Thomé, S. Enríquez, R. Iglesias-Prieto, X.
Herna, P. E. Thome, S. Enrí, R. Iglesias-Prieto, Photosynthesis and light utilization in the
Caribbean coral Montastraea faveolata recovering from a bleaching event. *Limnology and*
Oceanography **51**, 2702–2710 (2006).

77. T. Scheufen, R. Iglesias-Prieto, S. Enríquez, Changes in the number of symbionts and
Symbiodinium cell pigmentation modulate differentially coral light absorption and
photosynthetic performance. *Frontiers in Marine Science* **4**, 309 (2017).

78. T. Scheufen, W. E. Krämer, R. Iglesias-Prieto, S. Enríquez, Seasonal variation modulates
coral sensibility to heat-stress and explains annual changes in coral productivity. *Scientific*
Reports **7**, 4937 (2017).

79. N. Kramer, R. Tamir, G. Eyal, Y. Loya, Coral Morphology Portrays the Spatial
Distribution and Population Size-Structure Along a 5–100 m Depth Gradient. *Front. Mar.*
Sci. **7** (2020).

80. J. P. Krall, G. E. Edwards, Relationship between photosystem II activity and CO2 fixation
in leaves. *Physiologia Plantarum* **86**, 180–187 (1992).

81. M. J. Fryer, J. R. Andrews, K. Oxborough, D. A. Blowers, N. R. Baker, Relationship between CO2 Assimilation, Photosynthetic Electron Transport, and Active O2 Metabolism in Leaves of Maize in the Field during Periods of Low Temperature1. *Plant Physiology* **116**, 571–580 (1998).

82. O. Hoegh-Guldberg, R. Jones, Photoinhibition and photoprotection in symbiotic dinoflagellates from reef-building corals. *Mar. Ecol. Prog. Ser.* **183**, 73–86 (1999).

83. M. Lesser, Depth-dependent photoacclimatization to solar ultraviolet radiation in the Caribbean coral Montastraea faveolata. *Mar. Ecol. Prog. Ser.* **192**, 137–151 (2000).

84. J. A. Raven, The cost of photoinhibition. *Physiologia Plantarum* **142**, 87–104 (2011).

85. X.-P. Yi, H.-S. Yao, D.-Y. Fan, X.-G. Zhu, P. Losciale, Y.-L. Zhang, W.-F. Zhang, W. S. Chow, The energy cost of repairing photoinactivated photosystem II: an experimental determination in cotton leaf discs. *New Phytologist* **235**, 446–456 (2022).

86. S. Gunell, T. Lempiäinen, E. Rintamäki, E.-M. Aro, M. Tikkanen, Enhanced function of non-photoinhibited photosystem II complexes upon PSII photoinhibition. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1864**, 148978 (2023).

87. C. Langdon, On the causes of interspecific differences in the growth-irradiance relationship for phytoplankton. II. A general review. *Journal of Plankton Research* **10**, 1291–1312 (1988).

88. N. Murata, S. Takahashi, Y. Nishiyama, S. I. Allakhverdiev, Photoinhibition of photosystem II under environmental stress. *Biochimica et Biophysica Acta - Bioenergetics* **1767**, 414–421 (2007).

89. S. I. Allakhverdiev, Y. Nishiyama, S. Takahashi, S. Miyairi, I. Suzuki, N. Murata, Systematic Analysis of the Relation of Electron Transport and ATP Synthesis to the Photodamage and Repair of Photosystem II in Synechocystis. *Plant Physiology* **137**, 263–273 (2005).

90. R. Iglesias-prieto, R. K. Trench, Acclimation and adaptation to irradiance in symbiotic dinoflagellates. 1.Responses of the photosynthetic unit to changes in photon flux density. *Marine Ecology Progress Series* **113**, 163–176 (1994).

91. A. Trebst, Function of β-Carotene and Tocopherol in Photosystem II. *Zeitschrift für Naturforschung C* **58**, 609–620 (2003).

92. A. Krieger-Liszkay, Singlet oxygen production in photosynthesis. *J Exp Bot* **56**, 337–346 (2005).

93. M. E. Warner, W. K. Fitt, G. W. Schmidt, Damage to photosystem II in symbiotic dinoflagellates: a determinant of coral bleaching. *Proceedings of the National Academy of Sciences of the United States of America* **96**, 8007–12 (1999).

94. S. Takahashi, S. Whitney, S. Itoh, T. Maruyama, M. Badger, Heat stress causes inhibition of the de novo synthesis of antenna proteins and photobleaching in cultured Symbiodinium. *Proceedings of the National Academy of Sciences of the United States of America* **105**, 4203–8 (2008).

95. S. Takahashi, M. Yoshioka-Nishimura, D. Nanba, M. R. Badger, Thermal Acclimation of the Symbiotic Alga Symbiodinium spp. Alleviates Photobleaching under Heat Stress. *PLANT PHYSIOLOGY* **161**, 477–485 (2013).

96. R. Hill, C. M. Brown, K. DeZeeuw, D. a Campbell, P. J. Ralph, Increased rate of D1 repair in coral symbionts during bleaching is insufficient to counter accelerated photo-inactivation. *Limnology and Oceanography* **56**, 139–146 (2011).

97. S. Takahashi, S. E. Milward, W. Yamori, J. R. Evans, W. Hillier, M. R. Badger, The solar action spectrum of photosystem II damage. *Plant Physiology* **153**, 988–993 (2010).

98. J. M. Pandolfi, C. E. Lovelock, A. F. Budd, Character Release Following Extinction in a Caribbean Reef Coral Species Complex. *Evolution* **56**, 479–501 (2002).

99. C. Prada, M. Gómez-Corrales, A. González, B. Kamel, M. Medina, N. Knowlton, D. Levitan, Speciation across depth gradients in reef corals. In Review [Preprint] (2024). https://doi.org/10.21203/rs.3.rs-4202515/v1.

100. H. E. Aichelman, B. E. Benson, K. Gomez-Campo, M. I. Martinez-Rugerio, J. E. Fifer, L. Tsang, A. M. Hughes, C. B. Bove, O. C. Nieves, A. M. Pereslete, D. Stanizzi, N. G. Kriefall, J. H. Baumann, J. P. Rippe, P. Gondola, K. D. Castillo, S. W. Davies, Cryptic coral diversity is associated with symbioses, physiology, and response to thermal challenge. *Science Advances* **11**, eadr5237 (2025).

101. D. Iluz, I. Alexandrovich, Z. Dubinsky, The enhancement of photosynthesis by fluctuating light. *Artifical Photosynthesis* (2012).

102. J. E. Faust, V. Holcombe, N. C. Rajapakse, D. R. Layne, The effect of daily light integral on bedding plant growth and flowering. *HortScience* **40**, 645–649 (2005).

103. S. W. Jeffrey, G. F. Humphrey, New spectrophotometric equations for determining chlorophylls a, b, c1 and c2 in higher plants, algae and natural phytoplankton. *Biochemie und Physiologie der Pflanzen* **167**, 191–194 (1975).

104. R. M. Vásquez-Elizondo, L. Legaria-Moreno, M. Á. Pérez-Castro, W. E. Krämer, T. Scheufen, R. Iglesias-Prieto, S. Enríquez, Absorptance determinations on multicellular tissues. *Photosynthesis Research* **132**, 311–324 (2017).

105. S. Enríquez, K. Sand-Jensen, Variation in Light Absorption Properties of *Mentha aquatica* L. as a Function of Leaf Form: Implications for Plant Growth. *International Journal of Plant Sciences* **164**, 125–136 (2003).

106. N. M. Cayabyab, S. Enríquez, Leaf photoacclimatory responses of the tropical seagrass *Thalassia testudinum* under mesocosm conditions: a mechanistic scaling-up study. *New Phytologist* **176**, 108–123 (2007).

107. L. A. González-Guerrero, R. M. Vásquez-Elizondo, T. López-Londoño, G. Hernán, R. Iglesias-Prieto, S. Enríquez, Validation of parameters and protocols derived from chlorophyll a fluorescence commonly utilised in marine ecophysiological studies. *Functional Plant Biol.*, doi: 10.1071/FP21101 (2021).

108. W. C. Dennison, R. S. Alberte, Photosynthetic Responses of Zostera marina L. (Eelgrass)
to in situ Manipulations of Light Intensity. *Oecologia* **55**, 137–144 (1982).

109. M. K. DeSalvo, S. Sunagawa, P. L. Fisher, C. R. Voolstra, R. Iglesias-Prieto, M. Medina,
Coral host transcriptomic states are correlated with Symbiodinium genotypes. *Molecular*
Ecology **19**, 1174–1186 (2010).

110. D. W. Kemp, D. J. Thornhill, R. D. Rotjan, R. Iglesias-Prieto, W. K. Fitt, G. W. Schmidt,
Spatially distinct and regionally endemic Symbiodinium assemblages in the threatened
Caribbean reef-building coral Orbicella faveolata. *Coral Reefs* **34**, 535–547 (2015).

111. A. M. Lewis, A. N. Chan, T. C. LaJeunesse, New Species of Closely Related
Endosymbiotic Dinoflagellates in the Greater Caribbean have Niches Corresponding to
Host Coral Phylogeny. *Journal of Eukaryotic Microbiology* **66**, 469–482 (2019).

112. T. C. LaJeunesse, W. K. W. Loh, R. Van Woesik, O. Hoegh-Guldberg, G. W. Schmidt, W.
K. Fitt, Low symbiont diversity in southern Great Barrier Reef corals, relative to those of
the Caribbean. *Limnology and Oceanography* **48**, 2046–2054 (2003).

113. T. C. Lajeunesse, R. K. Trench, Biogeography of two species of Symbiodinium
(Freudenthal) inhabiting the intertidal sea anemone Anthopleura elegantissima (Brandt).
https://doi.org/10.2307/1542872 **199**, 126–134 (2016).

114. R. Zardoya, E. Costas, V. López-Rodas, A. Garrido-Pertierra, J. M. Bautista, Revised
dinoflagellate phylogeny inferred from molecular analysis of large-subunit ribosomal RNA
gene sequences. *Journal of Molecular Evolution* **41**, 637–645 (1995).

115. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv arXiv:1303.3997 [Preprint] (2013). https://doi.org/10.48550/arXiv.1303.3997.

116. V. Avila-Magaña, B. Kamel, M. DeSalvo, K. Gómez-Campo, S. Enríquez, H. Kitano, R. V.
Rohlfs, R. Iglesias-Prieto, M. Medina, Elucidating gene expression adaptation of
phylogenetically divergent coral holobionts under heat stress. *Nat Commun* **12**, 5731
(2021).

117. E. F. Camp, T. Kahlke, B. Signal, C. A. Oakley, A. Lutz, S. K. Davy, D. J. Suggett, W. P.
Leggat, Proteome metabolome and transcriptome data for three Symbiodiniaceae under
ambient and heat stress conditions. *Sci Data* **9**, 153 (2022).

118. R. A. Levin, V. H. Beltran, R. Hill, S. Kjelleberg, D. McDougald, P. D. Steinberg, M. J. H.
van Oppen, Sex, Scavengers, and Chaperones: Transcriptome Secrets of Divergent
Symbiodinium Thermal Tolerances. *Mol Biol Evol* **33**, 2201–2215 (2016).

119. C. M. Brown, D. A. Campbell, J. E. Lawrence, Resource dynamics during infection of
Micromonas pusilla by virus MpV-Sp1. *Environmental Microbiology* **9**, 2720–2727
(2007).

120. C. Six, Z. V. Finkel, A. J. Irwin, D. A. Campbell, Light Variability Illuminates Niche-
Partitioning among Marine Picocyanobacteria. *PLoS ONE* **2**, e1341 (2007).

121. P. Ewels, M. Magnusson, S. Lundin, M. Käller, MultiQC: summarize analysis results for
multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

122. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

123. N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol* **34**, 525–527 (2016).

124. L. Fu, B. Niu, Z. Zhu, S. Wu, W. Li, CD-HIT: accelerated for clustering the next-
generation sequencing data. *Bioinformatics* **28**, 3150–3152 (2012).

125. C. Camacho, G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, T. L. Madden,
BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421 (2009).

126. J. Kirk, *Light and Photosynthesis* (Cambridge University Press, New York, NY, 2011;
https://www.cambridge.org/core/books/light-and-photosynthesis-in-aquatic-
ecosystems/C19B28AE07B1CDEBDA5593194DE4E304).

127. A. D. Jassby, T. Platt, Mathematical formulation of the relationship between photosynthesis
and light for phytoplankton. *Limnology and Oceanography* **21**, 540–547 (1976).

1112

1113
Acknowledgments

We thank UNAM and ICMyL for staff support, facility assistance, SAMMO for data, S Enriquez for laboratory space and mesocosm facility. V Avila-Magana, CT Galindo-Martinez, NS Locatelli, CC Osborne for assistance in laboratory work. T López-Londono for assistance in modeling. VA Gómez-Chavez for Figure S1. Collection permit No. PPF/DGOPA-020/2018 and CITES permit No. MX96989, MX102782 to RIP and SE.

Funding:

This project has received funding from The Pennsylvania State University Start-up (RIP). NSF grant (NSF OCE-1537959) to I.B.B.

Author contributions:

Conceptualization: RIP, KGC.

Data curation: KGC.

Formal analysis: KGC and RIP.

Funding acquisition: RIP, SE, TCL, IBB.

Investigation and methodology: KGC performed the experiments analyze and integrated data. IMR contributed to fieldwork, phenotypic descriptions and light systems. MAGR engineered the light system. RGP to transcriptomic computational work and interpretations. MGS to translational laboratory work and interpretations. KET to algal identification laboratory work and interpretations. TLC and IBB to analysis and interpretations.

Project administration: RIP.

Resources: RIP, SE, TCL, IBB, MGSM.

Software: MAGR.

Supervision: RIP.

Validation: KGC, RIP.

Visualization: KGC, RIP.

Writing - original draft: KGC, RIP.

Writing - review & editing: KGC, IMR, MAGR, RGP, MGSM, KET, TCL, IBB, SE, RIP.

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data will be available in the main text or the supplementary materials. All detailed data, code, and materials used in the analyses will be available for purposes of reproducing or extending the analyses.

1151
1152   **Figures and Tables**
1153
1154

1155
Fig. 1. Phenotyping after acclimation to light conditions. (**A**) The general equation of
light attenuation in seawater ($DLI_{(z)} = DLI_{(o)} \times e^{(-Kd \times z)}$; Eq. 1) maps the four
experimental DLI conditions to their corresponding depths (5–35 m, assuming K_d
= 0.07 m^{-1}), illustrated alongside a depth gradient. (**B**) Maximum photochemical
efficiency (F_v/F_m at dusk) as a function of DLI after 4+ weeks of acclimation
illustrating a linear trend. Adjustments in photoprotection shown as (**C**) diurnal
changes in photochemical efficiency (maximum F_v/F_m at dawn/dusk, and effective
$\Delta F/F_m{}'$ during daylight) and (**D**) non-photochemical quenching (*NPQ*) measured
hourly across light conditions. Symbols represent mean ± SE (n = 16). Shaded
areas show bootstrapped 95% confidence intervals.

Fig. 2. Optical reorganization and metabolic rates for light harvesting across light conditions. (**A**) Estimated absorbance (*De*) from reflectance measurements indicating reduced absorption under high light (n = 16) and (photo) qualitative changes in tissue pigmentation. (**B**) Pigment-specific absorption coefficient (a*$_{Chla}$) showing the non-linear increase in absorption efficiency in the coral tissue as pigment content declines, consistent with the package effect. (**C**) Photosynthetic efficiency (α), (**D**) maximum photosynthetic rate (*P$_{max}$* normalized to host area, and *P$_{sym}$* normalized to symbiont cell) showing no significant changes in *P$_{max}$* across light regimes (n = 8). (**E**) Dark respiration rates (*R$_D$*) (n = 8), and (**F**) maximum photochemical efficiency of PSII (*F$_v$/F$_m$*) as a function of the minimum quantum requirement for photosynthesis (MQR, Φ^{-1}; mol photons mol^{-1} O$_2$). Lower MQR values indicate higher photosynthetic efficiency or fewer photons required per molecule of O$_2$ evolved. DLI-4 corals achieved MQR values approaching the theoretical minimum (~8 quanta O$_2^{-1}$), indicating near-maximal photosynthetic efficiency and delineating the lower boundary of photoacclimatory capacity for the symbiosis. With increasing irradiance (toward DLI-32), both *F$_v$/F$_m$* and quantum efficiency declined, reflecting the increasing energetic cost of sustaining photosynthesis under high light. Shaded areas show bootstrapped 95% confidence intervals; symbols are means ± SE.

1187

Fig. 3. **Diurnal PSII activity and PSII half-life.** (**A**) PSII photochemical efficiency
measured hourly across a 12 h photoperiod in the presence (+CAP; empty circles)
and absence (−CAP; filled circles) of the protein synthesis inhibitor
chloramphenicol. Blocking *de novo* protein synthesis revealed net PSII damage
accumulation, while control samples showed oscillations with characteristic daily
photodamage and repair cycles. (**B**) Relative photochemical efficiency (+CAP/-
CAP) decay showing proportional rate of photodamage with increased light
intensity, reflecting the rate of PSII inactivation in the absence of repair. Dashed
lines illustrate the time of day when 50% of PSII reaction centers are photo-
inactivated. In DLI-4 PSII $t_{1/2}$ exceeded the 12 h photoperiod. Curves show four-
parameter logistic fits (mean ± 95% CI).

Fig. 4. PSII regulation in zooxanthellate corals across light conditions. (**A**) PSII large
pigment–protein complex with its core reaction center D1 (encoded by *psbA* gene).
D1 provides key ligands to the oxygen-evolving complex, harbors the QB
plastoquinone binding site, and is the primary target of photodamage. (**B**) Power
law decay of PSII half-life ($t_{1/2}$, h) with increasing DLI. (**C**) PSII turnover rate
(day^{-1}) accelerated significantly under higher DLI. (**D**) Quantitative immunoblot of
the D1 protein, showing patterns across light conditions. Lanes include a negative
control (lane1, 0.00 pmol), quantification standards (lanes2–6, 0.10-2.00 pmol),
and samples loaded by total protein (lanes7–10, total protein^{-1}) and Chl*a* content
(11–14, mg Chl*a*$^{-1}$). Quantification of D1 protein normalized to total protein (**E**;
pmol D1 protein μg total protein^{-1}) and Chl*a* (**F**; pmol D1 protein mg Chl*a*$^{-1}$),
showing a similar power law decay with light exposure (n = 3 immunoblots). (**G**)
Transcriptional abundance (transcripts per million; TPM) of PSII complex genes
(*psbA, psbB, psbC, psbD*). Transcript levels showed no systematic relationship
with DLI, in contrast to the exponential decline in D1 protein abundance shown in
panels E and F, indicating post-transcriptional regulation. (**H**) PSII half-life ($t_{1/2}$, h)
when acclimated corals (DLI) were changed to an alternative light condition (DLI-
destination). Lines show a power law fit to experimental PAR (black line and
colored circles; this study) and natural sunlight measurements (blue line and grey
circles; data from (*5*)). Shaded areas show bootstrapped 95% confidence intervals;
symbols are means ± SE. (**I**) Photodamage accumulation after the change in light
condition. Results are shown as the relative change in F_v/F_m (*relative-F_v/F_m* =
F_v/F_m-destination / F_v/F_m-acclimation) when coral fragments were exposed to *EEE*
(*EEE* = DLI-destination - DLI-acclimation); the linear prediction is consistent with
prior findings (*57*).

1227
1228

1229
Fig. 5. Bio-optical model of *photosynthetic usable energy supply* (*PUES*). (**A**) Modeled
profiles of *PUES* across depth for a range of vertical attenuation coefficients of
downwelling irradiance (K_d = 0.04–0.40 m^{-1}), representing varying water column
optical properties. *PUES* represents the photosynthetic energy available to the
coral host after subtracting the depth-dependent energetic cost of PSII repair. Each
curve represents a distinct K_d value, color-coded from clear to turbid conditions.
(**B**) Optimal depth (*Depth$_{opt}$*) at which *PUES* is maximized, as a function of Kd.
This bio-optical state represents the depth at which the balance between
photosynthetic output and PSII repair cost is optimized, maximizing the energy
potentially available for host metabolic demands.
1240

Table 1. Table of terms. Parameters used to characterize coral holobiont phenotypes.

Descriptors	Parameters and notations	Units
PSII	Maximum photochemical efficiency (F_v/F_m)	Dimensionless
	Effective photochemical efficiency ($\Delta F/Fm'$)	Dimensionless
	Maximum excitation pressure over PSII (Q_m)	Dimensionless
	PII half-life ($PSII_{1/2}$)	Hours
	PSII D1 protein abundance per protein	pmol D1 µg total protein^{-1}
	PSII D1 protein abundance per Chl$a$	pmol D1 µg Chla^{-1}
	PII-complex transcript abundance	Transcripts per million
Structural and optical traits	Chl$a$ density per area	mg Chl$a$ m^{-2}
	Chl$a$ specific absorption coefficient (a*$_{Chla}$)	m^2 mg Chla^{-1}
	Algal symbionts per area of host tissue	Cells cm^2
	Chl$a$ per per cell (Ci)	Pg Chl$a$ sym^{-1}
	Absorptance (A)	Dimensionless
	Absorbance (De 675 nm)	Dimensionless
	Chl$a$ specific absorption coefficient (a*$_{Chla}$)	m^2 mg Chla^{-1}
	Symbiont specific absorption coefficient (a*$_{Sym}$)	m^2 sym^{-1}
Metabolic rates and P efficiency	Photosynthetic efficiency (α)	mol O$_2$ mol incident quanta^{-1}
	Quantum yield of photosyntheis (Φ_{O2})	mol O$_2$ mol absorbed quanta^{-1}
	Minimum Quantum Requirement ($1/\Phi_{O2}$)	photons absorbed per O$_2$ evolved
	Maximum photosynthetic rate ($Pmax$)	µmol O$_2$ m^{-2} s^{-1}
	Saturation irradiance (E_k)	µmol quanta m^{-2} s^{-1}
	Compensation irradiance (E_c)	µmol quanta m^{-2} s^{-1}
	Dark respiration (R_D)	µmol O$_2$ m^{-2} s^{-1}
	Light-enhanced respiration (R_L)	µmol O$_2$ m^{-2} s^{-1}
	Photosynthetic rate per symb cell ($Psym$)	µmol O$_2$ m^{-2} s^{-1}