1 **Gene Expansion and Regulatory Rewiring Shape Sex-Biased Evolution of**
2 **the Mouse Submandibular Gland Secretome**

3 **AUTHORS**
4 Luane Jandira Bueno Landau[1], Shikha Jain[2], Nathan Griffin[3], Achisha Saikia[4], Jill M.
5 Kramer[2], Sarah Knox[3], Stefan Ruhl[2], Omer Gokcumen[1]*
6
AFFILIATIONS
[1]Department of Biological Sciences, University at Buffalo, The State University of
New York, NY 14260, USA.
[2]Department of Oral Biology, School of Dental Medicine, University at Buffalo, The
State University of New York, NY, 14214, USA.
[3]Department of Cell & Tissue Biology, University of California, San Francisco, CA,
94143, USA.
[4]Department of Genetics, Genomics and Bioinformatics, Jacobs School of Medicine
and Biomedical Sciences, University at Buffalo, The State University of New York,
NY 14203, USA.
17
18 *correspondence:
19 Omer Gokcumen: omergokc@buffalo.edu

ABSTRACT

Mammalian saliva plays essential roles in digestion, immunity, and host-microbiome interactions, yet its protein composition varies across species and sexes. The evolutionary mechanisms underlying this molecular diversity remain poorly understood. Here, we compared mouse and human salivary gland secretomes at genomic, transcriptomic, and proteomic levels to understand how saliva composition evolves. We performed RNA-seq analysis of the major mouse salivary glands (parotid, submandibular, and sublingual), liver and pancreas from both sexes, compared them with reanalyzed previously published human salivary gland transcriptomes, and integrated them with proteome data of mouse and human whole saliva. We found that evolution of gene expression in mouse salivary glands is driven by rapid gene turnover and sexual dimorphism. In the submandibular and sublingual glands, respectively, 68% and 73% of expression from genes encoding secreted proteins derives from lineage-specific genes that lack one-to-one human orthologs. Mouse submandibular gland shows striking sexual dimorphism, with 1537 tissue specific sex-biased genes, five times higher than in the liver, a classic model of sex-biased expression. These genes cluster in regions shaped by recent gene duplication, such as the kallikrein gene cluster, a mouse-specific expansion that accounts for ~16.4% of male-biased submandibular expression. Our analyses suggest that this bias arises through regulatory changes that are expanded by gene duplication, including the spread of a testosterone-associated regulatory motif and the expansion of a shared chromatin domain that promotes coordinated gene regulation. Our results reveal how lineage-specific gene duplication and regulatory rewiring drive rapid, sex-specific evolution of the mammalian salivary gland secretome.

INTRODUCTION

Diet and pathogen exposure are major environmental drivers of mammalian evolution (Price et al. 2012; Enard et al. 2016). Because the oral cavity represents the body's first interface with the external environment, saliva plays a central role in mediating these pressures, balancing protection against pathogens with responsiveness to dietary variation. Saliva also contributes to oral homeostasis by lubricating and cleansing the mouth, protecting tooth enamel, enabling taste perception, and initiating digestion (Ruhl 2012; Pajic et al. 2022). These functions are carried out by a complex mixture of proteins secreted primarily by the three major salivary glands, parotid, submandibular, and sublingual glands (Dawes et al. 2015; Chibly et al. 2022). Importantly, the salivary gland secretome, i.e., the entirety of proteins secreted by the salivary glands, reflects adaptive tuning to species-specific ecological contexts, with protein repertoires shaped by dietary specialization (Pajic et al. 2019) and pathogen exposure (Xu et al. 2016). Together, the integration of immune, digestive, and sensory functions makes saliva an exceptional system for investigating how environmental pressures drive the evolution of mammalian genomes.

Previous studies suggested that saliva-related genes undergo rapid evolution, mainly to accommodate environmental and dietary changes which accompany speciation (Pajic et al. 2019). Recent locus-specific comparative genomic studies showed that genes expressed in the salivary glands that encode secreted salivary proteins differ substantially between mice and humans (Pajic et al. 2025). For instance, analyses of the secretory calcium-binding phosphoprotein (SCPP) locus indicated that multiple human salivary genes lack orthologs in mice. Proteomic analyses also have revealed that the composition of saliva proteins differs markedly between humans and mice (Blanchard et al. 2015; De Sousa□Pereira et al. 2015), and comparative studies further show substantial interspecies differences in their major components (Karn et al. 2013). These findings suggest that human and mouse salivary glands differ greatly in both gene content and expression profiles, supporting the view that the salivary gland secretome evolves rapidly in response to environmental and pathogenic pressures. Despite these insights, a systematic, secretome-wide investigation of the evolutionary dynamics and regulatory mechanisms underlying these differences has not yet been conducted.

In contrast to humans, the salivary glands in many other mammalian species are different between sexes. Sexual dimorphism in salivary glands has been well documented in rodents (Ikemoto and Matsushima 1984), and pronounced sex differences have been shown in salivary glands of mice (Chai et al. 1993), which are not known to occur in humans. Generally, sex-biased genes across multiple tissues in mammals are poorly conserved across species (Naqvi et al. 2019). In particular, the mouse submandibular gland exhibits striking sex-specific differences in cellular composition, secretory activity, and hormonal regulation (Hatakeyama et al. 1987;

Konttinen et al. 2010; Mukaibo et al. 2019). Although substantial progress has been
made in understanding saliva composition and its role in oral biology in both species,
the evolutionary basis of lineage- and sex-specific variation in the salivary gland
secretome remains underexplored.

Here, we investigate the mechanisms underlying the rapid diversification of the
salivary gland secretome. Using comparative analyses between humans and mice,
we assess the conservation of genes highly expressed in salivary glands and
determine the contribution of lineage-specific genes to the salivary gland secretome.
We further compare male and female mice to identify the genomic basis of sex-
biased expression in the submandibular gland. By integrating transcriptomic,
proteomic, and genomic data, we delineate the evolutionary processes driving
salivary diversity and sexual dimorphism.

RESULTS

The highest proportion of gene expression in mouse submandibular and sublingual glands derives from lineage-specific genes

To investigate differences in salivary gene expression across commonly used mouse models and compare them to humans, we performed transcriptomic profiling of the parotid, submandibular, and sublingual glands, along with the pancreas and liver as reference tissues, in CD1 and C57BL/6 strains (n = 3 per sex per strain; **Tables S1**). Both CD1 and C57BL/6 strains are widely used in laboratory studies of salivary glands, and were included as a representative of an inbred and outbred mouse strain, respectively, to strengthen the broader applicability of our findings (Nguyen et al. 2009; Gervais et al. 2015; Hsieh et al. 2017). In parallel, we conducted mass spectrometry-based proteomic profiling of whole saliva from ten C57BL/6 mice, five males and five females (**Table S2**). Principal component analysis (PCA) of the transcriptomic data shows that samples cluster strongly by tissue of origin, with clear separation among the three salivary glands (**Figure S1**), supporting the robustness of the dataset.

To characterize gland-specific expression patterns and contribution to saliva proteome, we calculated mean transcript abundance herein estimated in transcripts per million (TPM). Each gland displayed a distinct transcriptional signature, typically dominated by a small subset of highly expressed, gland-specific genes that code for secreted proteins, consistent with previous work (Gao et al. 2018). We then identified the 30 most highly expressed genes encoding secreted proteins in each mouse gland (see **Methods**; **Figure 1a, Table S3**). These genes account for 69.3%, 64.9%, and 35.6% of total transcript abundance in the parotid, submandibular, and sublingual glands, respectively. When considering only transcripts encoding secreted proteins, these top 30 genes explain an even larger fraction of expression: 95.7%, 95.9%, and 87.0% in the respective glands.

Next, we categorized mouse salivary gland genes based on their evolutionary relationships with human genes, either as one-to-one orthologs, other orthologous relationships (i.e., one-to-many or many-to-many; see **Methods**), or lineage-specific genes (**Figure 1a**). We applied the same classification to human salivary gland transcriptomes from our previous study (Saitou et al. 2020), enabling direct comparison between species (**Figure 1b, Table S4**). The proportion of expression originating from one-to-one orthologs that code for secreted proteins are remarkably low for sublingual and submandibular glands, approximately 8% and 11%, respectively (**Figure 1c**). In contrast, the parotid gland displays a higher proportion of expression of one-to-one orthologs (77.58%), suggesting greater evolutionary conservation in its core secretory functions.

We recently hypothesized, based on a locus-specific analysis, that salivary glands evolve faster than other tissues because of shifting dietary and pathogenic

evolutionary pressures (Pajic et al. 2025). Here, we tested this hypothesis more
comprehensively at the transcriptome-wide level by comparing the expression levels
of one-to-one orthologs between human and mouse, which are expected to maintain
comparable expression levels across species (Kryuchkova-Mostacci and Robinson-
Rechavi 2016). This expectation was met for genes that code for secreted proteins in
pancreas and liver, which showed strong correlation of expression between human
and mouse transcriptomes (Spearman's ρ = 0.64, p = 1.68 × 10^{-147} and ρ = 0.78, p =
8.55 × 10^{-267}, respectively). In contrast, genes that code for secreted proteins in the
salivary glands exhibited weaker mouse-human correlations (parotid: ρ = 0.575, p =
5.90 × 10^{-112}; submandibular: ρ = 0.58, p = 2.71 × 10^{-114}; sublingual: ρ = 0.606, p =
2.27 × 10^{-127}; **Figure S2**).

To determine whether these transcriptomic differences for genes encoding secreted
proteins are also reflected at the proteomic level, we performed gel electrophoresis
of whole saliva (**Figure 1d**; **Figure S3**). The resulting protein and glycoprotein
profiles revealed striking qualitative and quantitative differences in banding patterns
between human and mouse saliva, with only a few bands appearing to match
between the two species. Particularly noteworthy is the observed shift in gel
electrophoretic mobility of the high-molecular-weight glycoprotein band from 150 kDa
in males to 130 kDa in females. This protein band was identified by mass
spectrometry (MS/MS) analysis as Muc19 (**Table S5**).

In parallel, we asked the extent to which the transcriptomic variation observed
translates into variation in the salivary proteome. To assess this, we compared gene
expression in the salivary glands of each mouse strain with the abundance of the
corresponding proteins in their saliva for 467 gene-protein pairs (see **Methods**). We
observed a significant positive correlation for each gland (parotid: ρ = 0.362, p = 7.29
× 10^{-16}; sublingual: ρ = 0.268, p = 3.76 × 10^{-9}; submandibular: ρ = 0.304, p = 1.81 ×
10^{-11}) (**Figure S4**). Notably, correlation strength increased among the most highly
expressed genes that code for secreted proteins, with stronger associations
observed when restricting the analysis to the top 100 transcripts per gland (parotid: ρ
= 0.492, p = 2.06×10^{-7}; sublingual: ρ = 0.528, p = 1.65×10^{-8}; submandibular: ρ =
0.582, p = 2.10×10^{-10}) (**Figure 1e**). These protein-level analyses support the notion
that transcript abundance in the salivary glands directly affects protein levels in
saliva, and that highly expressed genes encoding secreted proteins have an
outsized influence on shaping the oral proteome.

Our combined transcriptomic and proteomic analyses support a model in which
mammalian salivary glands function as specialized organs that produce a limited set
of highly abundant secreted proteins. Our data further support the hypothesis that
the genes coding for secreted proteins in the salivary glands show faster turnover
than those in liver and pancreas, a phenomenon more pronounced among the top
expressed genes encoding secreted proteins. Transcriptomic data from individual
salivary glands accurately predict their relative contributions to the oral proteome,

consistent with previous observations in humans (Saitou et al. 2020). Overall, our
results reveal fundamental differences in expression profiles of genes coding for
secreted proteins in the salivary glands of humans and mice, consistent with a rapid
evolution   of   the   salivary   gland   secretome.

Figure 1. Most mouse genes coding for secreted proteins in salivary glands are lineage-specific. **(a)**
Bubble plots showing the top 30 most highly transcribed genes that code for secreted proteins in each

major mouse salivary gland. Bubble size represents mean transcript abundance herein estimated in transcripts per million (TPM), and colors denote the type of orthology: one-to-one (green), lineage-specific (maroon), and other orthologous relationships (including many-to-many and one-to-one) (gray). **(b)** Same analysis as in A, here for human salivary glands. (**c**) Proportion of total gene expression (mean TPM) of genes that code for secreted proteins attributed to each orthology category in mouse and human. **(d)** Saliva proteins separated by SDS-PAGE and stained with Coomassie blue and Periodic acid Schiff's stain for pooled whole saliva of males (M, n=5) and females (F, n=5) for mice and humans. Pink bands are glycosylated proteins revealed by periodic acid Schiff stain. Note the shift in gelelectrophoretic mobility of the high-molecular-weight glycoprotein band from 150 kDa in males to 130 kDa in females. MW denotes molecular weight markers. **(e)** Correlation between gene expression (Log_{10}(1+Mean TPM)) and protein abundance (Log_{10}(1+Mean iBAQ)) for genes that code for secreted proteins in the submandibular gland. The right panel highlights the top 100 most highly expressed genes, for which RNA and protein expression are more concordant. ρ denotes Spearman's correlation. *denotes $p<0.05$.

The submandibular gland is the major contributor to sexual dimorphism in saliva

Previous studies have documented sexual dimorphism in the postnatal development and cellular composition of the mouse submandibular gland, most notably the male-specific expansion of granular convoluted tubules (ducts) (Caramia 1966; Pardini and Taga 1996). These morphological differences are known to influence salivary protein output and gene expression (Treister et al. 2005; Mukaibo et al. 2019). Thus, we hypothesize that the submandibular gland contributes to sexual dimorphism in the salivary proteome. Analysis of our mass spectrometry dataset identified 368 and 152 proteins with significantly male- and female-biased protein abundances, respectively ($p_{adj}<0.05$ and |log_2FoldChange| > 0.5, **Table S6**). To determine the transcriptional basis of this protein-level dimorphism, we next examined whether sex-biased gene expression across individual salivary glands contributes to the observed differences in salivary protein abundance.

To systematically address this hypothesis, we performed comparative transcriptomic analyses of sex-biased gene expression across mouse salivary glands, liver, and pancreas (**Table S7**). The submandibular gland exhibited the largest number of tissue-specific differentially expressed genes (DEGs, FoldEnrichment>1, $p_{adj}<0.05$) between sexes (1,537 genes) with other salivary glands exhibiting markedly fewer tissue-specific DEGs (**Figure 2a**), consistent with previous reports (Mukaibo et al. 2019). What distinguishes our analysis is the direct cross-organ comparison. We found that the submandibular gland, analyzed within the same individuals, harbors more than fivefold sex-biased DEGs than liver and ninety-sixfold more than pancreas. These results place the submandibular gland among the most sexually dimorphic tissues in mice, clearly exceeding even the liver, which is widely regarded as the canonical model for studying sexual dimorphism in mammalian gene expression (Lai et al. 2016; Nikkanen et al. 2022). Further, a direct comparison of the extent of sexual differentiation in submandibular transcriptomes shows a dramatic difference between mice and humans (**Figure 2b, Table S8**).

Next, we tested if most sexually dimorphic saliva proteins originate from genes with
sex-biased expression in the submandibular gland. Protein abundance in saliva is
influenced by many factors, including RNA abundance, translational efficiency,
posttranslational modifications, and peptide degradation. Despite this complexity, we
found that 70 of the 507 sexually dimorphic salivary proteins are encoded by genes
that are differentially expressed in the submandibular gland, compared with only one
gene in the parotid gland and one in the sublingual gland (**Figure 2c**). These results
show that the submandibular gland is the primary contributor to the sexual
dimorphism observed in the protein composition of mouse saliva.
240
To further validate these results, we used immunohistochemical staining with well-
characterized antibodies to examine the expression of nerve growth factor (*Ngf*), one
representative sex-biased protein, which showed pronounced sexual dimorphism in
both submandibular RNA-seq and salivary proteome analyses (Bie et al. 2010;
Wong et al. 2019). Examination of submandibular gland sections from male and
female mice revealed clear immunohistochemical differences between sexes
(**Figure 2d**), supporting that pronounced sexual dimorphism at the gene expression
and secreted protein levels, are also manifested at the cellular level.
249
In addition to transcriptional differences, post-translational modifications, such as
glycosylation, may further contribute to sex-specific differences in the biochemical
and functional properties of salivary proteins. In particular, sialic acids are negatively
charged terminal glycan residues on most mammalian glycan chain and are
commonly present on glycans of secreted glycoproteins, most prominently on
mucins, where they influence the biophysical and tribological properties of these
molecules and play important roles in their interaction with microbes and viruses
(Varki and Schauer 2009; Hiremath and Wagner 2026). Consistent with this
possibility, we observed differential expression of several genes involved in
sialylation pathways, including the sialyltransferases *St3gal1*, *St3gal4*, *St3gal5*,
St6galnac1, and *St6galnac2*, between males and females (**Figure S5**). Notably, in
contrast to mice, in humans none of these sialyltransferases are differentially
expressed between sexes in the submandibular gland.
263
To test whether these transcriptional differences correspond to differences in protein
glycosylation, we investigated the prominent glycoprotein band in mouse saliva that
shifts in electrophoretic mobility from ~150 kDa in male saliva to ~130 kDa in female
saliva (see **Figure 1d**). Analysis by MS/MS of the major protein present in these
bands excised from the gel identified it as mucin Muc19. Lectin blotting using PNA,
revealed strong binding to the band in male saliva, but lack of binding to the band in
females, whereas lectin blotting with MAL-II showed strong binding to the band in
females and weak binding to the band in male saliva. Together, these results
indicate presence of unsialylated core O-glycans (Gal□1-3GalNAc) on male Muc19
and a higher proportion terminal α2,3-linked sialylation on female Muc19 (**Figure
2e**). The higher degree of Muc19 sialylation in female mice is consistent with the

observed electrophoretic shift of the Muc19 band from ~150 kDa in male saliva to
~130 kDa in female saliva due to the added electronegativity conferred by the higher
number of negatively charged sialic acid molecules on the female mucin glycan
chains. These results suggest that, in addition to sex-specific differences affecting
the transcription of genes encoding secreted proteins in the submandibular gland
directly, sex-specific post-translational modifications also contribute substantially to
the sexual dimorphism observed in mouse saliva, and are caused by sex-specific
transcription differences in modifying enzymes, thereby affecting secreted protein
products indirectly in a sex-specific manner.

Together, these findings demonstrate that the submandibular gland, by far the most
sexually dimorphic tissue in our dataset, is the primary contributor to the strong
sexual dimorphism observed in the salivary proteome, with additional contributions
from sex-specific post-translational modifications.

Figure 2. The submandibular gland is the major contributor to sexual dimorphism of the salivary gland
secretome. **(a)** Upset plot showing that the submandibular (SM) gland harbors more exclusive DEGs
between mouse sexes than other major salivary glands (parotid [PAR], sublingual [SL]) or non-
salivary tissues (liver [LIV], pancreas [PANC]). Sex-linked genes (*Ddx3y*, *Eif2s3y*, *Kdm5d*, *Uty*, *Xist*)
that are differentially expressed in all tissues are indicated with an arrow. **(b)** Volcano plot of

differentially expressed genes in the mouse (left) and human (right) submandibular gland. Arrows indicate the total number of significantly upregulated genes in females (pink) and males (blue). Pink dots represent genes significantly upregulated in females, and blue dots represent genes significantly upregulated in males. Significance is measured in $\text{Log}_{10}(p_{adj})$. **(c)**. Correlation plot of protein abundances in males and females quantified in $\text{Log}_{10}(1+\text{Mean iBAQ})$. Each dot represents a differentially abundant protein. Female protein abundance is shown on the y-axis and male protein abundance on the x-axis. The dashed line indicates one-to-one correspondence. Colored points denote sexually dimorphic proteins that are also differentially expressed genes in each gland. The inset panel shows the number of differentially abundant proteins whose genes are also differentially expressed in each gland, with the highest contribution observed in the submandibular gland. **(d)** Immunofluorescence staining showing differences in expression patterns between female (top) and male (bottom) submandibular glands. NGF (red). DAPI marks nuclei (blue). **(e)** Sex-biased expression of the sialyltransferase *St3gal1* is associated with differences in Muc19 glycosylation. Left: Boxplot showing transcript abundance of *St3gal1* (TPM) in male and female mice. One female outlier (TPM = 443) is shown as a triangle and was kept for statistical calculations. Right: SDS-PAGE and lectin blot analysis of Muc19 in pooled male (M) and female (F) saliva samples. PNA detects unsialylated core O-glycans, whereas MAL-II detects terminal α2,3-linked sialic acids. Differences in lectin binding and electrophoretic mobility indicate higher sialylation of female Muc19, consistent with increased *St3gal1* expression.

Klk gene family expansion underlies sex-biased expression in mice

The profound sexual dimorphism in the mouse submandibular gland is not recapitulated in humans (**Figure 2b**). This observation motivates a closer examination of the evolutionary processes shaping sex-biased gene expression in the mouse submandibular gland. In particular, we asked whether this extensive sex bias might reflect coordinated regulation of clusters of genes within specific genomic regions or a broad, genome-wide effect.

To address this, we first searched for physically clustered genes that may be co-regulated to express in a sex-specific manner. Specifically, we performed a genomic clustering analysis of sex-biased DEGs in the mouse submandibular gland using 100-kb windows (see **Methods**; **Table S9**). We found that the Klk gene family forms the largest cluster of sex-biased genes in the genome and is a significant outlier relative to other clusters ($p < 0.001$; **Figure 3a**). Fourteen members of the kallikrein (Klk) family (*Klk15, Klk1b8, Klk1b1, Klk1b9, Klk1b11, Klk1b26, Klk1b27, Klk1b21, Klk1b22, Klk1b16, Klk1b24, Klk1b3, Klk1b4, and Klk1b5*) exhibit strong male-biased expression, whereas one member (*Klk1*) shows female-biased expression. Altogether, *Klk* genes make up ~16.4% of total gene expression in male submandibular glands, compared with only 2.5% in female ones (**Figure 3b**).

We then asked whether the clustering of sex-biased genes can also be observed at the proteomic level. Of the Klk genes showing sex-biased expression, 11 out of 15 also display concordant sex-biased differences in protein abundance in saliva. Using genomic coordinates of genes encoding differentially abundant proteins, we observed that the *Klk* genes exhibiting sex-biased protein abundance are clustered on chromosome 7 (**Figure 3a**), consistent with the transcriptomic clustering pattern. Together, these results indicate that the kallikrein locus represents a genome-wide

outlier cluster of coordinated, sex-biased regulation at both the transcriptomic and
proteomic levels.

Previous studies have already shown that *Klk* genes exhibit male-biased expression
in the mouse submandibular gland (Stopka et al. 2016; Mukaibo et al. 2019) and in
saliva (Penschow et al. 1991; Karn and Laukaitis 2011; Blanchard et al. 2015;
Moustardas et al. 2021). Single-cell transcriptomic analyses further indicate that *Klk*
genes are specifically expressed in granular convoluted tubule (GCT) and acinar
cells (Horeth et al. 2021), which are approximately 5- and 1.5-fold more abundant in
males, respectively (Pardini and Taga 1996). This raises the question of whether
differences in cell abundance between male and female submandibular glands alone
can explain the observed sex-biased expression. If expression per cell were
equivalent between sexes, differences in cell-type abundance could, in principle,
account for higher *Klk* expression in males. However, when we quantified overall
expression levels, we found that male-female differences exceeded by at least 5.5-
fold for all differentially expressed *Klk*s, indicating that the observed expression
differences cannot be explained by cell-type proportions alone (**Figure 3c**).

One remarkable exception is that the observed male-biased pattern for *Klk1b* genes
is not observed in *Klk1*. In fact, we found that *Klk1* is expressed significantly higher in
female submandibular glands despite the fact that *Klk1b* genes were shown to be
duplicates of *Klk1* (Lawrence et al. 2010; Pavlopoulou et al. 2010)*,* and that *Klk1* is
also known to be expressed in acinar and GCT cells which are much more abundant
in males. To verify this transcriptional level observation at the protein and cellular
levels, we performed immunohistochemical staining of both male and female
submandibular gland sections for the female-biased *Klk1* gene, one of the few *Klk*
family members for which a robust antibody is available (**Figure 3d**). Consistent with
the gene expression pattern, protein abundance was significantly higher in female
granular convoluted tubules (**Figure S6**), resulting in greater overall *Klk1* expression
in females despite the higher abundance of these cells in males. This observation
suggests evolution of complex regulatory mechanisms in the Klk locus, distinct for
Klk1 and its paralogs.

Together, these results indicate that differences in cellular composition alone are
insufficient to explain sex-biased *Klk* expression and suggest additional genetic or
epigenetic regulation that drive this dimorphism.

Figure 3. The *Klk* gene family forms the largest cluster of sex-biased transcripts and proteins in mouse submandibular glands, independent of differences in cell abundance. **(a)** Genomic distribution of genes encoding for sex-biased proteins (top, green) and sex-biased transcripts (bottom, magenta) along mouse chromosome 7 in the submandibular gland. Points represent differentially abundant proteins or differentially expressed genes, plotted by genomic position and log2 fold difference between sexes. The shaded region highlights the *Klk* locus. Density tracks below each scatter plot show kernel-smoothed spatial density of significant genes or proteins along the chromosome 7. Blue and pink points indicate genes whose sex bias at the transcript level is also supported by proteomic differences in males and females, respectively. **(b)** Relative expression of *Klk* genes in male and female submandibular glands. Boxes show distribution across biological replicates; points represent individual samples. ** indicates *p* < 0.001. **(c)** Comparison of female versus male *Klk* gene expression levels. All *Klk1b* paralogs show >5-fold higher expression in males, indicating that sex-biased

384 expression is not explained by differences in glandular cell composition. **(d)** Immunofluorescence
385 staining of submandibular glands from female (top) and male (bottom). Klk1 protein is shown in green;
386 nuclei are counterstained with DAPI (blue), illustrating stronger Klk1 expression in females.

Sex-specific *Klk* expression is shaped by hormonal regulation, and lineage-specific 3D genomic architecture

To investigate the genetic basis of male-biased expression at the *Klk* locus, we first
generated a gene tree incorporating gene family expansion and contraction
estimates inferred by CAFE 5 and obtained from Ensembl (Mendes et al. 2021). We
found that placental mammals, including other rodents, harbor up to 15 *Klk* genes,
whereas mice and rats harbor approximately 17-25, indicating a burst of duplications
in the murid lineage (**Figure S7**). Specifically, there are 13 additional tandemly
duplicated copies in mice compared to the human genome (**Figure 4a**), all
duplicated from *Klk1* (Lawrence et al. 2010; Pavlopoulou et al. 2010) and with male-
biased expression.

Next, we asked whether the large cluster of differentially expressed genes containing
Klk genes in the submandibular gland share common regulatory sequences that
could underlie their coordinated male-specific expression. To address this, we
performed a *de novo* motif enrichment analysis on this cluster, identifying
overrepresented sequence motifs located near genes (see **Methods**). We found an
enrichment ($p = 1 \times 10^{-42}$) for a conserved CTGATCCTGTTC motif located within 30
bp of the transcription start site in all sex-biased *Klk* genes except *Klk15* (**Figure 4a**;
see Methods; **Tables S10** and **S11**). Previous work showed that this motif is a
binding site for Ghrl2, a developmental transcription factor whose activity is sex-
biased and correlated with testosterone levels (Paltoglou et al. 2017; Reese et al.
2019). Consistent with this, kallikreins are well documented as androgen-regulated
genes in the mouse submandibular gland (Van Leeuwen et al. 1987; Kurabuchi et al.
2001; Lawrence et al. 2010). In contrast, the motif is absent from *Klk* genes that do
not show sex-biased expression. Moreover, none of the human *KLK* genes present
this motif at the orthologous locus (**Table S11**). Given that the *Klk1b* paralogs arose
through lineage-specific duplications of an ancestral *Klk1* gene (Lawrence et al.
2010; Pavlopoulou et al. 2010), our results support an evolutionary scenario in which
the CTGATCCTGTTC motif emerged in the ancestral murine *Klk1* gene before
subsequent gene expansion. Subsequent duplications may have increased gene
expression through this regulatory mechanism, thereby amplifying male-biased
expression in the submandibular gland.

Previous work in humans identified two distinct topologically associating domains
(TADs) that structure the regulatory landscape of the *KLK* gene family, both
demarcated by CTCF binding sites (Khoury et al. 2020). We therefore asked whether
the lineage-specific duplications in mice resulted in the formation of a new or
expanded the existing topologically associating domain (TAD). To address this, we
analyzed publicly available Hi-C, ATAC-seq, and ChIP-seq datasets, including

histone modification profiles, from the mouse B-cell lymphoma cell line (see
Methods). Consistent with the human architecture, we identified two TADs at the
mouse *Klk* locus of which the second TAD is markedly expanded, encompassing 15
genes, including all those that show sex-biased expression (**Figure 4b**). Chromatin
accessibility (ATAC-seq), histone modification (H3K27ac), and transcriptional anchor
protein binding (Ctcf, Rad21) profiles delineate the borders of these TADs.

Although the TAD boundaries in mice largely mirror those observed in humans and
encompass all sex-biased *Klk* genes, it remains unclear how *Klk1* retains both the
regulatory motif and its position within the same TAD. To explore whether sex-
specific chromatin organization might contribute to this regulatory difference, we
compared TAD boundaries between male and female cells. Using publicly available
Hi-C data from male Sertoli cells and female granulosa cells, we reconstructed TAD
structures and compared their boundaries (**Figure 4c**). This analysis revealed a
slight boundary shift that coincides with the region showing male-specific expression,
suggesting that *Klk1* in females may be regulated within a different TAD context.
These findings indicate that regulatory control at this locus is more complex than
previously appreciated, and that the data presented here are insufficient to fully
explain the expression pattern of *Klk1*.

Together, these results suggest that lineage-specific *Klk* gene duplications have
reshaped the ancestral TAD structure shared with humans (**Figure 4d**), expanding it
in mice in a sex-specific manner.

Figure 4. Expansion of the mouse *Klk* locus is associated with sex-biased expression and reorganization of local 3D genome structure. **(a)** Schematic comparison of the *KLK/Klk* locus organization in humans (top) and mice (bottom). Circles represent individual *KLK/Klk* genes ordered by genomic position and transcriptional orientation. Genes showing female- or male-biased expression are indicated in pink and blue, respectively. Orange bars denote putative transcription factor binding motifs shared across the expanded mouse cluster. The mouse lineage shows extensive

tandem duplication of *Klk* genes compared with the more compact human locus. **(b)** Hi-C contact map
of the mouse *Klk* region generated from B-cell lymphoma cell line, revealing two adjacent topologically
associating domains (TADs). The left TAD predominantly contains non–sex-biased genes, whereas
the right TAD encompasses the expanded cluster of sex-biased *Klk* paralogs. Gene annotations are
shown below the contact map together with chromatin accessibility and regulatory marks (ATAC-seq,
H3K4me3, CTCF, and RAD21), highlighting active promoters and architectural features associated
with TAD boundaries. **(c)** Hi-C contact maps of the mouse *Klk* locus reconstructed from male Sertoli
cells (top) and female granulosa cells (bottom). Dashed vertical lines indicate shifts in TAD
boundaries between the two cell types. Genes located within these alternative domain configurations
are highlighted, suggesting that differences in higher-order chromatin organization may influence sex-
specific regulation of subsets of *Klk* paralogs. **(d)** Conceptual model illustrating how tandem gene
duplication at the *Klk* locus may have been accompanied by expansion and restructuring of the
surrounding TAD landscape.

DISCUSSION AND CONCLUSION

In this study, we investigated the evolutionary forces shaping the composition of the salivary gland secretome in mice. We show that most highly expressed genes encoding secreted saliva proteins in mice are lineage-specific and lack direct human orthologs. We further identify the submandibular gland as the focal point of sex-specific expression. The kallikrein (*Klk*) gene family in mice accounts for ~16.4% of male-biased expression, and we discuss how lineage-specific gene family expansion, emergence of novel regulatory motifs, and topological clustering contribute to the pronounced sex differences in expression. Together, the findings illustrate how gene duplication and regulatory rewiring drive sex-biased gene expression and contribute to the rapid evolution of the mammalian salivary gland secretome in different lineages.

When we examined sex differences in mouse saliva, we found that the submandibular gland is the main contributor to the protein differences observed between males and females, consistent with previous results (Treister et al. 2005; Mukaibo et al. 2019). This male bias in submandibular expression is particularly notable when compared with the liver, the most studied tissue for sex differences in mice (Nikkanen et al. 2022), where the degree of dimorphism is lower. We further found that this bias can be traced to the lineage-specific gene duplication clusters identified in our analysis, with the kallikrein (*Klk*) cluster standing out as a single gene family accounting for ~16.4% of male-biased expression in the submandibular gland. The sex-biased expression of kallikrein genes is well documented in the literature, aligning with the patterns we observe here (Karn and Laukaitis 2011; Stopka et al. 2016; Moustardas et al. 2021). Specifically, our results support a model in which the expression of a hypothetical ancestral *Klk* gene in the murine lineage was subsequently amplified through a burst of tandem duplications within the same topologically associating domain (TAD), a process that could have both accompanied and facilitated developmental shifts in the male submandibular gland. This mechanistic framework suggests a broader evolutionary trend: rapid regulatory change across multiple genes can emerge through *de novo* motif evolution at a single locus, followed by its propagation via gene duplication within the same 3D genomic domain (**Figure 5**). A similar trend for motif evolution has recently been observed at the primate amylase locus (Karageorgiou et al. 2025)

Figure 5. Proposed model for the evolutionary trajectory of mouse kallikrein (*Klk*) genes in salivary glands. Gene duplication created multiple *Klk* paralogs, which subsequently acquired new regulatory

motifs. These regulatory changes enabled functional diversification, including interaction with Ghrl2.
Further duplications occurred within a conserved topologically associating domain (TAD), promoting
coordinated regulation of the expanded gene set. This combination of gene family expansion and
regulatory rewiring led to increased male-biased expression in the submandibular gland.

The pronounced male-biased expression of kallikreins in saliva raises questions
about their phenotypic impact in male mice. A prevailing hypothesis proposes that
the presence of sex-specific proteins in saliva in male mice is related to improved
healing through wound licking behavior after male-male aggression (Hutson et al.
1979; Stopka et al. 2016). While this hypothesis may be valid for the higher
expression of wound healing related saliva proteins in male mice, such as epidermal
growth factor and fibroblast growth factor (Hirata and Orth 1979; Brown et al. 2020),
it may not extend to the kallikrein gene family. An alternative hypothesis has been
posited by Karn and Laukaitis where sex-biased expression of proteins in rodent
saliva is explained by either signaling male dominance among rival males or mate
identification through pheromone-like mechanisms (Karn and Laukaitis 2011). Our
results cannot confirm or falsify either of these hypotheses, but they may provide a
mechanistic base for formulating testable hypotheses approaching the questions
from a molecular genetic level.

While we propose a hypothesis of co-regulation among male-biased *Klk* genes, our
results do not fully explain the expression pattern of the only female-biased *Klk* gene
(*Klk1*). Previous studies suggest that additional regulatory mechanisms contribute to
the regulation of *Klk* transcripts, including microRNA-mediated silencing (Kurihara et
al. 2011), highlighting the regulatory complexity of this locus. Moreover, *Klk1* has
been reported to be regulated by estrogen in mice through mechanisms distinct from
those described for other *Klk* family members (Iwasaki et al. 2023). Future studies
integrating genomic, regulatory, and three-dimensional genome architecture data
from salivary glands of the same individuals ideally using a single-cell resolution will
be necessary to validate our proposed model of co-evolution among regulatory
motifs, chromatin organization, gene duplications, and cell-type composition.

The biological functions and physiological relevance of the extensive species- and
sex-specific differences in salivary gene and protein expression remain largely
unknown. While addressing these functions was beyond the scope of the present
study, our results provide a comprehensive catalogue of candidate genes and
proteins that can now be used to test hypotheses about the roles of salivary gland-
derived proteins in physiology, immunity, and behavior. One novel observation is the
sex-specific glycosylation of the Muc19 protein in mouse saliva, suggesting that
differences in glycosylation machinery, particularly the differential expression of
sialyltransferases between males and females, contribute to sex-specific
modification of secreted mucins. This stands in contrast to humans, where MUC19
expression is low in salivary glands and no clear sexual dimorphism has been
reported. Notably, genetic variation in human *MUC19* has been linked to complex
introgression patterns in hominins and signatures of local selection in Indigenous

American populations (Villanea et al. 2025), marking this gene as a hotspot of
adaptive evolution. Together, our observations provide a foundation for future work
aimed at dissecting the regulatory basis of species-specific expression, the
molecular mechanisms underlying sex-specific glycosylation, and the evolutionary
processes shaping Muc19 variation across mammals.

Of importance for biomedical research, our findings raise more fundamental
questions about the suitability of the mouse as a standard model for studying human
salivary gland biology, a concern that has also been noted previously (Maruyama et
al. 2019). Focusing on salivary glands and their secreted products, we show that
mice and humans differ fundamentally in both the expression patterns and
composition of salivary proteins, including their post-translational modifications. With
respect to disease relevance in humans, we found that several highly expressed
Sjögren's-related genes (Horeth et al. 2021), including *Dcpp1*, *Scgb2b26*, and
Scgb2b27, are mouse-specific and lack human counterparts, thus highlighting the
importance of validating translational studies involving mice salivary glands and
saliva in human models. In addition, our results underscore the importance of
explicitly accounting for sex as a biological variable in studies of oral and salivary
gland biology, particularly in mouse models.

Overall, this study, although limited to a comparison between two species,
contributes to the emerging view that saliva exhibits considerable evolutionary
plasticity, enabling accelerated evolution under selective pressures (Ruhl 2012; Pajic
et al. 2019; Pajic et al. 2025) and offers novel additional mechanistic hypotheses to
explain these trends. Furthermore, our results are consistent with the hypothesis of
developmental systems drift (True and Haag 2001), whereby similar phenotypic traits
across species evolve through distinct genetic or gene-regulatory mechanisms.
Although saliva performs broadly conserved functions across mammals, primarily
related to digestion, protection of oral tissues, microbial homeostasis, and pathogen
defense, our results suggest that these functions may be achieved through different
regulatory architectures across species. In this context, the murine *Klk* locus
provides a useful model for investigating how gene duplication, regulatory motif
evolution, and 3D genome organization can shape the evolution of sex-biased gene
regulation and the emergence of novel sexually dimorphic traits. Expanding similar
analyses across multiple mammalian species will be crucial for uncovering general
principles governing the evolution of salivary gland gene regulation and function.

METHODS

Sample collection for RNA-seq

A total of 12 mice (3 months old) were used, consisting of three biological replicates for each strain-sex combination: 3 female CD1, 3 male CD1, 3 female C57BL/6, and 3 male C57BL/6 (**Figure S8**). CD1 and C57BL/6 mice were selected as representative outbred and inbred strains, respectively, as both are widely used in laboratory studies of salivary glands (Nguyen et al. 2009; Gervais et al. 2015; Hsieh et al. 2017). Mouse samples were provided by UCSF. Animals were euthanized following the approved protocol AN201280-00I. From each mouse, we collected five glands: the three major salivary glands (parotid, sublingual, and submandibular glands), as well as the pancreas and liver, which were used as reference tissues.

RNA-sequencing and analysis of mouse transcriptomes and differential analysis

Total RNA was extracted from each tissue, and RNA-seq libraries were prepared by Azenta Life Sciences (GENEWIZ) using their Standard/Strand-Specific RNA-Seq service with poly(A) selection for mRNA enrichment, followed by sequencing on an Illumina® NovaSeq™ platform (2×150 bp). Raw reads were processed with Trim Galore (Krueger et al. 2023), which uses cutadapt for adapter trimming and FastQC for quality assessment, to remove low-quality bases and adapter sequences. Cleaned reads were then aligned to the *Mus musculus* reference genome (GRCm39) using HISAT2 (Kim et al. 2019), and gene-level quantification was performed with featureCounts (Liao et al. 2014), which assigns aligned reads to annotated genes and produces a matrix of raw read counts per gene.

Six parotid samples (two male and one female CD1, and three female C57BL/6) exhibited high expression of genes associated with muscle function and contraction, which is unusual in parotid gland transcriptomes. These samples were excluded from downstream analyses, as their expression profiles suggested possible contamination during tissue dissection. Next, we measured differential gene expression between sexes for each tissue using DESeq2 (Love et al. 2014), applying a significance threshold of $p_{adj} < 0.05$ and |log2 fold change| > 1. DESeq2 models raw read counts for each gene and compares expression levels between sexes, estimating differential expression while accounting for biological variability. Counts are internally normalized by library size and sequencing depth to allow accurate comparisons across samples, and significantly differentially expressed genes (DEGs) were identified for each tissue. To assess sample quality and confirm proper clustering by biological variables, we performed principal component analysis (PCA) across all glands and separately for the three major salivary glands (**Figure S1**).

Orthology and Secreted Gene Identification

We first sought to determine for all genes in each tissue the human orthologs and
which were mouse specific. For this, we used an orthology reference table provided
by   The   Jackson   Laboratory   (JAX)
(https://www.informatics.jax.org/downloads/reports/HMD_HumanPhenotype.rpt),
which lists mouse-human orthologous genes. Using this table, we grouped genes
based on the number of orthologous relationships between species. Specifically,
genes were classified as one-to-one when a single mouse gene mapped to a single
human gene; one-to-many when a gene in one species mapped to multiple genes in
the other; and many-to-many when multiple genes in both species were linked
through orthology relationships. Genes without any annotated orthologs in the other
species were classified as mouse- or human-specific.

Next, we aimed to characterize the contribution of highly expressed genes to salivary
secretion by identifying secreted proteins and building an expression profile for each
gland. Transcript abundance was calculated in transcripts per million (TPM) using
the standard formula (Wagner et al. 2012). TPM values were computed for all genes
in the three major salivary glands (parotid, sublingual, and submandibular). All genes
in each gland were annotated as secreted or non-secreted based on the UniProt
mouse   proteome   (https://www.uniprot.org/proteomes/UP000000589)   (UniProt
Consortium 2025) and human from Human Protein Atlas ([proteinatlas.org)](proteinatlas.org) (Uhlén et
al. 2019). For the mouse proteome, specific annotations for salivary proteins were
not available; therefore, we classified all genes as secreted or non-secreted based
on general proteome annotations from UniProt. Combined with the orthology
annotations described above, these data were used to generate Figure 1. We
identified the 30 most highly expressed genes that code for secreted proteins in each
salivary gland and plotted their expression levels (in TPM) to visualize gland-specific
expression patterns using packcircle (https://mbedward.r-universe.dev/packcircles).

Analysis of human transcriptomes

For humans, we reanalyzed publicly available RNA-seq data from salivary glands
(parotid, submandibular, and sublingual) generated by Saitou et al. (2020). Raw
sequencing reads were processed as described above for mouse, using the human
reference genome (hg38). Expression levels were quantified as transcripts per
million (TPM), and mean TPM values were calculated for each gland. For liver and
pancreas tissues, transcript data in TPMs was downloaded from GTEx and filtered
for the same genes present in our dataset for salivary glands. TPMs were then
recalculated for only those genes, and mean TPMs were calculated using all
individuals. For orthology classification, please see the description of orthology
classification above. Differential expression analysis was performed using DESeq2
(Love et al. 2014) using raw gene counts only for submandibular gland samples.

Correlation analysis of human-mouse transcriptomes

To assess the similarity of gene expression patterns between mouse and human
salivary glands, we performed cross-species correlations of gene expression (mean
TPM values) for each gland, focusing on one-to-one orthologs for genes with TPM>2
in each tissue. Because genes encoding secreted proteins and non-secreted
proteins are subject to distinct evolutionary pressures, and secreted proteins are
often expected to evolve and diverge more rapidly (Winter et al. 2004; Feyertag et al.
2016), we focused our analysis on genes encoding secreted proteins, which
constitute the primary functional components of saliva. For the mouse, we used the
annotated transcriptomes generated in this study for all glands. For the human data,
we used the transcriptomic profiles from Saitou et al. (2020) for the major salivary
glands (parotid, submandibular, and sublingual) and from GTEx for the pancreas and
liver. Spearman correlation coefficients (ρ) were calculated for each comparison.
Finally, we note that these analyses involve data from different cohorts and species
and are therefore subject to potential batch effects; as such, these results should be
interpreted with caution. Because of these technical limitations, we avoided direct
comparisons of human and mouse expression data as a primary analytical
approach. Nevertheless, we identified several one-to-one orthologs that exhibit
dozens- to hundreds-fold differences in expression between human and mouse
salivary glands (**Table S3 and S4)**, differences that are unlikely to be explained
solely by batch effects and therefore warrant further investigation.

Mouse saliva collection

Saliva samples were collected from 4-month-old male (n=5) and female (n=5)
C57BL/6 mice. Mice were anesthetized with isoflurane and administered the
sialogogue pilocarpine dissolved in sterile saline via intraperitoneal (i.p.) injection.
Pilocarpine was delivered at a final dose of 4.53mg/kg (concentration 0.68mg/mL;
injection volume 200uL per 30g body weight). Saliva collection commenced 3
minutes post-injection and continued for 10 minutes using a pipette placed in the oral
cavity. Samples were immediately snap-frozen and stored at -80 degrees Celsius for
subsequent protein analysis.

Human saliva collection

Saliva samples were collected from humans (male and female, n=5 each) using the
passive drooling technique as previously described (Thamadilok et al. 2020).

SDS-PAGE

The 8-16% gradient pre-cast Novex Tris-glycine gel (LifeTechnologies; catalog no.
XP08162BOX) was used to study the protein profiling of saliva samples based on
the molecular size of the protein. Meanwhile, the samples were prepared for gel
electrophoresis by mixing with a 2X sample buffer followed by boiling at 100^0C for 10
min. The samples (~15 µg of protein/well) were loaded into the wells, and the gel

was run on an electrophoresis system (XCell SureLock Mini-Cell , Invitrogen) at 170
V for approximately 40 minutes. Subsequently, the gel was prepared for gel staining.

Coomassie blue-PAS (Periodic acid Schiff) Staining

Protein bands were first stained using the Coomassie blue staining procedure. In
brief, the procedure included three steps: a) fixation using fixing solution for 15 min
twice, b) staining in Coomassie stain blue (R-250) solution for 1 hr., and c)
destaining for 30 min thrice. After the destaining step, the gel was stained
subsequently using the PAS staining procedure as described (Thamadilok et al.
2020). Stained gels were digitized using the GE Healthcare ImageScanner III.

Protein identification from excised gel bands by LC-MS/MS

Protein bands of interest were excised from Coomassie blue-stained SDS-PAGE
gels corresponding to the ~150 kDa (male) and ~130 kDa (female) regions using a
clean surgical blade and processed for protein identification by mass spectrometry.
Gel slices were subjected to in-gel digestion with trypsin following standard
protocols. Peptides were extracted and analyzed by liquid chromatography coupled
to tandem mass spectrometry (LC-MS/MS). The resulting spectra were processed
using Spectronaut (Biognosys AG, version 20.0) and searched against the UniProt
Mus musculus protein database. Protein identifications were filtered using a false
discovery rate (FDR) threshold of 1% at both the peptide and protein levels.

Lectin Blotting

Proteins were separated by SDS-PAGE under reducing conditions using 8-16%
gradient gels and subsequently transferred to nitrocellulose membranes. For lectin
detection, membranes were blocked in Tris-buffered saline containing 0.1% Tween-
20 (TBST) supplemented with 2% Polyvinyl alcohol (PVA) for 1 h at room
temperature. Membranes were then incubated with biotinylated lectins diluted in
blocking buffer. Peanut agglutinin (PNA) was used to detect exposed Galβ1–
3GalNAc (T-antigen) O-glycan structures, while Maackia amurensis lectin II (MAL-II)
was used to detect α2,3-linked sialic acid residues. Following lectin incubation,
membranes were washed extensively with TBST and incubated with Alexa fluor-488-
conjugated neutravidin. Lectin binding was detected by enhanced
chemiluminescence and imaged with a digital imaging system.

Protein digestion

Saliva specimens were processed following the E3 protocol described recently
(Martin et al. 2024). In brief, around 20 µL of crude saliva was first mixed with 100 µL
SDS buffer that contained 5% (w/v) sodium dodecyl sulfate (SDS), 100 mM Tris-HCl
solution (pH 8.0) and final concentration of 10 mM tris(2-carboxyethyl) phosphine
(TCEP) and 40 mM 2-chloroacetamide (CAA), and then boiled at 95 □C for 10 min.
Next, the lysate was mixed with 4x volume of 80% acetonitrile (CAN) aqueous

solution and transferred to E3filters (CDS Analytical, Oxford, PA). Following a
centrifugation step, the E3filters were washed with 200 µL of 80% ACN aqueous
solution three times. For protein digestion, 1 mg of trypsin (Promega, WI) in a 200 µL
of 50 mM triethylammonium bicarbonate (TEAB) buffer was added to E3filters and
incubated at 37 □C for 16-18 h with gentle shaking (300 rpm). To elute peptides, the
E3filters were first centrifuged at 3,000 rpm and then flushed with two elution buffers,
200 ml of 0.2% formic acid in water, and 200 ml of 0.2% formic acid in 50% ACN,
respectively. The eluents were dried in SpeedVac, and then subjected to desalting
cleanup using C18 based StageTips (CDS Analytical, Oxford, PA). Dried peptide
samples were stored under -80 □C until further analysis.

LC-MS/MS analysis

The LC-MS/MS analysis was performed using an UltiMate 3000 RSLCnano system
coupled with an Orbitrap Eclipse mass spectrometer and a FAIMS Pro Interface
(Thermo Scientific). The dried peptides were resuspended into 13 µl of LC buffer A
(0.1% formic acid in water) and 10 µl of it was then loaded onto a trap column
(PepMap100 C18, 300 µm × 2 mm, 5 µm; Thermo Scientific) followed by separation
on an analytical column (PepMap100 C18, 50 cm × 75 µm i.d., 3 µm; Thermo
Scientific). A linear LC gradient flowing at 250 nL/min was applied from 1% to 25%
mobile phase B (0.1% formic acid in ACN) over 125 min, followed by an increase to
32% mobile phase B over 10 min. The column was washed with 95% mobile phase
B for 5 min, followed by equilibration with mobile phase A for 15 min. For the ion
source settings, the spray voltage was set to 1.9 kV, funnel RF level at 50%, and
heated capillary temperature at 275°C. The MS data were acquired in Orbitrap at
120K resolution, followed by MS/MS acquisition in data-independent mode following
a protocol described previously (Elsayyid et al. 2025). The MS scan range (m/z) was
set to 380-985, maximum injection time was 246 ms, and normalized AGC target
was 100%. For MS/MS analysis, the spectra were acquired with Orbitrap at
resolution of 30K. The isolation mode was Quadrupole, the isolation window was 8
m/z, and the window overlap was 1 m/z. The collision energy was 30%, AGC Target
was 400K, and normalized AGC target was 800%. For FAIMS analysis, a 3-CV
experiment (–40|–55|–75) was applied.

Proteome quantitation and data analysis

The MS data were processed using Spectronaut software (version 20.0), a library-
free directDIA+ workflow, and the UniProt protein databases (17,141 sequences;
reviewed only). Briefly, the settings for Pulsar and library generation include:
Trypsin/P as specific enzyme; peptide length from 7 to 52 amino acids; allowing 2
missed cleavages; toggle N-terminal M turned on; Carbamidomethyl on C as fixed
modification; Oxidation on M and Acetyl at protein N-terminus as variable
modifications; FDRs at PSM, peptide and protein level all set to 0.01; Quantity MS
level set to MS2, and cross-run normalization turned on.

Differential protein abundance between male and female samples was assessed
using the built-in statistical framework in Spectronaut (version 20.0). The protein LFQ
method was set to Automatic (MaxLFQ); Quantity MS level was set to MS2, and the
Quantity type was Area. The Candidates output from Spectronaut was used for
quantitation analysis. Proteins were considered significantly differentially abundant
between sexes if they met the thresholds of |fold change| > 1.5 and q-value < 0.05.

Transcriptome-Proteome Integration and Correlation Analysis

Proteomic data were filtered to retain only proteins with a single corresponding gene
annotation to avoid ambiguity in gene-protein assignments. For each protein, mean
intensity-based absolute quantification (iBAQ) values were calculated separately for
males, females, and across all samples. For visualization and correlation analyses,
expression values were transformed as $\log_{10}(1 + \text{mean iBAQ})$. To enable
transcriptome-proteome comparisons, we first filtered the transcriptomic dataset to
genes annotated as secreted proteins (see description above). Proteins whose
corresponding genes were present in this filtered transcriptomic dataset were then
retained, resulting in 467 matched gene-protein pairs. Spearman correlation
analyses were then performed using these matched protein-transcript pairs. It is
important to note that each salivary gland contributes only partially to the overall
proteome composition of mixed whole saliva, the oral fluid analyzed in this study.
Therefore, strong correlations between glandular transcriptomes and the whole-
saliva proteome are not necessarily expected, given the distinct spectrum of proteins
secreted by each gland.

Furthermore, several highly expressed salivary gland transcripts were not detected
by mass spectrometry, consistent with known technical limitations of proteomic
assays for certain salivary proteins (Malaker 2024). Conversely, some proteins
detected in saliva at high abundance, such as serum albumin, are not synthesized
by salivary glands but instead enter whole saliva through plasma exudation in the
oral cavity, as reported previously (Oppenheim and Hay 1972; Ruhl 2012).

Immunohistochemistry analysis

Submandibular glands were manually dissected and immediately fixed in 4%
paraformaldehyde (PFA) at 4°C overnight. Following fixation, tissues were
cryoprotected using a graded sucrose approach (10% sucrose w/v followed by 25%
sucrose w/v in PBS). Tissues were then embedded in optimal cutting temperature
(OCT) compound (Fisher Scientific, Cat # 4585) and cryosectioned at a thickness of
16µM. Sections were permeabilized with 0.5% Triton X-100 in PBS for 20 minutes at
room temperature. Following permeabilization, sections were blocked for 1 hour at
room temperature in a buffer containing 10% donkey serum (Jackson
ImmunoResearch, Cat# 017-000-121) and 5% BSA (Fisher BioReagents, Cat#
BP9700100) in PBS.

Sections were incubated with primary antibodies, Kallikrein 1 polyclonal antibody
(1:300; Invitrogen, Cat# PA5-115481) and Nerve Growth Factor (NGF) polyclonal
antibody (1:300; Sigma-Aldrich, Cat# N6655), diluted in antibody diluent overnight at
4°C. Following washing with 0.05% Tween-20 in PBS, antibodies were detected
using Cy2-conjugated secondary Fab fragment antibodies (1:300; The Jackson
Laboratory) incubated for 2 hours at room temperature. Nuclei were counterstained
with Hoechst 33342 (1:1000; Anaspec Inc. Cat# AS-83218). Slides were mounted
using Fluoromount-G (SouthernBiotech, Cat# 01100-01) and imaged using a Zeiss
LSM 900 confocal microscope with a 40 X oil immersion objective.

Quantification of KLK-1 and NGF expression was performed using Fiji software.
Regions of interest (ROIs) were manually delineated around salivary gland ductal
structures identified by distinct morphology. The Mean Fluorescence Intensity (MFI)
was calculated for each ductal ROI. Results are reported as MFI (a.u.).

Clustering of DEGs, and motif enrichment analysis

We hypothesized that co-regulated gene clusters contribute to the differences in
gene expression observed between sexes. To test this, we focused on the
submandibular gland for sex differences, as this tissue showed the highest numbers
of DEGs. Using bedtools, we grouped DEGs into genomic clusters defined by a 100
kb window, such as the following example:

```
832 bedtools  cluster  -i  SM_sex_DEGs.sorted.bed  -d  100000  >
833 SM_sex_DEGs.clustered_100kb.bed
```

To evaluate whether DEGs were more clustered than expected by chance, we
performed a permutation analysis. Specifically, we randomly sampled 10,000
subsets of 1615 genes (matching the observed DEG count) from the total set of
expressed genes and applied the same clustering procedure to each subset. From
these permutations, we built an expected distribution of the number of genes per
cluster size and compared it to the observed data (**Table S9**).

Next, we focused on the kallikrein (Klk) gene cluster in the mouse submandibular
gland. Promoter regions for genes within the cluster were extracted and analyzed for
motif enrichment using HOMER (Hypergeometric Optimization of Motif EnRichment)
(Heinz et al. 2010). HOMER performs *de novo* motif discovery by identifying
sequence motifs that are significantly enriched in a target set of sequences relative
to a background set using a hypergeometric framework, and compares discovered
motifs to known transcription factor binding motifs in curated databases (including
HOMER and JASPAR libraries). For each gene in the cluster, sequences spanning
−400 bp upstream to +100 bp downstream of the transcription start site (TSS) were
analyzed. De novo motif discovery was performed using motif lengths of 8, 10, and
12 bp (**Table S10**), according to the following code

```
851   findMotifs.pl "$gene_list_in cluster" mouse "out_folder/" \
852           -start -400 -end 100 \
853           -len 8,10,12 \
854           -p 4
```

To further confirm the motifs identified in the enrichment analysis and evaluate
whether they are present across other Klk genes in mouse and in their human
orthologs, we searched for these motifs across all Klk gene sequences in both
species. Gene sequences for all annotated Klk genes were downloaded from the
GRCm39 mouse genome assembly and the corresponding human genome
assembly in FASTA format. Extracted sequences included the full gene regions
encompassing coding sequences as well as 5′ and 3′ untranslated regions (UTRs).
The short motif sequence identified by HOMER (CTGATCCTGTTC) was queried
against these gene sequences using BLASTN with the blastn-short task, which is
optimized for short query sequences, according to the following example:

```
865   blastn -task blastn-short -query $motif.fa -subject $genes.fa
866   -outfmt 6 -out out.blastn.out
```

This approach allowed us to identify exact or near-exact matches of the motifs within
Klk gene sequences (**Table S11**).

Evolution of the *Klk* genes, chromatin conformation and Topological
Associated Domains Visualization

To investigate the evolution of the Kallikrein genes and gene expansions-
contractions, we used CAFE 5 (Mendes et al. 2021). To investigate whether the sex-
biased expression of Kallikrein (*Klk*) genes could be explained by differences in
genome architecture and transcriptional regulation, we analyzed publicly available
epigenomic datasets from the ENCODE project. Specifically, we visualized the Klk
locus (chr7:43650000-44300000, mm10) using Hi-C contact maps at 25kb resolution
generated   from   B-cell   lymphoma   cell   line
(https://www.encodeproject.org/files/ENCFF076GYR/) using HiCExplorer (Wolff et al.
2020). We further identified topologically associating domains (TADs) from the Hi-C
data and overlaid them onto the figure. The code used for TAD calling and
visualization is provided below.

```
882   hicFindTADs \
883     --matrix ENCFF651SGY.mcool::/resolutions/25000 \
884     --chromosomes chr7 \
885     --minDepth 75000 \
886     --maxDepth 250000 \
887     --step 50000 \
888     --correctForMultipleTesting fdr \
889     --thresholdComparisons 0.05 \
890     --outPrefix TADs_heart_chr7_25kb_Local
```

```
891
892  WIN2="chr7:43650000-44300000"
893
894  hicPlotTADs --tracks 25kbTADs_selec.ini --region $WIN2 \
895    --outFileName 25kbTADs_selec.svg --dpi 300
896
```

To explore potential regulatory features underlying the observed expression
patterns, we overlaid ATAC-seq data
(https://www.encodeproject.org/experiments/ENCSR451NAE/) representing
chromatin accessibility, chromatin immunoprecipitation sequencing (ChIP-seq) data
for the architectural protein Ctcf
(https://www.encodeproject.org/experiments/ENCSR000ERM/) and for histone
modification H3K27me3
(https://www.encodeproject.org/experiments/ENCSR000CGK/), and for Rad21
(https://www.encodeproject.org/experiments/ENCSR000ERK/) from the same cell
line.

To investigate sex-specific differences in chromatin organization at the kallikrein
(Klk) locus, we analyzed publicly available Hi-C datasets generated from granulosa
(female) and Sertoli (male) cells (Lindeman et al., 2021). These datasets were
obtained from the 4DN Data Portal (https://data.4dnucleome.org/) under accession
4DNEXDK2R8M5 and 4DNEX99DGLEF. Multi-resolution cooler (.mcool) files were
downloaded for each dataset (Sertoli: 4DNFIB7RFFBB; granulosa:
4DNFIZ59STGB). These cell types were selected because they represent sex-
specific gonadal cell lineages, providing a model to examine potential sex-dependent
chromatin architecture. Hi-C contact maps were extracted at 25 kb resolution, and
topologically associating domains (TADs) were independently identified for each
dataset using HiCExplorer as described above. The resulting TAD boundaries were
then compared between male and female datasets to assess differences in higher-
order genome organization at the Klk locus.

ACKNOWLEDGMENTS

We thank Alber Aqil, Charikleia Karagerogiou, Kendra Scherr, Vincent Lynch, and Petar Pajic for helpful discussions and insights. We thank Yanbao Yu for assistance with proteomic analyses and mass spectrometry. We thank Azenta Life Sciences for assistance with RNA-seq library preparation and sequencing. This work utilized computational resources provided by the Center for Computational Research at the University at Buffalo. This work was supported by the National Science Foundation (NSF) (2049947) (to O.G.) and the National Institute of General Medical Sciences (R35-GM156519) (to O.G.).

DATA AVAILABILITY

DATA AVAILABILITY

RNA-seq data generated in this study will be deposited in the NCBI Gene
Expression Omnibus (GEO) prior to publication, and accession numbers will be
provided upon acceptance. Human salivary gland RNA-seq data were obtained
from Saitou et al. 2020 and are publicly available under the corresponding accession
numbers. Proteomic data generated in this study will be deposited in an appropriate
public repository (e.g., ProteomeXchange/PRIDE) prior to publication. All scripts
used for data processing and analysis will be made publicly available upon
acceptance.

REFERENCES

Bie B, Zhang Z, Cai Y-Q, Zhu W, Zhang Y, Dai J, Lowenstein CJ, Weinman EJ, Pan ZZ. 2010. Nerve Growth Factor-Regulated Emergence of Functional δ-Opioid Receptors. *J. Neurosci.* 30:5617–5628.

Blanchard AA, Ezzati P, Shamshurin D, Nistor AC, Leygue E, Wilkins JA, Myal Y. 2015. Towards further defining the proteome of mouse saliva. *Proteome Sci.* 13:10.

Brown CT, Nam K, Zhang Y, Qiu Y, Dean SM, Dos Santos HT, Lei P, Andreadis ST, Baker OJ. 2020. Sex-dependent Regeneration Patterns in Mouse Submandibular Glands. *J. Histochem. Cytochem.* 68:305–318.

Caramia F. 1966. Ultrastructure of mouse submaxillary gland. *J. Ultrastruct. Res.* 16:505–523.

Chai Y, Klauser DK, Denny PA, Denny PC. 1993. Proliferative and structural differences between male and female mouse submandibular glands. *Anat. Rec.* 235:303–311.

Chibly AM, Aure MH, Patel VN, Hoffman MP. 2022. Salivary gland function, development, and regeneration. *Physiol. Rev.* 102:1495–1552.

Dawes C, Pedersen AML, Villa A, Ekström J, Proctor GB, Vissink A, Aframian D, McGowan R, Aliko A, Narayana N, et al. 2015. The functions of human saliva: A review sponsored by the World Workshop on Oral Medicine VI. *Arch. Oral Biol.* 60:863–874.

De Sousa-Pereira P, Cova M, Abrantes J, Ferreira R, Trindade F, Barros A, Gomes P, Colaço B, Amado F, Esteves PJ, et al. 2015. Cross-species comparison of mammalian saliva using an LC–MALDI based proteomic approach. *PROTEOMICS* 15:1598–1607.

Elsayyid M, Tanis JE, Yu Y. 2025. Simple In-Cell Processing Enables Deep Proteome Analysis of Low-Input Caenorhabditis elegans. *Anal. Chem.* 97:9159–9167.

Enard D, Cai L, Gwennap C, Petrov DA. 2016. Viruses are a dominant driver of protein adaptation in mammals. *eLife* 5:e12469.

Feyertag F, Berninsone PM, Alvarez-Ponce D. 2016. Secreted proteins defy the expression level–evolutionary rate anticorrelation. *Mol. Biol. Evol.*:msw268.

Gahmberg CG, Andersson LC. 1982. Role of Sialic Acid in the Mobility of Membrane Proteins Containing O-Linked Oligosaccharides on Polyacrylamide Gel Electrophoresis in Sodium Dodecyl Sulfate. *Eur. J. Biochem.* 122:581–586.

Gao X, Oei MS, Ovitt CE, Sincan M, Melvin JE. 2018. Transcriptional profiling reveals gland-specific differential expression in the three major salivary glands of the adult mouse. *Physiol. Genomics* 50:263–271.

Gervais EM, Desantis KA, Pagendarm N, Nelson DA, Enger T, Skarstein K, Liaaen Jensen J, Larsen M. 2015. Changes in the Submandibular Salivary Gland Epithelial Cell Subpopulations During Progression of S jögren's Syndrome-Like Disease in the NOD / S hi L t J Mouse Model. *Anat. Rec.* 298:1622–1634.

Hatakeyama S, Sashima M, Suzuki A. 1987. A sexual dimorphism of mucous cells in the submandibular salivary gland of rat. *Arch. Oral Biol.* 32:689–693.

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK. 2010. Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Mol. Cell* 38:576–589.

Hirata Y, Orth DN. 1979. Concentrations of Epidermal Growth Factor, Nerve Growth Factor, and Submandibular Gland Renin in Male and Female Mouse Tissue and Fluids*. *Endocrinology* 105:1382–1387.

Hiremath IS, Wagner CE. 2026. Mechanisms for Physicochemical Interaction between Mucus and Respiratory Viruses. *ACS Biomater. Sci. Eng.* 12:13–30.

Horeth E, Oyelakin A, Song E-AC, Che M, Bard J, Min S, Kiripolsky J, Kramer JM, Sinha S, Romano R-A. 2021. Transcriptomic and Single-Cell Analysis Reveals Regulatory Networks and Cellular Heterogeneity in Mouse Primary Sjögren's Syndrome Salivary Glands. *Front. Immunol.* 12:729040.

Hsieh LS, Wen JH, Miyares L, Lombroso PJ, Bordey A. 2017. Outbred CD1 mice are as suitable as inbred C57BL/6J mice in performing social tasks. *Neurosci. Lett.* 637:142–147.

Hutson JM, Niall M, Evans D, Fowler R. 1979. Effect of salivary glands on wound contraction in mice. *Nature* 279:793–795.

Ikemoto S, Matsushima Y. 1984. Sex and strain differences of mouse variant salivary proteins. *Differentiation* 26:55–58.

Iwasaki T, Tokumori M, Matsubara M, Ojima F, Kamigochi K, Aizawa S, Ogoshi M, Kimura AP, Takeuchi S, Takahashi S. 2023. A regulatory mechanism of mouse kallikrein 1 gene expression by estrogen. *Mol. Cell. Endocrinol.* 577:112044.

Karageorgiou C, Ruhl S, Gokcumen O. 2025. Convergent evolution through independent rearrangements in the primate amylase locus. Available from: http://biorxiv.org/lookup/doi/10.1101/2025.08.14.670395

Karn R, Chung A, Laukaitis C. 2013. Shared and Unique Proteins in Human, Mouse and Rat Saliva Proteomes: Footprints of Functional Adaptation. *Proteomes* 1:275–289.

Karn RC, Laukaitis CM. 2011. Positive Selection Shaped the Convergent Evolution of Independently Expanded Kallikrein Subfamilies Expressed in Mouse and Rat Saliva Proteomes.Welch JJ, editor. *PLoS ONE* 6:e20979.

Khoury A, Achinger-Kawecka J, Bert SA, Smith GC, French HJ, Luu P-L, Peters TJ, Du Q, Parry AJ, Valdes-Mora F, et al. 2020. Constitutively bound CTCF sites maintain 3D chromatin architecture and long-range epigenetically regulated domains. *Nat. Commun.* 11:54.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* 37:907–915.

Konttinen Y, Stegaev V, Mackiewicz Z, Porola P, Hänninen A, Szodoray P. 2010. Salivary glands - `an unisex organ'?: Salivary glands -'an unisex organ'? *Oral*

1027   *Dis.* 16:577–585.

1028   Krueger F, James F, Ewels P, Afyounian E, Weinstein M, Schuster-Boeckler B,
1029   Hulselmans G, Sclamons. 2023. FelixKrueger/TrimGalore: v0.6.10 - add
1030   default   decompression   path.   Available   from:
1031   https://zenodo.org/record/5127898

1032   Kryuchkova-Mostacci N, Robinson-Rechavi M. 2016. Tissue-Specificity of Gene
1033   Expression Diverges Slowly between Orthologs, and Rapidly between
1034   Paralogs.Ouzounis CA, editor. *PLOS Comput. Biol.* 12:e1005274.

1035   Kurabuchi S, Hosoi K, Gresik EW. 2001. Androgen Regulation of the Cellular
1036   Distribution of the True Tissue Kallikrein mK1 in the Submandibular Gland of
1037   the Mouse. *J. Histochem. Cytochem.* 49:801–802.

1038   Kurihara K, Nakanishi N, Tomomura A. 2011. Interference of kallikrein 1b26
1039   (klk1b26) translation by microRNA specifically expressed in female mouse
1040   submandibular glands: an additional mechanism for sexual dimorphism of
1041   klk1b26 protein in the glands. *Biol. Sex Differ.* 2:13.

1042   Lai C-W, Chen H-L, Tsai T-C, Chu T-W, Yang S-H, Chong K-Y, Chen C-M. 2016.
1043   Sexually Dimorphic Expression of eGFP Transgene in the Akr1A1 Locus of
1044   Mouse Liver Regulated by Sex Hormone-Related Epigenetic Remodeling. *Sci.*
1045   *Rep.* 6:24023.

1046   Lawrence MG, Lai J, Clements JA. 2010. Kallikreins on Steroids: Structure,
1047   Function, and Hormonal Regulation of Prostate-Specific Antigen and the
1048   Extended Kallikrein Locus. *Endocr. Rev.* 31:407–446.

1049   Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose
1050   program for assigning sequence reads to genomic features. *Bioinformatics*
1051   30:923–930.

1052   Lindeman RE, Murphy MW, Agrimson KS, Gewiss RL, Bardwell VJ, Gearhart MD,
1053   Zarkower D. 2021. The conserved sex regulator DMRT1 recruits SOX9 in
1054   sexual cell fate reprogramming. *Nucleic Acids Res.* 49:6144–6164.

1055   Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and
1056   dispersion for RNA-seq data with DESeq2. *Genome Biol.* 15:550.

1057   Malaker SA. 2024. Glycoproteomics: Charting new territory in mass spectrometry
1058   and glycobiology. *J. Mass Spectrom.* 59:e5034.

1059   Martin KR, Le HT, Abdelgawad A, Yang C, Lu G, Keffer JL, Zhang X, Zhuang Z,
1060   Asare-Okai PN, Chan CS, et al. 2024. Development of an efficient, effective,
1061   and economical technology for proteome analysis. *Cell Rep. Methods*
1062   4:100796.

1063   Maruyama C, Monroe M, Hunt J, Buchmann L, Baker O. 2019. Comparing human
1064   and mouse salivary glands: A practice guide for salivary researchers. *Oral*
1065   *Dis.* 25:403–415.

1066   Mendes FK, Vanderpool D, Fulton B, Hahn MW. 2021. CAFE 5 models variation in
1067   evolutionary rates among gene families.Robinson P, editor. *Bioinformatics*
1068   36:5516–5518.

1069   Moustardas P, Yamada-Fowler N, Apostolou E, Tzioufas AG, Turkina MV, Spyrou G.
1070   2021. Deregulation of the Kallikrein Protease Family in the Salivary Glands of

the Sjögren's Syndrome ERdj5 Knockout Mouse Model. *Front. Immunol.* 12:693911.

Mukaibo T, Gao X, Yang N, Oei MS, Nakamoto T, Melvin JE. 2019. Sexual dimorphisms in the transcriptomes of murine salivary glands. *FEBS Open Bio* 9:947–958.

Naqvi S, Godfrey AK, Hughes JF, Goodheart ML, Mitchell RN, Page DC. 2019. Conservation, acquisition, and functional impact of sex-biased gene expression in mammals. *Science* 365:eaaw7317.

Nguyen CQ, Sharma A, Lee BH, She J-X, McIndoe RA, Peck AB. 2009. Differential gene expression in the salivary gland during development and onset of xerostomia in Sjögren's syndrome-like disease of the C57BL/6.NOD-Aec1Aec2mouse. *Arthritis Res. Ther.* 11:R56.

Nikkanen J, Leong YA, Krause WC, Dermadi D, Maschek JA, Van Ry T, Cox JE, Weiss EJ, Gokcumen O, Chawla A, et al. 2022. An evolutionary trade-off between host immunity and metabolism drives fatty liver in male mice. *Science* 378:290–295.

Oppenheim FG, Hay DI. 1972. Further studies of human serum albumin in oral fluid. *Helv. Odontol. Acta* 16:22–26.

Pajic P, Landau L, Gokcumen O, Ruhl S. 2025. Saliva Protein Genes in Humans were Shaped During Primate Evolution.Sohail M, editor. *Genome Biol. Evol.* 17:evaf165.

Pajic P, Pavlidis P, Dean K, Neznanova L, Romano R-A, Garneau D, Daugherity E, Globig A, Ruhl S, Gokcumen O. 2019. Independent amylase gene copy number bursts correlate with dietary preferences in mammals. *eLife* 8:e44628.

Pajic P, Shen S, Qu J, May AJ, Knox S, Ruhl S, Gokcumen O. 2022. A mechanism of gene evolution generating mucin function. *Sci. Adv.* 8:eabm8757.

Paltoglou S, Das R, Townley SL, Hickey TE, Tarulli GA, Coutinho I, Fernandes R, Hanson AR, Denis I, Carroll JS, et al. 2017. Novel Androgen Receptor Coregulator GRHL2 Exerts Both Oncogenic and Antimetastatic Functions in Prostate Cancer. *Cancer Res.* 77:3417–3430.

Pardini LC, Taga R. 1996. Stereological Study of the Sexual Dimorphism in Mouse Submandibular Glands. *Okajimas Folia Anat. Jpn.* 73:119–124.

Pavlopoulou A, Pampalakis G, Michalopoulos I, Sotiropoulou G. 2010. Evolutionary History of Tissue Kallikreins.DeSalle R, editor. *PLoS ONE* 5:e13781.

Penschow JD, Drinkwater CC, Haralambidis J, Coghlan JP. 1991. Sites of expression and induction of glandular kallikrein gene expression in mice. *Mol. Cell. Endocrinol.* 81:135–146.

Price SA, Hopkins SSB, Smith KK, Roth VL. 2012. Tempo of trophic evolution and its impact on mammalian diversification. *Proc. Natl. Acad. Sci.* 109:7008–7012.

Reese RM, Harrison MM, Alarid ET. 2019. Grainyhead-like Protein 2: The Emerging Role in Hormone-Dependent Cancers and Epigenetics. *Endocrinology* 160:1275–1288.

Ruhl S. 2012. The scientific exploration of saliva in the post-proteomic era: from database back to basic function. *Expert Rev. Proteomics* 9:85–96.

Saitou M, Gaylord EA, Xu E, May AJ, Neznanova L, Nathan S, Grawe A, Chang J, Ryan W, Ruhl S, et al. 2020. Functional Specialization of Human Salivary Glands and Origins of Proteins Intrinsic to Human Saliva. *Cell Rep.* 33:108402.

Stopka P, Kuntová B, Klempt P, Havrdová L, Černá M, Stopková R. 2016. On the saliva proteome of the Eastern European house mouse (Mus musculus musculus) focusing on sexual signalling and immunity. *Sci. Rep.* 6:32481.

Thamadilok S, Choi K-S, Ruhl L, Schulte F, Kazim AL, Hardt M, Gokcumen O, Ruhl S. 2020. Human and Nonhuman Primate Lineage-Specific Footprints in the Salivary Proteome.Nowick K, editor. *Mol. Biol. Evol.* 37:395–405.

Treister NS, Richards SM, Lombardi MJ, Rowley P, Jensen RV, Sullivan DA. 2005. Sex-related Differences in Gene Expression in Salivary Glands of BALB/c Mice. *J. Dent. Res.* 84:160–165.

True JR, Haag ES. 2001. Developmental system drift and flexibility in evolutionary trajectories. *Evol. Dev.* 3:109–119.

Uhlén M, Karlsson MJ, Hober A, Svensson A-S, Scheffel J, Kotol D, Zhong W, Tebani A, Strandberg L, Edfors F, et al. 2019. The human secretome. *Sci. Signal.* 12:eaaz0274.

UniProt Consortium. 2025. UniProt: the Universal Protein Knowledgebase in 2025. *Nucleic Acids Res.* 53:D609–D617.

Van Leeuwen BH, Penschow JD, Coghlan JP, Richards RI. 1987. Cellular basis for the differential response of mouse kallikrein genes to hormonal induction. *EMBO J.* 6:1705–1713.

Varki A, Schauer R. 2009. Sialic Acids. In: Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Bertozzi CR, Hart GW, Etzler ME, editors. Essentials of Glycobiology. 2nd ed. Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press. Available from: http://www.ncbi.nlm.nih.gov/books/NBK1920/

Villanea FA, Peede D, Kaufman EJ, Añorve-Garibay V, Chevy ET, Villa-Islas V, Witt KE, Zeloni R, Marnetto D, Moorjani P, et al. 2025. The *MUC19* gene: An evolutionary history of recurrent introgression and natural selection. *Science* 389:eadl0882.

Wagner GP, Kin K, Lynch VJ. 2012. Measurement of mRNA abundance using RNA-seq data: RPKM measure is inconsistent among samples. *Theory Biosci.* 131:281–285.

Winter EE, Goodstadt L, Ponting CP. 2004. Elevated Rates of Protein Secretion, Evolution, and Disease Among Tissue-Specific Genes. *Genome Res.* 14:54–61.

Wolff J, Rabbani L, Gilsbach R, Richard G, Manke T, Backofen R, Grüning BA. 2020. Galaxy HiCExplorer 3: a web server for reproducible Hi-C, capture Hi-C and single-cell Hi-C data analysis, quality control and visualization. *Nucleic Acids Res.* 48:W177–W184.

Wong HLX, Qin H, Tsang SW, Zuo X, Che S, Chow CFW, Li X, Xiao H, Zhao L, Huang T, et al. 2019. Early life stress disrupts intestinal homeostasis via NGF-

1159   TrkA signaling. *Nat. Commun.* 10:1745.

1160   Xu D, Pavlidis P, Thamadilok S, Redwood E, Fox S, Blekhman R, Ruhl S,

1161   Gokcumen O. 2016. Recent evolution of the salivary mucin MUC7. *Sci. Rep.*

1162   6:31791.