# A NAPE–LRRK2 metabolic axis controls lysosomal homeostasis in Parkinson's disease

Francesca Palese[1,2*], Carmela Giachino[3], Sylvie Syan[1], Giuliana Ottonello[2], Giulia Sciandrone[4], Carolina Filipponi[4], Michele Lai[4], Andrea Armirotti[2], Michela Deleidi[3], Chiara Zurzolo[1,5*]

[1] Institut Pasteur, Université Paris Cité, CNRS UMR 3691, Membrane Traffic and Pathogenesis, F-75015 Paris, France

[2] Analytical Chemistry Facility, Istituto Italiano di Tecnologia, Via Morego 30, 16163 Genova, Italy.

[3] Institut Imagine, INSERM UMR1163 Paris Cite' University, 24 boulevard du Montparnasse, 75015, Paris, France

[4] Retrovirus Center, Department of Translational Research and New Technologies in Medicine and Surgery, University of Pisa, Pisa, Italy

[5] Department of Molecular Medicine and Medical Biotechnology, University of Naples Federico II, Naples, Italy

**Email:** Francesca.palese@pasteur.fr

Chiara.zurzolo@pasteur.fr

**Author Contributions:** Conceptualization: FP; Methodology: FP, CG, SS, GO, GS; Investigation: FP; Visualization: FP; Supervision: FP, ML, MD, AA; Funding acquisition: FP, CZ; Writing – original draft: FP; Writing – review & editing: FP, MD, CZ.

**Competing Interest Statement:** No competing interests.

**Keywords:** N-acylphosphatidylethanolamines (NAPE), LRRK2, Lysosomes, Lipid signaling, Parkinson's disease

**Abstract**

*N*-acylphosphatidylethanolamines (NAPEs) are atypical glycerophospholipids that accumulate in response to cellular stress, yet their roles beyond serving as precursors of fatty-acid ethanolamines (FAEs) remain largely unexplored. Here, we identify NAPEs as endogenous regulators of leucine-rich repeat kinase 2 (LRRK2), a master controller of lysosomal homeostasis and a genetic driver of Parkinson's disease. We show that increasing NAPE synthesis or blocking their hydrolysis inhibits LRRK2 kinase activity, enhances lysosomal function, and promotes the clearance of α-synuclein aggregates. Conversely, cells in which NAPE hydrolysis is enhanced display increased LRRK2 activation and lysosomal dysfunction. Importantly, in induced pluripotent stem cell–derived dopaminergic neurons carrying the LRRK2-G2019S variant, pharmacological inhibition of NAPE-PLD – the enzyme that degrades NAPEs – restores lysosomal activity and favors the clearance of preformed α-synuclein fibrils. Together, our findings identify NAPEs as previously unrecognized lipid regulators of LRRK2 signaling and lysosomal function, revealing a metabolic axis with therapeutic potential in Parkinson's disease.

**Main Text**

**Introduction**

*N*-acylphopshpatidylethanolamines (NAPEs) are low-abundance glycerophospholipids present in the membranes of all eukaryotic cells (1). They are synthesized through the transfer of an acyl group from phosphatidylcholine (PC) to phosphatidylethanolamine (PE) by calcium-independent phospholipase A and acyltransferases (PLAATs), which maintain basal NAPE levels (2), or by the calcium-dependent phospholipase PLA2G4E, which is activated in a stimulus-dependent manner to markedly increase NAPE production under pathological conditions (3, 4). NAPEs are hydrolyzed by the NAPE-specific phospholipase D (NAPE-PLD) (5, 6), generating fatty-acid ethanolamines (FAEs) and phosphatidic acid (PA).

FAEs mediate diverse physiological and pathological processes, including neurotransmission (7, 8), pain (9), energy homeostasis (10, 11) and inflammation (12, 13), through receptors such as cannabinoid receptors, TRPV1, GPR119 and PPAR-α. Consequently, NAPEs have traditionally been viewed mainly as precursors of FAEs rather than molecules with intrinsic biological activity. However, NAPE production is dynamically regulated by physiological (7, 14) and pathological stimuli (15–17) including ischemia (18), excitotoxicity (19) and neurotoxic insults (20–22). Under pathological conditions, NAPE levels often exceed those of FAEs, suggesting functions independent of downstream FAE signaling. Consistent with this idea, NAPEs have been shown to stabilize cellular membranes (23, 24), regulate membrane fusion (25) and lipid raft organization (26), thereby influencing protein association at the membranes (27, 28). Notably, NAPEs accumulate in the brains of mouse models of Parkinson's disease (PD) (21, 29), where they exert neuroprotective effects against 6-OHDA-induced toxicity (29). In contrast, circulating NAPE levels are decreased in the plasma of female PD patients (30).

These observations suggest that NAPEs may participate in adaptive lipid response to neuronal stress in PD. However, the molecular mechanisms through which NAPEs influence neuronal survival and disease-relevant pathways remain largely unknown.

One potential link between NAPE signaling and PD pathogenesis is the regulation of membrane-associated signaling proteins that control endolysosomal trafficking. Among there, leucine-rich

repeat kinase 2 (LRRK2) has emerged as a central regulator of endolysosomal dynamics and
lysosomal function whose dysregulation is strongly linked to PD (31). Mutations in
the *LRRK2* gene represent one of the most common genetic causes of PD, accounting for ~1-2%
of all cases worldwide but reaching up to 30-40% of cases in certain founder populations (32).
Previous work   recruitment (28). Here, we investigate the molecular mechanism underlying
NAPE-mediated neuroprotection, focusing on LRRK2 regulation. We find that NAPE
accumulation — either through enhanced synthesis or inhibition of their hydrolysis — reduces
LRRK2 kinase activity and improves lysosomal function, thereby promoting the clearance of α-
synuclein pre-formed fibrils (PFFs). Conversely, overexpression of the hydrolyzing enzyme
NAPE-PLD increases LRRK2 protein abundance and kinase activity, impairs lysosomal function,
and decreases α-synuclein PFF clearance.

Importantly, in induced pluripotent stem cell-derived dopaminergic neurons carrying the
hyperactivating LRRK2-G2019S variant, pharmacological inhibition of NAPE-PLD restores
lysosomal activity and enhances α-synuclein PFF clearance. Together, these findings identify
NAPEs as previously unrecognized regulators of LRRK2 activity and lysosomal function, with
direct implications for PD pathogenesis, and highlight NAPE-PLD inhibitors as promising
therapeutic candidates.

**Results**
89
***NAPEs accumulation potentiates lysosomal activity via LRRK2 inhibition***

To investigate the effects of NAPE accumulation in a neuronal context, we genetically modulated
NAPE metabolism in catecholaminergic SH-SY5Y cells using two approaches: (i) overexpression
of the synthesizing enzyme PLA2G4E and (ii) knockdown of the hydrolyzing enzyme NAPE-PLD
(**Fig. 1A**). Using a CRISPR-Cas9 strategy we obtained a partial knockdown (KD), resulting in an
approximatively 30% reduction in *NAPE-PLD* mRNA (**Fig. 1B**), and a 30-50% decrease in protein
levels (depending on the technique used for the measurement, **Fig. 1C, D**). This led to a 30%
increase in total NAPE levels (**Fig. 1E**). Consistent with a hypomorphic condition, no significant
compensatory changes were observed in *PLA2G4E* mRNA or protein (**Fig. S1 A-C**).

In parallel, SH-SY5Y cells overexpressing (OE) the synthesizing enzyme PLA2G4E (33)
displayed a massive upregulation of both mRNA (~$3x10^4$-fold) and protein (~30-fold) levels **(Fig
1F-H**). In line with a strong potentiation of the synthetic pathway, these cells exhibited a 70-fold
increase in total NAPE content (**Fig. 1 I**). Notably, NAPE-PLD protein levels were downregulated
in these conditions (**Fig. S1 E, F**), likely reflecting an adaptive response aimed at limiting
excessive FAE production.

Previous studies in NAPE-PLD knockout mouse brains and in NAPE-PLD silenced neuronal cells
showed a reduced LRRK2 fraction associated with membranes (28); however, the underlying
mechanisms and functional consequences of this observation were not investigated. Here, we
show that both NAPE-PLD KD and PLA2G4E OE reduce LRRK2 abundance in neuronal cells
(**Fig. 2 A**). Moreover, the LRRK2 kinase activity measured by western blot using phosphorylation
of Rab8a at T72 — a LRRK2-specific phosphorylation site — was lowered in both NAPE-PLD KD
and PLA2G4E OE cells compared with WT SH-SY5Y cells (**Fig. 2 C**). Furthermore, we observed
diminished LRRK2 localization at lysosomes (**Fig. 2B, D**) in both NAPE-PLD KD and PLA2G4E
OE cells relative to WT cells. Given that LRRK2 can be recruited to lysosomes in response to

lysosomal stress or damage (34, 35) this is consistent with improved lysosomal health under
conditions of NAPE accumulation.

LRRK2 is an established regulator of lysosomal degradative activity. LRRK2 knockout or kinase
inhibition has been shown to increase lysosomal proteolytic activity, whereas pathogenic,
hyperactive LRRK2 mutations associated with PD impair lysosomal degradative activity (36–38).
In accordance with an inhibitory function of LRRK2 on lysosomal activity, we observed increased
lysosomal cathepsin B activity in cells where NAPEs accumulate and LRRK2 is decreased in a
NAPE-dose-dependent manner (+20% in NAPE-PLD KD cells and + 31% in PLA2G4E OE cells;
**Fig. 2E**). Interestingly, treatment with the calcium ionophore ionomycin - which is reported to
stimulate the production of NAPEs (39)(**Fig. S2A**)- further improved lysosomal activity in both
NAPE-enriched cell lines (**Fig. S2B**).

Since increased lysosomal enzyme activity can also occur as a compensatory response to
impaired autophagosome-lysosome fusion, we assessed autophagic flux to avoid
misinterpretation. Cells were treated with the autophagosome-lysosome fusion inhibitor
Bafilomycin A1, and p62-Lamp1 colocalization was measured. Under baseline conditions we did
not observe any significant difference among cell lines (**Fig. S3A**). As expected for autophagic
flux, Bafilomycin A1 treatment resulted in increased accumulation of p62-Lamp1-positive
structures in all conditions (**Fig. S3A**), indicating that autophagosome-lysosome fusion is
functional in our cell models. These results support the conclusion that NAPE accumulation
enhances lysosomal activity rather than reflecting impaired autophagic flux.

Furthermore, no differences were observed in TFEB nuclear localization or total levels across cell
lines (**Fig. S3B, C**), indicating that NAPE-mediated lysosomal protection operates independently
of TFEB activation, the master transcriptional regulator of lysosomal biogenesis (40).

***NAPEs-enriched cells clear exogenous pre-formed $\alpha$-synuclein fibrils more efficiently***

We next investigated the effect of NAPE accumulation under pathological conditions, specifically
following exposure to exogenous pre-formed $\alpha$-synuclein ($\alpha$-syn) fibrils (PFFs). $\alpha$-Syn PFFs are
well known to induce lysosomal stress and activate LRRK2 kinase activity in neurons (41, 42),
thereby establishing a negative feedback loop on lysosomal activity. Interestingly, following
overnight exposure of WT, NAPE-PLD KD and PLA2G4E OE cells to $\alpha$-syn PFFs we detected a
~2-fold increase in LRRK2 protein levels in WT cells, whereas the increase was very modest in
NAPE-enriched cells (+4% and +34% in NAPE-PLD KD and PLA2G4E OE cells, respectively;
**Fig. 3 A**). These data support NAPE-mediated negative regulation of LRRK2 protein levels, in
addition to reduced LRRK2 recruitment to lysosomes (**Fig. 3 B**). Furthermore, consistent with
their enhanced lysosomal activity, both NAPE-PLD KD and PLA2G4E OE cells showed fewer $\alpha$-
syn aggregates after overnight PFF treatment, followed by a 6-h washout period, compared with
WT SH-SY5Y cells (**Fig. 3 C**).

***Lysosomal activity and $\alpha$-synuclein fibrils clearance are impaired in NAPE-PLD-***
***overexpressing cells***

To strengthen these findings, we next generated SH-SY5Y cells overexpressing NAPE-PLD. In
these cells, *NAPE-PLD* mRNA and protein levels increased by approximately $10^3$-fold and 5-fold
respectively (**Fig. 4 A-D**), a condition expected to markedly enhance NAPE hydrolysis at cellular
membranes under basal condition (43)(**Fig. 4 A**). In line with the non-rate limiting role of NAPE-

PLD in FAE biosynthesis (44), no major compensatory effects were detected in PLA2G4E
expression levels (**Fig. S4 A-C**).

In contrast to NAPE-enriched cells, NAPE-PLD OE cells exhibited elevated LRRK2 protein levels
and kinase activity (**Fig. 4 E, F**). These cells displayed markedly reduced cathepsin B activity
relative to WT (77% of WT levels; **Fig. 4H**), consistent with LRRK2's known suppression of
lysosomal hydrolases. Notably, ionomycin treatment -which boosts NAPE production and, in
NAPE-PLD-overexpressing cells, is expected to promote robust FAEs generation- failed to
rescue cathepsin B activity (**Fig. 4H**). This suggests that lysosomal impairment is driven by
altered NAPE levels rather than by FAE or phosphatidic acid byproducts. Importantly, the
lysosomal impairment observed in NAPE-PLD OE cells was reversed by treatment with LEI-401
(a NAPE-PLD inhibitor) (**Fig. S5 A**). Furthermore, when LEI-401 and ionomycin treatments were
combined, lysosomal activity reached 136% of WT levels, consistent with maximal NAPE
accumulation (**Fig. 4H**). In line with our observation of elevated LRRK2 activity in NAPE-PLD OE
cells, overnight treatment with the LRRK2 pharmacological inhibitor MLi-2, not only improved
lysosomal activity in WT cells, but it also rescued the lysosomal impairment detected in NAPE-
PLD OE cells (**Fig. S6**), supporting the conclusion that NAPE levels regulate lysosomal activity
through modulation of LRRK2 abundance and kinase activity.

Following Bafilomycin A1 treatment, NAPE-PLD OE cells showed reduced p62-Lamp1
colocalization compared to WT cells (**Fig. S7A**), indicating impaired autophagosome-lysosome
fusion consistent LRRK2 overexpression (45). Moreover, these cells exhibited elevated TFEB
levels in both cytosolic and nuclear compartments (**Fig. S7B-E**), suggesting a compensatory
response attempting restore lysosomal homeostasis.

Finally, following overnight treatment with α-syn PFFs, LRRK2 protein levels increased further in
NAPE-PLD OE cells (~3-fold relative to untreated WT cells and ~1.5-fold relative to WT cells
treated with α-syn; **Fig. 5A**), in contrast to NAPE-enriched cells, where the α-syn-induced
increase in LRRK2 levels was contained. Consistent with reduced lysosomal degradative
capacity, the number of α-syn aggregates after a 6-hour washout period were higher in NAPE-
PLD OE SH-SY5Y cells than in WT cells (**Fig. 5 B**). Together, these results identify a previously
unrecognized role for NAPEs in regulating lysosomal activity and the handling of protein
aggregates in neurons through modulation of LRRK2 activity.

***NAPE-PLD inhibition restores lysosomal activity and α-synuclein fibril clearance in***
***G2019S-LRRK2 dopaminergic neurons***

We next tested the therapeutic potential of NAPE-PLD inhibition in human iPSC-derived
dopaminergic neurons (hDANs) carrying the PD-linked LRRK2-G2019S mutation and their
isogenic control line. This mutation is a gain-of-function variant that increases LRRK2 kinase
activity (46), resulting in impaired lysosomal function (47). Between differentiation day 21-24,
when hDANs express neuronal markers (TUJ1 and MAP2) and the dopaminergic marker tyrosine
hydroxylase (TH; **Fig. S8 A, B**), cells were treated overnight with the NAPE-PLD inhibitor LEI-401
prior to measuring lysosomal activity using the Magic Red Cathepsin B assay. Consistent with
LRRK2-mediated lysosomal impairment, G2019S-LRRK2 neurons displayed fewer active
lysosomes than isogenic controls, an effect rescued by LEI-401 treatment (**Fig. 6 A, B**). These
findings indicate that increasing NAPE levels through NAPE-PLD inhibition counteracts LRRK2
hyperactivation and restore lysosomal function.

We next assessed whether this rescue translated into improved handling of pathological α-syn.
hDANs were treated with α-syn PFFs in the presence or absence of LEI-401. Both G2019S-
LRRK2 neurons and isogenic controls efficiently internalized PFFs after 48 h incubation (**Fig. 6
C**), with comparable baseline uptake between genotypes. Strikingly, LEI-401 treatment
significantly reduced the number of intracellular α-syn aggregates per cell in both mutant and
control neurons compared to untreated conditions, demonstrating enhanced aggregate
clearance.  Together, these results establish NAPE-PLD inhibition as a mechanism to restore
lysosomal function and promote pathological protein clearance in human dopaminergic neurons.
These findings identify NAPEs as upstream modulators of LRRK2-dependent lysosomal
dysfunction and highlight NAPE-PLD as a promising therapeutic target for Parkinson's disease
and related neurodegenerative disorders characterized by impaired lysosomal proteostasis.

210
211   **Discussion**
212
In the present study, we identify NAPEs as previously unrecognized lipid regulators of LRRK2
signaling and lysosomal function in neuronal cells and human dopaminergic neurons. By
genetically and pharmacologically manipulating NAPE synthesis and hydrolysis, we define a lipid-
kinase axis in which NAPE accumulation (i) restrains LRRK2 kinase activity, (ii) preserves
lysosomal degradative capacity, and (iii) promotes α-synuclein aggregate clearance, whereas
NAPE depletion has the opposite effect. These findings provide a mechanistic framework linking
stress-induced NAPE production to endolysosomal homeostasis and suggest that NAPE-PLD is a
viable target to counteract LRRK2 hyperactivation in PD.

LRRK2 is increasingly recognized as a stress-responsive regulator of lysosomes, recruited to
enlarged or damaged lysosomes where it phosphorylates Rab GTPases and constrains
lysosomal degradative activity (36, 48–50). Our data place NAPEs upstream of this pathway:
elevating NAPEs by PLA2G4E overexpression or NAPE-PLD knockdown reduces LRRK2
abundance and Rab8a T72 phosphorylation and enhances lysosomal cathepsin B activity,
whereas NAPE-PLD overexpression increases LRRK2 protein and kinase output and impairs
lysosomal function. These manipulations support a model in which NAPE levels set a "lipid tone"
that tunes LRRK2 activation and, consequently, lysosomal capacity (35, 36, 38). Mechanistically,
this aligns with prior observations that NAPE accumulation reduces LRRK2 partitioning to
membrane fractions, whereas enhanced NAPE hydrolysis promotes membrane association (28).
NAPEs stabilize and remodel membranes (23, 24), regulate fusion (25), and alter lipid raft
organization (26), so changes in local NAPE content could influence LRRK2 recruitment to
membrane microdomains supporting its activation. Unaltered TFEB responses in NAPE-enriched
models suggest direct membrane effects, stabilizing fusion curvature or disrupting LRRK2-Rab
raft recruitment. Our data indeed do not yet discriminate whether NAPEs act directly on LRRK2
and its scaffolds or indirectly through altered Rab or organelle organization, but the tight
correlation between NAPE levels, LRRK2 activity, and lysosomal performance supports a
functionally integrated module.

NAPE-PLD hydrolyzes NAPEs into bioactive FAEs such as OEA and PEA, as well as
phosphatidic acid (PA), both of which can influence lysosomal function (51, 52). FAEs can
promote lysosomal biogenesis via PPAR-α–TFEB signaling (53, 54),while PA modulates
membrane properties in a concentration-dependent manner (55). However, our data argue that
NAPEs themselves are the primary drivers of the observed effects. In NAPE-PLD-overexpressing
cells, ionomycin-induced NAPE synthesis, which increases FAEs and PA, fails to restore

lysosomal activity, whereas pharmacological NAPE accumulation via LEI-401 robustly enhances
it. Although LRRK2 inhibition (MLi-2) partially rescues lysosomal defects, maximal restoration
requires combined NAPE elevation, indicating both LRRK2-dependent and -independent
contributions. Together, these findings support a model in which NAPEs act as stress-inducible
membrane lipids with autonomous regulatory functions, effectively constraining LRRK2 activity
and tuning lysosomal competence under proteotoxic conditions.

Pathogenic LRRK2 mutations, particularly G2019S, enhance kinase activity and drive lysosomal
dysfunction, impaired autophagy, and α-synuclein accumulation (38, 47). While LRRK2 kinase
inhibitors can partially restore lysosomal function and reduce α-synuclein burden (56), our results
extend this paradigm by identifying NAPEs as endogenous modulators of this axis. NAPE
accumulation attenuates PFF-induced LRRK2 upregulation and enhances α-synuclein clearance
in neuronal cells, whereas NAPE-PLD overexpression exacerbates both phenotypes. Importantly,
in human iPSC-derived dopaminergic neurons carrying the G2019S mutation, NAPE-PLD
inhibition restores lysosomal activity and reduces aggregate burden to levels comparable to
isogenic controls. These findings integrate LRRK2 hyperactivation and α-synuclein pathology into
a unified lipid-sensitive framework and suggest that modulation of NAPE metabolism can
normalize lysosomal function even in a genetically sensitized context.

From a therapeutic perspective, NAPE-PLD inhibition emerges as a promising strategy to restore
lysosomal proteostasis. By increasing endogenous NAPE levels, this approach enhances
lysosomal function and promotes α-synuclein clearance while potentially avoiding some of the
limitations associated with direct LRRK2 kinase inhibition. Small-molecule NAPE-PLD inhibitors
effectively elevate NAPEs without compensatory increases in FAEs (51), supporting their utility as
pharmacological tools. At the same time, given the broader roles of NAPEs and FAEs in brain
physiology, including regulation of feeding, reward, and anxiety (57, 58), careful in vivo studies
will be required to define therapeutic windows and assess long-term effects. Nonetheless,
targeting a stress-responsive lipid pathway offers a conceptually distinct and potentially
complementary strategy to address convergent lysosomal defects observed in both genetic and
sporadic PD (56).

While our study provides a robust framework for NAPE-mediated regulation of LRRK2 and
lysosomal function, the study has some limitations. First, most mechanistic experiments were
performed in SH-SY5Y-derived neuronal cells, which only partially recapitulate mature nigral
dopaminergic neuron biology. The rescue of lysosomal activity and α-synuclein handling in
human G2019S dopaminergic neurons argues for disease relevance, but confirmation in in vivo
(e.g., G2019S knock-in mice or α-synuclein PFF–injected rodents) will be important. Second, we
focused on cathepsin B activity and aggregate counts; future studies should expand to broader
lysosomal hydrolases, lysosomal pH, and TFEB/MiT–TFE transcriptional programs modulated by
LRRK2 (36, 59). Finally, the molecular interface between NAPEs and LRRK2 remains to be
defined; which NAPE species are responsible, whether NAPEs alter Rab/scaffold recruitment (35,
48, 50), and how NAPE dynamics interact with other lipid pathways, including
glucocerebrosidase-dependent sphingolipid metabolism. Addressing these questions will clarify
how a stress-inducible membrane lipid precisely controls a disease-critical kinase and may reveal
new opportunities to fine-tune lysosomal function in neurodegeneration.

287
288 **Materials and Methods**
289

### Chemicals

MLi-2 LRRK2 inhibitor was purchased from Selleckchem (S9694). Bafilomycin A1 was purchased by Sigma-Aldrich (SML1661). Lei-401 NAPE-PLD inhibitor was kindly donated by Professor Van der Stelt (Leiden University), DOPE-N-Nonadecanoyl standard was purchased by Larodan (38-9091). Ionomycin was purchased by Sigma-Aldrich (I3909).

### Culture of Cell lines and treatments

Human neuroblastoma cell line SH-SY5Y (SK-N-SH-derived, hereafter "neuronal cells") were cultured in RPMI1640 media (Euroclone, ECB2000L) and supplemented with 10% fetal calf serum (FCS) (Eurobio Scientific, CVFSVF00-01) and 1% penicillin-streptomycin (PS, Gibco, 15140-122; 100 units/mL final concentration). Cells were maintained at 37°C in a humidified $CO_2$ incubator, and passaged using 0.05% Trypsin-EDTA (Gibco, 25300-054) when they reached 90% confluence. Cells were counted before each experiment, and seeded on autoclaved, 12 mm glass coverslips (uncoated) (Epredia, CB00120RA120MNZ0) for fixed-cell imaging, or on 35 mm glass-bottom microdishes (Ibidi GmbH, 80807-90) for live-cell imaging. Cells between passages 3 and 10 were used for experiments. MLi-2 LRRK2 inhibitor (Selleckchem) was dissolved in DMSO and diluted in full growth media to a final concentration of 10nM and incubated over-night. Bafilomycin A1 (Sigma-Aldrich, 500X) was dissolved in fresh full growth media and incubated for 4h prior to further analysis. LEI-401 was dissolved in in DMSO and diluted in full growth media to a final concentration of 1 μM and incubated over-night. Ionomycin was diluted in full growth media to a final concentration of 5 μM and incubated 30 minutes.

### Generation of NAPE-PLD knock-down cell lines

The NAPE-PLD Human Gene Knockout Kit (CRISPR, Origene, KN409877) was used following manufacturer's instructions. Lipofectamine™ 3000 Transfection Reagent (Thermo Fisher Scientific, L3000001) was used following standard protocol in OptiMEM media supplemented with 1% PS and 1% FCS. One day before transfection, SH-SY5Y cells were seeded in a six-well plate to reach 80-90% confluence at the time of transfection. Before transfection, culture medium was aspirated and 1715 μL of transfection medium was added. A mixture of donor DNA (10 μL per well), gRNA (20 μL per well), P3000 reagent (2 μL per well), and Lipofectamine 3000 (3 μL per well) was prepared in transfection medium (250 μl each) and incubated (15 min, room temperature). Transfection was performed by dropwise addition of the transfection mixture to the cells. Then, 6 h post-transfection, the culture medium was replenished with full growth media for additional 48 h. Cells were selected in the medium containing 1 μg/mL puromycin. Clonal cell lines were selected by clonal dilution. Three clones were used throughout this study.

### Generation of PLA2G4E-overexpressing cell lines

The pcDNA5/TO vector harboring EGFP-FL-PL2G4E was kindly donated by Professor Uyama (Kagawa University, (33)) and transfected into SH-SY5Y cells using 5 μg of DNA per well and Lipofectamine 3000 in OptiMEM media for 6h as described above. After 48 h from transfection, cells were selected with the medium containing 500 μg/mL Geneticin (Gibco, 10131-027). Clonal cell lines were generated by clonal dilution and maintained in the medium containing geneticin. Three of the clones expressing EGF-PLA2G4E were used throughout this study.

### Generation of NAPE-PLD-overexpressing cells

Lentiviral particles were produced by transient transfection of HEK293T cells using
polyethyleneimine (PEI) (Sigma, #306185). Cells were seeded in 12-well plates at a density of 1.7
× 10□ cells per well and incubated for 24 hours to reach 70–80% confluency. Transfection
mixtures were prepared in DMEM without FBS: the DNA mix contained 400 ng psPAX2
packaging plasmid (Addgene #12260), 350 ng VSV-G pseudotyping plasmid (Addgene #14888),
300 ng pRSV-Rev plasmid (Addgene #12253), and 650 ng NAPE-PLD-pLenti-C-mGFP-P2A-Puro
transfer plasmid (OriGene #PS100093) (Fig. 1). A second mixture contained PEI diluted 1:50 in
DMEM without FBS. The two mixtures were combined at a 1:1 ratio, vortexed gently, and
incubated at room temperature for 20 minutes to allow PEI–DNA complex formation. Cell culture
medium was replaced with fresh complete medium (DMEM + 10% FBS), and the transfection
mixture was added dropwise to each well (total volume 100 µL per well). Cells were incubated at
37°C, 5% $CO_2$ for 48 hours. Supernatants were then harvested, centrifuged at 3,000 rpm (1,100
× g) for 15 minutes at 4°C to remove cellular debris, and the clarified lentiviral particle-containing
supernatant was aliquoted, snap-frozen in liquid nitrogen, and stored at –80°C. Viral titers were
not formally quantified but estimated as sufficient for downstream transduction based on pilot
GFP expression assays.

Stable SH-SY5Y NAPE-PLD-overexpressing cells were generated by transduction with the
harvested lentiviral supernatant. SH-SY5Y cells were seeded in 12-well plates at 2 × 10□ cells
per well and incubated for 24 hours. Medium was replaced with lentiviral supernatant
supplemented with polybrene (Sigma, #H9268; final 8 µg/mL, 1:100 dilution from 0.8 mg/mL
stock). Plates underwent spinoculation at 1,100 × g for 1 hour at 37°C to enhance transduction
efficiency. Following spinoculation, plates were incubated for an additional 2 hours at 37°C, after
which the medium was replaced with fresh DMEM/F-12 (1:1) supplemented with 2% FBS. Cells
were recovered for 72 hours before initiating selection. Puromycin selection was performed in two
sequential rounds: first at 5 µg/mL for 5 days, followed by 10 µg/mL for 7 days, with medium
changes every 2–3 days. Transduction efficiency and selection success were confirmed by
visualizing GFP fluorescence under a fluorescence microscope (Operetta CLS, Revvity, Milan).
Selected polyclonal populations exhibited >95% GFP positivity and were expanded for
downstream experiments.

***Cell culture experiments with hiPSC lines***

Dopaminergic neurons were generated from neural progenitor cells (NPCs) according to an
established protocol (60). Briefly, NPCs at 80% confluency were dissociated using Accutase for 5
minutes at 37 °C and collected via centrifugation (1100 rpm, 5 min). Cells were then seeded onto
matrigel-coated 6-well plates (1:10 dilution in DMEM/F12) at a density of 1.0 × 10□ cells/well.
Differentiation was initiated at day 0 by transitioning to differentiation medium. On day 6, the cells
were passaged and re-seeded at 3.0 × 10□ cells/well. Neuronal maturation began on day 8 with
the addition of 0.5 µM PMA, which was subsequently withdrawn on day 10. On day 14, the
dopaminergic neurons were harvested and re-plated at a density of 2.6 × 10□ cells/cm, according
to the experimental design planned. On day 21, cells were treated overnight with 1□µM LEI-401
and subsequently tested with magic red assay. Cells were treated with α-syn PFFs in the
presence of vehicle or 1$\mu$M LEI-401 for 48 h, followed by fixation and immunofluorescence
analysis.

***Preparation of α-syn aggregates***

α-Syn aggregates were prepared as described before (61). Briefly, human wild-type α-syn was
purified from *Escherichia coli* BL21 (DE3) with RP-HPLC. Fibrils were then conjugated with Alexa
Fluor™ 647 fluorophore (used in imaging experiments) (Invitrogen) using manufacturer's labelling
kit or not tagged with a fluorophore. Fibrils were kept at -80°C for long-term storage. Prior to
exposure to cells, fibrils were diluted in growth medium at a working concentration of 500 nM and
sonicated (BioBlock Scientific, Vibra Cell 7504) for 5 minutes at an amplitude of 80%, pulsed for 5
seconds "on" and 2 seconds "off". Sonicated fibrils were then added on cells directly without the
addition of any intracellular delivery agents for designated time points.

### *Immunocytochemistry*

Immunofluorescence on cells were performed using the standard protocol. Cells were fixed with
4% paraformaldehyde (PFA [Electron Microscopy Sciences, 15710]) for 20 minutes at room
temperature (RT), followed by three washes with 1X DPBS. Cells were incubated in 1X DPBS-Tx
(0.1% Triton-X100 [Sigma Aldrich, 9002-93-1] in 1X DPBS) for 3 minutes at RT, followed by
incubation with blocking solution of 2% bovine serum albumin (BSA [Sigma Aldrich, A9647])
prepared in 1X DPBS for 1h at RT. Cells were then incubated with primary antibody overnight at
4°C. The following primary antibodies were used: rabbit anti-NAPE-PLD (Sigma Aldrich,
HPA024338, 1:200), rabbit anti-PLA2G4E (Proteintech, 18088-1-AP, 1:400), rabbit anti-LRRK2
(Abcam, ab133474; 1:500) mouse anti-LAMP1 (Developmental Studies Hybridoma Bank, H4A3;
1:100), guinea pig anti-p62 (Progen, GP62-C; 1:500), rabbit anti-TFEB (Cell Signaling
Technology, 4240; 1:100), chicken anti-MAP2 (BioLegend, 82250, 1:10,000), rabbit anti-TH (Pel-
Freez, P40101, 1:1,000), and mouse anti-βIII-tubulin (BioLegend, 801202, 1:1,000). The next
day, cells were washed three times with DPBS, followed by incubation with respective secondary
antibodies (all from Thermo fisher Scientific, dilution of 1:500 in blocking solution) for 1h at RT.
Cells were then washed three times with 1X DPBS, counterstained for nuclei with DAPI (1:1000
in PBS [Sigma-Aldrich, D9542]) and mounted on glass slides with Aqua-Poly/Mount
(Polysciences Inc., 18606-20). Slides were imaged at least a day after mounting of coverslips.

### *Magic red CTSB assay*

Intracellular CTSB activity was measured in SH-SY5Y cells and dopaminergic iNeurons using the
Magic Red™ CTSB assay kit from Abcam, ab270772, according to the manufacturer's protocol.
Cells were plated in glass-bottom matteks (Ibidi GmbH, 80807-90) the day before the experiment
to reach a 70-80% confluence for neuronal cells, or two weeks before for iNeurons. Culture
medium was gently removed and replaced with pre‐warmed medium containing Magic Red
substrate and Hoechst 33342 (Abcam, AB270772, 1:100). Ionomycin was diluted in the same
media containing Magic Red to a final concentration of $1\mu$M. Cells were incubated at 37 °C, 5%
$CO_2$ for 30 min, protected from light. Following incubation, media was replaced with Phenol-free
media and immediately live-imaged using a Nikon Eclipse Ti2 microscope with a spinning disk
confocal set-up (Yokogawa) equipped with a live-cell incubation chamber that maintained the
samples at 37 °C in a humidified atmosphere of 5% $CO_2$ throughout imaging. Cells were imaged
using a 100X oil immersion objective (1.45 numerical aperture). Optical sections of 0.5 $\mu$m
interval were taken to acquire the whole volume of the cells. Exposure time and gain were kept
constant across all experimental conditions within each experiment.

### *Immunoblotting*

Cell pellets were homogenized in 100 µl of radioimmunoprecipitation assay (RIPA) buffer
consisting of 50 mM Tris-HCl (pH 7.4), 1% TritonX-100, 0.5% sodium-deoxycholate, 0.1%
sodium dodecyl sulphate, 150 mM sodium chloride and 2 mM ethylenediaminetetraacetic acid,
supplemented with 1X protease inhibitor (cOmplete Mini, EDTA-free; Sigma, 11836170001).
Protein concentrations were measured using Bradford's method, following manufacturer's
instructions (Thermo Fisher Scientific). Proteins (20 µg) were denatured in SDS (8%) and β-
mercaptoethanol (5%) at 95 °C for 10 minutes. After separation by SDS-PAGE on a 4-20% gel
(Biorad, 4568094), proteins were electro-transferred to nitrocellulose membranes. The
membranes were blocked with Tropix I-Block buffer (Applied Biosystems, Thermo Fisher
Scientific, T2015) and incubated overnight with primary antibodies (anti-NAPE-PLD, Sigma-
Aldrich, HPA024338, 1:500; anti-PLA2G4E, Proteintech, 18088-1-AP, 1:1000; anti Rab8a,
Abcam, Ab188574, 1:500; anti-pT72Rab8a, Abcam, Ab230260, 1:1000; anti-GAPDH antibody
Sigma-Aldrich, G9545, 1:5000; anti-Actin, MP Biomedicals, 691001, 1:1000) in blocking buffer,
followed by incubation with horseradish peroxidase-conjugated secondary antibody (Millipore,
1:5000) in TBS containing 0.1% Tween-20 at room temperature for 1 h. Finally, proteins were
visualized using an ECL kit (Thermo Fisher Scientific, 34580) and chemiluminescence images
were acquired using GE Amersham Imager AI680 analyzer.

*mRNA isolation and qPCR*

Total RNA was prepared from cell pellets using the Ambion PureLink RNA minikit (Invitrogen,
Thermo Fisher Scientific, 12183018A) as directed by the supplier. Samples were treated with
DNase (PureLink DNase, Invitrogen, Thermo Fisher Scientific, 12185010) and cDNA synthesis
was carried out using the high-capacity cDNA reverse transcription kit (Applied Biosystems,
Thermo Fisher Scientific 4368814) according to the manufacturer's protocol using purified RNA
(0.5-1 µg). First-strand cDNA was amplified using the iTaq universal SYBR Green supermix (Bio-
Rad, 1725124). Copy numbers of cDNA targets were quantified at the cycle threshold.
Quantitative PCR was performed in 384-well PCR plates and run at 95°C for 10 min, followed by
40 cycles, each cycle consisting of 15 s at 95°C and 1 min at 60°C, using a CFX 384-Real Time
System (Bio-Rad). The BestKeeper software (62) was used to determine expression stability and
geometric mean of three different housekeeping genes (Actin, GAPDH and HPRT). ΔCt values
were calculated by subtracting the Ct value of the geometric mean of these housekeeping genes
from the Ct value for the genes of interest. The relative quantity of genes of interest was
calculated by the expression 2–ΔΔCt. Results are reported as fold induction over control.

*Lipid extraction and LC/MS analyses*

NAPE levels were measured as previously described (29). Briefly, snap-frozen cell pellets were
homogenized in a chloroform:methanol mixture (1:2 v/v, 2 mL) containing DOPE-N-
Nonadecanoyl (Larodan, #38-9091) as internal standards. Lipids were extracted twice with
chloroform (0.6 mL) and the organic phases were washed with water (0.6 mL), collected, and
dried under nitrogen stream. The organic extracts from cells were reconstituted in a mix of
methanol/chloroform (9:1 v/v, 0.1 mL). Targeted analysis of the samples was carried out on a
Waters Acquity UPLC system coupled with a Xevo TQ-MS triple quadrupole mass spectrometer.
NAPEs were separated on a reversed-phase BEH C18 column (2.1 x 50 mm, 1.7 µm), whose
temperature was kept at 55°C, and eluted at a flow rate of 0.450 mL/min using the following
gradient conditions: A = 10 mM ammonium formate in acetonitrile/water (60:40 v/v), B = 10 mM
ammonium formate in isopropyl alcohol/acetonitrile (90:10 v/v);  0.0-1.0 min 40% B, 1.0-7.0 min

40 to 100% B, 7.0-8.0 min at 100% B. The column was then reconditioned at 40% B for 1 min.
The total run time was 9 min and the injection volume was set to 5 μL. The mass spectrometer
operated in the positive ESI mode and analytes were quantified by multiple reaction monitoring
(MRM), following the transitions reported in Table 2. The capillary and the cone voltages were set
at 3 kV and 30 V for all transitions, respectively. The source temperature was set to 120 °C.
Desolvation and cone gas flows ($N_2$) were set to 800 and 50 l/h, respectively. Desolvation
temperature was set at 450°C. Data were acquired by MassLynx software and quantified by
TargetLynx software. Quantification of NAPE species was performed by means of a six-point
calibration curve where standards were prepared by spiking the only non-endogenous synthetic
NAPE reference analytical standard we could purchase, DOPE-N-19 (18:1/18:1/N19:0) in matrix
at six different concentrations, from 1 to 500 nM.

As this LC/MS-MS protocol does not allow one to differentiate sn-1 from sn-2 substituents,
individual NAPE species are designated below as NAPE (X:Y-N-acyl, **Table 2**), where X is the
total number of carbon atoms and Y the total number of double bonds in the sn-1 and sn-2
chains.

### *Fixed-cell microscopy*

Images were acquired using a Nikon Eclipse Ti2 microscope with a spinning disk confocal set-up
(Yokogawa) equipped with four lasers (wavelength in nm): 405, 488, 561 and 640 nm. Samples
were imaged using a 100X oil immersion objective (1.45 numerical aperture) with a 1x zoom. The
entire cell volume was imaged for all samples, with optical sections of 0.5 $\mu$m.

Super-resolution images were acquired using Zeiss LSM 780 Elyra SIM set-up (Carl Zeiss,
Germany) using Plan-Apochromat 63×/1.4 oil objective with a 1.518 refractive index oil (Carl
Zeiss). 16-bit images were acquired in "frame-fast" mode between wavelengths, with 32.0 $\mu$m
grid size. Optical thickness was set at 0.133 $\mu$m. Raw images were processed using the SIM
processing tool of Zen black software.

### *Quantification and Statistical Analyses*

Images were processed for analysis using FIJI software (). Colocalization analysis was performed
using the JACoP plug-in FIJI (63). 3D images were used for colocalizing two channels with
object-based detection, and thresholding of objects was performed manually.

No prior power analysis was done to measure the sample size. Graphs were plotted using
GraphPad prism 10.0, and appropriate statistical tests were performed on raw data. For all
datasets, an initial normality distribution test was performed, and non-parametric tests were
performed for any datasets that did not satisfy normal distribution. Statistical tests performed are
mentioned in the figure legends. All the experiments were performed for three independent
biological replicates.

**Acknowledgments**

We are grateful to Prof. Toro Uyama and Prof. Natsuo Ueda for generously providing the EGFP-FL-PL2G4E construct, and to Prof. Mario van der Stelt for providing the NAPE-PLD inhibitor LEI-401. We thank Prof. Takashi Nonaka for providing the α-synuclein fibrils. We are grateful to Dr. Valeria Valente and Dr. Ranabir Chakraborty for critical reading of the manuscript; we also thank Dr. Ranabir Chakraborty for assistance with super-resolution image processing. We thank Dr. Audrey Salles for technical assistance with super-resolution microscopy. We also thank all members of the Membrane Traffic and Pathogenesis Unit at the Institut Pasteur for insightful discussions. Finally, we thank Reine Bouyssie, from the administrative staff of the Membrane Traffic and Pathogenesis Unit, for her continued support.

The present study was funded by MSCA-IF-Fellowship LipiSyn – 101064077 (FP) and Fondation Alzheimer Young Researcher Program, 990987 (FP), France Parkinson - Soutien de l'Association France Parkinson AAP 2021 (CZ), Don Explore AD - Programme Explore de l'Institut Pasteur (CZ), Agence Nationale de la Recherche ANR-20-CE13-0032-01 (CZ), ANR-23-CE16-0012-03 UnProSec (CZ), Fondation pour la Recherche Médicale FRM - EQU202103012692 (CZ), FRMMND202310017892 (CZ), Fondation Alzheimer AAP 2024 (CZ).

**References**

1.  D. Coulon, L. Faure, M. Salmon, V. Wattelet, J.-J. Bessoule, Occurrence, biosynthesis and functions of N-acylphosphatidylethanolamines (NAPE): not just precursors of N-acylethanolamines (NAE). *Biochimie* **94**, 75–85 (2012).

2.  J. Liu, *et al.*, A biosynthetic pathway for anandamide. *Proc. Natl. Acad. Sci. U. S. A.* **103**, 13345–13350 (2006).

3.  S. M. K. Rahman, *et al.*, Formation of N-acyl-phosphatidylethanolamines by cytosolic phospholipase A2ε in an ex vivo murine model of brain ischemia. *Biochim. Biophys. Acta Mol. Cell Biol. Lipids* **1867**, 159222 (2022).

4.  Y. Ogura, W. H. Parsons, S. S. Kamat, B. F. Cravatt, A calcium-dependent acyltransferase that produces N-acyl phosphatidylethanolamines. *Nat. Chem. Biol.* **12**, 669–671 (2016).

5.  Y. Okamoto, J. Morishita, K. Tsuboi, T. Tonai, N. Ueda, Molecular characterization of a phospholipase D generating anandamide and its congeners. *J. Biol. Chem.* **279**, 5298–5305 (2004).

6.  P. Magotti, *et al.*, Structure of human N-acylphosphatidylethanolamine-hydrolyzing phospholipase D: regulation of fatty acid ethanolamide biosynthesis by bile acids. *Structure* **23**, 598–604 (2015).

7.  H. Cadas, E. di Tomaso, D. Piomelli, Occurrence and biosynthesis of endogenous cannabinoid precursor, N-arachidonoyl phosphatidylethanolamine, in rat brain. *J. Neurosci. Off. J. Soc. Neurosci.* **17**, 1226–1242 (1997).

535    8.    T. Zimmermann, *et al.*, Impaired anandamide/palmitoylethanolamide signaling in
536           hippocampal glutamatergic neurons alters synaptic plasticity, learning, and emotional
537           responses. *Neuropsychopharmacol. Off. Publ. Am. Coll. Neuropsychopharmacol.* **44**, 1377–
538           1388 (2019).

539    9.    J. M. Walker, S. M. Huang, N. M. Strangman, K. Tsou, M. C. Sañudo-Peña, Pain modulation
540           by release of the endogenous cannabinoid anandamide. *Proc. Natl. Acad. Sci. U. S. A.* **96**,
541           12198–12203 (1999).

542    10.    M. Igarashi, *et al.*, Dietary oleic acid contributes to the regulation of food intake through
543           the synthesis of intestinal oleoylethanolamide. *Front. Endocrinol.* **13**, 1056116 (2022).

544    11.    N. V. DiPatrizio, D. Piomelli, Intestinal lipid-derived signals that sense dietary fat. *J. Clin.*
545           *Invest.* **125**, 891–898 (2015).

546    12.    S. D. Skaper, *et al.*, N-Palmitoylethanolamine and Neuroinflammation: a Novel Therapeutic
547           Strategy of Resolution. *Mol. Neurobiol.* **52**, 1034–1042 (2015).

548    13.    J. Lo Verme, *et al.*, The nuclear receptor peroxisome proliferator-activated receptor-alpha
549           mediates the anti-inflammatory actions of palmitoylethanolamide. *Mol. Pharmacol.* **67**,
550           15–19 (2005).

551    14.    M. P. Gillum, *et al.*, N-acylphosphatidylethanolamine, a gut- derived circulating factor
552           induced by fat ingestion, inhibits food intake. *Cell* **135**, 813–824 (2008).

553    15.    C. Janfelt, *et al.*, Visualization by mass spectrometry of 2-dimensional changes in rat brain
554           lipids, including N-acylphosphatidylethanolamines, during neonatal brain ischemia. *FASEB*
555           *J. Off. Publ. Fed. Am. Soc. Exp. Biol.* **26**, 2667–2673 (2012).

556    16.    V. Natarajan, P. C. Schmid, H. H. Schmid, N-acylethanolamine phospholipid metabolism in
557           normal and ischemic rat brain. *Biochim. Biophys. Acta* **878**, 32–41 (1986).

558    17.    D. Luptakova, *et al.*, Membrane depolarization and aberrant lipid distributions in the
559           neonatal rat brain following hypoxic-ischaemic insult. *Sci. Rep.* **8**, 6952 (2018).

560    18.    H. H. Hansen, C. Ikonomidou, P. Bittigau, S. H. Hansen, H. S. Hansen, Accumulation of the
561           anandamide precursor and other N-acylethanolamine phospholipids in infant rat models
562           of in vivo necrotic and apoptotic neuronal death. *J. Neurochem.* **76**, 39–46 (2001).

563    19.    H. S. Hansen, *et al.*, Characterization of glutamate-induced formation of N-
564           acylphosphatidylethanolamine and N-acylethanolamine in cultured neocortical neurons. *J.*
565           *Neurochem.* **69**, 753–761 (1997).

566    20.    M. Saito, *et al.*, Ethanol alters lipid profiles and phosphorylation status of AMP-activated
567           protein kinase in the neonatal mouse brain. *J. Neurochem.* **103**, 1208–1218 (2007).

568    21.    A. Basit, S. Pontis, D. Piomelli, A. Armirotti, Ion mobility mass spectrometry enhances low-
569           abundance species detection in untargeted lipidomics. *Metabolomics Off. J. Metabolomic*
570           *Soc.* **12**, 50 (2016).

571    22.    H. H. Hansen, S. H. Hansen, A. Schousboe, H. S. Hansen, Determination of the phospholipid
572           precursor of anandamide and other N-acylethanolamine phospholipids before and after
573           sodium azide-induced toxicity in cultured neocortical neurons. *J. Neurochem.* **75**, 861–871
574           (2000).

575    23.    M. J. Swamy, P. K. Tarafdar, R. K. Kamlekar, Structure, phase behaviour and membrane
576           interactions of N-acylethanolamines and N-acylphosphatidylethanolamines. *Chem. Phys.*
577           *Lipids* **163**, 266–279 (2010).

578    24.    C. P. Lafrance, J. E. Blochet, M. Pézolet, N-acylphosphatidylethanolamines: effect of the N-
579           acyl chain length on its orientation. *Biophys. J.* **72**, 2559–2568 (1997).

580    25.    T. Shangguan, C. C. Pak, S. Ali, A. S. Janoff, P. Meers, Cation-dependent fusogenicity of an
581           N-acyl phosphatidylethanolamine. *Biochim. Biophys. Acta* **1368**, 171–183 (1998).

582    26.    X.-M. Li, M. Ramakrishnan, H. L. Brockman, R. E. Brown, M. J. Swamy, N-Myristoylated
583           Phosphatidylethanolamine: Interfacial Behavior and Interaction with Cholesterol.
584           *Langmuir ACS J. Surf. Colloids* **18**, 231–238 (2002).

585    27.    D.-C. Chiu, Y.-T. Cho, H. Lin, J. M. Baskin, Photoaffinity Labeling Reveals a Role for the
586           Unusual Triply Acylated Phospholipid N-Acylphosphatidylethanolamine in Lactate
587           Homeostasis. *J. Am. Chem. Soc.* **147**, 33386–33394 (2025).

588    28.    F. Palese, S. Pontis, N. Realini, D. Piomelli, NAPE-specific phospholipase D regulates LRRK2
589           association with neuronal membranes. *Adv. Pharmacol.* **90**, 217–238 (2021).

590    29.    F. Palese, S. Pontis, N. Realini, D. Piomelli, A protective role for N-
591           acylphosphatidylethanolamine phospholipase D in 6-OHDA-induced neurodegeneration.
592           *Sci. Rep.* **9**, 15927 (2019).

593    30.    Z. Hamid, *et al.*, Gender specific decrease of a set of circulating N-acylphosphatidyl
594           ethanolamines (NAPEs) in the plasma of Parkinson's disease patients. *Metabolomics Off. J.*
595           *Metabolomic Soc.* **15**, 74 (2019).

596    31.    H. M. Bailey, M. R. Cookson, How Parkinson's Disease-Linked LRRK2 Mutations Affect
597           Different CNS Cell Types. *J. Park. Dis.* **14**, 1331–1352 (2024).

598    32.    C. Klein, A. Westenberger, Genetics of Parkinson's disease. *Cold Spring Harb. Perspect.*
599           *Med.* **2**, a008888 (2012).

600    33.    S. S. Binte Mustafiz, *et al.*, The role of intracellular anionic phospholipids in the production
601           of N-acyl-phosphatidylethanolamines by cytosolic phospholipase A2. *J. Biochem. (Tokyo)*
602           **165**, 343–352 (2019).

34.　E. Kania, *et al.*, LRRK2 phosphorylation status and kinase activity regulate
(macro)autophagy in a Rab8a/Rab10-dependent manner. *Cell Death Dis.* **14**, 436 (2023).

35.　T. Eguchi, *et al.*, LRRK2 and its substrate Rab GTPases are sequentially targeted onto
stressed lysosomes and maintain their homeostasis. *Proc. Natl. Acad. Sci. U. S. A.* **115**,
E9115–E9124 (2018).

36.　N. Yadavalli, S. M. Ferguson, LRRK2 suppresses lysosome degradative activity in
macrophages and microglia through MiT-TFE transcription factor inhibition. *Proc. Natl.*
*Acad. Sci. U. S. A.* **120**, e2303789120 (2023).

37.　M. Madureira, N. Connor-Robson, R. Wade-Martins, LRRK2: Autophagy and Lysosomal
Activity. *Front. Neurosci.* **14**, 498 (2020).

38.　J. Obergasteiger, *et al.*, Kinase inhibition of G2019S-LRRK2 enhances autolysosome
formation and function to reduce endogenous alpha-synuclein intracellular inclusions. *Cell*
*Death Discov.* **6**, 45 (2020).

39.　S. S. Binte Mustafiz, *et al.*, Intracellular Ca2+-dependent formation of N-acyl-
phosphatidylethanolamines by human cytosolic phospholipase A2ε. *Biochim. Biophys. Acta*
*Mol. Cell Biol. Lipids* **1864**, 158515 (2019).

40.　M. Palmieri, *et al.*, Characterization of the CLEAR network reveals an integrated control of
cellular clearance pathways. *Hum. Mol. Genet.* **20**, 3852–3866 (2011).

41.　S. MacIsaac, *et al.*, Neuron-autonomous susceptibility to induced synuclein aggregation is
exacerbated by endogenous Lrrk2 mutations and ameliorated by Lrrk2 genetic knock-out.
*Brain Commun.* **2**, fcz052 (2020).

42.　E.-J. Bae, *et al.*, LRRK2 kinase regulates α-synuclein propagation via RAB35
phosphorylation. *Nat. Commun.* **9**, 3465 (2018).

43.　Y. Okamoto, *et al.*, Mammalian cells stably overexpressing N-
acylphosphatidylethanolamine-hydrolysing phospholipase D exhibit significantly decreased
levels of N-acylphosphatidylethanolamines. *Biochem. J.* **389**, 241–247 (2005).

44.　K. Ahn, *et al.*, Discovery and characterization of a highly selective FAAH inhibitor that
reduces inflammatory pain. *Chem. Biol.* **16**, 411–420 (2009).

45.　R. Wallings, N. Connor-Robson, R. Wade-Martins, LRRK2 interacts with the vacuolar-type
H+-ATPase pump a1 subunit to regulate lysosomal function. *Hum. Mol. Genet.* **28**, 2696–
2710 (2019).

46.　A. B. West, *et al.*, Parkinson's disease-associated mutations in leucine-rich repeat kinase 2
augment kinase activity. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 16842–16847 (2005).

47. J. Schapansky, *et al.*, Familial knockin mutation of LRRK2 causes lysosomal dysfunction and
accumulation of endogenous insoluble α-synuclein in neurons. *Neurobiol. Dis.* **111**, 26–35
(2018).

48. J. H. Kluss, *et al.*, Lysosomal positioning regulates Rab10 phosphorylation at LRRK2+
lysosomes. *Proc. Natl. Acad. Sci. U. S. A.* **119**, e2205492119 (2022).

49. L. Bonet-Ponce, *et al.*, LRRK2 mediates tubulation and vesicle sorting from lysosomes. *Sci.*
*Adv.* **6**, eabb2454 (2020).

50. X. Wang, *et al.*, Rab12 is a regulator of LRRK2 and its activation by damaged lysosomes.
*eLife* **12**, e87255 (2023).

51. B. Castellani, *et al.*, Synthesis and characterization of the first inhibitor of N-
acylphosphatidylethanolamine phospholipase D (NAPE-PLD). *Chem. Commun.* **53**, 12814–
12817 (2017).

52. P. Rivera, *et al.*, Localization of peroxisome proliferator-activated receptor alpha (PPARα)
and N-acyl phosphatidylethanolamine phospholipase D (NAPE-PLD) in cells expressing the
Ca(2+)-binding proteins calbindin, calretinin, and parvalbumin in the adult rat
hippocampus. *Front. Neuroanat.* **8**, 12 (2014).

53. S. Wójtowicz, A. K. Strosznajder, M. Jeżyna, J. B. Strosznajder, The Novel Role of PPAR
Alpha in the Brain: Promising Target in Therapy of Alzheimer's Disease and Other
Neurodegenerative Disorders. *Neurochem. Res.* **45**, 972–988 (2020).

54. Y. S. Kim, *et al.*, PPAR-α Activation Mediates Innate Host Defense through Induction of
TFEB and Lipid Catabolism. *J. Immunol.* **198**, 3283–3295 (2017).

55. P. Holland, *et al.*, HS1BP3 negatively regulates autophagy by modulation of phosphatidic
acid levels. *Nat. Commun.* **7**, 13889 (2016).

56. A. Sanyal, H. S. Novis, E. Gasser, S. Lin, M. J. LaVoie, LRRK2 Kinase Inhibition Rescues
Deficits in Lysosome Function Due to Heterozygous GBA1 Expression in Human iPSC-
Derived Neurons. *Front. Neurosci.* **14**, 442 (2020).

57. E. D. Mock, *et al.*, Discovery of a NAPE-PLD inhibitor that modulates emotional behavior in
mice. *Nat. Chem. Biol.* **16**, 667–675 (2020).

58. E. Leishman, K. Mackie, S. Luquet, H. B. Bradshaw, Lipidomics profile of a NAPE-PLD KO
mouse provides evidence of a broader role of this enzyme in lipid metabolism in the brain.
*Biochim. Biophys. Acta* **1861**, 491–500 (2016).

59. T. Kuwahara, T. Iwatsubo, The Emerging Functions of LRRK2 and Rab GTPases in the
Endolysosomal System. *Front. Neurosci.* **14**, 227 (2020).

60. P. Reinhardt, *et al.*, Derivation and expansion using only small molecules of human neural
progenitors for neurodegenerative disease modeling. *PloS One* **8**, e59252 (2013).

61.    T. Nonaka, S. T. Watanabe, T. Iwatsubo, M. Hasegawa, Seeded aggregation and toxicity of
672            {alpha}-synuclein and tau: cellular models of neurodegenerative diseases. *J. Biol. Chem.*
673            **285**, 34885–34898 (2010).

674    62.    M. W. Pfaffl, A. Tichopad, C. Prgomet, T. P. Neuvians, Determination of stable
675            housekeeping genes, differentially regulated target genes and sample integrity:
676            BestKeeper--Excel-based tool using pair-wise correlations. *Biotechnol. Lett.* **26**, 509–515
677            (2004).

678    63.    S. Bolte, F. P. Cordelières, A guided tour into subcellular colocalization analysis in light
679            microscopy. *J. Microsc.* **224**, 213–232 (2006).

682 **Figures and Tables**

**Figure 1. Generation of catecholaminergic SH-SY5Y neuronal cells accumulating NAPEs.**
(**A**) Schematic representation of the experimental strategy. Stable SH-SY5Y cell lines were generated either by knocking down NAPE-PLD using CRISP-Cas9 approach or by overexpressing PLA2G4E, both manipulations led to NAPE accumulation at the cell membranes. (**B**) *NAPE-PLD* mRNA levels in NAPE-PLD-knockdown (KD; violet bars) and wild-type (WT; blue bars) cells. (**C**) Western blot analysis of NAPE-PLD protein levels in KD and WT cells. Top, representative blot; bottom, densitometric quantification normalized to glyceraldehyde 3-phosphate dehydrogenase (GAPDH). (**D**) Immunofluorescence analysis of NAPE-PLD (gray) with representative images (**D1, D2**) and quantification. Nuclei were counterstained with DAPI. (**E**) Total NAPE levels in KD and WT cells. (**F**) *PLA2G4E* mRNA levels in PLA2G4E-overexpressing (OE; pink bars) and WT (blue bars) cells. (**G**) Western blot analysis of PLA2G4E protein levels in OE and WT cells. Top, representative blot; bottom, densitometric quantification normalized to GAPDH. (**H**) Immunofluorescence analysis of PLA2G4E (gray) with representative images (**H1, H2**) and quantification. Nuclei were counterstained with DAPI. (**I**) Total NAPE levels in OE and WT cells. Error bars represent SEM. *$P < 0.05$, **$P < 0.01$ ***$P < 0.001$, ****$P < 0.0001$ Student's *t*-test with Welch's correction. Scale bars, 10 μm.

**A1.** SH-SY5Y WT
LRRK2 DAPI

**A2.** NAPE-PLD KD

**A3.** PLA2G4E OE

WT  KD  OE
Total Rab8a
pT72Rab8a
Actin

**A**
LRRK2 Fluorescence Intensity (% WT)
300
200
100
0
100%
67%  71%
****
****
SH-SY5Y WT    NAPE-PLD KD    PLA2G4E OE

**B**
Manders Overlap Coefficient (Lamp1 with LRRK2)
0.20
0.15
0.10
0.05
0.00
****
****
SH-SY5Y WT    NAPE-PLD KD    PLA2G4E OE

**C**
pT72Rab8a/Tiot Rab8a (%WT)
150
100
50
0
*
*
SH-SY5Y WT    NAPE-PLD KD    PLA2G4E OE

**D1.** SH-SY5Y WT
LAMP1 LRRK2 DAPI

**D2.** NAPE-PLD KD

**D3.** PLA2G4E OE

LAMP1 LRRK2

**E**
Magic Red Fluorescence Intensity (%WT)
131%
120%
100%
****
*
SH-SY5Y WT    NAPE-PLD KD    PLA2G4E OE

**E1.** SH-SY5Y WT

**E2.** NAPE-PLD KD

**E3.** PLA2G4E OE

**Figure 2. Neuronal cells accumulating NAPEs downregulate LRRK2 protein and potentiate lysosomal activity.** (**A**) Immunofluorescence analysis of LRRK2 (gray) with representative images (**A1–A3**) and quantification in wild-type (WT; **A1**, blue), NAPE-PLD knockdown (KD; **A2**, violet), and PLA2G4E-overexpressing (OE; **A3**, pink) SH-SY5Y cells. Nuclei were counterstained with DAPI. (**B**) Quantification of the colocalization between LAMP1-positive structures and LRRK2, expressed as Manders' overlap coefficient for the fraction of LAMP1 signal overlapping with LRRK2. (**C**) Western blot analysis of Rab8a and phosphorylated Rab8a (pT72-Rab8a) in KD and OE compared with WT cells. Top, representative blot; bottom, densitometric quantification normalized to actin. (**D**) Representative super-resolution immunofluorescence images of LAMP1 (magenta) and LRRK2 (green) in WT (**D1**), NAPE-PLD KD (**D2**), and PLA2G4E OE (**D3**) cells. Nuclei were stained with DAPI (gray). Scale bar, 5 μm. (**E**) Fluorescence intensity quantification and representative images of the Magic Red–cathepsin B assay, indicating lysosomal degradative capacity per cell, in WT (**E1**), NAPE-PLD KD (**E2**), and PLA2G4E OE (**E3**) cells. Representative images are displayed using the FIRE lookup table (LUT) to indicate fluorescence intensity. Scale bars, 10 μm. Mean values are indicated in the graphs; error bars represent SEM. *$P < 0.05$, ****$P < 0.0001$; one-way ANOVA with Tukey's multiple-comparison test.

719
720
**Figure 3. Neuronal cells accumulating NAPEs prevent LRRK2 accumulation following α-synuclein treatment and enhance aggregate clearance.** (**A**) Immunofluorescence analysis of LRRK2 (gray) with representative images (**A1-A4**) and quantification in WT (**A1, A2**), NAPE-PLD-knockdown (KD; **A3**), and PLA2G4E-overexpressing (OE; **A4**) cells treated or not with α-syn (**A2-A4**). (**B**) Quantification of the colocalization between LAMP1-positive structures and LRRK2, expressed as Mander's overlap coefficient for the fraction of LAMP1 signal overlapping with LRRK2. (**C**) Quantification and representative images of α-syn aggregate number per cell in WT (**C1**), NAPE-PLD KD (**C2**) and PLA2G4E OE (**C3**) cells. LAMP1 is shown in yellow, α-syn in pink, and nuclei were counterstained with DAPI (blue). Scale bar, 10 μm. Mean values are indicated in the graphs; error bars represent SEM. ***P < 0.001, ****P < 0.0001; one-way ANOVA with Tuckey's multiple-comparison test.

**Figure 4. NAPE-PLD overexpression upregulates LRRK2 protein and impairs lysosomal activity.** (**A**) Schematic representation of the generation of stable SH-SY5Y cell lines overexpressing NAPE-PLD, leading to reduced NAPE levels at cellular membranes. (**B**) *NAPE-PLD* mRNA levels in NAPE-PLD-overexpressing (OE; green) and wild-type (WT; blue) cells. (**C**) Western blot analysis of NAPE-PLD protein levels in OE and WT cells. Top, representative blot; bottom, densitometric quantification normalized to GAPDH. (**D**) Immunofluorescence analysis of NAPE-PLD (gray) with representative images (**D1, D2**) and quantification. (**E**) Immunofluorescence analysis of LRRK2 (gray) with representative images (**E1, E2**) and quantification in WT (**E1**) and NAPE-PLD OE (**E2**) cells. Nuclei were counterstained with DAPI (blue). (**F**) Western blot analysis of Rab8a and phosphorylated Rab8a (pT72-Rab8a) in NAPE-PLD OE and WT cells. Top: representative blot; bottom, densitometric quantification normalized to actin. (**G**) Schematic representation of the experimental strategy. Ionomycin treatment induces intracellular $Ca^{2+}$ release, promoting NAPE production and degradation in NAPE-PLD OE cells. NAPE-PLD activity was inhibited by overnight treatment with the NAPE-PLD inhibitor LEI-401. Ionomycin was administered together with Magic Red reagent for 30 min before image acquisition. (**H**) Fluorescence intensity quantification and representative images of the Magic Red – cathepsin B assay, indicating lysosomal degradative capacity per cell, in untreated WT (**H1,** white bar and blue dots) and NAPE-PLD OE cells (**H4,** white bar and green dots), ionomycin-treated WT (**H2,** blue bar and blue dots)  and NAPE-PLD OE cells (**H5,** green bar and green dots) or WT (**H3,** blue bar white dots) and NAPE-PLD OE cells (**H6**, green bar and white dots) treated with ionomycin and LEI-401. Representative images are displayed using the FIRE lookup table (LUT) to indicate fluorescence intensity. Scale bars, 10 μm. Mean values are indicated in the graphs; error bars represent SEM. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001; Student's *t*-test with Welch's correction (**B-F**) and two-way ANOVA with Tuckey's multiple-comparisons test (**H**).

**Figure 5. NAPE-PLD overexpression enhances LRRK2 accumulation following α-synuclein treatment and impairs aggregate clearance.** (**A**) Immunofluorescence analysis of LRRK2 (gray) with representative images (**A1-A3**) and quantification in WT (**A1, A2**), and NAPE-PLD OE (**A3**) SH-SY5Y cells treated or not with α-synuclein (α-syn; **A2, A3**). (**B**) Quantification and representative images of α-syn aggregate number per cell in WT (**B1**) and NAPE-PLD OE (**B2**) cells. α-Syn is shown in pink, and nuclei were counterstained with DAPI. Scale bar, 10 µm. Mean values are indicated in the graphs; error bars represent SEM. *P<0.05, ***P < 0.001, ****P < 0.0001; one-way ANOVA with Tuckey's multiple-comparison test (**A**) and Student's *t*-test with Welch's correction (**B**).

Figure labels:

**A1.** Healthy DAN - Vehicle  **A2.** Healthy DAN - LEI-401  **A3.** G2019S DAN - Vehicle  **A4.** G2019S DAN - LEI-401

DAPI Magic Red

**B** — MR + Lysosomes N/Cell (%Healthy DAN + Veh)

Healthy DAN + Veh, Healthy DAN + LEI-401, G2019S DAN + Veh, G2019S DAN + LEI-401

**C** — α-Synuclein Aggregates N/Cell

Healthy DAN + Veh, Healthy DAN + LEI-401, G2019S DAN + Veh, G2019S DAN + LEI-401

**D1.** Control DAN - Vehicle  **D2.** Control DAN – LEI-401  **D3.** G2019S DAN – Vehicle  **D4.** G2019S DAN – LEI-401

DAPI
Lamp1
α-Syn

α-Syn

**Figure 6. NAPE-PLD inhibition restores lysosomal activity and α-synuclein fibril clearance in G2019S-LRRK2 dopaminergic neurons.** (**A**) Representative images of the Magic Red – cathepsin B assay, indicating lysosomal degradative capacity per cell, in healthy dopaminergic neurons (DANs; **A1, A2**) and G2019S-LRRK2 mutant DANs (**A3, A4**) treated (**A2, A4**) or not (**A1, A3**) with the NAPE-PLD inhibitor LEI-401. (**B**) Fluorescence intensity quantification of the Magic Red assay in healthy DANs (green) and G2019S-LRRK2 mutant DANs (gray). (**C, D**) Quantification and representative images of α-synuclein (α-syn) aggregate number per cell in healthy DANs (**D1, D2**) and G2019-LRRK2 mutant DANs (**D3, D4**) treated (**D2, D4**) or not (**D1, D3**) with LEI-401. α-Syn is shown in pink, LAMP1 yellow, and nuclei were counterstained with DAPI. Scale bars, 10 μm. *P < 0.05, ****P < 0.0001; two-way ANOVA with Tuckey's multiple-comparison test.

784 **Table 1:** *Primers used in the study.*

| Gene | Forward primer | Reverse primer |
|------|----------------|----------------|
| *NAPE-PLD* | GGACATCAGTGTGAATCCTTACAG | GGACATCAGTGTGAATCCTTACAG |
| *PLA2G4E* | GATGCCTGGAACCTGTCACACA | TCACTACCGTGGTCTCCAGGAT |
| *GAPDH* | GTCTCCTCTGACTTCAACAGCG | ACCACCCTGTTGCTGTAGCCAA |
| *ACTINB* | CACCATTGGCAATGAGCGGTTC | AGGTCTTTGCGGATGTCCACGT |
| *HPRT* | CATTATGCTGAGGATTTGGAAAGG | CTTGAGCACACAGAGGGCTACA |

785
786

787 ***Table 2:*** *Parent, quantifier and qualifier ions (as m/z) tracked for each of the 14 NAPEs under*
788 *investigation. The corresponding observed retention time (in minutes) is also indicated.*

| ID | NAME | Parent m/z | Fragment m/z (Qn) | Fragment m/z (Qf) | Rt (min) |
|----|------|-----------|------------------|------------------|----------|
| 1 | PE (P50:1-N-acyl) | 940.8 | 282.3 | 561.5 | 5.03 |
| 2 | PE (P52:2-N-acyl) | 966.8 | 282.3 | 587.5 | 5.03 |
| 4 | PE (P54:5-N-acyl) | 988.8 | 282.3 | 609.5 | 4.79 |
| 5 | PE (P54:4-N-acyl) | 990.8 | 282.3 | 611.5 | 5.00 |
| 7 | PE (P50:6-N-acyl) | 1014.8 | 308.3 | 609.5 | 4.81 |
| 10 | PE (P50:5-N-acyl) | 1016.8 | 308.3 | 611.5 | 5.01 |
| 11 | PE (P50:5-N-acyl) | 1016.8 | 310.3 | 609.5 | 5.00 |
| 20 | PE (P50:6-N-acyl) | 1014.8 | 282.3 | 635.5 | 4.92 |
| 21 | PE (P50:6-N-acyl) | 1014.8 | 310.3 | 607.5 | 4.92 |
| 22 | PE (50:6-N-acyl) | 1030.8 | 282.3 | 651.5 | 4.84 |
| 23 | PE (50:6-N-acyl) | 1030.8 | 310.3 | 623.5 | 4.82 |
| 24 | PE (50:4-N-acyl) | 1034.8 | 310.3 | 627.5 | 5.10 |
| 25 | PE (P58:6-N-acyl) | 1042.8 | 310.3 | 635.5 | 4.98 |
| 26 | PE (58:6-N-acyl) | 1058.8 | 310.3 | 651.5 | 5.03 |

789
790