1

OVERLAP IN NEURAL REPRESENTATIONS OF COORDINATED WRIST AND FINGER MOVEMENTS IN HUMAN

MOTOR CORTEX

A.M.X. Emonds[1,2], E.V. Okorokova[1,3], G.H. Blumenthal[4,5], J.L. Collinger[4,5,6], S.J. Bensmaia[1,7], L.E.
Miller[8,9,10,11], J.E. Downey[1,2], and A.R. Sobinov[1*]

[1] Department of Organismal Biology and Anatomy, University of Chicago, Chicago, IL, USA
[2] Department of Neurological Surgery, University of Chicago, Chicago, IL, USA
[3] Department of Neurological Surgery, University of California, Davis, CA, USA
[4] Rehab Neural Engineering Labs, University of Pittsburgh, Pittsburgh, PA, USA
[5] Department of Physical Medicine and Rehabilitation, University of Pittsburgh, Pittsburgh, PA, USA
[6] Department of Bioengineering, University of Pittsburgh, Pittsburgh, PA, USA
[7] Neuroscience Institute, University of Chicago, Chicago, IL, USA
[8] Department of Neuroscience, Feinberg School of Medicine, Northwestern University, Chicago, IL,
USA
[9] Department of Physical Medicine and Rehabilitation, Feinberg School of Medicine, Northwestern
University, Chicago, IL, USA
[10] Department of Biomedical Engineering, McCormick School of Engineering, Northwestern
University, Evanston, IL, USA
[11] Shirley Ryan AbilityLab, Chicago, IL, USA
28
29 [*] Correspondence: an.sobinov@gmail.com
30

ABSTRACT

Dexterous hand function underlies many essential human activities, from tool use to expression
through gestures. Coordinated digit movements are enabled by the intricate musculature of the
hand and forearm, which also imposes mechanical coupling between the digits and wrist,
constraining their independent control. It remains unclear whether motor cortex inherits these
constraints in its activity or encodes digit and wrist independently. To address this problem, we
asked individuals with intracortical microelectrode arrays implanted in motor cortex to attempt
flexion and extension of individual digits, either in isolation or in combination with attempted wrist
movements. We could accurately decode which digit was moving based on cortical recordings, and
channels selective for digit identity were arranged somatotopically across the recording arrays.
Nevertheless, the activity during flexion or extension overlapped between digits, and movement
direction of a given digit could be reliably inferred by a decoder trained on movements of other digits.
This directional signal was largely invariant to the digit's initial posture. The population axis
describing digit movement direction was aligned with the axes associated with wrist flexion-
extension or pronation-supination. This alignment persisted during simultaneous wrist and digit
movements, which complicated efforts to control them individually. However, by decoding wrist and
digit motion from activity orthogonal to the shared direction axis, a participant was able to achieve
continuous control of virtual hand movements with improved speed and reduced unintended
movements. Together, the results identify both a code for digit identity and a low-dimensional
flexion-extension signal which is shared across the digits and wrist. This arrangement is consistent
with muscle-like biomechanical constraints on motor cortical activity, which must be accounted for
to improve coordinated BCI control.

INTRODUCTION

The hand endows us with an ability to interact with the world around us, from communicating through gestures to grasping a cup of coffee or playing music. This versatility is enabled by the intricate musculature of the hand and forearm and the disproportionately large regions of sensorimotor cortex devoted to hand control (Penfield and Rasmussen, 1950; Rizzolatti and Luppino, 2001; Baldwin *et al.*, 2018; Sobinov and Bensmaia, 2021). Yet, the musculoskeletal architecture that makes dexterity possible also imposes strong biomechanical constraints: the digits share major muscle groups whose bodies are located in the forearm, and tendon connections can transmit force across fingers (Kilbreath and Gandevia, 1994; Reilly and Schieber, 2003; Schieber and Santello, 2004; Duinen and Gandevia, 2011). Moreover, many muscles that actuate the digits cross the wrist, and changes in wrist posture alter the muscle lengths and patterns required to generate digit forces and motion (Adewuyi, Hargrove and Kuiken, 2017; Beringer *et al.*, 2020). Predictably, digit-actuating muscles are also active during wrist movements. These mechanics limit the independent control of individual degrees of freedom (Kilbreath and Gandevia, 1994; Zatsiorsky, Li and Latash, 2000; Reilly and Schieber, 2003; Schieber and Santello, 2004; Duinen and Gandevia, 2011). Despite this evidence, an implicit assumption in brain-computer interfaces treats each joint independently, rather than taking interactions into account (Aggarwal *et al.*, 2013; Vaskov *et al.*, 2018; Shah *et al.*, 2023; Jude *et al.*, 2026). It remains unclear to what extent motor cortex reflects these mechanics or encodes more independent control signals in BCIs.

A large body of work has shown that motor cortical activity contains information about fine hand kinematics (Schieber, 2001; Capaday *et al.*, 2013; Goodman *et al.*, 2019; Okorokova *et al.*, 2020). Finger and wrist movements activate often overlapping regions within motor cortex (Aggarwal *et al.*, 2008; Branco *et al.*, 2019; Chehade and Gharbawie, 2023), such that individual neurons within motor cortex are typically involved in the movement of multiple digits (Schieber and Hibbard, 1993; Georgopoulos *et al.*, 1999; Acharya *et al.*, 2007; Aggarwal *et al.*, 2008; Kirsch, Rivlis and Schieber, 2014; Branco *et al.*, 2019). This overlap is commonly interpreted as evidence for distributed, mixed selectivity rather than segregation of digit-specific neuronal groups. Despite the apparent somatotopic overlap, decoding analyses suggest partial separability between wrist- and digit-related activity (Mohamed and Aharonson, 2021), indicating that wrist and digit signals should also be separable during simultaneous wrist and digit movements (Nason *et al.*, 2021; Mender *et al.*, 2024). Importantly, apparent interactions between representations could arise not only from fixed mechanical coupling but reflect functional coupling—statistics of hand use arising from the correlated joint kinematics during everyday behavior (Ejaz, Hamada and Diedrichsen, 2015). Whether neural activity is fundamentally organized around independent digit- and wrist-specific commands, or instead around other activity profiles constrained by biomechanics or use, remains unclear—particularly during movements in which wrist and digits are engaged together, as they typically are during natural grasping and manipulation.

This question is not only fundamental for understanding motor control but also has practical consequences for neurotechnology. People with tetraplegia view restoration of arm and hand movement as fundamental to an improved quality of life (Anderson, 2004; Collinger, Boninger, *et al.*, 2013). Intracortical brain-computer interfaces (iBCIs) have enabled functional arm control and simple grasping behaviors (Collinger, Wodlinger, *et al.*, 2013; Ajiboye *et al.*, 2017; Flesher *et al.*, 2021; Rastogi *et al.*, 2021), but more complete restoration of hand function will require coordinated control of individual digits together with the wrist. Although high-density electrocorticography (ECoG) and intracortical recordings have enabled accurate decoding of finger movements in both offline and

online settings (Kubánek *et al.*, 2009; Hotson, McMullen, Matthew S Fifer, *et al.*, 2016; Irwin *et al.*,
2017; Jorge *et al.*, 2020; Nason *et al.*, 2021; Guan *et al.*, 2023; Nason-Tomaszewski *et al.*, 2023;
Vargas-Irwin *et al.*, 2024; Willsey *et al.*, 2025), achieving high-dimensional and fully independent
control remains out of our grasp; systems supporting many degrees of freedom are under active
development (Wodlinger *et al.*, 2015; Hotson, McMullen, Matthew S Fifer, *et al.*, 2016; Nason *et al.*,
2021; Shah *et al.*, 2023; Dekleva and Collinger, 2025). Moreover, adding coordinated wrist control
may induce gain adaptation, reducing effective bandwidth or otherwise altering individual
movement representations (Rasmussen, Schwartz and Chase, 2017). Together, these limitations
raise the possibility that understanding the constraints intrinsic to motor cortical population
activity—and not just decoder design—may be crucial for extracting hand control signals for BCI.

Here, we examined whether human motor cortical activity reflects coupling between digits and wrist
or natively supports independent control signals. Using intracortical recordings from participants
with tetraplegia, we analyzed population activity during attempted digit flexion or extension,
performed either in isolation or concurrently with wrist movements. We asked how digit identity and
movement direction are encoded and spatially organized across the arrays, quantified overlap
between digit- and wrist-related signals, and exploited this structure to build decoders that enabled
continuous control of a virtual hand.

RESULTS

We recorded motor cortical activity using intracortical microelectrode arrays (Neuroport Array,
Blackrock Neurotech, UT, USA) implanted in the motor cortex (Figure 1A) of three individuals with
tetraplegia due to cervical spinal cord injury who attempted to make finger and wrist movements
following audiovisual instructions (Figure 1B). We examined how this cortical activity represents
both digit identity (ID) and the direction of digit motion, as well as the interaction of the
representation of digit motion with that of wrist motion.

Movement of individual digits

We investigated the neural activity underlying individual finger movements by asking participants to
mimic the movements of a virtual hand, which they observed on a monitor (Figure 1B). Trials started
with the participants imagining their hand and wrist in a neutral posture. An auditory instruction
specifying a random digit and movement direction was followed two seconds later by an auditory
"go" cue and the movement of the virtual hand, which the participant was instructed to mimic (Figure
1C). A subsequent "relax" cue instructed them to return to the neutral posture. We built linear
decoders on neural activity from each channel, averaged within 100ms windows, only selecting
channels with a mean firing rate above 0.5 Hz. Neural activity allowed accurate classification of digit
identity starting from 125ms, 125ms, and 265ms after audio cue end (1.08s, 1.08s, and 940ms
before the go cue) in C1, C2, and P5, respectively (Figure 1D), with accuracy peaking after the go cue.
In participant C1, discriminability of the digits generally decreased from thumb to pinky (Figure 1E),
but in other participants the accuracy was similar for all digits, although P5 only imagined thumb,
index and middle digit movements. Overall, digit identity was strongly represented in the neural
activity of all participants.

Figure 1. Decoding digit movement. A| Implant locations for all participants. **B|** Experimental setup. **C|** Trial progression.
D| Time course of digit classification accuracy for each participant at each time point for decoders trained on both flexion
and extension trials. Each thin trace corresponds to a different session, and thick—to average. Gray bar above the traces
corresponds to the points where prediction significantly exceeded chance performance of shuffled distribution (pale pink
line). Red line illustrates chance level. Gray shaded region represents the range in which audio cues ended. **E|** Digit identity
confusion matrices constructed at the peak accuracy and averaged across sessions. **F|** Modulation of electrodes by digit

identity (significance by Kruskal-Wallis test, see **Supp. Figure 1**C and details in Methods) on C1 lateral motor array during
single digit movement, averaged across sessions. Pink corresponds to facilitation of activity and blue, to inhibition. Non-
significant channels are gray. Anterior-most medial corner of the array is black. (C1 n=5, C2 n=6, P5 n=3) **G|** Similarity of
spatial modulation maps for each subject. Since these correlations were calculated using composite maps built across
all sessions, the diagonal represents a comparison between identical maps.

We hypothesized that differences in discriminability might reflect the spatial organization of
modulated channels on the arrays. If so, digits that were more frequently confused would be
expected to activate highly overlapping regions of motor cortex. We made modulation maps by
computing the firing rate of a channel relative to its average firing rate during the movement phase of
the trial (Figure 1F, Supp. Figure 1A). In all participants, we focused the spatial analysis on the more
active array—lateral in C1 and C2, and medial in P5—because the other array was mostly silent
(mean firing rate < 0.5 Hz). We selected only the channels with modulation that differed consistently
across digits (Kruskal-Wallis test, p<0.05). The positively modulated units were somatotopically
organized in all participants, most saliently in C1 (Figure 1F), such that movements of neighboring
digits recruited activity in nearby channels more than distant ones. For each participant, we
quantified the similarity between the modulation maps for individual digits by computing the spatial
correlation between them (Figure 1G). We identified significant correlations by comparing them with
correlations obtained from shuffled data pooled across all conditions (Supp. Figure 1B). The patterns
of spatial correlation in all participants qualitatively matched the decoding confusion matrices
(Figure 1E and Figure 1G, respectively). The confusion in decoding middle, ring, and pinky in C1
(Figure 1E, left) was reflected in high spatial correlation between the maps for those digits (Figure
1G, left). At the same time, high discriminability of the thumb was reflected in low or negative spatial
correlations. In C2, confusion between the thumb and index was reflected in the spatial correlations
of their associated maps. We did not see these effects in P5, most likely because fewer digits were
tested across fewer sessions, lowering the total number of significantly modulated electrodes.
Overall, spatial correlations between neighboring digits were slightly higher than between digits that
were far apart, suggesting not only spatial organization of neural activity corresponding to each digit,
but also a certain degree of somatotopy across digits.

Having characterized the distinct patterns of activity corresponding to digit identity, we next
examined how digit movement direction is encoded in the motor cortex. We isolated trials
corresponding to the movement of a single digit and built decoders to indicate whether that digit was
flexed or extended (Figure 2A). In participant C2, movement direction was represented much more
weakly than in the other two participants, and signals were consistently weak for all digits (**Error!**
Reference source not found. Supp. Figure 2A). In the other two participants, C1 and P5, movement
direction information appeared at a similar time as digit identity (1.12s and 820ms before movement,
respectively). The encoding of movement direction did not strongly depend on the starting posture
of the movement. Decoders trained on the "move" phase—where flexion and extension started from
neutral posture—largely generalized to the "relax" phase, where movement started at one of the
extremes of the range of motion, and vice versa (same starting posture mean accuracy 91%, different
starting posture mean 84%, p=0.0026, Figure 2B; see Supp. Figure 3 for temporal generalization
heatmaps).

Given the biomechanical coupling
between digits, we assessed whether
movement direction of one digit could be
decoded from the neural activity
corresponding to movement of other
digits. We built decoders of movement
direction on trials with all but one digit and
compared them to decoders built on only
that digit (Figure 2C; for elaboration, see
Supp. Figure 4). The decoders tested on
the untrained digits performed better than
chance, and not significantly worse than
the same-digit decoders (p=0.058).
Flexion or extension of each digit evoked
activity in channels widely spread over the
arrays of the participants (Supp. Figure
2BC). Overall, these results indicate
strong digit movement direction encoding
in motor cortical activity that generalizes
between postures of the digit, as well as
across digits.

Neural overlap between wrist and digits

Digit movement rarely happens in isolation
from wrist movements, which are typically
needed to optimize hand orientation, as
when the hand approaches an object (Yan,
Sobinov and Bensmaia, 2022). Also, the
powerful extrinsic finger muscles that
cross the wrist create substantial wrist

Figure 2. Movement direction generalizes between digits. A| Average classification of digit movement direction—flexion or extension—for individual digits. Lines and shading follow the conventions defined in Figure 1D. (C1 n=5, C2 n=2, P5 n=3) **B|** Generalization of digit movement direction encoding for different starting postures. Decoders were trained on the 100ms window of activity during movement that maximized classification accuracy. Different colors correspond to different subjects (C1: purple, C2: pink, P5: green). Thin lines connect performance during the same session. **C|** Generalization of movement direction across digits during movement phase in subject C1. The extension-flexion classifier was trained and tested on single digits ("same digit") and trained on all but the tested digit ("held out"). Each dot represents average accuracy in a single session. Colors as in B. Thin lines connect sessions. Horizontal solid red line corresponds to chance level.

torque that must be counteracted by wrist extensor muscles. Furthermore, altered wrist posture
changes the force production capacity of the extrinsic hand muscles. The extent to which these
complex mechanics are represented within neural activity is an important basic neuroscientific
question with significant implications for iBCI control.

We began by characterizing the neural activity during isolated wrist movements. Participants
attempted to flex, extend, pronate, or supinate their hand starting from the neutral posture (Figure
3A). Linear classifiers based on neural activity could distinguish wrist orientation throughout these
trials, beginning 73ms, 413ms, and 373ms after mean audio cue end (1.20s, 880ms, and 920ms
before the movement cue) for C1, C2, and P5, respectively (Figure 3B). Classification accuracy was

generally lower in C2 than in C1 and P5, but no consistent confusion pattern appeared across
participants (Figure 3C).

Figure 3. Wrist movements. A| Schematic of a representative trial. **B|** Time course of wrist orientation classification
accuracy for subject C1 (n=13), C2 (n=4), and P5 (n=3) based on the neural activity averaged in 100ms window at each time
point. Lines and shading follow the conventions defined in Figure 1D. **C|** Wrist posture confusion matrix at peak accuracy
from B, averaged across sessions.

We then looked at the interaction between neural activity for isolated wrist and digit movements by
comparing movement axes in neural space. This analysis was performed with data from C1 and P5,
who had strong representation of digit movement direction (Figure 2, Supp. Figure 3C) and wrist
movement direction (Supp. Figure 5) throughout the trials. We defined a digit movement direction
axis in neural activity space as the vector running from neural activity during extension through the
neutral posture to activity during flexion. We identified digit movement direction axes both for each
digit individually and for all digits combined, and computed angles between the axes for each digit
and the common axis (Figure 4A). In C1, the vectors for individual digits aligned closely with the
common digit movement axis, which was expected given that digit movement direction generalized
between digits (Figure 2C, purple dots). Predictably, this was less pronounced in P5, whose digit
movement direction generalization, while significant, was weaker than that of the other participants

(Figure 2C, green dots). This was possibly due to a learned compensatory strategy P5 had developed
that relied upon residual movement ability in the index finger and thumb.

Similarly, we defined neural axes for
wrist extension to flexion (EF) and
supination to pronation (SP), and
calculated angles between those axes
and the common digit axis from data
collected on the same day. Wrist EF and
SP neural axes were orthogonal in
participant C1 (Figure 4B, brown and
orange distributions), which was
expected since they correspond to
orthogonal directions of movement and
are largely actuated by different
muscles. Wrist EF and digit EF axes were
aligned (Figure 4B, purple), but wrist SP
and digit EF were orthogonal (Figure 4C,
purple). One possible explanation is that
finger movements predominantly evoke
flexion and extension torques at the
wrist, with little contribution to pronation
or supination. In participant P5, the EF
and SP neural axes were aligned, but
pointed in opposite directions. Although
the sources of these interactions in the
neural space are unclear, the neural
activity underlying movements of the
wrist and digits is clearly linked.

Simultaneous movement of wrist and digits

Given the interaction between wrist and
digit encoding during independent
movements, we asked whether the
motor cortex would separate them when
participants attempted to move the wrist
and digits simultaneously. After two
sequential instructional cues—first wrist
and then digits—a single tone cued
participants to attempt movement of
both wrist and a specified finger at the
same time (Figure 5A). The different wrist

Figure 4. Overlap between digit and wrist neural axes. A| Distribution of angles between a common neural digit movement direction axis and the same axis computed for each digit independently (C1 n=6 and P5 n=2 for T, I, M, and all; C1 n=1 and P5 n=1 for R, P). "all" corresponds to angles between models computed on different data folds. Black dots represent the median and bars depict the interquartile range (IQR). A dotted red line at 90˚ denotes the vector angle of orthogonality. T: thumb; I: index; M: middle; R: ring; P: pinky. B| Distribution of angles between wrist extension-flexion (EF) axis and wrist supination-pronation (SP) axis and common digit EF axis. (C1 ext/flex n=5, pro/sup n=6; P5 n=2) C| Same as B for the wrist supination-pronation axis.

orientations, digit identities, and digit movement directions were randomized and interleaved. While
the timing of digit-identity information was similar to digit-only trials (1.00s and 960ms before

movement and 205ms and 245ms after mean audio cue end for C1 and C2, respectively, Figure 5B)
and peak digit identity classification accuracies were similar (Supp. Figure 6A, p=0.41), classification
of digit identity was poorer during the preparatory period during simultaneous movements (Supp.
Figure 6B, p<0.0003). A similar pattern was observed for the wrist: similar latency (2.24s and 1.60s
before movement, 33ms and 673ms after mean audio cue end for C1, C2, Figure 5C) but reduced
overall accuracy (Supp. Figure 6C, p=0.0023), particularly during preparation (Supp. Figure 6D,
p=0.0038).

To further probe finger and wrist interactions, we evaluated whether wrist orientation affected digit
identity signals in the motor cortex. We trained decoders of digit identity on one wrist orientation and
tested them on other orientations (Figure 5D). Surprisingly, wrist orientation had no effect on the digit
decoding accuracy in either participant (p=0.45, breakdown by session and training/testing wrist
orientation in Supp. Figure 6EF)

Figure 5. Simultaneous wrist and finger movements. A| Schematic of an example trial. **B|** Time course of digit identity
classification accuracy for subjects C1 (purple) and C2 (pink) based on the neural activity at each time point. Lines and
shading follow the conventions defined in Figure 1D. **C|** B, for wrist orientation classification. **D|** Mean digit classification
accuracies within the same wrist orientation ("same wrist") and across wrist orientations ("other wrist") for each
participant (colors as in B and C). Wilcoxon rank sum test p>0.05 within each session and with all sessions pooled.
Breakdown by session in Supp. Figure 6E. Breakdown by train and test wrist orientation pair in Supp. Figure 6F. (C1 n=5, C2

We then investigated whether the combined movements of the digits and wrist affected the
previously discovered neural geometry. For that, we computed movement direction axes
corresponding to neural spaces for digit extension to flexion (digit EF), wrist extension to flexion (wrist
EF), and wrist supination to pronation (wrist SP) and calculated angles between those axes.
Consistent with our expectations from previous experiments, the SP axis was fully orthogonal to the
digit and wrist EF axes (Figure 5E). The wrist EF axis remained aligned with the digit EF axis (Figure
5F), even for the dataset containing all possible combinations of wrist and digit movements,
underscoring the anatomical and functional connection between wrist and digit flexion-extension
movements.

However, this large overlap could impact our ability to decode desired movements during BCI
control and lead to unintended movements. It would also prevent us from training decoders on
separate sets of digit and wrist movements, forcing exploration of all possible combinations. As a
solution, we removed the shared axis between wrist and digits from neural activity and extracted
task information from the remaining neural space. Specifically, we removed the component of
activity aligned with the common digit axis. Even in this new, reduced neural space, we could decode
which digit was moved and its direction of motion (Figure 5GH). Furthermore, the space retained
reliable information about intended wrist movement (Figure 5I). Restricting the analysis to this neural
subspace allowed for reliable decoding of movement intent and could be completed without the
need to sample all possible combinations of wrist and digit movements.

Online decoding

Building on the principles learned in offline analyses, we implemented an online paradigm to allow
participant C1 to control a virtual robotic hand. We started by training Kalman filter-based position
and velocity decoders as a participant observed 10-20 repetitions of thumb, index, or middle finger
flexion or extension. The participant was then asked to move the digit to the instructed target posture
within 15 seconds while maintaining the other digits in the neutral position, before returning all digits
to the neutral posture (Figure 6A and Supp. Figure 7A, green and gray regions). Testing first with
decoders trained on the full neural space, C1 achieved a success rate above 80% (move success
94% (281/300), return success 81% (218/268), Supp. Figure 7AB). However, movement times were
long (median times of 2.5s, IQR 1.8-4.7s to move to the target and 4.4s, IQR 2.9-7.2s to return, Supp.
Figure 7A). We next built decoders on the neural space without the common movement direction
axis. Under this condition, C1 could achieve target postures and return to neutral significantly more
quickly (to the target: median 2.2s, IQR 1.5-4.2s, p=0.0073, Figure 6B purple swarms; to return to
neutral: 3.5s, IQR 2.8-5.6s, p=0.0122, Figure 6B black swarms; full traces in Figure 6A). Furthermore,
it improved the success rate of movements to the target to 95% (199/210) and of returns to 96%
(191/198) (Figure 6C).

To investigate wrist movement control, we asked the participant to extend, flex, pronate or supinate
their hand, then return to neutral posture. We trained Kalman filter-based position and velocity
decoders on common-axis-subtracted neural space to determine whether the orthogonal subspace
contained enough information for wrist control. On average (median), C1 achieved the target
postures within 1.9s (IQR 0.9-5.2s) and returned to neutral within 1.8s (IQR 1.1-3.0s, Figure 6D). This
was on par with performance in full neural space (p=0.73 and p=0.15 Figure 6E, orange and black
swarms, respectively; Supp. Figure 7CD). C1 achieved an 86% (85/99) success rate during
movement and 93% (74/80) during return (Supp. Figure 7E).

Figure 6. Online control. A| Position during successful individual finger movement trials completed using a shared axis-subtracted decoder, averaged by target (C1 n=3) and time-warped to the median stage length across successful trials. Target window is shaded green. Gray regions represent error threshold for non-target digits and for target digit during return. Red and orange shaded regions surrounding traces depict the standard error of the mean. **B|** Distributions of time to digit "move" (purple swarms) or "relax" (black swarms) phase success for C1 using a decoder built on the full neural space ("full") versus a decoder built on the common axis-subtracted neural subspace ("partial"). Median represented by thicker and IQR represented by thinner horizontal lines. **C|** Trial success rates during movement (purple trace) and relaxation (gray trace) phases for individual finger movements in A, broken down by digit. **D|** Extension/flexion (top row) and pronation/supination (bottom row) during successful wrist movement trials completed using a shared axis-subtracted decoder, averaged by target (C1 n=2). Interpolation and target window shading as in A. Orange regions surrounding traces depict the standard error of the mean. **E|** B, for wrist experiments. Movement phase swarms now depicted in orange. **F|** Distributions of time to digit "move" or "relax" phase success for C1 using a common axis-subtracted decoder while performing individual finger movements ("solo," purple/black swarms) versus simultaneous wrist and finger movements ("joint," green/black swarms).

Lastly, we explored the ability of the participant to make concurrent wrist and digit movements by
training digit movement decoders on digit-only trials and a wrist decoder on wrist-only trials after
subtracting the common extension-to-flexion axis. The participant was asked to flex or extend a
single digit and orient the wrist at the same time. They successfully completed 83% (45/54) of
movement phases and 69% (29/42) of returns, indicating that C1 can selectively flex and extend
digits during wrist movements (Supp. Figure 7FG). Although we could not directly enforce the
concurrency of the movements beyond instructing the participant, movement traces show that all
joints moved toward the target posture together (Supp. Figure 7H). Due to weak wrist decoder
performance on the day of testing and unbeknownst to the participant, however, the decoder
received assistance from the computer with wrist orientation. As in the separate control
experiments, C1 moved the target finger relatively quickly (1.5s, IQR 1.2-2.6s during movement and
4.1s, IQR 3.0-6.6s during relaxation), but stabilizing the other digits and wrist took an additional 1.8
and 2.5 seconds (Supp. Figure 7H, see Supp. Figure 7I for corresponding wrist traces). By subtracting
the common axis from the full neural population activity, we derived decoders that could be trained
independently on each movement group. This approach also reduced unintended digit and wrist
movements, bringing the time to success to the level of separate digit movements (p=0.099, Figure
6F purple and green swarms), reduced the dependence of finger-movement representations on
concurrent wrist movement, and ultimately enabled C1 to control wrist and digit movements
simultaneously.

DISCUSSION

Dexterous hand function relies on neural control signals operating a musculoskeletal system that
couples the digits and wrist through shared tendons and multiarticular muscles. Here we show that
there is a structured overlap across representations of digit and wrist movements in motor cortex as
well, challenging an often implicit assumption that joints are represented independently. These
shared subspaces contained the direction of movement of each joint, consistent with anatomical
organization of wrist and digit musculature.

Movements of individual digits are rarely fully independent: attempting to move or produce force
with one finger typically evokes unintentional motion and/or force in other digits (Kilbreath and
Gandevia, 1994; Hager-Ross and Schieber, 2000; Zatsiorsky, Li and Latash, 2000; Lang and
Schieber, 2004). Peripherally, this interdependence is explained at least in part by shared
musculotendinous anatomy. Multi-tendoned extrinsic hand muscles—such as *flexor digitorum*
profundus and *flexor digitorum superficialis*—send tendons to multiple digits, with incomplete
functional subdivision across fingers. Connective tissue links can also transmit forces between
digits (Reilly and Schieber, 2003; Schieber and Santello, 2004; Duinen and Gandevia, 2011; Lemelin
and Diogo, 2016). Coupling is also evident within the spinal neural circuits that contribute to hand
actions: spinal interneurons contribute to activation of multiple hand muscles and are thought to
support coordinated multi-muscle activation patterns during precision grip (Takei and Seki, 2010;
Takei *et al.*, 2017). One might expect that, upstream, the movement of separate joints would be
abstracted and independent from each other. However, in our data from motor cortex, we observed
strong interactions between neural signals associated with digit flexion and extension, with
movement direction generalizing across digits. This instead supports the view that motor cortical
population activity embeds some properties of the biomechanical constraints on digit control.

We further found that the direction of digit movement (flexion versus extension) was agnostic to the
digit's starting postures. This is notable given that, during grasping, hand-related motor cortical
activity carries prominent posture-related signals (Goodman *et al.*, 2019; Agudelo-Toro *et al.*, 2024),
which are more informative than velocity. In contrast, reaching-related activity in M1 reflects arm
movement velocity (Moran and Schwartz, 1999; Paninski *et al.*, 2004; Wang *et al.*, 2007; Weber *et al.*,
2011). However, even when posture-related modulation is strong, it does not fully obscure
directional information. Ultimately, this posture-tolerant directional signal is consistent with the
peripheral organization of digit control: flexion and extension of individual fingers are driven
predominantly by the same flexor and extensor muscles, even though their mechanics vary with
posture. Appropriately, our Kalman filter-based linear decoder performs best when trained on (and
predicting) both joint angle velocities and positions.

We observed a somatotopic organization of digit identity across the recording arrays, which may
seem surprising given extensive evidence that digit representations in human motor cortex overlap
substantially (Donoghue, Leibovic and Sanes, 1992; Schieber and Hibbard, 1993; Dechent and
Frahm, 2003; Rathelot and Strick, 2006). Importantly, strong overlap does not preclude somatotopy
(Hluštík *et al.*, 2001; Indovina and Sanes, 2001). While we also observed broad overlap across digit-
related activity, by subtracting the average activation signal, we revealed digit-specific hotspots of
activity. These findings extend prior work demonstrating spatial organization of digit-related signals
in human motor cortex, including ECoG and high-density surface recordings, and show that
penetrating recordings can resolve digit-specific spatial organization in humans (Miller *et al.*, 2009;
Wang *et al.*, 2009; Hotson, McMullen, Matthew S. Fifer, *et al.*, 2016; Shelchkova *et al.*, 2023).

Digit movements are often accompanied by the movement of the wrist (Yan, Sobinov and Bensmaia,
2022). Complementing digit decoding, we could also decode intended wrist movements from
activity in the motor cortex. The strength of this representation varied between participants,
however, consistent with sensitivity to recording site and the specific neural populations sampled
by each implant (Vargas-Irwin *et al.*, 2010; Ajiboye *et al.*, 2012). The wrist flexion-extension axis
showed strong alignment with the shared digit flexion-extension (EF) axis, and this alignment was
consistently greater than that between digit EF and wrist supination-pronation (SP). This is
consistent with peripheral biomechanics: the largest extensors and flexors of the digits cross the
wrist, generating a torque that needs to be counterbalanced by the wrist muscles. Similarly,
changing the wrist posture affects the muscle activity producing digit movements and grasping (Seo
et al., 2008; Beringer *et al.*, 2020; Popp *et al.*, 2023).

These effects differ between wrist movements—supination-pronation of the wrist is produced by
muscles that do not cross the wrist and should interact less with the musculature that actuates the
digits. However, we did observe interaction between the associated activity in one of the
participants. Unexpectedly, the digit EF axis aligned with wrist EF in the same direction rather than
the opposite direction, as might be anticipated from purely compensatory wrist torques during finger
flexion. A possible explanation is that the shared population dimension that we read from the neural
activity largely reflects the digit flexion-extension muscle drive. These would flex during wrist flexion
and extend during wrist extension and thus naturally covary with wrist flexion-extension, producing
the observed alignment in the neural space.

Although we did not see an effect of wrist orientation on digit identity decoding, we consistently
observed reduced decoding accuracy of the digits and wrist in preparation for and execution of
simultaneous movements. One explanation is limited bandwidth in the sampled motor cortical
population: as task complexity increases, the depth of modulation related to particular features
within the task may be reduced, an effect seen in monkeys making 2D and 3D virtual reaches
(Rasmussen, Schwartz and Chase, 2017; Ferguson and Cardin, 2020). This effect parallels
normalization computations widely observed in sensory systems, where responses to a feature can
depend on overall stimulus strength, e.g., brightness or volume (Carandini and Heeger, 2012).

Finally, by accounting for the shared structure of digit-related neural activity, we built decoders that
improved the ability of our participant to control multiple digits at the same time: flexing or extending
one digit while not moving the others. This result highlights how the overlap in neural spaces can be
managed rather than treated purely as noise. This performance is encouraging, but it does not yet
achieve the precision, stability, or dimensionality of truly dexterous hand control. Our results
complement and extend other efforts in restoring hand and arm control (Collinger, Wodlinger, *et al.*,
2013; Ajiboye *et al.*, 2017; Nason *et al.*, 2021; Willsey *et al.*, 2025) by focusing on the neural activity
underlying simultaneous control of many digits and the wrist. Given the constraints of the recorded
signal induced by the spatial organization of informative channels, achieving fully dexterous control
will likely require larger spatial sampling, including caudal M1 hidden in the central sulcus, which
contains circuitry more directly linked to distal musculature thought to be critical for individual finger
movement (Lemon and Griffiths, 2005; Rathelot and Strick, 2009). It may also benefit from
recordings across multiple cortical areas that carry complementary information about hand
movements (Guan *et al.*, 2023).

Here, using intracortical microelectrode arrays implanted in the motor cortex of human participants,
we characterized the representation of individuated digit and wrist movements both during isolated
and simultaneous movement attempts. We found somatotopic organization of digit identity across
the arrays alongside robust digit- and wrist-related signals. We also identified a shared low-
dimensional axis in the motor cortex describing digit and wrist flexion-extension that can confound
independent decoding. Exploiting orthogonal dimensions enabled better digit BCI performance,
including during simultaneous wrist movements. This decoding approach should ultimately enable
skilled dexterous control of a prosthetic hand for people with upper limb motor impairments.

METHODS

Participants

Participants gave informed consent prior to participation in a clinical trial (NCT01894802) conducted
under an investigational device exemption from the US Food and Drug Administration and with
approval from the Investigational Review Boards at the Universities of Chicago and Pittsburgh.
Participants C1 and C2 had C4-level ASIA D spinal cord injuries resulting in paralysis of their
dominant right hand. Participant P5 had C8-level ASIA A spinal cord injury resulting in a paralysis of
the dominant right hand. All three participants had two 96-channel microelectrode arrays (Blackrock
Neurotech, Salt Lake City, Utah) implanted in the left motor cortex. Participant C1 had arrays
implanted 35 years after injury occurred and 41-57 months before the experiments reported here.
Participant C2 had arrays implanted 4 years after injury occurred and 11-27 months before the

experiments reported here. Participant P5 had arrays implanted 6 years after the injury and 3-5
months before the experiments reported here.

Classification and significance testing

To identify the presence of the information about the condition (e.g., which digit was being moved,
or wrist posture assumed), we have trained linear discriminant analysis (LDA) models on each
session separately. We have used five-fold stratified cross validation to avoid overfitting. The thin
lines in plots like Figure 1D represent classification accuracy traces across a single session. The
mean accuracy across sessions is illustrated with a thicker line.

To determine significance, a null distribution of raw classification accuracies (*null distribution$_t$*) was
created at each time point by training 100 models per session on trials with randomized target
assignments and testing on held-out trials from the same session (80/20 split with five folds of cross-
validation). The mean classification accuracy trace across sessions was computed, and significance
at each time point was determined as

$$p_t = 1 - normcdf\big(a_t, mean(null\ distribution_t), std(null\ distribution_t)\big)$$

where a_t represents the mean classification accuracy at time t and p_t represents the p-value at time
t. To compensate for multiple comparisons, classification accuracy was considered to be
significantly above chance where the Benjamini & Hochberg-adjusted p<0.05 (pink horizontal line in
e.g. Figure 1D). Specific stage durations varied slightly across sites, as is evident in the stage timing
differences between C1/C2 and P5 in Figure 1D and Figure 2A.

Measuring appearance of conditional information relative to mean prompt completion time

The instruction cue marker in all experiments represents the time at which the audio prompt began
to play, not the time at which the entirety of the audio prompt had been delivered. As such, the
appearance of information about the task conditions in all cases was occasionally reported relative
to the mean prompt completion time for each experiment—the time of the instruction cue plus the
mean duration of the audible portion of the prompt as determined from the relevant .wav files.

Classification generalization matrices

To produce time generalization matrices like the one depicted in Supp. Figure 3C, a series of linear
classifiers were trained on a sliding window of 100ms of neural activity beginning at time x and used
to predict the task condition, e.g., digit movement direction, for each time point over the entirety of
the time course (y). Training and testing data were 80/20 split across trials with five folds of cross-
validation. Supp. Figure 3C and Supp. Figure 5A-D represent mean heatmaps of temporal
generalization across sessions.

To emphasize the negative deviation from chance signifying decoded movement in the opposite
direction in two-class models, Supp. Figure 3C and Supp. Figure 5A-D are presented in units of
adjusted accuracy, where $adjusted\ accuracy = 2 \cdot (raw\ accuracy - 0.5)$. An adjusted accuracy
of 0 means that classification is at chance level. 1 means classification is always consistent with
trial type. -1 means classification is always opposite the trial type.

To determine the threshold of significance conveyed in each heatmap colorbar, 100 linear classifiers
with randomized targets were trained on a sliding window of 100ms of neural activity beginning at

time x and used to predict e.g. digit movement direction over the entirety of the time course. Two-
tailed significance at each time point in a heatmap of adjusted accuracies was determined as

$$p_{x,y} \; = \; 2 \cdot \left(1 - normcdf\left(abs(a_{x,y}),\; mean(null\; distribution_{x,y}),\; std(null\; distribution_{x,y}) \right) \right)$$

where $a_{x,y}$ represents the adjusted accuracy at train time x and test time y and $p_{x,y}$ represents the
p-value at train time x and test time y. As before, the threshold of significance was taken as the
minimum across all x and y of the values of $a_{x,y}$ producing a Benjamini & Hochberg-adjusted p<0.05.

Digit and wrist classification axes

To identify the axes in the neural space defining the extension to flexion movement of digits, we first
defined the periods corresponding to neutral, flexion, and extension of a digit. For neutral, we used
the first 25% of the presentation phase (before completion of the audible instruction). For flexion, we
used the first 50% of the movement phase in flexion trials, and the first 50% of the relaxation phase
in extension trials. Similarly, for extension, we used the first 50% of the movement in extension trials,
and 50% of relaxation in flexion trials. Flexion during movement or relaxation was assigned a
kinematic value of 1, extension was assigned a kinematic value of -1, and time points pre-instruction
were assigned a kinematic value of 0. Then, we trained an LDA model on those periods, extracting
the neural axis from it.

Models derived from the Individual Finger Movement data (Figure 2) were trained on (1) thumb
extension-flexion trials only, (2) index extension-flexion trials only, (3) middle extension-flexion trials
only, (4) ring extension-flexion trials only, (5) pinky extension-flexion trials only, or (6) all-digit
extension-flexion. Models derived from the Wrist Movement data (Figure 3) were trained on (7) wrist
extension-flexion trials only or (8) wrist supination-pronation trials only (with pronation arbitrarily
being analogous to flexion, i.e. having a kinematic value of 1, and supination therefore being
analogous to extension, i.e. having a kinematic value of -1, in training).

To identify the alignment between those neural spaces, we then calculated the vector angle between
the coefficients for every possible pair of models. Because we generated 5 extension-flexion
regression models per category (5 folds of cross-validation), there were 25 vector angles per cross-
categorical model pairing and 10 vector angles per within-category model pairing. The violin plots in
Figure 4 represent the estimated probability density distribution of angles per categorical model
pairing, the open dots depict the median, and the bars show the standard deviation. Colors are
redundant and included only for ease of interpretation.

Calculation of electrode modulation maps

We computed the mean peri-event time histogram in 20ms bins for each motor channel across a
two second period centered on the start of movement. We then identified the response window
during which the absolute difference between average FR_{digit} (across repetitions) and $mean(FR_{all})$
(across repetitions) is at a maximum. Then,

$$modulation\; value = \frac{FR_{digit} - mean(FR_{all})}{mean(FR_{all})}$$

We repeated this process using 1s of the relaxation stage and computed the correlation between the
activation map during movement and the activation map during relaxation. Channels where
$mean(FR_{all}) < 0.5$ spikes/second were ignored and treated as NaNs.

Modulation map significance

At each electrode on each array, we performed two Kruskal-Wallis tests on z-scored responses of
the channel with digit identity and digit extension-flexion as factors (for Individual Finger Movement
sessions) or one Kruskal-Wallis test with wrist orientation as the factor (for Wrist Movement
sessions); the response data for each trial consists of the maximum z-scored response during the
first 50% of the movement stage (when considering modulation significance during movement) or
the maximum z-scored response during the first 50% of the relaxation stage (when considering
modulation significance during relaxation). Spatial electrode modulation significance masks were
created for movement during each session (Supp. Figure 1C, Supp. Figure 2C) by thresholding on the
confidence (i.e. p-value) of the Kruskal-Wallis test for a given factor. Modulation was considered to
be significant if p<0.05. Only significantly modulated electrodes were considered in the creation of
Figure 1F and supplementary modulation maps. To assess the similarity of the positive modulation
in array activity maps representative of individual fingers, the correlation between each pair of
significance-masked maps (thumb vs. index, thumb vs. middle, etc.) was calculated for each
session. These were compared against the distribution of correlations between trial-randomized
maps within sessions. Wilcoxon rank sum tests were performed to establish the significance of the
difference between distributions of correlations.

Protocol for online control of individual fingers

At the start of each session, a regression decoder was trained on neural activity during 10 to 20
repeated observations of automated individual flexions and extensions of the thumb, index, and
middle finger. During observation, the participant was instructed to observe and attempt or imagine
performing the instructed movements. During online control, the participant was given 15s to move
the audibly prompted target digit to its target position while holding all other fingers at the rest
position (movement phase) and 15s to relax all digits to the neutral position once the target position
was achieved (relaxation phase). To achieve a successful movement phase, the target digit had to
remain within 30% of the target angle and each of the non-target digits had to remain within 30% of
the neutral angle for 0.25s. To achieve a successful relaxation phase, the target digit and all non-
target digits had to remain within 30% of the neutral angle for 0.25s. Trials proceeded to the
relaxation phase only if the movement phase was successful. The hand avatar was automatically
reset to the neutral position between each trial. To produce the trial-averaged traces in Figure 6A and
Supp. Figure 7A, movement and relaxation phases were interpolated to the median duration of
successful movement and relaxation phases across all trials.

Protocol for online control of wrist movements

A second decoder, trained on neural activity during 10 to 20 repeated observations of automated
wrist-only movements (neutral, flexion, extension, pronation, supination), was combined with the
decoder previously derived from individual finger movement observations to produce a decoder
capable of simultaneous prediction of individual digit position and wrist roll and pitch. During wrist-
only online control, brain control of finger postures was deactivated, limiting the participant's
control space to wrist roll and pitch. As in digit-only online control, the participant was given 15s to
assume the audibly prompted wrist orientation and 15s to relax to the neutral wrist position. To

achieve successful movement or relaxation, wrist roll and pitch both had to be within 30% of their
target values. The hand avatar was automatically reset to the neutral position between each trial. To
produce the trial-averaged traces in Figure 6D and Supp. Figure 7C, movement and relaxation
phases were interpolated to the median duration of successful movement and relaxation phases
across all trials.

Protocol for simultaneous online control of fingers and wrist

For simultaneous online control of fingers and wrist, brain control of both wrist roll and pitch and
finger postures was enabled. During simultaneous online control, the participant was given 25s to
move the target digit and wrist to the audibly prompted target positions while holding all other fingers
at the rest position (movement phase) and 25s to relax all digits and the wrist to their respective
neutral positions once the target positions were achieved (relaxation phase). As in online control of
individual fingers, digit and wrist angles had to remain within 30% of their targets for 0.25s in order
to achieve movement or relaxation phase success. To produce the trial-averaged traces in Supp.
Figure 7HI, movement and relaxation phases were interpolated to the median duration of successful
movement and relaxation phases across all trials.

Online decoders

For online finger position decoding, a Kalman filter-based regression decoder was used. Principal
component analysis (PCA) was performed on the 256 channels of neural activity recorded during
observation and the first N components that could explain 90% of the variance in the data were used
in training. The coefficient transformation matrix was stored for later application to incoming neural
activity during online prediction. We extracted the angular position descriptive of each finger's
movement over the course of the observation period (ranging from -1 for full extension to +1 for full
flexion) and combined those traces with the time courses of the first N PCs to produce a model of
the common digit extension-flexion axis. After subtracting the common axis from neural activity, we
trained a digit position-estimating Kalman filter. We also found the gradient of each of the five
angular position traces to produce five corresponding angular velocity traces and combined those
traces with the common axis-subtracted time courses of the first N PCs to produce a digit velocity-
estimating Kalman filter. During online control, the estimated position of each digit was taken as the
mean of the prediction from the position-estimating filter and the compounded position estimate
from the velocity-estimating filter.

A similar procedure was repeated for online wrist control, focused on extracting the roll and pitch of
the wrist using position- and velocity-estimating Kalman filters. The number and identities of the PCs
explaining 90% of the variance in the data were reevaluated using the combined set of individual
finger and wrist observation data. The common digit extension-flexion axis was recalculated in the
new PC space and subtracted from neural activity prior to training and during online prediction.

Subtracting the common extension-flexion axis from neural activity

629 The axis describing the common digit extension-flexion neural activity was calculated as

$$A_{common} = \frac{A_{raw} \cdot C_{common}}{|C_{common}|^2} \cdot C_{common}$$

631 where A_{raw} is the PC-transformed incoming neural signal, C_{common} are the estimated coefficients
632 of the common digit extension-flexion axis, and $\cdot$ represents the dot product of vectors. In decoders

633 with shared axis subtraction, the vector A_{common} was subtracted from the PC-transformed
634 incoming neural signal prior to Kalman filter training and prediction.

AUTHOR CONTRIBUTIONS

636 AMXE, ARS, and SJB designed the study with input from other authors. AMXE, EVO, GHB, and ARS
637 performed the experiments. AMXE and ARS performed the analysis. AMXE wrote the first draft of the
638 manuscript. AMXE and ARS wrote the paper with input from all authors.

COMPETING INTERESTS

640 The authors declare no competing interests.

ACKNOWLEDGEMENTS

642 The authors would like to thank the participants for their time. The authors would like to thank
643 Sensorimotor Bionics Group at the University of Chicago and interuniversity Cortical Bionics
644 Research Group for insightful discussions and technical support. This work was supported by the
645 National Institute of Neurological Disorders and Stroke, R35 NS122333, R01 NS131953 and R01
646 NS130302.

DATA AVAILABILITY

648 All data generated or analyzed during this study as well as analysis and plotting scripts are going to
649 be made available on DABI upon publication.

CODE AVAILABILITY

651 All analysis and plotting scripts are going to be made available in a public repository
652 https://github.com/SobinovLab/finger_individuation

REFERENCES

Acharya, S. *et al.* (2007) "Towards a Brain-Computer Interface for Dexterous Control of a Multi-Fingered Prosthetic Hand," *2007 3rd International IEEE/EMBS Conference on Neural Engineering*. *2007 3rd International IEEE/EMBS Conference on Neural Engineering*, Kohala Coast, HI, USA: IEEE, pp. 200–203. Available at: https://doi.org/10.1109/CNE.2007.369646.

Adewuyi, A.A., Hargrove, L.J. and Kuiken, T.A. (2017) "Resolving the effect of wrist position on myoelectric pattern recognition control," *Journal of NeuroEngineering and Rehabilitation*, 14(1), p. 39. Available at: https://doi.org/10.1186/s12984-017-0246-x.

Aggarwal, V. *et al.* (2008) "Asynchronous Decoding of Dexterous Finger Movements Using M1 Neurons," *IEEE Transactions on Neural Systems and Rehabilitation Engineering*, 16(1), pp. 3–14. Available at: https://doi.org/10.1109/TNSRE.2007.916289.

Aggarwal, V. *et al.* (2013) "State-based decoding of hand and finger kinematics using neuronal ensemble and LFP activity during dexterous reach-to-grasp movements," *Journal of Neurophysiology*, 109(12), pp. 3067–3081. Available at: https://doi.org/10.1152/jn.01038.2011.

Agudelo-Toro, A. *et al.* (2024) "Accurate neural control of a hand prosthesis by posture-related activity in the primate grasping circuit," *Neuron*, 0(0). Available at: https://doi.org/10.1016/j.neuron.2024.09.018.

Ajiboye, A.B. *et al.* (2012) "Prediction of Imagined Single-Joint Movements in a Person with High Level Tetraplegia," *IEEE transactions on bio-medical engineering*, 59(10), pp. 2755–2765. Available at: https://doi.org/10.1109/TBME.2012.2209882.

Ajiboye, A.B. *et al.* (2017) "Restoration of reaching and grasping movements through brain-controlled muscle stimulation in a person with tetraplegia: a proof-of-concept demonstration.," *Lancet (London, England)*, 389(10081), pp. 1821–1830. Available at: https://doi.org/10.1016/S0140-6736(17)30601-3.

Anderson, K.D. (2004) "Targeting Recovery: Priorities of the Spinal Cord-Injured Population," *Journal of Neurotrauma*, 21(10), pp. 1371–1383. Available at: https://doi.org/10.1089/neu.2004.21.1371.

Baldwin, M.K.L. *et al.* (2018) "Representations of fine digit movements in posterior and anterior parietal cortex revealed using long-train intracortical microstimulation in macaque monkeys," *Cerebral Cortex*, 28(12), pp. 4244–4263. Available at: https://doi.org/10.1093/cercor/bhx279.

Beringer, C.R. *et al.* (2020) "The effect of wrist posture on extrinsic finger muscle activity during single joint movements," *Scientific Reports*, 10(1), p. 8377. Available at: https://doi.org/10.1038/s41598-020-65167-x.

Branco, M.P. *et al.* (2019) "Encoding of kinetic and kinematic movement parameters in the sensorimotor cortex: A Brain-Computer Interface perspective," *European Journal of Neuroscience*, 50(5), pp. 2755–2772. Available at: https://doi.org/10.1111/ejn.14342.

Capaday, C. *et al.* (2013) "On the functional organization and operational principles of the motor cortex," *Frontiers in Neural Circuits*, 7. Available at: https://doi.org/10.3389/fncir.2013.00066.

Carandini, M. and Heeger, D.J. (2012) "Normalization as a canonical neural computation," *Nature*
Reviews Neuroscience, 13(1), pp. 51–62. Available at: https://doi.org/10.1038/nrn3136.

Chehade, N.G. and Gharbawie, O.A. (2023) "Motor actions are spatially organized in motor and
dorsal premotor cortex," *eLife*. Edited by J. Diedrichsen, J.I. Gold, and W. Vanduffel, 12, p. e83196.
Available at: https://doi.org/10.7554/eLife.83196.

Collinger, J.L., Boninger, M.L., *et al.* (2013) "Functional priorities, assistive technology, and brain-
computer interfaces after spinal cord injury," *The Journal of Rehabilitation Research and*
Development, 50(2), p. 145. Available at: https://doi.org/10.1682/JRRD.2011.11.0213.

Collinger, J.L., Wodlinger, B., *et al.* (2013) "High-performance neuroprosthetic control by an
individual with tetraplegia," *The Lancet*, 381(9866), pp. 557–564. Available at:
https://doi.org/10.1016/S0140-6736(12)61816-9.

Dechent, P. and Frahm, J. (2003) "Functional somatotopy of finger representations in human primary
motor cortex," *Human Brain Mapping*, 18(4), pp. 272–283. Available at:
https://doi.org/10.1002/hbm.10084.

Dekleva, B.M. and Collinger, J.L. (2025) "Using transient, effector-specific neural responses to gate
decoding for brain–computer interfaces," *Journal of Neural Engineering*, 22(1), p. 016036. Available
at: https://doi.org/10.1088/1741-2552/adaa1f.

Donoghue, J.P., Leibovic, S. and Sanes, J.N. (1992) "Organization of the forelimb area in squirrel
monkey motor cortex: representation of digit, wrist, and elbow muscles," *Experimental Brain*
Research, 89(1). Available at: https://doi.org/10.1007/BF00228996.

Duinen, H. van and Gandevia, S.C. (2011) "Constraints for control of the human hand," *The Journal*
of Physiology, 589(23), pp. 5583–5593. Available at: https://doi.org/10.1113/jphysiol.2011.217810.

Ejaz, N., Hamada, M. and Diedrichsen, J. (2015) "Hand use predicts the structure of representations
in sensorimotor cortex.," *Nature neuroscience*, 18(7), pp. 1034–40. Available at:
https://doi.org/10.1038/nn.4038.

Ferguson, K.A. and Cardin, J.A. (2020) "Mechanisms underlying gain modulation in the cortex,"
Nature reviews. Neuroscience, 21(2), pp. 80–92. Available at: https://doi.org/10.1038/s41583-019-
0253-y.

Flesher, S.N. *et al.* (2021) "A brain-computer interface that evokes tactile sensations improves
robotic arm control," *Science*, 372(6544), pp. 831–836. Available at:
https://doi.org/10.1126/science.abd0380.

Georgopoulos, A.P. *et al.* (1999) "Neural coding of finger and wrist movements," *Journal of*
Computational Neuroscience, 6(3), pp. 279–288. Available at:
https://doi.org/10.1023/a:1008810007672.

Goodman, J.M. *et al.* (2019) "Postural Representations of the Hand in the Primate Sensorimotor
Cortex.," *Neuron*, p. 566539. Available at: https://doi.org/10.1016/j.neuron.2019.09.004.

Guan, C. *et al.* (2023) "Decoding and geometry of ten finger movements in human posterior parietal
cortex and motor cortex," *Journal of Neural Engineering*, 20(3), p. 036020. Available at:
https://doi.org/10.1088/1741-2552/acd3b1.

Hager-Ross, C. and Schieber, M.H. (2000) "Quantifying the independence of human finger
movements: Comparisons of digits, hands, and movement frequencies," *Journal of Neuroscience*,
20(22), pp. 8542–8550. Available at: https://doi.org/10.1523/jneurosci.20-22-08542.2000.

Hluštík, P. *et al.* (2001) "Somatotopy in Human Primary Motor and Somatosensory Hand
Representations Revisited," *Cerebral Cortex*, 11(4), pp. 312–321. Available at:
https://doi.org/10.1093/cercor/11.4.312.

Hotson, G., McMullen, D.P., Fifer, Matthew S, *et al.* (2016) "Individual finger control of a modular
prosthetic limb using high-density electrocorticography in a human subject," *Journal of Neural
Engineering*, 13(2), p. 026017. Available at: https://doi.org/10.1088/1741-2560/13/2/026017.

Hotson, G., McMullen, D.P., Fifer, Matthew S., *et al.* (2016) "Individual Finger Control of the Modular
Prosthetic Limb using High-Density Electrocorticography in a Human Subject," *Journal of neural
engineering*, 13(2), p. 026017. Available at: https://doi.org/10.1088/1741-2560/13/2/026017.

Indovina, I. and Sanes, J.N. (2001) "On Somatotopic Representation Centers for Finger Movements
in Human Primary Motor Cortex and Supplementary Motor Area," *NeuroImage*, 13(6), pp. 1027–
1034. Available at: https://doi.org/10.1006/nimg.2001.0776.

Irwin, Z.T. *et al.* (2017) "Neural control of finger movement via intracortical brain–machine interface,"
Journal of Neural Engineering, 14(6), p. 066004. Available at: https://doi.org/10.1088/1741-
2552/aa80bd.

Jorge, A. *et al.* (2020) "Classification of Individual Finger Movements Using Intracortical Recordings
in Human Motor Cortex," *Neurosurgery*, 87(4), pp. 630–638. Available at:
https://doi.org/10.1093/neuros/nyaa026.

Jude, J.J. *et al.* (2026) "Restoring rapid natural bimanual typing with a neuroprosthesis after
paralysis," *Nature Neuroscience*, pp. 1–10. Available at: https://doi.org/10.1038/s41593-026-
02218-y.

Kilbreath, S.L. and Gandevia, S.C. (1994) "Limited independent flexion of the thumb and fingers in
human subjects.," *The Journal of Physiology*, 479(3), pp. 487–497. Available at:
https://doi.org/10.1113/jphysiol.1994.sp020312.

Kirsch, E., Rivlis, G. and Schieber, M.H. (2014) "Primary Motor Cortex Neurons during Individuated
Finger and Wrist Movements: Correlation of Spike Firing Rates with the Motion of Individual Digits
versus Their Principal Components," *Frontiers in Neurology*, 5, p. 70. Available at:
https://doi.org/10.3389/fneur.2014.00070.

Kubánek, J. *et al.* (2009) "Decoding flexion of individual fingers using electrocorticographic signals in
humans," *Journal of Neural Engineering*, 6(6), p. 066001. Available at: https://doi.org/10.1088/1741-
2560/6/6/066001.

Lang, C.E. and Schieber, M.H. (2004) "Reduced Muscle Selectivity During Individuated Finger Movements in Humans After Damage to the Motor Cortex or Corticospinal Tract," *Journal of Neurophysiology*, 91(4), pp. 1722–1733. Available at: https://doi.org/10.1152/jn.00805.2003.

Lemelin, P. and Diogo, R. (2016) "Anatomy, Function, and Evolution of the Primate Hand Musculature," in T.L. Kivell et al. (eds.) *The Evolution of the Primate Hand*. New York.

Lemon, R.N. and Griffiths, J. (2005) "Comparing the function of the corticospinal system in different species: Organizational differences for motor specialization?," *Muscle and Nerve*, 32(3), pp. 261–279. Available at: https://doi.org/10.1002/mus.20333.

Mender, M.J. *et al.* (2024) "Functional Electrical Stimulation and Brain-Machine Interfaces for Simultaneous Control of Wrist and Finger Flexion." Neuroscience. Available at: https://doi.org/10.1101/2024.08.11.607263.

Miller, K.J. *et al.* (2009) "Decoupling the Cortical Power Spectrum Reveals Real-Time Representation of Individual Finger Movements in Humans," *Journal of Neuroscience*, 29(10), pp. 3132–3137. Available at: https://doi.org/10.1523/JNEUROSCI.5506-08.2009.

Mohamed, A.-K. and Aharonson, V. (2021) "Four-class BCI discrimination of right and left wrist and finger movements," *IFAC-PapersOnLine*, 54(21), pp. 91–96. Available at: https://doi.org/10.1016/j.ifacol.2021.12.016.

Moran, D.W. and Schwartz, A.B. (1999) "Motor cortical representation of speed and direction during reaching," *Journal of Neurophysiology*, 82, pp. 2676–2692.

Nason, S.R. *et al.* (2021) "Real-time linear prediction of simultaneous and independent movements of two finger groups using an intracortical brain-machine interface," *Neuron*, 109(19), pp. 3164-3177.e8. Available at: https://doi.org/10.1016/j.neuron.2021.08.009.

Nason-Tomaszewski, S.R. *et al.* (2023) "Restoring continuous finger function with temporarily paralyzed nonhuman primates using brain–machine interfaces," *Journal of Neural Engineering*, 20(3), p. 036006. Available at: https://doi.org/10.1088/1741-2552/accf36.

Okorokova, E.V. *et al.* (2020) "Decoding hand kinematics from population responses in sensorimotor cortex during grasping," *Journal of Neural Engineering*, 17(4), p. 046035. Available at: https://doi.org/10.1088/1741-2552/ab95ea.

Paninski, L. *et al.* (2004) "Spatiotemporal Tuning of Motor Cortical Neurons for Hand Position and Velocity," *Journal of Neurophysiology* [Preprint]. Available at: https://doi.org/10.1152/jn.00587.2002.

Penfield, W. and Rasmussen, T. (1950) *The cerebral cortex of man: a clinical study of localization of function*. New York: Macmillan.

Popp, W.L. *et al.* (2023) "Effects of wrist posture and stabilization on precision grip force production and muscle activation patterns," *Journal of Neurophysiology* [Preprint]. Available at: https://doi.org/10.1152/jn.00420.2020.

Rasmussen, R.G., Schwartz, A. and Chase, S.M. (2017) "Dynamic range adaptation in primary motor
cortical populations," *eLife*, 6, p. e21409. Available at: https://doi.org/10.7554/eLife.21409.

Rastogi, A. *et al.* (2021) "The Neural Representation of Force across Grasp Types in Motor Cortex of
Humans with Tetraplegia," *eNeuro*, 8(1). Available at: https://doi.org/10.1523/ENEURO.0231-
20.2020.

Rathelot, J.A. and Strick, P.L. (2006) "Muscle representation in the macaque motor cortex: An
anatomical perspective," *Proceedings of the National Academy of Sciences of the United States of
America*, 103(21), pp. 8257–8262. Available at: https://doi.org/10.1073/pnas.0602933103.

Rathelot, J.A. and Strick, P.L. (2009) "Subdivisions of primary motor cortex based on cortico-
motoneuronal cells," *Proceedings of the National Academy of Sciences of the United States of
America*, 106(3), pp. 918–923. Available at: https://doi.org/10.1073/pnas.0808362106.

Reilly, K.T. and Schieber, M.H. (2003) "Incomplete functional subdivision of the human
multitendoned finger muscle flexor digitorum profundus: An electromyographic study," *Journal of
Neurophysiology*, 90(4), pp. 2560–2570. Available at: https://doi.org/10.1152/jn.00287.2003.

Rizzolatti, G. and Luppino, G. (2001) "The Cortical Motor System," *Neuron*, 31(6), pp. 889–901.
Available at: https://doi.org/10.1016/S0896-6273(01)00423-8.

Schieber, M.H. (2001) "Constraints on somatotopic organization in the primary motor cortex,"
Journal of Neurophysiology, 86(5), pp. 2125–2143. Available at:
https://doi.org/10.1152/jn.2001.86.5.2125.

Schieber, M.H. and Hibbard, L.S. (1993) "How somatotopic is the motor cortex hand area?,"
Science, 261(5120), pp. 489–492. Available at: https://doi.org/10.1126/science.8332915.

Schieber, M.H. and Santello, M. (2004) "Hand function: Peripheral and central constraints on
performance," *Journal of Applied Physiology*, 96(6), pp. 2293–2300. Available at:
https://doi.org/10.1152/japplphysiol.01063.2003.

Seo, N.J. *et al.* (2008) "Wrist strength is dependent on simultaneous power grip intensity.,"
Ergonomics, 51(10), pp. 1594–1605. Available at: https://doi.org/10.1080/00140130802216925.

Shah, N.P. *et al.* (2023) "A brain-computer typing interface using finger movements," *2023 11th
International IEEE/EMBS Conference on Neural Engineering (NER)*. *2023 11th International
IEEE/EMBS Conference on Neural Engineering (NER)*, Baltimore, MD, USA: IEEE, pp. 1–4. Available
at: https://doi.org/10.1109/NER52421.2023.10123912.

Shelchkova, N.D. *et al.* (2023) "Microstimulation of human somatosensory cortex evokes task-
dependent, spatially patterned responses in motor cortex," *Nature Communications*, 14(1), p. 7270.
Available at: https://doi.org/10.1038/s41467-023-43140-2.

Sobinov, A.R. and Bensmaia, S.J. (2021) "The neural mechanisms of manual dexterity," *Nature
Reviews Neuroscience*, pp. 1–17. Available at: https://doi.org/10.1038/s41583-021-00528-7.

Takei, T. *et al.* (2017) "Neural basis for hand muscle synergies in the primate spinal cord,"
Proceedings of the National Academy of Sciences, 114(32), pp. 8643–8648. Available at:
https://doi.org/10.1073/pnas.1704328114.

Takei, T. and Seki, K. (2010) "Spinal Interneurons Facilitate Coactivation of Hand Muscles during a
Precision Grip Task in Monkeys," *Journal of Neuroscience*, 30(50), pp. 17041–17050. Available at:
https://doi.org/10.1523/JNEUROSCI.4297-10.2010.

Vargas-Irwin, C.E. *et al.* (2010) "Decoding Complete Reach and Grasp Actions from Local Primary
Motor Cortex Populations," *Journal of Neuroscience*, 30(29), pp. 9659–9669. Available at:
https://doi.org/10.1523/JNEUROSCI.5443-09.2010.

Vargas-Irwin, C.E. *et al.* (2024) "Gesture encoding in human left precentral gyrus neuronal
ensembles." Neuroscience. Available at: https://doi.org/10.1101/2024.08.23.608325.

Vaskov, A.K. *et al.* (2018) "Cortical Decoding of Individual Finger Group Motions Using ReFIT Kalman
Filter," *Frontiers in Neuroscience*, 12. Available at:
https://www.frontiersin.org/articles/10.3389/fnins.2018.00751 (Accessed: August 11, 2023).

Wang, W. *et al.* (2007) "Motor Cortical Representation of Position and Velocity During Reaching,"
Journal of Neurophysiology, 97(6), pp. 4258–4270. Available at:
https://doi.org/10.1152/jn.01180.2006.

Wang, W. *et al.* (2009) "Human Motor Cortical Activity Recorded with Micro-ECoG Electrodes During
Individual Finger Movements," *Conference proceedings : ... Annual International Conference of the*
IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society.
Conference, 2009, pp. 586–589. Available at: https://doi.org/10.1109/IEMBS.2009.5333704.

Weber, D.J. *et al.* (2011) "Limb-state information encoded by peripheral and central somatosensory
neurons: Implications for an afferent interface," *IEEE Transactions on Neural Systems and*
Rehabilitation Engineering, 19(5), pp. 501–513. Available at:
https://doi.org/10.1109/TNSRE.2011.2163145.

Willsey, M.S. *et al.* (2025) "A high-performance brain–computer interface for finger decoding and
quadcopter game control in an individual with paralysis," *Nature Medicine*, 31(1), pp. 96–104.
Available at: https://doi.org/10.1038/s41591-024-03341-8.

Wodlinger, B. *et al.* (2015) "Ten-dimensional anthropomorphic arm control in a human brain-
machine interface: difficulties, solutions, and limitations.," *Journal of neural engineering*, 12(1), p.
016011. Available at: https://doi.org/10.1088/1741-2560/12/1/016011.

Yan, Y., Sobinov, A.R. and Bensmaia, S.J. (2022) "Prehension kinematics in humans and macaques,"
Journal of Neurophysiology [Preprint]. Available at: https://doi.org/10.1152/jn.00522.2021.

Zatsiorsky, V.M., Li, Z.-M. and Latash, M.L. (2000) "Enslaving effects in multi-finger force
production," *Experimental Brain Research*, 131(2), pp. 187–195. Available at:
https://doi.org/10.1007/s002219900261.

Supp. Figure 1. A| Modulation by digit identity (thumb, index, middle, ring, and pinky, red to yellow borders) in C2's lateral motor array during solitary digit movement, averaged across sessions. Modulation by digit identity in P5's medial motor array during solitary digit movement, averaged across sessions. Non-significant channels are gray. Anterior-most medial corner of arrays is black. **B|** Distributions (across sessions) of correlation of the modulation of array activity maps of the annotated pair of digits (T, thumb; I, index; M, middle; R, ring; P, pinky) in C1, C2, and P5. p-values represent the result of a significance test between the correlation distribution for trial-randomized maps (mean per subject presented as red dotted line; C1: -0.17 , C2: -0.18, P5: -0.33) and the denoted pair of maps (R:P, ring vs. pinky). Top and bottom edges of each box are 75th and 25th percentiles. Line mid-box is median. Whiskers present maximum and minimum non-outliers. Outliers are represented as separate dots. * p<0.05, ** p<0.01, *** p<0.001. (C1 n=5, C2 n=6, P5 n=3) **C|** Kruskal-Wallis confidence in significance of electrode modulation by digit identity during movement in the lateral array (C1, C2) or medial array (P5) during individual finger movement sessions. p = (100-confidence)/100. See Methods for details. Anterior-most medial corner of arrays is pink.

Supp. Figure 2. A| Classification of digit movement direction – flexion or extension – for individual digits. Colored bars above traces correspond to points significantly deviating from chance (red line). Pale pink line depicts chance significance threshold. (C1 n=5, C2 n=2, P5 n=3). **B|** Digit movement direction modulation array activity maps for digit extension (left), digit flexion (center), and digit extension minus flexion (right) in C1, C2, and P5. Non-significant channels are gray. Anterior-most medial corner of arrays is black. **C|** Kruskal-Wallis confidence in significance of electrode modulation by digit extension-flexion during movement in the lateral array (C1, C2) or medial array (P5) during individual finger movement sessions. p = (100-confidence)/100. See Methods for details. Anterior-most medial corner of arrays is pink.

Supp. Figure 3. A| The placement of the linear classifier's extension-flexion hyperplane in neural space is such that the flexion during relaxation in an extension trial (pink dotted line) is more similar to the flexion during movement in a flexion trial (blue solid line) than it is to the extension during movement in an extension trial (pink solid line). **B|** Consequently, movement occurring during the relaxation phase is reliably classified as opposite the trial type (green coloration on dotted lines). **C|** Temporal generalization heatmaps of digit movement direction for C1, C2, and P5, averaged across digits. Color corresponds to the classification of the instructed movement direction (purple) or movement in the opposite direction (green). In C1 and P5, any single finger flexion or extension is a composite movement of both flexion and extension.

Supp. Figure 4. A| Generalization of movement direction across digits during movement phase in subjects C1, C2, and P5. The extension-flexion classifier was trained on a single digit and tested on the same digit ("same digit") or a different digit ("other digit") or trained on a pooled dataset containing all digits and tested on held-out examples of all digits ("all digits"). Each dot represents accuracy in a single session for a specific training and testing digit combination. Dot color corresponds to session. Horizontal solid red line illustrates chance level. * p<0.05, ** p<0.01, *** p<0.001. B| Average digit extension-flexion classification accuracy across sessions for all combinations of training and testing digit identity in C1, C2, and P5. A strong diagonal suggests that extension-flexion decoding is most successful when training and testing on the same digit, but classification accuracies well above chance (50%) for other train/test digit pairs provide evidence for a shared extension-flexion classification axis across digits.

Supp. Figure 5. A| Temporal generalization heatmap of wrist extension-flexion movement direction for C1 and P5. Color
corresponds to instructed (orange) or opposite (blue) direction classification. A wrist orientation classifier trained on the
movement stage or the relaxation stage correctly identifies the motion occurring during relaxation as the inverse of the
motion occurring during movement. **B|** Same as A for supination-pronation axis. **C|** Distinct phases in A are not visible in
C1 or P5 when performing analysis using non-complementary wrist orientations (left in subject pair: neutral-extension,
right in subject pair: neutral-flexion). **D|** Distinct phases in B are not visible in C1 or P5 when performing analysis using non-
complementary wrist orientations (left in subject pair: neutral-pronation, right in subject pair: neutral-supination).

Supp. Figure 6. A| Distribution of maximum accuracy of digit classification during movement in digit-only trials and
simultaneous digit and wrist trials (C1: purple, C2: pink). **B|** Distribution of maximum accuracy of digit classification during
the preparatory period in digit-only trials and simultaneous digit and wrist trials. Colors as in A. **C|** Same as A for wrist
orientation classification. **D|** Same as B for wrist orientation classification. **E|** Digit classification training and testing at
different wrist orientations, color-coded by session. The wide span of digit classification accuracies is due to large
variations in overall performance across sessions, not due to deficits specific to training/testing wrist combinations within
a session. Consistent and mixed wrist swarms are not significantly different within any single session. (C1 n=5, C2 n=2) **F|**
Accuracy of digit classification in C1 when trained within one wrist orientation and tested on another for C1 and C2.

Supp. Figure 7. A| Position during successful individual finger movement trials completed using a decoder without shared axis subtraction, averaged by target (C1 n=4) and interpolated to the median stage length across successful trials. Target window is shaded green. Gray regions represent error threshold for non-target digits and for target digit during return. Red and orange shaded regions surrounding traces depict the standard error of the mean. **B|** Trial success rates during movement (purple trace) and relaxation (gray trace) phases for trials in A, broken down by digit. **C|** Extension/flexion (top row) and pronation/supination (bottom row) during successful wrist movement trials completed using a decoder without shared axis subtraction, averaged by target (C1 n=1). Interpolation and target window shading as in A. Orange regions surrounding traces depict the standard error of the mean. Median time to movement success 2.3s (IQR 0.9-3.3s). Median time to relaxation success 2.8s (IQR 1.4-4.4s). **D|** Same as B, but for wrist control trials in C. Move success 90% (36/40). Relax success 91% (31/34). **E|** Same as D, but for wrist control trials completed using a decoder with shared axis subtraction. **F|** Digit success rates during the movement phase of simultaneous wrist and finger movements, grouped by wrist motion and broken down by digit. **G|** F, for the relaxation phase of simultaneous wrist and finger movements. **H|** Digit position during successful simultaneous wrist and finger movement trials completed using a decoder with shared axis subtraction, averaged by target (C1 n=1) and interpolated to the median stage length across successful trials. Colors as in A. **I|** Wrist movements during successful simultaneous wrist and finger trials, with brain control of wrist at 20%.