Chromatin priming and co-factor availability shape lineage response to the neuronal pioneer factor ASCL1 in pluripotency

Jethro Lundie-Brown[1], Rosalind Drummond[1], John-Poul Ng-Blichfeldt[1], Roberta Azzarelli[1,3]* and Anna Philpott[1,2]*

1 Cambridge Stem Cell Institute, University of Cambridge, Cambridge, CB2 0AW, UK

2 Department of Oncology, University of Cambridge, Cambridge, CB2 0AH, UK

3 Department of Pharmacology, UCL School of Pharmacy, 29-39 Brunswick Square, London, WC1N 1AX, UK

* Co-corresponding authors

HIGHLIGHTS

• Ectopic ASCL1 drives non-neuronal transcriptional responses in naïve and formative pluripotent cells

• ASCL1 occupies distinct, predominantly non-neuronal genomic targets in pluripotent cells due to differential chromatin accessibility

• ASCL1 pioneer activity is locus- and cell type-specific and predicted by histone acetylation status

• Co-expression of ASCL1 with *Phox2* homeodomain cofactors potentiates neuronal lineage acquisition in pluripotent cells

SUMMARY

Transcription factors often act within defined developmental windows, yet how naïve pluripotent cells acquire competence to execute specific transcription factor-driven fate programmes remains unclear. Pioneer transcription factors that engage target sites in closed chromatin to initiate gene expression programmes often act at the top of hierarchies in cell identity transitions. However, we show that the ability of ASCL1 to induce a coherent neuronal programme emerges only after exit from pluripotency, coincident with progressive chromatin remodelling and accumulation of permissive histone marks at neuronal ASCL1 target sites. Binding analysis reveals that although ASCL1 can access a subset of neuronal loci in mESCs and EpiLCs, ASCL1 is preferentially diverted to non-neuronal sites, resulting in divergent transcriptional responses. Increasing global histone acetylation enhances activation of individual neuronal genes but is insufficient to drive full neuronal differentiation. In contrast, co-expression of the homeodomain transcription factor PHOX2B redirects ASCL1 towards neuronal targets while suppressing inappropriate programmes in mESCs. These findings demonstrate that ASCL1 pioneer activity is highly context-dependent and that developmental priming of chromatin is essential for appropriate lineage specification.

INTRODUCTION

Development requires precise control over the generation of diverse cell types. Achieving the correct spatial and temporal coordination of growth, differentiation, and morphogenesis requires cells to regulate their competence to respond to inductive cues. However, the mechanisms by which cells acquire and maintain this competence remain poorly understood.

Mouse embryonic stem cells (mESCs) have proven an invaluable tool for studying competence for differentiation. The transition from naïve to formative pluripotency has revealed insights into the reorganisation of chromatin and transcription factor availability that prepares cells for subsequent state transitions.[1–5] However, the effect of these transitions on competence for differentiation in response to lineage-specifying transcription factors (TFs) remains incompletely understood. ESCs are widely used in forward programming strategies, in which TF overexpression drives direct conversion to specific cell types in vitro.[6–8] These approaches typically require prior priming through altered culture conditions or embryoid body formation, suggesting that naïve pluripotent cells are not inherently competent to respond to lineage-specifying TFs. Understanding how such TFs act in naïve pluripotent cells would therefore clarify both the biology of this foundational cell state and the epigenetic basis by which competence for TF-driven cell fate change is established.

Competence to respond to TFs is largely directed by the epigenetic landscape within a cell. The epigenetic state of naïve pluripotency is distinct from other tissues and reflects their capacity for multilineage differentiation and their transient nature in development.[9–12] Naïve pluripotent cells typically exhibit higher levels of chromatin accessibility compared with differentiated cells.[9,13,14] Although open chromatin does not necessarily determine enhancer activity,[13] active enhancers usually display a local depletion of nucleosome occupancy.[15] Many enhancers in early development are accessible but maintained in a poised state marked by the presence of both activation-associated H3K4me1 and repressive H3K27me3, rather than having the H3K27ac signature of active enhancers.[15–18] The ability of a cell to interpret and respond to lineage-specifying TFs is therefore determined partially by the availability of TFs and their recruitment to these regulatory elements, and partially by the chromatin states that permit or restrict their binding and activity.[19–21]

Recent work on cell fate reprogramming shows that certain TFs display pioneer activity, an ability to bind their preferred motifs in nucleosome-dense chromatin and initiate opening and activation of target gene expression.[22,23] Pioneer factors often sit at the apex of transcriptional networks to drive an entire lineage program during both development and reprogramming,[24,25] in some cases directly establishing competence for differentiation.[22,26,27] However, the simple distinction between pioneer and non-pioneer TFs has been challenged,[28] as some pioneer factors display different preferences for open and closed chromatin across cell types and at different loci.[29–32] Their activity is also frequently modulated by cofactors that refine or redirect their function, and TFs often cooperate to control cell fate in development and reprogramming.[33–37]

ASCL1, a proneural bHLH TF, drives both fibroblast-to-neuron conversion[38,39] and
neural stem cell differentiation *in vivo* and *in vitro*.[40,41] ASCL1 has been described as a
pioneer factor due to its ability to bind nucleosomal DNA directly.[42] However, it also binds
partially accessible chromatin and recruits chromatin remodelling enzymes.[32] Evidence
from somatic reprogramming has further suggested that competence for ASCL1-
mediated conversion correlates with a specific epigenetic signature at its binding sites,
[38,39] yet how epigenetic constraints shape the developmental acquisition of this
competence remains largely unexplored. ASCL1-mediated differentiation therefore
represents a tractable system for dissecting how the epigenetic landscape of early
developmental states determines competence to respond to lineage-specifying pioneer
TFs.[7,43–45]

Here, we investigate when and how cells acquire competence to respond to
ASCL1 during the exit from pluripotency. By integrating RNA-seq, ASCL1 ChIP-seq, and
chromatin accessibility mapping across naïve mESCs, epiblast-like cells (EpiLCs), and
neuroectoderm (NE) states, we show that ASCL1 binding is highly context-dependent:
key neuronal target loci are inaccessible and transcriptionally refractory in pluripotent
cells, while genes associated with divergent pathways are readily activated. We find that
the onset of competence to respond to ASCL1 with a coherent neuronal programme
correlates with progressive chromatin remodelling and acquisition of permissive histone
modifications at neuronal ASCL1 target sites. The ability of ASCL1 to drive neurogenesis
is therefore dependent on developmental priming of chromatin that is established after
the exit from pluripotency. We further show that the barrier to activation in mESCs for
neuronal ASCL1 target genes can be partially overcome by elevating levels of histone
acetylation, or co-expressing homeodomain TFs that bias ASCL1 activity towards
neuronal targets. Together, these findings establish that ASCL1 pioneer factor activity is
constrained in an early developmental context and reframe our understanding of how
competence for somatic differentiation in response to transcriptional cues is encoded in
the epigenome.

RESULTS

ASCL1 activates distinct transcriptional profiles in pluripotent, epiblast-like and neuroectodermal cells

Our previous work has shown that naïve mouse embryonic stem cells (mESCs) do not respond to ASCL1 overexpression by activating a neurogenic programme.[44] Here we set out to understand why ASCL1 fails to drive neuronal differentiation in mESCs and investigate what, if any, is the transcriptional response to ASCL1 in mESCs and epiblast-like cells (EpiLC) compared to the response in neuroectodermal (NE) cells. To this end, we developed an mESC-based system with inducible ASCL1 activation (iASCL1 mESCs). HA-tagged ASCL1 was fused with the ligand-binding domain of the glucocorticoid receptor (LBD-GR),[41,46,47] enabling conditional control by the addition of dexamethasone (Dex) to the culture medium (Fig. 1A and Fig. S1B).[48,49] The inducible ASCL1-GR transgene (iASCL1), driven by a constitutively active promoter, was inserted at the *Rosa26* locus (see Methods) (Fig. 1A). Constitutive expression of iASCL1 was confirmed in iASCL1 mESCs (Fig. S1C). iASCL1 mESCs can be propagated in standard 2iLIF pluripotency conditions,[50] where cells express pluripotency markers, NANOG and OCT4, in the absence of Dex (Fig. S1A).

Culture of mESCs in N2B27 media without 2iLIF spontaneously gives rise to anterior neural ectoderm cells (NE) (Fig. S1D).[51,52] Expression of SOX1, an early marker of NE,[53,54] was detected at low levels from days 2 and 3 and at high levels throughout cultured cells from day 4 (Fig. S1D–E). This was concomitant with a loss of pluripotency marker OCT4 between days 3 and 4 (Fig. S1D) and followed by PAX6 and sparse TUBB3 expression from day 5, indicating spontaneous but limited neuronal differentiation from NE by this later stage (Fig. S1D). NE cells at this stage do not express endogenous ASCL1.[55] mESC to NE differentiation proceeded via a NANOG-negative, OTX2-positive state resembling EpiLCs in the transition from naïve to formative pluripotency,[5,56] but this identity was transient and was lost by day 2 (Fig. S1E). The EpiLC identity can be stabilised with activin A and FGF2,[3,5] enabling interrogation of the formative state without spontaneous differentiation (Fig.1B–C and Fig.S1F). Together, mESCs, EpiLCs and NE cells constitute an *in vitro* model of the transition from pluripotency through to neuroectoderm after gastrulation (Fig.1B & C).

We next characterised the response to iASCL1 across these cell types. In NE cells, activation of iASCL1 produced limited TUBB3-positive cells within 24 hours (Fig. 1D) but by 48 hours, neuronal differentiation was widespread and TUBB3 expression was robust compared to the sparse, spontaneous neuronal differentiation observed in NE cells without activated iASCL1 (Fig. 1D). EpiLCs displayed sparse TUBB3 expression only after 48 hours of activated iASCL1, with morphology that indicates less advanced neuronal differentiation (Fig. 1D). As expected,[44] mESCs failed to upregulate TUBB3 with or without iASCL1 over this period (Fig. 1D). This confirmed that competence for differentiation in response to ASCL1 activity is acquired during the exit from pluripotency. To exclude the possibility that resistance to iASCL1 in mESCs is attributable simply to inhibitors present in the media, we also assessed the response to iASCL1 in variants of 2iLIF lacking either

the ERK inhibitor (Chiron-LIF) or the GSK3β inhibitor (PD-LIF), as well as in serum-LIF
which also maintains pluripotency of mESCs[57–59] (Fig. S1G). ASCL1-driven expression of
target genes (*Tubb3* and *Hes5*) was observed exclusively in NE cells and not in these
variant 2iLIF conditions, indicating that neither ERK inhibition nor GSK3β inhibition alone
was responsible for suppressing neuronal differentiation (Fig. S1G).

During ASCL1-mediated differentiation, neural progenitor cells undergo rapid cell
cycle exit[60–62] and dismantle the progenitor regulatory network.[60–63] We asked whether
pluripotent cells responded in a similar way, despite the failure to induce neuronal
differentiation. We found that 24 hours of iASCL1 activation in 2iLIF mESCs resulted in a
significant reduction in the number of NANOG- and OCT4-positive mESCs (from 90–
100% to 55–60%, $p < 0.05$) (Fig. 1E) and a decrease in the number of EdU-positive cycling
cells (from 65% to 40%) (Fig. S1H). This is consistent with previous reports of ASCL1
activity accelerating loss of pluripotency genes in differentiating mESCs.[7] These results
indicate biological activity of iASCL1 in mESCs and suggest that the failure of mESCs to
differentiate into neurons in response to ASCL1 is not caused by intrinsic resistance to
ASCL1-mediated transcriptional activation.

To investigate the transcriptional response to iASCL1 genome-wide in mESC,
EpiLC and NE, we performed RNA-seq in cells exposed to 0, 6 or 24 hours of iASCL1
activity before collection (Fig. 1B and 1F). ASCL1 altered the expression of numerous
genes in all three cell types (Fig. 1G and Fig. S1I–K) demonstrating its transcriptional
activity in all three contexts. Moreover, the greatest number of upregulated differentially
expressed genes (DEGs) after 24 hours of ASCL1 activation was seen in mESCs (mESC:
780; EpiLC: 310; NE: 419) (Fig. 1G). Fewer genes were downregulated than upregulated
in all cell types (Fig. 1G), consistent with ASCL1's established predominant role as an
activator rather than a repressor.[41] Subsequent analysis therefore focused on
upregulated DEGs. A set of 63 "universal" ASCL1-responsive DEGs upregulated in all
three contexts was identified, including known ASCL1- and Notch-related targets, *Dll3*,
Id1, and *Id3*. However, the majority of ASCL1-upregulated DEGs after 24 hours were cell
type-specific, including 235 genes that were only upregulated by ASCL1 in NE, 114 in
EpiLC and 601 in mESC (Fig. 1H–I).

To better understand the functional identity of genes activated by ASCL1 in each
cell type, we performed gene ontology (GO) analysis on upregulated DEGs from mESCs,
EpiLCs and NE cells (Fig. 1J–L). iASCL1-responsive DEGs in NE were associated with
neurogenesis-related terms including "forebrain development" and "regulation of neuron
differentiation" (Fig. 1L), whereas ASCL1-responsive genes in mESC and EpiLC, were
associated with terms including "placenta development", "hemopoiesis", "muscle cell
differentiation", and "mesenchyme development" (Fig. 1J–K), indicating activation of
divergent programmes not associated with neurogenesis. Notably, ASCL1-activated
genes associated with "central nervous system neuron differentiation" were upregulated
in EpiLCs, yet these cells displayed only limited TUBB3-expression (Fig. 1D and 1K). This
indicates that at least some of the 235 NE-only ASCL1 responsive genes must be
essential for neuronal differentiation; their resistance to activation in response to ASCL1

in mESC and EpiLC likely contributes to the limited competence of these cells to undergo
ASCL1-mediated neuronal differentiation.

The list of DEGs identified after 24 hours of iASCL1 activity will also include targets
of secondary effectors in addition to genes that are directly targeted by ASCL1. It is
possible that ASCL1 directly activates the same target genes in all cell types, and
subsequent transcriptional responses diverge due to downstream effects. One
advantage of our iASCL1 system is the ability to capture early transcriptional changes;
we therefore also identified ASCL1-responsive DEGs after 6 hours of iASCL1 activation in
each cell type (Fig. S1L–O). As expected, fewer DEGs were identified after 6 hours than at
24 hours, but ASCL1-responsive DEGs were already cell type-specific at this earlier
timepoint (Fig. S1L) and GO analysis of these 6-hour DEGs also revealed that neuronal-
associated terms were only highly represented in NE (Fig. S1M–O). Together, these data
support a model for distinct ASCL1 activity in these three cell types that is largely shaped
by differences in the initial cellular context, rather than as a result of downstream
divergent responses to ASCL1 activation.

ASCL1 binding is cell-type specific and explains divergent transcriptional responses

To activate its target genes, ASCL1 must bind to *cis*-regulatory elements (CREs)
and initiate transcription. ASCL1 has been described as an on-target pioneer TF, and
previous reports have claimed that its binding sites are similar in both developmental and
reprogramming contexts.[38,39] We therefore asked whether ASCL1 binds distinct loci in
mESC, EpiLC and NE and whether this explains the divergent transcriptional responses
we observed. We used ChIP-seq to probe genome-wide ASCL1 binding (Fig. S2A–C) and
focused our analysis on the top 10,000 peaks from each cell type (Fig. S2D). This resulted
in a total consensus set of 17,255 unique ASCL1 binding sites bound in at least one of
mESC, EpiLC or NE (Fig.2A). Grouping these peaks revealed striking cell type-specific
binding patterns for ASCL1 (Fig. 2A). EpiLCs and NE have more ASCL1-binding sites in
common (3,181) than either mESC and EpiLC (1,353) or mESC and NE (402). A further
3,972 ASCL1-binding sites are shared in all three cell types (Fig. 2A). Distinguishing
ASCL1 binding sites at promoters, introns, or distal intergenic regions revealed similar
proportions in mESCs, EpiLCs and NE (Fig. S2E). Motif enrichment analysis across all
ASCL1 site clusters confirmed that binding sites in all three cell types share a similar but
not identical central E-box motif (CAGCTG), rather than harbouring cell type-specific E-
box variants (Fig. S2F). Preferential binding of bHLH TFs to E-box variants can be altered
by formation of distinct dimers,[64] but the similarity of motifs discovered in our ASCL1
ChIP-seq suggests that this is not a primary determinant of cell type-specific ASCL1
activity in mESC, EpiLC and NE.

We next asked whether these divergent binding profiles of ASCL1 could explain
the divergent transcriptional profiles. We therefore integrated our ChIP-seq and RNA-seq
data to determine the relationship between ASCL1 binding and ASCL1-responsive
differentially expressed genes, identifying all ASCL1 binding sites within 100 kb upstream
or downstream of gene transcription start sites.[65,66] We found that a large majority of the
most upregulated ASCL1-responsive DEGs in each cell type were proximal to at least one

ASCL1 binding site (mESC: 685/780; EpiLC: 291/310; NE: 385/419) (Fig. S2G).
Furthermore, differentially expressed neuronal genes that are only expressed in NE were
associated with NE-specific ASCL1 binding sites, while non-neuronal mESC-specific
ASCL1-responsive DEGs (Fig. 2A, labelled genes) were associated with mESC-specific
binding sites.

To confirm the *in vivo* neuronal activity of the NE-specific ASCL1 regulatory
elements, we compared these ASCL1 binding sites to the location of validated enhancers
from the VISTA database.[67] Gene expression of NE-specific DEGs and their associated
NE-specific ASCL1 binding sites is consistent with *in vivo* activity of enhancers
associated with these sites in the developing mouse forebrain, hindbrain and spinal cord
(Fig. 2B–B'' and Fig. 2C–C''). These data indicate that ectopic ASCL1 in NE is recruited to
physiologically relevant ASCL1 binding sites, even before endogenous ASCL1 has been
activated.

NE cells undergo a strong upregulation of the neuronal programme on ASCL1
activation (Fig. 1D and L). To understand more about the mechanisms that underpin
ASCL1-mediated neurogenesis, we next focused more closely on the 235 NE-specific
ASCL1-responsive DEGs (Fig. 1H) and found that 208 of them were associated with an
ASCL1 binding site within ± 100kb. As some of these genes were linked to multiple ASCL1
binding sites, this corresponded to 653 ASCL1 binding sites proximal to the 208 bound
and NE-specific target genes (Fig. 2D). Since ASCL1 failed to activate neurogenesis in
mESCs and EpiLCs, we then asked whether these 653 binding sites were also bound by
ASCL1 in mESCs or EpiLCs or not. Surprisingly, 455 of these sites associated with genes
only upregulated in NE were also bound by ASCL1 in either mESC or EpiLC, while only 198
of these sites were exclusively bound by ASCL1 in NE (Fig. 2D'). However, where ASCL1
was found to be bound to specific sites in two or more cell types, average ChIP-seq signal
at the shared sites was significantly higher in NE than in mESC and EpiLC (Fig. 2E). Taken
together, the ChIP-seq data suggest that ASCL1 binding, although present at some of the
correct neuronal target genes, is lower in mESC and EpiLC and binding at non-neuronal
sites is favoured. This may underlie the reduced competence of mESC and EpiLC to
respond to ASCL1 by executing a neuronal differentiation programme.

Context-dependent binding has been reported for various TFs, including pioneer
factors,[68] but ASCL1 has been described as targeting the same sites in development and
reprogramming.[38] To further explore context-dependent ASCL1 activity, we curated a
range of published ASCL1 ChIP-seq datasets from *in vitro* neural stem cells (NSCs),[38,41]
ex vivo adult neural progenitors,[69] mouse embryonic fibroblasts (MEFs),[38,70] mESCs,[45] and
embryonic neural tube[71] (Fig. S2H) and compared with our ASCL1 binding sites from
mESC, EpiLC and NE. Correlation analysis comparing the similarity of ASCL1 binding
sites across these datasets revealed that ASCL1 indeed displays cell type-specific
occupancy of binding sites, even across cell types that undergo ASCL1-mediated
neuronal differentiation (e.g. NSCs and MEFs). Looking more closely at strength of
binding at cell type-specific peaks, ASCL1 binding in our ASCL1 NE ChIP-seq was higher
than in mESC or EpiLC at ASCL1 binding sites that are also found in NSC and embryonic

neural tube from published datasets[41,71] (Fig. S2I). This analysis suggests that the pioneer
model of ASCL1 activity where ASCL1 accesses the same sites in a variety of chromatin
states across cell types does not extend to explaining binding differences across cell
types in very early development.[38] Instead, the data indicate that its binding is context-
dependent and that competence for ASCL1-mediated neuronal differentiation is
underpinned by binding at specific sets of neurogenic regulatory elements that is
restricted in pluripotent cells.

Cell type-specific chromatin pre-patterning correlates with distinct binding patterns

Having established that ASCL1 occupies distinct binding sites in mESCs, EpiLCs,
and NE, we next investigated chromatin features that may explain these differences.
Pluripotent cells are characterised by having globally low levels of repressive
heterochromatin,[9,72] so it was striking that ASCL1 fails to bind its NE-specific sites in
mESCs. We sought to better understand whether chromatin accessibility might explain
distinct ASCL1 binding profiles. To examine ASCL1 binding site accessibility across cell
types, we performed ATAC-seq on mESCs, EpiLCs, and NE that had been exposed to
either 6 or 24 hours of iASCL1 activity before collection (Fig. 3A–B and Fig. S3A–D). We
also took untreated samples at the same final time point (No ASCL1) (Fig. 3A–C) to
measure chromatin accessibility that is independent of ASCL1 activity. Surprisingly,
given ASCL1's affinity for nucleosomal DNA,[42] we found that 80–87% of ASCL1 binding
sites were accessible in each cell type even in the absence of iASCL1 activity (Fig. S3E–
G). Moreover, differences in chromatin accessibility between cell types in the absence of
ASCL1 closely mirrored the cell type-specific ASCL1 binding patterns described above
(Fig. 3C). For example, mESC-only ASCL1 binding sites are more likely to be accessible
in mESC than those same sites in EpiLC or NE in the absence of ASCL1 activation, with a
similar pattern seen for other cell type-specific binding sites (Fig. 3C and Fig. S3H). This
indicates preferential recruitment of ASCL1 to binding sites that are already accessible
and that these vary between cell types.

We focused our analysis on the previously defined 653 ASCL1 binding sites
associated with NE-specific ASCL1 target genes and asked whether these sites were
open or closed in NE in the absence of iASCL1 activation. Consistent with the global
pattern described above, 493 (75%) of these 653 ASCL1 binding sites were open in NE
even without iASCL1 activity (Fig. 3D and 3F). Therefore, most of the binding sites linked
to NE-specific target genes are open in NE independently of ASCL1 (Fig. 3G and Fig. S3I).
The remaining 160 (25%) of these 653 ASCL1 binding sites were inaccessible in the
absence of ASCL1 (Fig. 3D and 3F). After 24 hours of iASCL1 activity, 598 (92%) of the 653
sites were open, indicating 105 of 160 initially closed sites had been rendered accessible
after 24 hours of ASCL1 activity (Fig. 3E–3F and 3H). Therefore, in NE the majority of
binding sites associated with NE-specific gene expression were open independently of
ASCL1 activity, while additional further sites were pioneered after ASCL1 activation (Fig.
3E–F and 3H).

Given this strong preference for ASCL1 to bind to accessible chromatin in NE, we
asked whether the 653 ASCL1 binding sites linked to NE-specific gene activation were or

were not accessible in mESCs. Surprisingly, we found that 55% of these sites were also
accessible in mESCs in the absence of ASCL1 (Fig. 4A–B). Since the genes associated
with these open sites are not responsive to ASCL1 in mESCs, chromatin accessibility
alone cannot fully account for the divergent transcriptional response to ASCL1 at these
loci. We therefore asked why transcriptional activation of these genes fails in mESC
despite the fact that many of the ASCL1 binding sites associated with them are open and
bound by ASCL1 (Fig. 4A–B). We reasoned that histone modifications might provide an
additional layer of gene regulation.

Histone H3 acetylation at lysine 27 (H3K27ac) and methylation of lysine 4
(H3K4me1) both correlate with transcriptional activity of enhancers.[73] Moreover, mESCs
have been shown to maintain enhancers associated with differentiation in a poised state,
marked by accessible, H3K4me1-rich chromatin but lacking H3K27ac.[15,17] Poised
enhancers are not active, but respond rapidly at the onset of differentiation.[16,17,74] To see
whether differential enrichment of these marks at ASCL1 binding sites in mESC and NE
might explain the cell type-specific patterns of ASCL1 target gene activation, we analysed
published histone ChIP-seq data[17,75] from 2iLIF mESCs, EpiLCs and anterior neural
precursor cells (AntNPCs) (Fig. S4A–D). AntNPCs are approximately equivalent to NE
cells and they have also not yet upregulated endogenous ASCL1 expression.[17,55] Globally,
we found that cell type-specific ASCL1 binding correlated with H3K27ac (Fig. S4B) and
H3K4me1 marks (Fig. S4C), consistent with preferential recruitment of ASCL1 to active
chromatin. Notably, there was no enrichment of H3K4me1 at NE-specific ASCL1 binding
sites in mESC or EpiLC, indicating that these sites are not maintained in a poised state in
these cells (Fig. S4C').

We then looked specifically at the 362 ASCL1 binding sites that were open in
mESCs, despite only being associated with ASCL1 transcriptional activity in NE (Fig. 4A–
B). Of these 362 open ASCL1 binding sites, 176 were not bound in mESCs, despite being
accessible (Fig. 4C). We saw that these sites displayed low levels of H3K27ac deposition
in mESC (Fig. 4D), consistent with their maintenance in inactive chromatin. On the other
hand, these sites displayed high levels of deposited H3K27ac in NE (AntNPC) (Fig. 4D).
This indicates that these sites are not bound or activated by ASCL1, despite their
accessibility, because they are not marked with active H3K27ac. The remaining 186 sites
were bound in mESC and open in mESC, even though they also displayed NE-specific
transcriptional upregulation of their associated genes in response to ASCL1 (Fig. 4C'). We
saw that H3K27ac was lower at these sites in ESC than in NE (AntNPC), indicating that
lower levels of pre-existing H3K27ac marks contribute to a failure of ASCL1 to activate
associated NE-specific genes in ESCs, even when ASCL1 is bound at these sites in both
cell types. Example NE-specific ASCL1 target genes (Fig. 4E) and H3K27ac marks at their
associated binding sites (Fig. 4F–G) highlight this pattern. For example, *Ank2* is bound
and accessible in mESC and NE, but displays a clear enrichment of H3K27ac at its ASCL1
binding site in NE (AntNPC) compared to mESC (Fig. 4F). By contrast, *Cyth1* and *Hes5* are
examples of genes that are accessible but not bound in mESC, and these sites show low
levels of H3K27ac in mESCs (Fig. 4G). These findings indicate that the NE-specific ASCL1

target genes are not activated in mESCs because they are either inaccessible to ASCL1
or accessible but not marked with active H3K27ac.

362   Modulating histone acetylation enhances ASCL1 transcriptional activity, but does
363   not facilitate neuronal differentiation in mESCs

Our epigenetic data suggest that pre-existing chromatin status, particularly
histone acetylation and chromatin accessibility, is the primary determinant of ASCL1
binding and cellular competence for ASCL1-mediated neuronal gene activation. We
therefore asked whether artificially enhancing histone acetylation could increase the
competence of mESCs to respond to iASCL1 (Fig. 5A). We treated mESCs and NE cells
with 10 nM trichostatin A (TSA), a class I/II histone deacetylase (HDAC) inhibitor that
broadly elevates H3K27 acetylation, and profiled the transcriptional response to iASCL1
activation by RNA-seq (Fig. 5B–C and Fig. S5A–B).

TSA treatment altered the transcriptome of both mESCs and NE cells with and
without iASCL1 (Fig. 5C and S5B–F). To assess whether TSA could alter the
responsiveness of NE-specific ASCL1 target genes in mESCs, we focused on this
previously defined set of genes (Fig. 2D). Several of these genes have established roles in
neurogenesis, including *Nkx6-1*, *Disp2*, and *Plxna2* (Fig. 5D–D''), and all three of these
were now upregulated by iASCL1 in mESCs when co-treated with TSA (Fig. 5D–D''). More
broadly, combined iASCL1+TSA treatment drove the highest expression of NE-specific
ASCL1 target genes in mESCs compared to iASCL1 or TSA alone, with 116 genes
significantly upregulated relative to the control in this condition (Fig. 5E–E').

However, TSA alone had a substantial independent effect on the mESC
transcriptome compared to untreated cells, upregulating 4,924 genes including 102 of
the previously identified NE-specific iASCL1 targets (Fig. S5D and Fig. 5E'), suggesting
that many of these changes reflect opportunistic activation by other factors in response
to broadly increased acetylation rather than directed ASCL1 activity.

Beyond analysis of the previously identified NE-specific targets described above,
broader transcriptional analysis of differentially expressed genes in mESCs revealed
many additional genes that responded to combined iASCL1+TSA treatment relative to
TSA alone (n = 458) (Fig. 5F and S5F). GO analysis of these genes returned terms including
"placenta development" and "mesenchyme development" as the top hits, alongside
"regulation of neurogenesis" (Fig. 5F'), indicating that co-treatment activates a largely
incoherent and generally non-neuronal transcriptional programme. Consistent with this,
co-treatment with iASCL1 and TSA failed to produce TUBB3-positive neurons within 48
hours (Fig. 5G). This is likely because global changes to the acetylation landscape caused
by TSA treatment expose additional sites primed for ASCL1 binding at non-neuronal
targets, generating conflicting alternative transcriptional programmes. Taken together,
these results show that globally increasing histone acetylation can sensitise individual
NE-specific target genes to ASCL1 in mESCs but is insufficient to redirect ASCL1 activity
towards a productive neuronal differentiation programme and away from divergent
pathways.

Homeodomain motifs are enriched at ASCL1 binding sites

We next asked whether we could use more targeted approaches to engage a neuronal ASCL1 programme in mESCs, which do not have an epigenetic landscape that favours directed neuronal differentiation. We reasoned that, despite the availability of open neuronal gene-associated ASCL1 binding sites, ASCL1 activation of neuronal target genes may be limited in mESCs because essential co-binding TFs that normally guide ASCL1 activity to these sites in an NE context are absent from mESCs. To identify cofactors capable of enhancing ASCL1 activity at genomic sites associated with neuronal genes, we examined potential cofactor motif enrichment in NE-specific ASCL1 binding sites compared against the set of universally bound sites (Fig. 6A), prioritising motifs associated with transcription factors found at higher levels in NE relative to mESCs before ASCL1 activation (Fig. 6A). We confirmed that these motifs were also enriched in the 653 sites associated with NE-specific iASCL1 target genes (Fig. S6B). This analysis identified Sox (HMG) and Homeodomain-containing TFs as the dominant motif families at NE-specific ASCL1 binding sites. In contrast, mESC-specific ASCL1 binding sites were enriched for motifs associated with nuclear receptor, zinc finger and AP2 TFs (Fig. S6A), indicating distinct cofactor environments in mESCs and NE.

SOX4 and SOX11 are known to act downstream of bHLH proteins in neuronal development,[76,77] while SOX1 is expressed in neuroectoderm, and SOX2 has known functions in both pluripotency and neural stem cells. Top candidate homeodomain cofactors identified in our analysis included PHOX2B, PHOX2A, and DLX3 (Fig. 6A and Fig. S6B). DLX3 has been reported to potentiate ASCL1 reprogramming of fibroblasts.[39] We therefore decided to focus on relatively unexplored homeodomain-containing transcription factors as potential ASCL1 cofactors in potentiating context-dependent coherent neuronal programming. We also included LMX1A as a representative homeodomain (HD) TF with a known role in neural development[78] that was not enriched in the motif analysis to test whether any functional effects we saw were more generalisable to other homeodomain-containing TFs.

Homeodomain TFs potentiate ASCL1-driven neuronal differentiation in mESCs

To investigate whether ASCL1 and HD TFs act cooperatively to drive neurogenesis, we used a doxycycline-inducible piggyBac expression system[79] to conditionally overexpress four candidate cofactors alone or together with iASCL1 in mESCs and NE cells (Fig. S6C–E). We first asked whether co-expression could activate previously unresponsive ASCL1 neuronal target genes in mESCs after 24 hours, focusing on *Zbtb18* and *Hdac11*, which are both associated with a proximal ASCL1 binding site containing a homeodomain TF motif. Co-expression of PHOX2A, PHOX2B, or DLX3 with ASCL1 increased expression of both genes in NE cells and further potentiated the effect of ASCL1 alone (Fig. 6D–E and Fig. S6F), while LMX1A showed less consistent effects across targets (Fig. S6G). Critically, although neither *Zbtb18* nor *Hdac11* responds to iASCL1 alone in mESCs, co-expression with PHOX2A, PHOX2B, or DLX3 was sufficient to upregulate both genes. Co-activation also increased expression of *Map2*, a downstream neuronal differentiation marker and NE-specific iASCL1 target gene, in mESCs (Fig. 6D–

E and Fig. S6–G). These results demonstrate that selected HD TFs can enhance ASCL1
responsiveness at individual neuronal target genes in otherwise resistant mESCs.

To investigate how co-expression of these cofactors with ASCL1 affects
transcriptional targets genome-wide in mESCs, we performed bulk RNA-seq after 24
hours on mESCs overexpressing PHOX2B with or without iASCL1 (Fig. 7A–B and Fig. S7A).
Firstly, we saw that the number of upregulated genes in iASCL1+PHOX2B mESCs (n = 757)
was greater than that observed in iASCL1 alone (n = 480) or PHOX2B alone (n = 240),
indicating a compound effect of both factors together (Fig. S7A). We then looked at how
the iASCL1+PHOX2B upregulated genes behaved across conditions (Fig. 7A). We saw
that these genes fit broadly into two clusters: those that have highest expression in the
iASCL1+PHOX2B-treated mESCs (Fig. 7A, Cluster 1) and those that have comparable
expression in mESCs treated with either iASCL1 alone or iASCL1+PHOX2B (Fig. 7A,
Cluster 2). This indicates that the combined activity of these two transcription factors
creates a synergistic new set of target genes that are otherwise silent. GO analysis on
both Cluster 1 (Fig. 7A') and Cluster 2 (Fig. 7A'') identified "regulation of neurogenesis",
but "axon guidance" was only enriched in Cluster 1. This could indicate that the genes
upregulated by combined activity of iASCL1 and PHOX2B are associated with a more
mature or coherent neuronal transcriptional programme.

ASCL1 activation in ESCs results in upregulation of divergent gene programmes
that may actively interfere with coherent neuronal programming. We therefore asked
whether previously identified ASCL1 target genes upregulated specifically in mESCs (Fig.
1H–I) displayed an altered response to ASCL1 when it was co-expressed with PHOX2B.
(Only 497 of the previously defined 601 mESC-specific iASCL1 targets were detected in
this RNA-seq experiment due to differences in library preparation and sequencing depth.)
A subset of these genes (178 out of 497) showed less activation in mESCs when iASCL1
activation was combined with PHOX2B activation (Fig. 7B) than with iASCL1 alone, and
these genes were associated with terms such as "regulation of hemopoiesis" and "T cell
proliferation" (Fig. 7B'). Therefore, PHOX2B may work with ASCL1 both to facilitate the
activation of neuronal target genes and also to suppress the activation of non-neuronal
targets that interferes with successful differentiation in mESCs.

Finally, we assessed whether these transcriptional changes we observed in
mESCs in response to iASCL1 and HD TFs translated into neuronal differentiation,
assayed by TUBB3 expression and cell morphology after a longer 72-hour period
compared to earlier analysis of more immediate iASCL1 effects (Fig. 1–5). In NE, which
already express some endogenous PHOX2B and PHOX2A (Fig. 6A), additional ectopic
PHOX2B and PHOX2A both induced a modest increase in neurogenesis compared to
control. However, neither PHOX2B nor PHOX2A co-expression with iASCL1 significantly
increased the proportion of TUBB3-positive cells relative to iASCL1 alone (PHOX2B: 3.0%
versus 7.7%, p > 0.05; PHOX2A: 9.2% versus 12.9%, p > 0.05) (Fig. 7D and 7G and Fig.
S7C–D). DLX3 and LMX1A did not increase TUBB3-positive cell numbers in NE even when
expressed alone or with iASCL1 (Fig. S7E–F), highlighting differences in the intrinsic
neurogenic capacity of individual HD TFs.

In mESCs, iASCL1 activation alone produced only sparse, diffuse TUBB3
expression after 3 days (Fig. 7C). PHOX2B and PHOX2A expression did not result in
TUBB3-positive cells in the absence of iASCL1 (Fig. 7C and 7F; Fig. S7B and S7D).
However, strikingly, co-expression of PHOX2B or PHOX2A with iASCL1 produced TUBB3-
positive cells with immature neuronal morphology (Fig. 7C and 7E–F; S7B and S7D). DLX3
and LMX1A failed to generate TUBB3-positive cells in mESCs with or without iASCL1 (Fig.
S7E–F), indicating that the ability to confer neuronal competence on ASCL1 in mESCs is
specific to PHOX2A and PHOX2B and not a general feature of homeodomain-containing
TFs.

DISCUSSION

Together, these data support a model in which ASCL1 is not an intrinsically
neuron-specific pioneer factor, but a highly context-dependent transcription factor
whose activity is constrained by both chromatin state and cofactor availability. Cell-
intrinsic changes during the exit from pluripotency are required for competence for
ASCL1-mediated neuronal differentiation and to channel ASCL1 activity away from
activating divergent programmes. However, this neuronal competence can be increased
by altering the global chromatin landscape or by providing additional cofactor TFs that
enhance neuronal target gene activation.

Many studies have used mESCs (and human iPSCs) for forward programming and
TF-mediated neuronal differentiation. However, it is important to note that many of them
incorporate embryoid body formation, or some other priming step.[7,80] Vainorius et al. use
ASCL1 to drive neuronal differentiation from mESCs, but with two important differences:
firstly, they cultured their mESCs in serum-LIF, in which ESCs are heterogeneous and
display epigenetic and transcriptional priming of many lineage commitment genes;[56,81]
secondly, cells were plated on Matrigel and differentiated in N2B27 after doxycycline
induction of ASCL1 expression. Thus, these cells have likely already transitioned through
formative pluripotency and are competent for ASCL1-mediated differentiation. Casey et
al. profiled genome-wide binding of ASCL1 in mESCs grown on feeder cells, and observed
a similar binding profile of ASCL1 to our mESC data, but did not investigate competence
for neuronal differentiation or the expression of non-neuronal genes.[45] We find that while
failing to upregulate a coherent neuronal programme, pluripotent mESCs are
nevertheless fully competent to respond transcriptionally to ectopic ASCL1, but in this
context non-neuronal heterologous gene programmes dominate; exit from pluripotency
is required to allow an appropriate ASCL1-specific developmental programme to be
engaged. This underscores the importance of chromatin remodelling to restrict the cell
identity during the early stages of embryonic development.[75,82,83]

Despite the fact that pluripotent cells have the ability to differentiate into any
embryonic cell type and are characterised by having globally low levels of repressive
heterochromatin,[9,72] it is perhaps surprising to find that chromatin accessibility played a
major role in determining ASCL1 binding in ESCs. Rather than a model with widespread
chromatin accessibility in pluripotent cells offering a "blank canvas" for highly permissive
binding of transcription factors, with an epigenetic environment progressively refined
during development, our data point to complex but distinct landscapes. Unlike
accessible sites bound by ASCL1 in NEs, sites accessible to ASCL1 binding in ESCs are
generally not associated with neuronal genes. Moreover, in the context of gene activation
in ESCs, EpiLCs and NEs, we see a limited ability of ASCL1 to act as a pioneer opening
chromatin. Recent data in human iPSC differentiation to neurons, has shown ASCL1
displays different modes of activity at different sites across the genome, sometimes
acting alone and sometimes recruiting SWI/SNF chromatin remodelling complexes to act
as a non-classical pioneer factor.[32] Our study adds to this model, showing that ASCL1
acts as both a pioneer and a non-pioneer TF to varying degrees across the genome and

between cell types. Moreover, there must be additional layers of regulation that determine whether a given binding site is accessible to ASCL1-mediated pioneering. Differential availability of chromatin remodelling complexes across the developmental time course may determine whether a given site can be pioneered by ASCL1, as has been described for SWI/SNF subunit switching in neuronal differentiation.[32,84]

Accessibility, though clearly important, is not the only factor that determines context-dependent ASCL1 binding and gene activation. Many ASCL1 binding sites near neuroectoderm-specific target genes were marked by H3K27ac in neural progenitors but not in mESCs or EpiLCs, even when accessibility is comparable at these sites across cell types. This is especially interesting in light of the prevailing view of mESCs being broadly unrestricted in terms of their epigenetic landscape and developmental potential. Instead, ASCL1 binding sites that are activated in NE are maintained in inactive chromatin in mESC and EpiLC and are not poised or primed. This likely reflects the fact that they are not the first regulatory elements recruited *in vivo* in the exit from pluripotency, as ASCL1 itself is expressed at later stages of embryo development.[85,86] Consistent with this, global elevation of histone acetylation using an HDAC inhibitor renders many previously unresponsive neuronal genes inducible by ASCL1 in mESCs, highlighting the importance of chromatin priming in establishing developmental competence. Even transient treatment with HDAC inhibition can have significant changes on genome architecture in mESCs,[87] and our transcriptomic data indicates that TSA treatment creates many additional non-neuronal ASCL1 target genes. HDAC inhibitors have been used to potentiate ASCL1-mediated reprogramming in the adult retina,[88] but it is possible that additional epigenetic constraints in these tissues limit the ectopic effects of induced ASCL1 activity.

To identify more targeted approaches to modulating ASCL1 activity in mESCs, we identified candidate co-factors by identifying specific motifs for neuroepithelial transcription factors found near to ASCL1 binding sites in NE target genes. We found that co-expression of ASCL1 with candidate homeobox TFs in mESCs allows activation of many previously NE-specific neuronal genes while simultaneously inactivating genes associated with divergent pathways. Interaction between HD TFs and bHLH factors, including ASCL1, has been previously shown in the diversification of cell types in the developing retina[89] cranial neural crest cells.[90] PHOX2A and PHOX2B have been implicated in gene regulatory networks with ASCL1 in the development of the central nervous system,[91–93] sympathetic neurons,[94,95] and in neuroblastoma.[96,97] ASCL1 expression typically precedes and is required for PHOX2 factor expression in these cases, so it is notable that we show simultaneous expression of both factors potentiated the direct programming capacity of ASCL1 in mESCs, where PHOX2 factors cannot be upregulated by ASCL1 activation directly. This suggests that the sequential logic of endogenous development can be bypassed by simultaneous cofactor delivery.

In summary, we demonstrate that despite their ability to differentiate into any embryonic cell type, the competence of ES cells to respond to the lineage-determining transcription factor ASCL1 is limited by a chromatin landscape in which non-neuronal

578 enhancers are favourable for ASCL1 binding, which instead directs ASCL1 activity to
579 divergent genes and pathways. Our findings illustrate the multifaceted controls that limit
580 reprogramming in different cellular contexts and provide a framework for improving
581 directed differentiation and reprogramming approaches by targeting competence rather
582 than transcription factor expression alone.

 FIGURE LEGENDS

Figure 1 – Inducible ASCL1 drives divergent and non-neuronal transcriptional programmes in mESC and EpiLC

A) Schematic of mouse embryonic stem cells with inducible ASCL1 (iASCL1 mESC).

B) Differentiation timeline and sample collection for IF (D–E) and RNA-seq (F–L). "+ ASCL1" indicates 5 µM dexamethasone treatment.

C) Phase-contrast images of mESC, EpiLC, and NE at day 2 without ASCL1, with schematic of in vivo approximates.

D) IF showing mESC, EpiLC, and NE ± iASCL1 for 24 h (top) or 48 h (bottom). TUBB3 (green) marks neuronal differentiation. Scale bar = 100 µm. Representative of 4 experiments.

E) IF of iASCL1 mESC ± 24 h iASCL1 stained for NANOG (yellow) and OCT4 (red) and percentage of positive cells (n=3; 8–10 images per experiment). Scale bar = 100 µm. * $p < 0.05$, t-test.

F) PCA of top 1000 variable genes across conditions; grey points denote Dex-treated control line with no transgene.

G) Number of differentially expressed genes (DEGs) at 6 h and 24 h relative to 0 h in each cell type (mESC blue, EpiLC yellow, NE pink; positive upregulated, negative downregulated).

H) Venn diagram of overlap between ASCL1-upregulated DEGs across cell types after 24 h iASCL1.

I) Heatmap of z-scored expression for DEGs upregulated by ASCL1 in at least one cell type; clusters grouped by cell-type specificity with selected genes indicated.

J–L) Top 5 GO terms for upregulated genes after 24 h ASCL1 in mESC (J), EpiLC (K), and NE (L).

Figure 2 – Context-dependent ASCL1 genomic occupancy underlies divergent transcriptional responses across developmental states

A) Top 10,000 ASCL1 binding sites from anti-HA tag ChIP-seq from each cell type (mESC, EpiLC, NE) collated to give 17,255 unique binding sites. ASCL1 binding sites are clustered by cell type-specific binding patterns. Mean RLE-normalised ChIP-seq signal from 3 or 4 replicates.

B–C) Examples of NE-specific target genes from RNA-seq (B and C) and NE-specific ASCL1 binding sites (B' and C'), which overlap with validated enhancers from the VISTA dataset (B'' and C''). *In situ* hybridisation from VISTA database shows enhancer activity in neural tube and brain at E11.5.

D) Categorising ASCL1 binding sites linked to NE-specific genes based on whether they are exclusively bound in NE (left, n = 198) or bound in at least one other cell type (right, n = 455). (D') Venn diagram for these categories. Example IGV tracks of ASCL1 ChIP-seq for each category: *Wnt5a*, *Plxna2*, *Inppl1*.

E) ChIP-seq signal at sites in each of the categories in (D). One-way ANOVA with repeated measures, Benjamini-Hochberg corrected pairwise comparisons: **** $p < 0.0001$.

Figure 3 – Chromatin accessibility dictates ASCL1 binding, with pioneer activity restricted to a minority of neuronal target sites

A) Schematic for ATAC-seq experiment. Samples were exposed to either no ASCL1, 6 hrs, or 24 hrs before collection at the same endpoint.

B) PCA for ATAC-seq samples

C) Heatmap of ASCL1 ChIP-seq (black, left) and ATAC-seq signal with No ASCL1 (green, right) clustered by cell type-specific binding patterns defined in Fig. 2A. Row order is preserved across the two heatmaps. Heatmaps centred on ASCL1 peak ± 1 kb.

D) Heatmaps of ASCL1 ChIP-seq (black) and ATAC-seq (No ASCL1) (green) for 653 ASCL1 sites linked to NE-specific DEGs defined in Fig. 2D. Peaks are clustered by whether they are open (top, n = 563) or closed in NE (No ASCL1) (bottom, n = 90). Heatmaps centred on ASCL1 peak ± 1 kb.

E) Heatmap of ATAC-seq (24hr ASCL1) for same 653 ASCL1 sites linked to NE-specific DEGs with row order preserved from (D). Heatmaps centred on ASCL1 peak ± 1 kb.

F) Donut plots to show percentage of 653 ASCL1 binding sites in D and E that are open (green) or closed (grey) in No ASCL1 ATAC-seq (top) or 24 hr ASCL1 ATAC-seq (bottom).

G) IGV tracks of ASCL1 binding sites that are open in NE (No ASCL1) ATAC-seq.

H) IGV tracks of ASCL1 binding sites that are closed in NE (No ASCL1) ATAC-seq but become open in NE (24 hr ASCL1) ATAC-seq.

Figure 4 – Histone acetylation status at shared accessible sites predicts productive ASCL1 binding and target gene activation

A) Heatmaps of ASCL1 ChIP-seq (black) and No ASCL1 ATAC-seq (green) from mESC at the 653 ASCL1 binding sites associated with NE-specific ASCL1 target genes defined in Fig. 3D. Heatmaps centred on ASCL1 peak ± 1 kb.

B) Donut plot of percentage of the 653 sites that are open (green) or closed (grey) in mESC (No ASCL1) ATAC-seq data.

C) Heatmaps of ASCL1 ChIP-seq (black) and ATAC-seq (No ASCL1) (green) from mESC for the 362 out of 653 sites that are open in mESC (No ASCL1) ATAC-seq data. Peaks are clustered based on whether are (C') bound in mESC (bottom, n = 186) or (C) not bound in mESC (top, n = 176). Heatmaps are centred on ASCL1 peak ± 2 kb.

D) H3K27ac ChIP-seq signal at 176 ASCL1 binding sites in (C); D') H3K27ac ChIP-seq signal at 186 ASCL1 binding sites in (C'). H3K27ac data is shown for mESC (blue) and AntNPC (pink) from Cruz-Molina et al., 2017[17]. Profiles are centred around ASCL1 binding sites ± 5 kb.

E) Normalised RNA-seq counts for representative NE-specific ASCL1 target genes (*Ank2*, *Cyth1*, *Hes5*) in mESC (blue) and NE (pink) exposed to 0, 6, or 24 hours of iASCL1 activity. "C" (grey) represents Dex-treated control line with no iASCL1 transgene. N = 4.

F–G) IGV tracks to show ASCL1 binding sites near the NE-specific target genes in (D) that are bound (F) or not bound (G) by ASCL1 in mESCs. mESC H3K27ac (blue), mESC No ASCL1 ATAC-seq (green), mESC ASCL1 ChIP-seq (black); NE (AntNPC) H3K27ac (blue), NE No ASCL1 ATAC-seq (green), NE ASCL1 ChIP-seq (black). Rectangles below tracks indicate a called peak. Peaks are highlighted in pink boxes.

Figure 5 – Global enhancement of histone acetylation partially unlocks neuronal gene expression by ASCL1 but fails to redirect differentiation in mESCs

A) Schematic of HDAC inhibition (HDACi) and acetylation.

B) Schematic of culture and treatment of mESC (blue) and NE (pink) with iASCL1 and/or 10 nM TSA for 24 hr (RNA and IF) or 48 hr (IF).

C) PCA plot for RNA-seq samples with mESC (blue) and NE (pink) treated with 24 hours iASCL1 (darker) or not (lighter). Simultaneously, samples were either treated with DMSO (circle) or 10 nM TSA (triangle).

D) Examples of previously identified NE-specific ASCL1-respsonsive genes that become ASCL1-responsive in mESC with ASCL1 + TSA treatment. D) *Nkx6-1*; D') *Disp2*; D'') *Plxna2*. Adjusted p-values from DESeq2 differential gene expression analysis. ns p > 0.05, ** p < 0.01, *** p < 0.001

E) Heatmap of z-scored RNA-seq counts for genes that were identified as NE-specific ASCL1 targets in Fig. 1H–I and have associated ASCL1 binding site (n = 208). Expression data from mESC: control, iASCL1 only, TSA only, iASCL1+TSA. E') genes of the 208 in the heatmap that are significantly upregulated in the comparisons listed: iASCL1+TSA versus control (n = 116); TSA only versus control (n = 102); iASCL1+TSA versus TSA only (n = 13).

F) Heatmap of z-scored RNA-seq counts for genes that are significantly upregulated in TSA+iASCL1 versus TSA only (n = 458). F') Gene ontology analysis for all genes in heatmap (F). Top 6 GO terms are listed. "Regulation of neurogenesis" highlighted with black arrow.

G) IF images of H3K27ac (magenta) and TUBB3-positive neurites (green) after 48 hours ±ASCL1 and ±TSA in mESC or NE. DAPI counterstain (blue). Scale bar = 100 μm.

Figure 6 – Homeodomain transcription factor motifs are enriched in NE-specific ASCL1 binding sites and co-expression with ASCL1 enables activation of previously NE-specific target genes

A) Plot of TFs scored by fold enrichment of motif in NE-specific peaks (x) against log fold change in expression comparing NE versus mESC (No iASCL1) from RNA-seq (Fig. 1). Coloured by TF family. Phox2b, Phox2a, and Dlx3are highlighted as shortlisted candidates.

B) Schematic for putative model of cooperativity between ASCL1 and Homeodomain (HD) TFs.

C) Schematic for culture and treatment of mESC (blue) and NE (pink) for 24 hrs: control (light), +iASCL1 only (dark), +cofactor (Dox induction) only, +iASCL1+cofactor

D–E) qPCR of NE-specific ASCL1 DEGs associated with ASCL1 binding sites that contain HD motif (*Hdac11* and *Zbtb18*) and NE-specific marker of neurogenesis (*Map2*). mESC (blue) and NE (pink). Dox used to induce overexpression of cofactors, either PHOX2A (D) or PHOX2B (E). Control, iASCL1 alone, Dox alone, or iASCL1 and Dox. N = 4 experiments, each performed in technical duplicate. Two-way ANOVA with Dunnett's correction for multiple comparisons to the control for each cell type.

Figure 7 – Genome-wide redirection of ASCL1 transcriptional output with PHOX2B co-expression

A) Heatmap of z-scored RNA-seq counts for genes that are significantly upregulated by iASCL1+PHOX2B relative to untreated control mESCs (n = 757). Genes are clustered into 2 groups: Cluster 1 shows highest enrichment in iASCL1+PHOX2B cells (n = 403); Cluster 2 shows comparable expression in iASCL1 alone and iASCL1+PHOX2B. A') GO analysis for genes in Cluster 1. A'') GO analysis for genes in Cluster 2. Top 6 terms are shown.

B) Heatmap of z-scored RNA-seq counts for genes that were previously identified as mESC-specific iASCL1 targets (Fig. 1H) (n = 497). Genes are clustered using k means = 3. B') GO analysis of genes in Cluster 1. Top 6 terms are shown.

C–D) Schematic for culture and treatment of mESC (C, blue) and NE (D, pink) before collection for immunofluorescence (IF) imaging. Cells were either untreated, iASCL1 only, PHOX2B only, or iASCL1+PHOX2B. TUBB3 (green), mCherry (magenta), DAPI (blue). mCherry and DAPI LUTs are adjusted separately for mESC and NE, but TUBB3 is consistent across conditions. Scale bar = 100 µm. Representative images from 2 experiments each done in duplicate.

E) Magnified inset from Fig. 7C. White arrow indicates neurite-like outgrowth. Scale bar = 100 µm.

F–G) Quantification of TUBB3-positive cells in cofactor differentiation assay in mESC and NE untreated, with ASCL1 alone, Dox alone, or ASCL1 and Dox. Each dot represents an image taken from a randomised position in the well. Statistics are performed using N = 4 for 4 wells, from 2 independent passages. Wilcoxon pairwise test with Benjamini–Hochberg post-hoc correction. * $p < 0.05$

 SUPPLEMENTARY FIGURE LEGENDS

Figure S1 – Related to Figure 1

A) iASCL1 mESC express OCT4 (green) and NANOG (red) in 2iLIF. Scale bar = 50 µm.

B) qPCR of ASCL1 target genes ± 24h Dex in iASCL1 mESC in 2iLIF. N = 3 experiments. **
$p < 0.01$, t-test.

C) Western blot to show HA-ASCL1-GR is specific to iASCL1 and levels are consistent
across differentiation conditions (mESC, EpiLC, and NE). Representative of 2 western
blots. Molecular weight markers are indicated (right).

D) IF of OCT4 (yellow), SOX1 (red) and PAX6 (magenta) in iASCL1 differentiated into NE
conditions on day 3-5. Spontaneous neuronal differentiation from day 4 onwards
indicated by TUBB3 (green). Scale bar = 100 µm. Representative of 2 experiments.

E) IF of NANOG (blue), OTX2 (yellow) and SOX1 (magenta), day 0-2 of NE differentiation.

F) IF for naïve pluripotency marker, NANOG (yellow), and formative markers, OCT6 (green)
and OTX2 (magenta) in iASCL1 cells on days 1-3 of EpiLC differentiation. Scale bar = 100
µm.

G) qPCR of ASCL1 target genes in 2iLIF variant media. N = 3 independent experiments.
Two-way ANOVA with Dunnett's post-hoc pairwise comparisons comparing control
versus +iASCL1 for 24 hours.

H) EdU-positive mESCs (yellow) with or without 24 hours of iASCL1 activity.
Quantification of positive cells from 8 images from 3 independent experiments. **** $p <$
0.0001, t-test.

I–K) Volcano plots to show upregulated genes in mESC (I), EpiLC (J) and NE (K) after 24
hours of iASCL1 relative to untreated cells in each of: mESC (blue), EpiLC (yellow), NE
(pink). Note: some genes appear above threshold for significance but are coloured grey
and excluded from analysis based on comparison between untreated cells and Dex-
treated control cells with no iASCL1-GR transgene to control for Dex-specific effects.

L) Venn diagram to show overlap in DEGs after 6 hours.

M–O) Volcano plots and GO analysis on upregulated genes in mESC (M), EpiLC (N) and
NE (O) after 24 hours of iASCL1 relative to untreated cells in each of: mESC (blue),
EpiLC (yellow), NE (pink).

Figure S2 – Related to Figure 2

A) ChIP-qPCR to validate the anti-HA ASCL1 ChIP-seq. N = 4 independent experiments.

B) Correlation matrix of ChIP-seq peak sets, including all iASCL1 samples and control cells that have no iASCL1-GR transgene.

C) PCA plot to show sample distances after normalisation (using RLE method and reads in peaks as library size).

D) Heatmap across replicates after normalisation at the top 10,000 peaks in the consensus set.

E) Annotation of top 10,000 ASCL1 binding sites in each cell type. Promoter regions are defined as ±1kb from TSS.

F) Top motif discovered by STREME in each of the clusters. E-value and percentage of target sites that contain the motif are reported next to each motif.

G) Number of differentially expressed genes (DEGs) in mESC (blue), EpiLC (yellow) and NE (pink) that have a binding site within ±100 kb. DEGs with no proximal ASCL1 binding site are grey.

H) Correlation matrix for publicly available ASCL1 ChIP-seq datasets, including mESC, EpiLC and NE from this study. Full GEO accession numbers are available in the Methods section.

I) Profiles of ChIP-seq signal in mESC, EpiLC, and NE from this study at peaks from neural stem cells (NSC),[41] and embryonic neural tube.[71]

Figure S3 – Related to Fig.3

A) Number of peaks called in each sample. Asterisk corresponds to the per-sample consensus peak set, comprising peaks called in at least 3 of the 4 replicates.

B) Fraction of reads in peaks (FRiP) for all samples.

C) TSS enrichment is the ratio of reads the align to TSS of known genes over the reads in 1000 bp regions flanking the TSS.

D) Percentage of reads that align to mitochondrial DNA per sample.

E) iASCL1 ChIP-seq (black) and No iASCL1 ATAC-seq (green) data in mESC at all 10,000 ASCL1 binding sites detected in mESC, split by whether they are open (top) or closed (bottom) in the No iASCL1 ATAC-seq data.

F) iASCL1 ChIP-seq (black) and No iASCL1 ATAC-seq (green) data in EpiLC at all 10,000 ASCL1 binding sites detected in EpiLC, split by whether they are open (top) or closed (bottom) in the No iASCL1 ATAC-seq data.

G) iASCL1 ChIP-seq (black) and No iASCL1 ATAC-seq (green) data in NE at all 10,000 iASCL1 binding sites detected in NE, split by whether they are open (top) or closed (bottom) in the No iASCL1 ATAC-seq data.

H) Donut plots for percentage of iASCL1 ChIP-seq peaks in cell type-specific clusters (Fig. 2A) that overlap with an ATAC-seq peak (green) or not (grey) in each cell type.

I) RNA-seq counts for NE-specific ASCL1-responsive DEGs highlight in Fig.3. As per RNA-seq experiment, samples were either treated with 0, 6, or 24 hours of iASCL1 and compared to a control line with no iASCL1 transgene for each cell type.

Figure S4 – Related to Fig.4

A) ASCL1 ChIP-seq as in Fig. 2 and 3.

B) H3K27ac signal in mESC and AntNPC from Cruz-Molina et al., 2017[17] plotted across ASCL1 binding sites (this study) split by cell type-specific clusters (Fig. 2A). Centred around ASCL1 binding site (500 bp) ± 5 kb.

C) H3K4me1 signal in mESC and EpiLC from Bleckwehl et al., 2021[75] plotted across ASCL1 binding sites (this study) split by cell type-specific clusters (Fig. 2A). Centred around ASCL1 binding site (500 bp) ± 5 kb. C') highlights the NE-specific iASCL1 binding sites.

Figure S5 – Related to Figure 5

A) Phase contrast to show morphology and density of mESC treated with nothing, +iASCL1 only, 10 nM TSA only, or +iASCL1 and +TSA.

B) PCA plot of PC2 and PC3 from RNA-seq.

C–F) Volcano plot to show differentially expressed genes in mESC in specific comparisons: C) iASCL1 versus control; D) TSA only versus control; E) iASCL1+TSA versus control; F) iASCL1+TSA versus TSA only. Significance thresholds: absolute log fold change > 0 and adjusted p-value < 0.01. ashr shrunken log fold changes are used for plotting.[98]

Figure S6 – Related to Figure 6

A) Plot of TFs scored by fold enrichment of motif in mESC-specific peaks (x) against log fold change in expression in mESC versus NE at t = 0. Coloured by TF family.

B) Number of ASCL1 binding sites that contain a DLX3 or PHOX2B motif.

C) piggyBac construct used for Dox-inducible overexpression. ITR, inverted terminal repeat; cHS4, chicken hypersensitive site 4 insulator; TRE3G, third-generation tetracycline-responsive promoter; COFACTOR is where candidate cofactors were cloned in; T2A, self-cleaving peptide; mCherry; pA, polyadenylation signal; pEF1α promoter; rtTA Tet-On activator; PuroR, puromycin resistance gene.

D) Titration of Dox treatment on pool of antibiotic-resistant cells. mCherry (red) indicates induction of the transgene.

E) qPCR of *Phox2b* transgene induction relative to *Tbp* housekeeping. N = 4 experiments

F–G) qPCR of NE-specific ASCL1 DEGs associated with ASCL1 binding sites that contain HD motif (*Hdac11* and *Zbtb18*) or *Map2*. Dox used to induce overexpression of either DLX3 (G) or LMX1A (H). Cells were untreated or treated with either iASCL1 alone, Dox alone, or iASCL1 and Dox. N = 4 experiments, each performed in technical duplicate. Two-way ANOVA with Dunnett's correction for multiple comparisons to each cell type's respective control.

Figure S7 – Related to Figure 7

A) Volcano plots for specific comparisons in mESC from RNA-seq (Fig. 7A–B): iASCL1 versus control; PHOX2B only versus control; iASCL1+PHOX2B versus control. Significance thresholds: absolute log fold change > 1 and adjusted p-value < 0.01. ashr shrunken log fold changes are used for plotting.[98]

B–C) Schematic for culture and treatment of mESC (C, blue) and NE (D, pink) before collection for immunofluorescence (IF) imaging. Cells were either untreated, iASCL1 only, PHOX2A only, or iASCL1+PHOX2A. TUBB3 (green), mCherry (magenta), DAPI (blue). mCherry and DAPI LUTs are adjusted separately for mESC and NE, but TUBB3 is consistent across conditions. Scale bar = 100 µm. Representative images from 2 experiments each done in duplicate.

D–F) Quantification of TUBB3-positive cells in cofactor differentiation assay in mESC and NE. Cofactors are PHOX2A (E), DLX3 (F) and LMX1A (G). Each dot represents an image taken from a randomised position in the well. Statistics are performed using N = 4 for 4 wells, from 2 independent passages. Wilcoxon pairwise test with Benjamini–Hochberg post-hoc correction. * p < 0.05

ACKNOWLEDGEMENTS

We would like to thank the Cambridge Stem Cell Institute Genomics Facility at the Jeffrey Cheah Biomedical Centre for all of their help with library preparation and sequencing; the Cambridge Stem Cell Institute Imaging Core Facility; Prof. Brian Hendrich and members of his lab group for their assistance with ChIP-seq and ATAC-seq; William Beckman, Lidiya Mykhaylechko, Sarah Gillen, Frances Connor, and all members of the Philpott lab for helpful discussions throughout; and Dr Mekayla Storer, Prof. Jason Carroll and Dr Srinjan Basu for their useful guidance.

This work was supported by a Wellcome Trust Investigator award (212253/Z/18/Z) to (A.P., J.L.-B., R.D., J.-P.N.-B.); J.L.-B. is funded by a Wellcome Trust studentship (218481/Z/19/Z); R.A. is supported by core funding from UCL School of Pharmacy and by the Royal Society grant (RGS\R2\242202).

AUTHOR CONTRIBUTIONS

Conceptualisation, R.A., A.P. and J.L.-B.; Methodology, R.A., J.L.-B., and J.-P.N.-B.; Investigation, J.L.-B., R.A., R.D., and J.-P.N.-B.; Formal analysis, Visualisation, Data curation, J.L.-B.; Writing – original draft, J.L.-B.; Writing – review & editing, J.L.-B., J.-P.N.-B., R.A., and A.P.; Supervision, R.A. and A.P.; Funding acquisition, R.A. and A.P.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contacts, Dr Roberta Azzarelli roberta.azzarelli@ucl.ac.uk or Prof. Anna Philpott ap113@cam.ac.uk.

Materials availability

All unique/stable reagents and cell lines generated in this study are available from the lead contact with a completed materials transfer agreement.

Data and code availability

Bulk RNA-seq, ChIP-seq and ATAC-seq data have been deposited at GEO and are publicly available as of the date of publication. Accession numbers are listed in the key resources table. Microscopy data reported in this paper will be shared by the lead contact upon request. All original code has been deposited at GitHub and is publicly available as of the date of publication. The DOI is listed in the key resources table. Any additional information required to reanalyse the data reported in this work is available from the lead contact upon request.

METHODS

Cell culture

Mouse embryonic stem cells (mESCs; ES-E14TG2a, ATCC #CRL-1821) were maintained as described previously.[99] Ascl1-GR-HA mESCs (iASCL1; R. Azzarelli, unpublished) were cultured in 2i/LIF medium: N2B27 basal medium (1:1 Neurobasal [Gibco, 21103-049] / DMEM-F12 [Gibco, D6421], N2 [in-house, Cambridge Stem Cell Institute], B-27 [Gibco, 17504044], 2 mM L-glutamine, 50 µM β-mercaptoethanol) supplemented with 100 U/mL leukaemia inhibitory factor (LIF; in-house), 3 µM CHIR-99021 (ABCR, AB253776) and 1 µM PD0325901 (ABCR, AB253775).

Cells were passaged every 2–3 days using Accutase (300 × g, 3 min) and seeded at 1.5×10^4 cells/cm^2 on 0.1% gelatin-coated 6-well plates. Stocks were cryopreserved at 1×10^6 cells/mL in 50% 2i/LIF / 40% KSR / 10% DMSO. Mycoplasma testing was performed every 3–6 months.

For lineage transitions, mESCs were plated on fibronectin (16.7 µg/mL) to generate epiblast-like cells (EpiLC) in N2B27 + FGF2 (12 ng/mL) + Activin A (20 ng/mL),[3,5] or on laminin (10 µg/mL) for primitive neuroectoderm (NE) differentiation in N2B27 + 0.1% sodium bicarbonate + 0.11% BSA + 20 µg/mL insulin.[99]

A gene fragment encoding N-terminal HA-tagged full-length ASCL1 fused to the glucocorticoid receptor ligand-binding domain (GR-LBD), followed by an IRES-GFP cassette, was synthesised with flanking attB1/attB2 sites (Twist Bioscience). Gateway LR Clonase II (Thermo Fisher) was used to insert the construct into a Rosa26-targeting destination vector (SP202) carrying appropriate homology arms.[100] E14 mESCs were co-transfected with the targeting vector, Rosa26-directed sgRNAs, and a Cas9 expression vector using the 4D Nucleofector (Lonza). Correctly targeted clones were isolated by FACS for constitutive GFP expression, expanded, and verified by PCR across both homology arms for monoallelic (heterozygous) insertion. All experiments used a single validated clonal line.

In iASCL1 mESCs, ASCL1 activity is post-translationally controlled: the HA-ASCL1-GR fusion protein is constitutively expressed but held inactive by HSP90 chaperone binding until displacement by 5 µM dexamethasone (Dex).[101]

Cofactor overexpression mESCs

PiggyBac-based, doxycycline (Dox)-inducible overexpression vectors were assembled from two published plasmids: the TRE3G-driven expression cassette was PCR-amplified (CloneAmp, Takara, 639298) from pPB-Ins-TRE3Gp-KRAB-dCas9-ecDHFR-IRES-GFP-EF1Ap-Puro (Addgene #183410), and the rtTA-Neo cassette was amplified from pPB-Ins-U6p-sgRNAentry-EF1Ap-TetOn3G-IRES-Neo (Addgene #183411). Both were cloned into a custom piggyBac-based vector (VB240515-1632gka, VectorBuilder) via EcoRV/PacI digestion and In-Fusion Snap Assembly (Takara, 638948). The dCas9-KRAB-U6-sgRNA module was excised with NotI/AsiSI and replaced with synthesised transgene sequences (PHOX2A, PHOX2B, LMX1A, and DLX3: Twist Bioscience; mCherry: VectorBuilder). Gel

purification used the Monarch kit (NEB, T1120); plasmids were propagated in Stellar
competent cells (Takara, 636763), purified with QIAprep Miniprep or Midiprep kits
(Qiagen), and sequence-verified by Nanopore sequencing (Plasmidsaurus).

For PHOX2A, PHOX2B, LMX1A, and DLX3 overexpression lines, mESCs were co-
transfected with PB-Dox-Cofactor plasmid and Super PiggyBac Transposase vector
(Stratech, PB210PA-1-SBI; 500 ng) to a total of 2 µg DNA, complexed with 3.5 µL
Lipofectamine 2000 (Invitrogen, 11668027) in 400 µL Opti-MEM™ and incubated on cells
for 4 h at 37°C. Medium was replaced with 2i/LIF, then selection in 0.5 µg/mL puromycin
(2i/LIF) initiated 48–72 h post-transfection and maintained until colonies were pooled.
Knockdown efficiency was confirmed by RT-qPCR after 1 µM Dox induction.

Immunofluorescence microscopy

Cells were fixed in 4% PFA (ThermoFisher, J19943-K2) for 20 min at room temperature,
permeabilised in 0.2% Triton-X100 (Sigma, X100)/PBS for 15 min, and blocked in 5%
serum / 0.01% Triton-X100 / PBS for 1 h. Primary antibodies (PBS + 2% serum + 0.01%
Triton-X100) were applied overnight at 4°C; secondary antibodies for 1 h at room
temperature. Washes used PBS-T (PBS + 0.01% Triton-X100); DAPI (1 µg/mL, Abcam,
ab228549) was added during the third wash (20 min). Images were acquired on a Leica
DMI6000 at 10× or 20×.

S-phase fraction was used as a proliferation proxy. Cells were treated with 5 µM Dex for
24 h; then 10 µM EdU (final; Click-iT EdU Kit, Alexa Fluor 488, Thermo Fisher, C10637) was
added for 1 h before fixation. EdU was detected per manufacturer's instructions. Note:
iASCL1 mESC constitutive GFP was negligible relative to AlexaFluor-488-labelled targets,
so the 488 channel was used without interference.

Image analysis

Nuclear intensities were quantified in CellProfiler v4.0.[102] Nuclei were segmented from
DAPI using the IdentifyPrimaryObjects module (Otsu two-class adaptive thresholding;
smoothing factors determined empirically) and filtered by size. Channel intensities within
each nuclear object were measured; binary positivity thresholds were set empirically.
Percentages of positive/negative nuclei were calculated from ≥10 images across 2–3
biological replicates.

For TUBB3 quantification (Figs 5–10), nuclear ROIs were expanded by 5 pixels to capture
peri-nuclear signal, which was used as a proxy for total TUBB3 (individual cell-level
quantification was not feasible in these morphologically complex monolayers).
Intensities were normalised to the control sample mean; statistical comparisons used
per-replicate means (N = 4 biological replicates), not individual nucleus counts.

RT-qPCR

973   RNA was extracted with the RNeasy Mini Kit (Qiagen, 74106) and stored at −80°C. cDNA
974   was synthesised with the QuantiTect Reverse Transcription Kit (Qiagen, 205314),
975   including DNase treatment; cDNA was diluted 1:5–1:10 prior to qPCR. Reactions (10 µL,

384-well) were run on a QuantStudio 7 Flex or ViiA 7 Real-Time PCR System. Primer efficiency was validated by serial dilution of a pooled mESC/EpiLC/NE cDNA reference; primers with an efficiency outside 85–115% were excluded. Quantification used the modified ΔΔCt method incorporating primer efficiency.[103] TBP and GAPDH served as housekeeping genes. Expression ratios were log-transformed prior to statistical testing to satisfy normality.

Bulk RNA-seq

RNA was extracted as above and treated with the DNA-free DNA Removal Kit (Invitrogen, AM1906). Poly(A) RNA was enriched using the NEBNext Poly(A) mRNA Magnetic Isolation Module (E7490); strand-specific libraries were prepared with the NEBNext Ultra II Directional RNA Library Prep Kit (E7760) and NEBNext Multiplex Oligos (E6440). Sequencing was performed as 50 bp paired-end reads on NovaSeq 6000 or NovaSeq X (Illumina).

For cofactor overexpression experiments, RNA-seq was performed by Plasmidsaurus (Illumina) using their in-house analysis pipeline; raw counts matrices were used for downstream analysis. Cells (~10^5 per condition) were collected in 50 µL DNA/RNA Shield (Zymo, R1100-50) after Accutase dissociation (300 × g, 3 min).

Computational analysis of RNA-seq

Raw BCL files were converted to FASTQ with bcl2fastq (Illumina). Adapter trimming and quality filtering (Phred ≥ 33) used Trim Galore; trimmed reads were pseudo-aligned to the mouse genome (mm39) with Kallisto.[104]

Kallisto output was imported into R with tximport (DESeq2 package).[105] Genes with fewer than 10 counts in fewer than 3 samples were removed. Differential expression analysis used DESeq2; DEGs (Fig. 1) were defined as $|\log_2 FC| > 0.585$ (≥1.5-fold) and adjusted p < 0.01. Fold change thresholds were > 0 for HDACi RNA-seq (Fig. 5) and > 1 for cofactor experiment (Fig. 7) to account for different variance within these datasets. Variance-stabilised transformation was used for PCA, distance matrix, and heatmap visualisation.[105] ashr was used for shrinkage estimation in volcano plots.[98] Gene Ontology enrichment used enrichGO (clusterProfiler).[106]

ChIP-seq

ChIP-seq was performed as described[107] with modifications. Cells were crosslinked in 1% formaldehyde (10 min, room temperature), quenched with 125 mM glycine, washed twice in ice-cold PBS, and pelleted (2,000 × g, 4 min). Pellets were flash-frozen on dry ice and stored at −80°C. Sequential lysis used: LB1 (50 mM HEPES-KOH pH 7.5, 140 mM NaCl, 1 mM EDTA, 10% glycerol, 0.5% NP-40, 0.25% Triton-X100; 10 min, ice, 2,000 × g), LB2 (10 mM Tris-HCl pH 8, 200 mM NaCl, 1 mM EDTA, 0.5 mM EGTA; 10 min, ice, 2,000 × g), and LB3 (10 mM Tris-HCl pH 8, 100 mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 0.1% Na-deoxycholate, 0.5% N-lauroylsarcosine). All buffers contained cOmplete protease inhibitor (Sigma, 4693159001).

Chromatin was sheared using a Bioruptor Plus (Diagenode, 4°C; 30 s on/off, 16 cycles in
2 rounds of 8) to yield 100–600 bp fragments (verified by agarose gel after heat reversal).
Chromatin concentration was determined by A280 on a NanoDrop (2 μL in 98 μL 0.1 M
NaOH). Per IP, 100 μg chromatin was diluted to 300 μL in LB3 + 1% Triton-X100 and
clarified (16,000 × g, 10 min); 10% was retained as input. IP was performed overnight with
anti-HA antibody (ab9110) pre-loaded on BSA-blocked Protein G Dynabeads (Invitrogen).
Beads were washed 10× in RIPA buffer (50 mM HEPES-KOH, 500 mM LiCl, 1 mM EDTA,
1% Igepal CA-630, 0.7% Na-deoxycholate) and once in TBS (20 mM Tris-HCl pH 7.6, 150
mM NaCl). DNA was eluted overnight at 65°C in 200 μL elution buffer (50 mM Tris-HCl pH
7.6, 10 mM EDTA, 1% SDS), which simultaneously reversed crosslinks.

Eluted DNA was diluted 1:1 in TE, then purified by sequential RNase A digestion (40
μg/mL, 37°C, 30 min), proteinase K digestion (400 μg/mL, 55°C, 2 h), phenol-chloroform-
isoamyl alcohol phase separation in MaXtract HD tubes (Qiagen), and ethanol
precipitation (200 mM NaCl, −20°C) with 1 μL GlycoBlue (20 μg/μL, Thermo Fisher,
AM9515) as co-precipitant. Purified DNA was resuspended in 20 μL 10 mM Tris-HCl pH
8. Enrichment was validated by qPCR at known ASCL1 target loci (percentage input from
10% fraction). Libraries were prepared using ThruPLEX DNA-Seq (Takara, R400675) at the
JCBC NGS Facility and sequenced on NovaSeq 6000 at 15 nM equimolar pooling.

Computational analysis of ChIP-seq data

Reads were trimmed with Trim Galore (Phred ≥ 33) and aligned to a composite
mm39/dm6 index with Bowtie2. *Drosophila* reads (spike-in) were discarded as
normalisation by spike-in was not applied; mm39-aligned reads were used for all
downstream analysis. Peaks were called with MACS3[108] using matched input as
background control. Specificity was confirmed by the near-absence of peaks in control
(no-transgene) samples; any such peaks were excluded from the consensus set. The
consensus peak set comprised peaks present in ≥2 of 4 replicates per condition. Reads-
in-peaks normalisation used RLE (DESeq2). Peaks were annotated to transcription start
sites within 100 kb using AnnotationHub[109] and Ensembl *Mus musculus* v108.

Published ENCODE mouse ASCL1 ChIP-seq datasets were downloaded via accession
numbers (Materials Table) and converted from mm10 to mm39 using UCSC liftOver. [110]
Inter-dataset Pearson correlation and Jaccard similarity were assessed from a binary
peak-presence matrix in DiffBind.

Motif analysis

1048 De novo motif discovery was performed using MEME-ChIP (MEME Suite, web server),[111]
1049 which applies STREME for improved sensitivity at centrally enriched motifs,[112] and
1050 independently with HOMER[113] (-size 150, -mask) for both de novo and known-motif
1051 enrichment.

ATAC-seq

ATAC-seq was performed using the Omni-ATAC protocol.[114,115] To synchronise multi-
timepoint collection, cells were seeded on sequential days. Cells were dissociated with

Accutase, counted with Trypan Blue, and 5×10^4 cells resuspended in ice-cold PBS were
pelleted and lysed in 50 µL RSB++ (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl$_2$,
0.1% NP-40, 0.1% Tween-20, 0.01% digitonin) for 3 min on ice. Nuclei were washed with
1 mL RSB+ (RSB++ without NP-40/digitonin), pelleted (500 × g, 10 min, 4°C), and
resuspended in 50 µL Tn5 transposition mix. Tagmentation was performed at 37°C for 30
min with 1,000 rpm mixing, then cleaned with Zymo DNA Clean & Concentrator (D4014)
and stored at −20°C.

Tagmented DNA was PCR-amplified with indexed i5/i7 primers, purified with the Qiagen
MinElute kit (eluted in 20 µL EB), and size-selected with a dual AMPure XP bead (#A63880)
strategy: (i) 0.55× beads added to remove large fragments (supernatant retained), then
(ii) 1.5× beads to capture the library; beads washed 2× in 80% ethanol and eluted in 20
µL EB. Fragment size distribution was confirmed by Bioanalyzer HS D1000 (Agilent,
#5067-5584). Libraries were pooled equimolarly and sequenced on NovaSeq X 25B
(PE100), targeting ~9×10^7 reads/sample.

Computational analysis of ATAC-seq

FASTQ files were generated with bcl2fastq2. A custom bash pipeline based on nf-core
ATAC-seq[116,117] and published best practices[118] was applied. Quality control used FastQC
followed by Trim Galore (default paired-end parameters). Reads were aligned to mm39
with Bowtie2[119] (--very-sensitive-local, --no-mixed, --no-discordant; maximum insert size
2,000 bp). SAMtools[120] was used to index, sort, and assess alignment statistics;
mitochondrial reads were removed. Duplicates were marked with Picard and removed;
insert size metrics were collected with Picard CollectInsertSizeMetrics. Peaks were
called with MACS3[108] in BAMPE mode (effective genome size 2,468,088,461).

Downstream analysis used R v4.5.0 in RStudio. Peaks overlapping ENCODE blacklisted
regions[121] or unplaced/random contigs were removed. DiffBind generated a consensus
peak set (peaks present in ≥3 of 4 replicates per condition); reads-in-peaks were TMM-
normalised with DiffBind. Bigwig files were produced with bamCoverage (deepTools) [122]
using TMM scale factors; per-condition averages used bamCompare. Visualisation used
IGV[123] and deepTools heatmaps.

References

1084

1. Rostovskaya, M., Stirparo, G.G., and Smith, A. (2019). Capacitation of human naïve
pluripotent stem cells for multi-lineage differentiation. Development *146*, dev172916.
https://doi.org/10.1242/dev.172916.

2. Kinoshita, M., Barber, M., Mansfield, W., Cui, Y., Spindlow, D., Stirparo, G.G., Dietmann, S.,
Nichols, J., and Smith, A. (2021). Capture of Mouse and Human Stem Cells with Features of
Formative Pluripotency. Cell Stem Cell *28*, 453-471.e8.
https://doi.org/10.1016/j.stem.2020.11.005.

3. Buecker, C., Srinivasan, R., Wu, Z., Calo, E., Acampora, D., Faial, T., Simeone, A., Tan, M.,
Swigut, T., and Wysocka, J. (2014). Reorganization of Enhancer Patterns in Transition from
Naive to Primed Pluripotency. Cell Stem Cell *14*, 838–853.
https://doi.org/10.1016/j.stem.2014.04.003.

4. Carbognin, E., Carlini, V., Panariello, F., Chieregato, M., Guerzoni, E., Benvegnù, D., Perrera,
V., Malucelli, C., Cesana, M., Grimaldi, A., et al. (2023). Esrrb guides naive pluripotent cells
through the formative transcriptional programme. Nat. Cell Biol. *25*, 643–657.
https://doi.org/10.1038/s41556-023-01131-x.

5. Hayashi, K., Ohta, H., Kurimoto, K., Aramaki, S., and Saitou, M. (2011). Reconstitution of the
Mouse Germ Cell Specification Pathway in Culture by Pluripotent Stem Cells. Cell *146*, 519–
532. https://doi.org/10.1016/j.cell.2011.06.052.

6. Nishimura, K., Nitta, T., Doi, K., and Takata, K. (2022). Rapid Conversion of Human Induced
Pluripotent Stem Cells into Dopaminergic Neurons by Inducible Expression of Two
Transcription Factors. Stem Cells Dev. *31*, 269–277. https://doi.org/10.1089/scd.2021.0363.

7. Vainorius, G., Novatchkova, M., Michlits, G., Baar, J.C., Raupach, C., Lee, J., Yelagandula, R.,
Wernig, M., and Elling, U. (2023). Ascl1 and Ngn2 convert mouse embryonic stem cells to
neurons via functionally distinct paths. Nat. Commun. *14*, 5341.
https://doi.org/10.1038/s41467-023-40803-y.

8. Thoma, E.C., Wischmeyer, E., Offen, N., Maurus, K., Sirén, A.-L., Schartl, M., and Wagner,
T.U. (2012). Ectopic Expression of Neurogenin 2 Alone is Sufficient to Induce Differentiation
of Embryonic Stem Cells into Mature Neurons. PLOS ONE *7*, e38651.
https://doi.org/10.1371/journal.pone.0038651.

9. Meshorer, E., and Misteli, T. (2006). Chromatin in pluripotent embryonic stem cells and
differentiation. Nat. Rev. Mol. Cell Biol. *7*, 540–546. https://doi.org/10.1038/nrm1938.

10. Peng, T., Zhai, Y., Atlasi, Y., ter Huurne, M., Marks, H., Stunnenberg, H.G., and
Megchelenbrink, W. (2020). STARR-seq identifies active, chromatin-masked, and dormant
enhancers in pluripotent mouse embryonic stem cells. Genome Biol. *21*, 243.
https://doi.org/10.1186/s13059-020-02156-3.

11. Sankar, A., Mohammad, F., Sundaramurthy, A.K., Wang, H., Lerdrup, M., Tatar, T., and Helin,
K. (2022). Histone editing elucidates the functional roles of H3K27 methylation and
acetylation in mammals. Nat. Genet. *54*, 754–760. https://doi.org/10.1038/s41588-022-
01091-2.

12. Voss, T.C., and Hager, G.L. (2014). Dynamic regulation of transcriptional states by chromatin
and transcription factors. Nat. Rev. Genet. *15*, 69–81. https://doi.org/10.1038/nrg3623.

13. Jonasson, M.E., and Buecker, C. (2025). The exit from naive pluripotency: a platform for the
study of enhancer mechanistics. Biochem. Soc. Trans. *53*, 1–15.
https://doi.org/10.1042/BST20253037.

14. Dupont, C., Chahar, D., Trullo, A., Gostan, T., Surcis, C., Grimaud, C., Fisher, D., Feil, R., and
Llères, D. (2023). Evidence for low nanocompaction of heterochromatin in living embryonic
stem cells. EMBO J. *42*, EMBJ2021110286. https://doi.org/10.15252/embj.2021110286.

15. Creyghton, M.P., Cheng, A.W., Welstead, G.G., Kooistra, T., Carey, B.W., Steine, E.J., Hanna,
J., Lodato, M.A., Frampton, G.M., Sharp, P.A., et al. (2010). Histone H3K27ac separates
active from poised enhancers and predicts developmental state. Proc. Natl. Acad. Sci. *107*,
21931–21936. https://doi.org/10.1073/pnas.1016071107.

16. Crispatzu, G., Rehimi, R., Pachano, T., Bleckwehl, T., Cruz-Molina, S., Xiao, C., Mahabir, E.,
Bazzi, H., and Rada-Iglesias, A. (2021). The chromatin, topological and regulatory properties
of pluripotency-associated poised enhancers are conserved in vivo. Nat. Commun. *12*,
4344. https://doi.org/10.1038/s41467-021-24641-4.

17. Cruz-Molina, S., Respuela, P., Tebartz, C., Kolovos, P., Nikolic, M., Fueyo, R., Ijcken, W.F.J.
van, Grosveld, F., Frommolt, P., Bazzi, H., et al. (2017). PRC2 Facilitates the Regulatory
Topology Required for Poised Enhancer Function during Pluripotent Stem Cell
Differentiation. Cell Stem Cell *20*, 689-705.e9. https://doi.org/10.1016/j.stem.2017.02.004.

18. Rugg-Gunn, P.J., Cox, B.J., Ralston, A., and Rossant, J. (2010). Distinct histone modifications
in stem cell lines and tissue lineages from the early mouse embryo. Proc. Natl. Acad. Sci.
107, 10783–10790. https://doi.org/10.1073/pnas.0914507107.

19. Kim, S., and Wysocka, J. (2023). Deciphering the multi-scale, quantitative cis-regulatory
code. Mol. Cell *83*, 373–392. https://doi.org/10.1016/j.molcel.2022.12.032.

20. Ong, C.-T., and Corces, V.G. (2011). Enhancer function: new insights into the regulation of
tissue-specific gene expression. Nat. Rev. Genet. *12*, 283–293.
https://doi.org/10.1038/nrg2957.

21. Ong, C.-T., and Corces, V.G. (2012). Enhancers: emerging roles in cell fate specification.
EMBO Rep. *13*, 423–430. https://doi.org/10.1038/embor.2012.52.

22. Iwafuchi-Doi, M., and Zaret, K.S. (2016). Cell fate control by pioneer transcription factors.
Development *143*, 1833–1837. https://doi.org/10.1242/dev.133900.

23. Zaret, K.S., and Mango, S.E. (2016). Pioneer transcription factors, chromatin dynamics, and
cell fate control. Curr. Opin. Genet. Dev. *37*, 76–81.
https://doi.org/10.1016/J.GDE.2015.12.003.

24. Tapscott, S.J., Davis, R.L., Thayer, M.J., Cheng, P.-F., Weintraub, H., and Lassar, A.B. (1988).
MyoD1: A Nuclear Phosphoprotein Requiring a Myc Homology Region to Convert
Fibroblasts to Myoblasts. Science *242*, 405–411. https://doi.org/10.1126/science.3175662.

25. Davis, R.L., Weintraub, H., and Lassar, A.B. (1987). Expression of a single transfected cDNA
converts fibroblasts to myoblasts. Cell *51*, 987–1000. https://doi.org/10.1016/0092-
8674(87)90585-X.

26. Zaret, K.S., and Carroll, J.S. (2011). Pioneer transcription factors: establishing competence
for gene expression. Genes Dev. *25*, 2227–2241. https://doi.org/10.1101/GAD.176826.111.

27. Freund, M.M., Harrison, M.M., and Torres-Zelada, E.F. (2024). Exploring the reciprocity
between pioneer factors and development. Development *151*, dev201921.
https://doi.org/10.1242/dev.201921.

28. Hansen, J.L., Loell, K.J., and Cohen, B.A. (2022). A test of the pioneer factor hypothesis
using ectopic liver gene activation. eLife *11*, e73358. https://doi.org/10.7554/eLife.73358.

29. Donaghey, J., Thakurela, S., Charlton, J., Chen, J.S., Smith, Z.D., Gu, H., Pop, R., Clement, K.,
Stamenova, E.K., Karnik, R., et al. (2018). Genetic determinants and epigenetic effects of
pioneer-factor occupancy. Nat. Genet. *50*, 250–258. https://doi.org/10.1038/s41588-017-
0034-3.

30. Cernilogar, F.M., Hasenöder, S., Wang, Z., Scheibner, K., Burtscher, I., Sterr, M., Smialowski,
P., Groh, S., Evenroed, I.M., Gilfillan, G.D., et al. (2019). Pre-marked chromatin and
transcription factor co-binding shape the pioneering activity of Foxa2. Nucleic Acids Res.
47, 9069–9086. https://doi.org/10.1093/nar/gkz627.

31. Mayran, A., Sochodolsky, K., Khetchoumian, K., Harris, J., Gauthier, Y., Bemmo, A.,
Balsalobre, A., and Drouin, J. (2019). Pioneer and nonpioneer factor cooperation drives
lineage specific chromatin opening. Nat. Commun. *10*, 3807.
https://doi.org/10.1038/s41467-019-11791-9.

32. Păun, O., Tan, Y.X., Patel, H., Strohbuecker, S., Ghanate, A., Cobolli-Gigli, C., Sopena, M.L.,
Gerontogianni, L., Goldstone, R., Ang, S.-L., et al. (2023). Pioneer factor ASCL1 cooperates
with the mSWI/SNF complex at distal regulatory elements to regulate human neural
differentiation. Genes Dev. *37*, 218–242. https://doi.org/10.1101/gad.350269.122.

33. Guillemot, F. (2007). Spatial and temporal specification of neural fates by transcription
factor codes. Development *134*, 3771–3780. https://doi.org/10.1242/DEV.006379.

34. Mazzoni, E.O., Mahony, S., Closser, M., Morrison, C.A., Nedelec, S., Williams, D.J., An, D.,
Gifford, D.K., and Wichterle, H. (2013). Synergistic binding of transcription factors to cell-
specific enhancers programs motor neuron identity. Nat. Neurosci. *16*, 1219–1227.
https://doi.org/10.1038/nn.3467.

35. Spitz, F., and Furlong, E.E.M. (2012). Transcription factors: from enhancer binding to
developmental control. Nat. Rev. Genet. *13*, 613–626. https://doi.org/10.1038/nrg3207.

36. Morgunova, E., and Taipale, J. (2017). Structural perspective of cooperative transcription
factor binding. Curr. Opin. Struct. Biol. *47*, 1–8. https://doi.org/10.1016/j.sbi.2017.03.006.

37. Srivastava, D., and Mahony, S. (2020). Sequence and chromatin determinants of
transcription factor binding and the establishment of cell type-specific binding patterns.
Biochim. Biophys. Acta BBA - Gene Regul. Mech. *1863*, 194443.
https://doi.org/10.1016/j.bbagrm.2019.194443.

38. Wapinski, O.L., Vierbuchen, T., Qu, K., Lee, Q.Y., Chanda, S., Fuentes, D.R., Giresi, P.G., Ng,
Y.H., Marro, S., Neff, N.F., et al. (2013). Hierarchical mechanisms for transcription factor-
mediated reprogramming of fibroblasts to neurons. Cell *155*, 621.
https://doi.org/10.1016/J.CELL.2013.09.028.

39. Wapinski, O.L., Lee, Q.Y., Chen, A.C., Li, R., Corces, M.R., Ang, C.E., Treutlein, B., Xiang, C.,
Baubet, V., Suchy, F.P., et al. (2017). Rapid Chromatin Switch in the Direct Reprogramming
of Fibroblasts to Neurons. Cell Rep. *20*, 3236–3247.
https://doi.org/10.1016/j.celrep.2017.09.011.

40. Castro, D.S., Martynoga, B., Parras, C., Ramesh, V., Pacary, E., Johnston, C., Drechsel, D.,
Lebel-Potter, M., Garcia, L.G., Hunt, C., et al. (2011). A novel function of the proneural factor
Ascl1 in progenitor proliferation identified by genome-wide characterization of its targets.
Genes Dev. *25*, 930–945. https://doi.org/10.1101/gad.627811.

41. Raposo, A.A.S.F., Vasconcelos, F.F., Drechsel, D., Marie, C., Johnston, C., Dolle, D., Bithell,
A., Gillotin, S., van den Berg, D.L.C., Ettwiller, L., et al. (2015). Ascl1 Coordinately Regulates
Gene Expression and the Chromatin Landscape during Neurogenesis. Cell Rep. *10*, 1544–
1556. https://doi.org/10.1016/j.celrep.2015.02.025.

42. Kagawa, W., and Kurumizaka, H. (2021). Structural basis for DNA sequence recognition by
pioneer factors in nucleosomes. Curr. Opin. Struct. Biol. *71*, 59–64.
https://doi.org/10.1016/j.sbi.2021.05.011.

43. Lundie-Brown, J., Puletti, F., Philpott, A., and Azzarelli, R. (2025). Cell fate acquisition and
reprogramming by the proneural transcription factor ASCL1. Open Biol. *15*, 250018.
https://doi.org/10.1098/rsob.250018.

44. Azzarelli, R., Gillen, S., Connor, F., Lundie-Brown, J., Puletti, F., Drummond, R., Raffaelli, A.,
and Philpott, A. (2024). Phospho-regulation of ASCL1-mediated chromatin opening during
cellular reprogramming. Development *151*, dev204329.
https://doi.org/10.1242/dev.204329.

45. Casey, B.H., Kollipara, R.K., Pozo, K., and Johnson, J.E. (2018). Intrinsic DNA binding
properties demonstrated for lineage-specifying basic helix-loop-helix transcription factors.
Genome Res. *28*, 484–496. https://doi.org/10.1101/GR.224360.117/-/DC1.

46. Talikka, M., Perez, S.E., and Zimmerman, K. (2002). Distinct Patterns of Downstream Target
Activation Are Specified by the Helix–Loop–Helix Domain of Proneural Basic Helix–Loop–
Helix Transcription Factors. Dev. Biol. *247*, 137–148.
https://doi.org/10.1006/DBIO.2002.0677.

47. Mazurier, N., Parain, K., Parlier, D., Pretto, S., Hamdache, J., Vernier, P., Locker, M.,
Bellefroid, E., and Perron, M. (2014). Ascl1 as a Novel Player in the Ptf1a Transcriptional
Network for GABAergic Cell Specification in the Retina. PLoS ONE 9, e92113.
https://doi.org/10.1371/journal.pone.0092113.

48. Gammill, L.S., and Sive, H. (1997). Identification of otx2 target genes and restrictions in
ectodermal competence during Xenopus cement gland formation. Development *124*, 471–
481. https://doi.org/10.1242/dev.124.2.471.

49. Meijsing, S.H., Elbi, C., Luecke, H.F., Hager, G.L., and Yamamoto, K.R. (2007). The Ligand
Binding Domain Controls Glucocorticoid Receptor Dynamics Independent of Ligand
Release. Mol. Cell. Biol. *27*, 2442. https://doi.org/10.1128/MCB.01570-06.

50. Ying, Q.-L., Wray, J., Nichols, J., Batlle-Morera, L., Doble, B., Woodgett, J., Cohen, P., and
Smith, A. (2008). The ground state of embryonic stem cell self-renewal. Nature *453*, 519–
523. https://doi.org/10.1038/nature06968.

51. Ying, Q.-L., Stavridis, M., Griffiths, D., Li, M., and Smith, A. (2003). Conversion of embryonic
stem cells into neuroectodermal precursors in adherent monoculture. Nat. Biotechnol. *21*,
183–186. https://doi.org/10.1038/nbt780.

52. Ying, Q.-L., and Smith, A.G. (2003). Defined Conditions for Neural Commitment and
Differentiation. In Methods in Enzymology Differentiation of Embryonic Stem Cells.
(Academic Press), pp. 327–341. https://doi.org/10.1016/S0076-6879(03)65023-8.

53. Wood, H.B., and Episkopou, V. (1999). Comparative expression of the mouse *Sox1*, *Sox2*
and *Sox3* genes from pre-gastrulation to early somite stages. Mech. Dev. *86*, 197–201.
https://doi.org/10.1016/S0925-4773(99)00116-1.

54. Zhao, S., Nichols, J., Smith, A.G., and Li, M. (2004). SoxB transcription factors specify
neuroectodermal lineage choice in ES cells. Mol. Cell. Neurosci. *27*, 332–342.
https://doi.org/10.1016/j.mcn.2004.08.002.

55. Gouti, M., Tsakiridis, A., Wymeersch, F.J., Huang, Y., Kleinjung, J., Wilson, V., and Briscoe, J.
(2014). In Vitro Generation of Neuromesodermal Progenitors Reveals Distinct Roles for Wnt
Signalling in the Specification of Spinal Cord and Paraxial Mesoderm Identity. PLOS Biol. *12*,
e1001937. https://doi.org/10.1371/journal.pbio.1001937.

56. Smith, A. (2017). Formative pluripotency: the executive phase in a developmental
continuum. Development *144*, 365–373. https://doi.org/10.1242/dev.142679.

57. Evans, M.J., and Kaufman, M.H. (1981). Establishment in culture of pluripotential cells from
mouse embryos. Nature *292*, 154–156. https://doi.org/10.1038/292154a0.

58. Martin, G.R. (1981). Isolation of a pluripotent cell line from early mouse embryos cultured in
medium conditioned by teratocarcinoma stem cells. Proc. Natl. Acad. Sci. *78*, 7634–7638.
https://doi.org/10.1073/pnas.78.12.7634.

59. Hayashi, K., Lopes, S.M.C. de S., Tang, F., and Surani, M.A. (2008). Dynamic Equilibrium and
Heterogeneity of Mouse Pluripotent Stem Cells with Distinct Functional and Epigenetic
States. Cell Stem Cell *3*, 391–401. https://doi.org/10.1016/j.stem.2008.07.027.

60. Beckman, W.F., Parkinson, L.M., Chaytor, L., and Philpott, A. (2025). Distinct proliferative
and neuronal programmes of chromatin binding and gene activation by ASCL1 are cell cycle
stage-specific. Dev. Camb. Engl. *152*, dev204816. https://doi.org/10.1242/dev.204816.

61. Vue, T.Y., Kollipara, R.K., Borromeo, M.D., Smith, T., Mashimo, T., Burns, D.K., Bachoo, R.M.,
and Johnson, J.E. (2020). ASCL1 regulates neurodevelopmental transcription factors and
cell cycle genes in brain tumors of glioma mouse models. Glia *68*, 2613.
https://doi.org/10.1002/GLIA.23873.

62. Castro, D.S., Martynoga, B., Parras, C., Ramesh, V., Pacary, E., Johnston, C., Drechsel, D.,
Lebel-Potter, M., Garcia, L.G., Hunt, C., et al. (2011). A novel function of the proneural factor
Ascl1 in progenitor proliferation identified by genome-wide characterization of its targets.
Genes Dev. *25*, 930–945. https://doi.org/10.1101/gad.627811.

63. Urbán, N., Berg, D.L.C. van den, Forget, A., Andersen, J., Demmers, J.A.A., Hunt, C., Ayrault,
O., and Guillemot, F. (2016). Return to quiescence of mouse neural stem cells by
degradation of a proactivation protein. Science *353*, 292–295.
https://doi.org/10.1126/science.aaf4802.

64. Durdu, S., Iskar, M., Isbel, L., Hoerner, L., Wirbelauer, C., Burger, L., Hess, D.,
Iesmantavicius, V., and Schübeler, D. (2025). Chromatin-dependent motif syntax defines
differentiation trajectories. Mol. Cell *85*, 2900-2918.e16.
https://doi.org/10.1016/j.molcel.2025.07.005.

65. Wang, S., Sun, H., Ma, J., Zang, C., Wang, C., Wang, J., Tang, Q., Meyer, C.A., Zhang, Y., and
Liu, X.S. (2013). Target analysis by integration of transcriptome and ChIP-seq data with
BETA. Nat. Protoc. *8*, 2502–2515. https://doi.org/10.1038/nprot.2013.150.

66. Maienschein-Cline, M., Zhou, J., White, K.P., Sciammas, R., and Dinner, A.R. (2012).
Discovering transcription factor regulatory targets using gene expression and binding data.
Bioinformatics *28*, 206–213. https://doi.org/10.1093/bioinformatics/btr628.

67. Kosicki, M., Baltoumas, F.A., Kelman, G., Boverhof, J., Ong, Y., Cook, L.E., Dickel, D.E.,
Pavlopoulos, G.A., Pennacchio, L.A., and Visel, A. (2025). VISTA Enhancer browser: an
updated database of tissue-specific developmental enhancers. Nucleic Acids Res. *53*,
D324–D330. https://doi.org/10.1093/nar/gkae940.

68. Li, L., Lai, F., Hu, X., Liu, B., Lu, X., Lin, Z., Liu, L., Xiang, Y., Frum, T., Halbisen, M.A., et al.
(2023). Multifaceted SOX2-chromatin interaction underpins pluripotency progression in
early embryos. Science *382*, eadi5516. https://doi.org/10.1126/science.adi5516.

69. Webb, A.E., Pollina, E.A., Vierbuchen, T., Urbán, N., Ucar, D., Leeman, D.S., Martynoga, B.,
Sewak, M., Rando, T.A., Guillemot, F., et al. (2013). FOXO3 Shares Common Targets with
ASCL1 Genome-wide and Inhibits ASCL1-Dependent Neurogenesis. Cell Rep. *4*, 477–491.
https://doi.org/10.1016/j.celrep.2013.06.035.

70. Wang, H., Keepers, B., Qian, Y., Xie, Y., Colon, M., Liu, J., and Qian, L. (2022). Cross-lineage
potential of Ascl1 uncovered by comparing diverse reprogramming regulatomes. Cell Stem
Cell *29*, 1491-1504.e9. https://doi.org/10.1016/j.stem.2022.09.006.

71. Borromeo, M.D., Meredith, D.M., Castro, D.S., Chang, J.C., Tung, K.-C., Guillemot, F., and
Johnson, J.E. (2014). A transcription factor network specifying inhibitory versus excitatory
neurons in the dorsal spinal cord. Development *141*, 2803–2812.
https://doi.org/10.1242/dev.105866.

72. Gaspar-Maia, A., Alajem, A., Meshorer, E., and Ramalho-Santos, M. (2011). Open chromatin
in pluripotency and reprogramming. Nat. Rev. Mol. Cell Biol. *12*, 36–47.
https://doi.org/10.1038/nrm3036.

73. Zhou, V.W., Goren, A., and Bernstein, B.E. (2011). Charting histone modifications and the
functional organization of mammalian genomes. Nat. Rev. Genet. *12*, 7–18.
https://doi.org/10.1038/nrg2905.

74. Rada-Iglesias, A., Bajpai, R., Swigut, T., Brugmann, S.A., Flynn, R.A., and Wysocka, J. (2011).
A unique chromatin signature uncovers early developmental enhancers in humans. Nature
470, 279–283. https://doi.org/10.1038/nature09692.

75. Bleckwehl, T., Crispatzu, G., Schaaf, K., Respuela, P., Bartusel, M., Benson, L., Clark, S.J.,
Dorighi, K.M., Barral, A., Laugsch, M., et al. (2021). Enhancer-associated H3K4 methylation
safeguards in vitro germline competence. Nat. Commun. *12*, 5771.
https://doi.org/10.1038/s41467-021-26065-6.

76. Bergsland, M., Werme, M., Malewicz, M., Perlmann, T., and Muhr, J. (2006). The
establishment of neuronal properties is controlled by Sox4 and Sox11. Genes Dev. *20*,
3475–3486. https://doi.org/10.1101/gad.403406.

77. Bergsland, M., Ramsköld, D., Zaouter, C., Klum, S., Sandberg, R., and Muhr, J. (2011).
Sequentially acting Sox transcription factors in neural lineage development. Genes Dev. *25*,
2453–2464. https://doi.org/10.1101/gad.176008.111.

78. Doucet-Beaupré, H., Gilbert, C., Profes, M.S., Chabrat, A., Pacelli, C., Giguère, N., Rioux, V.,
Charest, J., Deng, Q., Laguna, A., et al. (2016). Lmx1a and Lmx1b regulate mitochondrial
functions and survival of adult midbrain dopaminergic neurons. Proc. Natl. Acad. Sci. U. S.
A. *113*, E4387-4396. https://doi.org/10.1073/pnas.1520387113.

79. Ng-Blichfeldt, J.-P., Drummond, R., Kausar, S., Lando, D., Barile, M., and Philpott, A. (2026).
Hierarchical TBX6-FOXC Regulatory Logic Controls Human Trunk Mesoderm Diversification.
Preprint at bioRxiv, https://doi.org/10.64898/2026.03.17.712295
https://doi.org/10.64898/2026.03.17.712295.

80. Aydin, B., Kakumanu, A., Rossillo, M., Moreno-Estellés, M., Garipler, G., Ringstad, N.,
Flames, N., Mahony, S., and Mazzoni, E.O. (2019). Proneural factors Ascl1 and Neurog2
contribute to neuronal subtype identities by establishing distinct chromatin landscapes.
Nat. Neurosci. *22*, 897–908. https://doi.org/10.1038/s41593-019-0399-y.

81. Marks, H., Kalkan, T., Menafra, R., Denissov, S., Jones, K., Hofemeister, H., Nichols, J., Kranz,
A., Francis Stewart, A., Smith, A., et al. (2012). The Transcriptional and Epigenomic
Foundations of Ground State Pluripotency. Cell *149*, 590–604.
https://doi.org/10.1016/j.cell.2012.03.026.

82. Zhang, J., Donahue, G., Gilbert, M.B., Lapidot, T., Nicetto, D., and Zaret, K.S. (2024). Distinct
H3K9me3 heterochromatin maintenance dynamics govern different gene programmes and
repeats in pluripotent cells. Nat. Cell Biol. *26*, 2115–2128. https://doi.org/10.1038/s41556-
024-01547-z.

83. Todd, C.D., Ijaz, J., Torabi, F., Dovgusha, O., Bevan, S., Cracknell, O., Lohoff, T., Clark, S.,
Argelaguet, R., Pierce, J., et al. (2025). Epigenetic priming of mammalian embryonic
enhancer elements coordinates developmental gene networks. Genome Biol. *26*, 214.
https://doi.org/10.1186/s13059-025-03658-8.

84. Keegan, S.E., Haskins, J., Simmonds, A.J., and Hughes, S.C. (2023). A chromatin
remodelling SWI/SNF subunit, Snr1, regulates neural stem cell determination and
differentiation. Dev. Camb. Engl. *150*, dev201484. https://doi.org/10.1242/dev.201484.

85. Guillemot, F., Lo, L.C., Johnson, J.E., Auerbach, A., Anderson, D.J., and Joyner, A.L. (1993).
Mammalian achaete-scute homolog 1 is required for the early development of olfactory and
autonomic neurons. Cell *75*, 463–476. https://doi.org/10.1016/0092-8674(93)90381-Y.

86. Guillemot, F., and Joyner, A.L. (1993). Dynamic expression of the murine Achaete-Scute
homologue Mash-1 in the developing nervous system. Mech. Dev. *42*, 171–185.
https://doi.org/10.1016/0925-4773(93)90006-J.

87. Paldi, F., Szalay, M.-F., Dufau, S., Di Stefano, M., Reboul, H., Jost, D., Bantignies, F., and
Cavalli, G. (2026). Transient histone deacetylase inhibition induces cellular memory of gene
expression and 3D genome folding. Nat. Genet. *58*, 404–417.
https://doi.org/10.1038/s41588-025-02489-4.

88. Jorstad, N.L., Wilken, M.S., Grimes, W.N., Wohl, S.G., Vandenbosch, L.S., Yoshimatsu, T.,
Wong, R.O., Rieke, F., and Reh, T.A. (2017). Stimulation of functional neuronal regeneration
from Müller glia in adult mice. Nat. 2017 5487665 *548*, 103–107.
https://doi.org/10.1038/nature23283.

89. Hatakeyama, J., and Kageyama, R. (2004). Retinal cell fate determination and bHLH factors.
Semin. Cell Dev. Biol. *15*, 83–89. https://doi.org/10.1016/j.semcdb.2003.09.005.

90. Kim, S., Morgunova, E., Naqvi, S., Goovaerts, S., Bader, M., Koska, M., Popov, A., Luong, C.,
Pogson, A., Swigut, T., et al. (2024). DNA-guided transcription factor cooperativity shapes
face and limb mesenchyme. Cell *187*, 692-711.e26.
https://doi.org/10.1016/j.cell.2023.12.032.

91. Pattyn, A., Simplicio, N., van Doorninck, J.H., Goridis, C., Guillemot, F., and Brunet, J.-F.
(2004). Ascl1/Mash1 is required for the development of central serotonergic neurons. Nat.
Neurosci. *7*, 589–595. https://doi.org/10.1038/nn1247.

92. Pattyn, A., Guillemot, F., and Brunet, J.F. (2006). Delays in neuronal differentiation in
Mash1/Ascl1 mutants. Dev. Biol. *295*, 67–75. https://doi.org/10.1016/J.YDBIO.2006.03.008.

93. Pattyn, A., Morin, X., Cremer, H., Goridis, C., and Brunet, J.-F. (1997). Expression and
interactions of the two closely related homeobox genes Phox2a and Phox2b during
neurogenesis. Development *124*, 4065–4075. https://doi.org/10.1242/dev.124.20.4065.

94. Stanke, M., Stubbusch, J., and Rohrer, H. (2004). Interaction of Mash1 and Phox2b in
sympathetic neuron development. Mol. Cell. Neurosci. *25*, 374–382.
https://doi.org/10.1016/j.mcn.2003.10.021.

95. Lo, L., Tiveron, M.-C., and Anderson, D.J. (1998). MASH1 activates expression of the paired
homeodomain transcription factor Phox2a, and couples pan-neuronal and subtype-specific
components of autonomic neuronal identity. Development *125*, 609–620.
https://doi.org/10.1242/dev.125.4.609.

96. Wang, L., Tan, T.K., Durbin, A.D., Zimmerman, M.W., Abraham, B.J., Tan, S.H., Ngoc, P.C.T.,
Weichert-Leahey, N., Akahane, K., Lawton, L.N., et al. (2019). ASCL1 is a MYCN- and LMO1-
dependent member of the adrenergic neuroblastoma core regulatory circuitry. Nat.
Commun. *10*, 5622. https://doi.org/10.1038/s41467-019-13515-5.

97. Parkinson, L.M., Gillen, S.L., Woods, L.M., Chaytor, L., Marcos, D., Ali, F.R., Carroll, J.S., and
Philpott, A. (2022). The proneural transcription factor ASCL1 regulates cell proliferation and
primes for differentiation in neuroblastoma. Front. Cell Dev. Biol. *10*.

98. Stephens, M. (2017). False discovery rates: a new deal. Biostatistics *18*, 275–294.
https://doi.org/10.1093/biostatistics/kxw041.

99. Mulas, C., Kalkan, T., von Meyenn, F., Leitch, H.G., Nichols, J., and Smith, A. (2019). Defined
conditions for propagation and manipulation of mouse embryonic stem cells. Development
146, dev173146. https://doi.org/10.1242/dev.173146.

100. Bressan, R.B., Dewari, P.S., Kalantzaki, M., Gangoso, E., Matjusaitis, M., Garcia-Diaz, C.,
Blin, C., Grant, V., Bulstrode, H., Gogolok, S., et al. (2017). Efficient CRISPR/Cas9-assisted
gene targeting enables rapid and precise genetic manipulation of mammalian neural stem
cells. Development *144*, 635–648. https://doi.org/10.1242/dev.140855.

101. Mattioni, T., Louvion, J.-F., and Picard, D. (1994). Chapter 16 Regulation of Protein
Activities by Fusion to Steroid Binding Domains. In Methods in Cell Biology Protein
Expression in Animal Cells., M. G. Roth, ed. (Academic Press), pp. 335–352.
https://doi.org/10.1016/S0091-679X(08)60611-1.

102. Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A., and
Goodman, A. (2021). CellProfiler 4: improvements in speed, utility and usability. BMC
Bioinformatics *22*, 433. https://doi.org/10.1186/s12859-021-04344-9.

103. Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time
RT–PCR. Nucleic Acids Res. *29*, e45. https://doi.org/10.1093/nar/29.9.e45.

104. Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic
RNA-seq quantification. Nat. Biotechnol. *34*, 525–527. https://doi.org/10.1038/nbt.3519.

105. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550.
https://doi.org/10.1186/s13059-014-0550-8.

106. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., et
al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The
Innovation *2*. https://doi.org/10.1016/j.xinn.2021.100141.

107. Schmidt, D., Wilson, M.D., Spyrou, C., Brown, G.D., Hadfield, J., and Odom, D.T. (2009).
ChIP-seq: using high-throughput sequencing to discover protein-DNA interactions. Methods
San Diego Calif *48*, 240–248. https://doi.org/10.1016/j.ymeth.2009.03.001.

108. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum,
C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS).
Genome Biol. *9*, R137. https://doi.org/10.1186/gb-2008-9-9-r137.

109. Morgan, M., and Shepherd, L. (2025). AnnotationHub: Client to access AnnotationHub
resources https://doi.org/10.18129/B9.bioc.AnnotationHub.

110. Hinrichs, A.S., Karolchik, D., Baertsch, R., Barber, G.P., Bejerano, G., Clawson, H.,
Diekhans, M., Furey, T.S., Harte, R.A., Hsu, F., et al. (2006). The UCSC Genome Browser

Database: update 2006. Nucleic Acids Res. *34*, D590–D598.
https://doi.org/10.1093/nar/gkj144.

111.   Bailey, T.L., Johnson, J., Grant, C.E., and Noble, W.S. (2015). The MEME Suite. Nucleic
Acids Res. *43*, W39–W49. https://doi.org/10.1093/nar/gkv416.

112.   Bailey, T.L. (2021). STREME: accurate and versatile sequence motif discovery.
Bioinformatics *37*, 2834–2840. https://doi.org/10.1093/bioinformatics/btab203.

113.   Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C.,
Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription
factors prime cis-regulatory elements required for macrophage and B cell identities. Mol.
Cell *38*, 576–589. https://doi.org/10.1016/j.molcel.2010.05.004.

114.   Buenrostro, J.D., Wu, B., Chang, H.Y., and Greenleaf, W.J. (2015). ATAC-seq: A Method for
Assaying Chromatin Accessibility Genome-Wide. Curr. Protoc. Mol. Biol. *109*, 21.29.1-
21.29.9. https://doi.org/10.1002/0471142727.mb2129s109.

115.   Corces, M.R., Trevino, A.E., Hamilton, E.G., Greenside, P.G., Sinnott-Armstrong, N.A.,
Vesuna, S., Satpathy, A.T., Rubin, A.J., Montine, K.S., Wu, B., et al. (2017). An improved
ATAC-seq protocol reduces background and enables interrogation of frozen tissues. Nat.
Methods *14*, 959–962. https://doi.org/10.1038/nmeth.4396.

116.   Ewels, P.A., Peltzer, A., Fillinger, S., Patel, H., Alneberg, J., Wilm, A., Garcia, M.U., Di
Tommaso, P., and Nahnsen, S. (2020). The nf-core framework for community-curated
bioinformatics pipelines. Nat. Biotechnol. *38*, 276–278. https://doi.org/10.1038/s41587-
020-0439-x.

117.   Patel, H., Espinosa-Carrasco, J., Langer, B., Ewels, P., Garcia, M.U., Syme, R., Peltzer, A.,
Talbot, A., Behrens, D., Gabernet, G., et al. (2023). nf-core/atacseq: [2.1.2] - 2022-08-07.
(Zenodo). https://doi.org/10.5281/zenodo.8222875
https://doi.org/10.5281/zenodo.8222875.

118.   Yan, F., Powell, D.R., Curtis, D.J., and Wong, N.C. (2020). From reads to insight: a
hitchhiker's guide to ATAC-seq data analysis. Genome Biol. *21*, 22.
https://doi.org/10.1186/s13059-020-1929-3.

119.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2.
Nat. Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

120.   Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A.,
Keane, T., McCarthy, S.A., Davies, R.M., et al. (2021). Twelve years of SAMtools and
BCFtools. GigaScience *10*, giab008. https://doi.org/10.1093/gigascience/giab008.

121.   Ogata, J.D., Mu, W., Davis, E.S., Xue, B., Harrell, J.C., Sheffield, N.C., Phanstiel, D.H.,
Love, M.I., and Dozmorov, M.G. (2023). excluderanges: exclusion sets for T2T-CHM13,
GRCm39, and other genome assemblies. Bioinformatics *39*, btad198.
https://doi.org/10.1093/bioinformatics/btad198.

122.   Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S.,
Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-
sequencing data analysis. Nucleic Acids Res. *44*, W160–W165.
https://doi.org/10.1093/nar/gkw257.

123.   Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and
Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.
https://doi.org/10.1038/nbt.1754.

1485

Figure 1:

Figure S1:

Figure 2:

A iASCL1-HA ChIP-seq

mESC EpiLC NE

(4,318) *Cebpa Cebpb Tnni2 Dppa1 Vgll1 Eno3*

(1,353)

(1,539)

(3,181)

(2,490) *Myt1 Hdac11 Plxna2 Onecut1 Map2 Wnt5a*

(402)

(3,972)

±1Kb ±1Kb ±1Kb

2 4 6 8 10 / 0 0 0 0 0

B RNA-seq *Myt1*

Norm. counts

C 0 6 24 C 0 6 24 C 0 6 24
mESC EpiLC NE

B' ChIP-seq

0–116

0–116

0–116

5 kb

VISTA: *hs2609* *Myt1*

B" VISTA: *hs2609*

E11.5

C RNA-seq *Hdac11*

Norm. counts

C 0 6 24 C 0 6 24 C 0 6 24
mESC EpiLC NE

C' ChIP-seq

0–526

0–526

0–526

5 kb

VISTA: *mm1314* *Hdac11*

C" VISTA: *mm1314*

E11.5

D ↑DEGs RNA-seq:

235 NE-specific ASCL1 DEGs → **653** ASCL1 sites linked to NE-specific DEGs → Are these binding sites *only* bound in NE?

D'

219 / 211 / 25 / 198

Bound in:
NE only
NE & mESC
NE & EpiLC
All

198 Bound in NE only

0–177 5kb

Wnt5a

455 Bound in NE & at least one other cell type: (236 in mESC, 430 in EpiLC)

0–607 5kb

Plxna2

0–333 5kb

Inppl1

E ChIP-seq signal

NE only *n = 198*   NE & other *n = 455*

Log2(norm. reads + 1)

**** **** ****   **** **** ****

mESC EpiLC NE

Figure S2:

Figure 3:

A

Day 0 1 2 3 4

ATAC
ATAC
ATAC

mESC	EpiLC	NE
No ASCL1	No ASCL1	No ASCL1
6hr ASCL1	6hr ASCL1	6hr ASCL1
24hr ASCL1	24hr ASCL1	24hr ASCL1

B

PC2 (16%) vs PC1 (54%)

mESC
No ASCL1
6 hr ASCL1
24 hr ASCL1
EpiLC
No ASCL1
6 hr ASCL1
24 hr ASCL1
NE
No ASCL1
6 hr ASCL1
24 hr ASCL1

C Accessibility at all ASCL1 binding sites

iASCL1 ChIP-seq
mESC EpiLC NE

ATAC-seq No iASCL1
mESC EpiLC NE

(4,318)
(1,353)
(1,539)
(3,181)
(2,490)
(402)
(3,972)

± 1kb ± 1kb ± 1kb   ± 1kb ± 1kb ± 1kb

0 40 80   0 10 20

Accessibility at 653 ASCL1 binding sites linked to NE-specific ASCL1 target genes

D

NE iASCL1 ChIP-seq | NE ATAC No iASCL1

Open in NE No iASCL1 n = 493

Closed in NE No iASCL1 n = 160

± 1kb ± 1kb

E

NE ATAC 24hr iASCL1

± 1kb

80 40 / 20 10

F

NE ATAC No ASCL1 — 75%

NE ATAC 24hr ASCL1 — 92%

Open / Closed

G *Example ASCL1 binding sites with ASCL1-independent accessibility in NE*

Hdac11   *Plxna2*   *Wnt5a*

NE iASCL1 ChIP-seq   0–528   0–607   0–176
NE ATAC No iASCL1   0–101   0–59   0–21
NE ATAC 24hr iASCL1   0–101   0–59   0–21

5kb 5kb 5kb

H *Example ASCL1 binding sites with ASCL1-altered accessibility in NE*

Myt1   *Elavl4*

NE iASCL1 ChIP-seq   0–116   0–155
NE ATAC No iASCL1   0–36   0–28
NE ATAC 24hr iASCL1 0–36   0–28

5kb 5kb

Figure S3:

A No. ATAC-seq peaks
B FRiP (%)
C TSS enrichment
D % chrM reads

Accessibility of ASCL1 binding sites in each cell type

E All ASCL1 peaks called in mESC

mESC iASCL1 ChIP-seq | mESC ATAC No iASCL1

Open (8,007)

Closed (1,993)

± 1kb | ± 1kb

F All ASCL1 peaks called in EpiLC

EpiLC iASCL1 ChIP-seq | EpiLC ATAC No iASCL1

Open (8,677)

Closed (1,323)

± 1kb | ± 1kb

G All ASCL1 peaks called in NE

NE iASCL1 ChIP-seq | NE ATAC No iASCL1

Open (8,099)

Closed (1,901)

± 1kb | ± 1kb

H % ASCL1 binding sites that overlap with ATAC-seq peak (No iASCL1)

	mESC	EpiLC	NE
mESC only binding sites	75%	42%	34%
NE only binding sites	28%	42%	62%
universal binding sites	85%	89%	89%

I RNA-seq counts for example iASCL1 target genes shown in Fig. 4

Myt1 | *Elavl4* | *Hdac11* | *Plxna2* | *Wnt5a*

iASCL1:
mESC: C 0 6 24
EpiLC: C 0 6 24
NE: C 0 6 24

Figure 4:

653 ASCL1 binding sites at NE-specific ASCL1 targets: **Are these sites _closed_ in mESC?** → 362 are open in mESC: **Why are these sites not activated?**

Figure S4:

A iASCL1 ChIP-seq

mESC   EpiLC   NE

(4,318)
(1,353)
(1,539)
(3,181)
(2,490)
(402)
(3,972)

± 1kb   ± 1kb   ± 1kb

0   40   80

B H3K27ac ChIP-seq:
Cruz-Molina et al., 2017

mESC   AntNPC ≈ NE

± 5kb   ± 5kb

0   3   6

C H3K4me1 ChIP-seq:
Bleckwehl et al., 2021

mESC   EpiLC

C'

± 5kb   ± 5kb

0   10   20

Figure 5:

A Histone deacetylase inhibition

TSA ⊣ HDAC ⟷ HAT

B

Day 0 1 2 3 4 5

RNA
IF   IF

RNA
IF   IF

Condition

mESC	mESC + 10 nM TSA
+ iASCL1	+ iASCL1 + 10 nM TSA
NE	NE + 10 nM TSA
+ iASCL1	+ iASCL1 + 10 nM TSA

C

RNA-seq

mESC   NE

No iASCL1
+10 nM TSA
24 hr iASCL1
+10 nM TSA

PC2: 18% variance
PC1: 60% variance

D *Nkx6-1*

Normalised counts

ns *** *** ***

iASCL1 − + − + − + − +
TSA − − + + − − + +

D' *Disp2*

ns ** *** ***

iASCL1 − + − + − + − +
TSA − − + + − − + +

D" *Plxna2*

ns ns *** ns

iASCL1 − + − + − + − +
TSA − − + + − − + +

E Previously identified NE-specific ASCL1 target genes
n = 208

mESC

iASCL1 − + − +
TSA − − + +

Tmem132d
Dlx2
Disp2
Dpysl4
Nos1
Wnt5a
Lhx1
Nppc
Nkx6-1
Nmrk1
Ptprn2
Tamalin
Camk2n1

Z-score
-2 0 2 4

E'

of 208 NE genes passing
significance threshold

▲ vs ●	iASCL1+TSA vs Control	116
▲ vs ●	TSA only vs Control	102
▲ vs ▲	iASCL1+TSA vs TSA only	13

F All genes up in ASCL1+TSA vs TSA only
n = 458

mESC

iASCL1 − + − +
TSA − − + +

Z-score
-2 0 2 4

F' GO analysis

- placenta development
- mesenchyme development
- regionalization
- morphogenesis of a branching structure
- negative regulation of cell development
- regulation of neurogenesis

Count: 0 10 20 30

p.adjust
4e-08
3e-08
2e-08
1e-08

G

DAPI TUBB3 H3K27ac

Figure S5:

Figure 6:

Figure S6:

A — TF expression log2 fold change (NE 0hr vs mESC 0hr) vs Fold enrichment of motif in peaks (NE only peaks vs universal peaks). Motif family: AP2 (green), Homeobox (magenta), NR (yellow), Other (grey), Zf (cyan). Labeled: Nr5a2, Nr5a2, Esrrb, Klf4, Klf5, Tfap2a, Hoxc13, Batf, Elf3, Elf4, Klf9, Junb, Gabpa, Sp2.

B

10,000 ASCL1 binding sites in NE	**653** ASCL1 binding sites NE-specific DEGs
1,492 sites with **Dlx3** motif	111 sites with **Dlx3** motif
372 sites with **Phox2b** motif	32 sites with **Phox2b** motif

C — piggyBac overexpression

ITR — cHS4 — TRE3G — *COFACTOR* — T2A — mCherry — pA — cHS4 — pEF1α — rtTA — T2A — PuroR — pA — cHS4 — ITR

D — iASCL1 mESC + piggyBac (*Phox2b*)

Dox: 0 µg/mL, 0.01 µg/mL, 0.1 µg/mL, 1 µg/mL; 24 hr, 48 hr

E — *Phox2b* (Transgene). ΔCt vs *Tbp*.
iASCL1: − + − + | − + − +
Cofactor: − − + + | − − + +
mESC (teal), NE (pink)

F — + DLX3 overexpression

Target gene: *Hdac11*, *Zbtb18*, *Map2*
iASCL1: − + − + | − + − +
Cofactor: − − + + | − − + +

G — + LMX1A overexpression

Hdac11, *Zbtb18*, *Map2*
iASCL1: − + − + | − + − +
Cofactor: − − + + | − − + +

Figure 7:

Figure S7: