Chromatin priming and co-factor availability shape lineage response to the neuronal pioneer factor ASCL1 in pluripotency

Jethro Lundie-Brown[1,3], Rosalind Drummond[1], John-Poul Ng-Blichfeldt[1], Roberta Azzarelli[1,3]* and Anna Philpott[1,2]*

1 Cambridge Stem Cell Institute, University of Cambridge, Cambridge, CB2 0AW, UK

2 Department of Oncology, University of Cambridge, Cambridge, CB2 0AH, UK

3 Department of Pharmacology, UCL School of Pharmacy, 29-39 Brunswick Square, London, WC1N 1AX, UK

* Co-corresponding authors

HIGHLIGHTS

• Ectopic ASCL1 drives non-neuronal transcriptional responses in naïve and formative pluripotent cells

• ASCL1 occupies distinct, predominantly non-neuronal genomic targets in pluripotent cells due to differential chromatin accessibility

• ASCL1 pioneer activity is locus- and cell type-specific and predicted by histone acetylation status

• Co-expression of ASCL1 with *Phox2* homeodomain cofactors potentiates neuronal lineage acquisition in pluripotent cells

SUMMARY

Transcription factors act within defined developmental windows, yet how naïve pluripotent cells acquire competence to execute specific transcription factor-driven fate programmes remains unclear. Pioneer transcription factors that engage target sites in closed chromatin to initiate gene expression programmes often act at the top of hierarchies in cell identity transitions. However, we show that the ability of ASCL1 to induce a coherent neuronal programme emerges only after exit from pluripotency, coincident with progressive chromatin remodelling and accumulation of permissive histone marks at neuronal ASCL1 target sites. Binding analysis reveals that although ASCL1 can access a subset of neuronal loci in mESCs and EpiLCs, ASCL1 is preferentially diverted to non-neuronal sites, resulting in divergent transcriptional responses. Increasing global histone acetylation enhances activation of individual neuronal genes but is insufficient to drive full neuronal differentiation. In contrast, co-expression of the homeodomain transcription factor PHOX2B redirects ASCL1 towards neuronal targets while suppressing inappropriate programmes in mESCs. These findings demonstrate that ASCL1 pioneer activity is highly context-dependent and that developmental priming of chromatin is essential for appropriate lineage specification.

INTRODUCTION

Development requires precise control over the generation of diverse cell types. Achieving the correct spatial and temporal coordination of growth, differentiation, and morphogenesis requires cells to regulate their competence to respond to inductive cues. However, the mechanisms by which cells acquire and maintain this competence remain poorly understood.

Mouse embryonic stem cells (mESCs) have proven an invaluable tool for studying competence for differentiation. The transition from naïve to formative pluripotency has revealed insights into the reorganisation of chromatin and transcription factor availability that prepares cells for subsequent state transitions.[1–5] However, the effect of these transitions on competence for differentiation in response to lineage-specifying transcription factors (TFs) remains incompletely understood. ESCs are widely used in forward programming strategies, in which TF overexpression drives direct conversion to specific cell types in vitro.[6–9] These approaches typically require prior priming through altered culture conditions[7] or embryoid body formation,[9] suggesting that naïve pluripotent cells are not inherently competent to respond to lineage-specifying TFs. Indeed, recent work showed that mESCs are refractory to directed differentiation induced by ASCL1,[10] but a mechanistic understanding of this phenomenon remains elusive. Understanding how such TFs act in naïve pluripotent cells would therefore clarify both the biology of this foundational cell state and the epigenetic basis by which competence for TF-driven cell fate change is established.

Competence to respond to TFs can be shaped by chromatin patterning.[11] Chromatin in naïve pluripotency is distinct from other tissues and reflects their capacity for multilineage differentiation and their transient nature in development.[12–15] Naïve pluripotent cells typically have higher levels of chromatin accessibility compared with differentiated cells.[12,16,17] Although open chromatin does not necessarily determine enhancer activity,[16] active enhancers usually exhibit local depletion of nucleosome occupancy.[18] Many enhancers in early development are accessible but maintained in a poised state marked by the presence of both activation-associated H3K4me1 and repressive H3K27me3, rather than the H3K27ac signature of active enhancers.[18–21] The ability of a cell to interpret and respond to lineage-specifying TFs is therefore determined partially by the availability of TFs and their recruitment to these regulatory elements, and partially by the chromatin states that permit or restrict their binding and activity.[22–24]

Certain TFs, however, have the ability to bind their preferred motifs in nucleosome-dense chromatin and initiate opening and activation of target gene expression.[25,26] These so-called pioneer factors often sit at the apex of transcriptional networks to drive an entire lineage program during both development and reprogramming,[27,28] in some cases directly establishing competence for differentiation.[25,29,30] The extent to which these pioneer TFs are subject to the same restrictions imposed by the chromatin landscape described above is not clear. Moreover, the simple distinction between pioneer and non-pioneer TFs has been challenged,[31] as some pioneer factors display different preferences for open and closed chromatin across cell types and at different loci.[32–35] Their activity is also

frequently modulated by cofactors that refine or redirect their function, and TFs often
cooperate to control cell fate in development and reprogramming.[11,36–39]

ASCL1, a proneural bHLH TF, drives both fibroblast-to-neuron conversion[40,41]
and neural stem cell differentiation *in vivo* and *in vitro*.[42,43] ASCL1 has been
described as a pioneer factor due to its ability to bind nucleosomal DNA directly.[44]
However, it also binds partially accessible chromatin and recruits chromatin
remodelling enzymes.[35] Evidence from somatic reprogramming has further
suggested that competence for ASCL1-mediated conversion correlates with a
specific epigenetic signature at its binding sites, [40,41] yet how epigenetic constraints
shape the developmental acquisition of this competence remains largely unexplored.
ASCL1-mediated differentiation therefore represents a tractable system for
dissecting how the epigenetic landscape of early developmental states determines
competence to respond to lineage-specifying pioneer TFs.[7,10,45,46]

Here, we investigate when and how cells acquire competence to respond to
ASCL1 during the exit from pluripotency. By integrating RNA-seq, ASCL1 ChIP-seq,
and chromatin accessibility mapping across naïve mESCs, epiblast-like cells
(EpiLCs), and neuroectoderm (NE) states, we show that ASCL1 binding is highly
context-dependent: key neuronal target loci are inaccessible and transcriptionally
refractory in pluripotent cells, while genes associated with divergent pathways are
readily activated. We find that the onset of competence to respond to ASCL1 with a
coherent neuronal programme correlates with progressive chromatin remodelling
and acquisition of permissive histone modifications at neuronal ASCL1 target sites.
The ability of ASCL1 to drive neurogenesis is therefore dependent on developmental
priming of chromatin that is established after the exit from pluripotency. We further
show that the barrier to activation in mESCs for neuronal ASCL1 target genes can be
partially overcome by elevating levels of histone acetylation, or co-expressing
homeodomain TFs that bias ASCL1 activity towards neuronal targets. Together,
these findings establish that ASCL1 pioneer factor activity is constrained in an early
developmental context and reframe our understanding of how competence for
somatic differentiation in response to transcriptional cues is encoded in the
epigenome.

RESULTS

An in vitro model of early development with inducible ASCL1 activity reveals changes in competence for neuronal differentiation

Our previous work has shown that naïve mouse embryonic stem cells (mESCs) do not respond to ASCL1 overexpression by activating a neurogenic programme.[10] Here we set out to understand why ASCL1 fails to drive neuronal differentiation in mESCs and investigate what, if any, is the transcriptional response to ASCL1 in mESCs and epiblast-like cells (EpiLC) compared to the response in neuroectodermal (NE) cells. To this end, we developed an mESC-based system with inducible ASCL1 activation (iASCL1 mESCs). HA-tagged ASCL1 was fused with the ligand-binding domain of the glucocorticoid receptor (LBD-GR),[43,47,48] enabling conditional control by the addition of dexamethasone (Dex) to the culture medium (Fig. 1A and Fig. S1B).[49,50] The inducible ASCL1-GR transgene (iASCL1), driven by a constitutively active promoter, was inserted at the *Rosa26* locus (see Methods) (Fig. 1A). Constitutive expression of iASCL1 was confirmed in iASCL1 mESCs (Fig. S1C). iASCL1 mESCs can be propagated in standard 2iLIF pluripotency conditions,[51] where cells express pluripotency markers, NANOG and OCT4, in the absence of Dex (Fig. S1A).

Culture of mESCs in N2B27 media without 2iLIF spontaneously gives rise to anterior neural ectoderm cells (NE) (Fig. S1D).[52,53] Expression of SOX1, an early marker of NE,[54,55] was detected at low levels from days 2 and 3 and at high levels throughout cultured cells from day 4 (Fig. S1D–E). This was concomitant with a loss of pluripotency marker OCT4 between days 3 and 4 (Fig. S1D) and followed by PAX6 and sparse TUBB3 expression from day 5, indicating spontaneous but limited neuronal differentiation from NE by this later stage (Fig. S1D). NE cells at this stage do not express endogenous ASCL1.[56] mESC to NE differentiation proceeded via a NANOG-negative, OTX2-positive state resembling EpiLCs in the transition from naïve to formative pluripotency,[5,57] but this identity was transient and was lost by day 2 (Fig. S1E). The EpiLC identity can be stabilised with activin A and FGF2,[3,5] enabling interrogation of the formative state without spontaneous differentiation (Fig.1B–C and Fig.S1F). Together, mESCs, EpiLCs and NE cells constitute an *in vitro* model of the transition from pluripotency through to neuroectoderm after gastrulation (Fig.1B & C).

We next characterised the response to iASCL1 across these cell types. In NE cells, activation of iASCL1 produced limited TUBB3-positive cells within 24 hours (Fig. 1D) but by 48 hours, neuronal differentiation was widespread and TUBB3 expression was robust compared to the sparse, spontaneous neuronal differentiation observed in NE cells without activated iASCL1 (Fig. 1D). EpiLCs displayed sparse TUBB3 expression only after 48 hours of activated iASCL1, with morphology that indicates less advanced neuronal differentiation (Fig. 1D). As expected,[10] mESCs failed to upregulate TUBB3 with or without iASCL1 over this period (Fig. 1D). This confirmed that competence for differentiation in response to ASCL1 activity is acquired during the exit from pluripotency. To exclude the possibility that resistance to iASCL1 in mESCs is attributable simply to inhibitors present in the media, we also

assessed the response to iASCL1 in variants of 2iLIF lacking either the ERK inhibitor
(Chiron-LIF) or the GSK3β inhibitor (PD-LIF), as well as in serum-LIF which also
maintains pluripotency of mESCs[58–60] (Fig. S1G). ASCL1-driven expression of target
genes (*Tubb3* and *Hes5*) was observed exclusively in NE cells and not in these
variant 2iLIF conditions, indicating that neither ERK inhibition nor GSK3β inhibition
alone was responsible for suppressing neuronal differentiation (Fig. S1G).

During ASCL1-mediated differentiation, neural progenitor cells undergo rapid
cell cycle exit[61–63] and dismantle the progenitor regulatory network.[61–64] We asked
whether pluripotent cells responded in a similar way, despite the failure to induce
neuronal differentiation. We found that 24 hours of iASCL1 activation in 2iLIF mESCs
resulted in a significant reduction in the number of NANOG- and OCT4-positive
mESCs (from 90–100% to 55–60%, $p < 0.05$) (Fig. 1E) and a decrease in the
number of EdU-positive cycling cells (from 65% to 40%) (Fig. S1H). This is
consistent with previous reports of ASCL1 activity accelerating loss of pluripotency
genes in differentiating mESCs.[7] These results indicate biological activity of iASCL1
in mESCs and suggest that the failure of mESCs to differentiate into neurons in
response to ASCL1 is not caused by intrinsic resistance to ASCL1-mediated
transcriptional activation.

177 ASCL1 activates distinct transcriptional profiles in pluripotent, epiblast-like
178 and neuroectodermal cells

To investigate the transcriptional response to iASCL1 genome-wide in mESC,
EpiLC and NE, we performed RNA-seq in cells exposed to 0, 6 or 24 hours of
iASCL1 activity before collection (Fig. 1B and 1F). ASCL1 altered the expression of
numerous genes in all three cell types (Fig. 1G and Fig. S1I–K) (Table S1)
demonstrating its transcriptional activity in all three contexts. Moreover, the greatest
number of upregulated differentially expressed genes (DEGs) after 24 hours of
ASCL1 activation was seen in mESCs (mESC: 780; EpiLC: 310; NE: 419) (Fig. 1G).
Fewer genes were downregulated than upregulated in all cell types (Fig. 1G),
consistent with ASCL1's established predominant role as an activator rather than a
repressor.[43] Subsequent analysis therefore focused on upregulated DEGs. A set of
63 "universal" ASCL1-responsive DEGs upregulated in all three contexts was
identified, including known ASCL1- and Notch-related targets, *Dll3*, *Id1*, and *Id3*.
However, the majority of ASCL1-upregulated DEGs after 24 hours were cell type-
specific, including 235 genes that were only upregulated by ASCL1 in NE, 114 in
EpiLC and 601 in mESC (Fig. 1H–I).

To better understand the functional identity of genes activated by ASCL1 in
each cell type, we performed gene ontology (GO) analysis on upregulated DEGs
from mESCs, EpiLCs and NE cells (Fig. 1J–L) (Table S2). iASCL1-responsive DEGs
in NE were associated with neurogenesis-related terms including "forebrain
development" and "regulation of neuron differentiation" (Fig. 1L), whereas ASCL1-
responsive genes in mESC and EpiLC, were associated with terms including
"placenta development", "hemopoiesis", "muscle cell differentiation", and
"mesenchyme development" (Fig. 1J–K), indicating activation of divergent
programmes not associated with neurogenesis. Notably, ASCL1-activated genes

associated with "central nervous system neuron differentiation" were upregulated in
EpiLCs, yet these cells displayed only limited TUBB3-expression (Fig. 1D and 1K).
This indicates that at least some of the 235 NE-only ASCL1 responsive genes must
be essential for neuronal differentiation; their resistance to activation in response to
ASCL1 in mESC and EpiLC likely contributes to the limited competence of these
cells to undergo ASCL1-mediated neuronal differentiation.

The list of DEGs identified after 24 hours of iASCL1 activity will also include
targets of secondary effectors in addition to genes that are directly targeted by
ASCL1. It is possible that ASCL1 directly activates the same target genes in all cell
types, and subsequent transcriptional responses diverge due to downstream effects.
One advantage of our iASCL1 system is the ability to capture early transcriptional
changes; we therefore also identified ASCL1-responsive DEGs after 6 hours of
iASCL1 activation in each cell type (Fig. S1L–O) (Table S1). As expected, fewer
DEGs were identified after 6 hours than at 24 hours, but ASCL1-responsive DEGs
were already cell type-specific at this earlier timepoint (Fig. S1L) and GO analysis of
these 6-hour DEGs also revealed that neuronal-associated terms were only highly
represented in NE (Fig. S1M–O) (Table S2). Together, these data support a model
for distinct ASCL1 activity in these three cell types that is largely shaped by
differences in the initial cellular context, rather than as a result of downstream
divergent responses to ASCL1 activation.

ASCL1 binding is cell-type specific and explains divergent transcriptional
responses

To activate its target genes, ASCL1 must bind to *cis*-regulatory elements
(CREs) and initiate transcription. ASCL1 has been described as an on-target pioneer
TF, and previous reports have claimed that its binding sites are similar in both
developmental and reprogramming contexts.[40,41] We therefore asked whether
ASCL1 binds distinct loci in mESC, EpiLC and NE and whether this explains the
divergent transcriptional responses we observed. We used ChIP-seq to probe
genome-wide ASCL1 binding (Fig. S2A–C) and focused our analysis on the top
10,000 peaks from each cell type (Fig. S2D). This resulted in a total consensus set
of 17,210 unique ASCL1 binding sites bound in at least one of mESC, EpiLC or NE
(Fig.2A). Grouping these peaks revealed striking cell type-specific binding patterns
for ASCL1 (Fig. 2A) (Table S3). EpiLCs and NE have more ASCL1-binding sites in
common (3,181) than either mESC and EpiLC (1,353) or mESC and NE (402). A
further 3,972 ASCL1-binding sites are shared in all three cell types (Fig. 2A).
Distinguishing ASCL1 binding sites at promoters, introns, or distal intergenic regions
revealed similar proportions in mESCs, EpiLCs and NE (Fig. S2E). Motif enrichment
analysis across all ASCL1 site clusters confirmed that binding sites in all three cell
types share a similar but not identical central E-box motif (CAGCTG), rather than
harbouring cell type-specific E-box variants (Fig. S2F). Preferential binding of bHLH
TFs to E-box variants can be altered by formation of distinct dimers,[65] but the
similarity of motifs discovered in our ASCL1 ChIP-seq suggests that this is not a
primary determinant of cell type-specific ASCL1 activity in mESC, EpiLC and NE.

We next asked whether these divergent binding profiles of ASCL1 could
explain the divergent transcriptional profiles. We therefore integrated our ChIP-seq
and RNA-seq data to determine the relationship between ASCL1 binding and
ASCL1-responsive differentially expressed genes, identifying all ASCL1 binding sites
within 100 kb upstream or downstream of gene transcription start sites (Table
S3).[66,67] We found that a large majority of the most upregulated ASCL1-responsive
DEGs in each cell type were proximal to at least one ASCL1 binding site (mESC:
685/780; EpiLC: 291/310; NE: 385/419) (Fig. S2G). Furthermore, differentially
expressed neuronal genes that are only expressed in NE were associated with NE-
specific ASCL1 binding sites, while non-neuronal mESC-specific ASCL1-responsive
DEGs (Fig. 2A, labelled genes) were associated with mESC-specific binding sites.

To confirm the *in vivo* neuronal activity of the NE-specific ASCL1 regulatory
elements, we compared these ASCL1 binding sites to the location of validated
enhancers from the VISTA database.[68] Gene expression of NE-specific DEGs and
their associated NE-specific ASCL1 binding sites is consistent with *in vivo* activity of
enhancers associated with these sites in the developing mouse forebrain, hindbrain
and spinal cord (Fig. 2B–B" and Fig. 2C–C"). These data indicate that ectopic
ASCL1 in NE is recruited to physiologically relevant ASCL1 binding sites, even
before endogenous ASCL1 has been activated.

NE cells undergo a strong upregulation of the neuronal programme on ASCL1
activation (Fig. 1D and L). To understand more about the mechanisms that underpin
ASCL1-mediated neurogenesis, we next focused more closely on the 235 NE-
specific ASCL1-responsive DEGs (Fig. 1H) and found that 208 of them were
associated with an ASCL1 binding site within ± 100kb. As some of these genes were
linked to multiple ASCL1 binding sites, this corresponded to 653 ASCL1 binding sites
proximal to the 208 bound and NE-specific target genes (Fig. 2D) (Table S4). Since
ASCL1 failed to activate neurogenesis in mESCs and EpiLCs, we then asked
whether these 653 binding sites were also bound by ASCL1 in mESCs or EpiLCs or
not. Surprisingly, 455 of these sites associated with genes only upregulated in NE
were also bound by ASCL1 in either mESC or EpiLC, while only 198 of these sites
were exclusively bound by ASCL1 in NE (Fig. 2D'). However, where ASCL1 was
found to be bound to specific sites in two or more cell types, average ChIP-seq
signal at the shared sites was significantly higher in NE than in mESC and EpiLC
(Fig. 2E). Taken together, the ChIP-seq data suggest that ASCL1 binding, although
present at some of the correct neuronal target genes, is lower in mESC and EpiLC
and binding at non-neuronal sites is favoured. This may underlie the reduced
competence of mESC and EpiLC to respond to ASCL1 by executing a neuronal
differentiation programme.

Context-dependent binding has been reported for various TFs, including
pioneer factors,[69] but ASCL1 has been described as targeting the same sites in
development and reprogramming.[40] To further explore context-dependent ASCL1
activity, we curated a range of published ASCL1 ChIP-seq datasets from *in vitro*
neural stem cells (NSCs),[40,43] *ex vivo* adult neural progenitors,[70] mouse embryonic
fibroblasts (MEFs),[40,71] mESCs,[46] and embryonic neural tube[72] (Fig. S2H) and
compared with our ASCL1 binding sites from mESC, EpiLC and NE. Correlation

analysis comparing the similarity of ASCL1 binding sites across these datasets
revealed that ASCL1 indeed displays cell type-specific occupancy of binding sites,
even across cell types that undergo ASCL1-mediated neuronal differentiation (e.g.
NSCs and MEFs). Looking more closely at strength of binding at cell type-specific
peaks, ASCL1 binding in our ASCL1 NE ChIP-seq was higher than in mESC or
EpiLC at ASCL1 binding sites that are also found in NSC and embryonic neural tube
from published datasets[43,72] (Fig. S2I). This analysis suggests that the pioneer model
of ASCL1 activity where ASCL1 accesses the same sites in a variety of chromatin
states across cell types does not extend to explaining binding differences across cell
types in very early development.[40] Instead, the data indicate that its binding is
context-dependent and that competence for ASCL1-mediated neuronal
differentiation is underpinned by binding at specific sets of neurogenic regulatory
elements that is restricted in pluripotent cells.

Cell type-specific chromatin pre-patterning correlates with distinct binding
patterns

Having established that ASCL1 occupies distinct binding sites in mESCs,
EpiLCs, and NE, we next investigated chromatin features that may explain these
differences. Pluripotent cells are characterised by having globally low levels of
repressive heterochromatin,[12,73] so it was striking that ASCL1 fails to bind its NE-
specific sites in mESCs. We sought to better understand whether chromatin
accessibility might explain distinct ASCL1 binding profiles. To examine ASCL1
binding site accessibility across cell types, we performed ATAC-seq on mESCs,
EpiLCs, and NE that had been exposed to either 6 or 24 hours of iASCL1 activity
before collection (Fig. 3A–B and Fig. S3A–D). We also took untreated samples at the
same final time point (No ASCL1) (Fig. 3A–C) to measure chromatin accessibility
that is independent of ASCL1 activity. Surprisingly, given ASCL1's affinity for
nucleosomal DNA,[44] we found that 80–87% of ASCL1 binding sites were accessible
in each cell type even in the absence of iASCL1 activity (Fig. S3E–G). Moreover,
differences in chromatin accessibility between cell types in the absence of ASCL1
closely mirrored the cell type-specific ASCL1 binding patterns described above (Fig.
3C). For example, mESC-only ASCL1 binding sites are more likely to be accessible
in mESC than those same sites in EpiLC or NE in the absence of ASCL1 activation,
with a similar pattern seen for other cell type-specific binding sites (Fig. 3C and Fig.
S3H). This indicates preferential recruitment of ASCL1 to binding sites that are
already accessible and that these vary between cell types.

We focused our analysis on the previously defined 653 ASCL1 binding sites
associated with NE-specific ASCL1 target genes and asked whether these sites
were open or closed in NE in the absence of iASCL1 activation (Table S4).
Consistent with the global pattern described above, 493 (75%) of these 653 ASCL1
binding sites were open in NE even without iASCL1 activity (Fig. 3D and 3F).
Therefore, most of the binding sites linked to NE-specific target genes are open in
NE independently of ASCL1 (Fig. 3G and Fig. S3I). The remaining 160 (25%) of
these 653 ASCL1 binding sites were inaccessible in the absence of ASCL1 (Fig. 3D
and 3F). After 24 hours of iASCL1 activity, 598 (92%) of the 653 sites were open,
indicating 105 of 160 initially closed sites had been rendered accessible after 24

hours of ASCL1 activity (Fig. 3E–3F and 3H). Therefore, in NE the majority of
binding sites associated with NE-specific gene expression were open independently
of ASCL1 activity, while additional further sites were pioneered after ASCL1
activation (Fig. 3E–F and 3H).

Given this strong preference for ASCL1 to bind to accessible chromatin in NE,
we asked whether the 653 ASCL1 binding sites linked to NE-specific gene activation
were or were not accessible in mESCs. Surprisingly, we found that 55% of these
sites were also accessible in mESCs in the absence of ASCL1 (Fig. 4A–B). Since
the genes associated with these open sites are not responsive to ASCL1 in mESCs,
chromatin accessibility alone cannot fully account for the divergent transcriptional
response to ASCL1 at these loci. We therefore asked why transcriptional activation
of these genes fails in mESC despite the fact that many of the ASCL1 binding sites
associated with them are open and bound by ASCL1 (Fig. 4A–B). We reasoned that
histone modifications might provide an additional layer of gene regulation.

Histone H3 acetylation at lysine 27 (H3K27ac) and methylation of lysine 4
(H3K4me1) both correlate with transcriptional activity of enhancers.[74] Moreover,
mESCs have been shown to maintain enhancers associated with differentiation in a
poised state, marked by accessible, H3K4me1-rich chromatin but lacking
H3K27ac.[18,20] Poised enhancers are not active, but respond rapidly at the onset of
differentiation.[19,20,75] To see whether differential enrichment of these marks at ASCL1
binding sites in mESC and NE might explain the cell type-specific patterns of ASCL1
target gene activation, we analysed published histone ChIP-seq data[20,76] from 2iLIF
mESCs, EpiLCs and anterior neural precursor cells (AntNPCs) (Fig. S4A–D).
AntNPCs are approximately equivalent to NE cells and they have also not yet
upregulated endogenous ASCL1 expression.[20,56] Globally, we found that cell type-
specific ASCL1 binding correlated with H3K27ac (Fig. S4B) and H3K4me1 marks
(Fig. S4C), consistent with preferential recruitment of ASCL1 to active chromatin.
Notably, there was no enrichment of H3K4me1 at NE-specific ASCL1 binding sites in
mESC or EpiLC, indicating that these sites are not maintained in a poised state in
these cells (Fig. S4C').

We then looked specifically at the 362 ASCL1 binding sites that were open in
mESCs, despite only being associated with ASCL1 transcriptional activity in NE (Fig.
4A–B). Of these 362 open ASCL1 binding sites, 176 were not bound in mESCs,
despite being accessible (Fig. 4C). We saw that these sites displayed low levels of
H3K27ac deposition in mESC (Fig. 4D), consistent with their maintenance in inactive
chromatin. On the other hand, these sites displayed high levels of deposited
H3K27ac in NE (AntNPC) (Fig. 4D). This indicates that these sites are not bound or
activated by ASCL1, despite their accessibility, because they are not marked with
active H3K27ac. The remaining 186 sites were bound in mESC and open in mESC,
even though they also displayed NE-specific transcriptional upregulation of their
associated genes in response to ASCL1 (Fig. 4C'). We saw that H3K27ac was lower
at these sites in ESC than in NE (AntNPC), indicating that lower levels of pre-existing
H3K27ac marks contribute to a failure of ASCL1 to activate associated NE-specific
genes in ESCs, even when ASCL1 is bound at these sites in both cell types.
Example NE-specific ASCL1 target genes (Fig. 4E) and H3K27ac marks at their

associated binding sites (Fig. 4F–G) highlight this pattern. For example, *Ank2* is
bound and accessible in mESC and NE, but displays a clear enrichment of H3K27ac
at its ASCL1 binding site in NE (AntNPC) compared to mESC (Fig. 4F). By contrast,
Cyth1 and *Hes5* are examples of genes that are accessible but not bound in mESC,
and these sites show low levels of H3K27ac in mESCs (Fig. 4G). These findings
indicate that the NE-specific ASCL1 target genes are not activated in mESCs
because they are either inaccessible to ASCL1 or accessible but not marked with
active H3K27ac.

389   Modulating histone acetylation enhances ASCL1 transcriptional activity, but
390   does not facilitate neuronal differentiation in mESCs

Our epigenetic data suggest that pre-existing chromatin status, particularly
histone acetylation and chromatin accessibility, is the primary determinant of ASCL1
binding and cellular competence for ASCL1-mediated neuronal gene activation. We
therefore asked whether artificially enhancing histone acetylation could increase the
competence of mESCs to respond to iASCL1 (Fig. 5A). We treated mESCs and NE
cells with 10 nM trichostatin A (TSA), a class I/II histone deacetylase (HDAC)
inhibitor that broadly elevates H3K27 acetylation, and profiled the transcriptional
response to iASCL1 activation by RNA-seq (Fig. 5B–C and Fig. S5A–B).

TSA treatment altered the transcriptome of both mESCs and NE cells with
and without iASCL1 (Fig. 5C and S5B–F) (Table S5). To assess whether TSA could
alter the responsiveness of NE-specific ASCL1 target genes in mESCs, we focused
on this previously defined set of genes (Fig. 2D). Several of these genes have
established roles in neurogenesis, including *Nkx6-1*, *Disp2*, and *Plxna2* (Fig. 5D–D''),
and all three of these were now upregulated by iASCL1 in mESCs when co-treated
with TSA (Fig. 5D–D''). More broadly, combined iASCL1+TSA treatment drove the
highest expression of NE-specific ASCL1 target genes in mESCs compared to
iASCL1 or TSA alone, with 116 genes significantly upregulated relative to the control
in this condition (Fig. 5E–E').

However, TSA alone had a substantial independent effect on the mESC
transcriptome compared to untreated cells, upregulating 4,924 genes including 102
of the previously identified NE-specific iASCL1 targets (Fig. S5D and Fig. 5E'),
suggesting that many of these changes reflect opportunistic activation by other
factors in response to broadly increased acetylation rather than directed ASCL1
activity.

Beyond analysis of the previously identified NE-specific targets described
above, broader transcriptional analysis of differentially expressed genes in mESCs
revealed many additional genes that responded to combined iASCL1+TSA treatment
relative to TSA alone (n = 458) (Fig. 5F and S5F). GO analysis of these genes
returned terms including "placenta development" and "mesenchyme development"
as the top hits, alongside "regulation of neurogenesis" (Fig. 5F'), indicating that co-
treatment activates a largely incoherent and generally non-neuronal transcriptional
programme. Consistent with this, co-treatment with iASCL1 and TSA failed to
produce TUBB3-positive neurons within 48 hours (Fig. 5G). This is likely because
global changes to the acetylation landscape caused by TSA treatment expose

additional sites primed for ASCL1 binding at non-neuronal targets, generating
conflicting alternative transcriptional programmes. Taken together, these results
show that globally increasing histone acetylation can sensitise individual NE-specific
target genes to ASCL1 in mESCs but is insufficient to redirect ASCL1 activity
towards a productive neuronal differentiation programme and away from divergent
pathways.

Homeodomain motifs are enriched at ASCL1 binding sites

We next asked whether we could use more targeted approaches to engage a
neuronal ASCL1 programme in mESCs, which do not have an epigenetic landscape
that favours directed neuronal differentiation. We reasoned that, despite the
availability of open neuronal gene-associated ASCL1 binding sites, ASCL1 activation
of neuronal target genes may be limited in mESCs because essential co-binding TFs
that normally guide ASCL1 activity to these sites in an NE context are absent from
mESCs. To identify cofactors capable of enhancing ASCL1 activity at genomic sites
associated with neuronal genes, we examined potential cofactor motif enrichment in
NE-specific ASCL1 binding sites compared against the set of universally bound sites
(Fig. 6A), prioritising motifs associated with transcription factors found at higher
levels in NE relative to mESCs before ASCL1 activation (Fig. 6A). We confirmed that
these motifs were also enriched in the 653 sites associated with NE-specific iASCL1
target genes (Fig. S6B). This analysis identified Sox (HMG) and Homeodomain-
containing TFs as the dominant motif families at NE-specific ASCL1 binding sites. In
contrast, mESC-specific ASCL1 binding sites were enriched for motifs associated
with nuclear receptor, zinc finger and AP2 TFs (Fig. S6A), indicating distinct cofactor
environments in mESCs and NE.

SOX4 and SOX11 are known to act downstream of bHLH proteins in neuronal
development,[77,78] while SOX1 is expressed in neuroectoderm. SOX2 is expressed in
both pluripotency and neural stem cells, so is unlikely to be the limiting factor for
ASCL1-mediated neuronal differentiation in mESCs. Top candidate homeodomain
cofactors identified in our analysis included PHOX2B, PHOX2A, and DLX3 (Fig. 6A
and Fig. S6B). DLX3 has been reported to potentiate ASCL1 reprogramming of
fibroblasts.[41] We therefore decided to focus on relatively unexplored
homeodomain⬚containing transcription factors as potential ASCL1 cofactors in
potentiating context-dependent coherent neuronal programming. We also included
LMX1A as a representative homeodomain (HD) TF with a known role in neural
development[79] that was not enriched in the motif analysis to test whether any
functional effects we saw were more generalisable to other homeodomain-containing
TFs.

Homeodomain TFs potentiate ASCL1-driven neuronal differentiation in mESCs

To investigate whether ASCL1 and HD TFs act cooperatively to drive
neurogenesis, we used a doxycycline-inducible piggyBac expression system[80] to
conditionally overexpress four candidate cofactors alone or together with iASCL1 in
mESCs and NE cells (Fig. S6C–E). We first asked whether co-expression could
activate previously unresponsive ASCL1 neuronal target genes in mESCs after 24
hours, focusing on *Zbtb18* and *Hdac11*, which are both associated with a proximal

ASCL1 binding site containing a homeodomain TF motif. Co-expression of PHOX2A, PHOX2B, or DLX3 with ASCL1 increased expression of both genes in NE cells and further potentiated the effect of ASCL1 alone (Fig. 6D–E and Fig. S6F), while LMX1A showed less consistent effects across targets (Fig. S6G). Critically, although neither *Zbtb18* nor *Hdac11* responds to iASCL1 alone in mESCs, co-expression with PHOX2A, PHOX2B, or DLX3 was sufficient to upregulate both genes. Co-activation also increased expression of *Map2*, a downstream neuronal differentiation marker and NE-specific iASCL1 target gene, in mESCs (Fig. 6D–E and Fig. S6F–G). These results demonstrate that selected HD TFs can enhance ASCL1 responsiveness at individual neuronal target genes in otherwise resistant mESCs.

To investigate how co-expression of these cofactors with ASCL1 affects transcriptional targets genome-wide in mESCs, we performed bulk RNA-seq on mESCs overexpressing PHOX2B with or without iASCL1 after 24 hours (Fig. 7A–B and Fig. S7A) (Table S6). Firstly, we saw that the number of upregulated genes in iASCL1+PHOX2B mESCs (n = 757) was greater than that observed in iASCL1 alone (n = 480) or PHOX2B alone (n = 240), indicating a compound effect of both factors together (Fig. S7A). We then looked at how the iASCL1+PHOX2B upregulated genes behaved across conditions (Fig. 7A). We saw that these genes fit broadly into two clusters: those that have highest expression in the iASCL1+PHOX2B-treated mESCs (Fig. 7A, Cluster 1) and those that have comparable expression in mESCs treated with either iASCL1 alone or iASCL1+PHOX2B (Fig. 7A, Cluster 2). This indicates that the combined activity of these two transcription factors creates a synergistic new set of target genes that are otherwise silent. GO analysis on both Cluster 1 (Fig. 7A') and Cluster 2 (Fig. 7A'') identified "regulation of neurogenesis", but "axon guidance" was only enriched in Cluster 1. This could indicate that the genes upregulated by combined activity of iASCL1 and PHOX2B are associated with a more mature or coherent neuronal transcriptional programme.

ASCL1 activation in ESCs results in upregulation of divergent gene programmes that may actively interfere with coherent neuronal programming. We therefore asked whether previously identified ASCL1 target genes upregulated specifically in mESCs (Fig. 1H–I) displayed an altered response to ASCL1 when it was co-expressed with PHOX2B. (Only 497 of the previously defined 601 mESC-specific iASCL1 targets were detected in this RNA-seq experiment due to differences in library preparation and sequencing depth.) A subset of these genes (178 out of 497) showed less activation in mESCs when iASCL1 activation was combined with PHOX2B activation (Fig. 7B) than with iASCL1 alone, and these genes were associated with terms such as "regulation of hemopoiesis" and "T cell proliferation" (Fig. 7B'). Therefore, PHOX2B may work with ASCL1 both to facilitate the activation of neuronal target genes and also to suppress the activation of non-neuronal targets that interferes with successful differentiation in mESCs.

Finally, we assessed whether these transcriptional changes we observed in mESCs in response to iASCL1 and HD TFs translated into neuronal differentiation, assayed by TUBB3 expression and cell morphology after a longer 72-hour period compared to earlier analysis of more immediate iASCL1 effects (Fig. 1–5). In NE, which already express some endogenous PHOX2B and PHOX2A (Fig. 6A),

additional ectopic PHOX2B and PHOX2A both induced a modest increase in
neurogenesis compared to control. However, neither PHOX2B nor PHOX2A co-
expression with iASCL1 significantly increased the proportion of TUBB3-positive
cells relative to iASCL1 alone (PHOX2B: 3.0% versus 7.7%, $p > 0.05$; PHOX2A:
9.2% versus 12.9%, $p > 0.05$) (Fig. 7D and 7G and Fig. S7C–D). DLX3 and LMX1A
did not increase TUBB3-positive cell numbers in NE even when expressed alone or
with iASCL1 (Fig. S7E–F), highlighting differences in the intrinsic neurogenic
capacity of individual HD TFs.

In mESCs, iASCL1 activation alone produced only sparse, diffuse TUBB3
expression after 3 days (Fig. 7C). PHOX2B and PHOX2A expression did not result in
TUBB3-positive cells in the absence of iASCL1 (Fig. 7C and 7F; Fig. S7B and S7D).
However, strikingly, co-expression of PHOX2B or PHOX2A with iASCL1 produced
TUBB3-positive cells with immature neuronal morphology (Fig. 7C and 7E–F; S7B
and S7D). DLX3 and LMX1A failed to generate TUBB3-positive cells in mESCs with
or without iASCL1 (Fig. S7E–F), indicating that the ability to confer neuronal
competence on ASCL1 in mESCs is specific to PHOX2A and PHOX2B and not a
general feature of homeodomain-containing TFs.

DISCUSSION

Together, these data support a model in which ASCL1 is not an intrinsically
neuron-specific pioneer factor, but a highly context-dependent transcription factor
whose activity is constrained by both chromatin state and cofactor availability. Cell-
intrinsic changes during the exit from pluripotency are required for competence for
ASCL1-mediated neuronal differentiation and to channel ASCL1 activity away from
activating divergent programmes. However, this neuronal competence can be
increased by altering the global chromatin landscape or by providing additional
cofactor TFs that enhance neuronal target gene activation.

Many studies have used mESCs (and human iPSCs) for forward
programming and TF-mediated neuronal differentiation. However, it is important to
note that many of them incorporate embryoid body formation, or some other priming
step.[7,9] Vainorius et al. use ASCL1 to drive neuronal differentiation from mESCs, but
with two important differences: firstly, they cultured their mESCs in serum-LIF, in
which ESCs are heterogeneous and display epigenetic and transcriptional priming of
many lineage commitment genes[57,81]; secondly, cells were plated on Matrigel and
differentiated in N2B27 after doxycycline induction of ASCL1 expression. Thus, these
cells have likely already transitioned through formative pluripotency and are
competent for ASCL1-mediated differentiation. Casey et al. profiled genome-wide
binding of ASCL1 in mESCs grown on feeder cells, and observed a similar binding
profile of ASCL1 to our mESC data, but did not investigate competence for neuronal
differentiation or the expression of non-neuronal genes.[46] We find that while failing to
upregulate a coherent neuronal programme, pluripotent mESCs are nevertheless
fully competent to respond transcriptionally to ectopic ASCL1, but in this context non-
neuronal heterologous gene programmes dominate; exit from pluripotency is
required to allow an appropriate ASCL1-specific developmental programme to be
engaged. This underscores the importance of chromatin remodelling to restrict the
cell identity during the early stages of embryonic development.[76,82,83]

Despite the fact that pluripotent cells have the ability to differentiate into any
embryonic cell type and are characterised by having globally low levels of repressive
heterochromatin,[12,73] it is perhaps surprising to find that chromatin accessibility
played a major role in determining ASCL1 binding in ESCs. Rather than a model with
widespread chromatin accessibility in pluripotent cells offering a "blank canvas" for
highly permissive binding of transcription factors, with an epigenetic environment
progressively refined during development, our data point to complex but distinct
landscapes. Unlike accessible sites bound by ASCL1 in NEs, sites accessible to
ASCL1 binding in ESCs are generally not associated with neuronal genes. Moreover,
in the context of gene activation in ESCs, EpiLCs and NEs, we see a limited ability of
ASCL1 to act as a pioneer opening chromatin. Recent data in human iPSC
differentiation to neurons, has shown ASCL1 displays different modes of activity at
different sites across the genome, sometimes acting alone and sometimes recruiting
SWI/SNF chromatin remodelling complexes to act as a non-classical pioneer
factor.[35] Our study adds to this model, showing that ASCL1 acts as both a pioneer
and a non-pioneer TF to varying degrees across the genome and between cell
types. Moreover, there must be additional layers of regulation that determine whether

a given binding site is accessible to ASCL1-mediated pioneering. Differential
availability of chromatin remodelling complexes across the developmental time
course may determine whether a given site can be pioneered by ASCL1, as has
been described for SWI/SNF subunit switching in neuronal differentiation.[35,84]

Accessibility, though clearly important, is not the only factor that determines
context-dependent ASCL1 binding and gene activation. Many ASCL1 binding sites
near neuroectoderm-specific target genes were marked by H3K27ac in neural
progenitors but not in mESCs or EpiLCs, even when accessibility is comparable at
these sites across cell types. This is especially interesting in light of the prevailing
view of mESCs being broadly unrestricted in terms of their epigenetic landscape and
developmental potential. Instead, ASCL1 binding sites that are activated in NE are
maintained in inactive chromatin in mESC and EpiLC and are not poised or primed.
This likely reflects the fact that they are not the first regulatory elements recruited *in*
vivo in the exit from pluripotency, as ASCL1 itself is expressed at later stages of
embryo development.[85,86] Consistent with this, global elevation of histone acetylation
using an HDAC inhibitor renders many previously unresponsive neuronal genes
inducible by ASCL1 in mESCs, highlighting the importance of chromatin priming in
establishing developmental competence. Even transient treatment with HDAC
inhibition can cause significant changes in genome architecture in mESCs,[87] and our
transcriptomic data indicates that TSA treatment creates many additional non-
neuronal ASCL1 target genes. HDAC inhibitors have been used to potentiate
ASCL1-mediated reprogramming in the adult retina,[88] but it is possible that additional
epigenetic constraints in these tissues limit the ectopic effects of induced ASCL1
activity.

To identify more targeted approaches to modulating ASCL1 activity in mESCs,
we identified candidate co-factors by identifying specific motifs for neuroepithelial
transcription factors found near to ASCL1 binding sites in NE target genes. We found
that co-expression of ASCL1 with candidate homeobox TFs in mESCs allows
activation of many previously NE-specific neuronal genes while simultaneously
inactivating genes associated with divergent pathways. Interaction between HD TFs
and bHLH factors, including ASCL1, has been previously shown in the diversification
of cell types in the developing retina[89] cranial neural crest cells.[90] PHOX2A and
PHOX2B have been implicated in gene regulatory networks with ASCL1 in the
development of the central nervous system,[91–93] sympathetic neurons,[94,95] and in
neuroblastoma.[96,97] ASCL1 expression typically precedes and is required for PHOX2
factor expression in these cases, so it is notable that we show simultaneous
expression of both factors potentiated the direct programming capacity of ASCL1 in
mESCs, where PHOX2 factors cannot be upregulated by ASCL1 activation directly.
This suggests that the sequential logic of endogenous development can be
bypassed by simultaneous cofactor delivery.

It is also possible that cell type-specific activity of ASCL1 is the result of
differential post-translational modification. Phosphorylation, for example, has been
shown to modulate the activity of various bHLH TFs in development and
reprogramming.[98–102] However, our recent work showed that a phosphorylation-
resistant ASCL1 mutant was not sufficient to drive neuronal differentiation in

mESCs,[10] so this is unlikely to explain why only NE cells undergo ASCL1-mediated
neuronal differentiation in these experiments.

In summary, we demonstrate that despite their ability to differentiate into any
embryonic cell type, the competence of ES cells to respond to the lineage-
determining transcription factor ASCL1 is limited by a chromatin landscape in which
non-neuronal enhancers are favourable for ASCL1 binding, which instead directs
ASCL1 activity to divergent genes and pathways. Our findings illustrate the
multifaceted controls that limit reprogramming in different cellular contexts and
provide a framework for improving directed differentiation and reprogramming
approaches by targeting competence rather than transcription factor expression
alone.

FIGURE LEGENDS

Figure 1 – Inducible ASCL1 drives divergent and non-neuronal transcriptional programmes in mESC and EpiLC

A) Schematic of mouse embryonic stem cells with inducible ASCL1 (iASCL1 mESC).

B) Differentiation timeline and sample collection for IF (D–E) and RNA-seq (F–L). "+ ASCL1" indicates 5 µM dexamethasone treatment.

C) Phase-contrast images of mESC, EpiLC, and NE at day 2 without ASCL1, with schematic of in vivo approximates.

D) IF showing mESC, EpiLC, and NE ± iASCL1 for 24 h (top) or 48 h (bottom). TUBB3 (green) marks neuronal differentiation. Scale bar = 100 µm. Representative of 4 experiments.

E) IF of iASCL1 mESC ± 24 h iASCL1 stained for NANOG (yellow) and OCT4 (red) and percentage of positive cells (n=3; 8–10 images per experiment). Scale bar = 100 µm. * $p < 0.05$, t-test.

F) PCA of top 1000 variable genes across conditions; grey points denote Dex-treated control line with no transgene.

G) Number of differentially expressed genes (DEGs) at 6 h and 24 h relative to 0 h in each cell type (mESC blue, EpiLC yellow, NE pink; positive upregulated, negative downregulated).

H) Venn diagram of overlap between ASCL1-upregulated DEGs across cell types after 24 h iASCL1.

I) Heatmap of z-scored expression for DEGs upregulated by ASCL1 in at least one cell type; clusters grouped by cell-type specificity with selected genes indicated.

J–L) Top 5 GO terms for upregulated genes after 24 h ASCL1 in mESC (J), EpiLC (K), and NE (L).

Figure 2 – Context-dependent ASCL1 genomic occupancy underlies divergent transcriptional responses across developmental states

A) Top 10,000 ASCL1 binding sites from anti-HA tag ChIP-seq from each cell type (mESC, EpiLC, NE) collated to give 17,210 unique binding sites. ASCL1 binding sites are clustered by cell type-specific binding patterns. Mean RLE-normalised ChIP-seq signal from 3 or 4 replicates.

B–C) Examples of NE-specific target genes from RNA-seq (B and C) and NE-specific ASCL1 binding sites (B' and C'), which overlap with validated enhancers from the VISTA dataset (B" and C"). *In situ* hybridisation from VISTA database shows enhancer activity in neural tube and brain at E11.5.

D) Categorising ASCL1 binding sites linked to NE-specific genes based on whether
they are exclusively bound in NE (left, n = 198) or bound in at least one other cell
type (right, n = 455). (D') Venn diagram for these categories. Example IGV tracks of
ASCL1 ChIP-seq for each category: *Wnt5a*, *Plxna2*, *Inppl1*.

E) ChIP-seq signal at sites in each of the categories in (D). One-way ANOVA with
repeated measures, Benjamini-Hochberg corrected pairwise comparisons: **** $p <$
0.0001.

676 Figure 3 – Chromatin accessibility dictates ASCL1 binding, with pioneer
677 activity restricted to a minority of neuronal target sites

A) Schematic for ATAC-seq experiment. Samples were exposed to either no ASCL1,
6 hrs, or 24 hrs before collection at the same endpoint.

B) PCA for ATAC-seq samples

C) Heatmap of ASCL1 ChIP-seq (black, left) and ATAC-seq signal with No ASCL1
(green, right) clustered by cell type-specific binding patterns defined in Fig. 2A. Row
order is preserved across the two heatmaps. Heatmaps centred on ASCL1 peak ± 1
kb.

D) Heatmaps of ASCL1 ChIP-seq (black) and ATAC-seq (No ASCL1) (green) for 653
ASCL1 sites linked to NE-specific DEGs defined in Fig. 2D. Peaks are clustered by
whether they are open (top, n = 493) or closed in NE (No ASCL1) (bottom, n = 160).
Heatmaps centred on ASCL1 peak ± 1 kb.

E) Heatmap of ATAC-seq (24hr ASCL1) for same 653 ASCL1 sites linked to NE-
specific DEGs with row order preserved from (D). Heatmaps centred on ASCL1 peak
± 1 kb.

F) Donut plots to show percentage of 653 ASCL1 binding sites in D and E that are
open (green) or closed (grey) in No ASCL1 ATAC-seq (top) or 24 hr ASCL1 ATAC-
seq (bottom).

G) IGV tracks of ASCL1 binding sites that are open in NE (No ASCL1) ATAC-seq.

H) IGV tracks of ASCL1 binding sites that are closed in NE (No ASCL1) ATAC-seq
but become open in NE (24 hr ASCL1) ATAC-seq.

699 Figure 4 – Histone acetylation status at shared accessible sites predicts
700 productive ASCL1 binding and target gene activation

A) Heatmaps of ASCL1 ChIP-seq (black) and No ASCL1 ATAC-seq (green) from
mESC at the 653 ASCL1 binding sites associated with NE-specific ASCL1 target
genes defined in Fig. 3D. Heatmaps centred on ASCL1 peak ± 1 kb.

B) Donut plot of percentage of the 653 sites that are open (green) or closed (grey) in
mESC (No ASCL1) ATAC-seq data.

C) Heatmaps of ASCL1 ChIP-seq (black) and ATAC-seq (No ASCL1) (green) from
mESC for the 362 out of 653 sites that are open in mESC (No ASCL1) ATAC-seq
data. Peaks are clustered based on whether are (C') bound in mESC (bottom, n =
186) or (C) not bound in mESC (top, n = 176). Heatmaps are centred on ASCL1
peak ± 2 kb.

D) H3K27ac ChIP-seq signal at 176 ASCL1 binding sites in (C); D') H3K27ac ChIP-
seq signal at 186 ASCL1 binding sites in (C'). H3K27ac data is shown for mESC
(blue) and AntNPC (pink) from Cruz-Molina et al., 2017[20]. Profiles are centred
around ASCL1 binding sites ± 5 kb.

E) Normalised RNA-seq counts for representative NE-specific ASCL1 target genes
(*Ank2*, *Cyth1*, *Hes5*) in mESC (blue) and NE (pink) exposed to 0, 6, or 24 hours of
iASCL1 activity. "C" (grey) represents Dex-treated control line with no iASCL1
transgene. N = 4.

F–G) IGV tracks to show ASCL1 binding sites near the NE-specific target genes in
(D) that are bound (F) or not bound (G) by ASCL1 in mESCs. mESC H3K27ac
(blue), mESC No ASCL1 ATAC-seq (green), mESC ASCL1 ChIP-seq (black); NE
(AntNPC) H3K27ac (blue), NE No ASCL1 ATAC-seq (green), NE ASCL1 ChIP-seq
(black). Rectangles below tracks indicate a called peak. Peaks are highlighted in
pink boxes.

Figure 5 – Global enhancement of histone acetylation partially unlocks
neuronal gene expression by ASCL1 but fails to redirect differentiation in
mESCs

A) Schematic of HDAC inhibition (HDACi) and acetylation.

B) Schematic of culture and treatment of mESC (blue) and NE (pink) with iASCL1
and/or 10 nM TSA for 24 hr (RNA and IF) or 48 hr (IF).

C) PCA plot for RNA-seq samples with mESC (blue) and NE (pink) treated with 24
hours iASCL1 (darker) or not (lighter). Simultaneously, samples were either treated
with DMSO (circle) or 10 nM TSA (triangle).

D) Examples of previously identified NE-specific ASCL1-respsonsive genes that
become ASCL1-responsive in mESC with ASCL1 + TSA treatment. D) *Nkx6-1*; D')
Disp2; D'') *Plxna2*. Adjusted p-values from DESeq2 differential gene expression
analysis. ns p > 0.05, ** p < 0.01, *** p < 0.001

E) Heatmap of z-scored RNA-seq counts for genes that were identified as NE-
specific ASCL1 targets in Fig. 1H–I and have associated ASCL1 binding site (n =
208). Expression data from mESC: control, iASCL1 only, TSA only, iASCL1+TSA. E')

genes of the 208 in the heatmap that are significantly upregulated in the
comparisons listed: iASCL1+TSA versus control (n = 116); TSA only versus control
(n = 102); iASCL1+TSA versus TSA only (n = 13).

F) Heatmap of z-scored RNA-seq counts for genes that are significantly upregulated
in TSA+iASCL1 versus TSA only (n = 458). F') Gene ontology analysis for all genes
in heatmap (F). Top 6 GO terms are listed. "Regulation of neurogenesis" highlighted
with black arrow.

G) IF images of H3K27ac (magenta) and TUBB3-positive neurites (green) after 48
hours ±ASCL1 and ±TSA in mESC or NE. DAPI counterstain (blue). Scale bar = 100
µm.

753   Figure 6 – Homeodomain transcription factor motifs are enriched in NE-
754   specific ASCL1 binding sites and co-expression with ASCL1 enables activation
755   of previously NE-specific target genes

A) Plot of TFs scored by fold enrichment of motif in NE-specific peaks (x) against log
fold change in expression comparing NE versus mESC (No iASCL1) from RNA-seq
(Fig. 1). Coloured by TF family. Phox2b, Phox2a, and Dlx3 are highlighted as
shortlisted candidates.

B) Schematic for putative model of cooperativity between ASCL1 and Homeodomain
(HD) TFs.

C) Schematic for culture and treatment of mESC (blue) and NE (pink) for 24 hrs:
control (light), +iASCL1 only (dark), +cofactor (Dox induction) only,
+iASCL1+cofactor

D–E) qPCR of NE-specific ASCL1 DEGs associated with ASCL1 binding sites that
contain HD motif (*Hdac11* and *Zbtb18*) and NE-specific marker of neurogenesis
(*Map2*). mESC (blue) and NE (pink). Dox used to induce overexpression of
cofactors, either PHOX2A (D) or PHOX2B (E). Control, iASCL1 alone, Dox alone, or
iASCL1 and Dox. N = 4 experiments, each performed in technical duplicate. Two-
way ANOVA with Dunnett's correction for multiple comparisons to the control for
each cell type.

773   Figure 7 – Genome-wide redirection of ASCL1 transcriptional output with
774   PHOX2B co-expression

A) Heatmap of z-scored RNA-seq counts for genes that are significantly upregulated
by iASCL1+PHOX2B relative to untreated control mESCs (n = 757). Genes are
clustered into 2 groups: Cluster 1 shows highest enrichment in iASCL1+PHOX2B
cells (n = 403); Cluster 2 shows comparable expression in iASCL1 alone and

iASCL1+PHOX2B. A') GO analysis for genes in Cluster 1. A'') GO analysis for genes
in Cluster 2. Top 6 terms are shown.

B) Heatmap of z-scored RNA-seq counts for genes that were previously identified as
mESC-specific iASCL1 targets (Fig. 1H) (n = 497). Genes are clustered using k
means = 3. B') GO analysis of genes in Cluster 1. Top 6 terms are shown.

C–D) Schematic for culture and treatment of mESC (C, blue) and NE (D, pink)
before collection for immunofluorescence (IF) imaging. Cells were either untreated,
iASCL1 only, PHOX2B only, or iASCL1+PHOX2B. TUBB3 (green), mCherry
(magenta), DAPI (blue). mCherry and DAPI LUTs are adjusted separately for mESC
and NE, but TUBB3 is consistent across conditions. Scale bar = 100 µm.
Representative images from 2 experiments each done in duplicate.

E) Magnified inset from Fig. 7C. White arrow indicates neurite-like outgrowth. Scale
bar = 100 µm.

F–G) Quantification of TUBB3-positive cells in cofactor differentiation assay in mESC
and NE untreated, with ASCL1 alone, Dox alone, or ASCL1 and Dox. Each dot
represents an image taken from a randomised position in the well. Statistics are
performed using N = 4 for 4 wells, from 2 independent passages. Wilcoxon pairwise
test with Benjamini–Hochberg post-hoc correction. * p < 0.05

SUPPLEMENTARY FIGURE LEGENDS

Figure S1 – Related to Figure 1

A) iASCL1 mESC express OCT4 (green) and NANOG (red) in 2iLIF. Scale bar = 50 µm.

B) qPCR of ASCL1 target genes ± 24h Dex in iASCL1 mESC in 2iLIF. N = 3 experiments. ** p < 0.01, t-test.

C) Western blot to show HA-ASCL1-GR is specific to iASCL1 and levels are consistent across differentiation conditions (mESC, EpiLC, and NE). Representative of 2 western blots. Molecular weight markers are indicated (right).

D) IF of OCT4 (yellow), SOX1 (red) and PAX6 (magenta) in iASCL1 differentiated into NE conditions on day 3-5. Spontaneous neuronal differentiation from day 4 onwards indicated by TUBB3 (green). Scale bar = 100 µm. Representative of 2 experiments.

E) IF of NANOG (blue), OTX2 (yellow) and SOX1 (magenta), day 0-2 of NE differentiation.

F) IF for naïve pluripotency marker, NANOG (yellow), and formative markers, OCT6 (green) and OTX2 (magenta) in iASCL1 cells on days 1-3 of EpiLC differentiation. Scale bar = 100 µm.

G) qPCR of ASCL1 target genes in 2iLIF variant media. N = 3 independent experiments. Two-way ANOVA with Dunnett's post-hoc pairwise comparisons comparing control versus +iASCL1 for 24 hours.

H) EdU-positive mESCs (yellow) with or without 24 hours of iASCL1 activity. Quantification of positive cells from 8 images from 3 independent experiments. **** p < 0.0001, t-test.

I–K) Volcano plots to show upregulated genes in mESC (I), EpiLC (J) and NE (K) after 24 hours of iASCL1 relative to untreated cells in each of: mESC (blue), EpiLC (yellow), NE (pink). Note: some genes appear above threshold for significance but are coloured grey and excluded from analysis based on comparison between untreated cells and Dex-treated control cells with no iASCL1-GR transgene to control for Dex-specific effects.

L) Venn diagram to show overlap in DEGs after 6 hours.

M–O) Volcano plots and GO analysis on upregulated genes in mESC (M), EpiLC (N) and NE (O) after 24 hours of iASCL1 relative to untreated cells in each of: mESC (blue), EpiLC (yellow), NE (pink).

Figure S2 – Related to Figure 2

A) ChIP-qPCR to validate the anti-HA ASCL1 ChIP-seq. N = 4 independent experiments.

B) Correlation matrix of ChIP-seq peak sets, including all iASCL1 samples and control cells that have no iASCL1-GR transgene.

C) PCA plot to show sample distances after normalisation (using RLE method and reads in peaks as library size).

D) Heatmap across replicates after normalisation at the top 10,000 peaks in the consensus set.

E) Annotation of top 10,000 ASCL1 binding sites in each cell type. Promoter regions are defined as ±1kb from TSS.

F) Top motif discovered by STREME in each of the clusters. E-value and percentage of target sites that contain the motif are reported next to each motif.

G) Number of differentially expressed genes (DEGs) in mESC (blue), EpiLC (yellow) and NE (pink) that have a binding site within ±100 kb. DEGs with no proximal ASCL1 binding site are grey.

H) Correlation matrix for publicly available ASCL1 ChIP-seq datasets, including mESC, EpiLC and NE from this study. Full GEO accession numbers are available in the Methods section.

I) Profiles of ChIP-seq signal in mESC, EpiLC, and NE from this study at peaks from neural stem cells (NSC),[43] and embryonic neural tube.[72]

Figure S3 – Related to Fig.3

A) Number of peaks called in each sample. Asterisk corresponds to the per-sample consensus peak set, comprising peaks called in at least 3 of the 4 replicates.

B) Fraction of reads in peaks (FRiP) for all samples.

C) TSS enrichment is the ratio of reads the align to TSS of known genes over the reads in 1000 bp regions flanking the TSS.

D) Percentage of reads that align to mitochondrial DNA per sample.

E) iASCL1 ChIP-seq (black) and No iASCL1 ATAC-seq (green) data in mESC at all 10,000 ASCL1 binding sites detected in mESC, split by whether they are open (top) or closed (bottom) in the No iASCL1 ATAC-seq data.

F) iASCL1 ChIP-seq (black) and No iASCL1 ATAC-seq (green) data in EpiLC at all 10,000 ASCL1 binding sites detected in EpiLC, split by whether they are open (top) or closed (bottom) in the No iASCL1 ATAC-seq data.

G) iASCL1 ChIP-seq (black) and No iASCL1 ATAC-seq (green) data in NE at all
10,000 iASCL1 binding sites detected in NE, split by whether they are open (top) or
closed (bottom) in the No iASCL1 ATAC-seq data.

H) Donut plots for percentage of iASCL1 ChIP-seq peaks in cell type-specific
clusters (Fig. 2A) that overlap with an ATAC-seq peak (green) or not (grey) in each
cell type.

I) RNA-seq counts for NE-specific ASCL1-responsive DEGs highlight in Fig.3. As per
RNA-seq experiment, samples were either treated with 0, 6, or 24 hours of iASCL1
and compared to a control line with no iASCL1 transgene for each cell type.

Figure S4 – Related to Fig.4

A) ASCL1 ChIP-seq as in Fig. 2 and 3.

B) H3K27ac signal in mESC and AntNPC from Cruz-Molina et al., 2017[20] plotted
across ASCL1 binding sites (this study) split by cell type-specific clusters (Fig. 2A).
Centred around ASCL1 binding site (500 bp) ± 5 kb.

C) H3K4me1 signal in mESC and EpiLC from Bleckwehl et al., 2021[76] plotted across
ASCL1 binding sites (this study) split by cell type-specific clusters (Fig. 2A). Centred
around ASCL1 binding site (500 bp) ± 5 kb. C') highlights the NE-specific iASCL1
binding sites.

Figure S5 – Related to Figure 5

A) Phase contrast to show morphology and density of mESC treated with nothing,
+iASCL1 only, 10 nM TSA only, or +iASCL1 and +TSA.

B) PCA plot of PC2 and PC3 from RNA-seq.

C–F) Volcano plot to show differentially expressed genes in mESC in specific
comparisons: C) iASCL1 versus control; D) TSA only versus control; E) iASCL1+TSA
versus control; F) iASCL1+TSA versus TSA only. Significance thresholds: absolute
log fold change > 0 and adjusted p-value < 0.01. ashr shrunken log fold changes are
used for plotting.[103]

Figure S6 – Related to Figure 6

898   A) Plot of TFs scored by fold enrichment of motif in mESC-specific peaks (x) against
899   log fold change in expression in mESC versus NE at t = 0. Coloured by TF family.

900   B) Number of ASCL1 binding sites that contain a DLX3 or PHOX2B motif.

C) piggyBac construct used for Dox-inducible overexpression. ITR, inverted terminal repeat; cHS4, chicken hypersensitive site 4 insulator; TRE3G, third-generation tetracycline-responsive promoter; COFACTOR is where candidate cofactors were cloned in; T2A, self-cleaving peptide; mCherry; pA, polyadenylation signal; pEF1⏑ promoter; rtTA Tet-On activator; PuroR, puromycin resistance gene.

D) Titration of Dox treatment on pool of antibiotic-resistant cells. mCherry (red) indicates induction of the transgene.

E) qPCR of *Phox2b* transgene induction relative to *Tbp* housekeeping. N = 4 experiments

F–G) qPCR of NE-specific ASCL1 DEGs associated with ASCL1 binding sites that contain HD motif (*Hdac11* and *Zbtb18*) or *Map2*. Dox used to induce overexpression of either DLX3 (F) or LMX1A (G). Cells were untreated or treated with either iASCL1 alone, Dox alone, or iASCL1 and Dox. N = 4 experiments, each performed in technical duplicate. Two-way ANOVA with Dunnett's correction for multiple comparisons to each cell type's respective control.

Figure S7 – Related to Figure 7

A) Volcano plots for specific comparisons in mESC from RNA-seq (Fig. 7A–B): iASCL1 versus control; PHOX2B only versus control; iASCL1+PHOX2B versus control. Significance thresholds: absolute log fold change > 1 and adjusted p-value < 0.01. ashr shrunken log fold changes are used for plotting.[103]

B–C) Schematic for culture and treatment of mESC (B, blue) and NE (C, pink) before collection for immunofluorescence (IF) imaging. Cells were either untreated, iASCL1 only, PHOX2A only, or iASCL1+PHOX2A. TUBB3 (green), mCherry (magenta), DAPI (blue). mCherry and DAPI LUTs are adjusted separately for mESC and NE, but TUBB3 is consistent across conditions. Scale bar = 100 µm. Representative images from 2 experiments each done in duplicate.

D–F) Quantification of TUBB3-positive cells in cofactor differentiation assay in mESC and NE. Cofactors are PHOX2A (D), DLX3 (E) and LMX1A (F). Each dot represents an image taken from a randomised position in the well. Statistics are performed using N = 4 for 4 wells, from 2 independent passages. Wilcoxon pairwise test with Benjamini–Hochberg post-hoc correction. * p < 0.05

METHODS

Cell culture

Mouse embryonic stem cells (mESCs; ES-E14TG2a, ATCC #CRL-1821) were maintained as described previously.[104] Ascl1-GR-HA mESCs (iASCL1; R. Azzarelli, unpublished) were cultured in 2i/LIF medium: N2B27 basal medium (1:1 Neurobasal [Gibco, 21103-049] / DMEM-F12 [Gibco, D6421], N2 [in-house, Cambridge Stem Cell Institute], B-27 [Gibco, 17504044], 2 mM L-glutamine, 50 µM β-mercaptoethanol) supplemented with 100 U/mL leukaemia inhibitory factor (LIF; in-house), 3 µM CHIR-99021 (ABCR, AB253776) and 1 µM PD0325901 (ABCR, AB253775).

Cells were passaged every 2–3 days using Accutase (300 × g, 3 min) and seeded at 1.5 × 10⬜ cells/cm² on 0.1% gelatin-coated 6-well plates. Stocks were cryopreserved at 1 × 10⬜ cells/mL in 50% 2i/LIF / 40% KSR / 10% DMSO. Mycoplasma testing was performed every 3–6 months.

For lineage transitions, mESCs were plated on fibronectin (16.7 µg/mL) to generate epiblast-like cells (EpiLC) in N2B27 + FGF2 (12 ng/mL) + Activin A (20 ng/mL),[3,5] or on laminin (10 µg/mL) for primitive neuroectoderm (NE) differentiation in N2B27 + 0.1% sodium bicarbonate + 0.11% BSA + 20 µg/mL insulin.[104]

A gene fragment encoding N-terminal HA-tagged full-length ASCL1 fused to the glucocorticoid receptor ligand-binding domain (GR-LBD), followed by an IRES-GFP cassette, was synthesised with flanking attB1/attB2 sites (Twist Bioscience). Gateway LR Clonase II (Thermo Fisher) was used to insert the construct into a Rosa26-targeting destination vector (SP202) carrying appropriate homology arms.[105] E14 mESCs were co-transfected with the targeting vector, Rosa26-directed sgRNAs, and a Cas9 expression vector using the 4D Nucleofector (Lonza). Correctly targeted clones were isolated by FACS for constitutive GFP expression, expanded, and verified by PCR across both homology arms for monoallelic (heterozygous) insertion. All experiments used a single validated clonal line.

In iASCL1 mESCs, ASCL1 activity is post-translationally controlled: the HA-ASCL1-GR fusion protein is constitutively expressed but held inactive by HSP90 chaperone binding until displacement by 5 µM dexamethasone (Dex).[106]

Cofactor overexpression mESCs

PiggyBac-based, doxycycline (Dox)-inducible overexpression vectors were assembled from two published plasmids: the TRE3G-driven expression cassette was PCR-amplified (CloneAmp, Takara, 639298) from pPB-Ins-TRE3Gp-KRAB-dCas9-ecDHFR-IRES-GFP-EF1Ap-Puro (Addgene #183410), and the rtTA-Neo cassette was amplified from pPB-Ins-U6p-sgRNAentry-EF1Ap-TetOn3G-IRES-Neo (Addgene #183411). Both were cloned into a custom piggyBac-based vector (VB240515-1632gka, VectorBuilder) via EcoRV/PacI digestion and In-Fusion Snap Assembly (Takara, 638948). The dCas9-KRAB-U6-sgRNA module was excised with NotI/AsiSI and replaced with synthesised transgene sequences (PHOX2A, PHOX2B, LMX1A, and DLX3: Twist Bioscience; mCherry: VectorBuilder). Gel purification used the

Monarch kit (NEB, T1120); plasmids were propagated in Stellar competent cells
(Takara, 636763), purified with QIAprep Miniprep or Midiprep kits (Qiagen), and
sequence-verified by Nanopore sequencing (Plasmidsaurus).

For PHOX2A, PHOX2B, LMX1A, and DLX3 overexpression lines, mESCs were co-
transfected with PB-Dox-Cofactor plasmid and Super PiggyBac Transposase vector
(Stratech, PB210PA-1-SBI; 500 ng) to a total of 2 μg DNA, complexed with 3.5 μL
Lipofectamine 2000 (Invitrogen, 11668027) in 400 μL Opti-MEM™ and incubated on
cells for 4 h at 37°C. Medium was replaced with 2i/LIF, then selection in 0.5 μg/mL
puromycin (2i/LIF) initiated 48–72 h post-transfection and maintained until colonies
were pooled. Overexpression efficiency was confirmed by RT-qPCR after 1 μM Dox
induction.

Immunofluorescence microscopy

Cells were fixed in 4% PFA (ThermoFisher, J19943-K2) for 20 min at room
temperature, permeabilised in 0.2% Triton-X100 (Sigma, X100)/PBS for 15 min, and
blocked in 5% serum / 0.01% Triton-X100 / PBS for 1 h. Primary antibodies (PBS +
2% serum + 0.01% Triton-X100) were applied overnight at 4°C; secondary
antibodies for 1 h at room temperature. Washes used PBS-T (PBS + 0.01% Triton-
X100); DAPI (1 μg/mL, Abcam, ab228549) was added during the third wash (20
min). Images were acquired on a Leica DMI6000 at 10× or 20×.

S-phase fraction was used as a proliferation proxy. Cells were treated with 5 μM Dex
for 24 h; then 10 μM EdU (final; Click-iT EdU Kit, Alexa Fluor 488, Thermo Fisher,
C10637) was added for 1 h before fixation. EdU was detected per manufacturer's
instructions. Note: iASCL1 mESC constitutive GFP was negligible relative to
AlexaFluor-488-labelled targets, so the 488 channel was used without interference.

Image analysis

Nuclear intensities were quantified in CellProfiler v4.0.[107] Nuclei were segmented
from DAPI using the IdentifyPrimaryObjects module (Otsu two-class adaptive
thresholding; smoothing factors determined empirically) and filtered by size. Channel
intensities within each nuclear object were measured; binary positivity thresholds
were set empirically. Percentages of positive/negative nuclei were calculated from
≥10 images across 2–3 biological replicates.

For TUBB3 quantification (Fig. 6), nuclear ROIs were expanded by 5 pixels to
capture peri-nuclear signal, which was used as a proxy for total TUBB3 (individual
cell-level quantification was not feasible in these morphologically complex
monolayers). Intensities were normalised to the control sample mean; statistical
comparisons used per-replicate means (N = 4, two separate passages, each
performed in duplicate), not individual nucleus counts.

RT-qPCR

RNA was extracted with the RNeasy Mini Kit (Qiagen, 74106) and stored at –80°C.
cDNA was synthesised with the QuantiTect Reverse Transcription Kit (Qiagen,
205314), including DNase treatment; cDNA was diluted 1:5–1:10 prior to qPCR.
Reactions (10 μL, 384-well) were run on a QuantStudio 7 Flex or ViiA 7 Real-Time

PCR System. Primer efficiency was validated by serial dilution of a pooled
mESC/EpiLC/NE cDNA reference; primers with an efficiency outside 85–115% were
excluded. Quantification used the modified ΔΔCt method incorporating primer
efficiency.[108] TBP and GAPDH served as housekeeping genes. Expression ratios
were log-transformed prior to statistical testing to satisfy normality.

Bulk RNA-seq

RNA was extracted as above and treated with the DNA-free DNA Removal Kit
(Invitrogen, AM1906). Poly(A) RNA was enriched using the NEBNext Poly(A) mRNA
Magnetic Isolation Module (E7490); strand-specific libraries were prepared with the
NEBNext Ultra II Directional RNA Library Prep Kit (E7760) and NEBNext Multiplex
Oligos (E6440). Sequencing was performed as 50 bp paired-end reads on NovaSeq
6000 or NovaSeq X (Illumina).

For cofactor overexpression experiments, RNA-seq was performed by
Plasmidsaurus (Illumina) using their in-house analysis pipeline; raw counts matrices
were used for downstream analysis. Cells (~10□ per condition) were collected in 50
µL DNA/RNA Shield (Zymo, R1100-50) after Accutase dissociation (300 × g, 3 min).

Computational analysis of RNA-seq

Raw BCL files were converted to FASTQ with bcl2fastq (Illumina). Adapter trimming
and quality filtering (Phred ≥ 33) used Trim Galore; trimmed reads were pseudo-
aligned to the mouse genome (mm39) with Kallisto.[109]

Kallisto output was imported into R with tximport (DESeq2 package).[110] Genes with
fewer than 10 counts in fewer than 3 samples were removed. Differential expression
analysis used DESeq2; DEGs (Fig. 1) were defined as |log□FC| > 0.585 (≥1.5-fold)
and adjusted p < 0.01. Fold change thresholds were > 0 for HDACi RNA-seq (Fig. 5)
and > 1 for cofactor experiment (Fig. 7) to account for different variance within these
datasets. Variance-stabilised transformation was used for PCA, distance matrix, and
heatmap visualisation.[110] ashr was used for shrinkage estimation in volcano plots.[103]
Gene Ontology enrichment used enrichGO (clusterProfiler).[111]

ChIP-seq

ChIP-seq was performed as described[112] with modifications. Cells were crosslinked
in 1% formaldehyde (10 min, room temperature), quenched with 125 mM glycine,
washed twice in ice-cold PBS, and pelleted (2,000 × g, 4 min). Pellets were flash-
frozen on dry ice and stored at –80°C. Sequential lysis used: LB1 (50 mM HEPES-
KOH pH 7.5, 140 mM NaCl, 1 mM EDTA, 10% glycerol, 0.5% NP-40, 0.25% Triton-
X100; 10 min, ice, 2,000 × g), LB2 (10 mM Tris-HCl pH 8, 200 mM NaCl, 1 mM
EDTA, 0.5 mM EGTA; 10 min, ice, 2,000 × g), and LB3 (10 mM Tris-HCl pH 8, 100
mM NaCl, 1 mM EDTA, 0.5 mM EGTA, 0.1% Na-deoxycholate, 0.5% N-
lauroylsarcosine). All buffers contained cOmplete protease inhibitor (Sigma,
4693159001).

Chromatin was sheared using a Bioruptor Plus (Diagenode, 4°C; 30 s on/off, 16
cycles in 2 rounds of 8) to yield 100–600 bp fragments (verified by agarose gel after
heat reversal). Chromatin concentration was determined by A280 on a NanoDrop (2

µL in 98 µL 0.1 M NaOH). Per IP, 100 µg chromatin was diluted to 300 µL in LB3 +
1% Triton-X100 and clarified (16,000 × g, 10 min); 10% was retained as input. IP
was performed overnight with anti-HA antibody (ab9110) pre-loaded on BSA-blocked
Protein G Dynabeads (Invitrogen). Beads were washed 10× in RIPA buffer (50 mM
HEPES-KOH, 500 mM LiCl, 1 mM EDTA, 1% Igepal CA-630, 0.7% Na-deoxycholate)
and once in TBS (20 mM Tris-HCl pH 7.6, 150 mM NaCl). DNA was eluted overnight
at 65°C in 200 µL elution buffer (50 mM Tris-HCl pH 7.6, 10 mM EDTA, 1% SDS),
which simultaneously reversed crosslinks.

Eluted DNA was diluted 1:1 in TE, then purified by sequential RNase A digestion (40
µg/mL, 37°C, 30 min), proteinase K digestion (400 µg/mL, 55°C, 2 h), phenol-
chloroform-isoamyl alcohol phase separation in MaXtract HD tubes (Qiagen), and
ethanol precipitation (200 mM NaCl, –20°C) with 1 µL GlycoBlue (20 µg/µL, Thermo
Fisher, AM9515) as co-precipitant. Purified DNA was resuspended in 20 µL 10 mM
Tris-HCl pH 8. Enrichment was validated by qPCR at known ASCL1 target loci
(percentage input from 10% fraction). Libraries were prepared using ThruPLEX
DNA-Seq (Takara, R400675) at the JCBC NGS Facility and sequenced on NovaSeq
6000 at 15 nM equimolar pooling.

Computational analysis of ChIP-seq data

Reads were trimmed with Trim Galore (Phred ≥ 33) and aligned to a composite
mm39/dm6 index with Bowtie2. *Drosophila* reads (spike-in) were discarded as
normalisation by spike-in was not applied; spike-ins were included as a
precautionary measure, but RLE normalisation was sufficient for comparison
between conditions. mm39-aligned reads were used for all downstream analysis.
Peaks were called with MACS3[113] using matched input as background control.
Specificity was confirmed by the near-absence of peaks in control (no-transgene)
samples; any such peaks were excluded from the consensus set. The consensus
peak set comprised peaks present in ≥2 of 4 replicates per condition. Reads-in-
peaks normalisation used RLE (DESeq2). Peaks were annotated to transcription
start sites within 100 kb using AnnotationHub[114] and Ensembl *Mus musculus* v108.

Published ENCODE mouse ASCL1 ChIP-seq datasets were downloaded via
accession numbers (Materials Table) and converted from mm10 to mm39 using
UCSC liftOver. [115] Inter-dataset Pearson correlation and Jaccard similarity were
assessed from a binary peak-presence matrix in DiffBind.

Motif analysis

1093 De novo motif discovery was performed using MEME-ChIP (MEME Suite, web
1094 server),[116] which applies STREME for improved sensitivity at centrally enriched
1095 motifs,[117] and independently with HOMER[118] (-size 150, -mask) for both de novo and
1096 known-motif enrichment.

ATAC-seq

1098 ATAC-seq was performed using the Omni-ATAC protocol.[119,120] To synchronise multi-
1099 timepoint collection, cells were seeded on sequential days. Cells were dissociated
1100 with Accutase, counted with Trypan Blue, and 5 × 10□ cells resuspended in ice-cold

PBS were pelleted and lysed in 50 µL RSB++ (10 mM Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl□, 0.1% NP-40, 0.1% Tween-20, 0.01% digitonin) for 3 min on ice. Nuclei were washed with 1 mL RSB+ (RSB++ without NP-40/digitonin), pelleted (500 × g, 10 min, 4°C), and resuspended in 50 µL Tn5 transposition mix. Tagmentation was performed at 37°C for 30 min with 1,000 rpm mixing, then cleaned with Zymo DNA Clean & Concentrator (D4014) and stored at –20°C.

Tagmented DNA was PCR-amplified with indexed i5/i7 primers, purified with the Qiagen MinElute kit (eluted in 20 µL EB), and size-selected with a dual AMPure XP bead (#A63880) strategy: (i) 0.55× beads added to remove large fragments (supernatant retained), then (ii) 1.5× beads to capture the library; beads washed 2× in 80% ethanol and eluted in 20 µL EB. Fragment size distribution was confirmed by Bioanalyzer HS D1000 (Agilent, #5067-5584). Libraries were pooled equimolarly and sequenced on NovaSeq X 25B (PE100), targeting ~9 × 10□ reads/sample.

Computational analysis of ATAC-seq

FASTQ files were generated with bcl2fastq2. A custom bash pipeline based on nf-core ATAC-seq[121,122] and published best practices[123] was applied. Quality control used FastQC followed by Trim Galore (default paired-end parameters). Reads were aligned to mm39 with Bowtie2[124] (--very-sensitive-local, --no-mixed, --no-discordant; maximum insert size 2,000 bp). SAMtools[125] was used to index, sort, and assess alignment statistics; mitochondrial reads were removed. Duplicates were marked with Picard and removed; insert size metrics were collected with Picard CollectInsertSizeMetrics. Peaks were called with MACS3[113] in BAMPE mode (effective genome size 2,468,088,461).

Downstream analysis used R v4.5.0 in RStudio. Peaks overlapping ENCODE blacklisted regions[126] or unplaced/random contigs were removed. DiffBind generated a consensus peak set (peaks present in ≥3 of 4 replicates per condition); reads-in-peaks were TMM-normalised with DiffBind. Bigwig files were produced with bamCoverage (deepTools)[127] using TMM scale factors; per-condition averages used bamCompare. Visualisation used IGV[128] and deepTools heatmaps.

ACKNOWLEDGEMENTS

We would like to thank the Cambridge Stem Cell Institute Genomics Facility at the Jeffrey Cheah Biomedical Centre for all of their help with library preparation and sequencing; the Cambridge Stem Cell Institute Imaging Core Facility; Prof. Brian Hendrich and members of his lab group for their assistance with ChIP-seq and ATAC-seq; William Beckman, Lidiya Mykhaylechko, Sarah Gillen, Frances Connor, and all members of the Philpott lab for helpful discussions throughout; and Dr Mekayla Storer, Prof. Jason Carroll and Dr Srinjan Basu for their useful guidance.

This work was supported by a Wellcome Trust Investigator award (212253/Z/18/Z) to (A.P., J.L.-B., R.D., J.-P.N.-B.); J.L.-B. is funded by a Wellcome Trust studentship (218481/Z/19/Z); R.A. is supported by core funding from UCL School of Pharmacy and by the Royal Society grant (RGS\R2\242202).

1143

AUTHOR CONTRIBUTIONS

1144
J.L.-B. performed most of the experimental work and bioinformatic analyses. R.A.
generated the iASCL1 mESC line with contribution from R.D. J.L.-B., R.D. and J.-
P.N.-B generated the piggyBac overexpression system for cofactor experiments.
R.A. and J.L.-B. designed the experiments. J.L.-B. prepared the figures with input
from R.A and A.P. R.A. and A.P. supervised the project and obtained funding. J.L.-
B., R.A. and A.P wrote the manuscript with input from R.D. and J.-P.N.-B.

1151

SUPPLEMENTAL MATERIAL

1152
Table S1 (differentially expressed genes per condition, relating to Fig. 1 and S1),
Table S2 (GO analysis, relating to Fig. 1 and S1), Table S3 (ASCL1 peaks linked to
proximal genes, relating to Fig. 2), Table S4 (653 NE-specific ASCL1 binding sites,
relating to Fig. 3 and Fig. 4), Table S5 (differentially expressed genes with HDACi
treatment, relating to Fig. 5), Table S6 (differentially expressed genes with PHOX2B
and ASCL1, relating to Fig. 7)

1159

RESOURCE AVAILABILITY

1160

Lead contact

1161
1162 Further information and requests for resources and reagents should be directed to
1163 and will be fulfilled by the lead contacts, Dr Roberta Azzarelli
1164 roberta.azzarelli@ucl.ac.uk or Prof. Anna Philpott ap113@cam.ac.uk.

Materials availability

1165
1166 All unique/stable reagents and cell lines generated in this study are available from
1167 the lead contact with a completed materials transfer agreement.

Data and code availability

1168
Bulk RNA-seq, ChIP-seq and ATAC-seq data have been deposited at GEO
(GSE327328) and are publicly available as of the date of publication. Published
ASCL1 ChIP-seq data were accessed from: GSM2576033, GSM1175113, E-MTAB-
2384, E-MTAB-2384, GSE43916, GSE43916, GSM1347006, GSE89209.
Microscopy data reported in this paper will be shared by the lead contact upon
request. All original code is available via GitHub (github.com/Philpott-
lab/ascl1_competence). Any additional information required to reanalyse the data
reported in this work is available from the lead contact upon request.

1177

DECLARATION OF GENERATIVE AI AND AI-ASSISTED TECHNOLOGIES IN THE WRITING PROCESS

1178
1179

During the preparation of this work the author(s) used Claude v1.2773.0 for proofreading of the manuscript and to assess R and bash code reproducibility. After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the published article.

REFERENCES

1. Rostovskaya, M., Stirparo, G.G., and Smith, A. (2019). Capacitation of human naïve pluripotent stem cells for multi-lineage differentiation. Development *146*, dev172916. https://doi.org/10.1242/dev.172916.

2. Kinoshita, M., Barber, M., Mansfield, W., Cui, Y., Spindlow, D., Stirparo, G.G., Dietmann, S., Nichols, J., and Smith, A. (2021). Capture of Mouse and Human Stem Cells with Features of Formative Pluripotency. Cell Stem Cell *28*, 453-471.e8. https://doi.org/10.1016/j.stem.2020.11.005.

3. Buecker, C., Srinivasan, R., Wu, Z., Calo, E., Acampora, D., Faial, T., Simeone, A., Tan, M., Swigut, T., and Wysocka, J. (2014). Reorganization of Enhancer Patterns in Transition from Naive to Primed Pluripotency. Cell Stem Cell *14*, 838–853. https://doi.org/10.1016/j.stem.2014.04.003.

4. Carbognin, E., Carlini, V., Panariello, F., Chieregato, M., Guerzoni, E., Benvegnù, D., Perrera, V., Malucelli, C., Cesana, M., Grimaldi, A., et al. (2023). Esrrb guides naive pluripotent cells through the formative transcriptional programme. Nat. Cell Biol. *25*, 643–657. https://doi.org/10.1038/s41556-023-01131-x.

5. Hayashi, K., Ohta, H., Kurimoto, K., Aramaki, S., and Saitou, M. (2011). Reconstitution of the Mouse Germ Cell Specification Pathway in Culture by Pluripotent Stem Cells. Cell *146*, 519–532. https://doi.org/10.1016/j.cell.2011.06.052.

6. Nishimura, K., Nitta, T., Doi, K., and Takata, K. (2022). Rapid Conversion of Human Induced Pluripotent Stem Cells into Dopaminergic Neurons by Inducible Expression of Two Transcription Factors. Stem Cells Dev. *31*, 269–277. https://doi.org/10.1089/scd.2021.0363.

7. Vainorius, G., Novatchkova, M., Michlits, G., Baar, J.C., Raupach, C., Lee, J., Yelagandula, R., Wernig, M., and Elling, U. (2023). Ascl1 and Ngn2 convert mouse embryonic stem cells to neurons via functionally distinct paths. Nat. Commun. *14*, 5341. https://doi.org/10.1038/s41467-023-40803-y.

8. Thoma, E.C., Wischmeyer, E., Offen, N., Maurus, K., Sirén, A.-L., Schartl, M., and Wagner, T.U. (2012). Ectopic Expression of Neurogenin 2 Alone is Sufficient to Induce Differentiation of Embryonic Stem Cells into Mature Neurons. PLOS ONE *7*, e38651. https://doi.org/10.1371/journal.pone.0038651.

9. Aydin, B., Kakumanu, A., Rossillo, M., Moreno-Estellés, M., Garipler, G., Ringstad, N., Flames, N., Mahony, S., and Mazzoni, E.O. (2019). Proneural factors Ascl1 and Neurog2 contribute to neuronal subtype identities by establishing distinct chromatin landscapes. Nat. Neurosci. *22*, 897–908. https://doi.org/10.1038/s41593-019-0399-y.

10. Azzarelli, R., Gillen, S., Connor, F., Lundie-Brown, J., Puletti, F., Drummond, R., Raffaelli, A., and Philpott, A. (2024). Phospho-regulation of ASCL1-mediated chromatin opening during cellular reprogramming. Development *151*, dev204329. https://doi.org/10.1242/dev.204329.

11. Srivastava, D., and Mahony, S. (2020). Sequence and chromatin determinants of
transcription factor binding and the establishment of cell type-specific binding patterns.
Biochim. Biophys. Acta BBA - Gene Regul. Mech. *1863*, 194443.
https://doi.org/10.1016/j.bbagrm.2019.194443.

12. Meshorer, E., and Misteli, T. (2006). Chromatin in pluripotent embryonic stem cells and
differentiation. Nat. Rev. Mol. Cell Biol. *7*, 540–546. https://doi.org/10.1038/nrm1938.

13. Peng, T., Zhai, Y., Atlasi, Y., ter Huurne, M., Marks, H., Stunnenberg, H.G., and
Megchelenbrink, W. (2020). STARR-seq identifies active, chromatin-masked, and
dormant enhancers in pluripotent mouse embryonic stem cells. Genome Biol. *21*, 243.
https://doi.org/10.1186/s13059-020-02156-3.

14. Sankar, A., Mohammad, F., Sundaramurthy, A.K., Wang, H., Lerdrup, M., Tatar, T., and
Helin, K. (2022). Histone editing elucidates the functional roles of H3K27 methylation
and acetylation in mammals. Nat. Genet. *54*, 754–760. https://doi.org/10.1038/s41588-
022-01091-2.

15. Voss, T.C., and Hager, G.L. (2014). Dynamic regulation of transcriptional states by
chromatin and transcription factors. Nat. Rev. Genet. *15*, 69–81.
https://doi.org/10.1038/nrg3623.

16. Jonasson, M.E., and Buecker, C. (2025). The exit from naive pluripotency: a platform for
the study of enhancer mechanistics. Biochem. Soc. Trans. *53*, 1–15.
https://doi.org/10.1042/BST20253037.

17. Dupont, C., Chahar, D., Trullo, A., Gostan, T., Surcis, C., Grimaud, C., Fisher, D., Feil,
R., and Llères, D. (2023). Evidence for low nanocompaction of heterochromatin in living
embryonic stem cells. EMBO J. *42*, EMBJ2021110286.
https://doi.org/10.15252/embj.2021110286.

18. Creyghton, M.P., Cheng, A.W., Welstead, G.G., Kooistra, T., Carey, B.W., Steine, E.J.,
Hanna, J., Lodato, M.A., Frampton, G.M., Sharp, P.A., et al. (2010). Histone H3K27ac
separates active from poised enhancers and predicts developmental state. Proc. Natl.
Acad. Sci. *107*, 21931–21936. https://doi.org/10.1073/pnas.1016071107.

19. Crispatzu, G., Rehimi, R., Pachano, T., Bleckwehl, T., Cruz-Molina, S., Xiao, C., Mahabir,
E., Bazzi, H., and Rada-Iglesias, A. (2021). The chromatin, topological and regulatory
properties of pluripotency-associated poised enhancers are conserved in vivo. Nat.
Commun. *12*, 4344. https://doi.org/10.1038/s41467-021-24641-4.

20. Cruz-Molina, S., Respuela, P., Tebartz, C., Kolovos, P., Nikolic, M., Fueyo, R., Ijcken,
W.F.J. van, Grosveld, F., Frommolt, P., Bazzi, H., et al. (2017). PRC2 Facilitates the
Regulatory Topology Required for Poised Enhancer Function during Pluripotent Stem
Cell Differentiation. Cell Stem Cell *20*, 689-705.e9.
https://doi.org/10.1016/j.stem.2017.02.004.

21. Rugg-Gunn, P.J., Cox, B.J., Ralston, A., and Rossant, J. (2010). Distinct histone
modifications in stem cell lines and tissue lineages from the early mouse embryo. Proc.
Natl. Acad. Sci. *107*, 10783–10790. https://doi.org/10.1073/pnas.0914507107.

22. Kim, S., and Wysocka, J. (2023). Deciphering the multi-scale, quantitative cis-regulatory
code. Mol. Cell *83*, 373–392. https://doi.org/10.1016/j.molcel.2022.12.032.

23. Ong, C.-T., and Corces, V.G. (2011). Enhancer function: new insights into the regulation
of tissue-specific gene expression. Nat. Rev. Genet. *12*, 283–293.
https://doi.org/10.1038/nrg2957.

24. Ong, C.-T., and Corces, V.G. (2012). Enhancers: emerging roles in cell fate specification.
EMBO Rep. *13*, 423–430. https://doi.org/10.1038/embor.2012.52.

25. Iwafuchi-Doi, M., and Zaret, K.S. (2016). Cell fate control by pioneer transcription
factors. Development *143*, 1833–1837. https://doi.org/10.1242/dev.133900.

26. Zaret, K.S., and Mango, S.E. (2016). Pioneer transcription factors, chromatin dynamics,
and cell fate control. Curr. Opin. Genet. Dev. *37*, 76–81.
https://doi.org/10.1016/J.GDE.2015.12.003.

27. Tapscott, S.J., Davis, R.L., Thayer, M.J., Cheng, P.-F., Weintraub, H., and Lassar, A.B.
(1988). MyoD1: A Nuclear Phosphoprotein Requiring a Myc Homology Region to
Convert Fibroblasts to Myoblasts. Science *242*, 405–411.
https://doi.org/10.1126/science.3175662.

28. Davis, R.L., Weintraub, H., and Lassar, A.B. (1987). Expression of a single transfected
cDNA converts fibroblasts to myoblasts. Cell *51*, 987–1000.
https://doi.org/10.1016/0092-8674(87)90585-X.

29. Zaret, K.S., and Carroll, J.S. (2011). Pioneer transcription factors: establishing
competence for gene expression. Genes Dev. *25*, 2227–2241.
https://doi.org/10.1101/GAD.176826.111.

30. Freund, M.M., Harrison, M.M., and Torres-Zelada, E.F. (2024). Exploring the reciprocity
between pioneer factors and development. Development *151*, dev201921.
https://doi.org/10.1242/dev.201921.

31. Hansen, J.L., Loell, K.J., and Cohen, B.A. (2022). A test of the pioneer factor hypothesis
using ectopic liver gene activation. eLife *11*, e73358. https://doi.org/10.7554/eLife.73358.

32. Donaghey, J., Thakurela, S., Charlton, J., Chen, J.S., Smith, Z.D., Gu, H., Pop, R.,
Clement, K., Stamenova, E.K., Karnik, R., et al. (2018). Genetic determinants and
epigenetic effects of pioneer-factor occupancy. Nat. Genet. *50*, 250–258.
https://doi.org/10.1038/s41588-017-0034-3.

33. Cernilogar, F.M., Hasenöder, S., Wang, Z., Scheibner, K., Burtscher, I., Sterr, M.,
Smialowski, P., Groh, S., Evenroed, I.M., Gilfillan, G.D., et al. (2019). Pre-marked
chromatin and transcription factor co-binding shape the pioneering activity of Foxa2.
Nucleic Acids Res. *47*, 9069–9086. https://doi.org/10.1093/nar/gkz627.

34. Mayran, A., Sochodolsky, K., Khetchoumian, K., Harris, J., Gauthier, Y., Bemmo, A.,
Balsalobre, A., and Drouin, J. (2019). Pioneer and nonpioneer factor cooperation drives
lineage specific chromatin opening. Nat. Commun. *10*, 3807.
https://doi.org/10.1038/s41467-019-11791-9.

35. Păun, O., Tan, Y.X., Patel, H., Strohbuecker, S., Ghanate, A., Cobolli-Gigli, C., Sopena,
M.L., Gerontogianni, L., Goldstone, R., Ang, S.-L., et al. (2023). Pioneer factor ASCL1
cooperates with the mSWI/SNF complex at distal regulatory elements to regulate human
neural differentiation. Genes Dev. *37*, 218–242. https://doi.org/10.1101/gad.350269.122.

36. Guillemot, F. (2007). Spatial and temporal specification of neural fates by transcription
factor codes. Development *134*, 3771–3780. https://doi.org/10.1242/DEV.006379.

37. Mazzoni, E.O., Mahony, S., Closser, M., Morrison, C.A., Nedelec, S., Williams, D.J., An,
D., Gifford, D.K., and Wichterle, H. (2013). Synergistic binding of transcription factors to
cell-specific enhancers programs motor neuron identity. Nat. Neurosci. *16*, 1219–1227.
https://doi.org/10.1038/nn.3467.

38. Spitz, F., and Furlong, E.E.M. (2012). Transcription factors: from enhancer binding to
developmental control. Nat. Rev. Genet. *13*, 613–626. https://doi.org/10.1038/nrg3207.

39. Morgunova, E., and Taipale, J. (2017). Structural perspective of cooperative transcription
factor binding. Curr. Opin. Struct. Biol. *47*, 1–8. https://doi.org/10.1016/j.sbi.2017.03.006.

40. Wapinski, O.L., Vierbuchen, T., Qu, K., Lee, Q.Y., Chanda, S., Fuentes, D.R., Giresi,
P.G., Ng, Y.H., Marro, S., Neff, N.F., et al. (2013). Hierarchical mechanisms for
transcription factor-mediated reprogramming of fibroblasts to neurons. Cell *155*, 621.
https://doi.org/10.1016/J.CELL.2013.09.028.

41. Wapinski, O.L., Lee, Q.Y., Chen, A.C., Li, R., Corces, M.R., Ang, C.E., Treutlein, B.,
Xiang, C., Baubet, V., Suchy, F.P., et al. (2017). Rapid Chromatin Switch in the Direct
Reprogramming of Fibroblasts to Neurons. Cell Rep. *20*, 3236–3247.
https://doi.org/10.1016/j.celrep.2017.09.011.

42. Castro, D.S., Martynoga, B., Parras, C., Ramesh, V., Pacary, E., Johnston, C., Drechsel,
D., Lebel-Potter, M., Garcia, L.G., Hunt, C., et al. (2011). A novel function of the
proneural factor Ascl1 in progenitor proliferation identified by genome-wide
characterization of its targets. Genes Dev. *25*, 930–945.
https://doi.org/10.1101/gad.627811.

43. Raposo, A.A.S.F., Vasconcelos, F.F., Drechsel, D., Marie, C., Johnston, C., Dolle, D.,
Bithell, A., Gillotin, S., van den Berg, D.L.C., Ettwiller, L., et al. (2015). Ascl1
Coordinately Regulates Gene Expression and the Chromatin Landscape during
Neurogenesis. Cell Rep. *10*, 1544–1556. https://doi.org/10.1016/j.celrep.2015.02.025.

44. Kagawa, W., and Kurumizaka, H. (2021). Structural basis for DNA sequence recognition
by pioneer factors in nucleosomes. Curr. Opin. Struct. Biol. *71*, 59–64.
https://doi.org/10.1016/j.sbi.2021.05.011.

45. Lundie-Brown, J., Puletti, F., Philpott, A., and Azzarelli, R. (2025). Cell fate acquisition
and reprogramming by the proneural transcription factor ASCL1. Open Biol. *15*, 250018.
https://doi.org/10.1098/rsob.250018.

46. Casey, B.H., Kollipara, R.K., Pozo, K., and Johnson, J.E. (2018). Intrinsic DNA binding
properties demonstrated for lineage-specifying basic helix-loop-helix transcription
factors. Genome Res. *28*, 484–496. https://doi.org/10.1101/GR.224360.117/-/DC1.

47. Talikka, M., Perez, S.E., and Zimmerman, K. (2002). Distinct Patterns of Downstream
Target Activation Are Specified by the Helix–Loop–Helix Domain of Proneural Basic
Helix–Loop–Helix Transcription Factors. Dev. Biol. *247*, 137–148.
https://doi.org/10.1006/DBIO.2002.0677.

48. Mazurier, N., Parain, K., Parlier, D., Pretto, S., Hamdache, J., Vernier, P., Locker, M.,
Bellefroid, E., and Perron, M. (2014). Ascl1 as a Novel Player in the Ptf1a Transcriptional

Network for GABAergic Cell Specification in the Retina. PLoS ONE *9*, e92113.
https://doi.org/10.1371/journal.pone.0092113.

49. Gammill, L.S., and Sive, H. (1997). Identification of otx2 target genes and restrictions in
ectodermal competence during Xenopus cement gland formation. Development *124*,
471–481. https://doi.org/10.1242/dev.124.2.471.

50. Meijsing, S.H., Elbi, C., Luecke, H.F., Hager, G.L., and Yamamoto, K.R. (2007). The
Ligand Binding Domain Controls Glucocorticoid Receptor Dynamics Independent of
Ligand Release. Mol. Cell. Biol. *27*, 2442. https://doi.org/10.1128/MCB.01570-06.

51. Ying, Q.-L., Wray, J., Nichols, J., Batlle-Morera, L., Doble, B., Woodgett, J., Cohen, P.,
and Smith, A. (2008). The ground state of embryonic stem cell self-renewal. Nature *453*,
519–523. https://doi.org/10.1038/nature06968.

52. Ying, Q.-L., Stavridis, M., Griffiths, D., Li, M., and Smith, A. (2003). Conversion of
embryonic stem cells into neuroectodermal precursors in adherent monoculture. Nat.
Biotechnol. *21*, 183–186. https://doi.org/10.1038/nbt780.

53. Ying, Q.-L., and Smith, A.G. (2003). Defined Conditions for Neural Commitment and
Differentiation. In Methods in Enzymology Differentiation of Embryonic Stem Cells.
(Academic Press), pp. 327–341. https://doi.org/10.1016/S0076-6879(03)65023-8.

54. Wood, H.B., and Episkopou, V. (1999). Comparative expression of the mouse *Sox1*,
Sox2 and *Sox3* genes from pre-gastrulation to early somite stages. Mech. Dev. *86*, 197–
201. https://doi.org/10.1016/S0925-4773(99)00116-1.

55. Zhao, S., Nichols, J., Smith, A.G., and Li, M. (2004). SoxB transcription factors specify
neuroectodermal lineage choice in ES cells. Mol. Cell. Neurosci. *27*, 332–342.
https://doi.org/10.1016/j.mcn.2004.08.002.

56. Gouti, M., Tsakiridis, A., Wymeersch, F.J., Huang, Y., Kleinjung, J., Wilson, V., and
Briscoe, J. (2014). In Vitro Generation of Neuromesodermal Progenitors Reveals Distinct
Roles for Wnt Signalling in the Specification of Spinal Cord and Paraxial Mesoderm
Identity. PLOS Biol. *12*, e1001937. https://doi.org/10.1371/journal.pbio.1001937.

57. Smith, A. (2017). Formative pluripotency: the executive phase in a developmental
continuum. Development *144*, 365–373. https://doi.org/10.1242/dev.142679.

58. Evans, M.J., and Kaufman, M.H. (1981). Establishment in culture of pluripotential cells
from mouse embryos. Nature *292*, 154–156. https://doi.org/10.1038/292154a0.

59. Martin, G.R. (1981). Isolation of a pluripotent cell line from early mouse embryos cultured
in medium conditioned by teratocarcinoma stem cells. Proc. Natl. Acad. Sci. *78*, 7634–
7638. https://doi.org/10.1073/pnas.78.12.7634.

60. Hayashi, K., Lopes, S.M.C. de S., Tang, F., and Surani, M.A. (2008). Dynamic
Equilibrium and Heterogeneity of Mouse Pluripotent Stem Cells with Distinct Functional
and Epigenetic States. Cell Stem Cell *3*, 391–401.
https://doi.org/10.1016/j.stem.2008.07.027.

61. Beckman, W.F., Parkinson, L.M., Chaytor, L., and Philpott, A. (2025). Distinct proliferative
and neuronal programmes of chromatin binding and gene activation by ASCL1 are cell
cycle stage-specific. Dev. Camb. Engl. *152*, dev204816.
https://doi.org/10.1242/dev.204816.

62. Vue, T.Y., Kollipara, R.K., Borromeo, M.D., Smith, T., Mashimo, T., Burns, D.K., Bachoo,
R.M., and Johnson, J.E. (2020). ASCL1 regulates neurodevelopmental transcription
factors and cell cycle genes in brain tumors of glioma mouse models. Glia *68*, 2613.
https://doi.org/10.1002/GLIA.23873.

63. Castro, D.S., Martynoga, B., Parras, C., Ramesh, V., Pacary, E., Johnston, C., Drechsel,
D., Lebel-Potter, M., Garcia, L.G., Hunt, C., et al. (2011). A novel function of the
proneural factor Ascl1 in progenitor proliferation identified by genome-wide
characterization of its targets. Genes Dev. *25*, 930–945.
https://doi.org/10.1101/gad.627811.

64. Urbán, N., Berg, D.L.C. van den, Forget, A., Andersen, J., Demmers, J.A.A., Hunt, C.,
Ayrault, O., and Guillemot, F. (2016). Return to quiescence of mouse neural stem cells
by degradation of a proactivation protein. Science *353*, 292–295.
https://doi.org/10.1126/science.aaf4802.

65. Durdu, S., Iskar, M., Isbel, L., Hoerner, L., Wirbelauer, C., Burger, L., Hess, D.,
Iesmantavicius, V., and Schübeler, D. (2025). Chromatin-dependent motif syntax defines
differentiation trajectories. Mol. Cell *85*, 2900-2918.e16.
https://doi.org/10.1016/j.molcel.2025.07.005.

66. Wang, S., Sun, H., Ma, J., Zang, C., Wang, C., Wang, J., Tang, Q., Meyer, C.A., Zhang,
Y., and Liu, X.S. (2013). Target analysis by integration of transcriptome and ChIP-seq
data with BETA. Nat. Protoc. *8*, 2502–2515. https://doi.org/10.1038/nprot.2013.150.

67. Maienschein-Cline, M., Zhou, J., White, K.P., Sciammas, R., and Dinner, A.R. (2012).
Discovering transcription factor regulatory targets using gene expression and binding
data. Bioinformatics *28*, 206–213. https://doi.org/10.1093/bioinformatics/btr628.

68. Kosicki, M., Baltoumas, F.A., Kelman, G., Boverhof, J., Ong, Y., Cook, L.E., Dickel, D.E.,
Pavlopoulos, G.A., Pennacchio, L.A., and Visel, A. (2025). VISTA Enhancer browser: an
updated database of tissue-specific developmental enhancers. Nucleic Acids Res. *53*,
D324–D330. https://doi.org/10.1093/nar/gkae940.

69. Li, L., Lai, F., Hu, X., Liu, B., Lu, X., Lin, Z., Liu, L., Xiang, Y., Frum, T., Halbisen, M.A., et
al. (2023). Multifaceted SOX2-chromatin interaction underpins pluripotency progression
in early embryos. Science *382*, eadi5516. https://doi.org/10.1126/science.adi5516.

70. Webb, A.E., Pollina, E.A., Vierbuchen, T., Urbán, N., Ucar, D., Leeman, D.S., Martynoga,
B., Sewak, M., Rando, T.A., Guillemot, F., et al. (2013). FOXO3 Shares Common Targets
with ASCL1 Genome-wide and Inhibits ASCL1-Dependent Neurogenesis. Cell Rep. *4*,
477–491. https://doi.org/10.1016/j.celrep.2013.06.035.

71. Wang, H., Keepers, B., Qian, Y., Xie, Y., Colon, M., Liu, J., and Qian, L. (2022). Cross-
lineage potential of Ascl1 uncovered by comparing diverse reprogramming regulatomes.
Cell Stem Cell *29*, 1491-1504.e9. https://doi.org/10.1016/j.stem.2022.09.006.

72. Borromeo, M.D., Meredith, D.M., Castro, D.S., Chang, J.C., Tung, K.-C., Guillemot, F.,
and Johnson, J.E. (2014). A transcription factor network specifying inhibitory versus
excitatory neurons in the dorsal spinal cord. Development *141*, 2803–2812.
https://doi.org/10.1242/dev.105866.

73. Gaspar-Maia, A., Alajem, A., Meshorer, E., and Ramalho-Santos, M. (2011). Open
chromatin in pluripotency and reprogramming. Nat. Rev. Mol. Cell Biol. *12*, 36–47.
https://doi.org/10.1038/nrm3036.

74. Zhou, V.W., Goren, A., and Bernstein, B.E. (2011). Charting histone modifications and
the functional organization of mammalian genomes. Nat. Rev. Genet. *12*, 7–18.
https://doi.org/10.1038/nrg2905.

75. Rada-Iglesias, A., Bajpai, R., Swigut, T., Brugmann, S.A., Flynn, R.A., and Wysocka, J.
(2011). A unique chromatin signature uncovers early developmental enhancers in
humans. Nature *470*, 279–283. https://doi.org/10.1038/nature09692.

76. Bleckwehl, T., Crispatzu, G., Schaaf, K., Respuela, P., Bartusel, M., Benson, L., Clark,
S.J., Dorighi, K.M., Barral, A., Laugsch, M., et al. (2021). Enhancer-associated H3K4
methylation safeguards in vitro germline competence. Nat. Commun. *12*, 5771.
https://doi.org/10.1038/s41467-021-26065-6.

77. Bergsland, M., Werme, M., Malewicz, M., Perlmann, T., and Muhr, J. (2006). The
establishment of neuronal properties is controlled by Sox4 and Sox11. Genes Dev. *20*,
3475–3486. https://doi.org/10.1101/gad.403406.

78. Bergsland, M., Ramsköld, D., Zaouter, C., Klum, S., Sandberg, R., and Muhr, J. (2011).
Sequentially acting Sox transcription factors in neural lineage development. Genes Dev.
25, 2453–2464. https://doi.org/10.1101/gad.176008.111.

79. Doucet-Beaupré, H., Gilbert, C., Profes, M.S., Chabrat, A., Pacelli, C., Giguère, N.,
Rioux, V., Charest, J., Deng, Q., Laguna, A., et al. (2016). Lmx1a and Lmx1b regulate
mitochondrial functions and survival of adult midbrain dopaminergic neurons. Proc. Natl.
Acad. Sci. U. S. A. *113*, E4387-4396. https://doi.org/10.1073/pnas.1520387113.

80. Ng-Blichfeldt, J.-P., Drummond, R., Kausar, S., Lando, D., Barile, M., and Philpott, A.
(2026). Hierarchical TBX6-FOXC Regulatory Logic Controls Human Trunk Mesoderm
Diversification. Preprint at bioRxiv, https://doi.org/10.64898/2026.03.17.712295
https://doi.org/10.64898/2026.03.17.712295.

81. Marks, H., Kalkan, T., Menafra, R., Denissov, S., Jones, K., Hofemeister, H., Nichols, J.,
Kranz, A., Francis Stewart, A., Smith, A., et al. (2012). The Transcriptional and
Epigenomic Foundations of Ground State Pluripotency. Cell *149*, 590–604.
https://doi.org/10.1016/j.cell.2012.03.026.

82. Zhang, J., Donahue, G., Gilbert, M.B., Lapidot, T., Nicetto, D., and Zaret, K.S. (2024).
Distinct H3K9me3 heterochromatin maintenance dynamics govern different gene
programmes and repeats in pluripotent cells. Nat. Cell Biol. *26*, 2115–2128.
https://doi.org/10.1038/s41556-024-01547-z.

83. Todd, C.D., Ijaz, J., Torabi, F., Dovgusha, O., Bevan, S., Cracknell, O., Lohoff, T., Clark,
S., Argelaguet, R., Pierce, J., et al. (2025). Epigenetic priming of mammalian embryonic
enhancer elements coordinates developmental gene networks. Genome Biol. *26*, 214.
https://doi.org/10.1186/s13059-025-03658-8.

84. Keegan, S.E., Haskins, J., Simmonds, A.J., and Hughes, S.C. (2023). A chromatin
remodelling SWI/SNF subunit, Snr1, regulates neural stem cell determination and
differentiation. Dev. Camb. Engl. *150*, dev201484. https://doi.org/10.1242/dev.201484.

85. Guillemot, F., Lo, L.C., Johnson, J.E., Auerbach, A., Anderson, D.J., and Joyner, A.L.
(1993). Mammalian achaete-scute homolog 1 is required for the early development of
olfactory and autonomic neurons. Cell *75*, 463–476. https://doi.org/10.1016/0092-
8674(93)90381-Y.

86. Guillemot, F., and Joyner, A.L. (1993). Dynamic expression of the murine Achaete-Scute
homologue Mash-1 in the developing nervous system. Mech. Dev. *42*, 171–185.
https://doi.org/10.1016/0925-4773(93)90006-J.

87. Paldi, F., Szalay, M.-F., Dufau, S., Di Stefano, M., Reboul, H., Jost, D., Bantignies, F.,
and Cavalli, G. (2026). Transient histone deacetylase inhibition induces cellular memory
of gene expression and 3D genome folding. Nat. Genet. *58*, 404–417.
https://doi.org/10.1038/s41588-025-02489-4.

88. Jorstad, N.L., Wilken, M.S., Grimes, W.N., Wohl, S.G., Vandenbosch, L.S., Yoshimatsu,
T., Wong, R.O., Rieke, F., and Reh, T.A. (2017). Stimulation of functional neuronal
regeneration from Müller glia in adult mice. Nat. 2017 5487665 *548*, 103–107.
https://doi.org/10.1038/nature23283.

89. Hatakeyama, J., and Kageyama, R. (2004). Retinal cell fate determination and bHLH
factors. Semin. Cell Dev. Biol. *15*, 83–89. https://doi.org/10.1016/j.semcdb.2003.09.005.

90. Kim, S., Morgunova, E., Naqvi, S., Goovaerts, S., Bader, M., Koska, M., Popov, A.,
Luong, C., Pogson, A., Swigut, T., et al. (2024). DNA-guided transcription factor
cooperativity shapes face and limb mesenchyme. Cell *187*, 692-711.e26.
https://doi.org/10.1016/j.cell.2023.12.032.

91. Pattyn, A., Simplicio, N., van Doorninck, J.H., Goridis, C., Guillemot, F., and Brunet, J.-F.
(2004). Ascl1/Mash1 is required for the development of central serotonergic neurons.
Nat. Neurosci. *7*, 589–595. https://doi.org/10.1038/nn1247.

92. Pattyn, A., Guillemot, F., and Brunet, J.F. (2006). Delays in neuronal differentiation in
Mash1/Ascl1 mutants. Dev. Biol. *295*, 67–75.
https://doi.org/10.1016/J.YDBIO.2006.03.008.

93. Pattyn, A., Morin, X., Cremer, H., Goridis, C., and Brunet, J.-F. (1997). Expression and
interactions of the two closely related homeobox genes Phox2a and Phox2b during
neurogenesis. Development *124*, 4065–4075. https://doi.org/10.1242/dev.124.20.4065.

94. Stanke, M., Stubbusch, J., and Rohrer, H. (2004). Interaction of Mash1 and Phox2b in
sympathetic neuron development. Mol. Cell. Neurosci. *25*, 374–382.
https://doi.org/10.1016/j.mcn.2003.10.021.

95. Lo, L., Tiveron, M.-C., and Anderson, D.J. (1998). MASH1 activates expression of the
paired homeodomain transcription factor Phox2a, and couples pan-neuronal and
subtype-specific components of autonomic neuronal identity. Development *125*, 609–
620. https://doi.org/10.1242/dev.125.4.609.

96. Wang, L., Tan, T.K., Durbin, A.D., Zimmerman, M.W., Abraham, B.J., Tan, S.H., Ngoc,
P.C.T., Weichert-Leahey, N., Akahane, K., Lawton, L.N., et al. (2019). ASCL1 is a MYCN-
and LMO1-dependent member of the adrenergic neuroblastoma core regulatory circuitry.
Nat. Commun. *10*, 5622. https://doi.org/10.1038/s41467-019-13515-5.

97. Parkinson, L.M., Gillen, S.L., Woods, L.M., Chaytor, L., Marcos, D., Ali, F.R., Carroll,
J.S., and Philpott, A. (2022). The proneural transcription factor ASCL1 regulates cell
proliferation and primes for differentiation in neuroblastoma. Front. Cell Dev. Biol. *10*.

98. Ali, F.R., Cheng, K., Kirwan, P., Metcalfe, S., Livesey, F.J., Barker, R.A., and Philpott, A.
(2014). The phosphorylation status of Ascl1 is a key determinant of neuronal

differentiation and maturation in vivo and in vitro. Dev. Camb. Engl. *141*, 2216–2224.
https://doi.org/10.1242/DEV.106377.

99. Ali, F., Hindley, C., McDowell, G., Deibler, R., Jones, A., Kirschner, M., Guillemot, F., and
Philpott, A. (2011). Cell cycle-regulated multi-site phosphorylation of Neurogenin 2
coordinates cell cycling with differentiation during neurogenesis. Development *138*,
4267–4277. https://doi.org/10.1242/DEV.067900.

100.   Ali, F.R., Marcos, D., Chernukhin, I., Woods, L.M., Parkinson, L.M., Wylie, L.A.,
Papkovskaia, T.D., Davies, J.D., Carroll, J.S., and Philpott, A. (2020). Dephosphorylation
of the Proneural Transcription Factor ASCL1 Re-Engages a Latent Post-Mitotic
Differentiation Program in Neuroblastoma. Mol. Cancer Res. MCR *18*, 1759–1766.
https://doi.org/10.1158/1541-7786.MCR-20-0693.

101.   Azzarelli, R., Hurley, C., Sznurkowska, M.K., Rulands, S., Hardwick, L., Gamper, I.,
Ali, F., McCracken, L., Hindley, C., McDuff, F., et al. (2017). Multi-site Neurogenin3
Phosphorylation Controls Pancreatic Endocrine Differentiation. Dev. Cell *41*, 274-286.e5.
https://doi.org/10.1016/J.DEVCEL.2017.04.004.

102.   Pereira, A., Diwakar, J., Masserdotti, G., Beşkardeş, S., Simon, T., So, Y., Martín-
Loarte, L., Bergemann, F., Vasan, L., Schauer, T., et al. (2024). Direct neuronal
reprogramming of mouse astrocytes is associated with multiscale epigenome
remodeling and requires Yy1. Nat. Neurosci. *27*, 1260–1273.
https://doi.org/10.1038/s41593-024-01677-5.

103.   Stephens, M. (2017). False discovery rates: a new deal. Biostatistics *18*, 275–294.
https://doi.org/10.1093/biostatistics/kxw041.

104.   Mulas, C., Kalkan, T., von Meyenn, F., Leitch, H.G., Nichols, J., and Smith, A. (2019).
Defined conditions for propagation and manipulation of mouse embryonic stem cells.
Development *146*, dev173146. https://doi.org/10.1242/dev.173146.

105.   Bressan, R.B., Dewari, P.S., Kalantzaki, M., Gangoso, E., Matjusaitis, M., Garcia-
Diaz, C., Blin, C., Grant, V., Bulstrode, H., Gogolok, S., et al. (2017). Efficient
CRISPR/Cas9-assisted gene targeting enables rapid and precise genetic manipulation
of mammalian neural stem cells. Development *144*, 635–648.
https://doi.org/10.1242/dev.140855.

106.   Mattioni, T., Louvion, J.-F., and Picard, D. (1994). Chapter 16 Regulation of Protein
Activities by Fusion to Steroid Binding Domains. In Methods in Cell Biology Protein
Expression in Animal Cells., M. G. Roth, ed. (Academic Press), pp. 335–352.
https://doi.org/10.1016/S0091-679X(08)60611-1.

107.   Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A., and
Goodman, A. (2021). CellProfiler 4: improvements in speed, utility and usability. BMC
Bioinformatics *22*, 433. https://doi.org/10.1186/s12859-021-04344-9.

108.   Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time
RT–PCR. Nucleic Acids Res. *29*, e45. https://doi.org/10.1093/nar/29.9.e45.

109.   Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal
probabilistic RNA-seq quantification. Nat. Biotechnol. *34*, 525–527.
https://doi.org/10.1038/nbt.3519.

110. Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*, 550. https://doi.org/10.1186/s13059-014-0550-8.

111. Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., Zhou, L., Tang, W., Zhan, L., et al. (2021). clusterProfiler 4.0: A universal enrichment tool for interpreting omics data. The Innovation *2*. https://doi.org/10.1016/j.xinn.2021.100141.

112. Schmidt, D., Wilson, M.D., Spyrou, C., Brown, G.D., Hadfield, J., and Odom, D.T. (2009). ChIP-seq: using high-throughput sequencing to discover protein-DNA interactions. Methods San Diego Calif *48*, 240–248. https://doi.org/10.1016/j.ymeth.2009.03.001.

113. Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E., Nusbaum, C., Myers, R.M., Brown, M., Li, W., et al. (2008). Model-based analysis of ChIP-Seq (MACS). Genome Biol. *9*, R137. https://doi.org/10.1186/gb-2008-9-9-r137.

114. Morgan, M., and Shepherd, L. (2025). AnnotationHub: Client to access AnnotationHub resources https://doi.org/10.18129/B9.bioc.AnnotationHub.

115. Hinrichs, A.S., Karolchik, D., Baertsch, R., Barber, G.P., Bejerano, G., Clawson, H., Diekhans, M., Furey, T.S., Harte, R.A., Hsu, F., et al. (2006). The UCSC Genome Browser Database: update 2006. Nucleic Acids Res. *34*, D590–D598. https://doi.org/10.1093/nar/gkj144.

116. Bailey, T.L., Johnson, J., Grant, C.E., and Noble, W.S. (2015). The MEME Suite. Nucleic Acids Res. *43*, W39–W49. https://doi.org/10.1093/nar/gkv416.

117. Bailey, T.L. (2021). STREME: accurate and versatile sequence motif discovery. Bioinformatics *37*, 2834–2840. https://doi.org/10.1093/bioinformatics/btab203.

118. Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X., Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. Mol. Cell *38*, 576–589. https://doi.org/10.1016/j.molcel.2010.05.004.

119. Buenrostro, J.D., Wu, B., Chang, H.Y., and Greenleaf, W.J. (2015). ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. Curr. Protoc. Mol. Biol. *109*, 21.29.1-21.29.9. https://doi.org/10.1002/0471142727.mb2129s109.

120. Corces, M.R., Trevino, A.E., Hamilton, E.G., Greenside, P.G., Sinnott-Armstrong, N.A., Vesuna, S., Satpathy, A.T., Rubin, A.J., Montine, K.S., Wu, B., et al. (2017). An improved ATAC-seq protocol reduces background and enables interrogation of frozen tissues. Nat. Methods *14*, 959–962. https://doi.org/10.1038/nmeth.4396.

121. Ewels, P.A., Peltzer, A., Fillinger, S., Patel, H., Alneberg, J., Wilm, A., Garcia, M.U., Di Tommaso, P., and Nahnsen, S. (2020). The nf-core framework for community-curated bioinformatics pipelines. Nat. Biotechnol. *38*, 276–278. https://doi.org/10.1038/s41587-020-0439-x.

122. Patel, H., Espinosa-Carrasco, J., Langer, B., Ewels, P., Garcia, M.U., Syme, R., Peltzer, A., Talbot, A., Behrens, D., Gabernet, G., et al. (2023). nf-core/atacseq: [2.1.2] - 2022-08-07. (Zenodo). https://doi.org/10.5281/zenodo.8222875 https://doi.org/10.5281/zenodo.8222875.

1603   123.   Yan, F., Powell, D.R., Curtis, D.J., and Wong, N.C. (2020). From reads to insight: a
1604   hitchhiker's guide to ATAC-seq data analysis. Genome Biol. *21*, 22.
1605   https://doi.org/10.1186/s13059-020-1929-3.

1606   124.   Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2.
1607   Nat. Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

1608   125.   Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O.,
1609   Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M., et al. (2021). Twelve years of
1610   SAMtools and BCFtools. GigaScience *10*, giab008.
1611   https://doi.org/10.1093/gigascience/giab008.

1612   126.   Ogata, J.D., Mu, W., Davis, E.S., Xue, B., Harrell, J.C., Sheffield, N.C., Phanstiel,
1613   D.H., Love, M.I., and Dozmorov, M.G. (2023). excluderanges: exclusion sets for T2T-
1614   CHM13, GRCm39, and other genome assemblies. Bioinformatics *39*, btad198.
1615   https://doi.org/10.1093/bioinformatics/btad198.

1616   127.   Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne,
1617   S., Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for
1618   deep-sequencing data analysis. Nucleic Acids Res. *44*, W160–W165.
1619   https://doi.org/10.1093/nar/gkw257.

1620   128.   Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G.,
1621   and Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.
1622   https://doi.org/10.1038/nbt.1754.

1623

Figure 1:

Figure 2:

A iASCL1-HA ChIP-seq

mESC | EpiLC | NE

(4,318) *Cebpa Cebpb Tnni2 Dppa1 Vgll1 Eno3*

(1,353)

(1,539)

(3,181)

(2,490) *Myt1 Hdac11 Plxna2 Onecut1 Map2 Wnt5a*

(402)

(3,927)

±1Kb ±1Kb ±1Kb

2 4 6 8 10
0 0 0 0 0

B RNA-seq *Myt1*

Norm. counts

C 0 6 24 C 0 6 24 C 0 6 24
mESC EpiLC NE

B' ChIP-seq

0–116

0–116

0–116

5 kb

VISTA: *hs2609* *Myt1*

B" VISTA: *hs2609*

E11.5

C RNA-seq *Hdac11*

Norm. counts

C 0 6 24 C 0 6 24 C 0 6 24
mESC EpiLC NE

C' ChIP-seq

0–526

0–526

0–526

5 kb

VISTA: *mm1314* *Hdac11*

C" VISTA: *mm1314*

E11.5

D ↑ DEGs RNA-seq:

235 NE-specific ASCL1 DEGs → **653** ASCL1 sites linked to NE-specific DEGs → Are these binding sites *only* bound in NE?

D'

219 | 211 | 25

Bound in:
- NE only
- NE & mESC
- NE & EpiLC
- All

198 Bound in NE only

0–177 5kb

Wnt5a

455 Bound in NE & at least one other cell type: (236 in mESC, 430 in EpiLC)

0–607 5kb

Plxna2

0–333 5kb

Inppl1

E ChIP-seq signal

NE only *n = 198*

NE & other *n = 455*

Log2(norm. reads + 1)

**** **** ****

mESC EpiLC NE

Figure 3:

A

Day 0 1 2 3 4

mESC
- No ASCL1
- 6hr ASCL1
- 24hr ASCL1

EpiLC
- No ASCL1
- 6hr ASCL1
- 24hr ASCL1

NE
- No ASCL1
- 6hr ASCL1
- 24hr ASCL1

B

PC2 (16%) vs PC1 (54%)

mESC
- No ASCL1
- 6 hr ASCL1
- 24 hr ASCL1

EpiLC
- No ASCL1
- 6 hr ASCL1
- 24 hr ASCL1

NE
- No ASCL1
- 6 hr ASCL1
- 24 hr ASCL1

C Acessibility at all ASCL1 binding sites

iASCL1 ChIP-seq: mESC, EpiLC, NE
ATAC-seq No iASCL1: mESC, EpiLC, NE

(4,318) (1,353) (1,539) (3,181) (2,490) (402) (3,972)

± 1kb

0 40 80   0 10 20

Accessibility at 653 ASCL1 binding sites linked to NE-specific ASCL1 target genes

D NE iASCL1 ChIP-seq | NE ATAC No iASCL1

E NE ATAC 24hr iASCL1

Open in NE No iASCL1 n = 493
Closed in NE No iASCL1 n = 160

± 1kb

80 40 / 20 10

F
NE ATAC No ASCL1 — 75%
NE ATAC 24hr ASCL1 — 92%

Open / Closed

G *Example ASCL1 binding sites with ASCL1-independent accessibility in NE*

Hdac11 | *Plxna2* | *Wnt5a*

NE iASCL1 ChIP-seq: 0–528 | 0–607 | 0–176
NE ATAC No iASCL1: 0–101 | 0–59 | 0–21
NE ATAC 24hr iASCL1: 0–101 | 0–59 | 0–21

5kb

H *Example ASCL1 binding sites with ASCL1-altered accessibility in NE*

Myt1 | *Elavl4*

NE iASCL1 ChIP-seq: 0–116 | 0–155
NE ATAC No iASCL1: 0–36 | 0–28
NE ATAC 24hr iASCL1: 0–36 | 0–28

5kb

Figure 4:

653 ASCL1 binding sites at NE-specific ASCL1 targets: **Are these sites *closed* in mESC?** → 362 are open in mESC: **Why are these sites not activated?**

Figure 5:

A Histone deacetylase inhibition

TSA ⊣ HDAC   HAT

B Day 0 1 2 3 4 5

RNA IF   IF

RNA IF   IF

Condition

mESC	mESC + 10 nM TSA
+ iASCL1	+ iASCL1 + 10 nM TSA
NE	NE + 10 nM TSA
+ iASCL1	+ iASCL1 + 10 nM TSA

C

mESC   NE

No iASCL1

+10 nM TSA

24 hr iASCL1

+10 nM TSA

RNA-seq

PC2: 18% variance

PC1: 60% variance

D *Nkx6-1*   **D'** *Disp2*   **D''** *Plxna2*

Normalised counts

ns *** *** ***   ns ** *** ***   ns ns *** ns

iASCL1 − + − +   − + − +
TSA   − − + +   − − + +

E Previously identified NE-specific ASCL1 target genes
n = 208

mESC

iASCL1 − + − +
TSA   − − + +

Tmem132d
Dlx2
Disp2
Dpysl4
Nos1
Wnt5a
Lhx1
Nppc
Nkx6-1
Nmrk1
Ptprn2
Tamalin
Camk2n1

Z-score
−2 0 2 4

E'

of 208 NE genes passing
significance threshold

▲ vs ●	iASCL1+TSA vs Control	116
▲ vs ●	TSA only vs Control	102
▲ vs ▲	iASCL1+TSA vs TSA only	13

F All genes up in ASCL1+TSA vs TSA only
n = 458

mESC

iASCL1 − + − +
TSA   − − + +

Z-score
−2 0 2 4

F' GO analysis

placenta development
mesenchyme development
regionalization
morphogenesis of a
branching structure
negative regulation of
cell development
regulation of
neurogenesis

p.adjust
4e-08
3e-08
2e-08
1e-08

Count
0 10 20 30

G

Figure 6:

Figure 7:

Figure S1:

Figure S2:

A ChIP-qPCR

Hes6 ORF — *Hes6* +2.5kb

Cell line: iA = iASCL1 C = Control

B iASCL1 — Control

C Normalised ChIP-seq counts

PC2 (14%) / PC1 (63%)

mESC | EpiLC | NE

D Top 10,000 ASCL1 binding sites in each cell type

mESC — EpiLC — NE

E Distal Intergenic / Downstream (<=300) / Other Intron / 1st Intron / Other Exon / 1st Exon / 3' UTR / 5' UTR / Promoter

F

ChIP-seq cluster (no. peaks)	Universal (3,927)	mESC only (4,318)	EpiLC only (1,539)	NE only (2,490)	mESC & EpiLC (1,353)	mESC & NE (402)	EpiLC & NE (3,181)
Top motif	CAGCTG	CACTG	CAGTG	CAGCTG	CACTG	CAGCTG	CAGCTG
E-value	7.5e-052	2.5e-047	2.5e-018	2.4E-028	1.8e-020	5.1e-004	3.6e-047
% sites	80.6%	75.6%	77.5%	76.7%	80.9%	78.4%	85.9%

G 24h iASCL1 DEGs with ASCL1 binding sites

95 / 685
19 / 291
34 / 385

mESC | NE | EpiLC | No binding site

H Pearson correlation between ASCL1 peak sets from published data

MEF_2 — Overexpression
mESCSerumLIF — Overexpression
mESC — Overexpression
EpiLC — *This study*
NE —
NSCs_18hr — Overexpression
NSCs — Overexpression
AdultNPCs — Endogenous
NSCs_30min — Overexpression
NeuralTubeE12.5 — Endogenous
MEF_1 — Overexpression

I ASCL1 ChIP-seq signal in mESC, EpiLC and NE at published peaks from different tissues

NSC (n = 7,387)

Neural tube E12.5 (n = 4,081)

Figure S3:

A No. ATAC-seq peaks
B FRiP (%)
C TSS enrichment
D % chrM reads

Accessibility of ASCL1 binding sites in each cell type

E All ASCL1 peaks called in mESC

mESC iASCL1 ChIP-seq | mESC ATAC No iASCL1

Open (8,007)

Closed (1,993)

± 1kb   ± 1kb

F All ASCL1 peaks called in EpiLC

EpiLC iASCL1 ChIP-seq | EpiLC ATAC No iASCL1

Open (8,677)

Closed (1,323)

± 1kb   ± 1kb

G All ASCL1 peaks called in NE

NE iASCL1 ChIP-seq | NE ATAC No iASCL1

Open (8,099)

Closed (1,901)

± 1kb   ± 1kb

H % ASCL1 binding sites that overlap with ATAC-seq peak (No iASCL1)

	mESC	EpiLC	NE
mESC only binding sites	75%	42%	34%
NE only binding sites	28%	42%	62%
universal binding sites	85%	89%	89%

I RNA-seq counts for example iASCL1 target genes shown in Fig. 4

Myt1 *Elavl4* *Hdac11* *Plxna2* *Wnt5a*

iASCL1:
mESC: C 0 6 24
EpiLC: C 0 6 24
NE: C 0 6 24

Figure S4:

Figure S5:

Figure S6:

Figure S7: