Animal collocation analysis 2.0: improving statistical inference and applications for cohort comparisons

Elliot Howard-Spink[1,2,*], Monika Mircheva[1], Judith M. Burkart[1,φ] & Simon W. Townsend[2,φ]

[1] Evolutionary Cognition Group, Department of Evolutionary Anthropology, University of Zurich, Zurich, Switzerland

[2] Comparative Communication & Cognition Group, Department of Evolutionary Anthropology, University of Zurich, Zurich, Switzerland

[φ] Co-senior authors

[*] Corresponding author: Elliot Howard-Spink (Elliot.howardspink@outlook.com)

ORCID IDs

E.H.S.: 0000-0003-3961-3982

M.M.: 0009-0006-5061-597X

J.B.: 0000-0002-6229-525X

S.T.: 0000-0003-1504-1801

Acknowledgements: This research was funded by the NCCR Evolving Language (Swiss NSF agreement Nr. 51NF40_180888) and the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 101001295). We would like to thank Stefan Gries and Alexandra Bosshard**,** whose previously published work has informed this research project.

Data & Code availability: Data and code associated with this manuscript can be found at **https://doi.org/10.17605/OSF.IO/6CGBM**

Abstract

Many animals communicate using sequences of signals, but identifying recurrent, non-random signal

combinations remains methodologically challenging. Collocation analyses are increasingly popular

approaches for detecting which signals animals combine at rates greater than expected by chance.

However, existing methods for animal collocation analysis face several limitations that reduce their

statistical rigour: they lack uncertainty estimates, fail to control for non-independence in sampled data, and

do not account for inflated family-wise error rates when identifying attraction among many different signal

types. These limitations restrict the broader applicability of animal collocation analysis, including preventing

robust comparisons of signal combination strength between cohorts (e.g. populations, sexes or age

classes). We adapt a novel form of Multiple Distinctive Collocation Analysis using Pearson residuals

(MDCA-Pr) that addresses these statistical limitations, and validate its use in animal communication

research in three ways: first, using numerous simulated datasets of different sizes and levels of signal

recombination; second, using simulated data to evaluate the performance of MDCA-Pr in intercohort

comparisons, and third, by demonstrating how MDCA-Pr can be applied to compare the vocal sequences

produced by male and female captive-living common marmosets (*Callithrix jacchus*). MDCA-Pr shows high

sensitivity, including at small sample sizes, and generally low false-positive rates, which we further reduce

by applying additional criteria for identifying attraction between signals. During intercohort comparisons,

MDCA-Pr is conservative, with low false-positive rates, and statistical power increases with sample size.

MDCA-Pr is a robust method for evaluating signal attraction in animal communication and enables accurate

intercohort comparison of animal signal combinations.

Key Words:

Signal Combinatoriality, Communication, Call Combination, Vocalization, Collocation Analysis

Statements & Declarations:

The authors declare no competing interests.

Significance Statement:

By assessing the performance of MDCA-Pr on simulated animal-like data, we demonstrate that this method

reliably detects signal combinations within and across animal cohorts, while overcoming statistical

limitations of previous collocation analyses. We present an analytical pipeline for applying MDCA-Pr to

animal signal data, including for intercohort comparisons, enabling identification and comparison of

combinatorial strategies across entire signal repertoires. We illustrate this approach by comparing call

combination strategies of male and female common marmosets when presented with food under

experimental conditions, finding similar combinatorial strategies between sexes. MDCA-Pr therefore

permits rigorous characterization of animal signal combinatoriality and opens avenues for investigating how

demographic, social, and group-level factors influence combinatorial patterns.

1. Introduction

Growing research suggests that animals combine communicative signals into non-randomly structured sequences (herein, *signal combinatoriality*; Engesser and Townsend 2019; Suzuki and Zuberbühler 2019), with increasing examples being reported across species of birds (e.g. Engesser et al. 2016, 2024; Suzuki et al. 2016; Spiess et al. 2022; Walsh et al. 2024), and mammals (e.g. Collier et al. 2017, 2020; Gu et al. 2023; Hedwig and Kohlberg 2024), particularly primates (Schlenker et al. 2016a; Bosshard et al. 2024; Berthet et al. 2025; Girard-Buttoz et al. 2025; Le Floch et al. 2026). Signal combinatoriality has been proposed to enhance the flexibility of animal communication, as smaller repertoires of signal units (e.g. calls, gestures etc.) can be combined into a much broader repertoires of combinations (Nowak et al. 2000). In some instances, the sequential order of units within combinations is important for encoding information between signallers and receivers (Engesser et al. 2016, 2019; Suzuki et al. 2016; Engesser and Townsend 2019; Leroux et al. 2023; Berthet et al. 2025; Girard-Buttoz et al. 2025); for example, wild chimpanzees (*Pan troglodytes*) have been documented producing 'Grunt→Hoo' combinations frequently in travelling and fusion contexts, whereas the reverse 'Hoot→Grunts' are used when stopping to feed and rest (Girard-Buttoz et al. 2025). Characterizing how animals combine signals is therefore crucial for understanding the structural flexibility and potential function of animal communication (Schlenker et al. 2016b&c, 2024). In addition, comparative research into animal signal combinatoriality represents one of a handful of avenues for understanding the evolution of human languages' extensive combinatoriality, including humans' dual generative systems of phonology and lexical syntax (Collier et al. 2014; Coye et al. 2018b; Fitch 2018; Townsend et al. 2018; Idsardi 2019; Zuberbühler 2019; Leroux and Townsend 2020).

Evaluating signal combinatoriality in animal communication systems requires robust methods to quantify the relative frequencies with which signals are combined, and whether these frequencies exceed those expected by random co-occurrence of signals over time. Derived from computational linguistics, collocation analysis represents one such method, and identifies whether specific pairs of sequence units (bigrams) co-occur more frequently (indicating 'attraction') or a less frequently ('repulsion') than expected based on the independent distribution of either signal type within a corpus of sample data. While several different forms of collocation analyses have been developed for analysis of human language corpora (Church et al. 1991;

Stefanowitsch and Gries 2003, 2005; Bartsch 2004; Nesselhauf 2005; Xiao and McEnery 2006; Evert 2009;

Lehecka 2015), Bosshard et al. (2022) validated two forms of collocation analysis for analysing signal

combinations within animal datasets (henceforth, 'animal collocation analysis'; methods include Multiple

Distinctive Collocation Analysis: MDCA, & Mutual Information Collocation Analysis: MICA). These forms of

animal collocation analysis allowed researchers to transition from qualitative assessment to quantitative

investigation of combinatoriality in animal signal repertoires. Consequently, several studies have applied

animal collocation analysis to identify significantly attracted signal combinations within animal vocalizations

(Leroux et al. 2021, 2022), multimodal signals (Wewhare and Krishnan 2024; Mine et al. 2024, 2025), and

during dyadic interactions (Demartsev et al. 2024; van Boekholt et al. 2025), as well as in the pairwise

structure of non-communicative sequential behaviours, such as animals' food item pairings during foraging

(Freymann et al. 2024). However, while animal collocation analysis has proven to be a valuable tool for

identifying re-occurring pairwise patterns in animals' sequential behaviours, current forms of animal

collocation analysis face several limitations that potentially compromise the validity of their outcomes (see

Gries 2022):

Firstly, existing methods for animal collocation analysis only yield point estimates of collocation strength,

and do not provide measures of uncertainty surrounding collocation strength estimates, such as confidence

intervals (limitation 1). In the absence of these measures, it is not possible to evaluate the precision of

estimated collocation strength, and this prevents thorough examination of the strength of evidence for

attraction or repulsion of signal combinations.

Secondly, all animal studies to date have reported significant attraction between signals using minimal

collocation-strength thresholds ('pbins', based on log-transformed binomial probability scores or mutual

information; Bosshard et al. 2022). However, no methods have been suggested for how to control for

inflated family-wise error rates when comparing increasingly more signal types within these analyses

(increasing the likelihood of type 1 errors for significant attraction or repulsion; limitation 2).

Thirdly, while existing methods for collocation analysis compare the frequency of signal co-occurrence

relative to what is expected if each signal type was distributed independently in a sample, they do not

account for whether nested relationships exist within sample data which could also influence the frequency

of signal co-occurrence (limitation 3). For example, pairwise signal transitions may be recorded from the

same bout, or same individual, meaning signal bigrams sampled for the analysis are not independent.

Currently, no methods have been proposed regarding how to account for this non-independence when

generating point-estimates in animal collocation analysis.

Finally, current methods for animal collocation analysis impose a trade-off between identifying directionality

in collocates (using MCDA, where A→B differs to B→A) or to maintaining higher sensitivity to rarer but

significant collocates (MICA, which is blind to the direction of transitions; Bosshard et al. 2022; limitation 4).

Analyses that can combine sensitivity and directionality are therefore needed, as directionality can be a

functionally-meaningful aspect of animals' signal combinations (Girard-Buttoz et al. 2025).

In an effort to overcome these obstacles when analysing human language corpora, Gries (2023) recently

proposed several adaptations to MDCA to improve the analysis' statistical rigour. Classical MDCA involves

iterative construction of 2x2 binomial arrays to estimate the probability of each type of sequence unit being

combined, and thus its collocation strength; however, this is computationally costly, and can be time

consuming. Therefore, Gries proposes quantifying collocation strength using Pearson residuals from a

singular array tallying all pairwise unit co-occurrences. Pearson residuals (*Pr*) are scaled comparisons of

the number of instances a particular pair of units co-occurred (*O*) compared to the number of times is

Expected (*E*) by chance given the frequency of either sequence unit type in the sample corpus (Liao et al.

2024):

$$Pr = (O - E)/\sqrt{E}$$

By estimating these residuals from one global table, MDCA by Pearson residuals (MDCA-Pr) considerably

reduces the computational burden of collocation analysis. This lends the analysis to bootstrapping, where

uncertainty estimates for the collocate strength for each pair of signal units can be generated, e.g.

confidence intervals for collocation strength (solving limitation 1; see Gries 2023). The size of confidence

intervals can be adjusted through Bonferroni correction, thus accounting for family-wise error rates that

have been inflated by multiple comparisons of different signal pairs (solving limitation 2; see Studies 1 & 2

below). Moreover, through block bootstrapping (resampling with replacement at highest hierarchical levels

in the corpus data, e.g. by resampling all bouts produced by a particular individual all at once), uncertainty

measures for collocation strength can be estimated whilst accounting for nested structure in the data

(solving limitation 3; see Gries 2023). As MDCA-Pr is also directional, this method may have the potential

to balance directionality and sensitivity (solving limitation 4). However, the performance of MDCA-Pr has

yet to be tested on the types of datasets collected by animal behaviour researchers, which are often small,

skewed, and contain far fewer element types than linguistic corpora.

In addition to these statistical improvements, MDCA-Pr may offer new avenues to explore questions that

are intractable with previously validated forms of animal collocation analysis, including whether signal

combinatoriality can be compared between cohorts of animals (e.g. different groups or populations of

animals, or different sex or age classes within a species). Emerging studies indicate that for some species,

signal combinations may differ between cohorts (usually animal populations; Schlenker et al. 2014; Girard-

Buttoz et al. 2022a), as may the contexts in which shared signal combinations are used (Schamberg et al.

2024), possibly reflecting cohort-specific communicative norms. However, these studies have

overwhelmingly focused on the *rates* with which signals are combined (irrespective of the signal units) or

the presence/absence of specific signal combinations. As of yet, no studies have characterized the extent

of intraspecific variation in combinatoriality across a species' entire signal repertoire, largely because of the

limited availability of analytical tools capable of supporting such comparisons. Previously validated

approaches to collocation analysis cannot fill this methodological gap, as they do not provide uncertainty

estimates for collocation strength. This makes it impossible to assess whether observed differences

between cohorts reflect valid behavioural differences, or if they are the product of trivial sampling error.

However, because MDCA-Pr can be easily bootstrapped to estimate confidence intervals, it may permit

formal testing for significant differences in collocation strength between cohorts. Given this potential, the

use of MDCA-Pr during intercohort comparison would also benefit from empirical validation.

Here, we take a three-pronged approach to validate the suitability of MDCA-Pr to analysing animal

behavioural data, including during intercohort comparison, using a mixture of synthetic vocal data and vocal

data taken from captive-living common marmosets (*Callithrix jacchus)*:

In Study 1, we evaluate the performance of MDCA-Pr on synthetic animal-like vocal datasets of different

sizes and levels of recombinatoriality, following a validation procedure from Bosshard et al. (2022; see

Section 2.1). We quantify the sensitivity (true-positive rate) and selectivity (false-positive rate) of MDCA-Pr

to evaluate its performance when identifying call attraction across these different types of animal datasets.

In Study 2, we evaluate the performance of MDCA-Pr when comparing synthetic vocal data from simulated

animal cohorts (Section 2.2). We assess the sensitivity with which MDCA-Pr can identify pre-programmed

differences in the probability of calls being combined between cohorts (the true-positive rate), and selectivity

of the analysis, including whether any call combinations with the same probabilities between cohorts were

erroneously identified as having different collocation strengths (the false-positive rate). Similarly to Study 1,

we evaluate MDCA-Pr during intercohort comparison on synthetic datasets of varying sizes and levels of

recombinatoriality, to build a comprehensive picture of the performance of MDCA-Pr given the different

types of datasets that animal behaviour researchers frequently encounter.

Finally, in Study 3, we extend beyond simulated data, by applying MDCA-Pr to compare combinatorial

strategies of captive-living male and female common marmosets in a feeding context (Section 2.3). This

study allows us to illustrate how MDCA-Pr can be applied to data from real interacting animals in a controlled

foraging context.

Together, this suite of analyses allows us to compare the performance of MDCA-Pr to previously-validated

forms of animal collocation analysis, and to evaluate its efficacy in identifying similarities and differences in

combinatorial behaviours between cohorts of animals. Additionally applying MDCA-Pr to marmoset call data

demonstrates its applicability to real-world data, and provides an initial showcase of how cohort-level

combinatoriality can be compared.

2. Methods & Results

Across all three studies, MDCA-Pr was performed using the method proposed by Gries (2023), using an

adapted code in R (v. 4.5.1; R Core Team 2022; for original code see Gries 2024). In each study, the

simulated and real call sequences were parsed into bigrams that described pairwise co-occurrence of calls.

Each unique type of bigram was tallied in a global table, and Pearson residuals were calculated for each

bigram type using the chisq.test() function from the stats package (base R v 4.5.1; R Core Team 2022).

Pearson residuals were calculated across 10,000 bootstraps of the data (blocked at the level of the

individual), and confidence intervals were estimated from this bootstrapped distribution for each bigram

type, where the original 95% confidence interval (alpha = 0.05) was adjusted via Bonferroni correction to

the number of comparisons being made (alpha$_{adjusted}$ = 0.05/Number of bigram types compared;

comparisons were counted as either each instance a bigrams' collocation strength was compared to zero

for a particular dataset (e.g. Study 1), or each instance a bigrams' collocation strength was compared

between two cohorts (e.g. Study 2)). Confidence intervals were estimated non-parametrically using upper

and lower quantiles of the bootstrapped distribution, and were constructed for two-tailed comparison (lower

percentile = 100 x alpha$_{adjusted}$/2; upper percentile = 100 x (1 – alpha$_{adjusted}$/2)). All scripts to run MDCA-Pr

on animal data can be found in our supplementary code.

2.1 Study 1 – validation of MDCA-Pr using simulated data

2.1.1 Simulating animal-like call data

Following Bosshard et al. 2022, we simulated four distinct datasets of animal-like calls. Two *Small* datasets

(N$_{Individuals}$ = 10; 16 bigrams per individual; total bigrams = 160), and two *Large* datasets (N$_{Individuals}$ = 20; 160

bigrams per individual; total bigrams = 3200). For simplicity, we assume that each bigram comes from its

own bout of 2 calls (i.e. for the *Small* dataset, 32 individual calls were produced in 16 pairs per individual)

and provide an example of how to analyse longer call bouts in Study 3 (Section 2.3).

For each size category, an *Exclusive* dataset was simulated that only contained four call types combined

into three distinct focal call bigrams: Peep→Howl, Howl→Peep and Huff→Puff (see Table 1). The *Small*

and *Large Exclusive* datasets functioned as a positive control, where calls within focal bigrams should be

identified as 'attracted' as no other competing call bigrams appear in the data.

Additionally, a *Recombinatorial* dataset was produced for each size category. These *Recombinatorial*

datasets included the same three focal call bigram types as before (each with a different probability of

occurring in the data; see Table 1). However, the *Recombinatorial* datasets also featured three additional

call types (Cough, Whistle and Hum), which, alongside Peep, Howl and Puff, were programmed to flexibly

combine into bigram types at lower probabilities (2% each). Some of these background combinations

included call types present in the focal call bigrams (e.g. Peep→Cough, or Hum→Puff), whereas others did

not (e.g. Cough→Hum). As the focal call bigrams occur at higher frequencies than other bigram types, focal

call bigrams should be identified as attracted call combinations (as found by other forms of animal

collocation analysis when run on data with the same bigram probability distributions; Bosshard et al. 2022).

In general, across the simulated datasets (*Small Exclusive*, *Small Recombinatorial*, *Large Exclusive*, *Large*

Recombinatorial), those that are smaller and more recombinatorial present a greater challenge for MDCA-

Pr to identify attraction between calls in the focal bigrams.

In addition, we ran MDCA-Pr on four datasets that were of equivalent size to each of these four datasets

and had the same numbers of call types; however, call bigrams were constructed by sampling call pairs

from a uniform random distribution. These randomized datasets offered a negative control, where no

significant attraction or repulsion should be identified between any call types.

2.1.2 Evaluating MDCA-Pr performance

MDCA-Pr was applied to each dataset independently. Following existing literature, we only refer bigrams

as call combinations if they are identified as experiencing significant levels of attraction (e.g., Leroux et al.

2022). To identify attracted call combinations across all bigram types, we followed a two-step procedure:

1. To identify call combinations that was unlikely to be driven by chance association in the data, we

initially identified attraction by determining which call bigram types had confidence intervals for collocate

strength that were wholly above zero.

2. After evaluating the results from step 1, we extended this initial assessment of attraction by

following a recent study that applied MDCA to animal communication data (van Boekholt et al. 2025). This

study reported that MDCA can identify many low-level attracted combinations in behavioural units,

potentially to frequencies that suggest instances false-positive attraction. The authors proposed using an

additional threshold to identify 'strongly attracted' bigrams, where attracted call combinations required a

collocation strength that surpassed two standard deviations of collocation strengths for all bigram types.

We evaluated whether a similar threshold would enhance the performance of MDCA-Pr using a post-hoc

analysis, where we reclassified bigram types as 'attracted' if their mean collocate strength was one standard

deviation higher than the mean of all collocate strengths. We chose to start at one standard deviation (rather

than two) to test the methodological benefits of a less conservative threshold for MDCA-Pr when accounting

for false-positive rates. If false-positive rates persisted under this threshold, our analysis lends itself to

calibration of a more suitable threshold that can address low-level false-positive rates across datasets of

different sizes and levels of recombinatoriality.

For each simulated dataset, we evaluated the performance of MDCA-Pr using measures of sensitivity and

selectivity. Sensitivity was measured by the number of true positives identified (more means higher

sensitivity), with a true positive being any instance where significant attraction was identified for the focal

call bigram types. Selectivity was measured by the number of false positives identified (lower means better

selectivity), with false positives categorized as any instance where significant attraction was detected

between non-focal call bigram types. We focused on attraction only, and did not examine significant

repulsion between call types (Gries 2023), as animal communication studies almost exclusively aim to

characterize which calls animals frequently combine, rather than those animals rarely or never combine.

2.1.3 Results

When identifying attracted call combinations solely based on the confidence intervals for collate strength,

MDCA-Pr demonstrated perfect selectivity and sensitivity across both *Small* and *Large Exclusive* datasets.

All three focal call bigram types were identified as significantly attracted, and no false-positive attractions

were identified (see Fig. 1. & Table 1; see Tables S1 & S2). As predicted, non-simulated combinations (e.g.

Peep→Puff) were significantly repulsed as these call pairings never occurred within the data.

Perfect selectivity was achieved for the *Small Recombinatorial* dataset, as no false-positive attraction was

identified. However, sensitivity was somewhat lower for the *Small Recombinatorial* dataset: only two focal

call bigram types were identified as significantly attracted (Peep→Howl, & Howl→Peep; Fig. 1; Tables 1 &

S3). For Huff→Puff, the focal call bigram type with the lowest combination probability, the confidence

interval for collocation strength overlapped zero.

Increasing the size of the *Recombinatorial* data improved sensitivity, as in the *Large Recombinatorial*

dataset, all focal call bigram types were identified as attracted (see Fig. 1 & Table S4). However, this came

at the expense of selectivity, as significant lower-level attraction was identified between four of the non-

focal call bigram types.

For each of the four datasets, we reclassified 'attraction' as any instance where a call bigram type had a

confidence interval for collocation strength that was wholly above zero, and a mean collocation strength

that exceeded one standard deviation of the collocation strengths across all bigram types. Following this

reclassification, the sensitivity of our analysis was unchanged, with attraction being identified at the same

rate as before across all focal bigram types. However, selectivity was improved after this threshold was

applied, with no false-positive attraction being identified in any of the datasets (see Fig. 1 & Table 1).

In the *randomized control* datasets, no significant attraction or repulsion was detected (both with and without

the addition of a 1SD filter), meaning MDCA-Pr is unlikely to identify significantly attracted call combinations

in completely randomized data (see Table 1 & Fig. S1).

Table 1 Simulated data and results of Study 1. All probabilities are rounded to 3.d.p, and collocation strengths presented to 3.s.f.

Focal call bigram types are in bold. Bigram probability indicates the likelihood that each type of bigram is simulated during the

production of data for each artificial individual. Mean collocate strengths are provided for focal call bigram types, alongside confidence

intervals. Collocate strengths for all other bigram types are reported in Tables S1-S4. Asterisks indicate significant attraction for a

given bigram type, including after applying the 1SD filter. Horizontal dashes indicate no significant attraction. For control datasets, no

attraction was detected even prior to applying the 1SD filter

Call Bigram type	Bigram Probability	MDCA-Pr Result								
		Small Dataset			Large Dataset			Control Datasets		
Exclusive		Mean Collocate Strength	1SD Upper Threshold	Attraction?	Mean Collocate Strength	1SD Upper Threshold	Attraction?	Small	Large	
Peep→Howl	**0.313**	8.61	6.32	*	39.2	28.3	*	-	-	
Howl→Peep	**0.438**	7.12	6.32	*	31.4	28.3	*	-	-	
Huff→Puff	**0.250**	9.57	6.32	*	42.5	28.3	*	-	-	
Recombinatorial										
Peep→Howl	**0.102**	4.81	1.55	*	16.9	5.99	*	-	-	
Peep → (Not Howl)	0.020			-			-	-	-	
Howl→Peep	**0.143**	3.84	1.55	*	18.5	5.99	*	-	-	
Howl→X (Not Peep)	0.020			-			-	-	-	
Huff→Puff	**0.082**	1.94	1.55	**-**	14.8	5.99	*	-	-	
Huff→X (Not Puff)	0.020			-			-	-	-	
X→X (Any other combination, e.g. Hum→Cough)	0.020			-			-	-	-	

Figure 1 MDCA-Pr on the four non-random simulated datasets from Study 1. Points indicate the mean collocation strength for a given call bigram type across bootstraps, and error bars indicate the confidence interval surrounding each mean (95% CI adjusted by Bonferroni correction; see Section 2). Focal call bigrams labelled in bold and plotted in green. Data plotted in blue indicates repulsion between two call types (CI wholly below zero). Data plotted in red indicates initial attraction for non-focal call bigrams (CI wholly above zero) prior to applying the +-1SD filter (indicated by the dashed horizontal lines). Zero is indicated by a solid horizontal line

2.2 Study 2 – intercohort comparisons using simulated data

2.2.1 Simulating call bigram data across cohorts

To evaluate whether MDCA-Pr could be used to compare combinatoriality across cohorts of animals, we simulated four classes of dataset, each containing three corpora of call bigrams. These corpora were of a fixed size within each class (*Small* & *Large*; following the size categorizations from Study 1) and contained

specific levels of recombinatoriality (see below). For each class, two corpora were generated using the

same probability distribution for all call bigram types (*Equivalent* cohorts 1 & 2) and a single corpus

generated from a different probability distribution (an *Unequivalent* cohort).

Two classes of datasets were *Recombinatorial*: one *Large* and one *Small*. For the *Equivalent*

Recombinatorial corpora in each class, we used the same probabilities for each call bigram type as the

Recombinatorial datasets used in Study 1. Whereas, for the *Unequivalent Recombinatorial* corpus in each

class, we reduced the probability of each focal bigram type occurring in the data by increasing amounts

(Peep→Howl: -5%; Huff→Puff: -7.5%; Howl→Peep: -10%; see Table 2). For each reduction in the

probability of a focal call bigram occurring in the data, all alternative transitions away from the first call in

the bigram received a small uniform increase in their likelihood of being produced (e.g., as Peep→Howl

was reduced by -5%, the bigrams Peep→Cough, Peep→Hum, Peep→Peep, Peep→Puff, & Peep→Whistle

all increased by 1% each). This ensured that the overall probability distribution across bigram types always

summed to 1. Whilst many bigram types therefore have different probabilities of occurring in the *Equivalent*

and *Unequivalent* corpora, we retain our use of 'focal bigram type' for Peep→Howl, Howl→Peep, and

Huff→Puff, as these bigram types exhibit the greatest differences between the two types of corpora.

We did not simulate any *Exclusive* classes of datasets, as any reductions in the probability of a focal call

bigram occurring in the data had to be compensated for by increasing the probability a related call bigram

being sampled during data simulation (i.e. a reduction in A→B led to an increase in A→C), thus we were

not able to maintain the exclusivity property when simulating an *Unequivalent* corpus.

Instead, we simulated two *Weakly Recombinatorial* classes of datasets, which had the same call types as

those in the *Exclusive* datasets from Study 1 (Howl, Peep, Huff & Puff), but in addition to the three types of

focal call bigrams, these calls were recombined into several non-focal bigram types at low levels (each with

a 1% probability of occurring; see Table 2 & Fig 2). As before, we simulated *Small Weakly Recombinatorial*

and *Large Weakly Recombinatorial* classes of data, each with two *Equivalent* corpora, and one

Unequivalent corpus. For the *Unequivalent* corpus, the probability of each focal call bigram occurring in the

data was reduced by the same amounts as in the *Recombinatorial* classes, and these probabilities were

distributed among other bigram types in the same way (see Table 2).

For all corpora, the frequency distributions for all call bigram types were visualized to ensure our simulated

corpora included sampling variation (at both the individual level, and between equivalent cohorts; see Figs.

S2-S4). This ensured that our simulations contained some stochastic variation, similarly to when data is

sampled from real-world animal populations.

2.2.2 Comparing simulated cohorts

For each dataset, MDCA-Pr was run on all three simulated corpora. We then made two comparisons:

1. A comparison of *Equivalent* cohort 1 and the *Unequivalent* cohort. This comparison allowed us

309   to detect whether imputed differences in call bigram probabilities between cohorts could be

310   detected (a measure of sensitivity).

311   2. A control comparison between *Equivalent* cohorts 1 and 2. This negative control comparison

312   allowed us to identify whether false-positive differences would be identified between cohorts

313   that had the same underlying probability of producing all call bigram types (a measure of

314   selectivity).

During these comparisons, significant differences in the collocation strength for a given call bigram type

were identified by non-overlapping confidence intervals between cohorts (where confidence intervals were

Bonferroni corrected to account for the number of pairwise comparisons between cohorts). Overlapping

confidence intervals were taken to indicate that there was no significant difference in collocation strength

between cohorts for a given bigram type. We compared all bigram types between cohorts, regardless of

whether bigrams were identified as significantly attracted in either corpus. This was necessary as for the

Unequivalent cohort, focal call bigram probabilities were reduced to be very similar to those of non-focal

call bigrams. Consequently, all bigram types may re-occur at similar rates in the *Unequivalent* corpus,

preventing any subset of bigrams being identified as significantly attracted.

2.2.3 Results

Across all classes of datasets, collocation strength did not significantly differ between *Equivalent* cohorts 1

and 2 for any bigram types, reflecting perfect selectivity of MDCA-Pr during this control intercohort

comparison (see Table 2; Fig S5 & Tables S5-S8).

When comparing *Equivalent* cohort 1 with the *Unequivalent* cohort, MDCA-Pr successfully identified

differences in collocation strengths for all focal bigram types in both *Large* data classes, and correctly

identified a number of differences in the probability of non-focal bigrams occurring in the two corpora (all

modified non-focal bigram types for the *Large Weakly Recombinatorial* dataset, and 6/15 modified non-

focal bigram types for the *Large Recombinatorial* dataset; see Table 2). Therefore, for *Large* datasets,

MDCA-Pr was sensitive to unequal probabilities of calls being combined between cohorts, including

consistent detection of differences as low as 5%.

For the *Small* datasets, sensitivity was lower during comparisons of the *Equivalent* cohort 1 and the

Unequivalent cohort. For the *Small Recombinatorial* dataset, only two focal bigram types were identified as

having different collocation strengths between the cohorts (Howl→Peep and Huff→Puff). Whereas, for the

Small Weakly Recombinatorial dataset, only one focal bigram type was identified differing in collocation

strength between cohorts (Huff→Puff). Thus, when datasets are *Small*, MDCA-Pr can be less sensitive to

intercohort differences in the probabilities of calls co-occurring, particularly when background recombination

is limited.

The sensitivity of our analysis during intercohort comparisons differed considerably between the *Small* and

Large datasets. As the size of these corpora differ substantially (*Small* = 160 bigrams; *Large* = 3200

bigrams), we ran additional analyses to evaluate whether more moderate increases in sample size would

improve sensitivity. Doubling the *Small* sample size (320 bigrams; either by doubling the number of

individuals, or the number of calls per individual), improved sensitivity during intercohort comparisons when

analysing *Weakly Recombinatorial* data (Double Calls = 2/3 focal bigram types identified as having different

collocation strengths; Double Individuals = 3/3; see Fig. S6) whereas sensitivity remained consistent or

improved for *Recombinatorial* data (Double Calls = 3/3; Double Individuals = 2/3).

Table 2 Simulated data and results of Study 2. Bigram probability indicates the likelihood that each type of bigram is simulated

during the production of each dataset. Asterisks indicate that a significant difference in collocation strength between cohorts for a

particular bigram type. Horizontal dashes indicate no significant difference. Specific values for collocate strengths for all call bigram

types are reported in Tables S5-S8

Call Bigram Type	Bigram Probability			MDCA Result: Equivalent		MDCA Result: Unequivalent	
	Equivalent Cohorts	Unequivalent Cohort	Treatment Difference	Small Dataset	Large Dataset	Small Dataset	Large Dataset
Weakly Recombinatorial							
Peep→Howl	**0.293**	**0.243**	**-0.050**	-	-	-	*
Peep→X (Not Howl)	0.010	0.035	+0.025	-	-	-	*All Transitions*
Howl→Peep	**0.418**	**0.318**	**-0.100**	-	-	-	*
Howl→X (Not Peep)	0.010	0.060	+0.050	-	-	*Howl→Puff*	*All Transitions*
Huff→Puff	**0.230**	**0.155**	**-0.075**	-	-	*	*
Huff→X (Not Puff)	0.010	0.048	+0.038	-	-	-	*All Transitions*
Recombinatorial							
Peep→Howl	**0.102**	**0.052**	**-0.050**	-	-	-	*
Peep→X (Not Howl)	0.020	0.030	+0.010	-	-	-	*Peep→Peep *Peep→Puff*
Howl→Peep	**0.143**	**0.043**	**-0.100**	-	-	*	*
Howl→X (Not Peep)	0.020	0.040	+0.020	-	-	-	*Howl→Howl *Howl→Puff*
Huff→Puff	**0.082**	**0.007**	**-0.075**	-	-	*	*
Huff→X (Not Puff)	0.020	0.035	+0.015	-	-	-	*Huff→Howl *Huff→Peep*
X→X (Any other combination, e.g. Hum→Cough)	0.020	0.020	0	-	-	*Cough→Peep*	-

Figure 2 MDCA results comparing *Equivalent* cohort 1 and the *Unequivalent* cohort for each dataset. The top row shows the *Weakly Recombinatorial* data classes, including **a.** *Small* and, **b.** *Large* corpora. The bottom row shows the *Recombinatorial* data classes, including **c.** *Small* and **d.** *Large* corpora. Significant differences between cohorts are identified by no overlap in confidence intervals for a given call bigram type (also see Table 2). Focal bigram types are marked in bold

2.3 Study 3 – marmoset vocal combinatoriality between sexes

2.3.1 Marmoset vocal data

To provide an example of how MDCA-Pr can be applied to intercohort comparisons of animal vocal data, we investigated whether male and female common marmosets differ in their call combinatoriality when in standardised context, in this case feeding. Common marmosets are a highly vocal primate species, possessing a broad repertoire of call types (Agamaite et al. 2015) that can be flexibly combined into a diverse array of sequences (Bosshard et al. 2024). As cooperative breeders, marmosets engage in active food sharing, and have various short and long-distance calls to signal food availability and individual's locations to conspecifics (Bosshard et al. 2024; Mircheva et al. 2026). These calls may signal individuals' own arousal, particularly for females, who get priority access to food (Box et al. 1999; Guerreiro Martins et al. 2019). However, calls may also indicate a general readiness to share food, especially for males, who are more proactive in ensuring food is shared equitably with partners (Mustoe et al. 2016). These discrepancies in food access and sharing between sexes may be reflected within the vocalizations used by either sex when encountering food; for example, males generally produce a higher number of food calls (Mircheva et al. 2026). However, it has not yet been examined whether these behavioural differences between sexes are reflected in the ways in which calls are combined.

We applied MDCA-Pr to an existing corpus of marmoset calls collected by Mircheva et al. (2026). Pairs of monkeys were recorded in two joint 1 x 1.8 x 2 m experimental chambers, where individuals were out of sight from each other (blocked by an opaque screen), but sounds (including vocalizations) could readily travel between either individual in the pair. One monkey of each pair was presented with different stimuli to create distinct behavioural contexts, including an unfamiliar modified rubber duck (inducing an unfamiliar context) or meal worms (inducing a feeding context). Only the individual who was presented with the stimuli was labelled as being in that context. After three minutes, the stimuli were removed, and either a new stimulus was presented to one individual in the pair, or the experiment ended. As data was collected from combinations of known individuals, data collection for this study was not blinded.

To identify the call combinations used by male and female common marmosets when in a feeding context, we sampled all calls produced by a particular individual during the time interval they (and not their partner)

were presented with the food stimulus. We restricted our sample to experimental sessions where the pair

of individuals were the breeding male and female of each group (2 sessions per pair). This controlled for

the effects of developmental stage and breeding status on the dyadic vocal interactions between pairs of

marmosets. In total, we initially sampled 10 individuals in pairs from 5 groups.

Our initial sample included 3007 calls (1695 for males; 1312 for females). These calls were part of eight

unique call types (see Table S9). Following existing literature on marmoset communication (Meshinska et

al. 2024; Bosshard et al. 2024), any calls which occurred within 0.5s of another call were classified as part

of the same bout. Additionally, following previous studies of marmoset call combinatoriality, we focused on

how marmosets combine different *types* of calls, and therefore condensed any repeats of the same call

type within a bout down to a single token in the data (Bosshard et al. 2024). Our analysis of marmoset call-

type combinations therefore also reflected Studies 1 & 2, where repeated calls were not included in

simulated data. Given repeated calls may be an important aspect of animal combinatoriality (Templeton et

al. 2005; Engesser et al. 2017; Berthet et al. 2019), an example applying MDCA-Pr to marmoset call data

that included repeats can be found in our supplementary materials. Call-type transitions were sampled from

bouts by constructing bigrams that describe all the internal neighbouring call-type combinations. For

example, the hypothetical bout '*A,B,C,A*' would be converted into three bigrams: $A \rightarrow B; B \rightarrow C; C \rightarrow A.$

Whilst this study did not collect new data from Marmoset monkeys, the data used were collected from

experiments approved by the Kantonales Veterinäramt Zürich, Switzerland (license number ZH 232/19),

and complied with Swiss law (Mircheva et al. 2026).

2.3.2 Applying MDCA-Pr to sex cohorts

MDCA-Pr was applied to the male and female call data. We excluded data for monkeys where fewer than

20 call type bigrams were sampled, representing a trade-off between maximizing the number of individuals

included in the data versus the vocal data that could be sampled per individual. This left four males and

three females from our initial sample, with a corpus of 733 recorded productions of non-repeating call types,

described via 488 pairwise transitions between call types (Males = 266 bigrams; Females = 222) across

246 bouts of vocalization. Note that, as call-type pairs in the same bout are not independent, the total raw

sample of call types is not a perfect duplication of pairwise transitions (see the example in the section above

that converted four call types into three transitions).

To ensure that the number of males and females sampled for each bootstrap was equal, we sampled with

replacement three individuals per sex per bootstrap. To account for uneven numbers of call-type bigrams

recorded from different individuals in the data, we modified the bootstrapping procedure to subsample equal

amounts of data from each individual monkey, using the smallest sample size for any individual monkey in

the data (22 call-type bigrams per individual). Subsampling prevented particularly voluble individuals from

being overrepresented in the bootstrapped datasets. Subsampling within individuals was conducted without

replacement, following the logic of methodological approaches used to control for unequal cluster sizes in

hierarchical or clustered resampling (e.g., Lee et al. 2019). Each bootstrap therefore had a relatively small

sample size of 66 call-type bigrams per sex.

For both sexes, we classified significantly attracted call types as those for which the confidence interval for

collocate strength was wholly above zero (confidence intervals were adjusted using the same method as

Study 2), and where the mean collocation strength for that call-type bigram was above one standard

deviation of all mean collocate strengths for a particular sex (as advised from Study 1). Where call types

were identified as attracted for both sexes, we evaluated whether collocate strength differed between sexes

(where non-overlapping confidence intervals between sexes indicated significant differences in collocate

strength; as advised by Study 2). Call-type combinations that were only found in one sex were termed either

male-biased or female-biased call-type combinations.

For each attracted combinations of call types, we ran a post-hoc analysis to review which individuals

produced each call-type combination in the underlying data. This allowed us to determine whether attracted

call types were being produced by many individuals (indicating either species-typical, or sex-biased

behaviours), or if they were produced by only one or a few individuals (thus, more likely reflecting

idiosyncratic call-type combinations of individuals in either cohort).

2.3.3 Results

We identified a total of five attracted call-type combinations (see Table 3). Two were shared between both

sexes: a combination of two long-distance call types (Food Phee→Phee), and a combination of two

mobbing call types (Tsik→Ekk). No significant difference in attraction strength was found between the sexes

for these shared call-type combinations.

Post-hoc analysis of the underlying data revealed that the Tsik→Ekk call-type combination was produced

by all individuals of both sexes, supporting that it is a species-typical behaviour. Post-hoc analysis also

identified that the Food Phee→Phee call-type combination was balanced between the sexes in the

underlying data; however, it was rare, and produced by one individual of either sex (once by a male, and

twice by a female). This combination of call types was likely identified as attracted given that sequences of

Food Phees only ever transitioned into sequences of Phees. Despite its rarity, the underlying data supports

equal usage by males and females.

We identified three female-biased call-type combinations: Food Peep→Food Phee; Ekk→Tsik; and

Phee→Short Phee. The Phee→Short Phee combination was produced by individuals of both sexes.

However, within the underlying data, Phee→Short Phee combinations were more commonly produced by

females ($N_{Transitions}$ = 16 for males; $N_{Transitions}$ = 24 for females), and males combined Phee calls with a greater

diversity of other call types ($N_{CallsTypes}$ = 4 for males, $N_{CallTypes}$ = 3 for females), supporting female-biased

call-type attraction. The Food Peep→Food Phee combination was rare, produced once by one female.

However, it was the only instance of a Food Phee following another call other than itself. For the female-

biased Ekk→Tsik call-type combination, analysis of the data revealed that both sexes produced Ekk→Tsik

call-types together (as reflected by the similar mean collocation strength between sexes). However, for

males, the confidence interval for this call-type transition overlapped with zero due to a greater uncertainty

around true collocate strength (meaning it was not a significant attraction). Whether both Ekk→Tsik and

Food Peep→Food Phee represent female-biased call-type combinations would benefit from analysis of a

larger dataset.

Table 3 Attracted call-type combinations identified via MDCA-Pr for male and female marmosets. The mean collocation strength
across bootstraps is shown for each call-type combination, alongside the confidence interval for collocation strength. Horizontal
dashes indicate that a particular call combination was not significantly attracted for a specific sex. All values rounded to 3.d.p.

Sex Category	Call combination	Male			Female		
		Mean	Lower CI	Upper CI	Mean	Lower CI	Upper CI
Both sexes	Tsik→Ekk	4.481	1.600	6.068	4.677	2.415	7.140
	Food Phee→Phee	3.746	2.180	5.222	3.008	1.711	6.030
Female-Biased	Ekk→Tsik	-	-	-	3.900	1.919	8.000
	Phee→Short Phee	-	-	-	4.556	3.015	6.724
	Food Peep→Food Phee	-	-	-	2.307	1.539	4.264

3. Discussion

Collocation analysis has proven to be a useful tool for identifying repeated combinations of signals within
animal communication. However current methods for animal collocation analysis lack important uncertainty
estimates for collocation strength; overlook the influence of inflated family-wise error induced by multiple
comparisons of signal pairs, and ignore nested structure within datasets, all of which reduce the assurance
with which animal signal combinations can be validly identified. Additionally, these statistical limitations limit
the diversity of questions which can be examined by existing forms of animal collocation analysis, including
by preventing cohort-level comparisons of combinatoriality across animals' full signal repertoires. Using
simulated animal-like data, we evaluated the performance of a recently developed form of collocation
analysis that can address these statistical limitations (MDCA-Pr), and validate whether it offers a suitable
alternative to existing forms of animal collocation analysis. We also evaluated the performance of MDCA-
Pr when comparing the combinatorial repertoire of simulated animal cohorts, before providing an example
of how this method can be applied to intercohort comparisons of vocal combinatoriality in real animals.

3.1 Study 1

Study 1 was used to evaluate the performance of MDCA-Pr on simulated datasets of different sizes and
levels of recombinatoriality. Sensitivity of the analysis was generally high across different types of data.

Only one false negative was detected in the *Small Recombinatorial* dataset (the focal bigram type with the

lowest combination probability; Huff→Puff, 8%), indicating that MDCA-Pr may only fail to identify call

combinations that are relatively weakly attracted if datasets are small and background recombinatoriality is

high. The selectivity of MDCA-Pr was also generally high, with false positives only being identified in one

dataset (*Large Recombinatorial*). Following a methodological adjustment suggested by van Boekholt et al.

(2025), where an additional filter for identifying attraction was implemented based on the standard deviation

of all bigrams, our analysis achieved perfect selectivity without any resultant costs to sensitivity. This high

selectivity was also reflected in our randomized control data, where no attracted call combinations were

detected.

MDCA-Pr therefore performed well on the analysis of animal-like datasets, including those of varying sizes

and levels of recombinatoriality. However, our analysis highlights the importance of carefully evaluating the

performance of linguistic methods when applied to animal-like data, as additional filters for identifying call

attraction were necessary for optimal MDCA performance. Overall, we report that MDCA-Pr offers a useful

approach for identifying attracted signal combinations within the communicative repertoires of animals,

using methods that possess greater statistical rigour than existing forms of animal collocation analysis.

However, we recommend that this method be implemented alongside additional filters that maintains the

selectivity with which attracted call combinations are identified (see Fig. 3).

3.2 Study 2

In Study 2, we evaluated the performance of MDCA-Pr when identifying differences in the combinatoriality

between cohorts of individuals. MDCA-Pr exhibited perfect selectivity during intercohort comparison, with

no false-positive differences being identified when corpora were generated using the same probability

distribution for bigram types. MDCA-Pr exhibited increasing sensitivity for larger sample sizes. Whilst

sensitivity was low for *Small* sample sizes, even moderate increases in sample size (doubling the *Small*

sample) markedly improved the analysis' sensitivity.

Our results validate that MDCA-Pr can detect intercohort differences in signal combinations, and is unlikely

to identify false-positive differences between cohorts. However, sample size is an important consideration

for ensuring MDCA-Pr is sensitive to more subtle differences between cohorts. Our simulated differences

in focal call combination probabilities were relatively small between the *Equivalent* and *Unequivalent*

cohorts (between 5-10%; see Table 2), and to detect these differences, several-hundred signal

combinations were required. Arguably, meaningful differences between the behaviours of real animal

cohorts could be expected to be greater than those we simulated, and in these instances, differences will

likely be more easily detected using small samples. Nevertheless, the analysis of small samples should be

expected to give conservative estimates of intercohort differences. To estimate minimal sample sizes

needed to maintain adequate power of MDCA-Pr during intercohort comparison, researchers could

consider simulating their own bespoke datasets of different sizes, tailored to the size of species' signal

repertoire and initial estimates of recombinatoriality. This may then permit evaluation of the minimal sample

sizes needed to afford reasonable sensitivity during intercohort comparison, following the protocol we have

outlined in Study 2.

3.3 Study 3

We followed up Studies 1 & 2 with an example of how MDCA-Pr can be applied to a corpus of vocal data

collected from real animals, by comparing the call-type combinations produced by male and female

common marmosets in a feeding context, where breeding partners were visually separated from each other

and only one dyad partner was presented with food. Our sample size was small for each bootstrap of the

data (66 call-type bigrams per sex), therefore the differences we identified are likely conservative (see Study

2). However, three female-biased call-type combinations were identified, as well as two call-type

combinations that were shared between sexes (Table 3). Post-hoc analysis revealed that the attracted call-

type combinations identified by MDCA-Pr traced how call transitions were distributed across the sexes in

the underlying data.

These results suggest that whilst some differences may exist in how male and female marmosets combine

call types, these differences are generally small. Whilst behavioural differences in feeding contexts have

been identified between the sexes of marmosets (see Section 2.3.1), both sexes engage in food sharing

with group members, meaning that within feeding contexts, both sexes may be expected to have largely

overlapping communicative goals. Moreover, the motivation to share food varies between individuals, such

as through individual-specific variation in hormonal signalling (Finkenwirth et al. 2016), and these individual-

level effects may further blur the predictability of sex on individual's motivation to communicate about food.

Before strong biological conclusions are made, an analysis of a larger sample of marmoset vocal data

would be beneficial, potentially via combining MDCA-Pr with other analytical tools to determine whether

these analyses converge on similar results (e.g. transition probability and positional bias models, see

Girard-Buttoz et al. 2022b; though the performance of these methods should also be evaluated when

comparing animal cohorts).

Importantly, Study 3 identified further considerations for how MDCA-Pr can be applied rigorously to analyse

animal call data. During intersex comparison, post-hoc analyses of underlying data were critical for

identifying whether attracted call-type combinations were produced by many individuals within a cohort

(thus suggesting a cohort-biased behaviour), or if they were produced by a specific subset of individuals

within a cohort (potentially suggesting individual-specific call-type combinations). We anticipate that this

additional methodological step is also important for characterizing signal combinatoriality in single

populations. Currently, all forms of collocation analysis pool data without consideration of the relative

contributions of different individuals within the underlying data. We recommend that in future applications

of any collocation analysis, the original data is reviewed to identify whether attracted signal combinations

are driven by the behaviours of many individuals, or potentially reflect the behaviour of a small subset of

individuals within a sample, and that conclusions are tempered accordingly.

Single Cohort	**Intercohort Comparison**

Data Collection:

DC1   Collect data from one target context, or balance sampling across different contexts (minimum total: 160 bigrams).

DC2   Collect data across many individuals, labelling which calls are within the same bout.

Data Collection:

Follow steps DC1 & DC2 but aim for larger sample sizes for both cohorts.

There is no minimum sample size, but small samples reduce statistical power and may not capture the full signal repertoire, weakening the strength of conclusions drawn from the data.

Vs

Bootstrapping Data:

B1   Run 1,000-10,000 bootstraps, using balanced block sizes at the highest hierarchical level in the data.

B2   If crossed variables are present (e.g., sex, age, classes), stratify bootstraps to sample each block equally where possible.

Bootstrapping Data:

Bootstrap data following steps B1 & B2, but for each cohort separately.

Apply MDCA-Pr to each bootstrapped dataset

Analysing Outputs:

A1   Calculate the mean collocation strength for each bigram type across bootstraps.

A2   Adjust the alpha levels used to construct confidence intervals (CI; A3) for mean collocation strength (based on the number of bigram types being compared):

$$\alpha_{Adjusted} = \alpha \ / \ \textbf{No. bigram types compared}$$

The default is a 95% CI ($\alpha = 0.05$).

A3   Construct a CI for each bigram type, using the $\alpha_{Adjusted}$ to sample the upper (U) and lower (L) percentiles of the bootstrapped distribution:

$$U = 100 \times (1 - (\alpha_{Adjusted} \ / \ 2))$$
$$L = 100 \times (\alpha_{Adjusted} \ / \ 2)$$

A4   Calculate 1SD across all mean collocation strengths for all bigram types (to act as an attraction filter).

A5   Identify signal combinations, where for a given bigram type, the CI is wholly above zero and the mean collocation strength is above the 1SD filter.

A6   Verify the number of individuals who produced each identified signal combination. The more individuals, the stronger the conclusion that the combination is shared across cohort members.

Analysing Outputs:

Perform step A1 for each cohort.

Perform step A2, where the number of comparisons is the total number of bigram types being compared between the two cohorts.

Perform step A3 for each bigram type in each cohort separately.

Perform steps A4 & A5 for each cohort separately, thus identifying each cohorts' signal combinations.

Perform step A6 to verify whether signal combinations are shared across members within cohorts (suggesting a cohort-general behaviour), or are confined to individuals (suggesting an individual-specific behaviour).

A7   Compare which signal combinations are present in both cohorts, versus those that differ between cohorts.

A8   Where cohorts share the same signal combinations, differences in attraction strength can be identified by non-overlapping CIs.

552   **Figure 3. Analytical pipeline for MDCA-Pr.** We provide a suggested analytical pipeline for applying MDCA-Pr, followed by further

553   steps to adapt this pipeline for intercohort comparison. R scripts to support implementation of MDCA-Pr can be found in this paper's

554   supplementary code.

3.4. General Discussion

Overall, we report that MDCA-Pr offers an accurate way to identify attracted signal combinations within animal data, and offers a suitable means to compare signal combinatoriality between cohorts of individuals. We provide a recommended analytical pipeline for applying MDCA-Pr to animal signal data, including during cohort comparison (Fig. 3). Within this pipeline, we provide two additional considerations for the collection and analysis of data when using MDCA-Pr:

First, researchers should be mindful of the contexts in which signals are sampled from animals during data collection. In our analysis of marmoset call data, we took a sample of data from a specific context (feeding). We advise that researchers evaluate signal combinatoriality separately across different contexts, as animals may reuse the same signals but combine them in context-specific ways (Berthet et al. 2025; Girard-Buttoz et al. 2025), and this may not be identifiable if data is pooled across contexts. However, if researchers do choose to pool data, we recommend that researchers ensure different contexts are sampled thoroughly and equitably. Similarly, if pooling data across contexts, bootstrapping should be stratified so that each context is consistently represented during error estimation. Stratified bootstraps may also be of use if other variables shared between compared cohorts need to be controlled. For example, when comparing sexes, ensuring that individuals with specific roles, ranks, or age classes are represented.

Second, across our studies we controlled for inflated type 1 error rates during multiple comparison using a Bonferroni correction. Bonferroni corrections can keep type 1 error rates around 5% including up to very high numbers of pairwise comparisons (Bland and Altman 1995). However, these corrections come at the cost of decreasing statistical power with increasing numbers of pairwise comparisons. Thus, when comparing many different bigram types, analyses may be more likely to result in type 2 errors (where valid attraction is not identified within a sample corpus, or a valid difference in attraction is not identified between cohorts). Given our analysis generally showed both high selectivity and sensitivity given reasonable sample sizes, we anticipate Bonferroni corrections will be sufficient in most cases, particularly when making similar numbers of comparisons to those in Studies 1 and 2. However, when working with small samples, or when making many more comparisons, other forms of family-wise error correction could be considered that have

less severe consequences for type 2 error rates (e.g. the Holm-Bonferroni method, the Šidák mothod, or

Tukey's method; see Midway et al. 2020).

In conclusion, we demonstrate that MDCA-Pr provides a robust method for identifying attracted signal

combinations in animal datasets, overcoming many of the statistical limitations associated with existing

approaches to animal collocation analysis. In addition, MDCA-Pr enables direct comparisons of signal

attraction across animal cohorts, offering a powerful tool for detecting intraspecific variation in the dynamics

of signal combinatoriality within animal communication systems. As with other forms of collocation analysis,

MDCA-Pr does not identify the behavioural responses elicited by signal combinations; consequently,

attracted signal combinations require further empirical investigation to determine their information content.

Nonetheless, as an initial analytical step, the enhanced statistical rigour of MDCA-Pr for identifying signal

attraction provides a strong foundation for future examinations of how signal combinatoriality contributes to

animal communication, including its potential roles in compositionality (Suzuki et al. 2016; Leroux et al.

2023; Berthet et al. 2025) and pragmatic inference (Schlenker et al. 2016a).

References

Agamaite JA, Chang C-J, Osmanski MS, Wang X (2015) A quantitative acoustic analysis of the vocal
597   repertoire of the common marmoset (*Callithrix jacchus*). J. Acoust. Soc. Am. 138:2906–2928.
598   https://doi.org/10.1121/1.4934268

599   Bartsch S (2004) Structural and Functional Properties of Collocations in English: A corpus study of lexical
600   and pragmatic constraints on lexical co-occurence. Gunter Narr Verlag, Tübingen

601   Berthet M, Mesbahi G, Pajot A, et al (2019) Titi monkeys combine alarm calls to create probabilistic
602   meaning. Sci. Adv. 5:eaav3991. https://doi.org/10.1126/sciadv.aav3991

603   Berthet M, Surbeck M, Townsend SW (2025) Extensive compositionality in the vocal system of bonobos.
604   Science 388:104–108. https://doi.org/10.1126/science.adv1170

605   Bland JMartin, Altman DG (1995) Multiple significance tests: the Bonferroni method. BMJ 310:170

606   Bosshard AB, Burkart JM, Merlo P, et al (2024) Beyond bigrams: call sequencing in the common
607   marmoset (*Callithrix jacchus*) vocal system. R. Soc. Open Sci. 11:240218.
608   https://doi.org/10.1098/rsos.240218

609   Bosshard AB, Leroux M, Lester NA, et al (2022) From collocations to call-locations : using linguistic
610   methods to quantify animal call combinations. Behav. Ecol. Sociobiol. 76:1–8.
611   https://doi.org/10.1007/s00265-022-03224-3

Box H, Yamamoto ME, Lopes FA (1999) Gender Differences in Marmosets and Tamarins: Responses to
Food Tasks. Int. J. Comp. Psychol. 12:59–70. https://doi.org/10.46867/C42G67

Briseño-Jaramillo M, Sosa-López JR, Ramos-Fernández G, Lemasson A (2022) Flexible use of contact
calls in a species with high fission–fusion dynamics. Phil. Trans. R. Soc. B 377:20210309.
https://doi.org/10.1098/rstb.2021.0309

Church K, Gale W, Hanks P, Hindle D (1991) Using Statistics in Lexical Analysis. In: Zernik U (ed) Lexical
Acquisition: Exploiting On-Line Resources to Build a Lexicon, 1st edn. Psychology Press, New
York, NY, pp 115–164

Collier K, Bickel B, Van Schaik CP, et al (2014) Language evolution: syntax before phonology? Proc. R.
Soc. B 281:20140263. https://doi.org/10.1098/rspb.2014.0263

Collier K, Radford AN, Stoll S, et al (2020) Dwarf mongoose alarm calls: investigating a complex non-
human animal call. Proc. R. Soc. B 287:20192514. https://doi.org/10.1098/rspb.2019.2514

Collier K, Townsend SW, Manser MB (2017) Call concatenation in wild meerkats. Anim. Behav. 134:257–
269. https://doi.org/10.1016/j.anbehav.2016.12.014

Coye C, Ouattara K, Arlet ME, et al (2018a) Flexible use of simple and combined calls in female
Campbell's monkeys. Anim Behav 141:171–181. https://doi.org/10.1016/j.anbehav.2018.05.014

Coye C, Townsend S, Lemasson A (2018b) From animal communication to linguistics and back: Insight
from combinatorial abilities in monkeys and birds. In: Boë L-J, Fagot J, Perrier P, Schwartz J-L
(eds) Origins of Human Language: Continuities and Discontinuities with Nonhuman Primates.
Peter Lang GmbH, Berlin

Demartsev V, Averly B, Johnson-Ulrich L, et al (2024) Mapping vocal interactions in space and time
differentiates signal broadcast versus signal exchange in meerkat groups. Phil. Trans. R. Soc. B
379:20230188. https://doi.org/10.1098/rstb.2023.0188

Engesser S, Holub JL, O'Neill LG, et al (2019) Chestnut-crowned babbler calls are composed of
meaningless shared building blocks. Proc. Nat. Acad. Sci. U.S.A. 116:19579–19584.
https://doi.org/10.1073/pnas.1819513116

Engesser S, Ridley AR, Townsend SW (2016) Meaningful call combinations and compositional processing
in the southern pied babbler. Proc. Nat. Acad. Sci. U.S.A. 113:5976–5981.
https://doi.org/10.1073/pnas.1600970113

Engesser S, Ridley AR, Townsend SW (2017) Element repetition rates encode functionally distinct
information in pied babbler 'clucks' and 'purrs.' Anim. Cogn. 20:953–960.
https://doi.org/10.1007/s10071-017-1114-6

Engesser S, Ridley AR, Watson SK, et al (2024) Seeds of language-like generativity in bird call
combinations. Proc. R. Soc. B 291:20240922. https://doi.org/10.1098/rspb.2024.0922

Engesser S, Townsend SW (2019) Combinatoriality in the vocal systems of nonhuman animals. WIREs
Cogn. Sci. 10:e1493. https://doi.org/10.1002/wcs.1493

Evert S (2009) Corpora and collocations. In: Lüdeling A, Kytö M (eds) Corpus Linguistics. An International
Handbook. Mouton de Gruyter, pp 1212–1248

Finkenwirth C, Martins E, Deschner T, Burkart JM (2016) Oxytocin is associated with infant-care behavior
and motivation in cooperatively breeding marmoset monkeys. Horm. Behav. 80:10–18.
https://doi.org/10.1016/j.yhbeh.2016.01.008

Fitch WT (2018) What animals can teach us about human language: the phonological continuity
hypothesis. Curr. Opin. Behav. Sci. 21:68–75. https://doi.org/10.1016/j.cobeha.2018.01.014

Freymann E, d'Oliveira Coelho J, Hobaiter C, et al (2024) Applying collocation and APRIORI analyses to
chimpanzee diets: Methods for investigating nonrandom food combinations in primate self-
medication. Am. J. Primatol. 86:e23603. https://doi.org/10.1002/ajp.23603

Girard-Buttoz C, Bortolato T, Laporte M, et al (2022a) Population-specific call order in chimpanzee
greeting vocal sequences. iScience 25:104851. https://doi.org/10.1016/j.isci.2022.104851

Girard-Buttoz C, Neumann C, Bortolato T, et al (2025) Versatile use of chimpanzee call combinations
promotes meaning expansion. Sci. Adv. 11:eadq2879. https://doi.org/ 10.1126/sciadv.adq2879

Girard-Buttoz C, Zaccarella E, Bortolato T, et al (2022b) Chimpanzees produce diverse vocal sequences
with ordered and recombinatorial properties. Commun. Biol. 5:410.
https://doi.org/10.1038/s42003-022-03350-8

Gries S (2024) Collostructional analysis: Computing the degree of association between words and
words/constructions v4.1. https://www.stgries.info/teaching/groningen/coll.analysis.r accessed
10th August 2025

Gries STh (2022) Toward more careful corpus statistics: uncertainty estimates for frequencies,
dispersions, association measures, and more. RMAL 1:100002.
https://doi.org/10.1016/j.rmal.2021.100002

Gries STh (2023) Overhauling Collostructional Analysis: Towards More Descriptive Simplicity and More
Explanatory Adequacy. Cogn. Semant. 9:351–386. https://doi.org/10.1163/23526416-bja10056

Gu H, Sun C, Gong L, et al (2023) Sex ratio potentially influence the complexity of social calls in
Himalayan leaf-nosed bat groups. Front. Ecol. Evol. 11:955540.
https://doi.org/10.3389/fevo.2023.955540

Guerreiro Martins EM, Moura AC d. A, Finkenwirth C, et al (2019) Food sharing patterns in three species
of callitrichid monkeys (*Callithrix jacchus, Leontopithecus chrysomelas, Saguinus midas*):
Individual and species differences. J. Comp. Psychol. 133:474–487.
https://doi.org/10.1037/com0000169

Hedwig D, Kohlberg A (2024) Call combination in African forest elephants *Loxodonta cyclotis*. PLoS ONE
19:e0299656. https://doi.org/10.1371/journal.pone.0299656

Idsardi WJ (2019) Some cautions regarding the phonological continuity hypothesis. Phil. Trans. R. Soc. B
375:20190050. https://doi.org/10.1098/rstb.2019.0050

Le Floch A, Girard-Buttoz C, Azaiez TS, et al (2026) Rule-based sequences in sooty mangabey vocal
communication. R. Soc. Open. Sci. 13(2):251944. https://doi.org/10.1098/rsos.251944

Lee D, Kim J, Skinner C (2019) Within-cluster resampling for multilevel models under informative cluster
size. Biometrika 106:965–972. https://doi.org/10.1093/biomet/asz035

Lehecka T (2015) Collocation and colligation. In: Handbook of Pragmatics. John Benjamins Publishing
Company, pp 1–20. https://doi.org/10.1075/hop.19.col2

Leroux M, Bosshard AB, Chandia B, et al (2021) Chimpanzees combine pant hoots with food calls into
larger structures. Anim. Behav. 179:41–50. https://doi.org/10.1016/j.anbehav.2021.06.026

Leroux M, Chandia B, Bosshard AB, et al (2022) Call combinations in chimpanzees: a social tool? Behav.
Ecol. 33:1036–1043. https://doi.org/10.1093/beheco/arac074

Leroux M, Schel, Anne M, Wilke, Claudia, et al (2023) Call combinations and compositional processing in
wild chimpanzees. Nat. Commun. 14:1–8. https://doi.org/10.1038/s41467-023-37816-y

Leroux M, Townsend SW (2020) Call combinations in great apes and the evolution of syntax. AB&C
7:131–139. https://doi.org/10.26451/abc.07.02.07.2020

Liao S, Gries STh, Wulff S (2024) Transfer five ways: applications of multiple distinctive collexeme
analysis to the dative alternation in Mandarin Chinese. CLLT. https://doi.org/10.1515/cllt-2024-
0033

Meshinska K, Burkart JM, Bell MBV, Wierucka K (2024) Echoes of self: Understanding acoustic structure
and informational content in common marmoset (*Callithrix jacchus*) phee sequences. bioRxiv.
https://doi.org/10.1101/2024.04.14.589400

Midway S, Robertson M, Flinn S, Kaller M (2020) Comparing multiple comparisons: practical guidance for
choosing the best multiple comparisons test. PeerJ 8:e10387. https://doi.org/10.7717/peerj.10387

Mine JG, Dees LC, Wilke C, et al (2025) Chimpanzee mothers, but not fathers, influence offspring vocal–
visual communicative behavior. PLoS Biol. 23:e3003270.
https://doi.org/10.1371/journal.pbio.3003270

Mine JG, Wilke C, Zulberti C, et al (2024) Vocal-visual combinations in wild chimpanzees. Behav. Ecol.
Sociobiol. 78:108. https://doi.org/10.1007/s00265-024-03523-x

Mircheva M, Brügger R, Burkart JM (2026) Who says what when? Patterns in captive common marmoset
(*Callithrix jacchus*) volubility. https://doi.org/10.64898/2026.01.24.701497

Mustoe AC, Harnisch AM, Hochfelder B, et al (2016) Inequity aversion strategies between marmosets are
influenced by partner familiarity and sex but not by oxytocin. Anim. Behav. 114:69–79.
https://doi.org/10.1016/j.anbehav.2016.01.025

Nesselhauf N (2005) Collocations in a Learner Corpus. John Benjamins B.V., Amsterdam, The
Netherlands

Nowak MA, Plotkin JB, Jansen VAA (2000) The evolution of syntactic communication. Nature 404:495–
498. https://doi.org/10.1038/35006635

R Core Team (2022) R: A language and environment for statistical computing. https://www.R-project.org/

Schamberg I, Surbeck M, Townsend SW (2024) Cross-population variation in usage of a call combination:
evidence of signal usage flexibility in wild bonobos. Anim. Cogn. 27:58.
https://doi.org/10.1007/s10071-024-01884-4

Schlenker P, Chemla E, Schel AM, et al (2016a) Formal monkey linguistics. Theor. Linguist. 42:1–90.
https://doi.org/10.1515/tl-2016-0001

Schlenker P, Chemla E, Schel AM, et al (2016b) Formal monkey linguistics: The debate. Theor. Linguist.
42:173–201. https://doi.org/10.1515/tl-2016-0010

Schlenker P, Chlema E, Arnold K, et al (2014) Monkey semantics: two "dialects" of Campbell's monkey
alarm calls. Linguist. Philos. 37:439–501. https://doi.org/10.1007/s

Schlenker P, Salis A, Leroux M, et al (2024) Minimal Compositionality *versus* Bird Implicatures: two
theories of ABC-D sequences in Japanese tits. Biol. Rev. brv.13068.
https://doi.org/10.1111/brv.13068

Spiess S, Mylne HK, Engesser S, et al (2022) Syntax-like Structures in Maternal Contact Calls of
Chestnut-Crowned Babblers (*Pomatostomus ruficeps*). Int. J. Primatol.
https://doi.org/10.1007/s10764-022-00332-9

Stefanowitsch A, Gries STh (2003) Collostructions: Investigating the interaction of words and
constructions. IJCL 8:209–243. https://doi.org/10.1075/ijcl.8.2.03ste

Stefanowitsch A, Gries STh (2005) Covarying collexemes. CLLT 1:1–43.
https://doi.org/10.1515/cllt.2005.1.1.1

Suzuki TN, Wheatcroft D, Griesser M (2016) Experimental evidence for compositional syntax in bird calls.
Nat. Commun. 7:10986. https://doi.org/10.1038/ncomms10986

Suzuki TN, Zuberbühler K (2019) Animal syntax. Curr. Biol. 29:R669–R671.
https://doi.org/10.1016/j.cub.2019.05.045

Templeton CN, Greene E, Davis K (2005) Allometry of Alarm Calls: Black-Capped Chickadees Encode
Information About Predator Size. Science 308:1934–1937.
https://doi.org/10.1126/science.1108841

Townsend SW, Engesser S, Stoll S, et al (2018) Compositionality in animals and humans. PLoS Biol.
16:e2006425. https://doi.org/10.1371/journal.pbio.2006425

van Boekholt B, Bosshard AB, Pika S (2025) Sequence organization of mother–infant interactions in
chimpanzees *(Pan troglodytes schweinfurthii)* in the wild. Proc. R. Soc. B. 292:20252271.
https://doi.org/10.1098/rspb.2025.2271

Walsh SL, Townsend SW, Engesser S, Ridley AR (2024) Call combination production is linked to the
social environment in Western Australian magpies (*Gymnorhina tibicen dorsalis*). Phil. Trans. R.
Soc. B. 379:20230198. https://doi.org/10.1098/rstb.2023.0198

Wewhare N, Krishnan A (2024) Individual-specific associations between warble song notes and body
movements in budgerigar courtship displays. BiO 13:bio060497.
https://doi.org/10.1242/bio.060497

Xiao R, McEnery T (2006) Collocation, Semantic Prosody, and Near Synonymy: A Cross-Linguistic
Perspective. Appl. Linguist. 27:103–129. https://doi.org/10.1093/applin/ami045

Zuberbühler K (2019) Syntax and compositionality in animal communication. Phil. Trans. R. Soc. B.
375:20190062. https://doi.org/10.1098/rstb.2019.0062