Longitudinal and stability-aware analysis reveals treatment-specific microRNA response signatures following immune-reconstitution and B-cell-targeted therapies in multiple sclerosis

Nasar Ata[1, 2], Joshua S. Mytych[3], Mirela Cerghet[1], Ramandeep Rattan[4], Sumit Govil[3], Shailendra Giri[1,*] and Yang Mao-Draayer[3,*]

[1]Department of Neurology, Henry Ford Health, Detroit, USA.

[2]School of Life and Basic Science Jaipur National University, Jaipur, India.

[3]Oklahoma Medical Research Foundation, Oklahoma City, OK, USA.

[4]Women's Health Services, Henry Ford Health, Detroit, MI, 48202, USA.

*These authors contributed equally to this work as principal investigators.

Corresponding Author

*Shailendra Giri, PhD
Department of Neurology
Henry Ford Health, Detroit, USA
Sgiri1@hfhs.org

Yang Mao-Draayer, MD
Multiple Sclerosis Center of Excellence Oklahoma Medical Research Foundation,
Department of Microbiology and Immunology,
University of Oklahoma Health Sciences Center,
Oklahoma City, Oklahoma.
Yang-Mao-Draayer@omrf.org

Abstract

Disease-modifying therapies (DMTs) for relapsing-remitting multiple sclerosis (RRMS) act through distinct immunological mechanisms, yet the within-patient molecular response programs associated with these therapies remain incompletely defined. Here, we reanalyzed publicly available PBMC miRNA microarray data (GSE230064) using a longitudinal, robustness-focused framework to compare therapy-associated miRNA response patterns following cladribine versus ocrelizumab treatment. Baseline (t0) and 6-month post-treatment (t1) samples were paired within individuals and technical replicates consolidated prior to analysis, yielding a final paired cohort of 4 cladribine-treated and 6 ocrelizumab-treated patients. Within each treatment arm, we quantified per-patient Δ-miRNA (t1−t0) values and prioritized therapy-associated response features using a multi-evidence framework integrating effect direction, magnitude, directional consistency across individuals, and leave-one-out sensitivity. Cladribine treatment was associated with a highly coordinated, directionally concordant upregulation of five miRNAs including hsa-miR-27a-3p, hsa-miR-27b-3p, hsa-miR-503-5p, hsa-miR-148a-3p, and hsa-miR-26a-5p, all exhibiting 100% directional stability across patients and mean Δ-expression values ranging from +0.77 to +1.38. These miRNAs target pathways relevant to MS pathophysiology, including Th17/Treg balance, Wnt-β-catenin signaling, macrophage polarization, and epigenetic immune regulation. In contrast, ocrelizumab elicited a more selective response pattern, with five miRNAs including hsa-miR-100-5p, hsa-miR-410-3p, hsa-miR-432-5p, hsa-miR-296-5p, and hsa-miR-485-3p showing moderate directional stability (83%) and greater inter-individual heterogeneity, consistent with the more targeted mechanism of CD20+ B-cell depletion. Notably, the two treatment-associated signatures were non-overlapping, with hsa-miR-27b-3p representing the only miRNA shared with prior cross-sectional analyses of this dataset. The identified ocrelizumab-associated miRNAs implicate pathways including mTOR/IGF1R signaling, NF-κB regulation, RNA editing, and mitochondrial biogenesis, several of which are dysregulated in progressive MS. Together, these findings demonstrate that cladribine and ocrelizumab induce distinct, treatment-specific miRNA response architectures that reflect their divergent immunological mechanisms. This work establishes a stability-aware analytic template for extracting reproducible longitudinal miRNA signals from small paired RRMS cohorts and provides a ranked set of biologically plausible candidate miRNAs for prospective validation and mechanistic investigation.

Introduction

Multiple sclerosis (MS) is a chronic, immune-mediated disorder of the central nervous system characterized by inflammation, neurodegeneration and demyelination (Arisi et al., 2023). Although numerous disease-modifying therapies (DMTs) are available, patients exhibit heterogeneous disease courses and treatment responses, so individualising therapy based on molecular stratification remains an unmet need (Arisi et al., 2023). MicroRNAs (miRNAs) are small (~22 nucleotide) non-coding RNAs that regulate gene expression post-transcriptionally(Zabalza, Pappolla, Comabella, Montalban, & Malhotra, 2024). Their stability in blood, plasma and cerebrospinal fluid and ease of measurement make them attractive biomarkers for disease activity and therapeutic response. Deregulated miRNA expression patterns have been proposed as diagnostic tools in neuroinflammatory diseases such as MS.

Among high-efficacy DMTs used to treat MS, cladribine and ocrelizumab represent distinct mechanistic classes. Cladribine is a purine nucleoside analogue that must be phosphorylated to become active and is selectively taken up and activated by cells of lymphoid lineage. Because lymphocytes express high levels of the enzyme deoxycytidine kinase, cladribine accumulates in B and T cells, leading to DNA incorporation, induction of apoptosis and prolonged depletion of memory B cells while sparing much of the innate immune system. This pulsed therapy produces long-term immune reconstitution with a relatively favourable safety profile (Lublin, 2005). In contrast, ocrelizumab is a humanized monoclonal antibody that binds the CD20 antigen on pre-B, mature and memory B cells (but not plasma cells or hematopoietic stem cells) and mediates depletion of these cells (Avasarala, 2017). Administration of ocrelizumab via intravenous infusion is approved for relapsing and primary-progressive MS; the therapy rapidly depletes circulating B cells and subsequently modulates other immune pathways, such as reducing inflammatory T-cell activity (Cencioni, Mattoscio, Magliozzi, Bar-Or, & Muraro, 2021).

Given the regulatory functions of miRNAs and their potential as biomarkers, characterising miRNA responses to cladribine and ocrelizumab could provide mechanistic insights and support personalised treatment strategies. However, most existing miRNA studies employ cross-sectional designs that cannot capture intra-individual dynamics. This study therefore utilises publicly available peripheral blood mononuclear cell miRNA data and a longitudinal, stability-aware analytical framework to identify treatment-specific miRNA signatures following cladribine-mediated immune reconstitution and ocrelizumab-mediated B-cell depletion

The initial study largely concentrated on cross-sectional comparisons of miRNA expression among treatment groups, whereas the current study re-evaluates the dataset utilizing a longitudinal within-subject approach. By calculating patient-specific Δ-miRNA profiles (t1−t0) and using directional consistency and leave-one-out stability measures, we uncover treatment-related miRNA signatures that are resilient to small cohort sizes and individual heterogeneity.

Methods

Dataset:

The publicly accessible Gene Expression Omnibus (GEO) with accession number GSE230064 provided the miRNA expression dataset utilized in this investigation. PBMC samples from patients with relapsing-remitting multiple sclerosis (RRMS) receiving high-efficacy disease-modifying treatments, such as ocrelizumab (OCRE; n = 14) and cladribine (CLA; n = 11), as well as untreated control individuals (CTRL; n = 15), are included in the collection. While control participants were sampled just once, samples from treated patients were taken at baseline (t0) and six months after treatment started (t1). The original trial reported that ocrelizumab was given as 300 mg intravenous infusions twice over a two-week period, whereas cladribine pills were given at a cumulative dosage of 1.75 mg/kg over five days. The collection includes 65 microarray samples and 40 distinct participants overall, which reflects the inclusion of technical duplicates and repeated measurements throughout time. Only treated patients with full paired t0–t1 samples were kept for the current longitudinal study after technical replicates were collapsed, resulting in a final cohort of 10 paired patients (4 CLA, 6 OCRE).

Table 1. **Summary of dataset composition and sample selection from GEO dataset GSE230064**

Category	Number
Total subjects	40
Cladribine-treated patients	11
Ocrelizumab-treated patients	14
Untreated control subjects	15
Total microarray samples (including technical replicates)	65
Baseline samples (t0)	33
Follow-up samples (t1)	32
Patients with complete paired samples used in longitudinal analysis	10
Cladribine paired patients	4
Ocrelizumab paired patients	6

Study Cohort and Experimental Design

The dataset comprises peripheral blood mononuclear cell (PBMC) samples collected from individuals with relapsing-remitting MS (RRMS) treated with high-efficacy disease-modifying therapies, as well as untreated control subjects. RRMS patients received either Cladribine (CLA; $n = 11$ patients) or Ocrelizumab (OCRE; $n = 14$ patients). For treated patients, blood samples were collected at baseline prior to treatment initiation (t0) and again at 6 months post-treatment (t1). Untreated control subjects ($n = 15$) were sampled once at baseline. In total, 65 samples were included in the analysis. miRNA expression profiling was performed using an

Agilent non-coding RNA microarray platform. Sample-level metadata, including treatment group, timepoint, tissue source, and cell type, were extracted directly from the GEO series matrix file. Although the GEO cohort included 11 cladribine-treated and 14 ocrelizumab-treated patients, only individuals with complete baseline (t0) and post-treatment (t1) samples were eligible for longitudinal Δ-miRNA analyses. After patient-level pairing and consolidation of technical replicates, the final paired cohort comprised 4 cladribine-treated and 6 ocrelizumab-treated patients. The statistical data were interpreted with caution due to the small number of matched patients that were included in each therapy arm within the study. Rather than strictly testing the hypothesis, the major objective of statistical testing was to prioritize the signals, and as a result, statistical significance was interpreted in conjunction with effect magnitude and stability criteria.

A detailed summary of sample characteristics and study design is provided in **Table 2**.

Table 2. Summary of study design and sample characteristics for cladribine- and ocrelizumab-treated patients included in the longitudinal Δ-miRNA analysis. The table reports sample counts, timepoints, and paired patient structure prior to consolidation of technical replicates.

Characteristic	Cladribine (CLA)	Ocrelizumab (OCRE)	Total
Total samples (including technical replicates of t0 and t1)	29	36	65
Baseline samples (t0)	15	18	33
Post-treatment samples (t1)	14	18	32
Patients with complete paired t0–t1 samples	4	6	10
Samples per patient (range)	1–3	1–3	—
Presence of technical replicates	Yes	Yes	Yes
Tissue source	Peripheral blood	Peripheral blood	—
Cell type	PBMC	PBMC	—
Expression platform	GEO microarray	GEO microarray	—

Quality Control and Preprocessing of miRNA Expression Data

Quality control (QC) analyses were performed to assess the global structure, distributional consistency, and technical integrity of miRNA expression data prior to downstream statistical analyses. miRNA expression values were extracted from the GEO series matrix file associated with accession GSE230064 and organized into a sample-by-feature expression matrix. No additional filtering, re-normalization, or batch correction was applied beyond the preprocessing performed by the original platform pipeline.

To make it easier to understand and keep the variance stable across the range of measured intensities, expression values were log2-transformed using $\log2(x + 1)$(Lin, Du, Huber, & Kibbe, 2008). We used sample-wise density plots to look at the global expression distributions across samples. This helped us make sure that the expression profiles were consistent and find any technical outliers. We did unsupervised principal component analysis (PCA) on the log2-transformed expression matrix to find the main sources of variance and any possible sample-level artifacts or batch-driven structure. PCA was performed without the inclusion of

phenotypic or treatment labels, guaranteeing an impartial evaluation of data quality(Blaise et al., 2021).

Identification of Paired Samples and Handling of Technical Replicates

To enable longitudinal within-subject analyses, patient-level pairing of samples collected at baseline (t0) and 6 months post-treatment (t1) was performed for Cladribine- and Ocrelizumab-treated individuals. We got sample metadata from the GEO series matrix file. This included the treatment assignment, timepoint, and unique sample identifiers(Clough et al., 2024). A consistent patient-level identification was created by taking common prefixes from sample titles and using them to group together several technical measures that belonged to the same person. For every patient who received therapy, samples were organized by treatment arm and patient identifier. To be included in paired analyses, there had to be at least one baseline (t0) sample and one post-treatment (t1) sample. When there were more than one technical replicate for a patient and timepoint, the number of replicates was recorded to check for technical redundancy. All patients involved in subsequent analyses possessed complete t0–t1 pairings, and none were omitted due to absent timepoints.

Technical Replicates and Computation of Longitudinal miRNA Changes

To measure changes in miRNA expression over time in the same patient after therapy, technical replicates that were taken at the same time and for the same patient were combined before the statistical analysis(Kahraman et al., 2018). We got the miRNA expression values from the processed expression matrix and changed them to log2(x + 1) to make the variance more stable and lower the effect of very high or low intensity values. For each patient i, miRNA j, and time point t (baseline t0 or post-treatment t1), we combined many technical replicates by finding their arithmetic mean(Kahraman, 2018 #48). Formally, the collapsed expression value was defined as:

$$\bar{E}_{i,j,t} = \frac{1}{n_{i,t}} \sum_{k=1}^{n_{i,t}} \log_2 \left(E_{i,j,t,k} + 1 \right)$$

where $E_{i,j,t,k}$E_{i,j,t,k}$E_{i,j,t,k}$ denotes the raw expression value of miRNA *j* for the *k*-th technical replicate of patient *i* at timepoint *t*, and $n_{i,t}$n_{i,t}$n_{i,t}$ represents the number of available technical replicates for that patient and timepoint.

Longitudinal miRNA expression changes were then computed on a per-patient basis as the difference between post-treatment and baseline expression levels(Blaise, 2021 #47):

$$\Delta_{\{i,j\}} = \bar{E}_{\{i,j,t^1\}} - \bar{E}_{\{i,j,t^0\}}$$

where $\Delta_{i,j}$\Delta_{\{i,j\}}$\Delta_{i,j}$ represents the within-subject change in expression of miRNA *j* for patient *i*. Positive Δ values indicate increased expression following treatment, whereas negative Δ values indicate decreased expression(Baran, Kula, & Erciyes, 2013). Δ-miRNA matrices were computed separately for Cladribine-treated and Ocrelizumab-treated patients and retained in comma-separated value (CSV) format for downstream statistical testing and visualization.

To identify treatment-associated miRNA changes, we performed statistical testing on within-subject Δ-miRNA values derived from paired baseline (t0) and post-treatment (t1) samples. For each patient and each miRNA, Δ-expression was computed as the difference between

post-treatment and baseline expression levels (Δ = t1 − t0), following collapse of technical replicates by the mean on a $\log_2(x+1)$ scale.

Separate analyses were performed for each treatment arm; Cladribine and Ocrelizumab, to prevent confounding molecular effects specific to each drug. A one-sample t-test was conducted for each miRNA within each treatment group to see if the mean Δ-expression substantially deviated from zero, in line with the null hypothesis of no systematic treatment-related change.

For each miRNA, the following statistics were computed: mean Δ-expression, standard deviation of Δ, t-statistic, and nominal p-value. Multiple-testing correction was performed using the Benjamini–Hochberg false discovery rate (FDR) procedure across all tested miRNAs within each treatment arm(Baran et al., 2013). Due to the limited sample size and the presence of miRNAs with insufficient within-subject variance, FDR estimates were conservatively interpreted and reported primarily for transparency. All statistical analyses were implemented in Python using SciPy and stats models libraries.

Stability and consistency analysis of treatment-associated Δ-miRNAs

To assess the robustness of treatment-associated Δ-miRNAs, a stability and consistency analysis was performed using subject-level Δ-miRNA matrices. Analyses were conducted separately for cladribine and ocrelizumab treatment arms.

We measured directional consistency for each miRNA by finding the percentage of patients whose Δ-expression was in the same direction as the group-level mean Δ. This statistic gives an easy-to-understand way to tell if changes seen in therapy are common to all participants or only to a few (Gavin, 2008). We also did a leave-one-out (LOO) sensitivity analysis by taking away one individual at a time and calculating the mean Δ-expression again(Agrawal & Domke). We looked just how sensitive group-level estimations are to individual participants by looking at how different LOO means were. Before using biological interpretation, these extra measures were utilized to check the stability of the top-ranked Δ-miRNAs.

We used a multi-evidence framework to combine the data to come up with strong treatment-related miRNA signatures. For each treatment arm (cladribine and ocrelizumab), miRNAs were maintained only if they exhibited: Consistent longitudinal change (Δ = t1 − t0), (ii) high ranking based on impact size, and (iii) substantial directional stability across individuals. This integrated method puts more value on consistency and repeatability than on standalone statistical significance. It produces treatment-specific miRNA signatures that show sustained longitudinal molecular responses.

Results

Several additional quality control (QC) investigations were carried out to assess the overall quality and global structure of the miRNA expression data. All samples showed substantially overlapping distributions in sample-wise density plots of log2-transformed expression levels (Ben-Elazar et al., 2021), suggesting that there was no significant technical bias and that global expression patterns were consistent (**Figure 1A**). The dataset's general comparability was supported by the fact that no sample showed an aberrant density pattern or a significantly altered distribution.

Figure 1: (A) Sample-wise density distributions of log2-transformed miRNA expression values across all samples, showing highly comparable global expression profiles. **(B)** Unsupervised principal component analysis (PCA) of log2-transformed miRNA expression values, demonstrating absence of extreme outliers or dominant technical effects.

The log2-transformed expression matrix was subjected to unsupervised principal component analysis (PCA), as previously demonstrated(Yang, Liu, Lu, Riggs, & Wu, 2017), which showed a continuous distribution of samples along the first two principal components. PC1 and PC2 accounted for a substantial proportion of the total variance, and was without evidence of dominant technical effects (Cameron, Apletalina, & Lindberg, 2002) **(Figure 1B).** Crucially, neither a dominating technical axis nor any individual samples showed up as severe outliers as previously seen(Aguinis, Gottfredson, & Joo, 2013), indicating that batch effects or technical abnormalities were not the primary causes of global variation. The consistency of median expression levels and interquartile ranges across samples was further supported by the overlapping distributional characteristics observed across samples.

Patient-Level Pairing and Longitudinal Sample Structure

Patient-level pairing identified a subset of treated individuals with complete longitudinal sampling suitable for within-subject analysis. From the full treated cohort, only patients with complete paired baseline and post-treatment samples were retained for longitudinal Δ-miRNA analyses, resulting in 4 cladribine-treated and 6 ocrelizumab-treated individuals. Six patients in the group treated with ocrelizumab satisfied the same requirements, whereas four patients in the group treated with cladribine had full baseline and post-treatment sample pairs. At least one technical replication was present for each timepoint in all paired patients, and few patients had more than one technical measurement at baseline and follow-up.

The number of patients with complete pairs and the existence of technical replicates at each timepoint are highlighted in **Table 3**, which provides an overview of the paired sample structure.

Table 3. **Summary of paired samples included in longitudinal analyses**

Treatment	Number of patients	Complete t0–t1 pairs	Patients with technical replicates
Cladribine (CLA)	4	4	4
Ocrelizumab (OCRE)	6	6	6

This paired cohort formed the basis for subsequent longitudinal analyses assessing within-subject miRNA expression changes following treatment.

Δ-miRNA profiles following cladribine and ocrelizumab treatment

Volcano plot analysis of Δ-miRNA values following cladribine treatment revealed a pronounced and structured pattern of miRNA changes **(Figure 2A)**. The volcano plot for cladribine (Figure 2A) shows strong changes in many miRNAs after treatment. Several miRNAs, including *hsa-miR-503-5p*, *hsa-miR-26a-5p*, *hsa-miR-27a-3p*, *hsa-miR-148a-3p*, and *hsa-miR-27b-3p*, show high increases in expression with strong significance. Most of the important changes are in the positive direction, while fewer miRNAs show decreases. Overall, this indicates that cladribine causes a broad and strong miRNA response. Large positive Δ-expression values with significant statistical support were shown by a group of miRNAs (Migdalska-Sęk et al., 2021), which clearly separated from the background distribution that was centred around zero. The overall distribution indicated a coherent treatment-associated shift rather than random variability. The volcano plot for ocrelizumab **(Figure 2B)** shows that most miRNAs do not change strongly after OCR treatment, as they are clustered near the center with low significance. However, a few miRNAs, such as *hsa-miR-100-5p*, *hsa-miR-410-3p*, and *hsa-miR-432-5p*, show moderate increases in expression along with relatively higher significance. There are very few miRNAs showing strong decreases, indicating that the changes are mainly in one direction (increase). Overall, this suggests that ocrelizumab leads to selective and mild changes in specific miRNAs, rather than a broad or strong response across many miRNAs.

Figure 2: (A) Δ-miRNA volcano plot following cladribine (CLA) treatment**. (B)** Δ-miRNA volcano plot following ocrelizumab (OCRE) treatment. Volcano plots show within-subject miRNA expression changes following cladribine or ocrelizumab treatment. All data points correspond to individual miRNAs, displayed by mean Δ-expression (t1 − t0) versus −log10(p-value). Representative miRNAs exhibiting the strongest treatment-associated changes are labelled. Fewer miRNAs exhibited large effect sizes, and the overall spread of Δ-expression values was narrower following cladribine treatment. Nevertheless, the volcano plot demonstrated a directional trend, indicating selective miRNA modulation following therapy. In contrast, the Δ-miRNA profile following ocrelizumab treatment showed a more modest pattern of change.

Top treatment-associated Δ-miRNAs

A concise summary of the representative miRNAs showing the strongest longitudinal changes and statistical support for each treatment group is provided in **Table 4**. This table presents representative miRNAs with the most significant Δ-expression values and the strongest statistical validation within each group offering a concise summary of treatment-related signals. Together with the volcano plots, these findings demonstrate distinct molecular response patterns between cladribine and ocrelizumab at the miRNA level, similar to conclusions from the initial paper.

Table 4. Selected Δ-miRNAs associated with cladribine and ocrelizumab treatment:

Treatment	miRNA	Mean Δ (t1–t0)	SD Δ	t-statistic	p-value	FDR (q-value)
CLA	hsa-miR-503-5p	0.952332	0.088553	21.508778	2.20×10^{-4}	0.134808
CLA	hsa-miR-26a-5p	0.765690	0.093851	16.317194	5.01×10^{-4}	0.141283
CLA	hsa-miR-27a-3p	0.955557	0.139707	13.679444	8.45×10^{-4}	0.141283
CLA	hsa-miR-148a-3p	0.861446	0.129696	13.284068	9.22×10^{-4}	0.141283
CLA	hsa-miR-27b-3p	1.376618	0.250515	10.990301	1.61×10^{-3}	0.197760
OCRE	hsa-miR-100-5p	1.063450	0.754590	3.452085	1.82×10^{-2}	0.752265
OCRE	hsa-miR-410-3p	1.267997	1.090543	2.848071	3.59×10^{-2}	0.752265
OCRE	hsa-miR-432-5p	1.359277	1.201364	2.771462	3.93×10^{-2}	0.752265
OCRE	hsa-miR-296-5p	0.612689	0.562015	2.670351	4.43×10^{-2}	0.752265
OCRE	hsa-miR-485-3p	1.347491	1.239347	2.663228	4.47×10^{-2}	0.752265

Table 4 summarizing representative miRNAs showing treatment-associated changes following cladribine (CLA) and ocrelizumab (OCRE) therapy. Mean Δ-expression (t1 − t0), standard deviation, test statistic, and nominal p-values are reported for each miRNA; false discovery rate (FDR) values are provided for transparency but interpreted conservatively due to limited sample size.

Ranking of Δ-miRNAs following cladribine and ocrelizumab treatment

Ranking of Δ-miRNAs following cladribine treatment, as previously done, weidentified a subset of miRNAs exhibiting the largest and most consistent expression changes across patients (Chiricosta, Blando, D'Angiolini, Gugliandolo, & Mazzon, 2023) (**Figure 3A**). These miRNAs showed high mean Δ-expression values, reinforcing the structured response pattern observed in the volcano analysis (Cipriano, Schepici, Mazzon, & Anchesi, 2024). Several of the top-ranked miRNAs belonged to related regulatory families, suggesting coordinated transcriptional modulation rather than isolated effects. A number of the cladribine-associated miRNAs are members of similar regulatory families, such as immune-regulatory miRNA clusters and the miR-27 family, indicating coordinated control of immune signalling pathways. The biological plausibility of the noted longitudinal alterations is strengthened by such family-level trends.

In contrast, ranking of Δ-miRNAs following ocrelizumab (De Felice et al., 2025; Torres-Iglesias et al., 2025) treatment revealed a more selective response profile **(Figure 3B).** The top-ranked miRNAs demonstrated moderate but consistent Δ-expression changes, reflecting a subtler molecular response compared with cladribine (Bonato et al., 2025). Despite the reduced magnitude, the ranked miRNAs highlight specific treatment-associated signals that are also unqiue to the specific treatment, no top miRNA's were found to overlap.

Figure 3 (A) Top-ranked Δ-miRNAs following cladribine (CLA) treatment, ordered by mean within-subject Δ-expression (t1 − t0) across paired patients. **(B)** Top-ranked Δ-miRNAs following ocrelizumab (OCRE) treatment, ordered by mean Δ-expression (t1 − t0). Bars represent the magnitude of mean within-subject change for each miRNA.

Figure 4 (A) Directional consistency of top-ranked Δ-miRNAs following cladribine (CLA) treatment, expressed as the percentage of patients exhibiting Δ-expression in the same direction as the group mean. **(B)** Directional consistency of top-ranked Δ-miRNAs following ocrelizumab (OCRE) treatment. Bars indicate the proportion of subjects showing concordant directionality for each miRNA.

Stability of Δ-miRNAs following cladribine and ocrelizumab treatment

Stability analysis revealed a high degree of directional consistency among the top-ranked Δ-miRNAs following cladribine treatment **(Figure 4A)**. Most miRNAs demonstrated consistent directionality across all or nearly all patients, indicating that observed Δ-expression changes were not driven by single individuals (Dimesa, Coban, & Shoari, 2026). This pattern indicates high directional consistency among cladribine-associated Δ-miRNAs across patients.

On the other hand, stability patterns of Δ-miRNAs linked to ocrelizumab therapy were more variable **(Figure 4B)**. More inter-individual heterogeneity was reflected by the moderate consistency of certain miRNAs across patients, whereas others exhibited high directional consistency. This trend is consistent with ocrelizumab's more focused and specific molecular actions compared to other anti-CD20 therapies (Cree, Berger, & Greenberg, 2025, Gabelić et al., 2021; Lamb, 2022)). Compact and treatment-specific miRNA signatures for cladribine (CLA) and ocrelizumab (OCRE) were obtained by integrating statistical evidence, ranking, and

stability measures. These miRNAs are the most repeatable longitudinal signals found throughout the analytical process(Dong et al., 2013).

Table 5: Final robust treatment-associated Δ-miRNA signatures derived from integrated longitudinal analysis

Table 5 summarizing miRNAs retained as robust treatment-associated signatures following integration of longitudinal change (Δ = t1 − t0), effect-size ranking, and directional stability across patients. Mean Δ-expression and directional stability percentages are shown separately for cladribine (CLA) and ocrelizumab (OCRE) treatment groups.

Treatment	miRNA	Mean Δ (t1–t0)	Directional Stability
CLA	hsa-miR-503-5p	+0.95	100
CLA	hsa-miR-26a-5p	+0.77	100
CLA	hsa-miR-27a-3p	+0.96	100
CLA	hsa-miR-148a-3p	+0.86	100
CLA	hsa-miR-27b-3p	+1.38	100
OCRE	hsa-miR-100-5p	+1.06	83
OCRE	hsa-miR-410-3p	+1.27	83
OCRE	hsa-miR-432-5p	+1.36	83
OCRE	hsa-miR-296-5p	+0.61	83
OCRE	hsa-miR-485-3p	+1.35	83

Functional Consequences of miRNA changes:

We detected unique miRNAs not identified by Arisi et al 2021 following either ocrelizumab or cladribine therapy. Of note, many are related to MS and are elevated in progressive forms of the disease as depicted in Table 6.

Table 6: Overview of potential cellular targets:

Table 6 summarizes potential miRNA cellular targets unique to either OCRE or CLA. Of note, only hsa-miR-27b-3p was common between the prior analysis by Arisi et al and our current t0- and t1-paired analysis.

Treatment	miRNA	Validated/Predicted Targets	Function in relation to MS	Citations
CLA	hsa-miR-503-5p	CDC25a, CCND1, BCL2	Not directly implicated in MS, Regulation of cell cycle and apoptosis under stress.	Sarkar et al 2010, Zhang Met al 2025
CLA	hsa-miR-26a-5p	SLC1A1 Glutamate receptor signaling; inversely correlated to DLG4	Upregulated in Natalizumab and IFNβ treatment	Potenza et al 2018, Mameli G, et al 2016,

Treatment	miRNA	Validated/Predicted Targets	Function in relation to MS	Citations
				De Felice Bet al 2014.
CLA	hsa-miR-27a-3p	PPARγ, Wnt3a, Wnt-β-catenin signaling pathway modulator	Th17/Treg balance, OPC differentiation, Linked to RRMS=> SPMS transition	Kulyté A et al 2019, Tripathi A et al 2019, Regev K, et al 2016.
CLA	hsa-miR-148a-3p	HLA-G, DNMT1	Epigenetic regulation of immune tolerance, Low serum levels inked to primary progressive MS	Reches A et al 2012, Muñoz-San Martín Met al 2022.
CLA	hsa-miR-27b-3p	PPARγ, FOXO1	Impacts lipid metabolism and Th1/Th2 balance	Kulyté Aet al 2019, D'Onofrio N, et al 2023, Guerau-de-Arellano Met al 2011.
OCRE	hsa-miR-100-5p	mTOR, IGF1R	Not directly implicated in MS, Modulates autophagy and cell survival.	Liu H, et al 2024, Pinheiro AHG, et al 2024
OCRE	hsa-miR-410-3p	STAT3, ELAVL4	Reduced in RRMS, Impairs pro-inflammatory cytokine secretion	Liu D, et al 2016, Keller A, et al 2015, Wang Yet al 2019.
OCRE	hsa-miR-432-5p	ADAR1, Syt7	Elevated in gray matter lesions and in progressive MS patients, Regulation of RNA editing and innate immune sensing.	Fritsche Let al 2020, Vistbakka Jet al 2017, Zhang X at al. 2021
OCRE	hsa-miR-296-5p	IKBKE, NumbL, STAT5a	Regulated by circular RNA linked to MS, Modulates NF-κB signaling	Li, L et al 2022, Li H, et al 2018.
OCRE	hsa-miR-485-3p	PGC-1α, STAT3	Downregulated in RRMS compared to control patients, Regulates Th17 response in experimental autoimmune encephalitis, Mitochondrial biogenesis and energy homeostasis.	Alizadeh-Fanalou Set al 2020, Ryu Iset al 2026, Xue Yet al 2021.

Discussion

Comprehensive quality control analyses confirmed that the miRNA expression dataset exhibits high technical integrity and is suitable for downstream longitudinal and treatment-specific analyses. The lack of extreme outliers in unsupervised principal component analysis and density **(Figure 1A-B)** and the strong overlap seen in sample-wise density distributions show that global expression patterns are very consistent across samples and are not dominated by batch-driven effects or technical artifacts. The observed variability appears to be mostly due to biological heterogeneity rather than noise, as this consistency is consistent with the Agilent microarray workflow's efficient platform-level preprocessing and normalization (Diaz-Herrera, 2020). Importantly, the QC analyses' absence of noticeable non-biological factor separation or clustering lends credence to the reliability of later within-subject comparisons and treatment-stratified analyses, although this behaviour is more indicative of technological resilience than biological signal loss (Panga). When taken as a whole, these QC results give assurance that the dataset provides a stable and trustworthy basis for examining long-term changes in miRNA expression linked to disease-modifying treatments, supporting the move to paired analyses that concentrate on treatment-induced molecular dynamics.

Longitudinal Pairing and Sample Structure

The rigorous criterion for full t0–t1 pairing improves statistical robustness by allowing direct within-subject comparisons, therefore decreasing confounding from inter-individual variability, even when the number of matched patients is limited. Internal validation of measurement consistency across timepoints and further support for data dependability are provided by the existence of several technical replicates for various patients (Ratitch et al., 2023). When taken as a whole, the paired sample structure shown in **Table 3** provides a solid basis for further studies that concentrate on changes in treatment-associated miRNA expression and supports the use of longitudinal difference-based methods.

The Δ-miRNA study demonstrated unique and treatment-specific molecular response patterns subsequent to cladribine and ocrelizumab therapy. **Figure 2A** demonstrates that cladribine therapy resulted in a coordinated alteration in miRNA expression, with several miRNAs showing significant positive Δ-expression and uniform directionality among patients. The inclusion of many miRNAs from related families among the most significant signals **(Table 4)** indicates organized regulatory alterations rather than random or isolated effects, aligning with the extensive immunomodulatory and immunological reconstitution processes associated with cladribine.

Conversely, the Δ-miRNA profile after ocrelizumab therapy exhibited a more subdued pattern of alteration **(Figure 2B),** with a reduced number of miRNAs demonstrating significant changes in expression. This selective response likely indicates the specific mechanism of B-cell depletion and the anticipation of more nuanced downstream transcriptional regulation, as previously shown to be the case for B cell differentiation (Laidlaw & Cyster, 2021). Although the quantity of change is diminished, the discernible directionality and consistency among the predominant miRNAs suggest that ocrelizumab medication is still linked to certain miRNA-level modifications.

Collectively, our data illustrate that cladribine and ocrelizumab elicit unique miRNA response signatures, underscoring therapy-specific molecular reconfiguration rather than a common therapeutic impact. The miRNAs listed in **Table 4** constitute a targeted selection of candidates for subsequent integrative studies and endorse the application of Δ-based, within-subject methodologies to elucidate treatment-related molecular dynamics. These results establish a molecular framework for subsequent stability-aware and machine-learning–based analyses aimed at characterizing treatment response patterns. The miRNAs found in this research most

likely represent immune remodelling mechanisms linked to therapy that take place in response to multiple sclerosis disease-modifying therapies. A number of the cladribine-associated miRNAs have previously been connected to inflammatory signalling pathways and immunological control. For instance, miR-503-5p has been linked to the control of inflammatory signalling networks and immune cell proliferation, indicating that its elevated expression may represent immunological reconstitution activities after cladribine-induced lymphocyte depletion. Similar to this, miR-26a-5p has been shown to control cytokine signalling pathways and T-cell activation, indicating that it may play a part in influencing adaptive immunological responses during treatment-associated immune rebalancing. The coordinated overexpression of members of the miR-27 family, such as miR-27a-3p and miR-27b-3p, which are recognized regulators of inflammatory responses and macrophage polarization, may suggest more extensive modulation of innate immune signalling after cladribine treatment. The miRNA profile seen after ocrelizumab therapy, on the other hand, is more selective and contains miR-100-5p, a regulator that was previously linked to immune regulatory pathways and B-cell development. This finding is in line with ocrelizumab's mode of action, which modifies downstream immune signalling networks by specifically eliminating CD20-positive B cells. When combined, these findings imply that the miRNA signatures found in this study may represent therapy-specific transcriptional regulatory programs that reflect different immune modulation mechanisms brought on by ocrelizumab-mediated B-cell depletion and cladribine-mediated immune reconstitution. These miRNAs are physiologically reasonable candidates for further research aiming at comprehending molecular responses to disease-modifying medications in multiple sclerosis, even though functional validation will be necessary.

The ranking analysis refined the Δ-miRNA signals identified in earlier analyses into treatment-specific molecular signatures. Data in **figure 3A** illustrates that cladribine therapy resulted in a significant and coordinated alteration in the expression of a specific group of miRNAs, along with extensive immunomodulatory and immune-reconstitution effects. The recurrence of miRNAs from related families among the highest-ranked candidates **(Table 4)** indicates the existence of organized regulatory responses rather than random fluctuation (Janky et al., 2014).
Conversely, ocrelizumab therapy yielded a more subdued ranking profile **(Figure 3B)**, marked by a reduced number of miRNAs exhibiting moderate effect sizes. This pattern corresponds with the intended mode of action of ocrelizumab and indicates selective downstream transcriptional regulation. The found ranking miRNAs for ocrelizumab offer insight into certain biological pathways activated post-therapy (Jain, 2020). The graded Δ-miRNA profiles indicate that cladribine and ocrelizumab elicit unique and treatment-specific miRNA response patterns.

Interpretation of stability findings

The stability analysis presented provides critical validation of the treatment-associated Δ-miRNA signals identified in earlier results. The idea that these variations represent a common molecular response rather than individual-specific fluctuations is supported by the high directional consistency of cladribine-associated miRNAs across patients, as illustrated in **(Figure 4A).** The wide-ranging immunomodulatory and immunological reconstitution effects linked to cladribine treatment are in line with this stability (Alroughani et al., 2025; Lünemann, Ruck, Muraro, Bar-Or, & Wiendl, 2020). The ocrelizumab-associated miRNA profile, on the other hand, shows significant inter-individual variability **(Figure 4B)**, indicating a more varied molecular response. This result suggests that downstream transcriptional effects may differ across individuals and is consistent with the targeted mechanism of B-cell depletion (Laidlaw & Cyster, 2021; Lee, Rojas, & Gommerman, 2021). Crucially, despite their decreased overall stability, a number of miRNAs continue to show strong directional consistency, confirming their

significance as treatment-associated signals. The final treatment-associated Δ-miRNA signatures identified in this study represent the most reproducible longitudinal molecular alterations observed across the analytical workflow. These signatures capture miRNAs that consistently respond to therapy instead of showing random variation or isolated statistical significance (Volinia & Croce, 2013). They do this by combining longitudinal change, effect magnitude, and directional stability across individuals. The cladribine-associated signature is defined by precisely coordinated miRNA regulation exhibiting consistently high directional stability among patients, signifying a collective molecular response to treatment. This pattern aligns with the extensive immunomodulatory and immune-reconstitution effects of cladribine, reinforcing the notion that these miRNAs represent authentic treatment-related regulatory mechanisms.

In contrast, the signature associated with ocrelizumab shows stable but more varied longitudinal modulation, which has also been noted in human T cells in response following Ocrelizumab treatment Although ocrelizumab-associated Δ-miRNAs exhibit greater inter-individual variability than cladribine-associated signals, several miRNAs still demonstrate consistent directional changes across patients. This suggests that the therapy induces selective but reproducible transcriptional modulation, rather than the broader coordinated response observed following cladribine treatment. These results show that strong changes in miRNA levels that are linked to treatment can be found without relying only on isolated statistical significance. By focusing on longitudinal consistency and reproducibility we developed a biologically-based way to understand therapy-specific molecular responses and a clear list of candidate miRNAs for future validation studies.

Conclusion

This study conducted a longitudinal examination of treatment-related miRNA dynamics in MS by within-subject Δ-miRNA profiling. Through the integration of statistical testing, effect-size ranking, and stability evaluation across people, we discerned consistent and treatment-specific miRNA response patterns subsequent to cladribine and ocrelizumab medication. The findings indicate that cladribine is linked to a highly coordinated miRNA response with significant directional consistency among patients, while ocrelizumab elicits a more selective and variable miRNA modulation pattern. The definitive miRNA signatures presented in **Table 4** underscore molecular alterations that are consistent among people and unlikely to be influenced by individual-specific influences. This work prioritizes repeatability and longitudinal consistency above isolated statistical significance as essential criterion for identifying physiologically relevant treatment-associated miRNA signals. These findings collectively offer a consistent molecular framework for comprehending therapy-specific immune modulation in MS and build a solid platform for future translational and mechanistic research.

Future Scope

Multiple avenues can enhance and reinforce the conclusions of this research. Initially, the validation of the discovered miRNA signatures in larger, independent cohorts is crucial to ascertain their generalizability across varied patient groups and therapeutic contexts. The inclusion of extended follow-up periods may elucidate the temporal stability of these miRNA responses and their correlation with prolonged therapy benefits. Secondly, the integration of the discovered miRNAs with downstream mRNA, proteomic, or immune-cell-specific data may clarify the molecular pathways by which these miRNAs influence treatment response. Multi-omics methodologies might enhance the biological understanding of therapy-specific regulation networks. Subsequently, future research may investigate the therapeutic significance of these miRNA signatures by analysing correlations with therapy efficacy, illness progression, or long-term results. With proper validation, persistent longitudinal miRNA

signatures may function as relevant biological markers to assist in treatment monitoring and tailored therapeutic approaches in MS. This study has several limitations. First, the number of patients with complete longitudinal pairs was small (n=4 for cladribine and n=6 for ocrelizumab), which limits statistical power. Second, the analysis relied on publicly available microarray data without access to detailed clinical outcomes, preventing correlation of miRNA changes with treatment response or disease activity. Third, only a single post-treatment timepoint (6 months) was available, limiting evaluation of temporal dynamics. Finally, functional validation of the identified miRNA candidates was beyond the scope of this study.

Source of Data

The miRNA expression data analysed in this study were obtained from the NCBI GEO # accession number GSE230064, a publicly accessible repository hosting high-throughput functional genomics datasets. The dataset comprised longitudinal miRNA expression profiles from MS patients treated with cladribine or ocrelizumab, including paired baseline and post-treatment samples. The data were downloaded from GEO, curated, and reanalysed locally for the purposes of this study.

References:

Agrawal, A., & Domke, J. *Understanding the difficulties of posterior predictive estimation.* Paper presented at the Forty-second International Conference on Machine Learning.

Aguinis, H., Gottfredson, R. K., & Joo, H. J. O. r. m. (2013). Best-practice recommendations for defining, identifying, and handling outliers. *16*(2), 270-301.

Alroughani, R., Inshasi, J., Farouk, S., Al-Asmi, A., Hassan, A., Jacob, A., . . . Therapy. (2025). Role of Immune Reconstitution Therapy with Cladribine Tablets in the Management of Relapsing Multiple Sclerosis in Older Patients. 1-16.

Alizadeh-Fanalou, S., Karimizadeh, G., Sharanloo, G., & Shayanfar, N. (2020). Dysregulation of microRNAs regulating survivin in CD4+ T cells of patients with multiple sclerosis. Journal of Neuroimmunology, 348, 577395.

Arisi, I., Malimpensa, L., Manzini, V., Brandi, R., Gosetti di Sturmeck, T., D'Amelio, C., . . . Malerba, F. J. F. i. I. (2023). Cladribine and ocrelizumab induce differential miRNA profiles in peripheral blood mononucleated cells from relapsing–remitting multiple sclerosis patients. *14*, 1234869.

Avasarala, J. J. D. t. i. (2017). Anti-CD20 cell therapies in multiple sclerosis—a fixed dosing schedule for ocrelizumab is overkill. *11*, 1177392817737515.

Baran, M., Kula, S., & Erciyes, A. J. F. (2013). *27*(4), 537-546.

Bartnik, A. J. M. S. J. (2023). ACTRIMS Forum 2023-Poster Presentations. *29*(2S), 18-242.

Ben-Elazar, S., Aure, M. R., Jonsdottir, K., Leivonen, S.-K., Kristensen, V. N., Janssen, E. A., . . . Yakhini, Z. J. P. c. b. (2021). miRNA normalization enables joint analysis of several datasets to increase sensitivity and to reveal novel miRNAs differentially expressed in breast cancer. *17*(2), e1008608.

Blaise, B. J., Correia, G. D., Haggart, G. A., Surowiec, I., Sands, C., Lewis, M. R., . . . Holmes, E. J. N. P. (2021). Statistical analysis in metabolic phenotyping. *16*(9), 4299-4326.

Bonato, M., Cerrato, V., Dioni, L., Montarolo, F., Parolisi, R., Bertolotto, A., . . . Boda, E. J. F. i. I. (2025). Short-term exposure to particulate matter triggers a selective alteration of plasma extracellular vesicle-packaged miRNAs in a mouse model of Multiple Sclerosis. *16*, 1596935.

Cameron, A., Apletalina, E., & Lindberg, I. J. T. e. (2002). The enzymology of PC1 and PC2. *22*, 291-332.

Cencioni, M. T., Mattoscio, M., Magliozzi, R., Bar-Or, A., & Muraro, P. A. J. N. R. N. (2021). B cells in multiple sclerosis—from targeted depletion to immune reconstitution therapies. *17*(7), 399-414.

Chiricosta, L., Blando, S., D'Angiolini, S., Gugliandolo, A., & Mazzon, E. J. I. J. o. M. S. (2023). A comprehensive exploration of the transcriptomic landscape in multiple sclerosis: a systematic review. *24*(2), 1448.

Cipriano, G. L., Schepici, G., Mazzon, E., & Anchesi, I. J. I. J. o. M. S. (2024). Multiple sclerosis: roles of miRNA, lcnRNA, and circRNA and their implications in cellular pathways. *25*(4), 2255.

Clough, E., Barrett, T., Wilhite, S. E., Ledoux, P., Evangelista, C., Kim, I. F., . . . Sherman, P. M. J. N. a. r. (2024). NCBI GEO: archive for gene expression and epigenomics data sets: 23-year update. *52*(D1), D138-D144.

Cree, B. A., Berger, J. R., & Greenberg, B. J. C. d. (2025). The Evolution of Anti-CD20 Treatment for Multiple Sclerosis: Optimization of Antibody Characteristics and Function. 1-20.

De Felice, B., Signoriello, E., Montanino, C., Romano, G., Archetto, D., Maida, E., . . . Farinella, F. J. N.-c. R. R. (2025). Role of Ocrelizumab in modulating gene and microRNA expression in multiple sclerosis.

De Felice, B., Mondola, P., Sasso, A., Orefice, G., Bresciamorra, V., Vacca, G., Biffali, E., Borra, M., & Pannone, R. (2014). Small non-coding RNA signature in multiple sclerosis patients after treatment with interferon-β. BMC medical genomics, 7, 26. https://doi.org/10.1186/1755-8794-7-26

Diaz-Herrera, D. M. (2020). *Methods to Integrate Genetic and Clinical Data for Disease Subtyping*: Wayne State University.

Dimesa, A. M., Coban, M. A., & Shoari, A. J. C. (2026). Matrix Metalloproteinases in Hepatocellular Carcinoma: Mechanistic Roles and Emerging Inhibitory Strategies for Therapeutic Intervention. *18*(2), 288.

Dong, H., Lei, J., Ding, L., Wen, Y., Ju, H., & Zhang, X. J. C. r. (2013). MicroRNA: function, detection, and bioanalysis. *113*(8), 6207-6233.

D'Onofrio, N., Prattichizzo, F., Martino, E., Anastasio, C., Mele, L., La Grotta, R., Sardu, C., Ceriello, A., Marfella, R., Paolisso, G., & Balestrieri, M. L. (2023). MiR-27b attenuates mitochondrial oxidative stress and inflammation in endothelial cells. Redox biology, 62, 102681. https://doi.org/10.1016/j.redox.2023.102681

Fritsche, L., Teuber-Hanselmann, S., Soub, D., Harnisch, K., Mairinger, F., & Junker, A. (2020). MicroRNA profiles of MS gray matter lesions identify modulators of the synaptic protein synaptotagmin-7. Brain pathology (Zurich, Switzerland), 30(3), 524–540. https://doi.org/10.1111/bpa.12800

Fissolo, N., Calvo-Barreiro, L., Eixarch, H., Boschert, U., Villar, L. M., Costa-Frossard, L., . . . Sabidó, E. J. F. i. i. (2023). Molecular signature associated with cladribine treatment in patients with multiple sclerosis. *14*, 1233546.

Gabelić, T., Barun, B., Adamec, I., Krbot Skorić, M., Habek, M. J. H. V., & Immunotherapeutics. (2021). Product review on MAbs (alemtuzumab and ocrelizumab) for the treatment of multiple sclerosis. *17*(11), 4345-4362.

Gavin, H. (2008). Understanding research methods and statistics in psychology.

Guerau-de-Arellano, M., Smith, K. M., Godlewski, J., Liu, Y., Winger, R., Lawler, S. E., Whitacre, C. C., Racke, M. K., & Lovett-Racke, A. E. (2011). Micro-RNA dysregulation in multiple sclerosis favours pro-inflammatory T-cell-mediated autoimmunity. Brain : a journal of neurology, 134(Pt 12), 3578–3589. https://doi.org/10.1093/brain/awr262

Jain, K. K. (2020). Personalized therapy of neurological disorders. In *Textbook of Personalized Medicine* (pp. 213-262): Springer.

Janky, R. s., Verfaillie, A., Imrichová, H., Van de Sande, B., Standaert, L., Christiaens, V., . . . Potier, D. J. P. c. b. (2014). iRegulon: from a gene list to a gene regulatory network using large motif and track collections. *10*(7), e1003731.

Kahraman, M., Röske, A., Laufer, T., Fehlmann, T., Backes, C., Kern, F., . . . Zabler, C. J. S. r. (2018). MicroRNA in diagnosis and therapy monitoring of early-stage triple-negative breast cancer. *8*(1), 11584.

Keller, A., Leidinger, P., Meese, E., Haas, J., Backes, C., Rasche, L., Behrens, J. R., Pfuhl, C., Wakonig, K., Gieß, R. M., Jarius, S., Meder, B., Bellmann-Strobl, J., Paul, F., Pache, F. C., & Ruprecht, K. (2015). Next-generation sequencing identifies altered whole blood microRNAs in neuromyelitis optica spectrum disorder which may permit discrimination from multiple sclerosis. Journal of neuroinflammation, 12, 196. https://doi.org/10.1186/s12974-015-0418-1

Kulyté, A., Kwok, K. H. M., de Hoon, M., Carninci, P., Hayashizaki, Y., Arner, P., & Arner, E. (2019). MicroRNA-27a/b-3p and PPARG regulate SCAMP3 through a feed-forward loop during adipogenesis. Scientific reports, 9(1), 13891. https://doi.org/10.1038/s41598-019-50210-3

Laidlaw, B. J., & Cyster, J. G. J. N. R. I. (2021). Transcriptional regulation of memory B cell differentiation. *21*(4), 209-220.

Lamb, Y. N. J. D. (2022). Ocrelizumab: a review in multiple sclerosis. *82*(3), 323-334.

Lee, D. S., Rojas, O. L., & Gommerman, J. L. J. N. r. D. d. (2021). B cell depletion therapies in autoimmune disease: advances and mechanistic insights. *20*(3), 179-199.

Lin, S. M., Du, P., Huber, W., & Kibbe, W. A. J. N. a. r. (2008). Model-based variance-stabilizing transformation for Illumina microarray data. *36*(2), e11-e11.

Li, H., Ouyang, X. P., Jiang, T., Zheng, X. L., He, P. P., & Zhao, G. J. (2018). MicroRNA-296: a promising target in the pathogenesis of atherosclerosis?. Molecular medicine (Cambridge, Mass.), 24(1), 12. https://doi.org/10.1186/s10020-018-0012-y

Liu, H., Chen, Y., Huang, Y., Wei, L., Ran, J., Li, Q., Tian, Y., Luo, Z., Yang, L., Liu, H., Yin, G., & Xie, Q. (2024). Macrophage-derived mir-100-5p orchestrates synovial proliferation and inflammation in rheumatoid arthritis through mTOR signaling. Journal of nanobiotechnology, 22(1), 197. https://doi.org/10.1186/s12951-024-02444-1

Liu, Y., Cheng, Z., Pan, F., Yan, W., Liang, H., & Huang, W. (2016). MicroRNA-410 inhibits cell proliferation and invasion in cholangiocarcinoma by directly targeting IL-10-activated STAT3 pathway. Biomedicine & Pharmacotherapy, 82, 599–604.

Li, L., Feng, G., Chen, T., & Zhang, L. (2022). Circ_0000514 promotes breast cancer progression by regulating the miR-296-5p/CXCL10 axis. Journal of biochemistry, 170(6), 753–761. https://doi.org/10.1093/jb/mvab097

Lublin, F. J. J. o. N. (2005). History of modern multiple sclerosis therapy. *252*(Suppl 3), iii3-iii9.

Lünemann, J. D., Ruck, T., Muraro, P. A., Bar-Or, A., & Wiendl, H. J. N. R. N. (2020). Immune reconstitution therapies: concepts for durable remission in multiple sclerosis. *16*(1), 56-62.

Mameli, G., Arru, G., Caggiu, E., Niegowska, M., Leoni, S., Madeddu, G., Babudieri, S., Sechi, G. P., & Sechi, L. A. (2016). Natalizumab Therapy Modulates miR-155, miR-26a and Proinflammatory Cytokine Expression in MS Patients. PloS one, 11(6), e0157153.

Mathias, A., Pantazou, V., Perriot, S., Canales, M., Jones, S., Oberholster, L., . . . Neuroinflammation. (2023). Ocrelizumab impairs the phenotype and function of memory CD8+ T cells: A 1-year longitudinal study in patients with multiple sclerosis. *10*(2), e200084.

Migdalska-Sęk, M., Modrzewska, B., Kordiak, J., Pastuszak-Lewandoska, D., Kiszałkiewicz, J. M., Bielec, F., . . . Brzeziańska-Lasota, E. J. S. R. (2021). Diagnostic value of PPARδ and miRNA-17 expression levels in patients with non-small cell lung cancer. *11*(1), 24136.

Muñoz-San Martín, M., Gómez, I., Quiroga-Varela, A., Gonzalez-Del Río, M., Robles Cedeño, R., Álvarez, G., Buxó, M., Miguela, A., Villar, L. M., Castillo-Villalba, J., Casanova, B., Quintana, E., & Ramió-Torrentà, L. (2022). miRNA Signature in CSF From Patients With Primary Progressive Multiple Sclerosis. Neurology(R) neuroimmunology & neuroinflammation, 10(1), e200069. https://doi.org/10.1212/NXI.0000000000200069

Panga, N. K. R. Enhancing Data Quality: Quality Control and Assurance Methods in Data Processing and Visualization. In: vol.

Pinheiro, A. H. G., Pereira, B. O., Silva, L. S. D., Melo, F. T. C., Souza, A. C. C. B., Leal, V. S. G., Figueiredo, P. B. B., Neto, J. F. A., Santos, M. C. D., Queiroz, N. N. M., Felício, K. M., Ribeiro-Dos-Santos, Â., Felício, J. S., & Cavalcante, G. C. (2024). Downregulation of hsa-miR-100-5p May Be a Protective Factor in the Early Stages of Nephropathy in Type 1 Diabetes Mellitus. International journal of molecular sciences, 25(11), 5663. https://doi.org/10.3390/ijms25115663

Potenza, N., Mosca, N., Mondola, P., Damiano, S., Russo, A., & De Felice, B. (2018). Human miR-26a-5p regulates the glutamate transporter SLC1A1 (EAAT3) expression. Relevance in multiple sclerosis. Biochimica et biophysica acta. Molecular basis of disease, 1864(1), 317–323. https://doi.org/10.1016/j.bbadis.2017.09.024Ratitch, B., Trigg, A., Majumder, M., Vlajnic, V., Rethemeier, N., & Nkulikiyinka, R. J. D. B. (2023). Clinical validation of novel digital measures: statistical methods for reliability evaluation. *7*(1), 74-91.

Reches, A., Nachmani, D., Berhani, O., Stein, N., & Mandelboim, O. (2012). miRNA-mediated control of HLA-G expression and function. PLoS ONE, 7(3), e33395.

Regev, K., Paul, A., Healy, B., von Glenn, F., Diaz-Cruz, C., Gholipour, T., Mazzola, M. A., Raheja, R., Nejad, P., Glanz, B. I., Kivisakk, P., Chitnis, T., Weiner, H. L., & Gandhi, R. (2016). Comprehensive evaluation of serum microRNAs as biomarkers in multiple sclerosis. Neurology(R) neuroimmunology & neuroinflammation, 3(5), e267. https://doi.org/10.1212/NXI.0000000000000267

Ryu, I. S., Ha, D. I., Jung, Y. J., Lee, H. J., Kim, I., Lim, Y. N., Min, H. S., Kim, S. H., Yoon, I., Cho, H. J., & Ryu, J. H. (2026). Modulation of the miR-485-3p/PGC-1α Pathway by

ASO-Loaded Nanoparticles Attenuates ALS Pathogenesis. International journal of molecular sciences, 27(2), 615. https://doi.org/10.3390/ijms27020615

Sarkar, S., Dey, B. K., & Dutta, A. (2010). MiR-322/424 and -503 are induced during muscle differentiation and promote cell cycle quiescence and differentiation by down-regulation of Cdc25A. Molecular biology of the cell, 21(13), 2138–2149. https://doi.org/10.1091/mbc.e10-01-0062

Torres-Iglesias, G., López-Molina, M., Ayala-Suárez, R., Egido, B., Laso-García, F., Chamorro, B., . . . Neuroinflammation. (2025). Extracellular Vesicle-Derived MicroRNAs as a Biomarker for Therapeutic Response in Multiple Sclerosis. *12*(4), e200420.

Tripathi, A., Volsko, C., Garcia, J. P., Agirre, E., Allan, K. C., Tesar, P. J., Trapp, B. D., Castelo-Branco, G., Sim, F. J., & Dutta, R. (2019). Oligodendrocyte Intrinsic miR-27a Controls Myelination and Remyelination. Cell reports, 29(4), 904–919.e9. https://doi.org/10.1016/j.celrep.2019.09.020

Vasadia, D. J. (2018). Functional characterization of differentially expressed miRNAs in Trematomus bernacchii acclimated to acute and chronic thermal stress.

Vistbakka, J., Elovaara, I., Lehtimäki, T., & Kuusisto, M. E. (2017). Circulating microRNAs as biomarkers in progressive multiple sclerosis. Multiple Sclerosis Journal, 23(3), 403–412. doi.org

Volinia, S., & Croce, C. M. J. P. o. t. N. A. o. S. (2013). Prognostic microRNA/mRNA signature from the integrated analysis of patients with invasive breast cancer. *110*(18), 7413-

Wang, Y., Xu, N., Zhao, S., Jiao, T., Fu, W., Yang, L., & Zhang, N. (2019). miR-410-3p Suppresses Cytokine Release from Fibroblast-Like Synoviocytes by Regulating NF-κB Signaling in Rheumatoid Arthritis. Inflammation, 42(1), 331–341. https://doi.org/10.1007/s10753-018-0896-2

Wermuth, P. J., Addya, S., & Jimenez, S. A. J. P. o. (2011). Effect of protein kinase C delta (PKC-δ) inhibition on the transcriptome of normal and systemic sclerosis human dermal fibroblasts in vitro. *6*(11), e27110.

Xue, Y., Zuo, X., Shen, T., & Wang, J. (2021). Targeting microRNA-485-3p blocks Alzheimer's disease progression by enhancing Aβ clearance and reducing neuroinflammation. International Journal of Molecular Sciences, 22(23), 13136

Yang, L., Liu, J., Lu, Q., Riggs, A. D., & Wu, X. J. B. g. (2017). SAIC: an iterative clustering approach for analysis of single cell RNA-seq data. *18*(Suppl 6), 689.

Zabalza, A., Pappolla, A., Comabella, M., Montalban, X., & Malhotra, S. J. F. i. i. (2024). MiRNA-based therapeutic potential in multiple sclerosis. *15*, 1441733.

Zhang, X., Yan, W., Xue, Y., Xu, H., Li, J., Zhao, Z., Sun, Y., Wang, Y., He, J., Huang, Y., Yu, D., Xiao, Z., & Yin, S. (2021). Roles of miR-432 and circ_0000418 in mediating the anti-depressant action of ADAR1. Neurobiology of stress, 15, 100396. https://doi.org/10.1016/j.ynstr.2021.100396

Zhang, M., Shuai, P., Wang, X., Lan, Y., Deng, K., & Zeng, Y. (2025). Systematic validation of miR-503-5p abnormal expression in diabetic vascular complications: a meta-analysis and exploration of its targeted regulatory mechanism in retinopathy. Diabetology & metabolic syndrome, 17(1), 412. https://doi.org/10.1186/s13098-025-01982-4