Using Cryogenic Electron Tomography (cryoET) to Determine Rubisco Polymerization Constants

in α-Carboxysomes

Wenxiang Cao[a,1], Kristy Rochon[b,1], Ryan H. Gray[c], Luke Oltrogge[d,e], David F. Savage[d,e], Enrique M. De

La Cruz[a,2,*], Lauren Ann Metskas[b,c,2,*]

a Department of Molecular Biophysics and Biochemistry, Yale University, New Haven, CT, United States
b Department of Biological Sciences, Purdue University, West Lafayette, IN, United States
c James Tarpo Jr. and Margaret Tarpo Department of Chemistry, Purdue University, West Lafayette, IN,
United States
d Department of Molecular and Cell Biology, University of California Berkeley, Berkeley, CA, United States
e Howard Hughes Medical Institute, University of California; Berkeley, CA, United States
1 W.C. and K.R. contributed equally to this work
2 To whom correspondence may be addressed. Email: enrique.delacruz@yale.edu or
lmetskas@purdue.edu.
*Co-corresponding authors

Enrique M. De La Cruz: PO Box 208114, 266 Whitney Avenue, New Haven, Connecticut, 06520; 203-
506-2090; enrique.delacruz@yale.edu
18
19 Lauren Ann Metskas: 240 Martin Jischke Dr., Purdue University, West Lafayette, Indiana, 47907; 765-
20 494-4909; metskas@purdue.edu
21

Abstract

Bacteria microcompartments (BMCs) are pseudo-organelles comprised of a self-assembling, semi-permeable protein shell, most commonly enclosing components of enzymatic pathways. α-Carboxysomes (α-CBs) are anabolic BMCs known for their role in sequestering Rubisco, the enzyme responsible for carbon fixation in plants, algae and bacteria, along with an upstream enzyme and an assembly protein. Rubisco has low selectivity for its substrate, CO_2, and has a slow enzymatic turnover rate, resulting in an inefficient metabolic pathway. Within the α-CB, Rubisco has been observed at a range of concentrations and with either a liquid-like assembly or a pseudo-lattice of polymerized fibrils. The biophysical origins of the fibril ultrastructure organization are unclear; however, it is only observed inside α-CBs. Quantitative knowledge of the binding constants and energies for assembly and maintenance of these fibrils is critical for understanding this organization and Rubisco regulation, but quantitative methods for *in situ* analysis of Rubisco polymerization have been lacking. Here, we present an approach to convert tomography-derived α-CB volumes and Rubisco particle positions into polymerization binding curves. We used this procedure to determine the Rubisco polymerization constants, including the nucleus size (n) and equilibrium polymerization constant (K_{pol}). The adopted modeling approach is consistent with *in situ* constraints, such as concentration-dependent binding interactions and confinement. This approach offers a powerful tool to evaluate both *in vitro* and potentially *in vivo* biomolecular interactions, both of Rubisco and of other proteins and polymers suitable for analysis by cryo-electron tomography.

Significance Statement

Cryogenic electron tomography (cryoET) is a powerful method to resolve structures of proteins in their

native environment at subnanometer-level resolution. Because tomography data retains spatial

relationships of all particles, it intrinsically contains information about component (e.g., protein) binding

interactions. Here, we use Rubisco polymerization in α-carboxysomes as a model system to demonstrate

that quantitative, biochemical binding analysis is possible with cryoET.

Introduction

Quantitative knowledge of the equilibrium binding affinities of (macro) molecular interactions is

crucial for understanding the regulation of key biological processes including enzyme catalysis, signal

transduction, gene expression, and the assembly of macromolecular complexes (1–4). Established

methods for determining molecular binding constants commonly rely on experimental data acquired *in*

vitro, under controlled experimental conditions (5, 6). Such *in vitro* studies are indispensable for

developing quantitative, mechanistic models, but they often have inherent limitations, such as restricted

concentration ranges, non-native chemical modifications (i.e., tags), and/or minimal buffer systems

substituting for uncharacterized physiological chemical environments. Studies rarely evaluate the effects

of physiological crowding (7, 8), and thus fail to replicate native physiological conditions.

Cryogenic electron tomography (cryoET) is a powerful imaging technique with resolutions ranging

from near-atomic (protein structures) to micron (cellular features) (9–11). While the method typically

provides slightly lower resolution than cryogenic electron microscopy (cryoEM) in structural biology

applications, it has the unique ability to preserve the spatial relationships between imaged particles. In

subtomogram averaging (STA), proteins are identified, extracted from the larger image, aligned and

averaged, and then replaced into the tomogram, enabling extrapolation of these spatial relationships (12,

13). Despite these capabilities, the most detailed analyses of proteins with cryoET typically focus on

structure, with more qualitative or morphological descriptions of protein environment, proximity to other

features, and ultrastructure arrangement.

Here, we present a method using cryoET to quantify binding interactions and polymerization

constants of a self-associating biopolymer *in situ*. We posit that if all particles in a defined and precisely

measured volume are accurately identified and their oligomeric states distinguished, biophysical binding

parameters such as polymerization constants can be calculated from STA data. In addition to providing

simultaneous insight into protein conformation and binding, this approach eliminates the need for complex

in vitro assay design while maximizing the data available in tomograms and extending polymerization

assay capabilities to systems not amenable to *in vitro* measurements.

We use Rubisco in α-carboxysomes (α-CBs), a bacterial microcompartment (BMC) responsible

for carbon fixation in *Halothiobacillus neapolitanus* (14–16), as a model system to validate this approach.

α-CBs are ~1 aL pseudo-organelles with a selectively permeable protein shell (17–20). The protein shell

assembles around an enzyme condensate, facilitated by a disordered scaffolding protein (21). α-CBs

encapsulate Rubisco with its upstream enzyme carbonic anhydrase, maximizing Rubisco turnover rates

presumably through substrate saturation (14, 19, 22–24). α-CB assembly and cargo sequestration rely

largely on Rubisco condensation through liquid-liquid phase separation (25). This process requires a

disordered accessory protein (CsoS2) interfacing with the shell to assemble the facets while also forming

interfaces with Rubisco to facilitate inclusion within the microcompartment (21). The shell is assembled

around the condensate.

Rubisco condensates with CsoS2 inside α-CBs assemble pseudo-lattices of polymer fibrils, a

behavior not replicated with purified components *in vitro* (Fig. 1A) (21, 26, 27). Recent studies have

indicated that BMC shells can separate redox states and that the interior of β-CBs becomes oxidized over

time, with unknown implications on the BMC's interior organization or function (28, 29).

In this study, we developed and validated a framework and procedure to quantitatively extract

polymerization binding constants directly from cryoET data, using Rubisco in α-CBs as a model system.

By converting STA volumes and particle positions into polymerization "binding" curves, we determined key

binding equilibrium constants governing Rubisco fibril assembly *in situ*. This approach not only addresses

a critical gap in our ability to quantify biomolecular assembly under native physiological conditions, offers

a complementary technique to traditional *in vitro* methods and extending the analytical applications of

cryoET beyond structural studies, but also provides a general method in analysis of biomolecule

polymerization with the theories already developed.

Results

Rubisco oligomeric states within α-CBs

Rubisco within α-CBs can adopt two oligomeric/polymer states (Fig. 1B): dense (disorganized or liquid-like) and ordered (lattice-like packing). Purified α-CBs can also have low Rubisco concentration (termed "sparse") as an experimental consequence of centrifugation (26). It is not known what determines whether Rubisco within α-CBs adopts the dense or ordered morphology at equivalent concentrations. We hypothesize that Rubisco polymerization could be affected by redox state. To test this, α-CBs were isolated from *H. neapolitanus* and treated with either DTT (reducing agent) or diamide (oxidizing agent) to alter their internal redox state before vitrification and cryoET analysis of polymerization.

Rubisco monomers, oligomers, and linear polymers were identified within intact α-CBs and STA performed out to sub-7Å resolution using published methods (Fig. 1C) (26, 30). Helices are well defined at this resolution, allowing precise estimation of position and orientation for each Rubisco complex (Fig. 1D). We used a numerical classification approach to calculate α-CB volume and determine the binding state of each Rubisco complex, using the positions and orientations generated by STA analysis (31). These individual Rubisco are classified as "bound" (oligomer or polymer; nuclei are treated as polymers; this is necessary for mass balance in the theory) or "unbound" (monomer for nucleus size $n = 2$, or monomer plus dimer for nucleus size $n = 3$; discussed below). The bound species can be categorized by the number (n) of contiguous subunits comprising the polymer. A Rubisco identified as part of a polymer chain can be further processed with STA to isolate the Rubisco-Rubisco subunit interface (Fig. 1E-G). By using a numerical analysis rather than image classification, the analyst controls the parameters and corresponding error rates and can cleanly establish the polymer length and connectivity of each Rubisco; it is also computationally more efficient.

Rubisco packing within α-CBs

The average volume of an untreated α-CB is $7 \pm 2 \times 10^{-4}$ µm³ corresponding to approximately a cube with 90 nm edge length or a sphere with 60 nm radius (Fig. 2A, Table 1). DTT treatment changed the average volume (Table 1), possibly through destabilization of the shell leading to selective loss of larger α-CBs (Fig. 2A, Table 1). Both chemical treatments resulted in populations with slightly higher

Rubisco concentrations (45% for DTT treatment, 26% for diamide treatment, Table 1), presumably due to

loss of less mechanically stable α-CBs during plunge-freezing. While the chemical treatments reduced the

number of low-concentration data points available for polymerization analysis, enough remained to

facilitate curve fits.

The concentration of inner Rubiscos (defined as those not adjacent to the α-CB shell) scaled

linearly with the total Rubisco concentration in untreated, DTT-treated, and diamide-treated samples (Fig.

S2A-C). On average, inner Rubisco concentrations are about 1.4-fold higher than their total Rubisco

concentrations for all three treated and untreated samples. The best fit lines to the inner Rubisco

concentrations to total Rubisco concentrations for all three α-CB samples have a positive *x*-axis intercept

(Fig. S2A-C, Table 1). This behavior indicates that inner Rubiscos are absent and all Rubisco is contained

within the shell region at low total Rubisco concentrations, consistent with tomograms of lower-

concentration α-CBs. This suggests that Rubisco near the shell may have different properties, interactions

or other behaviors compared to the α-CB interior.

Rubisco polymer chain length distribution within α-*carboxysomes.*

Rubisco forms polymer chains in all α-CB populations imaged (Fig. 1B). Most of the polymers

observed in the data are dimers, and the concentrations of longer chains decay with the chain length (Fig.

3). The Rubisco in dimers observed are not randomly contacting each other; rather, the interfaces

observed between adjacent, continuous subunits in the Rubisco dimers are the same as those found in

the longer polymers (Experimental section).

The length distribution of an equilibrium polymer follows an exponential decay (32) (see Eq. 6 in

Theory section). Histograms of Rubisco chain lengths inside all untreated, DTT treated and diamide

treated α-CBs were fitted to an exponential (Fig. 3). The best fit to all the polymer lengths including

dimers ($j \geq 2$ Rubisco complexes; black smooth line) does not fit the data well, especially at longer

polymer lengths. The observed dimer population deviates from the exponential decay defining the

polymer population distribution, indicating that dimers are thermodynamically distinct from polymers of

three or more Rubisco complexes. In contrast, the best fit to the polymer length distribution for trimers and

longer polymers ($j \geq 3$ Rubisco complexes; gray smooth line) fits the data well for all samples (Fig. 3).

This exponential length distribution behavior indicates (Eq. 6) that the internal Rubisco interactions in

polymers of three or more Rubisco complexes ($j \geq 3$) are thermodynamically equivalent. We interpret this

as an indication that the nucleus for Rubisco polymerization is three contiguous Rubisco complexes ($n =$

3).

A previous thermodynamic theory analysis (33) has shown that the chain length exponential

decay constant (κ) is directly proportional to the total free energy of the polymer ($G_j^{o\prime}$ for a polymer

composed of j Rubisco complexes) and the free energy change of the polymer ($\Delta G_{j \to j+1}^{o\prime}$) resulting from

elongation by one subunit, in turn, can be linked to the polymerization constant (K_{pol}; also referred to as

the "critical concentration" for polymerization (34)) and the free monomer concentration in equilibrium with

polymer (A_1; Eq. 6 in the Theory section) according to:

$$\frac{\Delta G_{j \to j+1}^{o\prime}}{k_B T} = \kappa = -\ln\left(\frac{A_1}{K_{pol}}\right)$$

The free energy change ($\Delta G_{j \to j+1}^{o\prime}$) of the polymer when elongated by one subunit is given by:

$$\Delta G_{j \to j+1}^{o\prime} = \kappa k_B T = k_B T\big(\ln(K_{pol}) - \ln(A_1)\big)$$ Eq. 10

The free energy change per subunit extension ($\Delta G_{j \to j+1}^{o\prime}$) is constant for given K_{pol} and A_{tot}, so the total

value of $\Delta G_{j \to j+i}^{o\prime} = i\kappa k_B T$ scales with the total number (i) of subunits incorporated into the polymer,

including those resulting from the annealing of existing polymers.

The best fit to the Rubisco polymer ($j \geq 3$) length distribution to an exponential (Fig. 3 and Table 2)

yields $\Delta G_{j \to j+1}^{o\prime}$ = +0.42 ± 0.03, +0.46 ± 0.08, and +0.49 ± 0.02 $k_B T$ for untreated, DTT treated and diamide

treated Rubisco complexes, indicating the Rubisco polymer chain internal stability is < $k_B T$ and

comparable for all three samples. We emphasize that this free energy is distinct from the standard Gibbs

free energy change ($\Delta G^{0\prime}_{pol}$) associated with subunit addition to the filament end (discussed below).

Rubisco polymerization mechanism within α-CBs

Because we determined the nucleus for Rubisco polymerization is a trimer ($n = 3$; Fig. 3), we

treat Rubisco trimers and longer polymers as "bound" when plotting the concentration-dependence of

polymer concentration (blue filled circles in Figs. 4 and S4 right panels). In this case, Rubisco dimers

belong to the unbound, 'monomer pool' (light blue filled circles in Figs. 4 and S4 right panels). To extract

the polymerization binding constants from this data, we can either fit the A_{tot} -dependence of bound

Rubisco alone or globally fit both unbound and bound Rubisco concentrations together. Both fitting

methods yield similar results, so we present only the global fits.

The best global fit of the data to a length unlimited nucleation model (Eq. 3 and 4) with a trimer

nucleus (n = 3) yields a K_{pol} = 541 $\pm$ 44 μM for untreated Rubisco, 762 $\pm$ 5 μM for DTT-treated Rubisco

and 730 $\pm$ 170 μM for diamide-treated Rubisco polymers. These polymerization constants are weak (K_{pol}

>500 μM), and associated with small standard Gibbs free energy changes for incorporation of a Rubisco

subunit at a polymer end ($\Delta G^{0'}_{pol}$) according to:

$$\Delta G^{0'}_{pol} = k_B T \ln K_{pol} > k_B T \ln(5 \times 10^{-4} \text{ M}) = -7.6 k_B T \qquad\qquad \text{Eq. 11}$$

The nucleation binding constants (K_{nuc}) are even weaker in all three cases, with values ranging from 10^9

to 10^{21} μM (Fig. 4, Table 3).

For comparison, we analyzed the data with five other polymerization models: length-limited and

length-unlimited isodesmic polymerization, length-limited and length-unlimited nucleated polymerization

with a dimer nucleus (n = 2; Fig. S4A-C, left), and length-limited nucleated polymerization with a trimer

nucleus (n = 3; Fig. S4D-F, right), together with length-unlimited n = 3 nucleated polymerization (Figs. 4

and S4D-F). In the case of isodesmic polymerization and dimer nucleation polymerization, unbound

Rubisco complexes are treated as "monomers" (light gray filled circles), and bound Rubisco complexes

are dimers and longer polymers (gray filled circles in Fig. S4A-C, left).

Isodesmic polymerization models do not fit the experimental data well for either length-limited or

length-unlimited models (Fig. S4A-C, left). Dimer and trimer nucleation models with unlimited

polymerization fit the corresponding data equally well (Fig. S4) and considerably better than with a length-

limited constraint. The results of these analyses are summarized in Table 3. Fitting such polymerization

data cannot determine the size of the nucleus (n) because the nucleus size was constrained when

classifying oligomers as bound or unbound, thus, relating fit quality to nucleus size would be circular

reasoning.

The best fits of the data to length-unlimited nucleated (n = 2 or 3) polymerization models yield

very weak nucleation constant (K_{nuc}) values with large uncertainties for all Rubisco samples (Table 3). The

K_{nuc} values also vary by many orders of magnitude among the three samples. Despite the large

uncertainties, the K_{nuc} values reflect and capture the sharpness of the monomer-polymer transition (i.e.,

inflection point) at the macromolecule critical concentration (Fig. S5). Abrupt transitions at the critical

concentration indicate that nucleation (K_{nuc}) is much weaker than subunit incorporation at a filament end

(K_{pol}). Self-assembling polymer systems with K_{pol} << K_{nuc} and abrupt transitions (i.e., actin filaments; (35))

are said to be "highly cooperative". Thus, Rubisco polymerization is highly cooperative, despite the

observed weak critical concentration. We note that despite the weak polymerization and large

uncertainties in the collected data, the best fits can readily distinguish changes in the critical

concentration for assembly under different conditions (Figs. 4 and S4).

Discussion

A novel method for evaluating protein-protein binding in STA

We demonstrate a novel approach to determine protein binding using volumetric imaging with

STA, a technique typically used for protein structure determination (36). This approach leverages the

intrinsic capability of STA to provide position and orientation information about proteins and other

biological macromolecules, using these data to determine the relative concentrations of bound (i.e.,

polymerized, in a complex, etc.) and unbound particles in their biological context. This bottom-up method

can be implemented as long as a protein can be clearly identified in both bound and unbound states for

accurate classification. It is agnostic to concentration, allows experiments under native conditions, and

does not require tagging or chemical modification of the target molecules. Concentrations can be

calculated either from an enclosing barrier or directly from the particles' occupied volume, either counting

the particles alone or including molecular crowding as nonspecific electron density within the particle-

occupied area (31).

Single-particle cryoEM has previously been used to estimate binding constants of purified

proteins by resolving the structures of bound and unbound states. In structure-based drug design,

cryoEM structures are used to select computational binding poses that correlate with compound potency,

with binding energies validated experimentally by surface plasmon resonance or estimated

computationally through energy minimization of predicted conformational changes upon substrate binding

(37, 38). While this approach has shown promise for drug discovery, physiological relevance remains a

concern: during plunge-freezing preparations, proteins are exposed to fluid shearing force, uncontrolled

osmolarity increases, and surface interactions. Furthermore, the high-resolution structures required for

atomic-level comparison preclude inclusion of the entire dataset, as less populated or heterogeneous

states would not resolve. CryoET mitigates some of these concerns by imaging proteins within an

enclosed system, such as a cell, microcompartment, or artificial lipid vesicle enclosing purified protein.

These environments shield the interior from nonphysiological conditions during sample preparation, while

supporting more complex three-dimensional analyses of protein organization and interactions.

Our model system, Rubisco in α-CBs, offers several advantages for the development of the

method used here to identify protein assembly and determine binding and polymerization constants.

Rubisco complexes are large and easily identifiable on a grid, and the presence of a visible shell with a

defined volume provides secondary validation of occupied volume calculations for determining Rubisco

concentration (31). The protein shell sequesters the interior chemically and mechanically over the sample

preparation timescale, preserving the native conditions. Rubisco's large size allows unambiguous particle

identification in bound and unbound forms, and its marked conformational stability allows STA of the full

population without the need for classification (30). Finally, the α-CB is relatively well characterized

biochemically and relative abundances of the contained proteins known, providing validation for

underlying calculations such as Rubisco concentration in the interior (39). Application of this new binding

analysis methodology to other systems is certainly possible, with noise and error rates ultimately

dependent upon the particle identification accuracy and final resolution of the STA dataset.

Rubisco polymerization in α-carboxysomes

All the three Rubisco polymerization datasets yield an weak (large in value) polymerization

constant K_{pol}, (i.e., overall dissociation equilibrium constant for a monomer to bind either end of a polymer

chain (40)), and an even weaker nucleation constant K_{nuc} (Table 3), consistent with the observed sharp

transitions near the inflection point for all three samples (Fig. 4, S4 and S5). This supports a mechanism

in which space confinement is a primary factor for Rubisco polymerization.

We tested four general models of macromolecule polymerization: isodesmic and nucleation

models, both with and without spatial constraints. In unlimited space, the length of a self-assembling

polymer chain is unconstrained and, in theory, can reach infinite lengths as the total subunit concentration

approaches infinity. Under confinement, the polymer chain length is limited by the accessible space,

geometry of the polymer chain, and the chain's material properties.

The average diameter of an untreated α-CB is equivalent to the linear length of 10 or 11 Rubiscos

complexes (see *Polymerization theory within a confined volume* in Theory of Protein Polymerization),

suggesting the volume may limit the maximum allowed Rubisco chain length. The polymerization models

must be modified to reflect this spatial constraint (Eq. 7). The compliance of the polymer can also

contribute to the achievable length. For example, if Rubisco polymers are very compliant, such that they

could bend and twist easily, the maximum allowed chain length could conceivably be longer than a rigid,

linear polymer. In our data, the maximum Rubisco chain length observed is equivalent to the α-CB

diameter (10 Rubisco complexes for untreated and DTT-treated and 11 for diamide-treated (Table 2)).

This suggests that Rubisco polymers are rigid on the observed length scales and the chain length is

limited by the accessible space within α-CBs.

All polymerization models tested here predict that the polymer chain length distribution at

equilibrium follows an exponential decay (Eq. 6). The observed Rubisco chain length distribution shows

that the dimer fractions deviate from the exponential distribution of the population, whereas trimers and

longer polymers adhere to this exponential population distribution (Fig. 3). This suggests Rubisco

polymerization follows a nucleation mechanism with nucleus size (*n*) of 3 Rubisco complexes.

Isodesmic polymerization models, in which nuclei are not required for polymer formation, do not

fit data well (Fig. S4A-C). Nucleation models with either *n* = 2 or *n* = 3 with confinement yield reasonable

fits to each corresponding data set, although the fits are worse than without confinement (Figs. 4 and S4)

as indicated from the sharpness of the transition from unbound to bound species (i.e., the critical

concentration inflection point).

The polymerization data were better fitted by the chain-length unlimited nucleation model (Figs. 4

and S4), suggesting that spatial confinement has limited effects on Rubisco polymerization. If the length

limit was achieved before subunit packing, the observed filament length distribution would shift toward a

uniform population with the spatially-limited length. Therefore, the Rubisco filament length distribution and

polymerization analyses together favor a length-unlimited nucleation mechanism with a trimer (n =3)

nucleus for Rubisco polymerization in α-CBs.

299   Previous α-CB assembly studies have returned conflicting results a tomography study observed

α-CB-sized Rubisco condensates in the cytoplasm that the authors interpreted as assembly intermediates

(41), suggesting that the Rubisco condensate would determine the α-CB size, while biochemical

experiments indicate that the accessory protein CsoS2 is the primary determinant of α-CB shell diameter

(42–44). We conclude that, while we cannot determine the details of α-CB assembly from these

polymerization fits, we can successfully determine that the maximal Rubisco chain length is an intrinsic

property of the condensate rather than imposed by the shell.

Disproportional abundance of Rubisco dimers

308   Rubisco dimers are prevalent in condensates (Fig. 1A) as well as in α-CBs *in vitro* and *in situ*.

309   Conversely, Rubisco trimers and longer polymers are found in low concentrations in solution, and are

310   primarily in α-CBs(21).

311   We observed a large fraction of Rubisco dimers in α-CBs. This abundance of dimers is

312   unexpected for a nucleus size (n) of 2 or 3, as oligomers shorter than the nucleus (n = 3) are typically

313   unstable and populated at low concentrations, and oligomers equal to the nucleus should behave as

314   polymers in the exponential length distribution (Fig. 3). The deviation of the dimer population from the

315   exponential, polymer length distribution (Fig. 3), suggests that they are thermodynamically distinct from

316   polymerized Rubisco fibrils.

317   The total dimer species is overpopulated, implying that it is more stable than any polymer

species, and raising the possibility that a fraction of this dimer population is polymerization-incompetent.

At least four cases can potentially account for this unexpected finding:

Case 1: *Ambiguity identifying and quantifying the Rubisco dimer population*: Two Rubisco monomers

positioned in an adjacent manner could be incorrectly classified as a dimer, yielding an "overcounting" of

the dimer population. (This would yield n=3 nucleus size, but with some dimers being monomers; see

Figs. 4 and S4D-F.) However, the cryoET method employed here retains sufficient spatial and structural

resolution to distinguish and confirm that the observed dimers are not randomly contacting each other but

adopt the same orientation as adjacent, contiguous subunits in longer polymers. Furthermore, this

packing effect should lead to similar "overcounting" of all observed oligomer and polymer species

detected. We thus can eliminate miscounting as the source of the observed abundant dimer population.

Case 2. <u>*A population of Rubisco dimers are polymerization-incompetent and on the polymerization*</u>

<u>*pathway*</u>.

A mechanism in which two distinct Rubisco dimer conformers – one capable of polymerizing (R_2) and

one not (R_2^*) – are linked by a reversible and on-pathway isomerization equilibrium (K_{isom}) can account for

the observed overabundance of Rubisco dimers (Scheme 3a).

$$K_{nuc} = K_{nuc1}\ K_{isom}\ K_{nuc2}$$

333 $$R\ +\ R \underset{\xrightarrow{\hspace{0.6cm}}}{\xleftarrow{K_{nuc1}}}\ R_2^* \underset{\xrightarrow{\hspace{0.6cm}}}{\xleftarrow{K_{isom}}}\ R_2 + R \underset{\xrightarrow{\hspace{0.6cm}}}{\xleftarrow{K_{nuc2}}}\ R_3 + R \underset{\xrightarrow{\hspace{0.6cm}}}{\xleftarrow{K_{pol}}}\ R_4 + R \underset{\xrightarrow{\hspace{0.6cm}}}{\xleftarrow{K_{pol}}}\ ......$$
                                            Scheme 3a

Given the resolution and orientation criteria for dimer classification (discussed in *Case 1*), the structural

difference between the two dimer conformers would need to have a negligible effect on the Rubisco

backbone, and likely be located at the subunit interface.

This polymerization pathway (Scheme 3a) defines a Rubisco trimer as the nucleus for polymerization

($n = 3$). The nucleation constant (K_{nuc}) is defined here in a step-wise manner with binding constants K_{nuc1},

K_{isom}, and K_{nuc2}, such that the overall nucleation equilibrium constant is given by the product of these

three step-wise binding constants ($K_{nuc} = K_{nuc1}K_{isom}K_{nuc2}$). This allows both the polymerization-incompetent

dimers (R_2^*) and the nucleus (R_3) to be expressed in terms of the polymerization-competent dimer (R_2)

concentration at equilibrium and the corresponding dissociation equilibrium constants (K_{isom} and K_{nuc2})

according to $[R_2^*] = [R_2]K_{isom}$ and $[R_3] = [R_2][R]/K_{nuc2}$.

Accordingly, the ratio of the total dimer concentration (i.e., $[R_{2,tot}] = [R_2^*] + [R_2]$) over the nucleus

trimer (R_3) concentration is given by (Eq. 12):

$$\frac{[R_2^*] + [R_2]}{[R_3]} = \frac{(1 + K_{isom})K_{nuc2}}{[R]}$$
                Eq. 12

This relationship indicates the observed dimer population ($R_{2,tot}$) will exceed that of the nucleus trimer (R_3)

when $(1 + K_{isom})K_{nuc2} > [R] \sim 500\ \mu M$ (i.e., the Rubisco monomer concentration in equilibrium with

polymer). Therefore, a weak K_{isom} (i.e., equilibrium largely favors polymerization-incompetent R_2*) and/or

weak K_{nuc2} (i.e., equilibrium favors R_2 over R_3) would generate an overpopulated dimer fraction.

In Scheme 3a, subunit addition to a dimer end (K_{nuc2}) could be weaker or tighter than addition to a

trimer end (K_{pol}). If the ends of a Rubisco dimer and trimer are identical, as may be expected for a 1-D

polymer, K_{nuc2} would be defined as K_{pol}, K_{nuc} would be defined as $K_{nuc1}K_{isom}$, and the nucleus size would

be a dimer (R_2, $n = 2$), as detailed by (Scheme 3b):

$$K_{nuc} = K_{nuc1}\ K_{isom}$$

$$R + R \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{nuc1}}} R_2^* \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{isom}}} R_2 + R \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{pol}}} R_3 + R \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{pol}}} R_4 + R \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{pol}}} \ ......$$

355   Scheme 3b

In this case, the nucleus R_2 population adheres to the exponential, polymer-length distribution (Fig.

3), and the polymerization incompetent dimer (R_2^*) contributes to the over-counted dimer fraction. As with

Scheme 3a, Eq. 12 with K_{nuc2} defined as K_{pol} predicts that the total observed dimer ($R_{2,tot}$) will be

overpopulated when K_{isom} is weak and largely favors R_2^* such that $(1 + K_{isom}) > 1$.

Case 3. <u>A population of Rubisco dimers is polymerization-incompetent and off-the polymerization</u>

<u>pathway</u>.

A mechanism with two distinct Rubisco conformers (R_2 and R_2*), in which the incompetent dimer (R_2^*)

is populated through a branched, off-pathway equilibrium can also potentially account for the observed

population of Rubisco dimers. The non-productive, polymerization-incompetent dimers can form either

through association of two Rubisco monomers (K_{dimer}; Scheme 4a) or isomerization (K_{isom}, Scheme 4b) of

a polymerization-competent Rubisco dimer (R_2) according to the following reaction pathways:

$$K_{nuc} = K_{nuc1}\ K_{nuc2}$$

$$R + R \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{nuc1}}} R_2 + R \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{nuc2}}} R_3 + R \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{pol}}} R_4 + R \underset{\xleftarrow{\hspace{1em}}}{\xrightarrow{K_{pol}}} \ ......$$

$$K_{dimer} \updownarrow$$

$$R_2^*$$

368   Scheme 4a

$$K_{nuc} = K_{nuc1}\,K_{nuc2}$$

$$R + R \;\underset{}{\overset{K_{nuc1}}{\rightleftharpoons}}\; R_2 + R \;\underset{}{\overset{K_{nuc2}}{\rightleftharpoons}}\; R_3 + R \;\underset{}{\overset{K_{pol}}{\rightleftharpoons}}\; R_4 + R \;\underset{}{\overset{K_{pol}}{\rightleftharpoons}}\; ......$$

$$K_{isom} \updownarrow$$

$$R_2^{*}$$

369   Scheme 4b

370   In Scheme 4a and 4b, if the nucleus size $n = 3$, the overall nucleation constant (K_{nuc}) can be defined

371   as $K_{nuc} = [R]^3/[R_3]$ or in a stepwise manner as $K_{nuc} = K_{nuc1}K_{nuc2}$, with $K_{nuc1} = [R]^2/[R_2]$. The non-productive

372   dimer concentration ($[R_2^{*}]$) is given by $[R_2^{*}] = [R]^2/K_{dimer}$ in Scheme 4a, and the following relationship holds

373   (Eq. 13a):

374
$$\frac{[R_2^{*}] + [R_2]}{[R_3]} = \frac{\left(\frac{1}{K_{nuc1}} + \frac{1}{K_{dimer}}\right)[R]^2}{\frac{[R]^3}{K_{nuc}}} = \frac{(K_{nuc1} + K_{dimer})K_{nuc2}}{[R]K_{dimer}} = \frac{\left(\frac{K_{nuc1}}{K_{dimer}} + 1\right)K_{nuc2}}{[R]} \qquad \text{Eq. 13a}$$

375   In Scheme 4b, $[R_2^{*}] = [R_2]/K_{isom} = [R]^2/(K_{nuc1}K_{isom})$, and the following relationship holds (Eq. 13b):

376
$$\frac{[R_2^{*}] + [R_2]}{[R_3]} = \frac{\left(\frac{1}{K_{nuc1}} + \frac{1}{K_{nuc1}K_{isom}}\right)[R]^2}{\frac{[R]^3}{K_{nuc}}} = \frac{(1 + K_{dimer})K_{nuc2}}{[R]K_{dimer}} = \frac{\left(\frac{1}{K_{dimer}} + 1\right)K_{nuc2}}{[R]} \qquad \text{Eq. 13b}$$

In both non-productive, off-pathway Rubisco dimer pathways, the observed dimer population will exceed

the nucleus population (R_3) in the polymer length distribution (Fig. 3) when K_{isom} or K_{dimer} favors R_2^{*} and/or

K_{nuc2} is weak and favors R_2 over R_3.

As explained above in the case of an on-pathway, non-productive dimer (Scheme 3b), if the ends of a

Rubisco dimer and trimer are identical, K_{nuc2} is defined as K_{pol}, K_{nuc} is defined as K_{nuc1}, and the nucleus

size would be a dimer (R_2, $n = 2$), according to:

$$R + R \;\underset{}{\overset{K_{nuc}}{\rightleftharpoons}}\; R_2 + R \;\underset{}{\overset{K_{pol}}{\rightleftharpoons}}\; R_3 + R \;\underset{}{\overset{K_{pol}}{\rightleftharpoons}}\; R_4 + R \;\underset{}{\overset{K_{pol}}{\rightleftharpoons}}\; ......$$

$$K_{dimer} \updownarrow$$

$$R_2^{*}$$

383   Scheme 4c

$$R + R \; \underset{\longleftarrow}{\overset{K_{nuc}}{\longrightarrow}} \; R_2 + R \; \underset{\longleftarrow}{\overset{K_{pol}}{\longrightarrow}} \; R_3 + R \; \underset{\longleftarrow}{\overset{K_{pol}}{\longrightarrow}} \; R_4 + R \; \underset{\longleftarrow}{\overset{K_{pol}}{\longrightarrow}} \; ......$$

$$\Big\updownarrow K_{isom}$$

$$R_2^{\;*}$$

384   Scheme 4d

In these two cases, the total observed dimer population will be over counted when K_{nuc} is weaker than

K_{dimer}, thereby favoring R_2^*, or K_{isom} favors R_2^*, as above. As in *Case 2*, such an effect could not alter

secondary or tertiary structure, or it would be visible in the structural biology analysis.

Case 4: <u>*A regulatory factor in α-carboxysomes binds and sequesters Rubisco dimers*</u>.

An off-pathway, polymerization-incompetent dimer (R_2^*; *Case 3*) could potentially originate from the

binding of a regulatory ligand (*L*), resulting in dimer sequestration. In this case, Schemes 4b and 4d

become Scheme 5a and 5b, respectively, and all other relationships hold.

$$K_{nuc} = K_{nuc1}\,K_{nuc2}$$

$$R + R \; \underset{\longleftarrow}{\overset{K_{nuc1}}{\longrightarrow}} \; R_2 + R \; \underset{\longleftarrow}{\overset{K_{nuc2}}{\longrightarrow}} \; R_3 + R \; \underset{\longleftarrow}{\overset{K_{pol}}{\longrightarrow}} \; R_4 + R \; \underset{\longleftarrow}{\overset{K_{pol}}{\longrightarrow}} \; ......$$

$$\Big\updownarrow K_{ligand}$$

$$R_2 L$$

393   Scheme 5a

$$R + R \; \underset{\longleftarrow}{\overset{K_{nuc}}{\longrightarrow}} \; R_2 + R \; \underset{\longleftarrow}{\overset{K_{pol}}{\longrightarrow}} \; R_3 + R \; \underset{\longleftarrow}{\overset{K_{pol}}{\longrightarrow}} \; R_4 + R \; \underset{\longleftarrow}{\overset{K_{pol}}{\longrightarrow}} \; ......$$

$$\Big\updownarrow K_{ligand}$$

$$R_2 L$$

394   Scheme 5b

395   It is possible that a small-appearing ligand could be invisible in the STA structures if its occupancy

396   was only once per complex, as the symmetry constraint in processing would diffuse its signal. While the

397   STA was confirmed with no symmetry imposed (C1), a small-appearing ligand would have too small a

398   signal to drive alignment.

 Free energy changes associated with Rubisco polymerization

The total free energy of a polymer chain ($G_j^{o'}$) scales with the total number of subunits (Eq. 10)

and thus increases when it elongates. The free energy change of a Rubisco fibril associated with

incorporation of one subunit ($\Delta G_{j \to j+1}^{o'} = \kappa k_B T$) is < 0.5 $k_B T$ (Eq. 10) for all Rubisco samples evaluated here.

A half $k_B T$ of energy change seems small since it is equal to the amount of the thermal energy associated

with one degree of freedom, but it is >1200-fold larger than that associated with actin filament elongation

($\Delta G_{j \to j+1}^{o'}$ = 0.0004 $k_B T$; (32)). A polymer with larger $\Delta G_{j \to j+1}^{o'}$ is thermodynamically and mechanically less

stable and more susceptible to spontaneous fragmentation due to thermal and applied forces (45–47).

Therefore, Rubisco polymer chains are considerably thermodynamically and mechanically less stable

than actin filaments.

We clarify here for general readers that the two free energy changes for subunit incorporation at

polymer ends discussed here, $\Delta G^{0'}_{pol}$ vs. $\Delta G_{j \to j+1}^{o'}$, are not equivalent. The term $\Delta G_{j \to j+1}^{o'}$ is the amount of

free energy added to a polymer when it extends one subunit, while $\Delta G^{0'}_{pol}$ is the free energy change for

the overall reaction of a monomer binding to polymer end. Both are material properties, but $\Delta G_{j \to j+1}^{o'}$ is an

intrinsic constant, inversely scaling with polymers' thermal and mechanical stabilities, while $\Delta G^{0'}_{pol}$ is an

intrinsic constant reflecting the feasibility for monomer incorporation at a polymer end.

The (standard) Gibbs free energy change $\Delta G^{0'}_{pol}$ is the negative change (loss) given by the

difference between the Gibbs free energy ($G^{0'}$) sum of a polymer with j subunits plus a free monomer

$\left(G^{0'}_j + G^{0'}_1\right)$ and a polymer with j + 1 subunits $\left(G^{0'}_{j+1}\right)$:

$$\Delta G^{0'}_{pol} = G^{0'}_{j+1} - \left(G^{0'}_j + G^{0'}_1\right) = k_B T \ln K_{pol} \qquad \text{Eq. 14}$$

It reflects the free energy change associated with subunit incorporation at a filament end and is

independent of the length of the elongating filament it is adding to. The polymerized subunit free energy is

lower than that of a free monomer in solution, so subunit elongation is associated with a negative Gibbs

free energy change.

In contrast, $\Delta G_{j \to j+1}^{o'} = G^{0'}_{j+1} - G^{0'}_j$ is defined as the (standard) free energy increase of the

polymer itself when it elongates by one subunit ($\Delta G_{j \to j+1}^{o'} = \kappa k_B$, Eq 10). A longer polymer (of a given

subunit type, such that the free energy change per subunit is constant) has a larger total free energy than

a shorter one and is less therefore stable.

The two free energy changes refer to different reacting components are related according to the

following, obtained by combining Eqs. 10 and 14:

$$\Delta G^{o\prime}_{j\to j+1} = k_B T\big(\ln(K_{pol}) - \ln(A_1)\big) = \Delta G^{0\prime}_{pol} - k_B T \ln(A_1) \qquad \text{Eq. 15}$$

431   or

$$\Delta G^{o\prime}_{j\to j+1} - \Delta G^{0\prime}_{pol} = -k_B T \ln(A_1) \qquad \text{Eq. 15}\prime.$$

This relationship means the difference in the polymer free energy ($\Delta G^{o\prime}_{j\to j+1}$) and the free energy

($\Delta G^{0\prime}_{pol}$) for subunit incorporation is determined by the monomer concentration in equilibrium with

polymer. The relationship between $\Delta G^{0\prime}_{pol}$ and $\Delta G^{o\prime}_{j\to j+1}$ (Eq. 15 and 15') can be independently obtained

directly from the Gibbs free energy calculation (described in the Supplement).

Relation of Rubisco chemical state and polymerization

One known chemical difference between cytoplasm and α-CBs is redox state: the cytoplasm is a

reducing environment, while the interior of α-CBs is oxidized (28, 29). We therefore tested whether redox

state could affect Rubisco polymerization. Using chemical treatments, we observed that the ordered

lattices were often larger and more prevalent in oxidized samples compared to reduced samples, but with

only subtle differences in K_{pol} (Figs. 4 and S4; Table 3). This finding emphasizes the importance of

quantifying binding rather than using morphology as a guide: the decreased volume of the DTT-treated α-

CBs likely affected this observation. α-CBs are purified without chemical reducing agents, and thus we

anticipate that the untreated sample should be mildly oxidized as has been previously seen (48). One

potential explanation is that the DTT treatment was not sufficient to fully shift the Rubisco polymerization,

either due to the inability to revert side-chain oxygen additions or due to the relatively short treatment time

tolerated by the α-CB shell.

If the Rubisco complexes exist in two populations within the α-CB (for example, reduced and

oxidized), and if both populations polymerize and have different K_{pol} values within the observed A_t range,

we would predict two inflection points in the data. Conversely, the sole inflection point observed here

indicates that either only one polymerizing species is present, or that two polymerizing species exist but

only one has a critical concentration within the observed range of A_t. To test this, we introduced a floating

parameter in the fitting program as the fraction of total Rubisco A_t that is either unable to polymerize at all

or polymerizes beyond the observed A_t range. Fitting to all the three datasets resulted in this parameter

being indistinguishable from zero, indicating that Rubisco polymerizes as a single species and most likely

that redox state does not affect intrinsic polymerization capability in these experiments. We note that a

case in which two species of Rubiscos exist but copolymerize is also consistent with the data.

For a system in which there is only one uniform polymerization species, α-CBs with the dense

morphology (Figure 1B) should have Rubisco concentrations slightly below the K_{pol}. However, a previous

study of Rubisco lattice formation found that while the ordered morphology consistently had Rubisco

concentrations above what we now know to be the K_{pol}, the dense morphology covered a range of

concentrations and could occur at Rubisco concentrations above this threshold(26).

Taken together, these findings suggest that changes to α-CB packing under different redox

conditions are likely occurring through the action of accessary proteins, such as CsoS2, rather than

through modification of the Rubisco itself (*Case 4* above).

Conclusion

In summary, we have developed a novel tomography-based method for measuring polymerization

in situ. In a model test of measuring Rubisco inside α-CBs, we were able to measure Rubisco K_{pol} and

K_{nuc} constants inside an incompletely characterized physiological system with multiple protein species

present, and at low affinities and high concentrations not typically amenable to in vitro study. We were

able to determine that polymerization proceeds through a nucleus, and that redox-based effects on

polymerization act through a secondary protein partner. This test case demonstrates the power of

combining robust biophysical calculation and modeling with the *in situ* capabilities of cryo-electron

tomography.

Materials and Methods (abridged)

α-CBs were purified from wild-type *H. neapolitanus* as previous described (56).

Negative stain imaging of condensate

Rubisco (0.5 μM) of L8S8 and the CsoS2 (2.7 μM) were mixed at room temperature for 1 hour. The sample was diluted 5x and Rubisco and CsoS2 condensates were visualized by TEM using a 2% (w/v) uranyl acetate stain. Imaging was performed on a JEOL 1200 EX transmission electron microscope.

Sample Conditions and Tomography

The α-CB sample was diluted with BSA-coated 10 nm gold fiducials in a 4:1 ratio. Samples were treated with 5 mM diamide or DTT for one hour (57), then plunge-frozen onto glow-discharged C-flat 2/2-300 copper TEM grids. Imaging was performed at the Pacific Northwest Center for Cryo-Electron Microscopy on a Titan Krios G4 with a Gatan K3 camera and BioQuantum energy filter. Tomograms were collected in SerialEM according to Supplementary Table 2. Frame alignment, 2D image processing, and tilt alignment were performed in etomo; CTF estimation in ctffind4; and tomogram reconstruction in novaCTF following previously published pipelines (58–61).

Subtomogram averaging

Particles were identified in SIRT-like filtered tomograms using a customized workflow designed to identify every Rubisco complex within each α-CB (26, 30). The general subtomogram averaging workflow was based upon a previous high-resolution approach originally derived from the Briggs group (9, 26). Subtomogram alignment and averaging was performed in Dynamo v.1.1.532, using the minimum number of iterations possible at each bin size and employing adaptive bandpass filtering throughout (62). Half-maps were sent to RELION5 for post-processing and final resolution calculations (63). Rigid body docking was performed in UCSF Chimera (64).

The Rubicso-Rubisco interface was generated by isolating all particles in trimers and averaging at the Rubisco-Rubisco interface. The model (Fig. 1E,F) showing the stacked Rubisco-Rubisco dimer was

generated by fitting the monomer maps into the masked interface volume. The interface (Fig. 1G) is the

STA volume generated by Dynamo.

Data Analysis

We used a previously published Matlab script package to determine Rubisco binding states (31).

Rubisco Filament Length Distribution Analysis

At equilibrium, the polymer length distribution is an exponential function of A_1 (Eq. 6) and the

unpolymerized monomer concentration A_1 is function of A_{tot} (Eq. 1, 3, and 7) and increases with A_{tot}, but

$A_1 \rightarrow K_{pol}$ as $A_{tot} \rightarrow \infty$. For a nucleation polymerization, polymers appear only when $A_{tot} > K_{pol}$, and in this

region, $A_1 \sim K_{pol}$ and is almost constant and changes little with A_{tot} (Fig. 4). Consequently, the polymer

length distribution (Eq. 6) changes very little with A_{tot}, and aggregating all polymer lengths across α-CBs

with different A_{tot} (Fig 3) is acceptable for a length distribution analysis.

Polymerization Analysis

521   The total Rubisco concentration-dependence of the monomer- and polymer- subunit

concentrations were globally fitted to the chain length limited and unlimited isodesmic and cooperative

nucleation-elongation polymerization models, (Eqs. 1-4 and 7) using Origin software, with the dimer

treated as monomer or polymer depending on n = 2 or 3. A non-polymerizing fraction (non_poly) was

included but was zero for all conditions. Implicit equations of the monomer concentration A_1 were solved

with a custom numerical routine based on a root-finding algorithm implementing the Newton-Raphson

method combined with the Bisect method (65). Simulations were done with Matlab. For length-limited fits,

m=10 (untreated, DTT) or 11 (diamide), with $A_1 \leq K_{pol}$ but K_{pol} and K_{nuc} unconstrained.

Data availability

Untreated carboxysome data were previously published and can be found at the Electron Microscopy

Data Bank under accession code EMD-27654 and the Electron Microscopy public Image Archive under

accession code EMPIAR-11125. The Rubisco unfiltered half-maps and full filtered map from both treated

data sets have been deposited in the Electron Microscopy Data Bank under accession codes EMD-76204

and EMD-76205. The complete set of subtomograms for each sample as well as exemplar reconstructed

tomograms and frames have been deposited in the Electron Microscopy Public Image Archive under

EMPIAR-13403. All other data that support this study are available from the corresponding authors upon

reasonable request.

Acknowledgements

We thank Keith Meyers, Morgan Stephens, and Izabella Soswa for their assistance aligning

tomograms and processing data for subtomogram averaging. Molecular graphics and analyses performed

with UCSF Chimera (64).

Authors' contributions

L.A.M. and E.M.D.L.C. conceived and supervised the project. L.O. and D.F.S. provided the

purified carboxysome samples and contributed images of the Rubisco/csoS2 condensate. L.A.M.

prepared the samples for imaging and collected the tilt stacks. K.R., R.H.G., and L.A.M. processed the

tomograms, conducted the STA, and carried out the biophysical processing of the particles. W.C. and

E.M.D.L.C. established the mathematical theory and models, and interpreted the binding results. K.R.,

W.C., L.A.M., and E.M.D.L.C. wrote the manuscript. W.C. and K.R. prepared figures. K.R. finalized the

structures and the EMDB/EMPIAR deposition. All authors contributed to the analysis and interpretation of

the data.

Funding

This work was supported by funding from the Ralph W. and Grace M. Showalter Research Trust

and National Science Foundation CAREER award 2439291 to L.A.M., National Institute of General

Medical Sciences (NIGMS) award R35 GM136656 to E.M.D.L.C., and Department of Energy Physical

Biosciences Program award number DE-SC0016240 to D.F.S. K.R. was funded in part by NIGMS F32

Postdoctoral Fellowship (F32 GM161096), and R.H.G. by NIGMS T32 Training Grant GM132024. D.F.S.

is an Investigator of the Howard Hughes Medical Institute. A portion of this research was supported by

562   NIGMS award R24GM154185 and performed at the Pacific Northwest Center for Cryo-EM (PNCC) with

563   assistance from Vamseedhar Rayaprolu. The contents of this publication are solely the responsibility of

564   the authors and do not necessarily represent the official views of the funding agencies.

Conflicts of Interest

The authors declare the following competing interests: L.A.M. has received intellectual property

licensing payments from Eli Lilly and Company totaling less than $5,000 annually, as well as research

funding from Eli Lilly and Company, as part of a consortium grant. This work is unrelated to the subject

matter of this manuscript. D.F.S. is a co-founder of Scribe Therapeutics and a scientific advisory board

member of Scribe Therapeutics and Mammoth Biosciences. All other authors declare no competing

interests.

Tables

Table 1. Rubisco packing properties inside α-CB.

	untreated	DTT-treated	Diamide-treated	Data
Packing (average values)				
α-CB Volume(10^{-4} μm^3)	6.9 ± 2.1	5.3 ± 1.4	8.5 ± 3.4	Fig. 2A
Rubisco complexes per α-CB	242 ± 73	260 ± 77	363.4 ± 160	Fig. 2B
Rubisco concentration (μM)	583 ± 106	822 ± 100	718 ± 139	Fig. 2C
Rubisco complexes per 10^{-3} μm^3	340 ± 6	494 ± 7	427 ± 7	Fig. 2D
Rubisco [inner] vs. [total] slope	1.48 ± 0.07	1.30 ± 0.06	1.38 ± 0.04	Fig. S2
[inner] vs. [total] *x*-intercept (μM)	148 ± 5	82 ± 22	68 ± 6	Fig. S2

575
576

Table 2. Rubisco fibril chain properties inside α-CB.

	untreated	DTT-treated	Diamide-treated	Data
Distance between adjacent fibrils (nm)	10.9 ± 0.7	10.3 ± 0.5	10.4 ± 0.6	cryoET
Maximum fibril chain length, m (Rubisco)	10	10	11	Fig. 3
length decay const., κ (+dimer; Rubisco^{-1})	1.3 ± 0.2	1.8 ± 0.2	1.2 ± 0.1	Fig. 3
length decay const., κ (–dimer; Rubisco^{-1})	0.42 ± 0.03	0.46 ± 0.08	0.49 ± 0.02	Fig. 3

577
578

Table 3. Rubisco polymerization properties inside α-CB.

	untreated	DTT-treated	Diamide-treated	Data
data: monomer vs. polymer with dimer				
Nucleation Polymerization (from fits), nucleation n = 2,				
K_{pol} (μM, un-limit)[*]	473 ± 16	654 ± 5	660 ± 13	Fig. 4
K_{nuc} (μM, un-limit)[*]	9×10^{17}	5×10^{20}	6×10^{6}	Fig. 4
K_{pol} (μM, limit)[*]	516 ± 10	678 ± 12	686 ± 10	Fig. S4
K_{nuc} (μM, limit)[*]	5×10^{4}	1×10^{5}	1×10^{5}	Fig. S4
Isodesmic Polymerization (doesn't fit, for comparison only, values from fits)				
K_{pol} (μM, un-limit)[*]	3705 ± 176	5433 ± 230	7405 ± 324	Fig. S4
K_{pol} (μM, limit)[*]	3705 ± 208	5433 ± 230	7404 ± 324	Fig. S4
data: monomer+dimer vs. polymer without dimer				
Nucleation Polymerization (from fits), nucleation n = 3,				
K_{pol} (μM, un-limit)[*]	541 ± 44	762 ± 5	730 ± 170	Fig. 4
K_{nuc} (μM, un-limit)[*]	6×10^{12}	2×10^{21}	9×10^{9}	Fig. 4
K_{pol} (μM, limit)[*]	593 ± 15	794 ± 21	762 ± 15	Fig. S4
K_{nuc} (μM, limit)[*]	6×10^{7}	2×10^{8}	2×10^{8}	Fig. S4

[*] un-limit: un-limited length model. limit: limited length model.

582
Figure 1. Aligned Rubisco Particles from Tomograms. A. Negative stain TEM micrograph of a
Rubisco-CsoS2 condensate. **B.** Orthoslice of a tomogram containing α-CBs with examples of Rubisco
packing: dense (top), ordered (middle), and sparse (bottom). **C.** Rubisco densities resulting from STA of
DTT treated (left, orange) and diamide treated (right, blue) samples. **D.** Docked crystal structure (PBD ID:
6UEW) in DTT-treated density (left, yellow) and diamide-treated density (right, blue). **E-G.** Composite
chains of STA results with docked crystal structures of (PDB ID: 6UEW) for untreated Rubisco (left, gray),
DTT-treated (middle, orange) and diamide-treated (right, blue). **G.** Isolated Rubisco-Rubisco interfaces
resolved using STA. Untreated (left, gray), DTT-treated (middle, orange) and diamide-treated (right, blue).

592
593
594 **Figure 2. Rubisco packing in α-CBs. A-C.** Comparisons of α-CB volume (**A**), number of Rubisco
595 complexes in α-CB (**B**) and Rubisco concentration (A_{tot}) in α-CBs (**C**) of untreated (black), DTT-treated
596 (magenta), and diamide-treated (blue) samples. Data are summarized in Table 1. **D.** Correlation between
597 the number of Rubisco complexes packed inside an α-CB vs. the volume.
598

Figure 3. Length distribution of Rubisco fibrils in α-CBs. A-C. Histogram of polymer length for untreated, DTT-treated or diamide-treated. The continuous black lines are the best fits of the data to exponential decays including the dimer population (the first bar). The continuous gray lines (starting lower) are the best fits excluding dimers. Data available in Table 2.

Figure 4. Rubisco polymerization analysis. A-C: Unbound (monomer+dimer, light blue filled circles) and bound (polymer of length $j \geq 3$ Rubisco complexes; blue filled circles) Rubisco concentrations change as a function of the total Rubisco concentration inside untreated, DTT-treated, and diamide-treated α-CB samples. The smooth red lines through the corresponding data points represent the best global fits with implicit nucleation-polymerization equations for unbound (Eq. 3) and bound (Eq. 4) and the nucleation size of $n = 3$ held, respectively. The dashed vertical lines in all panels indicate the positions of K_{pol}.

Supplementary Materials:

616 *Morphological analysis*

Table S1A: Manual morphology classification of diamide-treated α-CBs			
Analyst	**# Sparse**	**# Dense**	**# Ordered**
1	**35**	**55**	**234**
2	**42**	**111**	**176**
3	**32**	**173**	**111**
Percentage	**11 +- 2 %**	**35 +- 18 %**	**54 +- 19 %**

617
618

Table S1B: Manual morphology classification of DTT-treated α-CBs			
Analyst	**# Sparse**	**# Dense**	**# Ordered**
1	**5**	**6**	**256**
2	**5**	**126**	**137**
3	**9**	**144**	**109**
Percentage	**2 +- 1 %**	**34 +- 28 %**	**63 +- 29 %**

619
620
Each analyst was instructed to classify morphology as sparse (>33% empty interior volume), ordered (>3
fibrils with >3 Rubisco subunits in one bundle), or dense (not ordered or sparse). Analysts needed to
search in three-dimensional space for Rubisco arrangements, and personal biases are strong: for
example, in DTT and diamide treated α-CBs (Supplementary Table 1A and B), analyst #1 identifies
patterns readily and classified most α-CBs as ordered, while analyst #3 struggled to classify α-CBs
without a strong central lattice and thus always found more dense than ordered morphologies. However,
analysts were not told what to expect from the datasets, nor were they allowed to see each other's
results. The morphology classification Table for untreated α-CBs was reported previously (26).
629
630 Many of the discrepancies in the morphology classification arise when α-CBs contain multiple short
631 chains, which can be difficult to identify manually (Supplemental Fig. S1)
632

633
634
635 **Figure S1. Fibril arrangement in representative α-CB.** Representative ambiguous DTT-treated α-CB
636 that contains multiple short fibrils arranged radially.
637
638

639
640 **Figure S2. Inner Rubisco concentration. A-C.** Correlations of inner Rubisco concentration vs. total
641 Rubisco concentration for untreated, DTT-treated, and diamide-treated in α-CBs. All three panels (**A-C**)
642 are plotted in the same *x*- and *y*-axis scales for easy visual comparison of the slopes of line fits (straight
643 lines through the data). **D.** Comparison of inner Rubisco concentrations for untreated (black), and DTT
644 treated (magenta) and diamide treated (blue) samples. Data in all panels are summarized in Table 1.
645

646
Figure S3. Simulation of maximum chain length (m) dependence of nucleation polymerization
647
curves. A. Total subunit concentration A_{tot} dependence of monomer A_1 and polymer A_{pol} Rubisco (dimer
648
and longer) concentrations calculated by solving the maximum chain length limited model (Eq. 7) for m =
649
5 (red), 10 (green), 20 (blue), 50 (cyan) and 100 (magenta) and chain length unlimited model (Eq. 3,
650
equivalent to $m = \infty$, thick black). The other parameter values: K_{pol}, K_{nuc}, and $n = 2$ are same for all and
651
indicated on the plot. The horizontal arrows with m above them indicate the curve shape trend with
652
increasing m value. The horizontal thin dashed line indicates K_{pol} where the monomer concentration
653
exceeding it, $A_1 > K_{pol}$, is thermally unfavorable. For the length unlimited nucleation model (black line), A_1
654
is always less than, but approaches K_{pol}. On the other hand, the length limited nucleation model can have
655
$A_1 > K_{pol}$ when A_{tot} is large enough. The curve trend shows when $m = 100$ (magenta line), the
656
polymerization curves of limited length model are almost the same as those of length unlimited model
657
(black line). **B.** The simulated curves are the same curves as those in panel **A** except confining $A_1 \leq K_{pol}$,
658
such that the region $A_1 > K_{pol}$ of all simulated monomer curves in Panel A have been cut off by the
659
horizontal thin dashed line. When confining $A_1 \leq K_{pol}$, the subunit concentration in polymer form also
660
confined, therefore, the polymer curves are also cut off. The purpose of Panel B is to show which part of a
661
simulated curve for a limited length polymerization model has the monomer concentration $A_1 > K_{pol}$ by
662
comparing the two panels.
663
664

Figure S4. Comparison different fitting model. A-C. Plots of the total Rubisco concentration dependence of unbound Rubisco concentration composed of monomers only (light gray filled circles) and bound Rubisco concentration in dimer and longer polymers (gray filled circles). The unbound and bound data are globally fitted with a length unlimited nucleation (n = 2, red lines), length unlimited isodesmic (green lines), length limited nucleation (n = 2, magenta lines) or length limited isodesmic (blue dashed lines) model in untreated, DTT-treated, and diamide-treated in α-CBs. The best global fit result by length limited isodesmic (blue dashed lines) model is basically the same as that by length unlimited isodesmic (green lines), and the two fit lines are on top one another for all three treated and untreated sample (A-C, Table 3). **D-F.** Total Rubisco concentration dependence of unbound Rubisco concentration in monomers and dimers (light blue filled circles) and bound Rubisco concentration in length ≥ 3 polymers (blue filled circles). The data are globally fitted with a length unlimited nucleation (n = 3, red lines) or length limited nucleation (n = 3, magenta lines) model in untreated, DTT- treated, and diamide-treated in α-CBs. In all fits, the unbound concentration A_1 (monomer only in n = 2 nucleation and isodesmic models, whereas monomer+dimer n = 3 nucleation model) is constrained as $A_1 < K_{pol}$. In all length limit models, the maximum chain length is 10 Rubisco complexes for untreated and DTT treated, and 11 Rubisco complexes for diamide treated. The magenta fit lines for nucleation with length limit in all panels stop and don't go all the way to the right as other fitting models do. It is because in the A_t region beyond stopping point, there is no $A_1 < K_{pol}$ solution in Eq. 7. All the fitting results are listed in Table 3 for comparison.

Figure S5. Polymerization inflection point. A large K_{nuc} value increases the sharpness of transition at the inflection point. The monomer (light filled gray circles) and polymer (including dimers, gray filled circles) Rubisco complexes are the same data as untreated Rubisco complexes in Fig. S4A. The smooth thick black lines are the global fits of the length unlimited model to the data with $n = 2$ (Table 3). The other smooth lines are simulations with the same n and K_{pol}, but $K_{nuc} = 10^5$ (magenta), 10^6 (red), 10^7 (blue) and 10^{10} (green) μM. The up and down arrows on the plot indicate the direction of simulated monomer (up) and polymer (down) lines approaching with increasing K_{nuc} values at the transition. The larger the K_{nuc} value is, the sharper the transition is. When $K_{nuc} = 10^{10}$ (green), the simulated lines are not much different from the global fitting lines (thick black) to the data.

Theory of Protein Polymerization

Protein polymerization can be described as a highly cooperative, concentration-dependent phase

transition, analogous to condensation or crystallization (49). This transition (between monomer and

polymer phases) occurs at a protein concentration called the critical concentration for polymerization.

When the total protein concentration is less than the critical concentration, the protein exists in the

monomeric form (phase 1). At total protein concentrations above the critical concentration, polymer forms

(phase 2).

Assembled polymer exists in a reversible equilibrium with monomers, which remain at a

concentration approximately equal to the 'critical concentration' such that the total polymer concentration

is equal to the protein concentration exceeding the critical concentration. For this reason, the critical

concentration is sometimes called the critical monomer concentration. The critical concentration of linear

polymers is also approximately equal to the overall equilibrium binding constant for monomer

incorporation at filament ends (32, 49). The critical concentration is constant under a defined set of

conditions and can be influenced by regulatory proteins, chemical modifications, temperature, and

solution conditions, including crowding (8).

The critical concentration is usually determined from the apparent inflection point by linear

extrapolation of the linear protein concentration-dependence of polymer and/or monomer concentrations

(49, 50). However, the inflection points of our Rubisco binding curves (Fig. 4) are difficult to determine

accurately by simple linear extrapolations from the regions before and after the critical concentration

because of intrinsic noise inherited from the image processing of tomograms. We, therefore, determine

the inflection points and critical concentrations from global fits of the monomer and polymer concentration

curves (Fig. 4 and supplementary Fig. S4). We also assess how various polymer assembly models,

specifically isodesmic and nucleation (with nucleus $n = 2$ and 3 Rubisco complexes) with limited length or

unlimited length, account for the experimental data.

General polymerization theory

The following reaction scheme is general for the growth (i.e., elongation) of a self-associating

linear polymer (40):

$$L_j + L_1 \underset{k_{pol,-}}{\overset{k_{pol,+}}{\rightleftharpoons}} L_{j+1}$$

(Scheme 1),

where L_j is a polymer chain consisting of j (=1, 2, …) subunits and L_1 is a monomer (or "polymer" chain

comprised of 1 subunit). This reaction is a reversible with an overall dissociation equilibrium constant

(K_{pol}), such that polymer L_j becomes L_{j+1} when a monomer incorporates and L_{j-1} when a single subunit

dissociates.

Linear polymer assembly often occurs through the formation of a "nucleus", defined as the

oligomer size that binds incoming subunits the same as long polymers (affinity K_{pol}; discussed below).

Oligomers smaller than the nucleus size are thermodynamically unstable, are thus populated at low

concentrations, and tend to rapidly dissociate. In cases where polymerization proceeds by a nucleation-

elongation mechanism, the nucleus is defined as an oligomer of n subunits (n >1):

$$\underbrace{L_1 + L_1 + \ldots + L_1}_{n} \overset{K_{nuc}}{\rightleftharpoons} L_n \quad n \geq 2$$

(Scheme 2),

where K_{nuc} is the overall dissociation equilibrium constant for nucleus formation. Once a nucleus is

formed, polymer elongation follows Scheme 1. In this case, $j \geq n$ in Scheme 1.

Isodesmic polymerization theory

When the nucleus size n = 1, polymerization does not proceed via a nucleation-elongation

mechanism; a monomer can directly bind another to form the polymer (Scheme 1). This is a special case

called isodesmic polymerization. In this case, Scheme 2 is part of Scheme 1 with $K_{nuc} = K_{pol}$ and can be

ignored.

According to established theories of isodesmic polymerization (40, 50), the monomer

concentration in equilibrium with stable polymer (chain length $i \geq 2$ subunit) is given by the following

relation:

$$A_1 = \frac{K_{pol}A_{tot}}{A_{tot} + \frac{K_{pol}}{2} + \sqrt{\left(A_{tot} + \frac{K_{pol}}{2}\right)^2 - A_{tot}^2}} \rightarrow \begin{cases} \sim A_{tot}, & A_{tot} \ll K_{pol} \\ K_{pol}, & A_{tot} \gg K_{pol} \end{cases}$$ Eq. 1,

where $A_1 = [L_1]$ is the monomer concentration in reversible equilibrium with polymers, A_{tot} is the total

concentration of all the polymerizing macromolecules, and K_{pol} is the polymerization equilibrium constant

(Scheme 1). The value of K_{pol} is the critical concentration for polymerization(51).

At equilibrium (Eq. 1), the free monomer concentration (A_1) is approximately equal to the total

macromolecule concentration ($A_1 \sim A_{tot}$) when $A_{tot} \ll K_{pol}$, and levels off when $A_{tot} \sim K_{pol}$, yielding a

curved, gradual transition around the value of K_{pol} in a plot of monomer concentration versus total

macromolecule concentration (52, 53) (Fig. S4). When $A_{tot} \gg K_{pol}$, the free monomer concentration (A_1)

infinitely approaches the constant K_{pol} value.

Because the total macromolecule concentration is the sum of monomeric (A_1) and polymerized

subunits (A_{pol}; $A_{tot} = A_1 + A_{pol}$), the concentration of polymerized subunits (A_{pol}) is given by:

$$A_{pol} = A_{tot} - A_1 = \sum_{j=2}^{\infty} jA_j = 2A_2 + 3A_3 + \cdots$$

$$= A_{tot}\left(1 - \frac{K_{pol}}{A_{tot} + \frac{K_{pol}}{2} + \sqrt{\left(A_{tot} + \frac{K_{pol}}{2}\right)^2 - A_{tot}^2}}\right) \rightarrow \begin{cases} \sim 0, & A_{tot} \ll K_{pol} \\ A_{tot} - K_{pol}, & A_{tot} \gg K_{pol} \end{cases}$$ Eq. 2

where $A_j = [L_j]$, j = 1, 2, 3 … is the concentration of polymers with chain length equal to i subunits. The

approximated expression (far right of Eq. 2) indicates the polymerized subunit concentration is negligible

($A_{pol} \approx 0$) when the total macromolecule concentration is much less than the polymerization equilibrium

constant ($A_{tot} \ll K_{pol}$), whereas it increases approximately linearly with the total macromolecule

concentration ($A_{pol} \approx A_{tot} - K_{pol}$) when it far exceeds the polymerization equilibrium constant ($A_{tot} \gg K_{pol}$).

This relationship yields K_{pol} as the center of transition in polymer concentration versus total

macromolecule concentration plots as well (52, 53). The transitions are gradual rather than abrupt for

isodesmic polymerization, because the transitions are not cooperative.

Cooperative nucleation-elongation polymerization theory

Cases of linear polymer assembly in which $n \geq 2$ monomers form a "nucleus" for polymerization

are cooperative (i.e., free monomers bind nuclei and polymers longer than nuclei more strongly than other

monomers) and are referred to as nucleation-elongation polymerization. In these cases, the equation for

monomer concentration (A_1) in reversible equilibrium with polymer is given by (40):

773
$$\left(1 - \frac{A_1}{K_{pol}}\right)^2 (A_t - A_1) - \frac{K_{pol}^n}{K_{nucl}}\left(\frac{A_1}{K_{pol}}\right)^n\left((n-1)\left(1 - \frac{A_1}{K_{pol}}\right) + 1\right) = 0 \qquad \text{Eq. 3}$$

774   where the nucleation constant (K_{nuc}) is the overall dissociation equilibrium constant for nucleus formation

775   (Scheme 2). Eq. 3 is an implicit polynomial equation of A_1 and it can be solved numerically for A_1, from

776   which the subunit concentration in all polymers is calculated from mass conservation according to $A_{pol} =$

777    $A_{tot} - A_1$, yielding:

778
$$A_{pol} = A_{tot} - A_1 = \sum_{j=n}^{\infty} j A_i = n A_n + (n+1)A_{n+1} + \cdots \qquad \text{Eq. 4.}$$

Eq. 4 treats non-polymer oligomers smaller than the nucleus as "monomers". These intermediates are

unstable and exist at low concentrations. Oligomers equal to or longer than the nucleus size are treated

as "polymer".

Eqs. 3 and 4 (or Eqs. 1 and 2 for isodesmic polymerization) can used to fit the total

macromolecule concentration (A_{tot})-dependence of both monomer (A_1) and polymer populations (A_{pol}).

When numerically solving Eq. 3 for A_1, $0 \leq A_1 \leq A_{tot}$ since A_1 is the monomer concentration. On the other

hand, a value of $A_1 \geq K_{pol}$, is singularity of Eq. 4 and the summation leading to Eq. 3 (54), so the

mathematically allowed solution of polynomial Eq. 3 must be $A_1 < K_{pol}$. Therefore, the combined

constraint for A_1 value is in the range $0 \leq A_1 < \min(K_{pol}, A_{tot})$, where the term $\min(K_{pol}, A_{tot})$ means the

smaller value of K_{pol} and A_{tot}.

Similar to the isodesmic case, when $A_{tot} << K_{pol}$, i.e., $A_1 < A_{tot} << K_{pol}$, $A_1 \sim A_{tot}$ (Eq. 3), indicating

that the free monomer concentration at equilibrium (A_1) increases proportion to the total macromolecule

concentration, but approaches the K_{pol} value when $A_{tot} >> K_{pol}$. Also similar to isodesmic polymerization,

the polymer subunit concentration (A_{pol}) is negligible when the total macromolecule concentration is low

($A_{tot} << K_{pol}$), and the total polymer concentration (A_{pol}) approaches $A_{pol} - K_{pol}$ when $A_{tot} >> K_{pol}$. Therefore,

K_{pol} is the observed critical concentration (i.e., inflection point) in the plots of monomer/polymer versus

total mscromolecule concentration (*Bull. Math. Biol.* **60,** 449–475 (1998)).

Polymerization theory within a confined volume

798   The polymerization theory above assumes unconstrained polymerization in an infinite accessible

volume. The total concentration of macromolecule and polymer length in infinite accessible volume is

unlimited. However, the length distribution remains an exponential decay (28,30). When polymerization

occurs under confinement, as is the case of Rubisco polymerization inside α-CBs, the maximum

concentration of polymerization macromolecules is limited by the exclusion volume of polymerization

units. Since the volume excluded by a single Rubisco is ~ $10 \times 10 \times 10 = 1000$ nm^3 (Table 2), and the

volume of a typical untreated α-CB is ~ 7×10^{-4} μm^3 = 7×10^5 nm^3 (Table 1), the maximum number of

Rubisco complexes that can fit into a typical α-CB is given by the ratio of the two volumes (i.e., N_{max} ~

7×10^5 nm^3/10^3 nm^3 = 700, which is equivalent to $700/(6.02 \times 10^{17}$ μmol$^{-1})/7 \times 10^{-4}$ μm^3 ~ 1700 μM).

However, the maximum observed Rubisco concentration (A_{tot}) in our untreated α-CB samples is ~785 μM

(Fig. 4A), less than half of the estimated theoretic value of 1700 μM, suggesting that not all the α-CB

volume is accessible to Rubisco.

810   The polymer length in a confined volume is naturally limited by the accessible volume as well.

Again in the case of Rubisco polymerization in α-CB, an untreated α-CB with typical volume of 7×10^{-4}

μm^3 is equivalent a 55 nm radius ball (see packing section), which is about the size of 5.5 Rubisco

complexes lined up (size of a Rubisco ~10 nm, see above), indicating that a maximum of only 10 – 11

polymerized Rubisco complexes can potentially occupy the diameter of a typical untreated α-CB. This is

precisely the observed maximal Rubisco chain length (10 – 11 Rubisco complexes) observed in the data

for all treated and untreated samples (Table 2). Therefore, confinement appears to contribute to the

observed Rubisco polymer distribution. Accordingly, the polymerization theory presented above must be

modified to consider confinement and subsequent geometric constraints.

819   Scheme 1 and 2 above still apply under confinement, but the maximum possible total

macromolecule concentration (A_{tot}) and length is limited (L_m). When A_{tot} is limited by confinement before

the polymer length is limited, the theories of isodesimic and nucleation polymerizations are the exactly the

same as above (Eq. 1-4), and the analysis remains the same.

In contrast, when confinement limits the polymer length before it limits A_{tot}, a different treatment is

needed, which we present here. Defining m as the maximum number of subunits in a polymer chain, then

L_m is a polymer with the maximum length and its concentration is $A_m = [L_m]$, which obeys the following

equation

827
$$\frac{dA_m}{dt} = k_{on,pol}A_1A_{m-1} - k_{off,pol}A_m$$
Eq. 5

828   where $k_{on,pol}$ and $k_{off,pol}$ are the average on- and off-rate constants for monomers with concentration A_1 to

829   incorporate at the either end. The right hand side of Eq. 5 lacks the terms linking A_{m+1} (see Ref (40)), the

830   concentration for L_{m+1}. Despite this difference, the general solution for the concentration of polymers with j

831   number of subunits is the same as that in an unlimited space (40), i.e. an exponential function of j

832   according to:

833
$$A_j = \frac{K_{pol}^n}{K_{nuc}}\left(\frac{A_1}{K_{pol}}\right)^j = \frac{K_{pol}^n}{K_{nuc}}e^{-\kappa j}, \qquad j = n, n+1, \dots, m$$
Eq. 6

834   Where polymerization constant (or critical concentration) $K_{pol} = \frac{k_{off,pol}}{k_{on,pol}}$ and the exponential decay

835   constant $\kappa = -\ln\left(\frac{A_1}{K_{pol}}\right)$. The total polymerizing macromolecule concentration A_{tot} is equal to the sum of

836   monomer concentration (A_1) and the polymerized subunits (A_{pol}) as follows:

837
$$A_{tot} = A_1 + \sum_{j=n}^{m} jA_j = A_1 + \frac{K_{pol}^n}{K_{nuc}}\sum_{j=n}^{m} j\left(\frac{A_1}{K_{pol}}\right)^j$$
Eq. 7

838
$$= A_1 + \frac{\frac{K_{pol}^n}{K_{nuc}}}{\left(1 - \frac{A_1}{K_{pol}}\right)^2}\left(\left(\frac{A_1}{K_{pol}}\right)^n\left(n - (n-1)\left(\frac{A_1}{K_{pol}}\right)\right) - \left(\frac{A_1}{K_{pol}}\right)^{m+1}\left(1 + m - m\left(\frac{A_1}{K_{pol}}\right)\right)\right)$$
Eq. 7'

839   Eq. 7 is a polynomial equation for the monomer concentration A_1. It has one and only one positive

840   solution in the region $0 < A_1 < A_{tot}$, which can be readily solved numerically for any given m ($> n$), K_{pol} and

841   K_{nuc} values. Therefore, Eq. 7 applies for polymerization under length confinement in a limited space,

842   where Eq. 3 applies above for polymerization in an unlimited space.

843   Under length confinement, A_1 is obtained by solving Eq. 7, the concentration of polymerized

844 subunits can be calculated using Eq. 4 with the summation up to m instead of infinity, or calculated by A_{pol}

845 $= A_{tot} - A_1$, as presented above for an unlimited space. For isodesmic polymerization in a confined volume

846 with a limited maximum chain length m, Eq. 7 can still be used with $K_{nuc} = K_{pol}$ and the summation starting

847 $n = 2$. Note that for isodesmic polymerization, the integer number of $n = 2$ in Eq. 7 reflects the summation

848 starting number because it doesn't need to form nuclei for polymerization or the number of molecules to

849 form nucleus is $n = 1$ (see above discussion), whereas, for nucleation polymerization ($n \geq 2$), the number

850 n in Eq. 7 means both summation starting and nucleus number.

851   Eq. 6 is the general expression for concentration of polymers with a given number of subunits (j)

852 for a length-unlimited polymer model as well (40). However, according to Eq. 6, in the monomer

853 concentration region of $A_1 \geq K_{pol}$, the polymer concentration A_j increases with the chain length j subunits,

854 according to Eq. 8:

855
$$A_{j+1} = \frac{K_{pol}^n}{K_{nuc}} \left(\frac{A_1}{K_{pol}} \right)^{j+1} = \frac{K_{pol}^n}{K_{nuc}} \left(\frac{A_1}{K_{pol}} \right)^{j} \left(\frac{A_1}{K_{pol}} \right) = A_j \left(\frac{A_1}{K_{pol}} \right) > A_j, \quad j = n, n+1, \ldots \qquad \text{Eq. 8.}$$

856 Longer polymers are thermodynamically less stable than shorter ones because statistically they have

857 potential sites for fragmentation (47). Accordingly, there should be fewer long polymers than short

858 polymers observed in the length distribution. Hence, for thermodynamically stable polymers $A_1 < K_{pol}$ is

859 required in both length-limited and -unlimited models. In the case of a length-unlimited polymer model,

860 substituting A_j from Eq. 6 into Eq. 4 for calculating A_{pol} will not converge if $A_1 \geq K_{pol}$, since A_{pol} approaches

861 infinity (i.e., $A_1 \geq K_{pol}$ is singularity of Eq. 4). Therefore, in a length-unlimited polymer model $A_1 < K_{pol}$ is not

862 only a thermodynamic requirement, but also a mathematical requirement.

863   For polymerization under polymer length confinement, $A_1 \geq K_{pol}$ is no longer a problem for the

864 polynomial equation Eq. 7. Thus, mathematically, $A_1 \geq K_{pol}$ can be a solution of Eq. 7. In this case, the

865 monomer concentration A_1 is limited only by A_{tot} since it should never be more than A_{tot}, but it can

866 potentially increase with A_{tot} continuously unlike that A_1 has a constant mathematical limit in an unlimited

867 space. Nevertheless, when $A_1 > K_{pol}$, Eq. 6 indicates that the polymer concentration A_j increase with its

868 length exponentially, i.e., the larger the chain length j, the more concentrated the polymer as noted above.

869 This is thermodynamically unfavorable. For this reason, we can still confine $A_1 < K_{pol}$ to conform to the

thermodynamic stability, as we do in this work (see below). When constraining $A_1 \neq K_{pol}$, the polynomial in

Eq. 7 can sum up to Eq. 7'. Eq. 7' is another form of Eq. 7 and can be useful in some cases. Since $A_1 =$

K_{pol} is a pseudo singularity point of Eq. 7', Eq. 7' would have problem at the point in programming and

care must be taken when using it.

When the maximum chain length m value gets longer (i.e., approaches infinity), the length limited

model should behave like a length-unlimited model. This prediction was confirmed using simulated

polymerizations with various maximum chain lengths and shared n, K_{pol} and K_{nuc} values (Fig. S3);

polymerization with $m = 50$ (cyan) is very close to the length-unlimited polymer, and essentially identical

when $m = 100$ (magenta).

Both the maximum polymer length (m) and the total macromolecule concentration (A_{tot}) are

limited by the accessible volume as discussed above. If A_{tot} reaches a limit first, the maximum polymer

chain length can't grow infinitely. Similarly, if the monomer concentration A_1 is limited to K_{pol}, i.e., $A_1 < K_{pol}$

under the thermodynamic stability consideration, all the polymer concentrations are also limited according

to Eq. 6, and then, the maximum allowed total macromolecule concentration A_{tot} in the length limited

model is also automatically limited (Fig. S3B) by the mass conservation of Eq. 7, i.e.,

885
$$A_{tot} = K_{pol} + \frac{K_{pol}^n}{K_{nuc}} \sum_{j=n}^{m} j \left(\frac{K_{pol}}{K_{pol}}\right)^j = K_{pol} + \frac{K_{pol}^n}{K_{nuc}} \frac{(m+n)(m-n+1)}{2}$$
Eq 9,

In the region of $A_1 < K_{pol}$, the simulated polymerizations in Fig. S3B are exactly the same as those in Fig.

S3A, but in the region $A_1 > K_{pol}$, the polymerizations in Fig.S3B have no data (Eq. 7 doesn't have $A_1 < K_{pol}$

solution). The longer the maximum allowed chain length m is, the larger the maximum allowed A_{tot} is (Eq.

9 and Fig. S3B). The maximum allowed chain length goes hand to hand with the maximum allowed A_{tot}

and when the maximum allowed chain length approaches infinity, so does the maximum allowed A_{tot}.

The mathematical models for polymerization presented here and use in this work (Eqs. 1-9) do

not consider polymer fragmentation and annealing, which can potentially affect polymerization and the

length distribution (55). However, we assume Rubisco polymer fragmentation and annealing occur much

less frequent than other linear polymers (e.g., actin) since they are relatively short by comparison.

Alternative Free Energy Proof

In Eq. 15 or 15' of the main manuscript, we obtain the relation of the two standard free energy changes from three different approaches, i.e., one from the solution (Eq. 6) of the kinetic equations associated with the general polymerization schemes Scheme 1 and 2, another one from the standard free energy change for binding ($\Delta G^{0'}_{pol}$, Eq. 11 and 14) and the third from a separated thermodynamic study of polymerization for polymer free energy increase for elongating one subunit ($\Delta G^{o'}_{j \to j+1}$, Eq. 10; (33)). This relationship between $\Delta G^{0'}_{pol}$ and $\Delta G^{o'}_{j \to j+1}$ (Eq. 15 and 15') can be independently obtained directly from the Gibbs free energy calculation. The free energy of a population of j-subunits-long polymer (G_j) scales with its concentration (A_j) according to (66):

$$G_j = G^{0'}_j + k_B T \ln\left(A_j\right)$$ 16·

At equilibrium, the free energy change ($\Delta G_{j \to j+1}$) associated with subunit incorporation (i.e. polymer of j subunits elongating to $j + 1$ subunits) is 0 (66), as given by:

$$\Delta G_{j \to j+1} = G_{j+1} - G_j = G^{0'}_{j+1} - G^{0'}_j + k_B T \ln\left(A_{j+1}\right) - k_B T \ln\left(A_j\right) = 0$$ 17

and can be rearranged to:

$$\Delta G^{o'}_{j \to j+1} = G^{0'}_{j+1} - G^{0'}_j = k_B T \ln\left(\frac{A_j}{A_{j+1}}\right) = k_B T \ln\left(\frac{A_j A_1}{A_{j+1}} \frac{1}{A_1}\right) = k_B T \ln\left(\frac{K_{pol}}{A_1}\right) = \Delta G^{0'}_{pol} - k_B T \ln(A_1)$$ 18

Therefore, Eq. 18 is Eq. 15, and it can be independently obtained directly from the Gibbs free energy as defined in college textbooks.

Extended Materials and Methods

α-Carboxysome Preparation

We purified α-CBs from wild-type *H. neapolitanus* (56). Briefly, wild-type *H. neapolitanus* cells

were grown in DSMZ-68 medium in a 10 L chemostat (pH 6.4, ambient air sparged, 30 °C), harvesting

cells every 2–3 days by 6000 × g centrifugation. A ~8 g cell pellet (resulting from 10–15 L of culture) was

chemically lysed and purified by centrifugations (12k × g 15 min, 40k × g 30 min) and sucrose gradient

(25 mL, 10–50% sucrose step gradient, centrifuged 105k × g 35 min). Quality and purity of sample was

analyzed by Coomassie stained SDS-PAGE and negative stain TEM. The final sample had an A280 of

roughly 10 AU and was stored at 4 °C in TEMB buffer (10 mM Tris pH 8, 1 mM EDTA, 10 mM $MgCl_2$, 20

mM $NaHCO_3$) until use.

Negative stain imaging of condensate

Rubisco (0.5 μM) of L8S8 and the CsoS2 (2.7 μM) were mixed at room temperature for 1 hour.

The sample was diluted 5x and Rubisco and CsoS2 condensates were visualized by negative-stain TEM.

Formvar/carbon-coated copper grids were prepared by glow discharge before sample application. The

grids were washed with deionized water several times before staining with 2% (w/v) uranyl acetate.

Imaging was performed on a JEOL 1200 EX transmission electron microscope.

Sample Conditions and Tomography

10 nm gold colloids (Aurion M-CEM-25512) were mixed with 5% BSA in PBS to passivate the

surface, then were buffer-exchanged into TEMB buffer and concentrated ~100x. The α-CB sample was

diluted with the BSA-coated gold in a 4:1 ratio. Samples were treated with 5 mM diamide or DTT for one

hour, because published work on *Salmonella* BMC shells has established that small compounds

penetrate the shell within that time (57). Samples were plunge-frozen onto glow-discharged C-flat 2/2-300

copper TEM grids using a Vitrobot. Imaging was performed at the Pacific Northwest Center for Cryo-

Electron Microscopy on a Titan Krios G4 with a Gatan K3 camera and BioQuantum energy filter.

Tomograms were collected in SerialEM according to Supplementary Table 2. Frame alignment, 2D image

processing, and tilt alignment were performed in etomo; CTF estimation in ctffind4; and tomogram

reconstruction in novaCTF following previously published pipelines (58–61).

Subtomogram averaging

Particles were identified in SIRT-like filtered tomograms using a customized workflow designed to

identify every Rubisco complex within each α-CB (26, 30). The estimated false negative and false positive

rates were 1–2% of total Rubisco particles within a α-CB when compared with manual identification in a

clear, high-defocus tomogram. Unfiltered map resolutions were not degraded by using every single

particle identified, further strengthening the evidence for the strong success rates in particle identification.

Resolution was calculated using a loose mask in Dynamo; final mask-corrected FSC was calculated in

RELION5 using a five-pixel extended mask derived from the low-pass filtered 6UEW PDB structure.

The general subtomogram averaging workflow was based upon a previous high-resolution

approach originally derived from the Briggs group (9, 26). Following particle identification, subtomograms

were re-extracted from 3D CTF-corrected weighted back projection tomograms. α-CBs were divided into

half-sets, and subtomogram alignment and averaging was performed in Dynamo v.1.1.532, using the

minimum number of iterations possible at each bin size and employing adaptive bandpass filtering

throughout (62). C4 symmetry was tested for accuracy, then imposed after searches were restricted to

<90-degree rotations. The best 80% of particles according to cross-correlation were used to generate the

reference for the next iteration, but all particles were aligned and participated in the next round. Masks

throughout were kept loose. When the resolution ceased to improve by at least 1 Fourier pixel per

iteration, the half-maps were sent to RELION5 for post-processing and final resolution calculations (63).

Rigid body docking was performed in UCSF Chimera (64).

The reference used for STA was first developed fully *de novo* utilizing a low-resolution average of

124 Rubisco complexes excluded from the dataset. Following a full STA procedure that brought the

resolution subnanometer, the *de novo* reference was compared to existing Rubisco structures and

confirmed to match the 6UEW structure in all visible helices and bulky side chains. We then performed a

second round of STA on the independent half-sets using the adaptive bandpass-filtered 6UEW structure,

which refined the resolution further.

The Rubicso-Rubisco interface was generated by isolating all particles in trimers. Averaged

particles were shifted within the box to center the interface and a mask was generated to isolate the

interface. The model (Fig. 1E,F) showing the stacked Rubisco-Rubisco dimer was generated by fitting the

monomer maps into the masked interface volume. The interface (Fig. 1G) is the STA volume generated

by Dynamo.

Data Analysis

We used a previously published Matlab script package to determine Rubisco binding states (31).

Briefly, the particle position and orientation were used to determine whether two adjacent Rubisco

complexes were likely to be bound, using the parameters of binding requiring an inter-particle distance of

9.936 nm and angle of 25°. We previously determined that Rubisco complexes immediately adjacent to

the shell appear to behave differently (26), and thus allow calculations both including and excluding these

particles from analysis (31).

Rubisco Filament Length Distribution Analysis

At equilibrium, the polymer length distribution is an exponential function of A_1 (Eq. 6). On the

other hand, the remaining unpolymerized monomer concentration A_1 is function of A_{tot} (Eq. 1, 3, and 7,

depending on the polymerization model), and increases with A_{tot}, but $A_1 \to K_{pol}$ as $A_{tot} \to \infty$.

Because the polymer length distribution function depends on A_1 (Eq. 6), and A_1 depends on A_{tot}

(Eqs. 1,3, and 7), different A_{tot} yield different polymer length distributions. Accordingly, using Eq. 6 to fit

the length distribution data to an exponential requires that all data be acquired at the same A_{tot}.

Nevertheless, for a nucleation polymerization, polymers appear only when $A_{tot} > K_{pol}$, and in this region,

$A_1 \sim K_{pol}$ and is almost constant and changes little with A_{tot} (Fig. 4). Consequently, the polymer length

distribution (Eq. 6) changes very little with A_{tot}, and aggregating all the polymer lengths inside different α-

CBs with different A_{tot} (Fig 3) is acceptable for a length distribution analysis. Since the exponential decay

constant $\kappa = -\ln\left(\frac{A_1}{K_{pol}}\right)$ changes little with A_{tot}, it can be used as a characteristic property of

polymerization, which can be compared among different macromolecules, as done previously(32).

Polymerization Analysis

The total Rubisco concentration-dependence of the monomer- and polymer- (including dimer for

$n = 2$) subunit concentrations were globally fitted to the chain length limited and unlimited isodesmic (Eqs.

1 and 2 for length unlimited; Eq. 7 with $K_{nuc} = K_{pol}$ for length limited) and $n = 2$ cooperative nucleation-

elongation (Eqs. 3 and 4 for length unlimited; Eq. 7 for length limited) polymerization models, whereas the

total Rubisco concentration-dependence of the monomer+dimer- and polymer- (excluding dimer) were

globally fitted to the chain length limited and unlimited $n = 3$ cooperative nucleation-elongation (Eqs. 3 and

4 for length unlimited; Eq. 7 for length limited) polymerization models. The fitting was performed using

Origin software (OriginLab, MA, USA; www.OriginLab.com). The fitting procedure was controlled by a

user-defined parameter (poly_switch) that fits each data set, unbound or bound, to a different equation,

with some parameters shared among the equations. When poly_switch = 0, the unbound species,

monomer concentration (A_1) or monomer+dimer concentration is fitted by solving Eq 1 (isodesmic), or Eq.

3 or Eq. 7 (nucleation); when poly_switch = 1, the polymer subunit concentration (A_{pol}) is fitted using A_{tot} –

A_1 with the monomer value (A_1) obtained from Eq. 1 or Eq. 3 or Eq. 7, respectively. Another parameter,

non_poly, was introduced in the Origin fitting program as the faction of the species that is not able to

polymerize in the observed A_{tot} range. In fitting, the combined monomer concentration is calculated as

non_poly*A_{tot} +A_1 and the A_1 of polymerization species is determined with the polymerization species (1 –

non_poly)* A_{tot}. The analyses of all three treated and untreated Rubisco polymerization show non_poly =

0 within the uncertainties.

The simulated data according to the four polymerization models were done with Matlab

(MathWorks, MA, USA; www.mathworks.com)

Eq. 3 and Eq. 7 are implicit equations of the monomer concentration A_1. It was solved for A_1 in

fitting procedures and simulations with a custom numerical routine based on a root-finding algorithm

implementing the Newton-Raphson method combined with the Bisect method (65).

In the polymer length limited analyses, the length $m = 10$ for untreated and DTT treated, but 11

for diamide treated, we also confined the monomer concentration $A_1 \leq K_{pol}$, but let K_{pol} and K_{nuc} values

unconstrained.

1025

References

1. K. A. Johnson, R. S. Goody, The Original Michaelis Constant: Translation of the 1913 Michaelis–Menten Paper. *Biochemistry* **50**, 8264–8269 (2011).

2. R. P. Bhattacharyya, A. Reményi, B. J. Yeh, W. A. Lim, Domains, Motifs, and Scaffolds: The Role of Modular Interactions in the Evolution and Wiring of Cell Signaling Circuits. *Annu. Rev. Biochem.* **75**, 655–680 (2006).

3. G. D. Stormo, D. S. Fields, Specificity, free energy and information content in protein–DNA interactions. *Trends Biochem. Sci.* **23**, 109–113 (1998).

4. J. A. Marsh, S. A. Teichmann, Structure, Dynamics, Assembly, and Evolution of Protein Complexes. *Annu. Rev. Biochem.* **84**, 551–575 (2015).

5. T. D. Pollard, A Guide to Simple and Informative Binding Assays. *Mol. Biol. Cell* **21**, 4061–4067 (2010).

6. I. Jarmoskaite, I. AlSadhan, P. P. Vaidyanathan, D. Herschlag, How to measure and evaluate binding affinities. *eLife* **9**, e57264 (2020).

7. A. Hromić-Jahjefendić, H. Krnjić, V. N. Uversky, "Effects of Molecular Crowding on the Structural, Dynamic, and Functional Properties of Biological Macromolecules: A General Overview" in *(Macro)Molecular Crowding: Life of the Pottage*, V. N. Uversky, Ed. (Springer Nature Switzerland, 2025), pp. 15–39.

8. K. B. Frederick, D. Sept, E. M. De La Cruz, Effects of solution crowding on actin polymerization reveal the energetic basis for nucleotide-dependent filament stability. *J. Mol. Biol.* **378**, 540–550 (2008).

9. F. K. M. Schur, *et al.*, An atomic model of HIV-1 capsid-SP1 reveals structures regulating assembly and maturation. *Science* **353**, 506–508 (2016).

10. J. Hutchings, V. Stancheva, E. A. Miller, G. Zanetti, Subtomogram averaging of COPII assemblies reveals how coat organization dictates membrane shape. *Nat. Commun.* **9**, 4154 (2018).

11. T. A. M. Bharat, *et al.*, Structure of the hexagonal surface layer on Caulobacter crescentus cells. *Nat. Microbiol.* **2**, 17059 (2017).

12. J. A. G. Briggs, Structural biology in situ--the potential of subtomogram averaging. *Curr. Opin. Struct. Biol.* **23**, 261–267 (2013).

13. W. Wan, J. A. G. Briggs, "Cryo-Electron Tomography and Subtomogram Averaging" in *Methods in Enzymology*, (Elsevier, 2016), pp. 329–367.

14. G. S. Espie, M. S. Kimber, Carboxysomes: cyanobacterial RubisCO comes in small packages. *Photosynth. Res.* **109**, 7–20 (2011).

15. B. D. Rae, B. M. Long, M. R. Badger, G. D. Price, Functions, Compositions, and Evolution of the Two Types of Carboxysomes: Polyhedral Microcompartments That Facilitate CO2 Fixation in Cyanobacteria and Some Proteobacteria. *Microbiol. Mol. Biol. Rev.* **77**, 357–379 (2013).

16. J. M. Shively, F. Ball, D. H. Brown, R. E. Saunders, Functional Organelles in Prokaryotes: Polyhedral Inclusions (Carboxysomes) of Thiobacillus neapolitanus. *Science* **182**, 584–586 (1973).

17. G. C. Cannon, *et al.*, Microcompartments in Prokaryotes: Carboxysomes and Related Polyhedra. *Appl. Environ. Microbiol.* **67**, 5351–5361 (2001).

18. J. M. Shively, G. L. Decker, J. W. Greenawalt, Comparative ultrastructure of the thiobacilli. *J. Bacteriol.* **101**, 618–627 (1970).

19. Z. Dou, *et al.*, CO2 fixation kinetics of Halothiobacillus neapolitanus mutant carboxysomes lacking carbonic anhydrase suggest the shell acts as a diffusional barrier for CO2. *J. Biol. Chem.* **283**, 10377–10384 (2008).

20. L. Reinhold, R. Kosloff, A. Kaplan, A model for inorganic carbon fluxes and photosynthesis in cyanobacterial carboxysomes. *Can. J. Bot.* **69**, 984–988 (1991).

21. L. M. Oltrogge, *et al.*, Multivalent interactions between CsoS2 and Rubisco mediate α-carboxysome formation. *Nat. Struct. Mol. Biol.* **27**, 281–287 (2020).

22. F. Cai, *et al.*, Advances in Understanding Carboxysome Assembly in Prochlorococcus and Synechococcus Implicate CsoS2 as a Critical Component. *Life Basel Switz.* **5**, 1141–1171 (2015).

23. G. C. Cannon, *et al.*, Organization of Carboxysome Genes in the Thiobacilli. *Curr. Microbiol.* **46**, 0115–0119 (2003).

24. N. M. Mangan, A. Flamholz, R. D. Hood, R. Milo, D. F. Savage, pH determines the energetic efficiency of the cyanobacterial CO2 concentrating mechanism. *Proc. Natl. Acad. Sci.* **113**, E5354–E5362 (2016).

25. H. Wang, *et al.*, Rubisco condensate formation by CcmM in β-carboxysome biogenesis. *Nature* **566**, 131–135 (2019).

26. L. A. Metskas, *et al.*, Rubisco forms a lattice inside alpha-carboxysomes. *Nat. Commun.* **13**, 4863 (2022).

27. T. Ni, *et al.*, Structure and assembly of cargo Rubisco in two native α-carboxysomes. *Nat. Commun.* **13**, 4299 (2022).

28. N. C. Hill, J. W. Tay, S. Altus, D. M. Bortz, J. C. Cameron, Life cycle of a cyanobacterial carboxysome. *Sci. Adv.* **6**, eaba1269 (2020).

29. C. A. Huffine, C. Fontana, A. Avramov, C. Sempeck, J. C. Cameron, Cyanobacteria form a procarboxysome-like structure in response to high CO2. [Preprint] (2024). Available at: https://www.biorxiv.org/content/10.1101/2024.06.28.601118v1 [Accessed 26 November 2024].

30. L. A. Metskas, R. Wilfong, G. J. Jensen, Subtomogram averaging for biophysical analysis and supramolecular context. *J. Struct. Biol. X* **6**, 100076 (2022).

31. F. Lee, *et al.*, A toolkit to characterize protein polymerization from cryo-electron tomography data. [Preprint] (2025). Available at: https://www.biorxiv.org/content/10.1101/2025.04.22.650024v1 [Accessed 3 December 2025].

32. W. Cao, T. E. Sladewski, A. T. Heaslip, E. M. De La Cruz, Bending stiffness of Toxoplasma gondii actin filaments. *J. Biol. Chem.* **301**, 108101 (2025).

33. A. S. Jermyn, W. Cao, W. A. Elam, E. M. De La Cruz, M. M. Lin, Directional allosteric regulation of protein filament length. *Phys. Rev. E* **101**, 032409 (2020).

34. L. Edelstein-Keshet, G. B. Ermentrout, Models for the length distributions of actin filaments: I. Simple polymerization and fragmentation. *Bull. Math. Biol.* **60**, 449–475 (1998).

35. F. Oosawa, *Thermodynamics of the polymerization of protein* (London ; New York : Academic Press, 1975).

36. "Cryo-Electron Tomography and Subtomogram Averaging" in *Methods in Enzymology*, (Academic Press, 2016), pp. 329–367.

37. W. Zhou, *et al.*, Cryo-EM structure-based selection of computed ligand poses enables design of MTA-synergic PRMT5 inhibitors of better potency. *Commun. Biol.* **5**, 1–15 (2022).

38. J. H. Van Drie, L. Tong, Cryo-EM as a powerful tool for drug discovery. *Bioorg. Med. Chem. Lett.* **30**, 127524 (2020).

39. Y. Sun, *et al.*, Decoding the Absolute Stoichiometric Composition and Structural Plasticity of α-Carboxysomes. *mBio* **13**, e0362921 (2022).

40. L. Edelstein-Keshet, G. B. Ermentrout, Models for the length distributions of actin filaments: I. Simple polymerization and fragmentation. *Bull. Math. Biol.* **60**, 449–475 (1998).

41. W. Dai, *et al.*, Visualizing Individual RuBisCO and Its Assembly into Carboxysomes in Marine Cyanobacteria by Cryo-Electron Tomography. *J. Mol. Biol.* **430**, 4156–4167 (2018).

42. T. Ni, *et al.*, Intrinsically disordered CsoS2 acts as a general molecular thread for α-carboxysome shell assembly. *Nat. Commun.* **14**, 5512 (2023).

43. L. M. Oltrogge, A. W. Chen, T. Chaijarasphong, J. B. Turnšek, D. F. Savage, α-Carboxysome Size Is Controlled by the Disordered Scaffold Protein CsoS2. *Biochemistry* **63**, 219–229 (2023).

44. R.-Q. Zhou, *et al.*, Structure and assembly of the α-carboxysome in the marine cyanobacterium Prochlorococcus. *Nat. Plants* 1–12 (2024). https://doi.org/10.1038/s41477-024-01660-9.

45. A. C. Schramm, *et al.*, Actin Filament Strain Promotes Severing and Cofilin Dissociation. *Biophys. J.* **112**, 2624–2633 (2017).

46. A. C. Schramm, G. M. Hocky, G. A. Voth, J.-L. Martiel, E. M. D. L. Cruz, Plastic Deformation and Fragmentation of Strained Actin Filaments. *Biophys. J.* **117**, 453–463 (2019).

47. J. P. Bibeau, *et al.*, Twist response of actin filaments. *Proc. Natl. Acad. Sci.* **120**, e2208536120 (2023).

48. W. B. Carpenter, *et al.*, Ratiometric Sensing of Redox Environments Inside Individual Carboxysomes Trapped in Solution. *J. Phys. Chem. Lett.* **13**, 4455–4462 (2022).

49. F. Oosawa, S. Asakura, *Thermodynamics of the Polymerization of Protein* (Academic Press, 1975).

50. J. Howard, *Mechanics of motor proteins and the cytoskeleton* (Sinauer Associates, 2001).

51. L. Edelstein-Keshet, G. B. Ermentrout, Models for the length distributions of actin filaments: I. Simple polymerization and fragmentation. *Bull. Math. Biol.* **60**, 449–475 (1998).

52. M. M. J. Smulders, *et al.*, How to distinguish isodesmic from cooperative supramolecular polymerisation. *Chem. Weinh. Bergstr. Ger.* **16**, 362–367 (2010).

53. K. M. Skillman, *et al.*, The unusual dynamics of parasite actin result from isodesmic polymerization. *Nat. Commun.* **4**, 2285 (2013).

54. L. Edelstein-Keshet, G. B. Ermentrout, Models for the length distributions of actin filaments: I. Simple polymerization and fragmentation. *Bull. Math. Biol.* **60**, 449–475 (1998).

55. E. Andrianantoandro, L. Blanchoin, D. Sept, J. A. McCammon, T. D. Pollard, Kinetic mechanism of end-to-end annealing of actin filaments1. *J. Mol. Biol.* **312**, 721–730 (2001).

56. A. I. Flamholz, *et al.*, Functional reconstitution of a bacterial CO2 concentrating mechanism in Escherichia coli. *eLife* **9**, e59882 (2020).

57. D. S. Trettel, W. Resager, B. M. Ueberheide, C. C. Jenkins, W. C. Winkler, Chemical probing provides insight into the native assembly state of a bacterial microcompartment. *Structure* **30**, 537-550.e5 (2022).

58. B. Turoňová, F. K. M. Schur, W. Wan, J. A. G. Briggs, Efficient 3D-CTF correction for cryo-electron tomography using NovaCTF improves subtomogram averaging resolution to 3.4Å. *J. Struct. Biol.* **199**, 187–195 (2017).

59. A. Rohou, N. Grigorieff, CTFFIND4: Fast and accurate defocus estimation from electron micrographs. *J. Struct. Biol.* **192**, 216–221 (2015).

60. D. N. Mastronarde, Dual-axis tomography: an approach with alignment methods that preserve resolution. *J. Struct. Biol.* **120**, 343–352 (1997).

61. T. Grant, N. Grigorieff, Measuring the optimal exposure for single particle cryo-EM using a 2.6 Å reconstruction of rotavirus VP6. *eLife* **4**, e06980 (2015).

62. D. Castaño-Díez, M. Kudryashev, M. Arheit, H. Stahlberg, Dynamo: a flexible, user-friendly development tool for subtomogram averaging of cryo-EM data in high-performance computing environments. *J. Struct. Biol.* **178**, 139–151 (2012).

63. S. H. Scheres, RELION: implementation of a Bayesian approach to cryo-EM structure determination. *J. Struct. Biol.* **180**, 519–530 (2012).

64. T. D. Goddard, C. C. Huang, T. E. Ferrin, Visualizing density maps with UCSF Chimera. *J Struct Biol* **157**, 281–7 (2007).

65. W. H. Press, Ed., *Numerical recipes: the art of scientific computing*, 3rd ed (Cambridge University Press, 2007).

66. D. Voet, J. G. Voet, *Biochemistry*, 4th Ed. (New York : J. Wiley & Sons, 2004).