S. cerevisiae Cwc15p Tunes the Spliceosome Active Site for 5' Splice Site Cleavage

Natalie J. Zeps[1], Giovanni Balice[1], Zach Freedman[1], Sierra Jones[2], Dennis Halterman[2,3], and Aaron A. Hoskins[1,4*]

[1]Department of Biochemistry, University of Wisconsin-Madison, Madison, WI 53706

[2]Department of Plant Pathology, University of Wisconsin-Madison, Madison, WI 53706

[3]USDA-ARS Vegetable Crops Research Unit, Madison, WI 53706

[4]Department of Chemistry, University of Wisconsin-Madison, Madison, WI 53706

*CORRESPONDING AUTHOR:

Aaron A. Hoskins, ahoskins@wisc.edu

ORCID:

Natalie J. Zeps, 0009-0001-0625-404X

Dennis Halterman, 0000-0002-2428-9393

Aaron A. Hoskins, 0000-0002-9777-519X

Running Title: Roles of Cwc15p in Pre-mRNA Splicing

KEYWORDS: pre-mRNA, splicing, yeast, spliceosome, RNA, *S. cerevisiae*

ABSTRACT (250 words)

Pre-mRNA splicing is an essential step in eukaryotic gene expression during which spliceosomes remove introns from nascent RNAs while ligating the adjacent exons. Spliceosomes are cellular nanomachines composed of five small nuclear (snRNA) components and dozens of proteins, most of which are highly conserved. Despite the high conservation of many splicing factors between *S. cerevisiae* and *H. sapiens*, several protein components of the *S. cerevisiae* spliceosome are not essential for growth under normal laboratory conditions. This is particularly surprising for nonessential factors whose conserved domains contact the spliceosome's catalytic core. Uncovering a function for these splicing factors can be challenging since they are not required for viability, may engage in functionally redundant interactions, and may display only weak phenotypes in the absence of secondary mutations in other spliceosome components. One such nonessential factor is the Cwc15 protein. Cwc15's highly conserved N-terminus directly contacts the U2/U6 di-snRNA within the spliceosome catalytic core; yet its precise role in splicing has not been defined in any organism. In this work, we use molecular genetics in *S. cerevisiae* combined with splicing reporter assays to study Cwc15p function. We propose that Cwc15p not only promotes active site stability during 5' splice site cleavage but also impacts structural transitions into and out of this spliceosome conformation. This function may be critical for splicing in *S. cerevisiae* under nonoptimal conditions, facilitating use of weak or alternate splice sites, and could have implications for proofreading of spliceosome active site formation.

 Graphical Abstract

Cwc15p Fine-Tunes the Spliceosome Active Site

 Created in BioRender. Hoskins, A. (2026) https://BioRender.com/9nrbm6r

Article Summary (100 word)

Pre-mRNA splicing is carried out by large macromolecular machines called

spliceosomes which are composed of several snRNAs and dozens of proteins. Despite

decades of study, the functions of many splicing factors such as *S. cerevisiae* Cwc15p

remain unknown. Cwc15p is highly conserved among eukaryotes and directly contacts

the spliceosome catalytic core. Here, we have used genetic and splicing reporter assays

to study the function of Cwc15p during splicing *in vivo*. We propose that Cwc15p both

stabilizes the spliceosome active site during 5' splice site cleavage and impacts

remodeling of that site.

INTRODUCTION

In eukaryotes, nuclear DNA genes frequently contain introns which must be removed from precursor mRNA (pre-mRNA) transcripts concomitant with ligation of the flanking exons for gene expression (**Fig. 1A**). This process, called pre-mRNA splicing, is carried out by a large macromolecular machine called the spliceosome. Spliceosomes assemble around introns from uridine-rich small nuclear RNAs (the U snRNAs; U1, U2, U4, U5, and U6) and dozens of protein components. The snRNAs and many proteins associate with introns in pre-assembled complexes called small nuclear ribonucleoproteins (snRNPs). Other proteins also join spliceosomes in either multi-subunit complexes lacking RNA components [such as the Prp19-containing complex or Nineteen Complex (NTC)] or as individual units. The processes of spliceosome assembly and correct identification of introns are complex and involve numerous ATP-dependent and -independent stages (Plaschka et al. 2019; Kastner et al. 2019).

Once assembled, spliceosomes contain RNA active sites composed of the U2 and U6 snRNAs, the pre-mRNA substrate, and metal ions (Fica et al. 2013; Senn et al. 2024; Yan et al. 2019). Despite being ribozymes at their catalytic heart, the RNAs require numerous proteins to scaffold and arrange them into the catalytic conformations needed for 5' splice site (5'SS) cleavage (1st chemical step of splicing) and exon ligation (2nd step). While some of these proteins, such as Prp8p (Grainger and Beggs 2005), are essential components of the splicing machinery, a number of other protein splicing factors are not required (or at least not under certain conditions). In particular, the NTC and its related complexes contain many proteins that are non-essential in *Saccharomyces cerevisiae* (yeast) for growth and pre-mRNA splicing under laboratory conditions (Hogg et al. 2010).

For several of these proteins, their non-essentiality is surprising given their locations near

the spliceosome active site and contacts with critical protein and RNA components in

cryo-EM structures. Network analysis of these structures suggested that non-essentiality

may be due to redundancy in splicing factor interactions—non-essential factors make

fewer unique interactions compared with essential factors (Kaur et al. 2022). This implies

that they primarily function in *S. cerevisiae* to fine-tune splicing reactions under certain

conditions or on certain substrates.

One nonessential component is yeast Cwc15p. Cwc15p is a small, 20 kDa protein

that was identified as complexed with Cef1 (or cdc5 in *Schizosaccharomyces pombe*) by

mass spectrometry of purified splicing factor complexes (Ohi et al. 2002). Despite being

non-essential in *S. cerevisiae*, Cwc15p is highly conserved with homologs present

throughout eukaryotes including in humans and plants (Ohi et al. 2002) (**Fig. S1**). Cwc15p

homologs are essential in *S. pombe* (Ohi et al. 2002), *Arabidopsis* (Slane et al. 2020),

and *Bos taurus* (Sonstegard et al. 2013). This suggests that organisms with more complex

splicing regulatory mechanisms, intron composition, and/or machineries than *S.*

cerevisiae (Sales-Lee et al. 2021) require Cwc15p for proper spliceosome function.

No detailed biochemical or molecular genetics work has been reported for analysis

of Cwc15p's function during pre-mRNA splicing. Cryo-EM structures of catalytic-stage

spliceosomes from a variety of organisms all show that Cwc15p lacks appreciable

secondary structure and threads its way throughout the spliceosome (**Fig. 1B**). Cwc15p's

highly conserved N-terminus (**Fig. S1**) inserts directly into the catalytic core of the

spliceosome, contacting the back of the catalytic U2/U6 snRNA duplex opposite of the

face on which catalytically essential metal ions are bound (**Fig. 1B**, boxed region). Large

portions of the remainder of Cwc15p are unresolved in available cryo-EM structures;

however, a tryptophan in a conserved region near the C-terminus (**Fig. S1**) is visible and

interacts with a hydrophobic pocket on Prp8p (the large protein responsible for scaffolding

the catalytic RNAs). Given these interactions with critical components of the spliceosome,

it is likely that Cwc15p influences pre-mRNA splicing despite being non-essential in *S.*

cerevisiae. In support of this, *S. cerevisiae cwc15Δ* is synthetically lethal with the *prp19-*

1 allele (Ohi et al. 2002), and in *S. pombe*, the *fsm32* mutant allele of *CWF15* (the *S.*

pombe homolog of *CWC15*) results in intron retention and splicing defects of the MCS2

transcript (Hoyos-Manchado et al. 2017). Additional evidence comes from studies of plant

Cwc15 proteins. The *Arabidopsis cwc15-1* allele causes splicing defects *in vivo*, although

these were not extensively characterized (Slane et al. 2020), and *S. tuberosum* (potato)

Cwc15p was recently found to be the target of the pathogen *Phytophthora infestans*

effector protein IPI-O1 (Sun et al. 2024). This interaction results in a change in splicing of

an immunity gene in potato, resulting in pathogen resistance.

Here, we use yeast molecular genetics to probe the function of Cwc15p during pre-

mRNA splicing. By testing genetic interactions between *cwc15Δ* and splicing factors and

using *in vivo* splicing reporter assays, we propose that Cwc15p stabilizes the active site

during 1st step catalysis while also facilitating its remodeling before and after this step.

MATERIALS AND METHODS:

<u>Yeast strain and plasmid construction</u>

The *CWC15* gene was deleted from yeast strains by replacement with a

hygromycin resistance cassette (HygR) through homologous recombination (Goldstein

and McCusker 1999). Gene deletions were confirmed by PCR.

Mutant *CWC15* and *SNR6* plasmids were constructed either by site-directed

mutagenesis using inverse PCR (Silva et al. 2023) or by purchasing plasmids containing

the appropriate DNAs from GENEWIZ (South Plainfield, NJ USA) or Genscript

(Pennington, NJ USA). All plasmids were sequenced before use by Sanger or whole-

plasmid next-generation DNA sequencing (Functional Biosciences; Madison, WI USA).

Yeast growth and transformation were carried out using standard protocols and media

(Guthrie et al. 1991). Yeast strains and plasmids are listed in **Sup. Tables S1** and **S2**,

respectively.

<u>Yeast growth assays at various temperatures</u>

Yeast cultures were grown overnight in liquid culture to saturation and then

adjusted to an OD_{600} of 0.5 in 10% (v/v) glycerol. Three consecutive 1:10 serial dilutions

were performed by transferring 100 µL of culture into 900 µL of diluent, resulting in final

dilutions of 1:1, 1:10, 1:100, and 1:1000. Diluted cultures were then spotted onto plates

and incubated at 16°C, 23°C, 30°C, or 37°C before imaging with an ImageQuant™ LAS

4000 imaging system (GE Healthcare/Cytiva). Assays were performed in at least

triplicate.

<u>ACT1-CUP1 assays</u>

Yeast strains transformed with ACT1–CUP1 reporter plasmids were cultured in -

Leu dropout (DO) media. Diluted cultures (OD_{600} = 0.5) were spotted onto -Leu DO plates

supplemented with 0–2.5 mM $CuSO_4$ (Lesser and Guthrie 1993; Carrocci et al. 2018) and

incubated at 30°C for 3 days before imaging.

RESULTS

<u>Genetic Interactions between *CWC15* and Spliceosomal ATPases</u>

To assay Cwc15p function, we first deleted the *CWC15* gene from *S. cerevisiae*

and studied yeast growth at various temperatures (**Fig. 1C**). Consistent with a recent

report by Henke-Schulz and colleagues (Henke-Schulz et al. 2024), we observed that

cwc15Δ yeast had no observable growth defect at 16, 23, or 30°C on YPD plates but grew

more slowly and were heat sensitive (temperature sensitive or *ts*) at 37°C. (We note that

the strength of the *ts* phenotype is strain dependent with some strains growing more

robustly at 37°C, *c.f.* **Figs. 1C** vs. **5E**, for example).

Given Cwc15p's location proximal to the spliceosome active site, we hypothesized

that genetic interactions may exist between *cwc15Δ* and the DEAH-box ATPases

responsible for remodeling the spliceosome during formation or disruption of the catalytic

center. We tested genetic interactions between *cwc15Δ* and *PRP2*, *PRP16*, and *PRP22*

(**Fig. 2A**). Prp2p is responsible for releasing U2 snRNP proteins from the branch site

(BS)/U2 snRNA duplex and proofreading the catalytic core during the final stages of active

site assembly (Schmitzova et al. 2023; Lardelli et al. 2010; Wlodaver and Staley 2014).

Prp16p is responsible for releasing proteins from the spliceosome required for 5'SS

cleavage in order to facilitate active site reorganization and for proofreading the 1st step

(Koodathingal et al. 2010; Chung et al. 2023; Tseng et al. 2011). Prp22p is responsible

for proofreading exon ligation as well as releasing the mRNA product and proteins needed

during the 2nd step (Mayas et al. 2006; Company et al. 1991; Chung et al. 2025).

The Prp2p^{Q548N} mutation is cold sensitive (*cs*), inhibits splicing *in vivo*, and likely

inhibits ATP binding or hydrolysis by the enzyme (Wlodaver and Staley 2014). Splicing

inhibition arises due to difficulties in disruption of U2/U6 di-snRNA interactions needed for

the final stages of catalytic site folding. Destabilizing mutations in these snRNAs or other

splicing factors involved in activation can suppress the *cs* phenotype while stabilizing

mutations can do the opposite (Wlodaver and Staley 2014; Carrocci et al. 2017; van der

Feltz et al. 2021). When we combined Prp2p^{Q548N} with *cwc15Δ*, we observed that yeast

were synthetic sick at 30°C and synthetic lethal at 16, 23, and 37°C (**Fig. 2B**). This

indicates that Cwc15p functions during spliceosome active site formation and that loss of

Cwc15p is detrimental if aberrant Prp2p activity is present. This phenotype could arise

due to Cwc15p functioning to destabilize the U2/U6 di-snRNA during activation and/or

stabilizing the product of Prp2p remodeling (*i.e.,* the 1st-step active site of the

spliceosome). We favor the latter mechanism because it may provide a function for the

highly conserved Cwc15p N-terminus (see Discussion).

We then tested genetic interactions with mutants in Prp16p and Prp22p.

Prp16p^{R686I} is *cs* at 16°C and *ts* at 37°C, and at the latter temperature can be suppressed

by the *prp8-156* allele which favors the exon ligation conformation of the active site (Query

and Konarska 2004; Hotz and Schwer 1998). Similarly, the *cs* phenotype can be

suppressed by *ecm2Δ*, which eliminates a nonessential splicing factor that makes

stabilizing interactions in the 5'SS conformation (van der Feltz et al. 2021). Prp22p^{T637A}

decouples ATP hydrolysis from RNA unwinding and causes a *cs* phenotype *in vivo*

(Schwer and Meszaros 2000). Prp22p^{T637A} is also synthetic lethal at certain temperatures

with loss of the 2nd-step splicing factor Fyv6p or alleles of *CEF1* that disrupt exon ligation

(Lipinski et al. 2023; Query and Konarska 2012). Combined, these observations support

a model in which Prp16p and Prp22p promote exit from the 1st and 2nd-step conformations

of the active site, respectively; Prp16p^{R686I} and Prp22p^{T637A} inhibit these transitions; and

that the effects of these mutations can be suppressed (or exacerbated) by changes in the

relative stabilities of the active site conformations (Liu et al. 2007).

Genetic interactions were less pronounced with $Prp16p^{R686I}$ and $Prp22p^{T637A}$

mutants relative to $Prp2p^{Q548N}$. Loss of Cwc15p in cells expressing $Prp16p^{R686I}$ caused a

slight suppression of the *cs* phenotype; although, $Prp16p^{R686I}$/*cwc15Δ* cells still grew

poorly at 16°C (**Fig. 2B**). At 37°C, Prp16p wild type ($Prp16p^{WT}$)/*cwc15Δ* also grew poorly

while $Prp16p^{R686I}$/*cwc15Δ* cells did not grow. Similarly, the *cwc15Δ ts* phenotype was

exacerbated by $Prp22p^{T637A}$ (**Fig. 2B**). However, the *cs* phenotype due to $Prp22p^{T637A}$ was

not suppressed by *cwc15Δ*. As with Prp2p, these results with Prp16p and Prp22p are

consistent with Cwc15p impacting spliceosome active site stability and in ways that

change the ATPase-dependence of transitions between conformations or formation of

transient, intermediate structures (Eysmont et al. 2019).

<u>Genetic Interactions Between *CWC15, PRP8, and SNR6*</u>

As mentioned above, different alleles or mutants of yeast Prp8p have been shown

to influence pre-mRNA splicing by promoting the formation or stability of the active site

for a given step at the expense of the other (*e.g.,* promoting the 1st step reaction but

inhibiting the 2nd step, **Fig. 3A**) (Liu et al. 2007). Given our results with the ATPases (**Fig.**

2), we next tested genetic interactions between *PRP8* 1st- or 2nd-step alleles and *cwc15Δ*.

We expressed Prp8p mutants ($Prp8p^{P986T}$, 2nd step allele (*prp8-161*); $Prp8p^{R1753K}$,

1st-step allele; and $Prp8p^{E1960K}$, 1st-step allele (*prp8-101*)) (Liu et al. 2007) in the presence

of *CWC15* or in a deletion background (**Fig. 3B**). Deletion of *CWC15* was able to suppress

the *ts* phenotype at 37°C for both 1st-step alleles ($Prp8p^{R1753K}$ and $Prp8p^{E1960K}$). In

addition, yeast containing the 2nd-step allele $Prp8p^{P986T}$ grew slightly better in the absence

of Cwc15p than in its presence at 37°C. Loss of *CWC15* was also able to suppress the

cs phenotype of the Prp8p^{E1960K} mutant at 16°C. When 1st- and 2nd-step alleles are

combined in the same *PRP8* gene, they can cancel each other's effects (Liu et al. 2007).

While the addition of the Prp8p^{P986T} mutation restored growth at 37°C for Prp8p^{R1753K} in

the presence of Cwc15p, this effect was lost and yeast became sicker at all temperatures

when *CWC15* was deleted.

These results are consistent with a role for Cwc15p in stabilizing the 1st step

conformation of the spliceosome active site at a stage that acts in concert with Prp8p 1st-

step alleles. When the active site is destabilized due to loss of Cwc15p, the impact of

PRP8 1st-step alleles is weakened and growth is restored at 37°C. Cwc15p could also

have some additional function during the 2nd-step or at other stages given the weaker

effects seen with the Prp8^{P986T} mutant and the synthetic *cs* phenotype observed with

Prp8p$^{P986T+R1753K}$ in *cwc15Δ* cells.

Since mutations in the U6 snRNA gene (*SNR6*) can also change the relative

efficiencies of the 1st and 2nd steps splicing, we then tested whether U6 mutants that favor

(U57A, U70A, A79U) or disfavor (U57C, G63C, U65A) the transition between the 1st and

2nd step also have genetic interactions with *cwc15Δ* (Liu et al. 2007; Query and Konarska

2004; Eysmont et al. 2019). These mutations also span different regions of the U6 snRNA

including the upper and lower telestem (U70A, A79U and G63C, U65A; respectively) and

the U2/U6 di-snRNA (U57A, U57C). The lower telestem is proposed to play a key role in

formation of a transitional spliceosome structure between active conformations (Eysmont

et al. 2019).

In this case, we see the strongest genetic interactions between U6-U57C and

cwc15Δ, which are synthetic lethal at 37°C and cause poor growth at 16°C relative to cells

with *CWC15* (**Fig. 3C**). U57C is thought to stabilize the 1st step conformation of the

spliceosome or disfavor transitions away from this state (Liu et al. 2007; Eysmont et al.

2019; Wlodaver and Staley 2014). We favor an interpretation that Cwc15p facilitates the

1st-to-2nd step transition and loss of Cwc15p in combination with U6-U57C makes this

transition more difficult. In combination with results from Prp8p and the ATPases, one

model for Cwc15p function could be that it both stabilizes the active site during 5'SS

cleavage and facilitates rearrangement of the active site at stages preceding or following

catalysis. The latter activity could occur by destabilization of the active site conformation

at those times and/or by impacting the formation or stability of structures that form during

remodeling.

Impact of *CWC15* Deletion on Yeast Growth with ACT1-CUP1 Splicing Reporters

To further assess Cwc15p's role in pre-mRNA splicing, we used the *in vivo* ACT1-

CUP1 splicing reporter assay (Lesser and Guthrie 1993). In this assay, copper-sensitive

yeast are transformed with a plasmid encoding a gene for the Cup1p copper resistance

protein with derivatives of the *ACT1* intron inserted into its coding sequence (**Fig. 4A**).

They are then grown on plates containing increasing concentrations of Cu^{2+}. Growth in

the presence of copper requires splicing of the *ACT1-CUP1* pre-mRNA and expression

of the Cup1p protein. The degree of resistance of the yeast to copper correlates with the

extent of splicing.

Using the WT ACT1-CUP1 reporter, which contains consensus yeast splice sites,

we observed no change in resistance to copper between WT and *cwc15Δ* strains and

both readily grew on plates containing the maximal copper concentration (2.5 mM) (**Fig.**

4B). This indicates a high splicing efficiency for the ACT1-CUP1 reporter even when

Cwc15p is absent under these conditions. This likely explains, at least in part, the lack of

a significant growth phenotype at 30°C for the deletion strain since the majority of yeast

introns contain consensus splice sites.

We next tested growth of yeast containing reporters with a variety of substitutions

at the 5' or 3'SS or BS sequence (**Fig. 4A**, **B**). In these cases, we saw changes in copper

tolerances for the WT *vs. cwc15Δ* strains for particular variants. For the 5'SS, we observed

slight increases in tolerance for G+1 substitutions (G1A, G1C) and a reduction in

tolerance for the U2C reporter. For substitutions at the branch site, we observed

reductions in tolerance for both the BS-C and BS-G substitutions as well as for a C260A

substitution at the 3' flanking position. We did not observe any changes in copper

tolerance for yeast with substitutions at the -3 (gAG) and -2 (UuG) positions of the 3'SS.

Overall, the results show that loss of *CWC15* impacts the 1st-step splicing reaction

and changes copper tolerances observed with reporters that are limiting for 5'SS

cleavage: G1A, G1C, BS-C, and C260A. In contrast, we did not observe changes in

tolerance for most reporters limiting for exon ligation: U2A, A3C, gAG, and UuG with the

exception of BS-G (Liu et al. 2007). Despite impacting growth due to the variant ACT1-

CUP1 reporters, the *cwc15Δ* allele does not behave identically to characterized Prp8 1st

or 2nd-step alleles since we observe changes at the G+1 position and do not observe

other characteristic changes in growth (*e.g.*, decreased copper tolerance for the A3C

reporter for 1st-step alleles and increases in copper tolerances for gAG and UuG reporters

for 2nd-step alleles) (Liu et al. 2007; Query and Konarska 2004).

<u>Genes Encoding Mutant Cwc15 Proteins Can Rescue Yeast Growth</u>

We next tested if conserved regions of Cwc15p were essential for rescuing the *ts*

phenotype (**Fig. 1C**). The most highly conserved region in Cwc15p is the N-terminus,

which inserts into the spliceosome active site between the U2 and U6 snRNAs and

contacts many key splicing factors (**Fig. 5A**, **B**). In a cryo-EM structure of the *S. cerevisiae*

spliced product spliceosome (P complex), Cwc15p residues Thr2 and Thr3 form hydrogen

bonding interactions with U6 snRNA nucleotides C84 and C85 (Senn et al. 2024). Arg6

also forms a cation-π stacking interaction with U2 snRNA G20, and Gly14 closely packs

into an interface with Prp46p. Additionally, conservation of Pro7 may be due to limiting

flexibility of the N-terminus and facilitating its threading through a small channel formed

between U2 snRNA G20/G21 and U6 snRNA A62/G63. We hypothesized that these

interactions could be disrupted using T2V/T3V substitutions which replace hydrophilic

with hydrophobic side chains that cannot hydrogen bond to U6 nucleotides, an

H5A/R6A/P7A/Q8A quadruple mutant (AAAA) that eliminates the cation-π interaction with

U2 and increases flexibility at position 7, or by introducing a side chain at the interface

with Prp46p with a G14A mutation.

In addition to the N-terminus, Cwc15 proteins also contain a conserved tryptophan

residue located in the C-terminal half. This tryptophan is flanked by charged amino acids

and fits into a hydrophobic pocket found on the Prp8 protein in the reverse transcriptase-

like domain (**Fig. 5C, D**). Therefore, we also made Cwc15p mutants in which the

tryptophan (Trp127) was substituted with a differently-shaped and charged residue

(glutamate or lysine), which would be less favorable to bury within this pocket.

We transformed a *cwc15Δ* yeast strain with TRP1/CEN plasmids expressing either

WT Cwc15p or mutants expressed from the endogenous promoter and selected for

growth on -trp plates. None of the mutants caused any growth defect at 30°C (not shown).

We then assayed growth at a higher temperature to test for rescue of the *ts* phenotype.

In these experiments, we found that the *ts* phenotype was more apparent at 39°C than

37°C. Yeast lacking Cwc15p and transformed with an empty pRS414 plasmid grew more

slowly than those containing a plasmid that expressed WT Cwc15p (**Fig. 5E**). This

indicates that a plasmid-borne copy of the WT *CWC15* gene can rescue the *ts* phenotype

under these conditions.

Surprisingly, all the mutants grew more quickly than the control containing the

empty plasmid and similarly to WT. This suggests that these conserved amino acids are

not essential for rescuing the *ts* phenotype. However, it was difficult to distinguish

additional differences between yeast expressing WT or any of the Cwc15p mutants. One

possibility is that the *ts* phenotype arises from loss of Cwc15p interactions besides those

shown in **Fig. 5.** Cwc15 has not been fully resolved in any spliceosome structure from

any organism—the structural environment of large portions of the protein are unknown.

For example, in the most highly resolved spliceosome structure from *S. cerevisiae*, 58%

of the Cwc15p peptide could not be modeled (Senn et al. 2024). The *ts* phenotype could

depend on spliceosomal interactions not yet structurally defined or Cwc15p's interactions

with other machineries such as the transcription and export (TREX) complex (Henke-

Schulz et al. 2024).

Genetic Interaction Between *CWC15* and Destabilizing U6 snRNA Mutants

Finally, we further tested the idea that Cwc15p stabilizes the active site by

introducing destabilizing mutations into the U6 snRNA in regions that contact the N-

terminus of Cwc15p (**Fig. 6A**). We introduced U6 A62U, G63U, C84G, or C85G mutations

since it was previously shown that these U6 variants have little effect on yeast growth by

themselves (McPheeters 1996). When the growth of *CWC15* and *cwc15Δ* yeast were

compared with these variants, we observed a synthetic sick interaction between *cwc15Δ*

and the U6 snRNA A62U mutation at 37°C. It is not obvious what structural explanation

could account for this. One possibility is that Cwc15p could prevent inappropriate wobble

pairing between U6 A62U and U2 nucleotides G20 or G21 in precatalytic or catalytic

spliceosomes (Senn et al. 2024; Galej et al. 2016; Rauhut et al. 2016). The conformation

of U2 G20 may be particularly important for proper active site formation (see **Sup. Fig.**

S2 and Discussion). For the remaining U6 snRNA mutations studied here, it is likely that

they were not as significantly destabilizing and lack observable Cwc15p-dependent

phenotypes in these assays.

DISCUSSION

In summary, our data support a model in which Cwc15p fine-tunes the spliceosome

active site during 5'SS cleavage and also impacts transitions between spliceosome

conformations. The notion that non-essential *S. cerevisiae* splicing factors like Cwc15p

perturb catalytic conformations of the spliceosome or enforce fidelity has previously been

discussed with respect to other components of the NTC and the Ecm2 and Fyv6 proteins

(Hogg et al. 2010; Senn et al. 2024; Lipinski et al. 2023; van der Feltz et al. 2021). Our

group has previously proposed that non-essentiality may be linked to the relatively low

number of unique connections proteins like Cwc15p make between essential factors and

built-in interaction redundancy in the spliceosome (Kaur et al. 2022). Despite this, many

non-essential splicing factors become critical at low or high temperatures or in certain

genetic backgrounds. This could indicate that their conservation is due to subtle

modulation of the splicing machinery or the need for additional structural stability under

non-optimal growth conditions.

Beyond *S. cerevisiae*, this modulation may be critical for regulating alternative

splicing and result in Cwc15 essentiality. One aspect of this model is potential adaptation

(or exploitation) of Cwc15 for changing splicing outcomes in response to cellular

conditions. For example, direct interaction between the *P. infestans* virulence protein IPI-

O1 and the potato host protein StCwc15 leads to relocalization of StCwc15 from the

nucleoplasm to the nucleolus and nuclear speckles (Sun et al. 2024). This interaction also

results in a change in splicing of the *RB* immunity gene pre-mRNA leading to a decrease

in the intron-retained isoform and an increase in the fully-spliced mRNA (Sun et al. 2024).

While the molecular mechanism for this switch has not yet been determined, other NTC

components are also subject to dynamic regulation including by post-translational

modification (Lu and Legerski 2007; Lleres et al. 2010; de Almeida and O'Keefe 2015).

This provides a plausible mechanism by which external factors could modulate Cwc15's

activity and, consequently, 5'SS cleavage efficiency.

Determining a more precise role for Cwc15p during splicing is challenging, and the

data are most consistent with the protein functioning in stabilizing the 1^{st}-step active site

during structural transitions as well as during catalysis. Yeast with Cwc15p show

increased tolerance to Cu^{2+} using ACT1-CUP1 splicing reporters harboring a 5'SS U2C

substitution and several BS substitutions (**Fig. 4**). Additionally, loss of Cwc15p

suppresses growth defects associated with Prp8p 1st-step alleles (**Fig. 3**) and results in a

growth defect in the presence of a likely destabilizing U6 snRNA mutation (**Fig. 6**). These

data not only suggest that Cwc15p stabilizes the 1st-step conformation of the spliceosome

but that this may also be particularly important when non-consensus substrates or mutant

splicing factors are present.

Among our results, genetic interactions between *cwc15Δ* and Prp2p^{Q5548N} and U6-

U57C mutations are particularly interesting. In both cases, the *CWC15* deletion has a

negative genetic interaction with these mutations. While interpretation of U6 U57C

mutations can be complex (Wlodaver and Staley 2014; McPheeters 1996; Query and

Konarska 2004; Liu et al. 2007; Eysmont et al. 2019), the data are consistent with Cwc15p

also facilitating structural transitions into and out of the 1st-step active site conformation.

How exactly this occurs is unclear but could involve directly destabilizing the RNA

structures that form the spliceosome active site, stabilizing transient active site

conformations, or favoring formation of the stable products of these transitions. The latter

possibility would provide a parsimonious explanation for some of our results: stabilization

of the 1st-step active site by Cwc15p facilitates both its formation by Prp2 and 5'SS

cleavage itself.

An intriguing possibility is that Cwc15's conserved N-terminus participates in the

above processes directly. Analysis of multiple cryo-EM structures of *S. cerevisiae*

spliceosomes reveals that the very N-terminus of Cwc15p is located in between the U2

and U6 snRNAs in the B* spliceosome (the stage immediately preceding 5'SS cleavage)

and in this same position in subsequently formed complexes (B*→C→C*→P→ILS

complex spliceosomes) (Senn et al. 2024; Wan et al. 2017; Fica et al. 2017; Wan et al.

2019; Wilkinson et al. 2021) (**Fig. S2**). Importantly, in activated spliceosomes (B^{act})

trapped prior to Prp2p activity and formation of B*, this same region of Cwc15p could not

be modeled between the snRNAs and does not adopt a specific, stable conformation (see

Fig. S2) (Bai et al. 2021). Further, the N-terminus of Cwc15 proteins could not be modeled

in structures of an off-pathway, B^{act}-like *S. cerevisiae* spliceosome assembled on an

excised intron (Li et al. 2026) (Max Wilkinson, personal communication); a human

spliceosome trapped while Prp2 was functioning (Schmitzova et al. 2023); or in *S. pombe*

and *Chaetomium thermophilum* spliceosomes trapped prior to the B* stage and in the

midst of being disrupted by the DHX35 ATPase (Soni et al. 2025; Li et al. 2025). Stable

insertion of the Cwc15 N-terminus in between the U2 and U6 snRNAs may be a

consequence of Prp2 activity and a marker of a correctly assembled active site.

Conversely, a dynamic Cwc15 N-terminus (or just lack of its insertion between U2 and

U6) may facilitate active site disruption and proofreading by DHX35. These putative

functions are evidently not essential in *S. cerevisiae* (see **Figs. 1, 5**), in which a

corresponding DHX35-like activity has also not yet been identified.

The involvement of Cwc15p in other steps in gene expression besides pre-mRNA

splicing complicates phenotypic analysis in our assays. In addition to splicing, Cwc15p is

also important for tethering the TREX complex to the NTC, which itself is involved in both

pre-mRNA splicing and transcription elongation (Henke-Schulz et al. 2024; Chanarat et

al. 2011). Further, loss of *CWC15* has a negative genetic interaction with deletion of the

general transcription factor TFIIS (*dst1*Δ) (Henke-Schulz et al. 2024). Given the high

degree of coupling between transcription and splicing (Carrocci and Neugebauer 2024),

it is likely that the growth phenotypes reported here have contributions from both

processes. Future deconvolution of Cwc15p function may benefit from both molecular

and genetic approaches to disentangle these processes as well as studies in model

organisms in which the protein is essential for viability.

DATA AND REAGENT AVAILABILITY

Strains and plasmids are available upon request. The authors affirm that all data

necessary for confirming the conclusions of the article are present within the article,

figures, and tables.

ACKNOWLEDGMENTS

We thank Joshua Paulson for help in preparation of microbial growth media. We

thank Laura Vanderploeg for help with figures. We thank Max Wilkinson (Memorial Sloan

Kettering Cancer Center) for helpful discussions about the structure of Cwc15 and for

making figures S2A and B.

AUTHOR CONTRIBUTIONS

NZ, DH, AAH: project conceptualization; NZ, ZF, AAH: media and reagent

preparation; NZ, GB, ZF, SJ, and AAH: data acquisition and analysis; DH, AAH: writing

and editing of the manuscript with input from NZ.

STUDY FUNDING

This work was supported by a grant from the USDA (HATCH WIS06060). Funding

for DH was provided by USDA-ARS CRIS 5090-21220-005-00D. Funding for SJ was

provided by the University of Wisconsin-Madison Summer Research Opportunities

Program.

CONFLICT OF INTEREST

AAH is carrying out sponsored research for Remix Therapeutics.

REFERENCES

Bai, R., R. Wan, C. Yan, Q. Jia, J. Lei *et al.*, 2021 Mechanism of spliceosome remodeling by the ATPase/helicase Prp2 and its coactivator Spp2. *Science* 371 (6525).

Carrocci, T.J., and K.M. Neugebauer, 2024 Emerging and re-emerging themes in co-transcriptional pre-mRNA splicing. *Mol Cell* 84 (19):3656–3666.

Carrocci, T.J., J.C. Paulson, and A.A. Hoskins, 2018 Functional analysis of Hsh155/SF3b1 interactions with the U2 snRNA/branch site duplex. *RNA* 24 (8):1028–1040.

Carrocci, T.J., D.M. Zoerner, J.C. Paulson, and A.A. Hoskins, 2017 SF3b1 mutations associated with myelodysplastic syndromes alter the fidelity of branchsite selection in yeast. *Nucleic Acids Res* 45 (8):4837–4852.

Chanarat, S., M. Seizl, and K. Strasser, 2011 The Prp19 complex is a novel transcription elongation factor required for TREX occupancy at transcribed genes. *Genes Dev* 25 (11):1147–1158.

Chung, C.S., C.K. Tseng, H.C. Chen, and S.C. Cheng, 2025 New mechanistic insights into Prp22-mediated exon ligation and mRNA release. *Nucleic Acids Res* 53 (16).

Chung, C.S., H.L. Wai, C.Y. Kao, and S.C. Cheng, 2023 An ATP-independent role for Prp16 in promoting aberrant splicing. *Nucleic Acids Res* 51 (20):10815–10828.

Company, M., J. Arenas, and J. Abelson, 1991 Requirement of the RNA helicase-like protein PRP22 for release of messenger RNA from spliceosomes. *Nature* 349 (6309):487–493.

de Almeida, R.A., and R.T. O'Keefe, 2015 The NineTeen Complex (NTC) and NTC-associated proteins as targets for spliceosomal ATPase action during pre-mRNA splicing. *RNA Biol* 12 (2):109–114.

Eysmont, K., K. Matylla-Kulinska, A. Jaskulska, M. Magnus, and M.M. Konarska, 2019 Rearrangements within the U6 snRNA Core during the Transition between the Two Catalytic Steps of Splicing. *Mol Cell* 75 (3):538–548 e533.

Fica, S.M., C. Oubridge, W.P. Galej, M.E. Wilkinson, X.C. Bai *et al.*, 2017 Structure of a spliceosome remodelled for exon ligation. *Nature* 542 (7641):377–380.

Fica, S.M., N. Tuttle, T. Novak, N.S. Li, J. Lu *et al.*, 2013 RNA catalyses nuclear pre-mRNA splicing. *Nature* 503 (7475):229–234.

Galej, W.P., M.E. Wilkinson, S.M. Fica, C. Oubridge, A.J. Newman *et al.*, 2016 Cryo-EM structure of the spliceosome immediately after branching. *Nature* 537 (7619):197–201.

Goldstein, A.L., and J.H. McCusker, 1999 Three new dominant drug resistance cassettes for gene disruption in Saccharomyces cerevisiae. *Yeast* 15 (14):1541–1553.

Grainger, R.J., and J.D. Beggs, 2005 Prp8 protein: at the heart of the spliceosome. *RNA* 11 (5):533–557.

Guthrie, C., G.R. Fink, J.N. Abelson, M.I. Simon, and ScienceDirect, 1991 *Guide to yeast genetics and molecular biology*. San Diego ; London: Academic Press.

Henke-Schulz, L., R. Minocha, N.H. Maier, and K. Strasser, 2024 The Prp19C/NTC subunit Syf2 and the Prp19C/NTC-associated protein Cwc15 function in TREX occupancy and transcription elongation. *RNA* 30 (7):854–865.

485   Hogg, R., J.C. McGrail, and R.T. O'Keefe, 2010 The function of the NineTeen Complex (NTC) in
486   regulating spliceosome conformations and fidelity during pre-mRNA splicing. *Biochem Soc Trans*
487   38 (4):1110–1115.

488   Hotz, H.R., and B. Schwer, 1998 Mutational analysis of the yeast DEAH-box splicing factor Prp16.
489   *Genetics* 149 (2):807–815.

490   Hoyos-Manchado, R., F. Reyes-Martin, C. Rallis, E. Gamero-Estevez, P. Rodriguez-Gomez *et al.*, 2017
491   RNA metabolism is the primary target of formamide in vivo. *Sci Rep* 7 (1):15895.

492   Kastner, B., C.L. Will, H. Stark, and R. Lührmann, 2019 Structural Insights into Nuclear pre-mRNA
493   Splicing in Higher Eukaryotes. *Cold Spring Harb Perspect Biol* 11 (11).

494   Kaur, H., C. van der Feltz, Y. Sun, and A.A. Hoskins, 2022 Network theory reveals principles of
495   spliceosome structure and dynamics. *Structure* 30 (1):190–200 e192.

496   Koodathingal, P., T. Novak, J.A. Piccirilli, and J.P. Staley, 2010 The DEAH box ATPases Prp16 and Prp43
497   cooperate to proofread 5' splice site cleavage during pre-mRNA splicing. *Mol Cell* 39 (3):385–
498   395.

499   Lardelli, R.M., J.X. Thompson, J.R. Yates, 3rd, and S.W. Stevens, 2010 Release of SF3 from the intron
500   branchpoint activates the first step of pre-mRNA splicing. *RNA* 16 (3):516–528.

501   Lesser, C.F., and C. Guthrie, 1993 Mutational analysis of pre-mRNA splicing in Saccharomyces cerevisiae
502   using a sensitive new reporter gene, CUP1. *Genetics* 133 (4):851–863.

503   Li, G.W., M.E. Wilkinson, and D.P. Bartel, 2026 The spliceosome assembles on excised linear introns to
504   protect them from degradation. *bioRxiv*:2026.2001.2021.700889.

505   Li, Y., P. Fischer, M. Wang, Q. Zhou, A. Song *et al.*, 2025 Structural insights into spliceosome fidelity:
506   DHX35-GPATCH1- mediated rejection of aberrant splicing substrates. *Cell Res* 35 (4):296–308.

507   Lipinski, K.A., K.A. Senn, N.J. Zeps, and A.A. Hoskins, 2023 Biochemical and genetic evidence supports
508   Fyv6 as a second-step splicing factor in Saccharomyces cerevisiae. *RNA* 29 (11):1792–1802.

509   Liu, L., C.C. Query, and M.M. Konarska, 2007 Opposing classes of prp8 alleles modulate the transition
510   between the catalytic steps of pre-mRNA splicing. *Nat Struct Mol Biol* 14 (6):519–526.

511   Lleres, D., M. Denegri, M. Biggiogera, P. Ajuh, and A.I. Lamond, 2010 Direct interaction between hnRNP-
512   M and CDC5L/PLRG1 proteins affects alternative splice site choice. *EMBO Rep* 11 (6):445–451.

513   Lu, X., and R.J. Legerski, 2007 The Prp19/Pso4 core complex undergoes ubiquitylation and structural
514   alterations in response to DNA damage. *Biochem Biophys Res Commun* 354 (4):968–974.

515   Mayas, R.M., H. Maita, and J.P. Staley, 2006 Exon ligation is proofread by the DExD/H-box ATPase
516   Prp22p. *Nat Struct Mol Biol* 13 (6):482–490.

517   McPheeters, D.S., 1996 Interactions of the yeast U6 RNA with the pre-mRNA branch site. *RNA* 2
518   (11):1110–1123.

519   Meng, E.C., T.D. Goddard, E.F. Pettersen, G.S. Couch, Z.J. Pearson *et al.*, 2023 UCSF ChimeraX: Tools
520   for structure building and analysis. *Protein Sci* 32 (11):e4792.

Ohi, M.D., A.J. Link, L. Ren, J.L. Jennings, W.H. McDonald *et al.*, 2002 Proteomics analysis reveals stable multiprotein complexes in both fission and budding yeasts containing Myb-related Cdc5p/Cef1p, novel pre-mRNA splicing factors, and snRNAs. *Mol Cell Biol* 22 (7):2011–2024.

Plaschka, C., A.J. Newman, and K. Nagai, 2019 Structural Basis of Nuclear pre-mRNA Splicing: Lessons from Yeast. *Cold Spring Harb Perspect Biol* 11 (5).

Query, C.C., and M.M. Konarska, 2004 Suppression of multiple substrate mutations by spliceosomal prp8 alleles suggests functional correlations with ribosomal ambiguity mutants. *Mol Cell* 14 (3):343–354.

Query, C.C., and M.M. Konarska, 2012 CEF1/CDC5 alleles modulate transitions between catalytic conformations of the spliceosome. *RNA* 18 (5):1001–1013.

Rauhut, R., P. Fabrizio, O. Dybkov, K. Hartmuth, V. Pena *et al.*, 2016 Molecular architecture of the Saccharomyces cerevisiae activated spliceosome. *Science* 353 (6306):1399–1405.

Sales-Lee, J., D.S. Perry, B.A. Bowser, J.K. Diedrich, B. Rao *et al.*, 2021 Coupling of spliceosome complexity to intron diversity. *Curr Biol* 31 (22):4898–4910 e4894.

Schmitzova, J., C. Cretu, C. Dienemann, H. Urlaub, and V. Pena, 2023 Structural basis of catalytic activation in human splicing. *Nature* 617 (7962):842–850.

Schwer, B., and T. Meszaros, 2000 RNA helicase dynamics in pre-mRNA splicing. *EMBO J* 19 (23):6582–6591.

Senn, K.A., K.A. Lipinski, N.J. Zeps, A.F. Griffin, M.E. Wilkinson *et al.*, 2024 Control of 3' splice site selection by the yeast splicing factor Fyv6. *Elife* 13.

Silva, D., G. Santos, M. Barroca, D. Costa, and T. Collins, 2023 Inverse PCR for Site-Directed Mutagenesis. *Methods Mol Biol* 2967:223–238.

Slane, D., C.H. Lee, M. Kolb, C. Dent, Y. Miao *et al.*, 2020 The integral spliceosomal component CWC15 is required for development in Arabidopsis. *Sci Rep* 10 (1):13336.

Soni, K., A. Horvath, O. Dybkov, M. Schwan, S. Trakansuebkul *et al.*, 2025 Structures of aberrant spliceosome intermediates on their way to disassembly. *Nat Struct Mol Biol* 32 (5):914–925.

Sonstegard, T.S., J.B. Cole, P.M. VanRaden, C.P. Van Tassell, D.J. Null *et al.*, 2013 Identification of a nonsense mutation in CWC15 associated with decreased reproductive efficiency in Jersey cattle. *PLoS One* 8 (1):e54872.

Sun, B., J. Huang, L. Kong, C. Gao, F. Zhao *et al.*, 2024 Alternative splicing of a potato disease resistance gene maintains homeostasis between growth and immunity. *Plant Cell* 36 (9):3729–3750.

Tseng, C.K., H.L. Liu, and S.C. Cheng, 2011 DEAH-box ATPase Prp16 has dual roles in remodeling of the spliceosome in catalytic steps. *RNA* 17 (1):145–154.

van der Feltz, C., B. Nikolai, C. Schneider, J.C. Paulson, X. Fu *et al.*, 2021 Saccharomyces cerevisiae Ecm2 modulates the catalytic steps of pre-mRNA splicing. *RNA* 27 (5):591–603.

Wan, R., R. Bai, C. Yan, J. Lei, and Y. Shi, 2019 Structures of the Catalytically Activated Yeast Spliceosome Reveal the Mechanism of Branching. *Cell* 177 (2):339–351 e313.

Wan, R., C. Yan, R. Bai, J. Lei, and Y. Shi, 2017 Structure of an Intron Lariat Spliceosome from
Saccharomyces cerevisiae. *Cell* 171 (1):120–132 e112.

Wilkinson, M.E., S.M. Fica, W.P. Galej, and K. Nagai, 2021 Structural basis for conformational equilibrium
of the catalytic spliceosome. *Mol Cell* 81 (7):1439–1452 e1439.

Wlodaver, A.M., and J.P. Staley, 2014 The DExD/H-box ATPase Prp2p destabilizes and proofreads the
catalytic RNA core of the spliceosome. *RNA* 20 (3):282–294.

Yan, C., R. Wan, and Y. Shi, 2019 Molecular Mechanisms of pre-mRNA Splicing through Structural
Biology of the Spliceosome. *Cold Spring Harb Perspect Biol* 11 (1).
567
568

FIGURES AND FIGURE LEGENDS

Figure 1. The transesterification steps in pre-mRNA splicing and location of
Cwc15p in the spliceosome. (**A**) The two transesterification steps in splicing involve
5'SS cleavage (1st step) and exon ligation (2nd step). (**B**) Location of Cwc15p in the *S.*
cerevisiae product (P) complex spliceosome (PDB 9DTR). This view shows the
spliceosome in grey cartoon representation with Cwc15p shown in spacefill and colored
with a gradient from blue (N-terminus) to red (C-terminus). Not all of the Cwc15p peptide
chain could be modeled in the structure. Prp8p and U6 snRNA are also shown in blue
and pink, respectively. A clipping mask was used to make the spliceosome active site and
Cwc15p more visible. This figure was generated using ChimeraX (Meng et al. 2023).
(**C**)Yeast without *CWC15* exhibit a *ts* phenotype at 37ºC. For panel C, yeast were grown
on YPD plates and imaged after the given number of days.
582

583
Figure 2. Genetic interactions between _CWC15_ and spliceosome DEAH-box ATPases. (**A**) Schematic showing spliceosome remodeling events promoted by Prp2p, Prp16p, and Prp22p which promote activation, the 1st-to-2nd step transition, and mRNA release, respectively. The mutations shown can inhibit these transitions and also impact pre-mRNA splicing fidelity. (**B**) Deletion of _CWC15_ causes a negative genetic interaction with Prp2p[Q548N]. Genetic interactions with Prp16p and Prp22p mutants were less apparent, although weak suppression of the Prp16p[R686I] could be observed at 16 and 23°C and Prp22p[T637A] strengthened the _ts_ phenotype caused by _cwc15Δ_ at 37°C. For panel B, yeast were grown on YPD plates and imaged after the given number of days.

Figure 3. Genetic interactions between *CWC15* and *PRP8* or *SNR6* (U6 snRNA). (A) 1st-step alleles (blue) of Prp8p or U6 snRNA promote 5'SS cleavage and the 1st-step configuration of the spliceosome at the expense of the 2nd step. 2nd-step alleles do the opposite (red). (**B**) Loss of *CWC15* can rescue the *ts* phenotype caused by Prp8 1st-step alleles. (**C**) Loss of *CWC15* is synthetic lethal with U6-U57C (1st-step allele) at 37°C and causes a *cs* phenotype at 16°C. For panels B and C, yeast were grown on YPD plates and imaged after the given number of days.

605
Figure 4. Loss of Cwc15p impacts Cu²⁺ tolerances of yeast expressing non-
consensus ACT1-CUP1 reporters. (**A**) Schematic of the ACT1-CUP1 reporter pre-
mRNA with the tested substitutions shown. (**B**) Loss of Cwc15p changes copper
tolerances for yeast expressing reporter pre-mRNAs with particular substitutions within
the 5'SS or BS but not for those expressing a reporter with consensus sites (WT). Bar
heights represent values obtained from a single replicate of the ACT1-CUP1 assay. Dots
represent values obtained for all three replicates. Yeast were grown at 30°C on -Leu DO
plates containing various [Cu²⁺] and imaged after 3 days.
614

615
Figure 5. Cwc15p mutants can complement a *cwc15Δ* growth defect. (**A**) Close-up
of Cwc15p N-terminus interactions with the U2/U6 di-snRNA within the active site of a P
complex spliceosome (PDB 9DTR). The active site can be identified by the presence of
Mg^{2+} and K$^+$ ions. (**B**) Sequence alignments and logos for Cwc15 from single and
multicellular model organisms and humans. The N-terminus of Cwc15 is highly conserved
among eukaryotes. (**C**) Close-up of the interaction of Cwc15p residue Trp127 with a
hydrophobic pocket on Prp8p (surface representation, hydrophobic surfaces shown in
yellow and hydrophilic in blue-green). (**D**) Trp127 is also highly conserved. (**E**) Growth
assays of *cwc15Δ* yeast expressing Cwc15p mutants relative to WT and empty plasmid
(pRS414) controls. Yeast were grown on -TRP DO plates and imaged after 2 days. Panels
A and C were created using ChimeraX (Meng et al. 2023). Sequence alignments and
logos for panels B and D were generated using MegAlign Pro (Lasergene, DNASTAR,
Inc.).

630
631 **Figure 6. Genetic interactions between potentially active site-destabilizing**
632 **mutations in the U6 snRNA and *CWC15*.** A negative genetic interaction is observed
633 between U6 snRNA A62U and *cwc15Δ* at 37oC. Yeast were grown on YPD plates and
634 imaged after 2 (30ºC), 3 (23 and 37ºC), or 6 days (16ºC). Panel A was created using
635 ChimeraX (Meng et al. 2023).

636
637
638