Tracing mobility among Eneolithic-Bronze Age Kurgan populations in the North Pontic steppe

Alexey G. Nikitin[1*], Virginie Renson[2], Svitlana Ivanova[3], Nadia C. Neff[4], Haruan Straioto[5], Sofiia Svyryd[1]

[1]Department of Biology, Grand Valley State University, Allendale, Michigan, United States of America

[2]Archaeometry Laboratory, University of Missouri Research Reactor, Columbia, MO, United States of America

[3]Institute of Archaeology, Ukrainian Academy of Sciences, Odesa, Ukraine

[4]Department of Anthropology, University of New Mexico, Albuquerque, NM, United States of America

[5]Museum of Archaeology and Ethnology, University of São Paulo, Brazil

*Corresponding author

E-mail: nikitin@gvsu.edu

Abstract

Five millennia ago, nomadic people from the North Pontic steppe left a profound impact on the

course of Eurasian prehistory. However, little is known about their mobility patterns within

their home region. To address this knowledge gap, we conducted a survey of the strontium

isotope landscape of people interred in the 4th-3rd millennium BCE burial mounds (kurgans) of

the western part of the North Pontic steppe. By analyzing the strontium signature in human

bone and dentin, we established strontium baseline values for the region. We subsequently

correlated enamel strontium ratios from 25 selected individuals with the baseline obtained and

with published strontium data across the North Pontic steppe. Enamel strontium ratios show

that some individuals interred in the northwest North Pontic fall within the regional baseline

range, whereas others overlap with values reported for the eastern North Pontic steppe. In

conjunction with carbon (δ^{13}C) and nitrogen (δ^{15}N) stable isotope data, we further determined

that some individuals interred in the western Pontic steppe either spent the later part of life in

the west Caspian steppe or were affected by physiological stress during lifetime. By integrating

our data with published isotopic datasets, we produced a first baseline heatmap of the North

Pontic steppe for the c. 4000-2000 BCE chronological period.

Introduction

In the Early Bronze Age (EBA, c. 3300-2200 BCE according to southeast European

chronology), a wave of innovations swept across the Eurasian continent, introducing new

linguistic features in the form of branches of Indo-European language, a developed pastoralist

economy, and a hierarchal social and ritual system. These changes were accompanied by a

change in genetic ancestry across Europe. The Anatolian Neolithic Farmer-based ancestry of

early farming communities of Europe was replaced by a genetic package founded on the

ancestry of Caucasus-Lower Volga migrants admixed with Neolithic North Pontic hunter-fisher-

foragers [1]. The nomadic people of the North Pontic region carrying this genetic "steppe"

package are referred to as the Kurgan people because they erected earth burial mounds

throughout the steppe and adjacent forest-steppe [2]. Although their genetic profiles are well

characterized, far less is understood about other aspects of their life history, including patterns

of mobility within their home range. We set out to address this knowledge deficiency by

conducting exploratory analysis of strontium isotope ratios from Eneolithic-EBA interments in

kurgans in the west portion of the Pontic steppe (the Odesa region of Ukraine) focusing on a set

of samples of high interpretive potential, to establish a foundation for future large-scale

isotopic studies of mobility in the region from which the Early Bronze Age "steppization" of

Europe was launched.

Strontium isotope analysis serves as a geochemical proxy for determining the

geographic origins of individuals buried in the ground, providing valuable insights into patterns

of mobility. When contextualized, this geochemical evidence can shed light on the social

significance of burial inclusion or exclusion. For instance, comparing $^{87}Sr/^{86}Sr$ values from dental

enamel with locally established baseline ranges can help distinguish between individuals from

local and non-local backgrounds. The presence of both local and non-local signatures among

the buried population suggests a community that embraced individuals from diverse

geographic origins. In this context, "local" denotes conformity to the average bioavailable

regional geographic zone, or, in our case, the isotopic compatibility with the regional

geochemical landscape within which the mobile steppe groups likely circulate. Conversely,

multiple non-local values may indicate a structured pattern of mobility.

The geological composition of the Odesa Region in Ukraine is a complex blend of

Precambrian crystalline basement rocks and younger sedimentary layers (Fig 1). The region,

part of the Black Sea Lowland, is influenced by the Black Sea Depression and the Scythian Plate.

The Odesa region is located near the Podolian (Buh-Dniester) domain, which, along with the

Azov domain, are the oldest components of the Ukrainian shield. The Cretaceous and

Quaternary sedimentary fluvio-colluvial deposits in the area mostly consist of loess and loess-

like loams. Pontian limestones from the Late Miocene (Upper Cenozoic) era are exposed in the

coastal areas of the region [3–6].

Fig 1. **Geological map of the north Pontic steppe region showing kurgans from which samples**

were obtained for the present study and the sites used in the comparative analysis. 1,

Mayaky; 2, Katarzhyno kurgans 1 and 2; 3, Revova kurgan 3; 4, Liubasha kurgan 2; 5, Dubynove

kurgan 1; 6, Usatove-Velykyj Kuyalnik; 7, Sugokleya kurgan; 8, Pishchanka kurgan; 9, Shakhta

Stepova kurgan group; 10, Vynohradne kurgan group; 11, Mariupol Neolithic necropolis. Sand,

yellow and beige – Neogene, Pliocene; light and darker brown – Neogene, Miocene; darkest

brown – Paleogene, Neocene; light pink – Archaean, Neoarchaean; dark pink – Archaean,

Eoarchean; crimson – Proterozoic, Paleoproterozoic; green – Lower Cretaceous. Geological map

by A. Grachev, geomap.land.kiev.ua, reproduced with permission. Basemap -

Template_europe_map. Wikimedia Commons CC BY-SA.3.0.

Carbon (δ^{13}C) and nitrogen (δ^{15}N) isotope analysis of bone collagen offers a

complementary perspective on an individual's life history because these systems reflect the

isotopic composition of mainly dietary protein, and secondarily, in varying proportions

depending on nutritional status, carbohydrates, and lipids consumed over the last 7-20 years of

life [7]. While ^{87}Sr/^{86}Sr ratios primarily indicate movement across different geological

landscapes, δ^{13}C and δ^{15}N values can help differentiate individuals whose diets may have been

shaped by different ecological zones, subsistence practices, or physiological conditions [8,9]. In

highly mobile pastoral groups such as those of the North Pontic steppe, δ^{13}C and δ^{15}N values of

collagen are therefore useful for assessing whether individuals with unusual strontium

signatures also had different dietary histories later in life, which can strengthen interpretations

of mobility and regional interaction.

In this study, we employ a multiproxy isotopic approach combining ^{87}Sr/^{86}Sr and δ^{13}C

and δ^{15}N isotope values of bone collagen to investigate the life histories of individuals interred

in seven Eneolithic-Bronze Age kurgans in the northwestern North Pontic region. To strengthen

the reliability of mineral-based isotopic interpretations, we also evaluated the preservation of

selected bioapatite samples using Attenuated Total Reflectance Fourier Transform Infrared

Spectroscopy (ATR-FTIR) prior to ^{87}Sr/^{86}Sr analysis. FTIR is performed to screen bones or teeth

for diagenetic alteration and confirm that the mineral structure is sufficiently preserved so the

measured strontium ratios reflect the individual's original biogenic signal rather than post-

burial contamination. The analyzed specimens were selected to explore factors linking

individuals within and between kurgans and to evaluate the extent to which such data can

inform reconstructions of mobility across the North Pontic steppe during the 4th-3rd millennium

BCE, thereby guiding future large-scale sampling strategies. This research specifically

investigates whether individuals buried in northwest North Pontic kurgans exhibit strontium

isotope ratios consistent with locally derived baseline values or instead reflect mobility within

or beyond the broader steppe throughout their lives. To address this, the study first establishes

$^{87}Sr/^{86}Sr$ baseline values from the available dataset and then evaluates individual mobility

patterns against this framework. This study further evaluates whether individual $\delta^{13}C$ and $\delta^{15}N$

values provide independent lines of evidence for mobility beyond that inferred from strontium

isotope analysis alone. Ultimately, we integrate the newly obtained $^{87}Sr/^{86}Sr$ data with

previously published results to establish the first regional isotopic baseline framework for the

North Pontic region.

Methods

Provenance of archaeological samples

Samples detailed in this report came from a chain of kurgans in the Odesa Region,

southern Ukraine (Fig 1). The Mayaky kurgan cemetery is situated on a promontory of the

Dniester Estuary on the Black Sea coast, while the other five kurgans are located inland, to the

north from Mayaky. Katarzhyno kurgans 1 and 2 were excavated in 1990-1991 by expeditions of

the Odesa Archaeological Protection Center, which existed under the Ukrainian Society for the

Protection of Historical and Cultural Monuments. Dubynove 1, Liubasha 2 and Revova 3 kurgans

were excavated in 2003 by an expedition of the Department of Archaeology of the North-

Western Black Sea Region of the Institute of Archaeology of the National Academy of Sciences

of Ukraine [10]. The kurgan cemetery at Mayaky was excavated by the Odesa Archaeological

Museum of the National Academy of Sciences of Ukraine in 1974-1975, 1986, and 2009 [11–

13]. The excavations were carried out under state permits and in accordance with Ukrainian

heritage law. Following excavation, the remains were accessioned into the Odesa

Archaeological Museum collection, where they are curated as part of the national

archaeological archive. Detailed description of the sites and burials can be found in [1,10–14],

and additional details are provided in S1 File.

Enamel, dentin, and cortical bone samples were obtained from 25 individuals

representing Eneolithic Usatove (3800-3100 BCE), Steppe Eneolithic (3950-3300 BCE), EBA

Yamna (3300-2500 BCE) and Catacomb (2800-2200 BCE), as well as the Middle-Bronze-Age

(MBA) Babyne (Multi-Cordoned Ware, 2100-1600 BCE) archaeological groups (Table 1, S1 File).

Five individuals came from the Dubynove kurgan, eight from the Katarzhyno kurgan 1, one from

the Katarzhyno kurgan 2, five from the Liubasha kurgan, four from the Revova kurgan, and one

each from the Mayaky kurgans 8 (M8.7) and 10 (M10.2.2).

Table 1. Strontium ^{87}Sr/^{86}Sr ratios produced in this study along with other available isotope

metrics for 26 Eneolithic-Bronze Age individuals from the northwestern North Pontic Region.

Specimen	^{14}C, δ^{13}C, δ^{15}N Lab ID	^{14}C, BCE (uncalBP)	Archaeological group	Sex, age	δ^{13}C (‰)	δ^{15}N (‰)	C:N	Sr Lab ID	Material	^{87}Sr/^{86}Sr	*2se* (ratio)/	Local

											2σ (local baseline)	

Dubynove kurgan 1

ID	Lab code	Date	Culture	Sex/age				ANK	Material	⁸⁷Sr/⁸⁶Sr	2σ	baseline
D1.8	Ki-11200	2620-2206 calBCE (3940±70)	Catacomb	Adult	n/a	n/a	n/a	ANK002	Long bone	0.71044	0.00001	
								ANK003	Cranial bone	0.71056	0.00001	
								ANK004	M1 enamel	0.71054	0.00001	Y
D1.9	PSUAMS-7791	2575-2348 calBCE (3970±35)	Catacomb	M, adult	-19.15	12.01	3.31	ANK005	M3 enamel	0.71061	0.00001	Y
D1.11	Ki-11202	2343-1905 calBCE (3720±70)	Catacomb	n/a	n/a	n/a	n/a	ANK008	Enamel	0.71000	0.00001	N
D1.12	Ki-11203	2580-2139 calBCE (3900±80)	Catacomb	M (?), adult	n/a	n/a	n/a	ANK009	Maxilla bone fragment	0.71042	0.00001	
								ANK010	Upper molar fragment	0.71038	0.00001	
								ANK011	Bone fragments	0.71037	0.00001	
D1.14	n/a	2100-1600 BCE	Babyne	M, immature, >2	n/a	n/a	n/a	ANK012	Dentin	0.71051	0.00001	
Local baseline for Dubynove: (human bone, dentin, *n*=6): 0.71030-0.71059										0.71045	0.00015	

Katarzhyno kurgan 1

K1.10	Ki-11376, Ki-9523, PSUAMS-10375	3951-3715 calBCE (5039±27, weighted mean)	Steppe Eneolithic	F, adult	-18.05	15.17	3.28	ANK013	Bone (metacarpal)	0.71025	0.00001	
K1.13	PSUAMS-4762, PSUAMS-9815	3018-2911 calBCE (4359±16, weighted mean)	Yamna	M, adult	-19.10 ±0.02	12.26 ±0.07	3.22 - 3.32	ANK015	Enamel	0.71089	0.00001	N (compatible with Sygokleya)
								ANK016	Enamel	0.71083	0.00001	N (compatible

												with Sygokleya)
K1.16		3300-2500 BCE	Yamna	n/a	n/a	n/a	n/a	ANK017	Bone	0.71020	0.00001	
								ANK018	Enamel	0.70982	0.00001	N (compatible with Shakhta Stepova and Vynohradne)
K1.17		2200-1600 BCE	Babyne	n/a	n/a	n/a	n/a	ANK019	Bone	0.71026	0.00001	
K1.18		n/a	Bronze Age	n/a	n/a	n/a	n/a	ANK020	Bone	0.71046	0.00001	
K1.21E	Beta-751999	2896-2701 calBCE (4215±21)	Yamna	F, adult	-19.50	11.35	3.4	ANK023	Bone	0.71021	0.00001	
								ANK026	Enamel	0.71006	0.00001	N
K1.21		2900-2700 BCE	Yamna	Subadult, ca. 9-11	n/a	n/a	n/a	ANK025	Primary M1 enamel	0.70987	0.00001	N (compatible with Shakhta Stepova and Vynohradne)
K1.21W		2900-2700 BCE	Yamna	M (?), adult				ANK028	Lateral incisor enamel	0.71009	0.00001	N
Katarzhyno kurgan 2												
K2.1	BETA-647662; Ki-17331	3341-3093 calBCE (4487±22); 3314-2537 calBCE (4270±90)	Yamna	F, adolescent, ca. 12-14	-19.10	12.40	3.1	ANK029	Bone	0.71023	0.00001	
								ANK030	M2 enamel	0.70990	0.00001	N (compatible with Shakhta Stepova and Vynohradne)
Local baseline for Katarzhyno 1 and 2 (human bone, dentin, *n*=5): 0.71017-0.71029										0.71023	0.00006	

										87Sr/86Sr	error	
Liubasha kurgan 2												
L4	n/a	2600-2100 BCE	Catacomb	Adult, ca. 175-197 cm tall	n/a	n/a	n/a	ANK032	Lateral incisor enamel	0.70962	0.00001	Y
L7		3300-2500 BCE	Yamna	n/a	n/a	n/a	n/a	ANK034	Bone	0.71028	0.00001	
L8	Ki-11177	2855-2291 calBCE (3990±70)	Yamna	n/a	n/a	n/a	n/a	ANK035	Long bone	0.71026	0.00001	
								ANK037	Bone	0.71037	0.00001	
								ANK038	Dentin	0.71035	0.00001	
L9	BETA-750090	2100-1600 BCE	Babyne	M, adult	-17.43	14.08	3.2	IS-1852	Long bone	0.71025	0.00001	
L12	Ki-11173, BETA-750091	2119-1624 calBCE (3520±80)	Babyne	M, adolescent, ca. 14	-18.80	13.05	3.0	ANK039	Long bone	0.71055	0.00001	
L16	PSUAMS-7790	3074-2888 calBCE (43330±35)	Yamna	M, adult	-19.36	11.75	3.34	ANK040	Bone	0.71033	0.00001	
								ANK041	Enamel	0.71042	0.00001	Y
Local baseline for Liubasha (human bone, dentin, *n*=7): 0.71014-0.71054										0.71036	0.0002	
Mayaky kurgan 8												
M8.7	n/a	3800-3100 BCE	Usatove	F, immature, ca. 3-4				ANK043	Dentin	0.70968	0.00001	
Mayaky kurgan 10												
M10.2.2	PSUAMS-7845	4336-4055 calBCE (5390±30)	Usatove	M, adult	-20.81	15.73	3.24	ANK044	Long bone	0.70962	0.00001	
Local baseline for Mayaky kurgan cemetery (human bone, dentin (*n*=2), *Bos taurus* and *Ovis/Capra* dentin, *n*=8): 0.70947-0.710979										0.70962	0.00018	
Revova kurgan 3												
R3.13	Ki-11172	2580-2207 calBCE (3940±60)	Catacomb	n/a	n/a	n/a	n/a	ANK046	Bone	0.71046	0.00001	

R3.14	Ki-11175	2139-1748 calBCE (3590±70)	Babyne	n/a	n/a	n/a	n/a	ANK047	Bone	0.71054	0.00001	
R3.15	Ki-11060	2458-2031 calBCE (3780±70)	Yamna	n/a	n/a	n/a	n/a	ANK048	Enamel	0.71028	0.00001	Y
R3.19	PSUAMS-4763	3711-3639 calBCE (4905±20)	Usatove	M, adult	-19.17	11.52	3.2	ANK050	Patella	0.71043	0.00001	
								ANK051	Cranial bone	0.71040	0.00001	
Local baseline for Revova (human bone, dentin, n=4): 0.71034-0.71058										0.71046	0.00012	

149 Ki- lab code [14]C dates are from [10]; K1.21E [14]C and diet isotopes, L9 and L12 diet isotopes – this

150 report, the rest of [14]C and diet isotope data are from [1].

FTIR analysis (quality control)

For 12 out of 32 bone and tooth samples for which sufficient material was available, we

assessed preservation of biogenic hydroxyapatite (bioapatite) using attenuated total

reflectance Fourier transform infrared spectroscopy (ATR-FTIR) prior to Sr isotopic analysis to

evaluate potential diagenetic alteration. Approximately 1 mg of powdered bone/tooth

previously treated with 2% NaOCl and 0.2 N acetic acid from each sample was analyzed using a

Thermo Scientific Niolet Summit iS5 FTIR spectrometer equipped with an Everest diamond ATR

crystal. The spectra were collected in the mid-infrared range (4000-400 cm^{-1}) at 4 cm^{-1}

resolution, with 16 scans averaged per spectrum. We generated background spectra prior to

the analytical session. All spectra were baseline-corrected before peak-height measurements.

The peak heights were measured relative to defined baselines following established protocols

for ATR-FTIR assessment of biogenic hydroxyapatite (e.g., [15,16]). Using the baseline corrected

spectra, we calculated Crystallinity indexes (CI), also referred to as infrared splitting factor

(IRSF), carbonate-to-phosphate ratios (C/P), and screened for potential calcite precipitation. All

CI and C/P values were evaluated relative to established ATR-FTIR preservation ranges for

archaeological bioapatite [15].

CI was used to evaluate crystal order and identify potential recrystallization of

bioapatite. CI is calculated as:

$$CI = (A_{565} + A_{605})/A_{595}$$

where A_{565} and A_{605} correspond to the heights of the phosphate V_4 doublet peaks, and A_{595}

corresponds to the height of the valley in between them. Elevated CI values can indicate a

larger crystal size associated with diagenetic-related recrystallization. However, bone and tooth

bioapatite CI values must be evaluated using separate criteria because of the naturally

occurring differences in crystal size between these two tissues, which relate to their formation

and their association with different proportions of organics. Our measured values were

compared with published ATR-FTIR ranges for modern, well-preserved archaeological

bioapatites.

The C/P ratios were calculated to assess carbonate incorporation and to detect potential

exogenous carbonate contamination from the depositional environment. The ratio was

determined using the formula:

$$C/P = A_{1415}/A_{1035}$$

where A_{1415} corresponds to the B-type carbonate V_3 band, and A_{1035} corresponds to the

phosphate V_3 band. Elevated C/P values can indicate the presence of exogenous carbonate.

All spectra were examined for the presence of secondary calcite formation which is

identified by diagnostic absorption bands near ~710 cm^{-1} and ~874 cm^{-1}. We recorded the

presence or absence of calcite qualitatively.

Samples that exhibited CI and C/P values that were within published archaeological

ranges and lacked calcite-associated absorption bands were considered suitable for subsequent

mineral-based isotopic analyses.

Strontium isotope analysis

$^{87}Sr/^{86}Sr$ ratios for specimen L9 (Lab ID IS-1852) were produced by Isobar Science,

Miami, FL, USA, following the methods in [17,18]. The rest of $^{87}Sr/^{86}Sr$ ratios was obtained at

the University of Missouri. First, a chemical pretreatment was applied using the following

sequence: soaking in a solution of 2% NaOCl, centrifugation, elimination of supernatant, rinsing

with mQ water, soaking in 0.2N acetic acid, centrifugation, elimination of supernatant, rinsing

with mQ water, drying. The samples were then placed in PFA vials and dissolved using 7M

Optima Grade HNO_3 at 110˚C for 24 hours. The solutions were then evaporated at 90˚C and the

dry residues were re-dissolved in 2ml of 7M HNO_3. The Sr extraction was conducted using

Eichrom Sr-spec™ resin and 2 ml-reservoir columns following a protocol adapted from[19]. The

Sr eluates were evaporated at 90˚C. The dry residues were dissolved in 0.25 ml 14N HNO_3 and

evaporated at 90 ˚C. The residues were dissolved in 0.05N HNO_3 before analysis on the multi

collector - inductively coupled plasma-mass spectrometer in operation at the University of

Missouri Research Reactor (Nu Plasma II, Nu Instruments).

The instrument was optimized daily to obtain a maximum intensity for ^{88}Sr (~ 7 V). The

sample and standard solutions were prepared to obtain approximately 200 ng g-1 Sr. The

isotopic standard SRM987 was measured multiple times at the beginning of each analytical

session and after every two samples. The average and standard deviation for the SRM987 is

0.71023 ± 0.00004 (2SD) (n = 80). The values were corrected for mass fractionation using the

0.1194 value for the ^{86}Sr/^{88}Sr natural ratio. The isobaric interference at mass 87 was corrected

using the 0.3857 value for the ^{87}Rb/^{85}Rb natural ratio. The isobaric interferences at masses 86

and 84 (^{86}Kr and ^{84}Kr) were corrected using the iterative method. The sample values were

corrected by standard bracketing using the ^{87}Sr/^{86}Sr ratio value of 0.710248 [20]. To control the

reproducibility of the entire process one duplicate (i.e., the entire procedure was applied twice

to the same sample), one replicate (i.e., second analysis of the same solution) and two aliquots

of the SRM1400 (Bone Ash) were analyzed. The values for SRM1400 were 0.71314 +- 0.00001

(2*se*) for each aliquot.

Radiocarbon dating and stable isotope analysis

Newly reported ^{14}C date for K1.21E as well as δ^{13}C and δ^{15}N isotope ratios for L9 and L12

were produced by BETA Radiocarbon lab, Miami, Fl following in-house protocols. The rest of

δ^{13}C and δ^{15}N isotope data came from published datasets and from current study (Table 1, S1

Table).

$\delta^{13}C$ and $\delta^{15}N$ of bone collagen were analyzed descriptively and compared across

regional groups using group means and standard deviations. No parametric tests were applied

to $\delta^{13}C$ and $\delta^{15}N$ data due to sample size constraints. Instead, patterning was evaluated in

relation to established regional isotopic ranges reported in the literature.

Statistical calculations were performed using online resources and Microsoft Excel for Mac

Version 16. Graphs were constructed using Excel.

Statistical analysis

Statistical calculations were performed using online resources

(https://www.socscistatistics.com) and Microsoft Excel for Mac Version 16. Graphs were

constructed using Microsoft Excel (Microsoft 365 for Mac, Microsoft Corp., Redmond, WA,

USA). Descriptive statistics, including mean, standard deviation (SD), and range, were calculated

for each isotopic group. Group-level summaries are reported as mean ±1 SD unless otherwise

specified.

Strontium baseline evaluation

According to [21], up to 100% of non-enamel strontium in buried remains can be

diagenetic, thus representing the soil environment of the burial site. Rather than representing

strictly local bioavailable strontium [22], and taking into account the results of the FTIR analysis,

ratios derived from bone and dentine were interpreted as population-averaged values

reflecting individuals' movements across the broader region surrounding each kurgan during

approximately the last 7–10 years of life, under broadly comparable dietary regimes.

Following [22], we estimated the baseline range for Dubynove, Katarzhyno 1/2,

Liubasha, and Revova kurgans as the average of bone and dentin strontium ratios at each site

±2SD. Strontium ratios from specimen K1.18 were excluded from baseline calculations due to

unclear archaeological context. We emphasize that the inland kurgan baseline represents a

population-averaged mobility envelope rather than a strictly environmental bioavailable

baseline. Because pastoralist groups likely circulated within a shared regional isotopic

catchment, bone/dentin values are interpreted here as reflecting the isotopic space of habitual

movement rather than a sedentary locality. The Mayaky kurgan cemetery baseline was

estimated using the dentin values from M8.7 and the bone values from M10.2.2 as well as

dentin from *Bos taurus* and *Ovis/Capra* [23], the latter reflecting local bioavailable strontium

rather than population-averaged human mobility signatures.

Enamel values were evaluated relative to the obtained bone- and dentine-based

averages, with concordance or divergence between tissues used to identify patterns of

residential stability versus mobility or migration. Individuals were classified as "local" when

enamel ^{87}Sr/^{86}Sr values fell within the population-averaged baseline range (±2SD) established

for the kurgan or regional cluster under consideration (Table 1).

We compared the baseline and enamel ratios of strontium obtained from our sample

selection to published strontium data from human bone, dentin and enamel from elsewhere in

the Pontic steppe. Comparison sites included Pishchanka, Shakhta Stepova, Sygokleya, and

Vynohradne from the east North Pontic steppe and the interred represented Steppe Eneolithic

and EBA including Yamna and Catacomb archaeological groups) [25].

To assess whether the distribution of enamel classifications (within vs. outside baseline)

differed significantly from dentin classifications, contingency tables were constructed and

evaluated using chi-square (χ^2) tests. Statistical significance was assessed at $\alpha = 0.05$. Given the

modest subgroup sample sizes, statistical results are interpreted cautiously and are used to

identify structured patterning rather than to infer population-level parameters.

Results

We used ATR-FTIR analysis to evaluate bioapatite preservation for 12 bone and tooth

samples prior to conducting $^{87}Sr/^{86}Sr$ isotope analysis (Table 2). The CI values ranged from 3.25

to 4.46 (mean ± SD = 3.72 ± 0.35). The values of 10 samples all fall within the published ranges

for both modern and well-preserved archaeological bioapatite and do not indicate extensive

diagenetic recrystallization. The C/P ratios ranged from 0.00 to 0.33 (mean ± SD = 0.12 ± 0.09).

All values were within the established ranges for archaeological bioapatite and do not suggest

the presence of any exogenous carbonate material. No samples exhibited diagnostic absorption

bands indicative of secondary calcite precipitation (~710 cm^{-1} and ~874 cm^{-1}). Two of the

samples had slightly elevated CI values. However, this likely reflects diagenetic crystal

maturation rather than secondary carbonate contamination. Therefore, we have interpreted

these as moderately recrystallized but chemically well-preserved, as further supported by the

acceptable C/P ratios and the absence of calcite in the spectra. Overall, CI and C/P values, along

with the absence of calcite-associated absorption bands, indicate minimal mineral diagenesis in

the analyzed specimens. All samples were therefore considered preserved enough for

subsequent $^{87}Sr/^{86}Sr$ analysis.

Table 2. Results of FTIR-ATR analysis.

Sample Number	Peak 1415 cm^{-1}	Peak 1035 cm^{-1}	C/P Ratio	Calcite Peak at 710 cm^{-1}	Peak 565 cm^{-1}	Peak 605 cm^{-1}	Peak 595 cm^{-1}	CI Value
ANK008	0.01	0.03	0.33	No	0.01	0.009	0.005	3.80
ANK013	0.04	0.32	0.13	No	0.092	0.079	0.045	3.80
ANK026	0.03	0.17	0.18	No	0.06	0.05	0.03	3.67
ANK029	0.03	0.33	0.09	No	0.097	0.079	0.043	**4.09***
ANK034	0.03	0.39	0.08	No	0.119	0.095	0.048	**4.46***
ANK038	0.01	0.04	0.25	No	0.004	0.003	0.002	3.50
ANK041	0.02	0.18	0.11	No	0.064	0.049	0.031	3.65
ANK046	0.03	0.4	0.08	No	0.115	0.09	0.059	3.47
ANK048	0.01	0.12	0.08	No	0.04	0.03	0.02	3.50
ANK050	0.01	0.05	0.20	No	0.007	0.006	0.004	3.25
ANK051	0.04	0.23	0.17	No	0.07	0.06	0.04	3.25
ANK051_dupl	0.02	0.12	0.17	No	0.037	0.033	0.019	3.68

* Values slightly outside of acceptable range.

We generated 37 new ^{87}Sr/^{86}Sr ratios, ranging from 0.70962 to 0.71089, with an average

of 0.71028±0.0003 (Table 1). Enamel samples D1.8, D1.9, L4, L16, and R3.15 of the thirteen

used in the analysis fell within this population-averaged range for the western North Pontic

region. Overall, the samples analyzed in this study fall within the 0.709–0.711 range previously

reported for human skeletal tissues from the North Pontic region [22,25–27], indicating

agreement with the currently available regional strontium isotope data.

Population-averaged baseline ranges based on bone and dentin were tightly constrained

within each kurgan cluster (Table 1), supporting their use for evaluating enamel-derived

mobility signals. No clear association between sex or age and mobility status was observed,

although the limited sample size precludes robust demographic inference. Several enamel

values fell within overlapping baseline ranges of multiple inland kurgans, reflecting the

geological homogeneity of large portions of the northwestern steppe and underscoring the

importance of regional rather than strictly site-specific interpretations.

Enamel strontium signature for the D1.8 and D1.9 Catacomb individuals from the

Dubynove kurgan were in the local for that kurgan range. The strontium isotopic composition of

the enamel from individual D1.8 was within the range defined for Dubynove and similar to that

of the bone from the same individual. Enamel sample from burial D1.9 was at the upper end of

the Dubynove baseline variation. We have interpreted this ratio to be within the local range,

considering the uncertainty in the yet unsampled baseline range. A late Catacomb individual

D1.11 was outside of the Dubynove kurgan baseline range and just outside of the upper end of

the variation of the Shakhta Stepova kurgan from the eastern Pontic steppe [25] (Figs 1,2).

Fig 2. **Enamel strontium ratios of individuals from western North Pontic steppe kurgans**

plotted against the baseline ranges of the North Pontic steppe established on bone and

dentin. The northwestern steppe baseline is represented by Dubynove, Revova, Liubasha,

Katarzhyno, and Mayaky (this report). The eastern steppe baseline is represented by Shakhta

Stepova, Vynohradne, Pishchanka, and Sugokleya [25].

Enamel sample from the Yamna R3.15 individual differed from the Revova kurgan range

but fell within the range of the Liubasha and Katarzhyno kurgans baseline. Enamel from the

Liubasha primary burial L16 was within the Liubasha kurgan range as well as the ranges for

Dubynove and Revova, while the enamel from the Liubasha's Catacomb burial L4 was below

this range but fell within that of the Mayaky Eneolithic archaeological site (Fig 2).

None of the enamel from the Katarzhyno 1 and 2 kurgan samples matched the range of

that kurgan group or that of the northwestern Pontic baseline established by the bone and

dentin strontium ratio ranges from the six kurgans from this report (Fig 2). The three Yamna

adult individuals (K1.16, K1.21E and K2.1) for whom both enamel and bone or dentin were

measured show a discrepancy between the tissues, which is indicative of migration between

the moment the enamel formed and the last 7-10 years of life. Except for K1.13, enamel

samples from burials at Katarzhyno kurgans were within the baseline ranges for the eastern

North Pontic steppe [25]. Enamel from Yamna individuals K1.16 adult and the K1.21 subadult as

well as the adolescent early Yamna K2.1 from the primary burial at the Katarzhyno 2 kurgan

were within the baseline variation of Shakhta Stepova and Vynohradne. Enamel strontium

ratios of the Yamna K1.21E and K1.21W adults did not overlap with any baseline range in Fig 2.

Enamel of individual K1.13 fell within the range of Sugokleya.

To explore whether stable dietary isotopes might provide additional insights into

mobility and possible affiliations with kurgan populations from the greater steppe region, we

compared the δ^{13}C and δ^{15}N values obtained in this study with published data for Eneolithic–

Bronze Age individuals from the eastern North Pontic steppe as well as the West Caspian

steppe [25,28–30]. Eneolithic-EBA δ^{13}C values from northwest Pontic steppe were significantly

different from the eastern Pontic steppe, while $\delta^{15}N$ values were not (Mann-Whitney Two-Tailed test z-scores = 3.45944/ -0.45691, p = 0.00054/0.64552, respectively). We thus plotted the means for these values separately on the graph Fig 3. As evident from the graph, and confirmed statistically, all but two Eneolithic-Bronze Age samples from our sample selection were within the western North Pontic steppe distribution.

Fig 3. **Diet isotope ratios of Eneolithic-Bronze Age populations of the North Pontic and West Caspian-North Caucasus steppe.** Isotope values used to construct the graph, and the corresponding references are listed in S1 Table of S1 File.

Of the two identical twins L9 and L12 from the Babyne archaeological group of the MBA, the isotopic composition of individual L12 falls well within the variation of Eneolithic-EBA populations of the eastern Pontic (joint $\chi^2(2)$ = 1.60, p = 0.45) and was also compatible with those of the northwest Pontic steppe (joint $\chi^2(2)$ = 4.25, p = 0.12). On the other hand, the isotopic composition of individual L9 situated well outside the variation of both comparison groups ($\chi^2(2)$ = 7.82, p = 0.02, and $\chi^2(2)$ = 60.15, p < 0.001, respectively), indicating a diet distinct from either reference population. At the same time, L9's $\delta^{13}C$ and $\delta^{15}N$ values lie within the isotopic ranges of both West Manych Yamna and Eneolithic steppe Maykop, with joint probabilities of p = 0.72 and p = 0.96 respectively, suggesting strong isotopic similarity to these west Caspian/ North Caucasus Piedmont Eneolithic-Bronze Age groups.

Steppe Eneolithic individual K1.10 from the primary burial at the Katarzhyno 1 kurgan was another outlier in terms of $\delta^{13}C$ and $\delta^{15}N$ values, showing a similar to the L9 trend. The

stable isotope values of K1.10 (δ^{13}C = −18.05‰; δ^{15}N = 15.16‰) are statistically incongruent

with Eneolithic-EBA populations of the eastern and western Pontic steppe ($\chi^2(2)$ = 12.49, p =

0.002, and $\chi^2(2)$ = 37.43, p $\ll$ 0.001, respectively), but fall within the isotopic ranges of both

West Manych Yamna and Eneolithic Caspian steppe ($\chi^2(2)$ = 0.52, p = 0.77, and $\chi^2(2)$ = 0.59, p =

0.75), suggesting affinity with the latter groups.

Discussion

Kurgan burial tradition began in the North Pontic steppe in the early 4[th] millennium BCE

[1,31]. Kurgans were re-used for hundreds of years thereafter. Most of the Eneolithic-EBA

kurgans in the North Pontic steppe contain multiple individual burials. The western North

Pontic kurgans studied in this report contained 5-21 burials of mostly single individuals. One of

the burials in the Katarzhyno kurgan 1 contained two adults and a subadult, being one of the

only two such burials in the northwest Pontic steppe among the 2000 surveyed intact burials

[32]. It is unknown at this point if the subadult is genetically related to the adults in the same

burial. Some kurgans functioned as burial places into the modern times [10].

Bone and dentin strontium values from the Bronze Age individuals interred in the five

inland kurgans in the present sample selection were mostly concordant and within a relatively

tight isotopic baseline range of 0.71017-0.71059 (Table 1). In contrast, their enamel strontium

ratios largely deviated from this baseline, suggesting childhood residence in geological

environments distinct from those represented during the final years of life. Yamna individuals

from the Katarzhyno kurgan group, which provided the bulk of enamel datapoints for this

study, show distinctive trends in the enamel strontium signature variation. Such as individual

K1.13 was isotopically distinct from the northwest Pontic baseline and the remaining

Katarzhyno individuals, with a strontium signature within the baseline range of the Sugokleya

kurgan in the eastern steppe. Individuals K1.16 and K2.1 show parallel shifts between enamel

and bone values, with closely matching enamel ratios that point to a shared area of origin

outside of the northwest Pontic. The K1.21 subadult displayed a similar enamel signature,

suggesting that the three grew up in an area geologically similar to that of the Pishchanka and

Vynohradne kurgans in the eastern steppe. The intermediate enamel values observed in

individuals K1.21E and K1.21W are best explained by mobility during enamel formation,

resulting in isotopic signatures that average residence across two geologically distinct regions.

The variability of enamel signatures from Katarzhyno may reflect seasonal mobility of the

pastoralist Yamna groups. At the same time, if they all followed the same mobility pattern, one

would expect that average the same regions at the same time, resulting in a similar isotopic

signature. Alternatively, the interred in the Katarzhyno kurgans may have originated from

different steppe regions with distinct mobility patterns, and comingling in the Katarzhyno

kurgans area.

In contrast, the enamel strontium signature of Yamna individual L16 from the primary

burial in the Liubasha kurgan suggest L16 to have been part of the locally established Yamna

group. Enamel strontium ratios from the Liubasha's burial L4 of the succeeding Catacomb

archaeological culture suggests L4 spent the middle childhood in an area geologically similar to

the coastal area around the Mayaky site. The analysis of archival excavation reports and

pertinent publications revealed a number of archaeological sites associated with the Catacomb

culture group along the Dniester Estuary and the adjacent Black Sea littoral. Thus, it is possible

that L4 may have been associated with a Catacomb population group that utilized these sites.

Strontium baseline for the inland kurgans in our study was overlapping, although the

ranges were somewhat dissimilar in breadth, likely due to limited number of baseline samples

from each kurgan. There was an overlap between the inland kurgan group in our selection and

the lower isotopic strontium ratio range of the Pishchanka kurgan in eastern Ukraine (Fig 2),

possibly due to similar geology, which around Pishchanka consists of Cenozoic sediments and

exposed Proterozoic plutonic rocks [25]. The area around the Sugokleya kurgan encompassed

most of the known bioavailable strontium variation in Ukraine, likely explained by having a

complex geology (Fig 1), including a unique combination of crystalline Precambrian and

sedimentary layers, tectonic features, igneous intrusions, and ore formation [35–37]. Adding to

the uniqueness of the area's geology is its association with the 24km Boltysh crater, thought to

had been formed by an asteroid that followed the Chicxulub impact at the Cretaceous-

Paleogene boundary ca. 66 MYA [38].

δ^{13}C and δ^{15}N values of all but two of the studied individuals from the inland kurgans

were within the dietary range of the Eneolithic-Bronze Age inhabitants of the North Pontic

steppe (Fig 3). δ^{13}C and δ^{15}N values of Eneolithic individual K1.10 and the MBA Babyne

individual L9 indicate that they spent the last part of their lives outside of the North Pontic,

possibly in the west Caspian area. Supporting a west Caspian dietary affiliation for some mobile

steppe individuals from the North Pontic, the Eneolithic Taraclia II.2.14 individual interred in

the western steppe, identified as a second cousin of Sharakhalsun burial 18 from the Manych

steppe [1], exhibited δ^{13}C and δ^{15}N values consistent with the west Caspian steppe range (Fig 3).

At the same time, $\delta^{13}C$ and $\delta^{15}N$ values of L12, the L9's identical twin, are within the North

Pontic steppe range. This suggests that L12, an adolescent, likely lived and died in the North

Pontic, while L9, who lived into adulthood, may have spent the last decade of life in the North

Caucasus-West Caspian area, but was buried near his likely birthplace in the western steppe in

the same kurgan as his twin brother. On the other hand, L9 could have sustained on a special

diet. Alternatively, L9's $\delta^{13}C$ and $\delta^{15}N$ values may reflect population-level dietary changes,

possibly associated with the North Atlantic cooling event at c. 2250 BCE/4200 BP [39], followed

by a rapid climate change c. 2250-1850 BCE (4200-3800 BP [40]).

Comparative diet isotope information for the Babyne archaeological group is limited.

Anthropological studies indicate that Babyne individuals experienced considerably higher levels

of physiological stress compared to steppe populations from the preceding chronological period

[41]. Nutritional stress has been shown to lead to elevated $\delta^{15}N$ ratios in humans [42] as seen in

L9, but the pattern is less consistent and smaller in magnitude for $\delta^{13}C$. If climate change were

the cause of shifts in $\delta^{13}C$ and $\delta^{15}N$ values, then these data could be used as a correlate of

environmental dynamism. The limited data available on the Babyne diet and the scarcity of

climatic data for the North Black Sea region during the MBA hinder a detailed evaluation of the

latter hypothesis.

Using data obtained in this report in combination with those available from literature,

we constructed first strontium baseline heatmap for the North Pontic region during the

Eneolithic-MBA chronological period (Fig 4). Additional analyses of strontium isotope ratios

across the North Pontic region, particularly through denser spatial sampling of both human

remains and environmental proxies, will expand this baseline framework, improve

discrimination between overlapping geological zones, and allow more precise reconstructions

of mobility patterns, seasonal movement, and population connectivity within the steppe

landscape that played a central role in continental-scale cultural and demographic

transformations. As a pivotal corridor linking the Carpathian Basin, the Balkans, the Caucasus,

and the Volga-Ural region, the North Pontic steppe served as a dynamic center where

populations interacted, merged, and reorganized. Expanding strontium-based provenance

studies here would therefore clarify not only patterns of movement, but also the processes

through which this region became a major arena of change in prehistoric Eurasia.

Fig 4. **A heatmap of ^{87}Sr/^{86}Sr baseline distribution in the North Pontic region during c. 4000-**

2000 BCE chronological period based on the current study and the data available from

literature. Verteba baseline represents values from water and snail shells from a ritual site of

Eneolithic Trypillia culture [43]. Values for Dubynove, Katarzhyno, Liubasha, Revova (this

report), Pishchanka, Shakhta Stepova, Sugokleya, and Vynohradne [25] are from human bone

and dentin. Values for Mayaky are from human bone and dentin as well as *Bos* and *Ovis/Capra*

dentin [23].

Acknowledgements

468   The authors thank the Odesa Archaeological Museum for providing access to the samples

469   studied in this report.

References

1.   Nikitin AG, Lazaridis I, Patterson N, Ivanova S, Videiko M, Dergachev V, et al. A genomic

history of the North Pontic Region from the Neolithic to the Bronze Age. Nature.

2025;639: 124–131. doi:10.1038/s41586-024-08372-2

2.   Gimbutas MA. Three Waves of the Kurgan People into Old Europe, 4500-2500 B.C.

Journal of Indo-European Studies. 1997;18: 240–268.

3.   Golturenko I, Manyuk V. Geological heritage of the south-western Ukraine (Odessa,

Mykolaiv and Kherson region). Journal of Geology, Geography and Geoecology. 2008;17:

24–28.

4.   Claesson S, Bibikova E, Shumlyanskyy L, Dhuime B, Hawkesworth CJ. The oldest crust in

the Ukrainian Shield – Eoarchaean U–Pb ages and Hf–Nd constraints from enderbites and

metasediments. Geological Society, London, Special Publications. 2015;389: 227–259.

doi:10.1144/SP389.9

5.   Galets'kyi] LS, editor. An atlas of the geology and mineral deposits of Ukraine. Toronto :

University of Toronto Press, 2006.; 2006. Available:

https://search.library.wisc.edu/catalog/9910033291802121

6.   Liventseva H. The Mineral Resources of Ukraine. Madrid; 2022 May.

doi:10.21028/hl.2022.05.17

7.   Ben-David M, Flaherty EA. Stable isotopes in mammalian research: a beginner's guide. J

Mammal. 2012;93: 312–328. doi:10.1644/11-MAMM-S-166.1

8.   Reitsema LJ. Beyond diet reconstruction: Stable isotope applications to human

physiology, health, and nutrition. American Journal of Human Biology. 2013;25: 445–456.

doi:10.1002/ajhb.22398

9.   Neff NC, Busquets-Vass G, Ray EE, Robinson M, Thompson AE, Mes J, et al. Nutritional

adaptations to early maize cultivation: Earliest isotopic evidence of maize-based animal

provisioning in the neotropics. Sci Adv. 2026.

10.   Ivanova SV, Petrenko VG, Vetchinnikova NE. Kurgans of ancient herdsmen from the

South Bug and Dniester interfluve. Odessa: KP OGT; 2005.

11.   Petrenko VG, Kaiser E. Kompleksnyy pamyatnik Mayaki: novyye izotopnyye daty i

voprosy khronologii nalichnykh kul'tur. Materialy po arkheologii Severnogo

Prichernomor'ya. 2011;12: 31–61.

12.   Zinkovsky K V., Patokova EF. Issledovaniya Mayakskogo mogil'nika v 1974 g. .

Arkheologicheskiye issledovaniya Severo-Zapadnogo Prichernomor'ya. Kiev: Naukova

Dumka; 1978. pp. 134–143.

13.   Patokova EF, Petrenko VG, Burdo NB, Polischuk LY. Pamyatniki Tripol'skoj kul'tury v

Severo-Zapadnom Prichernomorje. Kiev: Naukova Dumka; 1989.

14.   Nikitin AG, Ivanova S, Culleton BJ, Potekhina I, Reich D. New radiocarbon and stable

isotope data from the Usatove culture site of Mayaky in Ukraine. SSRN Electronic Journal.

2023. doi:10.2139/ssrn.4236123

15.   France CAM, Sugiyama N, Aguayo E. Establishing a preservation index for bone, dentin,

and enamel bioapatite mineral using ATR-FTIR. J Archaeol Sci Rep. 2020;33: 102551.

doi:10.1016/j.jasrep.2020.102551

16.   Lopes C de CA, Limirio PHJO, Novais VR, Dechichi P. Fourier transform infrared

spectroscopy (FTIR) application chemical characterization of enamel, dentin and bone.

Appl Spectrosc Rev. 2018;53: 747–769. doi:10.1080/05704928.2018.1431923

17.   Pourmand A, Dauphas N. Distribution coefficients of 60 elements on TODGA resin:

Application to Ca, Lu, Hf, U and Th isotope geochemistry. Talanta. 2010;81: 741–753.

doi:10.1016/j.talanta.2010.01.008

18.   Pourmand A, Prospero JM, Sharifi A. Geochemical fingerprinting of trans-Atlantic African

dust based on radiogenic Sr-Nd-Hf isotopes and rare earth element anomalies. Geology.

2014;42: 675–678. doi:10.1130/G35624.1

19.   De Muynck D, Huelga-Suarez G, Van Heghe L, Degryse P, Vanhaecke F. Systematic

evaluation of a strontium-specific extraction chromatographic resin for obtaining a

purified Sr fraction with quantitative recovery from complex and Ca-rich matrices. J Anal

At Spectrom. 2009;24: 1498. doi:10.1039/b908645e

20.   Thirlwall MF. Long-term reproducibility of multicollector Sr and Nd isotope ratio analysis.

Chemical Geology: Isotope Geoscience section. 1991;94: 85–104. doi:10.1016/0168-

9622(91)90002-E

21.   Budd P, Montgomery J, Barreiro B, Thomas RG. Differential diagenesis of strontium in

archaeological human dental tissues. Applied Geochemistry. 2000. doi:10.1016/S0883-

2927(99)00069-4

22.   Gerling C, Kozak O, Liubichev M, Rudich T, Schultze E. Investigating the origins of the

Cherniakhiv community at Voitenki, East Ukraine, through strontium isotope analysis.

Praehistorische Zeitschrift. 2025. doi:10.1515/pz-2024-2048

23.   Petrenko V, Gerling C, Kaiser E. Majaki – ein komplexes Denkmal der Usatovo-Kultur.

Isotopendaten werfen neue Fragen auf. In: Hansen S, Wagner M, editors. Eurasia Antiqua

Zeitschrift für Archäologie Eurasiens. Bonn: Rudolf Habelt Verlag; 2018. pp. 45–74.

24.   Tukey JW. Exploratory Data Analysis. Reading, MA: Addison-Wesley; 1977.

25.   Gerling C. Prehistoric Mobility and Diet in the West Eurasian Steppes 3500 to 300 BC. De

Gruyter; 2015. doi:10.1515/9783110311211

26.   Ventresca Miller AR, Johnson J, Makhortykh S, Gerling C, Litvinova L, Andrukh S, et al. Re-

evaluating Scythian lifeways: Isotopic analysis of diet and mobility in Iron Age Ukraine.

PLoS One. 2021;16: e0245996. doi:10.1371/journal.pone.0245996

27.   Ventresca Miller AR, Johnson JA, Makhortykh S, Litvinova L, Taylor T, Rolle R, et al.

545   Mobility and diet in the Iron Age Pontic forest-steppe: A multi-isotopic study of urban

546   populations at Bel'sk. Archaeometry. 2019;61: 1399–1416. doi:10.1111/arcm.12493

547   28.   Hollund HI, Higham T, Belinskij A, Korenevskij S. Investigation of palaeodiet in the North

548   Caucasus (South Russia) Bronze Age using stable isotope analysis and AMS dating of

549   human and animal bones. J Archaeol Sci. 2010;37: 2971–2983.

550   doi:10.1016/j.jas.2010.08.009

551   29.   Knipper C, Reinhold S, Gresky J, Berezina N, Gerling C, Pichler SL, et al. Diet and

552   subsistence in Bronze Age pastoral communities from the southern Russian steppes and

553   the North Caucasus. Biehl PF, editor. PLoS One. 2020;15: e0239861.

554   doi:10.1371/journal.pone.0239861

555   30.   Shishlina NI, Zazovskaya EP, van der Plicht J, Hedges REM, Sevastyanov VS, Chichagova

556   OA. Paleoecology, Subsistence, and 14 C Chronology of the Eurasian Caspian Steppe

557   Bronze Age. Radiocarbon. 2009;51: 481–499. doi:10.1017/S0033822200055879

558   31.   Nikitin AG, Ivanova S. Eneolithic sanctuaries and Bronze Age kurgans in the Pontic Steppe

559   as markers of climatic shifts. 28th EAA Annual Meeting, Budapest, Hungary, 2022 .

560   Prague: European Association of Archaeologists; 2022. p. 20.

561   32.   Yarovoy E V. Skotovodcheskoye naseleniye Severo-Zapadnogo Prichernomor'ya epokhi

562   rannego metalla. Dissertatsiya na soiskaniye uchenoy stepeni doktora istoricheskikh nauk

563   po spetsial'nosti 07.00.06 - Arkheologiya. Doctorate Dissertation, Moskovskiy

564   gosudarstvennyy universitet im. M.V. Lomonosova. 2000.

33. Toshchev GN. Odesskiy kurgan: Vzglyad cherez desyatiletiya. Donetskiy arkheologicheskiy

sbornik. 2006;12: 62–69.

34. Dobrovolskij A V. Raskopki kurgana v predmest'i Odessy - "Slobodka-Romanovka."

Zapiski Odesskogo Obshestva Istorii i Drevnostej. 1915;32: 123–147.

35. Manyuk V V. Objects of Geological Heritage in the Kirovohrad Region as Basis for

Geotourism Development. In: Emelyanov IG, et al., editors. Proceedings of the National

Museum of Natural History. Kyiv: National Academy of Sciences of Ukraine; 2015. pp.

10–16.

36. Kurylo S, Uher P, Broska I, Lyzhachenko N, Bondarenko S, Gieré R. Fine-grained petalite

and spodumene dykes in the Stankuvatske Li-deposit, Ukrainian Shield: products of

tectono–metamorphic recrystallisation. Mineral Mag. 2022;86: 863–882.

doi:10.1180/mgm.2022.100

37. Baran AM, Vysotsky OB, Dovbush TI, Kovtun OV, Stepanyuk LM, Yaskevych TB. The

Uranium-Lead Monazite Isotopic Age of Granitoids of the Kropyvnytskyi Massif, on the

Example of the Subottsi Village Open Pit (Ingul Domain of the Ukrainian Shield).

Geochemistry and ore formation. 2024;45: 54–63. doi:10.15407/gof.2024.45.054

38. Pickersgill AE, Mark DF, Lee MR, Kelley SP, Jolley DW. The Boltysh impact structure: An

early Danian impact event during recovery from the K-Pg mass extinction. Sci Adv.

2021;7. doi:10.1126/sciadv.abe6530

39.   Bond G, Showers W, Cheseby M, Lotti R, Almasi P, DeMenocal P, et al. A pervasive

585   millennial-scale cycle in North Atlantic Holocene and glacial climates. Science (1979).

586   1997. doi:10.1126/science.278.5341.1257

587   40.   Mayewski PA, Rohling EE, Curt Stager J, Karlén W, Maasch KA, Meeker LD, et al. Holocene

588   climate variability. Quat Res. 2004;62: 243–255. doi:10.1016/j.yqres.2004.07.001

589   41.   Kozak A. Burials of Sugoklea Barrow. Palaeopathological Aspect of the Study. Vestnik

590   Volgogradskogo gosudarstvennogo universiteta Serija 4 Istorija Regionovedenie

591   Mezhdunarodnye otnoshenija. 2020; 256–274. doi:10.15688/jvolsu4.2020.4.17

592   42.   Fuller BT, Fuller JL, Sage NE, Harris DA, O'Connell TC, Hedges REM. Nitrogen balance and

593           $\delta15$ N: why you're not what you eat during nutritional stress. Rapid Communications in

594   Mass Spectrometry. 2005;19: 2497–2506. doi:10.1002/rcm.2090

595   43.   Pokutta D, Tkachuk T, Tóth P. Strontium in pots: An experimental study on the

596   determination of strontium isotope ratios (^{87}Sr/^{86}Sr) and provenance of prehistoric

597   vessels from Verteba Cave, Ukraine (Tripolye culture). In: Dębiec M et al., editor. From

598   Farmers To Heroes? Archaeological Studies In Honor Of Sławmir Kadrow.

599   Universitätsforschungen zur prähistorischen archäologie 376. Bonn: Verlag Dr. Rudolf

600   Habelt, GmbH.; 2022. pp. 223–232.

Supporting Information

603   S1 File. Description of sites and burials.

604   S1 Table. Stable isotopes of carbon and nitrogen used to construct the graph in Fig 3.

Кропивницький **7**

Дніпро **8** **9**

Запоріжжя

5

4

3

2

Миколаїв

Херсон

10

11

6 Одеса

1

20 km