Congenital aphantasia reveals frontotemporal and cingulate structural alterations underlying conscious access to imagery

Yusaku Takamura[1,2], Romain Delsanti[1,3], Laurent Cohen[1], Paolo Bartolomeo[1,†], Jianghao Liu[1,4,†,*]

1. Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, Inserm, CNRS, AP-HP, Hôpital de la Pitié-Salpêtrière, F-75013 Paris, France

2. Neurorehabilitation Research Center, Kio University, Nara 635-0832, Japan

3. Cyceron, Caen Normandie University, GIP Cyceron, Campus Jules Horowitz, bvd Henri Becquerel, BP 5229, 14074, Caen, France

4. Corporate Research, Dassault Systèmes, 78140 Vélizy-Villacoublay, France

†shared senior authorship

*Correspondence to: jianghaolouisliu@gmail.com

Abstract

Congenital aphantasia is characterized by a lifelong absence of voluntary visual imagery despite preserved visual knowledge, offering a natural model for dissociating sensory representation from conscious imagery. Functional imaging evidence suggests that this dissociation may arise from altered top-down interactions between higher-order control systems and high-level visual cortex, but its structural correlates remain unknown. Here, using diffusion and structural MRI in 18 individuals with congenital aphantasia and 18 matched visualisers, we tested two competing accounts of aphantasia: one predicting structural differences in visual pathways, the other predicting differences in higher-order associative networks. Across complementary analyses of white-matter tract microstructure, functional-ROI tractography, graph-theoretic network organization, and cortical morphometry, aphantasia was associated with selective structural differences in frontotemporal and cingulate white-matter tracts — including the uncinate fasciculus, posterior interparietal callosal fibers, and dorsal cingulum — and in frontotemporal cortical regions, including the anterior insula, anterior prefrontal cortex, and medial temporal cortex. By contrast, we found no reliable group differences in early visual cortex, major visual tracts, or the direct structural connections of the core imagery network. Congenital aphantasia therefore exhibits a selective structural phenotype centered on frontotemporal and cingulate systems, sparing the principal visual pathways. These findings suggest that higher-order systems supporting integration and conscious access — rather than visual representations themselves — constitute the primary structural substrate of absent conscious imagery in congenital aphantasia.

Introduction

Congenital aphantasia, characterized by a lifelong absence of voluntary visual mental imagery, affects an estimated 2–4% of the population (Zeman, 2024) and presents a striking paradox (Liu & Bartolomeo, 2025; Arcangeli & Bartolomeo, 2025): aphantasic individuals can accurately retrieve visual properties of objects from memory and perform normally on many visual imagery tasks, yet report little or no subjective imagery experience (Siena & Simons, 2024; Liu & Bartolomeo, 2023; Pounder et al., 2022; Suggate et al., 2026). This dissociation suggests that access to visual knowledge can be preserved even when the phenomenological experience of imagery is absent (Liu & Bartolomeo, 2023). Congenital aphantasia therefore provides a useful model for understanding what higher-order mechanisms are necessary to transform internally generated representations into conscious imagery experience (Liu, 2026, in press).

Beyond the absence of visual imagery, congenital aphantasia has been associated with a broader cognitive and affective profile consistent with a neurodevelopmental condition (Sokolowski & Levine, 2023). Aphantasic individuals retrieve fewer phenomenological details from autobiographical events and show reduced hippocampal-occipital functional connectivity during memory recall (Zeman et al., 2020; Dawes et al., 2022; Monzel et al., 2024b). Difficulties in domains such as face recognition and emotional awareness have also been reported (Dance et al., 2023; Monzel et al., 2024a; Zeman et al., 2020) and intrusive memories are less likely to take a visual form in aphantasics (Keogh et al., 2023). Together with the suggested co-occurrence of aphantasia with neurodevelopmental conditions primarily affecting higher-order rather than sensory systems (Dance et al., 2021; Milton et al., 2021; Mawtus et al., 2024), these findings suggest that aphantasia may involve higher-order systems extending beyond visual representation alone. Whether this broader profile is associated with a common structural substrate remains unknown.

Structural neuroimaging offers one way to test whether the functional alterations reported in aphantasia have an anatomical basis. In neurodevelopmental conditions, functional alterations often co-occur with white-matter microstructural differences, suggesting that atypical connectivity may arise from altered axonal development rather than acquired disruption — as seen for example in congenital prosopagnosia, ADHD, and autism spectrum conditions (Parlatini et al., 2023; Thomas et al., 2009; Travers et al., 2012). The suggested association of congenital aphantasia with some of these conditions, together with reports of altered autobiographical memory, face recognition, and emotional awareness (Zeman et al., 2020; Dawes et al., 2022; Dance et al., 2023; Monzel et al., 2024a), raises the possibility that aphantasia is associated with structural differences in fronto-temporal and cingulate systems. Characterizing these structural correlates may therefore illuminate both the stability of the condition across the lifespan and its broader cognitive profile (Liu, 2026; Zeman et al., 2020).

Neuroimaging studies of aphantasia (Milton et al., 2021; Liu et al., 2025a; Chang et al., 2025; Liu et al., 2025b; Cabbai et al., 2024; Montabes De La Cruz et al., 2024) have yielded two competing accounts. One view holds that aphantasia arises from alterations in visual pathways or early visual cortex, reflecting reduced sensory strength of internally generated representations (Keogh & Pearson, 2018). Consistent with this possibility, aphantasic individuals show altered decoding during imagery in V1, suggesting that imagery representations may not reach the perceptual-like format typically associated with vivid imagery experience (Chang et al., 2025). An alternative account emphasizes disrupted top-down modulation from higher-order brain regions (Liu, 2026; Liu & Bartolomeo, 2025). During imagery tasks, despite preserved activity in high-level visual areas, aphantasic individuals show reduced functional coupling between the fusiform imagery node (FIN)—a region of fusiform cortex selectively activated during visual imagery (Bartolomeo et al., 2026; Liu et al., 2025b; Spagna et al., 2021)—and left prefrontal regions (Liu et al., 2025b), including the anterior prefrontal cortex (aPFC) and frontal eye field (FEF). Higher-order attention and salience regions, including FEF and anterior insula, have been proposed to contribute to the top-down amplification of internally generated representations and their segregation from external input (Corbetta & Shulman, 2002). This view is further supported by reduced representational overlap between imagery and perception in high-level visual regions (Liu et al., 2025b) and orbitofrontal cortex (OFC; Liu et al., 2025a), as well as altered frontoparietal-temporal coupling and reduced neural signal complexity (Noble et al., 2026). These accounts make different structural predictions: a local visual-pathway account predicts differences in posterior visual tracts, including the vertical occipital fasciculus (VOF), inferior longitudinal fasciculus (ILF), and inferior fronto-occipital fasciculus (IFOF), whereas a higher-order account predicts differences in fronto-temporal and cingulate pathways.

Here, we tested these competing predictions using diffusion-weighted and T1-weighted MRI. We combined four complementary approaches—tractometry, functional region-of-interest (fROI)-based tractography, graph-theoretic network analysis, and surface-based cortical morphometry—to characterize potential group differences across tract, network, and cortical scales (see analysis pipeline in Fig. S1).

Results

Atlas-based tractometry reveals selective microstructural differences

We quantified fractional anisotropy (FA) along atlas-defined major white-matter tracts, providing a tract-profile measure of white-matter microstructure. We assessed major tracts associated with visual processing, including the VOF, IFOF, and ILF, as well as tracts associated with higher-order functions, including fronto-parietal attention pathways (Superior longitudinal fasciculus: SLFs, Arcuate fasciculus: AF), medial fronto-temporal pathways (Uncinate fasciculus: UF, ventral cingulum), and cognitive-control pathways (dorsal cingulum).

Figure 1 and Table S1 show the tractometry results. No significant FA differences were detected in the visual tracts examined. In contrast, group differences were observed in several higher-order associative pathways. Relative to visualizers, aphantasic individuals showed reduced FA in both the left and right UF, localized to the anterior temporal segment (Fig. 1A). Reduced FA was also observed in the posterior interparietal corpus callosum. By contrast, aphantasic individuals showed increased FA across much of the bilateral dorsal cingulum (Fig. 1B). No significant clusters were identified in the remaining tracts.

To further characterize these findings, we examined axial diffusivity (AD), radial diffusivity (RD), and neurite orientation dispersion and density imaging (NODDI) metrics at the affected segments (Fig. S2 and Table S1). These follow-up analyses were consistent with microstructural differences rather than fiber geometry alone, such as crossing-fiber artefact.

The right SLF-II and right ventral cingulum showed marginal reductions in FA (Fig. S3); however, these effects may be confounded by fiber geometry and were therefore not interpreted further.

Figure 1. Tractometry reveals selective white-matter microstructural differences in aphantasia. (A) Tracts showing reduced fractional anisotropy (FA) in aphantasic individuals relative to visualizers (Vis. > Aph.): left uncinate fasciculus (UF), right UF, and posterior

interparietal corpus callosum. Upper panels show tract profiles (mean ± SEM); lower panels show tract reconstructions from a representative participant, color-coded by segment position (nodes 1-100). Red lines indicate aphantasic individuals and blue lines indicate visualizers. Shaded gray regions mark segments showing significant group differences (p < 0.05, FWE-corrected). **(B)** Tracts showing increased FA in aphantasic individuals relative to visualizers (Aph. > Vis.): left and right dorsal cingulum. Upper panels show tract profiles (mean ± SEM); lower panels show representative tract reconstructions, color-coded by segment position. Red lines indicate aphantasic individuals and blue lines indicate visualizers. Shaded gray regions and black bars on the x axis mark significant segments (p < 0.05, FWE-corrected). Thin lines show individual participant profiles; thick lines show group means.

Graph-theoretic network analysis reveals disrupted segregation and inter-network hubness

We next applied graph-theoretic analysis to examine whether the tract-level differences extended to the structural organization of large-scale brain networks. We generated a 400 × 400 whole-brain structural connectivity matrix based on the Schaefer 17-network parcellation (see analysis pipeline in Fig. 2A). From this weighted undirected graph, we computed node-level metrics to characterize network organization. Given the study hypothesis, analyses focused on nodes within the visual, attention, salience, and default mode networks.

Group differences were observed in three higher-order network metrics (Fig. 2B–D; Table S2). In aphantasia, the left FEF, a key node of the dorsal attention network (DAN B in Fig. 2A), showed reduced node strength, indicating weaker overall structural connectivity at this node. The left anterior insula, part of the salience/ventral attention network (VAN A), and the left dorsolateral prefrontal cortex (dlPFC) of the default mode network (Default B) showed decreased clustering coefficient, indicating reduced local network segregation in salience and cognitive-control regions. By contrast, the right dorsomedial prefrontal cortex (dmPFC) of the default mode network (Default A) showed increased participation coefficient, indicating stronger inter-network hubness in aphantasic individuals. In addition, the right somatomotor network showed reduced participation coefficient in aphantasic individuals. Control analyses using alternative parcellations confirmed that these findings were robust across atlas options (Fig. S5A–B).

Tracing direct structural connections to these nodes showed that the left anterior insula was connected primarily via the AF and IFOF (Fig. S5C), whereas dmPFC connectivity involved dorsal cingulum pathways. These anatomical relationships are considered further in the Discussion.

By contrast, no significant differences were observed in the visual cortex nodes examined, in keeping with the absence of tractometry differences in the major visual pathways.

Figure 2. Structural network analysis reveals reduced local segregation and increased inter-network hubness in congenital aphantasia. (A) Analysis pipeline. Whole-brain tractography streamlines were weighted using spherical-deconvolution informed filtering of tractograms (SIFT2) and normalized by ROI volume to generate a 400 × 400 structural connectivity matrix based on the Schaefer 17-network parcellation. Node-level graph-theoretic metrics were then computed from the weighted undirected graph. **(B)** Node strength was reduced in the left frontal eye field (FEF; dorsal attention network B, DAN B) in aphantasic individuals relative to visualizers, indicating weaker overall structural connectivity at this node. **(C)** Clustering coefficient was reduced in the left anterior insula (salience/ventral attention network A, Sal/VAN A) and left dorsolateral prefrontal cortex (dlPFC; Default B), indicating reduced local network segregation in salience and association-network regions. **(D)** Participation coefficient was increased in the right dorsomedial prefrontal cortex (dmPFC; Default A) in aphantasia, indicating greater inter-network hubness at this node. Blue indicates Vis. > Aph.; dark red indicates Aph. > Vis. Boxplots show median and interquartile range; *p < 0.05, **p < 0.01, ***p < 0.001, FWE-corrected.

No difference found in white-matter connections of the core imagery network

We next characterized the white-matter connections of a core imagery network whose functional properties are altered in aphantasia. We defined fROIs on the basis of our 7T fMRI results (Liu et al., 2025b): the FIN, left aPFC, left OFC, and right anterior intraparietal sulcus (aIPS). For each of the four ROIs (Fig. S4A) in each group, we identified streamlines intersecting its projection on the grey/white-matter interface and quantified tract structure using two indices: volume-normalized streamline count (Fig. S4B) and streamline proportion (Fig. S4C) within anatomically defined bundles. No significant between-group differences were observed for either index in any tract or ROI after correction for multiple comparisons. Bayes factor (BF) analysis further provided some moderate evidence for the absence of a group difference, thus consistent with an equivalence, in volume-normalized streamline count in the VOF connecting the FIN (BF = 0.32), the major white-matter bundle linking dorsal and ventral visual cortex. Together, these findings indicate no reliable alteration in the direct structural connections examined within the core imagery network.

Cortical thickness: reduced in the left aPFC, increased in limbic regions

We finally examined cortical thickness across the whole brain. Prior work has linked prefrontal and early visual cortical thickness to individual differences in imagery vividness (Bergmann et al., 2016a, 2016b), including a relationship between smaller V1 and stronger imagery (Bergmann et al., 2016b). We therefore examined whether aphantasia is associated with thicker or larger V1.

No significant differences in cortical thickness or volume were observed in any early visual cortex region, with BFs yielding moderate evidence of group equivalence in both the calcarine cortex (Table S3) and V1 (Table S4). By contrast, whole-brain analysis revealed a single significant prefrontal cluster of reduced thickness in the left aPFC in aphantasia (Fig. 3A). Cluster statistics are reported in Table S5. Aphantasic individuals also showed greater cortical thickness in multiple medial temporal limbic regions, including left anterior parahippocampal gyrus and entorhinal cortex; right entorhinal cortex, retrosplenial cortex, posterior cingulate cortex. All statistical details are shown in Table S3. NODDI analysis identified higher NDI values in bilateral medial temporal regions and the left precuneus in aphantasia (Fig. S6), consistent with microstructural differences in these medial temporal regions.

To provide functional context for the observed thickness differences, we performed meta-analytic decoding of the t-map using Neurosynth (Fig. 3B). The left aPFC cluster showed strong associations with the terms "intentions" (r = 0.26), "actions" (r = 0.25), and "near (space)" (r = 0.25). Regions of increased thickness in aphantasia showed associations with anatomical terms including hippocampus, parahippocampus, and amygdala (all rs > 0.14), providing convergent functional annotation for the identified regions.

Figure 3. Regional differences in cortical thickness between aphantasic individuals and visualizers. (A) Whole-brain t map of the group comparison. Blue indicates regions of reduced cortical thickness and red indicates regions of increased cortical thickness in aphantasia, with whole-cortex FDR correction. Boxplots show individual participant values (dots), median, and interquartile range. *p < 0.05, **p < 0.01, ***p < 0.001. Vis., visualizers; Aph., aphantasia. aPFC: anterior prefrontal cortex; PoCG: post central gyrus; PHG: parahippocampal gyrus; EC: entorhinal cortex; ITG: inferior temporal gyrus; RSC: retrosplenial cortex; PCC: posterior cingulate cortex. **(B)** Functional term associations derived from meta-analytic decoding of the t map using Neurosynth. Blue terms are associated with regions of reduced cortical thickness in aphantasia, and red terms with regions of increased cortical thickness. Term size reflects the strength of the Pearson correlation with the corresponding meta-analytic activation map.

Multivariate logistic regression

As an exploratory analysis, we used penalized multivariate logistic regression to identify the most parsimonious combinations of structural features associated with group separation.

Separate models were fitted for FA, cortical thickness, and graph-theoretical measures, using only features showing group differences in the univariate analyses. The exploratory models showed strong in-sample discrimination, with all areas under the curve (AUCs) > 0.92. Across models, the strongest independent predictors of aphantasia were reduced FA in the right uncinate fasciculus and posterior interparietal corpus callosum, increased FA in the left dorsal cingulum, lower clustering coefficient in the left anterior insula, lower node strength in the left FEF, reduced left aPFC thickness, and greater anterior parahippocampal and retrosplenial cortical thickness. Full model statistics are reported in the Supplementary Materials.

Discussion

Congenital aphantasia has been proposed to reflect reduced functional coupling between higher-order prefrontal systems and high-level visual cortex, rather than a deficit in sensory representation alone (Liu & Bartolomeo, 2025). Here, we report structural findings that are, to our knowledge, the first direct anatomical evidence consistent with this account. Across tractometry, network analysis, and cortical morphometry, aphantasia was associated with a selective pattern of structural differences centered on higher-order fronto-temporal and cingulate systems, whereas we detected no reliable group differences in early visual cortex or in the major visual tracts examined.

Four converging observations characterize this structural phenotype. First, the UF and posterior interparietal corpus callosum showed reduced FA in aphantasia, implicating fronto-temporal and interhemispheric associative pathways. This pattern is consistent with prior functional findings of reduced coupling between OFC and anterior temporal regions in aphantasia (Liu et al., 2025a), two territories linked by the UF, potentially reflecting degraded binding between temporal mnemonic content and OFC affective and perceptual processing (Coad et al., 2020; Thomas et al., 2015). It may also be relevant to the broader behavioral profile reported in aphantasia, including altered autobiographical memory, face recognition, and emotional awareness (Zeman et al., 2020; Dawes et al., 2022; Monzel et al., 2024b), although direct structure-function relationships remain to be tested. Future studies should therefore assess these structural features alongside objective measures of episodic memory fidelity, face recognition, and affective processing.

Second, increased FA and neurite density in the dorsal cingulum are consistent with altered organization of a cognitive-control pathway. One possibility is that altered top-down regulation affects the conscious accessibility of internally generated representations, in a manner analogous to models of neurofunctional and psychiatric disorders (Mary et al., 2020; Kennis et al., 2015; Sasikumar & Strafella, 2021). Another is that these changes reflect compensatory reliance on semantically guided or non-visual control processes in aphantasia. Distinguishing between these possibilities will require task-based and longitudinal evidence.

Third, at the network level, aphantasia was associated with reduced local segregation in left anterior insula and altered connectivity metrics in frontal association regions, together with increased inter-network hubness in right dmPFC. These findings point to altered organization of attentional, salience, and default-mode systems. Given prior proposals implicating anterior insula and frontal control regions in the top-down regulation of internally generated representations, these network-level differences may affect conscious imagery construction at one or multiple stages (Liu, 2026): mediating inward attentional shifts (Nobre & Gresch, 2025), binding multiple features (Treisman, 2007), and gating conscious access (Huang et al., 2021). Along with the anterior cingulate cortex, insular alterations have also been proposed as a mechanism underlying some cases of acquired aphantasia, via reduced interoceptive and self-referential integration (Scholz et al., 2026).

Fourth, cortical thickness was selectively reduced in the left aPFC, a region which was found to be functionally disconnected in aphantasia (Liu & Bartolomeo, 2025), and whose grey-matter volume has previously been linked to imagery vividness in visualizers (Bergmann et al., 2016b). By contrast, cortical thickness was increased in medial temporal limbic regions, including the entorhinal (Moser et al., 2014) and retrosplenial cortex (Vann et al., 2009), critical for spatial navigation, episodic memory and imagined scene construction. Together with the tract and network findings, this pattern localizes the observed structural differences primarily to fronto-temporal and cingulate systems.

Across all four analyses, this structural dissociation is consistent with a recent multiple-stage model of conscious imagery and aphantasia (Liu, 2026), in which imagery experience depends on at least three broad operations: generation of internal visual representations, integration of distributed visual features into coherent object- or scene-like visual content (see also Scholz et al., 2025; Arcangeli & Bartolomeo, 2025), and amplification of that content for conscious access. Within this framework, the observed connectivity differences may affect integration and conscious access while leaving much of the sensory representational substrate relatively preserved.

In summary, this study provides the first comprehensive characterization of structural brain differences in congenital aphantasia, revealing a selective pattern centered on higher-order fronto-temporal and cingulate systems. By contrast, we found little evidence for major structural differences in early visual cortex or in its principal white-matter tracts, although the status of local U-shaped fibers (Guevara et al., 2020) remains to be determined. These findings argue that the structural basis of conscious imagery lies not in sensory representations themselves, but in the higher-order integrative and control systems that enable access to them.

Methods

Participants

Eighteen individuals with self-reported congenital aphantasia (mean ± SD age, 31.79 ± 12.11 years; 11 female) and 18 typical visualizers (32.29 ± 9.53 years; 15 female) were recruited. Ten aphantasic participants and five visualizers had also participated in a previous 7 Tesla fMRI study of mental imagery (Liu et al., 2025b). The sample size of 18 per group reflects the recruitment constraints inherent to congenital aphantasia, which affects an estimated 2–4% of the population. This is comparable to landmark structural MRI studies of analogous congenital conditions (Thomas et al., 2009). The convergent multimodal design and consistently large effect sizes (Cohen's d > 1.0) further support the sensitivity of the reported findings.

All participants completed the Vividness of Visual Imagery Questionnaire (VVIQ; Marks, 1973), which comprises 16 items rated from 1 ("no image at all") to 5 ("perfectly clear and as vivid as normal vision"), for a total score ranging from 16 to 80. All aphantasic participants had VVIQ scores below 22 (mean ± SD, 17.06 ± 2.10), and a lifelong absence of voluntary visual imagery was confirmed by structured interview conducted by J.L. No widely accepted objective diagnostic marker of congenital aphantasia is currently available (Bouyer et al., 2025). All visualizers had VVIQ scores above 50 (63.67 ± 9.54). All participants were right-handed, had normal or corrected-to-normal vision and hearing, and reported no history of neurological or psychiatric disorder. The two groups were matched for age, sex, and years of education. The study was approved by the institutional review board of INSERM (protocol C13-41), and all participants provided written informed consent in accordance with the Declaration of Helsinki.

Image acquisition

All MRI data were acquired at the Centre de NeuroImagerie de Recherche (CENIR), Paris Brain Institute, on a Siemens 3T MAGNETOM Prisma scanner equipped with a 64-channel head coil. Diffusion-weighted imaging (DWI) data were acquired across four sequences (398 volumes total), using two phase-encoding directions (anterior-to-posterior, AP; posterior-to-anterior, PA), each with 98-99 uniformly distributed diffusion directions at two b values (b = 1,500 and 3,000 s/mm²). In addition, 28 non-diffusion-weighted (b = 0) volumes were acquired. High-resolution T1-weighted anatomical images were acquired in the same session (TR = 2,400 ms, TE = 2.22 ms, flip angle = 9°).

DWI and T1-weighted data processing

Grey matter/white matter segmentation

Anatomical T1-weighted images were preprocessed using the DeepPrep 25.1.0 pipeline (Ren et al., 2025), which combines deep-learning-based and established neuroimaging tools for robust

tissue segmentation. Processing included motion correction using FreeSurfer (v7.2.0; RRID: SCR_001847), bias-field correction using SimpleITK (v2.3.0), skull stripping and tissue segmentation using FastSurferCNN from FastSurfer (v1.1.0), and cortical surface reconstruction using FastCSR (v1.0.0). The preprocessed T1-weighted image served as the anatomical reference for subsequent diffusion-weighted analyses.

Diffusion-weighted image preprocessing

Diffusion-weighted images were preprocessed using QSIPrep v1.0.2 (Cieslak et al., 2021) with default parameters. Sequences with the same phase-encoding polarity were merged before preprocessing. The pipeline included MP-PCA denoising, Gibbs-ringing removal, intensity normalization across b = 0 images, N4 bias-field correction, and correction for head motion, eddy currents, and susceptibility-induced distortion using FSL's eddy and TOPUP, with outlier replacement enabled. Final interpolation used Jacobian modulation, and the data were resampled to each participant's native AC-PC space at 1.25 mm isotropic resolution.

Fiber reconstruction

Preprocessed DWI data were reconstructed using QSIRecon v1.0.1 (Cieslak et al., 2021) with the *mrtrix_multishell_msmt_ACT-hsvs* workflow. Brain masks from *antsBrainExtraction* were used throughout reconstruction. Multi-tissue response functions were estimated using the Dhollander algorithm, and white-matter fiber orientation distributions (FODs) were computed using multi-tissue constrained spherical deconvolution and intensity-normalized with *mtnormalize*. These FODs were used for tractography, whereas anatomical constraints were derived from T1-weighted hybrid surface-volume segmentation (hsvs) to generate five-tissue-type images. Whole-brain tractography was then performed using anatomically constrained tractography (ACT), with streamlines constrained to initiate and terminate at the grey-white matter interface. SIFT2 was applied to estimate per-streamline weights, which were retained for subsequent bundle segmentation and connectivity analyses.

Bundle segmentation

White-matter fascicles of interest were identified using pyAFQ (Kruper et al., 2021) within QSIRecon, following the *mrtrix_multishell_msmt_pyafq_tractometry* workflow without re-running ACT. In addition to the standard major fiber bundles included in pyAFQ, several custom tracts were defined for hypothesis-driven analyses using region-of-interest (ROI) definitions from prior studies. These included superior longitudinal fasciculus subdivisions (Thiebaut de Schotten et al., 2011) (SLF I, II, IIIi) and ventral cingulum pathways containing posterior cingulate and medial temporal projections (Warrington et al., 2020). All ROIs were transformed into each participant's native AC-PC space. Association fibers were constrained to remain within the hemisphere, and bundle refinement was performed using iterative outlier removal and streamline clustering.

Tractometry

For each segmented bundle, tract profiles were extracted by resampling streamlines to 100 equidistant nodes using the pyAFQ workflow. Node-wise diffusion and microstructural metrics were then computed as weighted averages across streamlines, including fractional anisotropy (FA), mean diffusivity (MD), radial diffusivity (RD), axial diffusivity (AD), and neurite orientation dispersion and density imaging (NODDI) parameters. FA is sensitive to overall fiber organization but does not distinguish between true microstructural differences and geometric effects such as crossing, fanning, or bending fibers (Sotiropoulos & Zalesky, 2019). To aid interpretation, we therefore also quantified NODDI-derived neurite density index (NDI) and orientation dispersion index (ODI), which provide more specific estimates of tissue microstructure (Zhang et al., 2012). NODDI was fitted using the AMICO implementation in QSIRecon, with parallel diffusivity fixed at 1.7×10^{-3} mm²/s and isotropic diffusivity at 3.0×10^{-3} mm²/s. NDI and ODI values were computed at each node along the tract profiles. Prior to group-level statistical analysis, tract profiles were smoothed along the tract using a 3-node full-width at half maximum (FWHM) kernel.

fROI fiber tracking

Four ROIs from a core mental imagery network were extracted from a previous 7 Tesla fMRI study of aphantasia (Liu et al., 2025b) : the fusiform imagery node (FIN), left anterior prefrontal cortex (aPFC), left orbitofrontal cortex (OFC), and right anterior intraparietal sulcus (aIPS). In aphantasic individuals, these ROIs showed altered functional connectivity or activity, motivating the analysis of their white-matter connections. Group-level ROIs were transformed into each participant's native AC-PC space. To avoid overestimation of ROI size, volumetric fROIs were projected onto the cortical surface using FreeSurfer (RRID:SCR_001847) and mapped to the grey-white matter interface (Meisler et al., 2024). The streamline search space was defined as the intersection between the projected fROIs and the grey-white matter interface mask. Streamlines intersecting this region were identified using MRtrix3 (mrcalc) with the default radial endpoint search radius of 2 mm. For each ROI, streamlines were extracted and quantified in native space using two indices: the proportion of streamlines assigned to each anatomically defined bundle, and the number of streamlines normalized by ROI volume. Bundles of interest were defined according to the anatomical location of each ROI. For the FIN, these included the long segment of the arcuate fasciculus (AF), posterior arcuate fasciculus (pAF), inferior fronto-occipital fasciculus (IFOF), inferior longitudinal fasciculus (ILF), and vertical occipital fasciculus (VOF). For the left aPFC and left OFC, the bundles of interest were the IFOF and uncinate fasciculus (UF). For the right aIPS, the bundles of interest were the pAF and SLF I, II, and III. Group differences were assessed using both normalized streamline counts and bundle-wise streamline proportions.

Graph-theoretic network analysis

Cortical parcellation was performed using the Schaefer 400-parcel, 17-network atlas (Schaefer et al., 2018), registered from template space to individual DWI space via T1-weighted normalization. Structural connectivity between parcel pairs was estimated by intersecting the endpoints of SIFT2-weighted streamlines with parcel ROIs. A streamline was assigned to a parcel pair if both endpoints fell within a 2 mm radius sphere centered on the corresponding parcel centroids. Connection strength was defined as the sum of SIFT2 weights across all streamlines linking a parcel pair, normalized by parcel volume to account for region-size differences. Symmetric adjacency matrices were generated for each participant and thresholded using consensus thresholding (de Reus & van den Heuvel, 2013), such that connections absent in at least 50% of participants in either group were set to zero. Graph-theoretic metrics were computed on weighted, undirected networks using the Brain Connectivity Toolbox (Rubinov & Sporns, 2010) in MATLAB R2024b (RRID: SCR_001622). The metrics examined were node strength, clustering coefficient, global efficiency, betweenness centrality, and participation coefficient. To assess robustness across parcellation schemes, analyses were repeated using the Schaefer 100-parcel and Gordon 333-parcel functional atlases.

Surface extraction and surface-based morphometry

Surface-based morphometry was conducted using the Computational Anatomy Toolbox (CAT12) implemented in SPM12 (Gaser et al., 2024). Cortical thickness was estimated using CAT12's default projection-based method, which computes the distance between the inner cortical surface (grey-white matter boundary) and the outer cortical surface (grey matter-cerebrospinal fluid boundary). Individual cortical thickness maps were smoothed with a 12 mm FWHM Gaussian kernel and resampled to the 32k surface before statistical analysis. Whole-brain statistical analysis was performed in CAT12. The resulting t maps were then used as input for Neurosynth (RRID:SCR_006798) decoding in BrainStat to compute Pearson correlations with meta-analytic activation maps. The 20 strongest term correlations were retained for visualization.

To examine the visual cortex specifically, we used two atlas-based approaches. Grey matter volume was extracted from occipital regions using the Neuromorphometrics atlas (RRID: SCR_005656) as implemented in CAT12, and cortical thickness was extracted from early visual regions using the HCP-MMP Glasser atlas. Grey matter volume measures were normalized by total intracranial volume to account for individual differences in head size.

Quantification of NODDI in grey matter and white-matter tracts

To complement cortical thickness analysis with microstructurally interpretable measures, NODDI metrics were also quantified in cortical grey matter. NODDI was fit using the AMICO implementation described above, and volumetric NDI and ODI maps were sampled onto the cortical surface uniformly between the pial and white matter boundaries using Nilearn's *vol_to_surf* function, then resampled to the fsLR-32k surface. The NODDI model has been

validated against histological measures in both grey and white matter, and ODI has been shown to be sensitive to microstructural changes including demyelination in post-mortem tissue.

Logistic regression

To identify parsimonious combinations of structural features associated with group membership while accounting for shared variance across measures, we used a two-step modelling approach. First, LASSO-penalized logistic regression (*glmnet* package in R) was applied separately within each metric domain: tractometry FA, cortical thickness, and graph-theoretic measures. Predictors comprised mean FA across significant tract nodes, cortical thickness values, and graph metrics identified in prior analyses. Separate domain-specific models were used to limit the predictor-to-sample ratio given the modest sample size (N = 36). Model selection was performed using 5-fold cross-validation with approximately balanced class distributions, and the regularization parameter was chosen using the 1-standard-error rule ($\lambda_1$se). In a second step, variables retained by LASSO were entered into Firth's penalized logistic regression (Firth, 1993) to obtain bias-reduced parameter estimates and odds ratios, thereby reducing small-sample bias and separation problems. Variables that caused non-convergence were excluded using a p-value-based stepwise procedure.

Statistical analysis

Bayes factors

To quantify evidence for and against group differences, we computed Bayes factors (BFs) in JASP (https://jasp-stats.org/; RRID: SCR_015823) using Bayesian t tests and Mann-Whitney U tests with default priors on effect size. BF10 > 3 was taken as moderate evidence for a group difference, whereas BF01 > 3 (equivalently, BF10 < 1/3) was taken as moderate evidence for the absence of a group difference (Keysers et al., 2020).

Tractometry

Group differences in tract profiles were assessed using one-dimensional threshold-free cluster enhancement (TFCE; E = 0.5, H = 2) along tract nodes, implemented in PALM with 10,000 permutations. TFCE identifies spatially extended effects without requiring an arbitrary cluster-forming threshold and assigns a family-wise-error-corrected p value to each node. To reduce spurious findings, only clusters comprising at least 3 consecutive significant nodes were reported. Statistical significance was defined as $p < 0.05$, FWE-corrected.

Graph-theoretic network analysis

Group differences in node-level graph-theoretic metrics were assessed using the non-parametric Brunner-Munzel test, with false-discovery-rate correction applied across all nodes within each metric. The same test was also used to assess group differences in SIFT2-weighted streamline counts between ROI pairs.

fROI fiber tracking

Group differences in filtered tracts within each fROI were assessed using the Mann-Whitney test, because the data included a substantial number of zero values and did not satisfy the assumptions of parametric testing.

Surface-based morphometry

Statistical analyses of cortical thickness were performed in BrainStat (Larivière et al., 2023) using a linear model with a one-sided test and false discovery rate (FDR) correction. To identify spatially coherent clusters of significant vertices, hierarchical density-based spatial clustering of applications with noise (HDBSCAN) was applied to the x-, y-, and z-coordinates of significant vertices, grouping those that were spatially adjacent. Only clusters containing at least 30 vertices were retained for interpretation, and Desikan killiany atlas was used to identify the anatomical region of surviving clusters.

Grey matter NODDI

Cortical grey matter NODDI metrics were compared between groups using random field theory as implemented in BrainStat (Larivière et al., 2023). Cluster-level inference used a cluster-forming threshold of $p < 0.05$ and a cluster-significance threshold of $p < 0.05$, FWE-corrected.

Contributions

Jianghao Liu: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Visualization, Writing – original draft, Writing – review & editing; **Yusaku Takamura**: Conceptualization, Data curation, Formal analysis, Methodology, Software, Visualization, Writing – original draft, Writing – review & editing; **Romain Delsanti**: Data curation, Writing – review & editing; **Laurent Cohen:** Resources, Writing – review & editing; **Paolo Bartolomeo:** Conceptualization, Funding acquisition, Project administration, Resources, Supervision, Writing – review & editing.

Materials & correspondence

Requests for materials and correspondence should be addressed to the lead contact Jianghao Liu (jianghaolouisliu@gmail.com).

Acknowledgement

We thank Dounia Hajhajate, Hanna Kavaliova, and Diane Lamarle for their help in collecting part of the data, Minye Zhan, Elena Grosso, and Benoît Béranger for their suggestions regarding the data analysis and Michel Thiebaut de Schotten for commenting on an early version of our results. The work of Jianghao Liu is supported by specific funding from Dassault Systèmes. The work of Yusaku Takamura is supported by JSPS KAKENHI Grants JP24KK0296 and JST-CRESTJPMJCR23P1. The work of Paolo Bartolomeo is supported by the Agence Nationale de la Recherche through ANR-16-CE37-0005 and ANR-10-IAIHU-06, by the Fondation pour la Recherche sur les AVC through FR-AVC-017, and by the Paris Brain Institute grant ViBER - Vision Beyond External Reality.

Supplementary materials

Supplementary Figure S1. Overview of the analysis pipeline. Three data streams were processed in parallel. Diffusion MRI data underwent preprocessing and fiber reconstruction to generate whole-brain tractography. Anatomical T1-weighted images were processed for grey matter/white matter (GM/WM) segmentation and cortical surface extraction. Functional ROIs from a previous 7T fMRI study of mental imagery (Liu et al., 2025b) were registered to each participant's native space. These inputs were then used for four complementary analyses: **(1) tractometry**, in which white-matter bundles were segmented and microstructural properties, including fractional anisotropy (FA), were quantified along tract profiles; **(2) fROI-based tractography**, in which functional ROIs were projected to the GM/WM interface to identify streamlines connecting regions of the core imagery network; **(3) graph-theoretic network analysis**, in which cortical parcellation and SIFT2-weighted streamline counts were used to construct structural connectivity matrices and derive node-level network metrics; and **(4) surface morphometry**, in which cortical thickness was estimated from the extracted cortical surfaces and compared between groups across the whole brain.

Axial and radial diffusivity and NODDI control analyses for tractometry findings

To clarify the potential biological basis of the FA differences, we examined axial diffusivity (AD) and radial diffusivity (RD), often interpreted as being more sensitive to axonal and myelin-related properties, respectively (Song et al., 2002), together with neurite orientation dispersion and density imaging (NODDI) metrics, including neurite density index (NDI) and orientation dispersion index (ODI) (Zhang et al., 2012), at all segments showing significant FA effects. Full statistics are reported in Table S1.

Uncinate fasciculus and posterior interparietal corpus callosum. FA reductions in the bilateral uncinate fasciculus and posterior interparietal corpus callosum were accompanied by marginal RD increases at largely overlapping segments (all $p < 0.1$, FWE-corrected), a pattern consistent with altered myelin-related microstructure. No significant group differences in NDI or ODI were observed at the same segments, suggesting that these FA reductions are unlikely to be explained by fiber-geometry effects such as crossing fibers.

Bilateral dorsal cingulum. The FA increase in the bilateral dorsal cingulum was accompanied, in the left hemisphere, by a significant RD decrease and AD increase, consistent with altered microstructural organization rather than myelin degradation. A significant increase in NDI was observed at the same segments showing elevated FA, with no change in ODI, suggesting that the FA increase is more likely to reflect greater neurite density than reduced fiber dispersion.

Marginal findings. Marginal FA reductions were observed in the right SLF-II (Fig. S3A; $p = 0.082$) and right ventral cingulum ($p = 0.079$). Both tracts also showed increased ODI at the same segments (Fig. S3B), suggesting that these marginal effects may be influenced by local fiber geometry rather than robust microstructural differences. We therefore did not interpret these findings further.

Supplementary Figure S2. NODDI microstructural profiles in white-matter tracts showing significant FA differences. (A) Neurite density index (NDI) tract profiles. **(B)** Orientation dispersion index (ODI) tract profiles. Profiles are shown for the bilateral uncinate fasciculus, posterior interparietal corpus callosum, and bilateral dorsal cingulum, the tracts identified in the main tractometry analysis. Red lines indicate aphantasic individuals and blue lines indicate visualizers. Thin lines show individual participant profiles; thick lines show group means ± SEM. Shaded grey regions and black bars indicate significant between-group differences (p < 0.05, FWE-corrected).

Supplementary Figure S3. Marginal FA reductions in the right SLF-II and right ventral cingulum are accompanied by increased orientation dispersion. (A) Fractional anisotropy (FA) tract profiles for the right SLF-II (p = 0.082, FWE-corrected) and right ventral cingulum (p = 0.079, FWE-corrected), with representative tract reconstructions shown below. Shaded grey regions and black bars indicate segments showing marginal between-group differences. Red lines indicate aphantasic individuals and blue lines indicate visualizers; thin lines show individual participant profiles and thick lines show group means ± SEM. **(B)** Orientation dispersion index (ODI) profiles for the same tracts. In both tracts, marginal FA reductions overlapped with increased ODI in aphantasic individuals, suggesting that these effects may be influenced by local fiber geometry rather than robust microstructural differences. These findings were therefore not interpreted further.

Supplementary Figure S4. White-matter connections of the core imagery network. (A) White-matter streamlines associated with each functional ROI of the core imagery network, shown on sagittal MRI sections: left fusiform imagery node (FIN), left orbitofrontal cortex (OFC), left anterior prefrontal cortex (aPFC), and right anterior intraparietal sulcus (aIPS). **(B)** Streamline counts normalized by ROI volume for each functional ROI. The FIN was connected primarily by the long segment of the arcuate fasciculus (AF), posterior arcuate fasciculus (pAF), and vertical occipital fasciculus (VOF). The left OFC was connected by both the inferior fronto-occipital fasciculus (IFOF) and uncinate fasciculus (UF), whereas the left aPFC was connected mainly by the IFOF. No direct structural connection was identified between the FIN and left aPFC in this analysis. The right aIPS was connected predominantly by the third branch of the superior longitudinal fasciculus (SLF-III). A nominal increase in normalized streamline count was observed in aphantasic individuals for the left AF associated with the FIN (uncorrected p = 0.039, U = 229, Cliff's d = -0.41), but this effect did not survive correction for multiple comparisons. Bayes factor analysis provided moderate evidence for no group difference in VOF streamline count associated with the FIN (BF = 0.32). **(C)** Proportion of streamlines assigned to each tract of interest for each ROI. Light bars indicate visualizers and dark bars indicate aphantasic individuals. Error bars show SEM and dots indicate individual participants. IFOF, inferior fronto-occipital fasciculus; ILF, inferior longitudinal fasciculus; AF, arcuate fasciculus long segment; pAF, posterior arcuate fasciculus; VOF, vertical occipital fasciculus; UF, uncinate fasciculus; SLF-I/II/III, superior longitudinal fasciculus branches I, II, and III.

Control analysis on atlas selection in network analysis

A. Gordon 333 Regions

Supplementary Figure S5. Control analyses confirm the robustness of network findings across parcellation schemes. (A) Gordon 333-region atlas. Convergent group differences were observed across graph-theoretic metrics, with aphantasic individuals showing reduced clustering coefficient in the left anterior insula (Vis. > Aph., blue) and increased participation coefficient in the right dorsomedial prefrontal cortex (dmPFC; Aph. > Vis., dark red). **(B)** Schaefer-Yeo 100-region atlas. Aphantasic individuals showed increased participation coefficient in bilateral dmPFC (Aph. > Vis.) and reduced participation coefficient in right lateral prefrontal cortex (Vis. > Aph.). **(C)** White-matter tracts connected to the left anterior insula node in the Schaefer-Yeo 400-region atlas. The inferior fronto-occipital fasciculus (IFOF) and arcuate fasciculus (AF) converge on the left anterior insula, the node showing reduced clustering coefficient in the main analysis. The left anterior prefrontal cortex (aPFC) and fusiform imagery node (FIN) are linked by distinct white-matter pathways, primarily IFOF and AF, respectively. Dark red indicates Aph. > Vis. and blue indicates Vis. > Aph. All group differences shown in panels A and B were significant at $p < 0.05$, FDR-corrected. The left frontal eye field (FEF) was not identified consistently across atlases, consistent with uncertainty in its parcellation boundaries.

Control analysis on NODDI gray matter

Supplementary Figure S6. Cortical NODDI microstructural differences between aphantasic individuals and visualizers. Cortical NODDI metrics were compared between groups using random field theory as implemented in BrainStat, with cluster-level FWE correction (p < 0.05). Aphantasic individuals showed increased neurite density index (NDI) in bilateral medial temporal regions, including the parahippocampal gyrus and entorhinal cortex, as well as in the left precuneus, and reduced NDI in the left dorsomedial prefrontal cortex. Orientation dispersion index (ODI), isotropic volume fraction (ISOVF), and mean diffusivity (MD) also showed significant group differences. Regions of reduced MD overlapped spatially with areas of increased NDI in medial temporal cortex, a pattern consistent with denser neurite packing and lower overall diffusivity in these regions.

Supplementary Table S1. Tract-level diffusion and NODDI differences between aphantasic individuals and visualizers. Significant group differences in tract-profile diffusion tensor imaging (DTI) and neurite orientation dispersion and density imaging (NODDI) metrics are shown for tracts identified in the tractometry analysis. Values indicate the start and end nodes of each significant cluster, followed by the FWE-corrected p value, test statistic, and Cohen's d in parentheses. Direction of effect is indicated separately for each metric column (Vis. > Aph. or Aph. > Vis.). Only clusters meeting the reporting threshold of at least 3 consecutive significant nodes are listed. For the main analyses, significance was defined as p < 0.05, FWE-corrected; marginal effects (p < 0.1, FWE-corrected) are reported in the text but are not included in this table.

| Tract | DTI | | | NODDI | |
	Fractional anisotropy (FA)	Axial diffusivity (AD)	Radial diffusivity (RD)	Neurite density index (NDI)	Orientation dispersion index (ODI)
	Vis > Aph		*Aph > Vis*		
right UF	77-84 (0.018; 3.43; 1.14)		78-83 (0.061; 2.50; 0.83)		
left UF	75-77 (0.048; 2.89; 0.96)		74-78 (0.085; 2.54; 0.85)		
interparietal corpus callosum	47-50 (0.039; 3.01; 1.00)		47-52 (0.071; 2.71; 0.90)		
	Aph > Vis	*Aph > Vis*	*Vis > Aph*	*Aph > Vis*	*Vis > Aph*
right dorsal cingulum	46-69(0.019; 2.85; 0.95)		40-76 (0.016; 2.96; 0.99)	44-63 (0.046; 2.17; 0.72)	
left dorsal cingulum	39-68 (0.003; 3.95; 1.32)	52-61 (0.068; 2.51; 0.84)	30-69 (0.004; 3.78; 1.26)	59-62 (0.047; 2.50; 0.83)	59-61(0.041; 3.00; 1.00)

Supplementary Table S2. Nodal graph-theoretic network differences between aphantasic individuals and visualizers. Significant group differences in nodal graph-theoretic metrics are reported for node strength, clustering coefficient, and participation coefficient. Group comparisons were performed using the Brunner-Munzel test with false discovery rate (FDR) correction within each metric. MNI coordinates (x, y, z) correspond to parcel centroids. Network affiliations are defined according to the Schaefer 400-parcel, 17-network atlas. **Vis. > Aph.** indicates lower metric values in aphantasic individuals, whereas **Aph. > Vis.** indicates higher metric values in aphantasic individuals. **B-M statistic** denotes the Brunner-Munzel statistic, and **Cliff's delta** the effect size. Significance threshold: $p < 0.05$, FDR-corrected.

Metric	Node (Network)	MNI coordinates (x, y, z)	B-M statistic	p (FDR corrected)	Cliff's delta
Vis. > Aph.					
Strength	Left FEF (DAN B)	25, -1, 55	-4.33	0.050	0.65
Clustering coefficient	Left anterior insula (Salience/VAN A)	-33, 18, 6	-7.18	<0.001	0.77
Clustering coefficient	Left dlPFC (Default B)	-42, 7, 48	-4.93	0.004	0.66
Participation coefficient	Right somatomotor (SomMot B)	9, -40, 68	-4.05	0.038	0.59
Aph. > Vis.					
Participation coefficient	Right dmPFC (Default A)	7, 42, 4	4.64	0.024	-0.66
Participation coefficient	Right dmPFC (Default A)	7, 54, 13	4.19	0.038	-0.63

Supplementary Table S3. Gray matter volume in occipital cortical regions. Regions were defined using the Neuromorphometrics atlas in CAT12. Gray matter volumes were normalized by total intracranial volume (TIV). No significant group differences were observed in any occipital cortical region. Bayes factors provided moderate evidence for an equal volume between the two groups in the left and right calcarine cortices. Values are reported as mean ± SD.

Region	Aphantasics ($\times 10^{-4}$ mm³/TIV)	Visualizers ($\times 10^{-4}$ mm³/TIV)	BF+0
Right calcarine cortex	20 ± 2.26	20 ± 2.70	**0.21***
Left calcarine cortex	20 ± 2.71	20 ± 2.35	**0.22***
Right inferior occipital gyrus	40 ± 2.50	40 ± 3.15	0.35
Left inferior occipital gyrus	30 ± 2.97	30 ± 3.74	0.33
Right lingual gyrus	50 ± 4.74	50 ± 3.72	0.75
Left lingual gyrus	50 ± 2.87	50 ± 3.65	0.59
Right middle occipital gyrus	30 ± 3.50	30 ± 2.84	**0.23***
Left middle occipital gyrus	40 ± 4.50	40 ± 2.94	**0.33***
Right occipital pole	20 ± 2.94	20 ± 2.67	0.42
Left occipital pole	20 ± 2.18	20 ± 2.21	**0.28***
Right occipital fusiform gyrus	20 ± 2.10	20 ± 2.13	0.92
Left occipital fusiform gyrus	20 ± 1.89	20 ± 2.10	0.37

Note: For all Bayesian tests, the alternative hypothesis specified smaller regional volume in visualizers than in aphantasics. ***BF10 < 0.33**, indicating moderate evidence for the null hypothesis.

Supplementary Table S4. Cortical thickness in early visual areas in aphantasic individuals and visualizers. Mean cortical thickness values (mm ± SD) for atlas-defined early visual regions are shown for the aphantasia and visualizer groups. Regions of interest were defined using the HCP-MMP atlas. No significant group differences were observed in any region. Bayes factors indicated moderate evidence for the absence of a group difference in 4 of the 6 regions ($BF_{+0} < 0.33$). l, left hemisphere; r, right hemisphere.

Region	Aphantasics (mm)	Visualizers (mm)	BF+0
lV1	1.83 ± 0.11	1.86 ± 0.13	**0.21***
rV1	1.83 ± 0.10	1.82 ± 0.15	0.37
lV2	1.94 ± 0.09	1.96 ± 0.11	**0.21***
rV2	1.94 ± 0.10	1.93 ± 0.13	0.37
lV3	2.10 ± 0.10	2.11 ± 0.13	**0.30***
rV3	2.03 ± 0.12	2.07 ± 0.11	0.17

Note: For all Bayesian tests, the directional alternative specified greater cortical thickness in aphantasia than in visualizers. $\text{*BF}_{10} < 0.33$.

Supplementary Table S5. Cortical thickness differences between aphantasic individuals and visualizers. Significant group differences in cortical thickness are reported for clusters comprising more than 30 vertices. Group comparisons were performed using a linear model with FDR correction. aPFC: anterior prefrontal cortex; PoCG: Post central gyrus; PHG: parahippocampal gyrus; EC: Entorhinal cortex; ACC: Anterior cingulate cortex; RSC: retrosplenial cortex; PCC: Posterior cingulate cortex.

Region	Cluster size	Peak t-value	Peak p-value (FDR)	MNI coordinates (x, y, z)	Cohen's D
Vis. > Aph.					
Left aPFC	52	4.68	0.000	-36, 54, 4	1.56
Left PoCG	49	3.73	0.005	-64, -14, 22	1.24
Aph. > Vis.					
Left PHG/EC	264	6.83	0.000	-25, -19, -23	2.28
Left ACC	44	3.61	0.005	-4, 18, 23	1.20
Right RSC	294	3.52	0.007	17, -53, 6	1.17
Right EC/IFG	145	4.49	0.000	32, -6, -38	1.50
Right PCC	110	4.41	0.001	5, -18, 29	1.47

Multiple regression model

We used a two-step modelling approach. First, LASSO-penalized logistic regression was applied separately within each metric domain to identify the most informative predictors. Second, the variables retained by LASSO were entered into Firth's penalized logistic regression to obtain bias-reduced parameter estimates and odds ratios.

For the tractometry model, LASSO logistic regression retained the left dorsal cingulum, right uncinate fasciculus, and posterior interparietal corpus callosum at $\lambda_1$se. In the Firth model, higher FA in the right uncinate fasciculus was marginally associated with lower odds of aphantasia (beta = −1.00; OR = 0.36, 95% CI = 0.07-1.00; p = 0.051), as was higher FA in the posterior interparietal corpus callosum (beta = −1.70; OR = 0.18, 95% CI = 0.02-0.61; p = 0.003). Higher FA in the left dorsal cingulum was significantly associated with increased odds of aphantasia (beta = 1.79; OR = 6.03, 95% CI = 1.79-51.40; p < 0.001). These multivariate results were broadly consistent with the tract-level univariate findings. Area under the curve (AUC) for this model was 0.96.

For the cortical thickness model, we selected the left anterior prefrontal cortex (aPFC) and right retrosplenial cortex (RSC) clusters, which were the largest clusters showing negative and positive effects, respectively, to avoid complete separation in the logistic regression model. LASSO logistic regression retained both clusters. In the final Firth model, reduced left aPFC thickness (beta = −1.81; OR = 0.16, 95% CI = 0.03–0.48; p < 0.001) and greater right RSC

thickness (beta = 1.52; OR = 4.58, 95% CI = 1.54–27.33; p = 0.004) were associated with higher odds of aphantasia. The area under the curve (AUC) for the cortical thickness model was 0.93. In addition, we examined the single-variable effect of the left parahippocampal gyrus/entorhinal cortex (PHG/EC) cluster, which showed the largest effect size. Greater left PHG/EC thickness contributed to group discrimination (beta = 2.99; OR = 19.90, 95% CI = 4.27–313.68; p < 0.001). The AUC for this single-variable model was 0.95.

For the graph-theoretical metrics model, given concerns that including all candidate predictors would result in an overly complex model relative to the sample size, we excluded the participation coefficient in the right dorsomedial prefrontal cortex (dmPFC, region 3) and the somatomotor cortex before applying LASSO, based on overlap among regions and their relatively lower theoretical relevance to aphantasia. LASSO regression retained four predictors spanning three network measures. In the final Firth model, a lower clustering coefficient in the left insula (β = −2.41; OR = 0.09, 95% CI = 0.001–0.47; p = 0.002), a lower clustering coefficient in the left prefrontal cortex (β = −0.89; OR = 0.41, 95% CI = 0.007–2.17; p = 0.336), and lower node strength in the left frontal eye field (FEF) (β = −1.32; OR = 0.27, 95% CI = 0.029–0.84; p = 0.022) were associated with higher odds of aphantasia. A higher participation coefficient in the right dmPFC was also associated with higher odds of aphantasia (β = 1.81; OR = 6.14, 95% CI = 1.29–216.11; p = 0.021), consistent with the corresponding univariate result. The area under the curve (AUC) for the graph-theoretical metrics model was 0.99.

References

Arcangeli, M., & Bartolomeo, P. (2025). The aphantasia paradox: A Sartrean update. *Brain*, *148*(9), 3023–3025. https://doi.org/10.1093/brain/awaf233

Bartolomeo, P., Liu, J., & Spagna, A. (2026). The Fusiform Imagery Node: Where vision meets concepts in the left temporal lobe. *Neuropsychologia*, *224*, 109398. https://doi.org/10.1016/j.neuropsychologia.2026.109398

Bergmann, J., Genç, E., Kohler, A., Singer, W., & Pearson, J. (2016a). Neural Anatomy of Primary Visual Cortex Limits Visual Working Memory. *Cerebral Cortex*, *26*(1), 43–50. https://doi.org/10.1093/cercor/bhu168

Bergmann, J., Genç, E., Kohler, A., Singer, W., & Pearson, J. (2016b). Smaller Primary Visual Cortex Is Associated with Stronger, but Less Precise Mental Imagery. *Cerebral Cortex*, *26*(9), 3838–3850. https://doi.org/10.1093/cercor/bhv186

Bouyer, L. N., Schwarzkopf, D. S., Saurels, B. W., & Arnold, D. H. (2025). Objective priming from pre-imagining inputs before binocular rivalry presentations does not predict individual differences in the subjective intensity of imagined experiences. *Cognition*, *256*, 106048. https://doi.org/10.1016/j.cognition.2024.106048

Cabbai, G., Racey, C., Simner, J., Dance, C., Ward, J., & Forster, S. (2024). Sensory representations in primary visual cortex are not sufficient for subjective imagery. *Current Biology*, *34*(21), 5073-5082.e5. https://doi.org/10.1016/j.cub.2024.09.062

Chang, S., Zhang, X., Cao, Y., Pearson, J., & Meng, M. (2025). Imageless imagery in aphantasia revealed by early visual cortex decoding. *Current Biology*, *35*(3), 591-599.e4. https://doi.org/10.1016/j.cub.2024.12.012

Cieslak, M., Cook, P. A., He, X., Yeh, F.-C., Dhollander, T., Adebimpe, A., Aguirre, G. K., Bassett, D. S., Betzel, R. F., Bourque, J., Cabral, L. M., Davatzikos, C., Detre, J. A., Earl, E., Elliott, M. A., Fadnavis, S., Fair, D. A., Foran, W., Fotiadis, P., … Satterthwaite, T. D. (2021). QSIPrep: An integrative platform for preprocessing and reconstructing diffusion MRI data. *Nature Methods*, *18*(7), 775–778. https://doi.org/10.1038/s41592-021-01185-5

Coad, B. M., Postans, M., Hodgetts, C. J., Muhlert, N., Graham, K. S., & Lawrence, A. D. (2020). Structural connections support emotional connections: Uncinate Fasciculus microstructure is related to the ability to decode facial emotion expressions. *Neuropsychologia*, *145*, 106562. https://doi.org/10.1016/j.neuropsychologia.2017.11.006

Corbetta, M., & Shulman, G. L. (2002). Control of goal-directed and stimulus-driven attention in the brain. *Nature Reviews Neuroscience*, *3*(3), Article 3. https://doi.org/10.1038/nrn755

Dance, C. J., Hole, G., & Simner, J. (2023). The role of visual imagery in face recognition and the construction of facial composites. Evidence from Aphantasia. *Cortex*, *167*, 318–334. https://doi.org/10.1016/j.cortex.2023.06.015

Dance, C. J., Jaquiery, M., Eagleman, D. M., Porteous, D., Zeman, A., & Simner, J. (2021). What is the relationship between Aphantasia, Synaesthesia and Autism? *Consciousness and Cognition*, *89*, 103087. https://doi.org/10.1016/j.concog.2021.103087

Dawes, A. J., Keogh, R., Robuck, S., & Pearson, J. (2022). Memories with a blind mind: Remembering the past and imagining the future with aphantasia. *Cognition*, *227*, 105192. https://doi.org/10.1016/j.cognition.2022.105192

de Reus, M. A., & van den Heuvel, M. P. (2013). The parcellation-based connectome: Limitations and extensions. *NeuroImage*, *80*, 397–404. https://doi.org/10.1016/j.neuroimage.2013.03.053

Firth, D. (1993). Bias reduction of maximum likelihood estimates. *Biometrika*, *80*(1), 27–38. https://doi.org/10.1093/biomet/80.1.27

Gaser, C., Dahnke, R., Thompson, P. M., Kurth, F., Luders, E., & the Alzheimer's Disease Neuroimaging Initiative. (2024). CAT: A computational anatomy toolbox for the analysis of structural MRI data. *GigaScience*, *13*, giae049. https://doi.org/10.1093/gigascience/giae049

Guevara, M., Guevara, P., Román, C., & Mangin, J.-F. (2020). Superficial white matter: A review on the dMRI analysis methods and applications. *NeuroImage*, *212*, 116673. https://doi.org/10.1016/j.neuroimage.2020.116673

Huang, Z., Tarnal, V., Vlisides, P. E., Janke, E. L., McKinney, A. M., Picton, P., Mashour, G. A., & Hudetz, A. G. (2021). Anterior insula regulates brain network transitions that gate conscious access. *Cell Reports*, *35*(5). https://doi.org/10.1016/j.celrep.2021.109081

Kennis, M., van Rooij, S. J. H., Tromp, D. P. M., Fox, A. S., Rademaker, A. R., Kahn, R. S., Kalin, N. H., & Geuze, E. (2015). Treatment Outcome-Related White Matter Differences in Veterans with Posttraumatic Stress Disorder. *Neuropsychopharmacology*, *40*(10), 2434–2442. https://doi.org/10.1038/npp.2015.94

Keogh, R., & Pearson, J. (2018). The blind mind: No sensory visual imagery in aphantasia. *Cortex*, *105*, 53–60. https://doi.org/10.1016/j.cortex.2017.10.012

Keogh, R., Wicken, M., & Pearson, J. (2023). *Fewer intrusive memories in aphantasia: Using the trauma film paradigm as a laboratory model of PTSD* (7zqfe_v1). PsyArXiv. https://doi.org/10.31234/osf.io/7zqfe

Keysers, C., Gazzola, V., & Wagenmakers, E.-J. (2020). Using Bayes factor hypothesis testing in neuroscience to establish evidence of absence. *Nature Neuroscience*, *23*(7), 788–799. https://doi.org/10.1038/s41593-020-0660-4

Kruper, J., Yeatman, J. D., Richie-Halford, A., Bloom, D., Grotheer, M., Caffarra, S., Kiar, G., Karipidis, I. I., Roy, E., Chandio, B. Q., Garyfallidis, E., & Rokem, A. (2021). Evaluating the Reliability of Human Brain White Matter Tractometry. *Aperture Neuro*, *1*. https://doi.org/10.52294/e6198273-b8e3-4b63-babb-6e6b0da10669

Larivière, S., Bayrak, Ş., Vos de Wael, R., Benkarim, O., Herholz, P., Rodriguez-Cruces, R., Paquola, C., Hong, S.-J., Misic, B., Evans, A. C., Valk, S. L., & Bernhardt, B. C. (2023).

BrainStat: A toolbox for brain-wide statistics and multimodal feature associations. *NeuroImage*, *266*, 119807. https://doi.org/10.1016/j.neuroimage.2022.119807

Liu, J. (2026). A neural model of conscious mental imagery and aphantasia. *Neuropsychologia*, *224*, 109392. https://doi.org/10.1016/j.neuropsychologia.2026.109392

Liu, J. (in press). Aphantasia as a window into unconscious internal processing. *Behavioral and Brain Sciences*. https://osf.io/preprints/psyarxiv/rc9y7_v1/

Liu, J., & Bartolomeo, P. (2023). Probing the unimaginable: The impact of aphantasia on distinct domains of visual mental imagery and visual perception. *Cortex*, *166*, 338–347. https://doi.org/10.1016/j.cortex.2023.06.003

Liu, J., & Bartolomeo, P. (2025). Aphantasia as a functional disconnection. *Trends in Cognitive Sciences*, *29*(11), 963–964. https://doi.org/10.1016/j.tics.2025.05.012

Liu, J., Cohen, L., Pallier, C., Dehaene, S., Zhan, M., & Bartolomeo, P. (2025a). *Face-, color-, and word-specific patches in the human orbitofrontal cortex*. bioRxiv. https://doi.org/10.1101/2025.04.22.650059

Liu, J., Zhan, M., Hajhajate, D., Spagna, A., Dehaene, S., Cohen, L., & Bartolomeo, P. (2025b). Visual mental imagery in typical imagers and in aphantasia: A millimeter-scale 7-T fMRI study. *Cortex*, *185*, 113–132. https://doi.org/10.1016/j.cortex.2025.01.013

Marks, D. F. (1973). Visual Imagery Differences in the Recall of Pictures. *British Journal of Psychology*, *64*(1), 17–24. https://doi.org/10.1111/j.2044-8295.1973.tb01322.x

Mary, A., Dayan, J., Leone, G., Postel, C., Fraisse, F., Malle, C., Vallée, T., Klein-Peschanski, C., Viader, F., de la Sayette, V., Peschanski, D., Eustache, F., & Gagnepain, P. (2020). Resilience after trauma: The role of memory suppression. *Science*, *367*(6479), eaay8477. https://doi.org/10.1126/science.aay8477

Mawtus, B., Renwick, F., Thomas, B. R., & Reeder, R. R. (2024). The Impact of Aphantasia on Mental Healthcare Experiences. *Collabra: Psychology*, *10*(1), 127416. https://doi.org/10.1525/collabra.127416

Meisler, S. L., Kubota, E., Grotheer, M., Gabrieli, J. D. E., & Grill-Spector, K. (2024). A practical guide for combining functional regions of interest and white matter bundles. *Frontiers in Neuroscience*, *18*. https://doi.org/10.3389/fnins.2024.1385847

Milton, F., Fulford, J., Dance, C., Gaddum, J., Heuerman-Williamson, B., Jones, K., Knight, K. F., MacKisack, M., Winlove, C., & Zeman, A. (2021). Behavioral and Neural Signatures of Visual Imagery Vividness Extremes: Aphantasia versus Hyperphantasia. *Cerebral Cortex Communications*, *2*(2), tgab035. https://doi.org/10.1093/texcom/tgab035

Montabes De La Cruz, B. M., Abbatecola, C., Luciani, R. S., Paton, A. T., Bergmann, J., Vetter, P., Petro, L. S., & Muckli, L. F. (2024). Decoding sound content in the early visual cortex of aphantasic participants. *Current Biology*, *34*(21), 5083-5089.e3. https://doi.org/10.1016/j.cub.2024.09.008

Monzel, M., Karneboge, J., & Reuter, M. (2024a). Affective processing in aphantasia and potential overlaps with alexithymia: Mental imagery facilitates the recognition of

emotions in oneself and others. *Biomarkers in Neuropsychiatry*, *11*, 100106. https://doi.org/10.1016/j.bionps.2024.100106

Monzel, M., Leelaarporn, P., Lutz, T., Schultz, J., Brunheim, S., Reuter, M., & McCormick, C. (2024b). Hippocampal-occipital connectivity reflects autobiographical memory deficits in aphantasia. *eLife*, *13*. https://doi.org/10.7554/eLife.94916.2

Moser, E. I., Roudi, Y., Witter, M. P., Kentros, C., Bonhoeffer, T., & Moser, M.-B. (2014). Grid cells and cortical representation. *Nature Reviews Neuroscience*, *15*(7), 466–481. https://doi.org/10.1038/nrn3766

Noble, C., Taylor, N. L., Milton, F., Fulford, J., Tan, J. B., O'Callaghan, C., Zeman, A., & Shine, J. M. (2026). Seeing through the static: Reduced imagery vividness in aphantasia is associated with impaired temporal lobe signal complexity. *Neuropsychologia*, *221*, 109322. https://doi.org/10.1016/j.neuropsychologia.2025.109322

Nobre, A. C., & Gresch, D. (2025). How the brain shifts between external and internal attention. *Neuron*, *113*(15), 2382–2398. https://doi.org/10.1016/j.neuron.2025.06.013

Parlatini, V., Itahashi, T., Lee, Y., Liu, S., Nguyen, T. T., Aoki, Y. Y., Forkel, S. J., Catani, M., Rubia, K., Zhou, J. H., Murphy, D. G., & Cortese, S. (2023). White matter alterations in Attention-Deficit/Hyperactivity Disorder (ADHD): A systematic review of 129 diffusion imaging studies with meta-analysis. *Molecular Psychiatry*, *28*(10), 4098–4123. https://doi.org/10.1038/s41380-023-02173-1

Pounder, Z., Jacob, J., Evans, S., Loveday, C., Eardley, A. F., & Silvanto, J. (2022). Only minimal differences between individuals with congenital aphantasia and those with typical imagery on neuropsychological tasks that involve imagery. *Cortex*, *148*, 180–192. https://doi.org/10.1016/j.cortex.2021.12.010

Ren, J., An, N., Lin, C., Zhang, Y., Sun, Z., Zhang, W., Li, S., Guo, N., Cui, W., Hu, Q., Wang, W., Wu, X., Wang, Y., Jiang, T., Satterthwaite, T. D., Wang, D., & Liu, H. (2025). DeepPrep: An accelerated, scalable and robust pipeline for neuroimaging preprocessing empowered by deep learning. *Nature Methods*, *22*(3), 473–476. https://doi.org/10.1038/s41592-025-02599-1

Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: Uses and interpretations. *NeuroImage, Computational Models of the Brain*, *52*(3), 1059–1069. https://doi.org/10.1016/j.neuroimage.2009.10.003

Sasikumar, S., & Strafella, A. P. (2021). The neuroimaging evidence of brain abnormalities in functional movement disorders. *Brain*, *144*(8), 2278–2283. https://doi.org/10.1093/brain/awab131

Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X.-N., Holmes, A. J., Eickhoff, S. B., & Yeo, B. T. T. (2018). Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cerebral Cortex*, *28*(9), 3095–3114. https://doi.org/10.1093/cercor/bhx179

Scholz, C. O., Monzel, M., Kvamme, T. L., Liu, J., & Silvanto, J. (2026). An integration model of mental imagery and aphantasia: Conceptual framework, neuromechanistic pathways,

and clinical implications. *Neuropsychologia*, *225*, 109401. https://doi.org/10.1016/j.neuropsychologia.2026.109401

Scholz, C. O., Monzel, M., & Liu, J. (2025). Absence of shared representation in the visual cortex challenges unconscious imagery in aphantasia. *Current Biology*, *35*(13), R645–R646. https://doi.org/10.1016/j.cub.2025.05.009

Siena, M. J., & Simons, J. S. (2024). Metacognitive Awareness and the Subjective Experience of Remembering in Aphantasia. *Journal of Cognitive Neuroscience*, *36*(8), 1578–1598. https://doi.org/10.1162/jocn_a_02120

Sokolowski, H. M., & Levine, B. (2023). Common neural substrates of diverse neurodevelopmental disorders. *Brain*, *146*(2), 438–447. https://doi.org/10.1093/brain/awac387

Song, S.-K., Sun, S.-W., Ramsbottom, M. J., Chang, C., Russell, J., & Cross, A. H. (2002). Dysmyelination Revealed through MRI as Increased Radial (but Unchanged Axial) Diffusion of Water. *NeuroImage*, *17*(3), 1429–1436. https://doi.org/10.1006/nimg.2002.1267

Sotiropoulos, S. N., & Zalesky, A. (2019). Building connectomes using diffusion MRI: Why, how and but. *NMR in Biomedicine*, *32*(4), e3752. https://doi.org/10.1002/nbm.3752

Spagna, A., Hajhajate, D., Liu, J., & Bartolomeo, P. (2021). Visual mental imagery engages the left fusiform gyrus, but not the early visual cortex: A meta-analysis of neuroimaging evidence. *Neuroscience & Biobehavioral Reviews*, *122*, 201–217. https://doi.org/10.1016/j.neubiorev.2020.12.029

Suggate, S. P., Milton, F., & Tree, J. (2026). Multimodal mental comparisons in those with and without aphantasia. *Neuropsychologia*, *222*, 109373. https://doi.org/10.1016/j.neuropsychologia.2026.109373

Thiebaut de Schotten, M., Dell'Acqua, F., Forkel, S. J., Simmons, A., Vergani, F., Murphy, D. G., & Catani, M. (2011). A lateralized brain network for visuospatial attention. *Nat Neurosci*, *14*(10), 1245–1246. (21926985). https://doi.org/10.1038/nn.2905

Thomas, C., Avidan, G., Humphreys, K., Jung, K., Gao, F., & Behrmann, M. (2009). Reduced structural connectivity in ventral visual cortex in congenital prosopagnosia. *Nature Neuroscience*, *12*(1), 29–31. https://doi.org/10.1038/nn.2224

Thomas, C., Avram, A., Pierpaoli, C., & Baker, C. (2015). Diffusion MRI properties of the human uncinate fasciculus correlate with the ability to learn visual associations. *Cortex: A Journal Devoted to the Study of the Nervous System and Behavior*, *72*, 65–78. https://doi.org/10.1016/j.cortex.2015.01.023

Travers, B. G., Adluru, N., Ennis, C., Tromp, D. P. M., Destiche, D., Doran, S., Bigler, E. D., Lange, N., Lainhart, J. E., & Alexander, A. L. (2012). Diffusion Tensor Imaging in Autism Spectrum Disorder: A Review. *Autism Research*, *5*(5), 289–313. https://doi.org/10.1002/aur.1243

Treisman, A. (2007). How the deployment of attention determines what we see. In *Progress in Psychological Science around the World. Volume 1 Neural, Cognitive and Developmental Issues.* Psychology Press.

Vann, S. D., Aggleton, J. P., & Maguire, E. A. (2009). What does the retrosplenial cortex do? *Nature Reviews Neuroscience*, *10*(11), 792–802. https://doi.org/10.1038/nrn2733

Warrington, S., Bryant, K. L., Khrapitchev, A. A., Sallet, J., Charquero-Ballester, M., Douaud, G., Jbabdi, S., Mars, R. B., & Sotiropoulos, S. N. (2020). XTRACT - Standardised protocols for automated tractography in the human and macaque brain. *NeuroImage*, *217*, 116923. https://doi.org/10.1016/j.neuroimage.2020.116923

Zeman, A. (2024). Aphantasia and hyperphantasia: Exploring imagery vividness extremes. *Trends in Cognitive Sciences*, *28*(5), 467–480. https://doi.org/10.1016/j.tics.2024.02.007

Zeman, A., Milton, F., Della Sala, S., Dewar, M., Frayling, T., Gaddum, J., Hattersley, A., Heuerman-Williamson, B., Jones, K., & MacKisack, M. (2020). Phantasia–the psychological significance of lifelong visual imagery vividness extremes. *Cortex*, *130*, 426–440. https://doi.org/10.1016/j.cortex.2020.04.003

Zhang, H., Schneider, T., Wheeler-Kingshott, C. A., & Alexander, D. C. (2012). NODDI: Practical *in vivo* neurite orientation dispersion and density imaging of the human brain. *NeuroImage*, *61*(4), 1000–1016. https://doi.org/10.1016/j.neuroimage.2012.03.072