Comprehensive mapping of Neurofibromin (NF1) expression in developing mouse brain

Vishal Lolam[1] and Achira Roy[1,*]

[1]Neurodevelopment Laboratory, Neuroscience Unit, Jawaharlal Nehru Centre for Advanced Scientific Research (JNCASR), India

9
10
11
12

*Correspondence: Achira Roy, MSc, PhD
                   Assistant Professor
                   Neurodevelopment Lab, Room: SBS-B201, Neuroscience Unit,
                   Jawaharlal Nehru Centre for Advanced Scientific Research (JNCASR),
                   Rachenahalli Lake Road, Jakkur PO, Bangalore, India. PIN: 560064
                   achira.roy@jncasr.ac.in

Short Title: NF1 expression in mouse brain

26
Keywords: NF1, neurodevelopment, brain expression, cerebellum, single-cell reanalysis

29
30
31
32
Total number of words: 4820

34
Number of Figures: 9
Number of Supplementary Figures: 3

37
38

Abstract:

Neurofibromin (NF1) is a critical negative regulator of the RAS-RAF-ERK pathway, mutations in which have been clinically implicated in various neurodevelopmental disorders. However, the lack of a high-resolution spatiotemporal map has obscured the understanding of why specific cell populations and developmental processes are uniquely vulnerable to *NF1* loss. In this study, we present a comprehensive atlas of NF1 expression in the developing mouse brain. Using *in situ* hybridization and immunohistochemistry, we characterized NF1 distribution from early embryonic stages through postnatal maturation. We further integrated these findings with single-nuclei RNA-sequencing (snRNA-seq) datasets from adult mouse brain to achieve higher resolution. Our results reveal a previously undocumented graded expression pattern of NF1 across various brain regions and cell lineages. This comprehensive study will not only help understand the fundamental role of NF1 in brain development but will also be pivotal in providing a framework for studying *NF1*-associated brain disorders.

Summary statement:

Graded expression of Neurofibromin (NF1) identified across embryonic and postnatal stages reveals unique sub-lineages of neural cells in different parts of the developing mouse brain.

Introduction:

Neurofibromatosis type 1 is one of the most common autosomal dominant neurogenetic disorders, affecting approximately 1 in 3,000 individuals worldwide (Nix et al. 2020). The condition is caused by mutations in the *NF1* gene, which encodes neurofibromin, a large multifunctional protein that serves as a critical negative regulator of the RAS-RAF-ERK pathway (Karaconji et al. 2019). NF1 consists of RAS-GTPase activating (GAP) domain, which facilitates the conversion of RAS-GTP (Active) to RAS-GDP (inactive), thereby negatively regulating RAS function (Martin et al. 1990). Since the RAS-RAF-ERK pathway is important for cell growth, proliferation and survival, NF1 acts as a brake on cellular growth and proliferation (Cichowski & Jacks 2001). In patients suffering from Neurofibromatosis type 1, this brake is lost, resulting in the formation of benign or malignant tumours in the brain and brainstem (Albers & Gutmann 2009; Lucas et al. 2022; Ayasa et al. 2024). Interestingly, a significant proportion of these pediatric patients develop neurodevelopmental disorders such as cortical malformations, epilepsy, hydrocephalus, autism spectrum disorder, intellectual disabilities, and movement disorders (Balestri et al. 2003; Sorrentino et al. 2021; Roth et al. 2019; Plasschaert et al. 2015; Hyman et al. 2005; Rietman et al. 2017; Lolam & Roy 2025; Báez-Flores et al. 2023). These clinical findings underscore the importance of NF1 in neurodevelopment beyond its canonical role in tumorigenesis.

Despite the profound impact of *NF1* mutations on the functioning of the central nervous system (CNS), our understanding of the spatiotemporal dynamics of *NF1* expression during brain development remains incomplete. Previous studies have established that NF1 is essential for embryonic development, with null mice dying in utero between embryonic days (E) 12.5 and 13.5 due to cardiovascular defects (Brannan et al. 1994; Jacks et al. 1994). In addition, studies done by knocking out *Nf1* in cultured cells and in various model systems have demonstrated altered neurodevelopmental processes (Lakkis et al. 1999; Zhu et al. 2001; Gutmann et al. 1999; Sanchez-Ortiz et al. 2014; Shin et al. 2012; Durkin et al. 2023). However, the lack of a high-resolution spatiotemporal map has limited our ability to pinpoint which developmental processes or cell populations are most vulnerable to *Nf1* loss.

Although there are some reports of NF1 expression in the brain (Daston et al. 1992; Su et al. 2007; Ryu et al. 2019; Zhu et al. 2001; Provenzano et al. 2014), no one has conducted a comprehensive high-resolution mapping of this protein in mammalian brain development. In this study, we present a high-resolution, comprehensive map of NF1 expression in the developing mouse brain. Using in situ hybridization and immunohistochemistry (IHC), we characterize the precise spatiotemporal distribution of NF1 across major brain regions and cell lineages from early embryonic stages through

109   postnatal maturation. Moreover, we used publicly available single-nuclei RNA-seq

110   (snRNA-seq) databases of the adult mouse brain to map NF1 expression across

111   different cell clusters, at a resolution much higher than achievable with classical

112   techniques alone. These results revealed a previously unreported graded expression

113   pattern of NF1.

Results:

NF1 is expressed in different cell lineages of the murine forebrain

Using IHC, we observed broad NF1 expression in different parts of the mouse forebrain, such as the neocortex, hippocampus, choroid plexus, thalamus, and hypothalamus (**Fig. 1A-P**). In the postnatal day (P)35 hypothalamus, NF1 strongly marked the paraventricular nucleus (PVN), suprachiasmatic nucleus (SCN), and supraoptic nucleus (SON), involved in regulating circadian rhythm, neurohormone release and fluid balance. Similar localization of *Nf1* mRNA was also observed at P35 and P0 using *in situ* hybridization (**Supplementary Fig. S1A-B"**). In embryonic ages E17.5, E14.5, and E12.5, NF1 protein expression was observed along the lateral and third ventricular linings, including neocortical and ventral telencephalic ventricular zone, cortical hem, and hippocampal primordium (**Fig. 1D-F**). Specifically, NF1 showed perinuclear expression in the neocortical plate and white matter as well as CA fields and hilus of hippocampus, but not dentate gyrus (DG) (**Fig. 1G-I**). By colabelling with different cell-type-specific telencephalic markers across development, we identified sets of cells that express the NF1 protein (**Fig. 2**). We observed strong perinuclear labelling of NF1 in excitatory pyramidal neocortical neurons from the peak neurogenic stage at E14.5 till developed cortical stage at P35 (**Fig. 2A-H,K,L**). This includes coexpression with markers specific to all mature pyramidal neurons (NeuN), deep layer-specific neurons (Ctip2), and upper layer neurons (Satb2). When checked for colocalization with the progenitors at the neocortical ventricular zone, NF1 was found perinuclearly in the Sox2$^+$ apical progenitors at E17.5, E14.5, and E12.5, but not in the Tbr2$^+$ intermediate progenitor cells (**Fig. 2I,J,M-P**). Further, NF1 was strongly expressed in the hippocampal CA1 and CA3 pyramidal neurons and hilar neurons but did not label any cells in the dentate gyrus (DG), as observed at P35, P0 and E17.5 (**Fig. 2Q-Y"**). These were determined by colabelling NF1 with NeuN to mark all differentiated hippocampal neurons across ages, and Ctip2, which prominently marks CA1 and developing DG at E17.5. Even at E14.5, weak NF1 expression was observed in the ventricular zone of cortical hem and hippocampal primordium/anlage (**Fig. 2Z-Z"**). E12.5 dorsal midline showed mild NF1 expression mostly concentrated along the ventricular lining (**Fig. 2AA-AA"**).

We also observed strong NF1 expression in different subtypes of cortical interneurons at P35, colabelling with Parvalbumin and Somatostatin (**Supplementary Fig. S2A-D"**). Interestingly, NF1 was observed to be present only in certain subsets of Olig2$^+$ oligodendrocyte populations in the major axonal tracts of P35 mouse forebrain, namely corpus callosum, hippocampal commissure, and anterior commissure (**Fig. 3A-F''**). This observation may suggest differential function of molecularly distinct subtypes of oligodendrocytes when NF1 levels are disrupted in the brain.

NF1 is expressed in different cell lineages of the murine cerebellum

Unlike the forebrain, a more restricted protein expression profile of NF1 was observed in the murine cerebellum across the developmental timeline (**Fig. 4A-J"**). Particularly, strong perinuclear NF1 expression was observed in mature and migrating Purkinje cells, colabelling with Calbindin (**Fig. 4E',F',G'; Fig. 5A-D"**). On the other hand, very low or no IHC signal of NF1 was observed at any stage of cerebellar granule cell development, when colabelled with NeuN (**Fig. 5E-H",M,N**). *Nf1* mRNA was also observed in the parts of cerebellum at P35 and P0 using *in situ* hybridization (**Supplementary Fig. S1C-D"**). Intriguingly, NF1 expression was observed throughout the development of the cerebellar neurons, be it in the migratory state or transitory state at the embryonic nuclear transitory zone (NTZ) or later when forming the deep cerebellar nuclei (DCN) located within the white matter (**Fig. 4E",F",G"H", I,I',J,J'**). It was coexpressed with NeuN, marking neurons at the DCN, from P3 to P35 (**Fig. 5I-K"**); NeuN+ cells were not present in the prospective DCN area at P0 (**Fig. 5L-L"**).

At embryonic stages, the cerebellar anlage has distinct proliferative zones, namely rhombic lip and cerebellar ventricular zone (CVZ) (Leto et al. 2016). At early cerebellar development (E12.5), CVZ is further subdivided molecularly into the Purkinje cell-producing zone and the Pax2$^+$ interneuron-producing zone. Later in development, the whole CVZ becomes potent to produce cerebellar interneurons. Perinuclear NF1 expression was observed around Sox2+ nuclei in the apical progenitors of the CVZ at E12.5 and E14.5 (**Fig. 6 A-B-B",D,E-E"**). However, no overlap was observed in rhombic lip-derived Pax6$^+$ cells at E12.5 or E14.5 (**Fig. 6C-C", F-F"**). We also found NF1 to mark differentiated Pax2$^+$ cells that have migrated out of the interneuron-producing side of the CVZ at E14.5 (**Fig. 4 I,I", Supplementary Fig. S3A-B"**). However, NF1 was expressed in only a small subset of Tbr2+ rhombic lip-derived unipolar brush cells at E14.5 (**Supplementary Fig. S3E-E"**).

Finally, similar to forebrain results, NF1 showed an interesting expression pattern in glial lineages of the cerebellum (**Fig. 7A-N"**). At both the cerebellar folium and the DCN region, NF1 showed perinuclear expression in a subset of Olig2$^+$ oligodendrocytes, as observed at P35 and P12 (**Fig. 7A-H"**). The same trend was observed in the subpopulations of Bergmann glia, which are unipolar astrocytes of the cerebellum (**Fig. 7I-N"**).

Taken together, here we report differential expression of NF1 in the mouse brain, specifically in glial populations, along development. This suggests the possibility of alternate regulatory mechanisms that eventually govern brain function.

NF1 is expressed in distinct cell layers of the eye and olfactory bulb

Beyond regular brain regions, NF1 was also found to be expressed in accessory brain regions like the eye and olfactory bulb, both of which are connected to neurosensation (**Fig. 8A-R"**). Expression of NF1 in the developing eye was substantially strong in the lens, retinal pigmented epithelium (RPE), and neuroretina. In the neuroretina, NF1 expression was distinct in the photoreceptor cell layer (PRC), and in the retinal ganglionic cells and Müller glial cells present in the ganglionic cell layer. A subset of NF1$^+$ cells in the retinal ganglionic cell (RGC) layer colocalized with Tbr2, which is known to be a key regulator of the intrinsically photosensitive RGCs (Chen et al. 2021). Sox2$^+$ zone of E14.5 and E17.5 neuroretina largely did not overlap with the NF1$^+$ layers, apart from the PRCs (**Fig. 8A,C-F"**). With respect to the developing olfactory bulb, NF1 was observed specifically in the mitral cell layer (MCL) of the main olfactory bulb (MOB), which are projection neurons critical for processing the sense of odours (Davison & Katz 2007). NF1 was also expressed in the accessory olfactory bulb (AOB) from P0 to P35, which is important for aggression and reproduction (Ennis & Holy 2015) (**Fig. 8G-J**). NF1 overlapped with NeuN in the MCL at P35 and P12 (**Fig. 8K,L,O-P"**). On the other hand, it did not have much overlap with Pax6$^+$ cells at P3 and P0, apart from posterior AOB (**Fig. 8M,N,Q-R"**).

Reanalysis of single nuclei RNAseq data reveals enriched *Nf1* expression in diverse neural cell types

To determine the expression of *Nf1* at higher resolution, we utilized publicly available single-nuclei RNAseq databases (Mortberg et al. 2023; Kozareva et al. 2021). In the adult mouse cortex, *Nf1* expression was identified in almost all the marked cortical clusters (**Fig. 9A-C**). However, detailed analysis revealed specific enrichment of *Nf1* transcripts in the lineages of cortical interneurons, oligodendrocytes, and excitatory neurons (**Fig. 9D**). In particular, *Nf1* expression was found to be highest in the oligodendrocyte lineage. Moreover, among different oligodendrocyte clusters, *Nf1* expression was found to be highest in differentiating oligodendrocytes (**Fig. 9D**). Based on our IHC data, all Olig2+ cells in the cortex were not positive for NF1. Hence, we looked at the percentage of cells positive for *Nf1* expression across different cortical clusters (**Fig. 9E**). Interestingly, oligodendrocyte precursor cells (OPCs) and differentiating OPCs showed a very high frequency of *Nf1*-expressing cells than mature oligodendrocytes. This data strongly suggests the importance of NF1 in oligodendrocyte development. Besides oligodendrocytes, cortical interneurons and excitatory pyramidal neurons also exhibit higher frequency and expression, corroborating our IHC findings.

We performed a similar analysis on adult cerebellum and identified an enriched expression of *Nf1* in CVZ-derivatives, such as cerebellar interneurons, Purkinje cells, Bergmann glia, and rhombic lip-derivative unipolar brush cell (**Fig. 9F-H**). Among these

clusters, the highest expression was found in the Bergmann glial cells (**Fig. 9I**).
Similarly, *Nf1* showed enriched expression in the oligodendrocyte lineage of the adult
neocortex, with a higher frequency of *Nf1*-expressing cells in the OPC cluster as
compared to mature oligodendrocytes (**Fig. 9H**). The frequency plot also demonstrated
that almost all cells in the Purkinje cell cluster were positive for *Nf1*, thus aligning with
our IHC expression data.
240
241   Overall, these reanalysis results not only validated our IHC data but also helped identify
242   *Nf1* expression in some neural cell types, such as differentiating oligodendrocytes,
243   which are difficult to capture using conventional IHCs. Our in-depth analysis also
244   uncovered the graded expression pattern of *Nf1* across diverse cell types in the murine
245   adult brain. This may suggest the differential role of *Nf1* in these diverse subtypes.
246

Discussion:

In this comprehensive study, we demonstrate the spatiotemporal expression of neurofibromin (NF1) in the murine brain across embryonic, perinatal, and adult developmental time points, using IHC, *in situ* hybridization, and reanalyses of publicly available single-nuclei RNA-seq datasets. To our knowledge, this is the first report of an NF1 expression study, not only across major brain regions but also along almost the entire span of development. Crucially, our findings reveal molecularly distinct neural subtypes in the forebrain and cerebellum defined by the presence or absence of NF1, suggesting a previously unappreciated heterogeneity in NF1-mediated signaling during neurodevelopment. The high degree of concordance between our protein (IHC) and transcriptomic (snRNA-seq) data establishes a robust framework for understanding the role of NF1 in neural development.

Cell-type-specific NF1 expression in the forebrain suggests roles in neuronal lineage progression and excitation-inhibition balance

In this study, we identify broad yet cell-type-specific NF1 expression across the developing and postnatal forebrain, including the neocortex, hippocampus, and major white matter tracts. NF1 demonstrated consistent perinuclear localization and strong agreement between protein and transcript levels. In the neocortex, it is expressed in excitatory pyramidal neurons across all layers and in $Sox2^+$ apical progenitors, but interestingly not in $Tbr2^+$ intermediate progenitors. This indicates a stage-specific role of NF1 during cortical neurogenesis. In parallel, NF1 is expressed in inhibitory interneurons, including $Parvalbumin^+$ and $Somatostatin^+$ populations. This suggests that NF1 is involved in both arms of cortical circuitry, raising the possibility that NF1 loss can disrupt excitation–inhibition balance. In the hippocampus, NF1 shows region-specific enrichment in the pyramidal neurons of CA fields and hilar neurons, but is notably absent from the dentate gyrus, suggesting subfield-specific functional roles. Besides, its expression is also observed in specific hypothalamic and thalamic nuclei of the diencephalon, as well as in the ventricular zone of the ventral telencephalon in younger ages. Together, these findings position NF1 as a key regulator across multiple forebrain cell types, with potential implications for lineage specification, neuronal integrity, circuit formation, and function.

Role of NF1 in cerebellar development and motor coordination

In the cerebellum, NF1 is expressed across multiple neuronal populations throughout development, including deep cerebellar nuclei (DCN) neurons, Purkinje neurons, and cerebellar interneurons across all examined stages. In the cerebellum, proper functioning of DCN neurons and associated circuits is critical for having regular motor coordination and function (Lackey et al. 2023; van der Heijden 2024). The persistent expression of NF1 identified in the DCN and developing neurons throughout our

experimental timeline highlights its importance in motor circuits. Mutations in *NF1* may impair the development or function of these neurons, resulting in motor deficits, as commonly seen in patients harbouring *NF1* mutations (Rietman et al. 2017). Similarly, defects in NF1 may affect the Purkinje cell- and cerebellar interneuron-related miscoordination. We also observed NF1 expression in subsets of Bergmann glia, cerebellar astrocytic cells that coordinate cell-cell communication, support development, and prevent neurological injury (Araujo et al. 2019; De Zeeuw & Hoogland 2015; Buffo & Rossi 2013). Having the knowledge of which subtype is affected in a disease condition may help identify specific therapeutic interventions.

Mosaic NF1 expression across neural lineages reveals previously unrecognized cellular heterogeneity

Our analyses reveal molecular heterogeneity within both neuronal and glial populations based on NF1 expression status, identifying NF1+ve and NF1−ve subpopulations across brain regions and suggesting a mosaic pattern of expression. In particular, distinct NF1+ve and NF1−ve subsets of Olig2$^+$ oligodendrocytes are observed in cerebellum and major forebrain white matter tracts, while only a subset of Bergmann glia expresses NF1. This reveals previously unappreciated diversity within glial lineages. These findings are supported both by IHC data and snRNA-seq analyses. In addition, transcriptomic data reveal NF1+ve and NF1−ve interneuron populations in the cortex and cerebellum, extending this heterogeneity to neuronal lineages. The functional significance of this differential expression remains unclear, but it suggests several possibilities: NF1 may mark distinct developmental trajectories, reflect differences in maturation state, or be dynamically regulated within lineages, defining transient cellular states. These observations raise key questions regarding the developmental origins of NF1+ve and NF1−ve populations, the mechanisms controlling NF1 expression, and the functional consequences of this heterogeneity, which may ultimately contribute to variability in NF1-associated disease phenotypes.

NF1 expression in sensory systems extends its role to photo-dependent and olfactory circuitry

Beyond the major regions of the brain, we report NF1 expression also in specific layers of the developing eye and olfactory bulb, offering new insights into sensory-related pathologies. Specifically, NF1 is expressed in the photoreceptor layer of the neuroretina, which is vital for visual phototransduction (Hussey et al. 2022). It is also identified in Müller glial cells, which provide support and guidance for the light to reach the photoreceptor layer (Franze et al. 2007), and retinal ganglion cells (RGCs), which finally transmit sensory information to the brain. The signal perceived by the eye is known to produce both visual and non-visual responses in different parts of the diencephalon (Peirson et al. 2018). While light information cascades through the

thalamic nuclei to the visual cortex for vision, it also triggers the hypothalamic nuclei
critical for circadian rhythm, sleep and neuroendocrine functions (Dumbell et al. 2016;
Trachtman 2010). Our study revealed NF1 expression specifically in the SCN, the
master circadian rhythm generator, and in the VPN and SON, neuroendocrine nuclei
involved in fluid balance and rhythmic production of hormones. Together, these strongly
suggest a potential role of NF1 in light sensation, transduction, and eventually the light-
dependent regulatory functions in the brain. Further work may reveal novel molecular
interactions between NF1 and the diencephalon regulating sleep and metabolic
homeostasis.
336
337 Similarly, the presence of NF1 expression in the mitral cells of the olfactory bulb and in
338 both parts of the AOB suggests an underexplored role for NF1 in odor and pheromone
339 processing and in pheromone-driven behaviors, such as reproductive and defensive
340 responses. The conservation of these patterns suggests that NF1 may be integral to
341 evolutionary ancient sensory circuits.
342
Implications for NF1-associated neurodevelopmental disorders and
tumorigenesis
This comprehensive atlas of NF1 expression across murine brain development provides
a framework for linking cellular context to NF1-associated neurodevelopmental
disorders and tumorigenesis. The widespread yet heterogeneous expression of NF1
across neuronal and glial populations suggests that its dysregulation may impact
multiple circuits and developmental processes, contributing to the clinical variability
observed in NF1 patients. The identification of NF1+ve and NF1−ve subpopulations
further raises the possibility of selective cellular vulnerability, which may influence both
disease manifestation and tumor initiation. These findings offer a basis for identifying
cells of origin and critical developmental windows in NF1-associated pathologies.
354
355 Overall, this work provides a resource for understanding NF1 function in the developing
356 brain and supports future efforts toward cell-type-specific and targeted therapeutic
357 strategies.
358
Limitations of the study
While this study provides a comprehensive spatiotemporal map of NF1 expression,
several limitations should be considered. First, our analyses are primarily descriptive
and based on IHCs, *in situ* hybridization, and reanalysis of existing snRNAseq datasets;
therefore, they do not directly address the functional consequences of NF1 loss in
specific cell types. Second, although strong concordance is observed across modalities,
differences in sensitivity and resolution between protein- and transcript-level
measurements may influence the detection of low-abundance or transient expression of

NF1 in different cellular lineages. In addition, our conclusions are drawn from murine
models, and although broadly informative, species-specific differences may limit direct
translation to human biology. The observed NF1+ve and NF1−ve heterogeneity across
cell populations also remains unresolved, particularly with respect to developmental
origin, temporal dynamics, and regulatory mechanisms. Addressing these limitations
through functional perturbation studies, higher-resolution temporal analyses, and cross-
species validation will be important for further elucidating the role of NF1 in brain
development and disease.
375

376 **Materials and methods:**

378 **Key resources table:**

Reagent type/ resource	Designation	Reference/ Source	Identifiers
Mouse	FVB/NJ	Gift from Prof Kathleen Millen, Seattle Children's Research Institute, USA	-
Antibody	Anti-Parvalbumin Antibody	Merck	Millipore (Cat # MAB1572, RRID:AB_2174013)
Antibody	Anti-Somatostatin Antibody, clone YC7	Merck	Sigma-Aldrich (Cat # MAB354, RRID: AB_2255365)
Antibody	Calbindin D28K Monoclonal Antibody (5A9)	Thermo Fisher Scientific	Invitrogen (Cat # MA5-47441, RRID:AB_2942423)
Antibody	SOX2 Monoclonal Antibody (20G5)	Thermo Fisher Scientific	Invitrogen (Cat # MA1-014, RRID:AB_2536667)
Antibody	Ki-67 Monoclonal Antibody (SolA15)	Thermo Fisher Scientific	Invitrogen (Cat # 14-5698-82, RRID:AB_10854564)
Antibody	Olig2 (F9R1H) Mouse Monoclonal Antibody	Cell Signaling Technology	CellSignalingTechnology (Cat # 39588)
Antibody	PAX2 Monoclonal Antibody (3C7)	Thermo Fisher Scientific	Invitrogen (Cat # H00005076-M01)
Antibody	PAX6 Monoclonal Antibody (GT9412)	Thermo Fisher Scientific	Invitrogen (Cat # MA5-50298, RRID:AB_3094028)
Antibody	Anti-NeuN Antibody, clone A60	Merck	Sigma-Aldrich (Cat # MAB377, RRID: AB_2298772)
Antibody	ALDH1L1 mouse monoclonal antibody, clone UMAB43	OriGene	Origene (Cat # UM500039, RRID:AB_2629053)
Antibody	Anti-Iba1 goat polyclonal antibody	Abcam	Abcam (Cat # ab5076, RRID:AB_2224402)
Antibody	Custom Rabbit polyclonal antibody	Neurodevelopment lab, India	

	against NF1		
Antibody	Anti-SATB1 + SATB2 antibody [SATBA4B10] - C-terminal	Abcam	Abcam (Cat # ab51502, RRID:AB_882455)
Antibody	Anti-Ctip2 antibody [25B6]	Abcam	Abcam (Cat # ab18465, RRID:AB_2064130)
Antibody	Anti-Tbr1 Antibody	Merck	Millipore (Ca t# AB2261, RRID:AB_10615497)
Antibody	EOMES Monoclonal Antibody (Dan11mag)	Thermo Fisher Scientific	Invitrogen (Cat # 14-4875-82, RRID:AB_11042577)
Antibody	FOXJ1 Monoclonal Antibody (2A5)	Thermo Fisher Scientific	Invitrogen (Cat # 14-9965-82, RRID:AB_1548835)
Antibody	Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488 -	Thermo Fisher Scientific	Invitrogen (Cat # A-11029, RRID:AB_2534088)
Antibody	Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 568 -	Thermo Fisher Scientific	Invitrogen (Cat # A-11036, RRID:AB_10563566)
Antibody	Goat anti-Rat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 647	Thermo Fisher Scientific	Invitrogen (Cat # A-21247, RRID:AB_141778)
Antibody	Goat anti-Chicken IgY (H+L) Secondary Antibody, Alexa Fluor™ 568	Thermo Fisher Scientific	Invitrogen (Cat # A-11041, RRID:AB_2534098)
Antibody	Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 488	Thermo Fisher Scientific	Invitrogen (Cat # A-11034, RRID:AB_2576217)
Antibody	Donkey anti-Goat IgG (H+L) Cross-Adsorbed Secondary	Thermo Fisher Scientific	Invitrogen (Cat # A-21447, RRID:AB_2535864)

	Antibody, Alexa Fluor™ 647		
Commercial assay/Kit	RNAscope™ 2.5 HD Reagent Kit-RED	Biotechne	ACD (Cat # 322350)
Probe	RNAscope™ Negative Control Probe	Biotechne	ACD (Cat # 320751)
Probe	RNAscope™ Positive Control Probe- Mm-Polr2a	Biotechne	ACD (Cat # 312471)
Probe	RNAscope Probe Mm Nf1 C1	Biotechne	ACD (Cat # 417351)
Software, Algorithm	Fiji	PMID: 22743772	
Software, Algorithm	10x Genomics Cell Ranger v10.0.0		Single-cell analyses
Software, Algorithm	Seurat v5	PMID: 37231261	Single-cell analyses

Mice:

For all the experiments FVB/NjJ mouse strain was used. Mice were housed and bred under pathogen-free conditions in the animal facility at Jawaharlal Nehru Centre for Advanced Scientific Research (JNCASR, India). All mice received food and water *ad libitum* and were subjected to a 12-hour light-dark cycle. Noon of the day of vaginal plug was designated as embryonic day 0.5 (E0.5). The day of birth was designated as postnatal day 0 (P0), which usually occurs between E19-20.

Sample preparation:

Brains from postnatal pups were harvested in phosphate buffer saline (PBS), then were fixed in 10% Formalin Solution neutral buffer (10% NBF) for 48-72 hrs for P35 and 2-6hrs for all embryonic and early postnatal ages, equilibrated in 30% (wt/vol) sucrose made in 1X PBS, and sectioned at 30 µm on a freezing microtome (Leica, Germany). These sections were then processed for *in situ* hybridization or immunohistochemical procedures.

Immunohistochemistry:

boiled in 10mM sodium citrate solution for antigen retrieval, blocked in 10% serum in PBS with 0.1% Triton X-100, and then incubated for 36-48hrs at 4°C with primary antibodies. The next day, sections were washed thrice in PBS, incubated with

appropriate species-specific secondary antibodies conjugated with Alexa 488, 568, or
647 fluorophores (Invitrogen) for 2 hrs at room temperature and then counterstained
with 4′,6-diamidino-2-phenylindole (DAPI) to visualize the nuclei. Sections were
coverslipped using Fluoromount G mounting medium (Fisher Scientific, USA).
Information regarding the antibodies used in this study is listed in the Key Resources
table above. Immunohistochemistry experiments were replicated with biologically
independent samples (minimum three). All attempts for replication were successful. No
outliers were encountered. Each antibody was validated for mouse and application
(IHC) by the respective manufacturer, and the data are publicly available. This was also
validated by us in our experiments, replicating published/expected expressions in
control tissue.
412
413 *In situ* hybridization:
414 Assays were run using commercially available probes from Advanced Cell Diagnostics
415 (ACD, Biotechne). Manufacturer-recommended protocols were used without
416 modification. Sections were probed using *Nf1* (Cat# 417351) and counterstained with
417 hematoxylin.
418
Imaging of brain sections:
Imaging of IHC brain sections was done using CellVoyager™ CQ1 Benchtop High-
Content Analysis System using CellPathfinder (Yokogawa, Japan), and later processed
in ImageJ software (NIH, Bethesda, Maryland, USA). *In situ* slides were visualized
under an Olympus VS 200 slide scanner brightfield microscope. Apart from minor
adjustments to contrast and brightness across the entire image, there was no further
image alteration. Pseudo-colouring was used to highlight the cells of interest. Figures
were prepared on Adobe Illustrator.
427
Reanalysis of single-nuclei transcriptomics datasets:
Using Single Cell Portal (singlecell.broadinstitute.org/single_cell), we conducted a
reanalysis of single-nuclei RNAseq data of adult cerebellum (SCP795) and adult cortex
(SCP2126), originally reported by (Mortberg et al. 2023) and (Kozareva et al. 2021),
respectively. The Seurat object was downloaded from the portal, and downstream
analysis was performed using Seurat v5 (Hao et al. 2024). UMAP clusters were
annotated to different neural cell types based on their transcriptomic signatures. We
specifically focused on examining Nf1 expression levels across diverse cell types.
Subsequently, we generated graphical plots with the help of ggplot2 package (Wickham
2016).
438

Author Contributions:
VL and AR contributed to the study conception, study design, data acquisition, data analysis, and interpretation. VL and AR wrote the first draft of the manuscript, reviewed it, and approved the final version. AR provided funding and infrastructural resources.

Acknowledgements:
We thank Prof Kathleen J Millen, Seattle Children's Research Institute, for gifting the FVB mouse line; Varun M. and Shruti Srivastava for technical support during image acquisition. Cerebellar lamination schematics in Fig. 5M and 5N have been created using the BioRender software (https://www.biorender.com/).

Funding:
This work was funded by the Ramalingaswami Re-entry Fellowship (BT/RLF/Re-entry/73/2020) from the Department of Biotechnology (DBT), India (AR); the Prime Minister Early Career research Grant (PM-ECRG, ANRF/ECRG/2024/001273/LS) from the Anusandhan National Research Foundation (ANRF), India (AR); Early Career Award from the International Brain Research Organization (IBRO, ECA-8126694738) (AR); and the intramural funding from the Jawaharlal Nehru Centre for Advanced Scientific Research, India (AR).

Ethics
Animal experimentation: All animal experimentation was conducted in accordance with the guidelines laid down by the Institutional Animal Ethics Committees (IAEC) of Jawaharlal Nehru Centre for Advanced Scientific Research, India (protocols AR 002 and AR 003).

Data/resource availability:
All relevant data and details on resources are provided in the article and its supplementary information.

Conflict of interest:
The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Diversity and inclusion statement
Preparation of this document has involved diversity and inclusion practices relevant to both the scientific content of the paper and authorship and attribution.

References:

Albers, A.C. & Gutmann, D.H., 2009. Gliomas in patients with neurofibromatosis type 1. *Expert Review of Neurotherapeutics*, 9(4), pp.535–539. Available at: https://doi.org/10.1586/ern.09.4.

Araujo, A.P.B., Carpi-Santos, R. & Gomes, F.C.A., 2019. The role of astrocytes in the development of the cerebellum. *Cerebellum (London, England)*, 18(6), pp.1017–1035. Available at: https://doi.org/10.1007/s12311-019-01046-0.

Ayasa, L.A. et al., 2024. Glioblastoma multiforme in a patient with neurofibromatosis type 1: a case report and review of literature. *Journal of Surgical Case Reports*, 2024(8), p.rjae517. Available at: https://doi.org/10.1093/jscr/rjae517.

Báez-Flores, J., Rodríguez-Martín, M. & Lacal, J., 2023. The therapeutic potential of neurofibromin signaling pathways and binding partners. *Communications Biology*, 6(1), p.436. Available at: http://dx.doi.org/10.1038/s42003-023-04815-0.

Balestri, P. et al., 2003. Malformations of cortical development in neurofibromatosis type 1. *Neurology*, 61(12), pp.1799–1801. Available at: https://doi.org/10.1212/01.WNL.0000099080.90726.BA.

Brannan, C.I. et al., 1994. Targeted disruption of the neurofibromatosis type-1 gene leads to developmental abnormalities in heart and various neural crest-derived tissues. *Genes & Development*, 8(9), pp.1019–1029. Available at: http://dx.doi.org/10.1101/gad.8.9.1019.

Buffo, A. & Rossi, F., 2013. Origin, lineage and function of cerebellar glia. *Progress in Neurobiology*, 109, pp.42–63. Available at: https://doi.org/10.1016/j.pneurobio.2013.08.001.

Chen, C.-K. et al., 2021. Characterization of Tbr2-expressing retinal ganglion cells. *The Journal of Comparative Neurology*, 529(15), pp.3513–3532. Available at: http://dx.doi.org/10.1002/cne.25208.

Cichowski, K. & Jacks, T., 2001. NF1 tumor suppressor gene function: narrowing the GAP. *Cell*, 104(4), pp.593–604. Available at: http://dx.doi.org/10.1016/s0092-8674(01)00245-8.

Daston, M.M. et al., 1992. The protein product of the neurofibromatosis type 1 gene is expressed at highest abundance in neurons, Schwann cells, and oligodendrocytes. *Neuron*, 8(3), pp.415–428. Available at: https://doi.org/10.1016/0896-6273(92)90270-N.

Davison, I.G. & Katz, L.C., 2007. Sparse and selective odor coding by mitral/tufted neurons in the main olfactory bulb. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 27(8), pp.2091–2101. Available at: https://doi.org/10.1523/JNEUROSCI.3779-06.2007.

De Zeeuw, C.I. & Hoogland, T.M., 2015. Reappraisal of Bergmann glial cells as modulators of cerebellar circuit function. *Frontiers in Cellular Neuroscience*, 9, p.246. Available at: https://doi.org/10.3389/fncel.2015.00246.

Dumbell, R., Matveeva, O. & Oster, H., 2016. Circadian clocks, stress, and immunity. *Frontiers in Endocrinology*, 7, p.37. Available at: http://dx.doi.org/10.3389/fendo.2016.00037.

Durkin, J. et al., 2023. Neurofibromin 1 regulates early developmental sleep in Drosophila.
Neurobiology of Sleep and Circadian Rhythms, 15(100101), p.100101. Available at:
http://dx.doi.org/10.1016/j.nbscr.2023.100101.

Ennis, M. & Holy, T.E., 2015. Anatomy and neurobiology of the main and accessory olfactory
bulbs. *Handbook of Olfaction and Gustation*, pp.157–182. Available at:
http://dx.doi.org/10.1002/9781118971758.ch8.

Franze, K. et al., 2007. Müller cells are living optical fibers in the vertebrate retina. *Proceedings*
of the National Academy of Sciences of the United States of America, 104(20), pp.8287–
8292. Available at: http://dx.doi.org/10.1073/pnas.0611180104.

Gutmann, D.H. et al., 1999. Haploinsufficiency for the neurofibromatosis 1 (NF1) tumor
suppressor results in increased astrocyte proliferation. *Oncogene*, 18(31), pp.4450–4459.
Available at: https://doi.org/10.1038/sj.onc.1202829.

Hao, Y. et al., 2024. Dictionary learning for integrative, multimodal and scalable single-cell
analysis. *Nature Biotechnology*, 42(2), pp.293–304. Available at:
https://doi.org/10.1038/s41587-023-01767-y.

van der Heijden, M.E., 2024. Converging and diverging cerebellar pathways for motor and
social behaviors in mice. *Cerebellum (London, England)*, 23(5), pp.1754–1767. Available
at: http://dx.doi.org/10.1007/s12311-024-01706-w.

Hussey, K.A., Hadyniak, S.E. & Johnston, R.J., Jr, 2022. Patterning and development of
photoreceptors in the human retina. *Frontiers in Cell and Developmental Biology*, 10,
p.878350. Available at: http://dx.doi.org/10.3389/fcell.2022.878350.

Hyman, S.L., Shores, A. & North, K.N., 2005. The nature and frequency of cognitive deficits in
children with neurofibromatosis type 1. *Neurology*, 65(7), pp.1037–1044. Available at:
https://doi.org/10.1212/01.wnl.0000179303.72345.ce.

Jacks, T. et al., 1994. Tumour predisposition in mice heterozygous for a targeted mutation in
Nf1. *Nature Genetics*, 7(3), pp.353–361. Available at: https://doi.org/10.1038/ng0794-353.

Karaconji, T. et al., 2019. Neurofibromatosis type 1: Review and update on emerging therapies.
Asia-Pacific Journal of Ophthalmology (Philadelphia, Pa.), 8(1), pp.62–72. Available at:
http://dx.doi.org/10.22608/APO.2018182.

Kozareva, V. et al., 2021. A transcriptomic atlas of mouse cerebellar cortex comprehensively
defines cell types. *Nature*, 598(7879), pp.214–219. Available at:
https://doi.org/10.1038/s41586-021-03220-z.

Lackey, E.P., Rey Hipolito, A.G. & Sillitoe, R.V., 2023. Motor circuit abnormalities during
cerebellar development. In *Contemporary Clinical Neuroscience*. Cham: Springer
International Publishing, pp. 129–154. Available at: http://dx.doi.org/10.1007/978-3-031-
23104-9_7.

Lakkis, M.M. et al., 1999. Neurofibromin deficiency in mice causes exencephaly and is a
modifier for Splotch neural tube defects. *Developmental Biology*, 212(1), pp.80–92.
Available at: https://doi.org/10.1006/dbio.1999.9327.

Leto, K. et al., 2016. Consensus paper: Cerebellar development. *Cerebellum (London,*
England), 15(6), pp.789–828. Available at: http://dx.doi.org/10.1007/s12311-015-0724-2.

Lolam, V. & Roy, A., 2025. Developmental mechanisms underlying pediatric epilepsy. *Frontiers*
in Neurology, 16, p.1586947. Available at: http://dx.doi.org/10.3389/fneur.2025.1586947.

Lucas, C.-H.G. et al., 2022. Multiplatform molecular analyses refine classification of gliomas
arising in patients with neurofibromatosis type 1. *Acta neuropathologica*, 144(4), pp.747–
765. Available at: https://doi.org/10.1007/s00401-022-02478-5.

Martin, G.A. et al., 1990. The GAP-related domain of the neurofibromatosis type 1 gene product
interacts with ras p21. *Cell*, 63(4), pp.843–849. Available at: http://dx.doi.org/10.1016/0092-
8674(90)90150-d.

Mortberg, M.A. et al., 2023. A single-cell map of antisense oligonucleotide activity in the brain.
Nucleic Acids Research, 51(14), pp.7109–7124. Available at:
https://doi.org/10.1093/nar/gkad371.

Nix, J.S., Blakeley, J. & Rodriguez, F.J., 2020. An update on the central nervous system
manifestations of neurofibromatosis type 1. *Acta neuropathologica*, 139(4), pp.625–641.
Available at: http://dx.doi.org/10.1007/s00401-019-02002-2.

Peirson, S.N. et al., 2018. Light and the laboratory mouse. *Journal of Neuroscience Methods*,
300, pp.26–36. Available at: https://doi.org/10.1016/j.jneumeth.2017.04.007.

Plasschaert, E. et al., 2015. Prevalence of Autism Spectrum Disorder symptoms in children with
neurofibromatosis type 1. *American Journal of Medical Genetics. Part B, Neuropsychiatric*
Genetics: The Official Publication of the International Society of Psychiatric Genetics,
168B(1), pp.72–80. Available at: http://dx.doi.org/10.1002/ajmg.b.32280.

Provenzano, G. et al., 2014. Hippocampal dysregulation of neurofibromin-dependent pathways
is associated with impaired spatial learning in engrailed 2 knock-out mice. *The Journal of*
Neuroscience: The Official Journal of the Society for Neuroscience, 34(40), pp.13281–
13288. Available at: http://dx.doi.org/10.1523/JNEUROSCI.2894-13.2014.

Rietman, A.B. et al., 2017. Motor problems in children with neurofibromatosis type 1. *Journal of*
Neurodevelopmental Disorders, 9(1), p.19. Available at: http://dx.doi.org/10.1186/s11689-
017-9198-5.

Roth, J., Ber, R. & Constantini, S., 2019. Neurofibromatosis type 1-related hydrocephalus:
Treatment options and considerations. *World Neurosurgery*, 128, pp.e664–e668. Available
at: https://doi.org/10.1016/j.wneu.2019.04.231.

Ryu, H.-H. et al., 2019. Enriched expression of NF1 in inhibitory neurons in both mouse and
human brain. *Molecular Brain*, 12(1), p.60. Available at: https://doi.org/10.1186/s13041-
019-0481-0.

Sanchez-Ortiz, E. et al., 2014. NF1 regulation of RAS/ERK signaling is required for appropriate
granule neuron progenitor expansion and migration in cerebellar development. *Genes &*
Development, 28(21), pp.2407–2420. Available at:
http://dx.doi.org/10.1101/gad.246603.114.

Shin, J. et al., 2012. Zebrafish neurofibromatosis type 1 genes have redundant functions in
tumorigenesis and embryonic development. *Disease Models & Mechanisms*, 5(6), pp.881–
894. Available at: http://dx.doi.org/10.1242/dmm.009779.

Sorrentino, U. et al., 2021. Epilepsy in NF1: Epidemiologic, genetic, and clinical features. A
monocentric retrospective study in a cohort of 784 patients. *Cancers*, 13(24), p.6336.
Available at: http://dx.doi.org/10.3390/cancers13246336.

Su, W. et al., 2007. Mice with GFAP-targeted loss of neurofibromin demonstrate increased
axonal MET expression with aging. *Glia*, 55(7), pp.723–733. Available at:
http://dx.doi.org/10.1002/glia.20501.

Trachtman, J.N., 2010. Vision and the hypothalamus. *Optometry (St. Louis, Mo.)*, 81(2),
pp.100–115. Available at: http://dx.doi.org/10.1016/j.optm.2009.07.016.

Wickham, H., 2016. Data Analysis. In *Use R!*. Cham: Springer International Publishing, pp. 189–
201. Available at: https://doi.org/10.1007/978-3-319-24277-4_9.

Zhu, Y. et al., 2001. Ablation of NF1 function in neurons induces abnormal development of
cerebral cortex and reactive gliosis in the brain. *Genes & Development*, 15(7), pp.859–876.
Available at: http://dx.doi.org/10.1101/gad.862101.

608
609

Figure legends:

Figure 1: NF1 is expressed in mouse forebrain

(A) Schematics showing representative coronal sections of the mouse forebrain at adult and embryonic stages. NF1 protein expression in whole brain (B-E), hippocampus (E-I), and neocortex (J-M) across descending ages P35, P0, E17.5, E14.5, and E12.5. (B-F) NF1 was expressed in distinct zones of thalamus (t) and hypothalamus (hy) in the diencephalon. In the hypothalamus, NF1 clearly marked the paraventricular nucleus (PVN), suprachiasmatic nucleus (SCN), and supraoptic nucleus (SON) (B). Dotted boxes in (B-E) mark insets of parts of telencephalon, namely the neocortex (nctx) and hippocampus (h). (G-I) NF1 expression was observed in CA1, CA3, and hilar cells of hippocampus at E17.5, P0, and P35, but not in the dentate gyrus (DG). (J-K) NF1 was also expressed along the lateral and third ventricular linings, encompassing midline structures, namely choroid plexus (cp), cortical hem (ch), hippocampal primordium (hp), and ventral telencephalon (vtel) across ages (arrows, H-K). (L-P) NF1 is strongly expressed in all mature pyramidal neurons perinuclearly and in the ventricular/subventricular progenitor zones of neocortex across development. P, posterior; A, anterior; hp, hippocampal primordium; cc, corpus callosum; wm, white matter; 3V, third ventricle; lv, lateral ventricle; mz, marginal zone; cp, cortical plate; sp, subplate layer; iz, intermediate zone; vz, ventricular zone; svz, subventricular zone. Scalebars: 500µm (B-F); 100µm (G-K); 50µm (L-P).

Figure 2: NF1 colocalizes with mature neurons and apical progenitors of forebrain

(A-D'') NF1 colocalizes with mature neuronal marker NeuN in P35 (A-A'') and P0 (C-C'') neocortical plates (cp); insets show colocalizations indicated as yellow in both ages (B-B'', D-D''). (E-H) NF1 was expressed in both deep and upper neocortical layers. Perinuclear NF1 expression surrounded the nuclear expression of transcription factors Ctip2 and Satb2, respectively demarcating deep and upper layers. (I,J) NF1 is present perinuclearly in Sox2$^+$ apical progenitors at the ventricular zone (vz) but not in Tbr2$^+$ intermediate progenitor cells. (K-N) NF1 was expressed in both the developing cortical plate marked by Ctip2 and Sox2$^+$ apical progenitor zone at E14.5, but not in the Tbr2$^+$ subventricular zone (svz). (O,P) NF1 was expressed in the Sox2$^+$ apical progenitor zone at E12.5. Arrows in J,N,P indicate perinuclear NF1 expression in the apical progenitors. (Q-Y'') NF1 was expressed predominantly in the CA fields (CA1, CA3) and hilus of P35, P0, and E17.5 hippocampi, but no or low expression was seen in the granular neurons of dentate gyrus (DG) across ages. Hippocampal neuron-dense regions were denoted by NeuN and Ctip2 expression. (Z-AA'') At E14.5 and E12.5, the dorsal midline of the telencephalon showed NF1 expression in choroid plexus (cp), ventricular lining, and a subset of mature neurons derived from the cortical hem (ch) and the hippocampal primordium (hp). nctx, neocortex; wm, white matter; 3V, third ventricle; LV, lateral

ventricle; mz, marginal zone; cp, cortical plate; sp, subplate layer; iz, intermediate zone; vz, ventricular zone; h, hippocampus. Scalebars: 100µm (Q-Q'', T-T'', W-W''); 50µm (A-A'', R-S'', U-V''); 25µm (B-C'', E-I, K,M,O, X-AA''); 12.5µm (D-D'', F,H,J, L,N,P).

Figure 3: NF1 is expressed in subsets of cortical oligodendrocytes
(A-A'') NF1 was expressed in a subset of Olig2$^+$ oligodendrocytes in the main three cortical commissures of the P35 mouse brain - corpus callosum (A-B''), anterior commissure (C-D''), and hippocampal commissure (E-F''). In a magnified view, cream dashed circles mark NF1$^+$Olig2$^+$ double-positive oligodendrocytes, while red dotted circles mark Olig2$^+$ oligodendrocytes that did not express NF1 (B-B'', D-D'', F-F''). Scalebars: 50µm (A-A'', C-C'', E-E''); 25µm (B-B'', D-D'', F-F'').

Figure 4: NF1 is expressed in the mouse cerebellum
(A-D) Schematics showing sagittal sections of the mouse forebrain at adult, early postnatal, and embryonic stages, demonstrating how a complex lobular laminated structure like the cerebellum develops from a simple fate-specified neuroepithelium. Boxes mark insets at P35, P12, P3 and P0 indicating magnified views of folium and cerebellar ventricular zone. (E-F'') P35 and P12 cerebella showed strong NF1 expression in the Purkinje cell layer (pcl), cells in the white matter (wm), and in the deep cerebellar nuclei (DCN). (G-H'') NF1 was also expressed in migrating Purkinje cells and in developing DCN at perinatal ages, P3 and P0. (I-J'''') At E14.5 and E12.5, NF1 was expressed very strongly in immature cerebellar neurons of the nuclear transitory zone (NTZ), and in differentiating neurons near the cerebellar ventricular zone (cvz) and/or extra cerebellar ventricular zone. NF1 was also expressed at weaker levels in migrating rhombic lip-derived cells. NF1 was expressed in the hindbrain choroid plexus (cp) across the developmental stages, from E12.5 to the adult stage. Mes, mesencephalon; rl, rhombic lip; ml, molecular layer; gcl, granular cell layer; egl, external granular cell layer; igl, internal granular cell layer. Scalebars: 200µm (E,F,G,H,I,J); 50µm (E'-F''); 20µm (G'-H'', I'-J'').

Figure 5: NF1 is expressed in cerebellar neuronal lineages across development
(A-D'') NF1 was expressed in Purkinje cells across development, and arrowheads show its colocalization with Calbindin (Calb). Insets show magnified views of one Calb$^+$ Purkinje cell across development and its colocalization with NF1. (E-H'') Colabelling NF1 with NeuN marking cerebellar granule cells showed no or low coexpression across development. (I-L'') NF1 expression was predominant in the mature neurons of deep cerebellar nuclei (DCN) across development, colocalizing with NeuN (cream dashed circles). NeuN$^+$ mature cells were not observed here at P0. (M,N) Schematics demonstrate morphology and differential lamination of developing (<P21) and developed (>P21) mouse cerebella. Mes, mesencephalon; rl, rhombic lip; ml, molecular

690 layer; gcl, granular cell layer; egl, external granular cell layer; igl, internal granular cell
691 layer; bv, blood vessels; cp, choroid plexus; bg, Bergmann glia. Scalebars: 25µm (A-B",
692 E-F"); 10µm (C-D", G-L").
693

Figure 6: NF1 is expressed in the embryonic cerebellar ventricular progenitors
(A,D) Schematics demarcate different germinal zones of cerebellar anlage at E14.5 and
E12.5. At E12.5, the cerebellar ventricular zone is subdivided into the zone producing
Purkinje cells (deep blue) and the interneuron-producing zone (light blue). (B-B", E-E")
NF1 is expressed perinuclearly in Sox2$^+$ progenitors in the cerebellar ventricular zone at
E12.5 and E14.5. However, NF1 was not expressed in Pax6$^+$ secondary progenitor cells
derived from the rhombic lip. Mes, mesencephalon; NTZ, nuclear transitory zone; 4V,
fourth ventricle; cvz, cerebellar ventricular zone. Scalebars: 20µm (B-F").
702

Figure 7: NF1 is expressed in cerebellar glial lineages
(A-H") NF1 was expressed in a subset of Olig2$^+$ oligodendrocytes in both folium (A-D")
and deep cerebellar nuclei (E-H") at mature (P35) and developing (P12) mouse
cerebella (arrowheads). In magnified views (B-B",D-D",F-F",H-H"), cream dashed circles
mark NF1$^+$Olig2$^+$ double-positive oligodendrocytes, while red dotted circles mark Olig2$^+$
oligodendrocytes that did not express NF1. (I-N") NF1 was also expressed in the soma
and projection of the Bergmann glia across development. Colocalization with Blbp is
shown using arrowheads. ml, molecular layer; gcl, granular cell layer; egl, external
granular cell layer; igl, internal granular cell layer; pcl, Purkinje cell layer; wm, white
matter. Scalebars: 25µm (A-A", C-C", E-E", G-G", I-I", K-K", M-M"); 10µm (B-B", D-D", F-
F", H-H", J-J", L-L", N-N").
714

Figure 8: Nf1 expression in murine optic cup and olfactory bulb
(A,B) Schematics showing zones and cell layers of neurosensory organs, eye and
olfactory bulb. (C,F") NF1 is expressed in the developing mouse eye, predominantly in
the lens (L) and neuroretina (NR), which includes the photoreceptor layer (PRC), retinal
ganglionic cells and Müller glial cells in the broad retinal ganglionic cell layer (RGC) at
E17.5 and E14.5. Magnified insets (D-D", F-F") show colocalization of NF1 and Tbr2 in
the lens and a subset of intrinsically photosensitive RGCs. NF1 also showed wide
coexpression with Sox2 in PRC and retinal pigmented epithelium (RPE), but not with
the interneurons of neuroretina. NF1 colocalized with Tbr2 in the lens and a subset of
(G-J) NF1 was expressed strongly across the postnatal development of the olfactory
bulb. It showed strong expression in the mitral cell layer (MCL) of the main olfactory
bulb (MOB) and also in the accessory olfactory bulb (AOB). (K,L,O-P") Colabelling with
NeuN at P35 and P12 identified subsets of mitral cells that were double positive for NF1
and NeuN (arrowheads). (M,N, Q-R") Strong coexpression with Pax6 was seen in
posterior AOB (pAOB) at P0 and P3, but not in the other parts of MOB. IN, interneurons;

aAOB, anterior accessory olfactory bulb; EPL, external plexiform layer; GCL, granule
cell layer; GL, glomerular layer. Scalebars: 200µm (G,H,K,L); 100µm (C-C", E-E",
I,J,M,N); 50µm (O-O", P-P"); 25µm (Q-R"); 20µm (D-D", F-F").
733
Figure 9: Reanalysis of single nuclei RNAseq data showing expression of *Nf1* in
adult mouse brain
(A,F) UMAP plot for single nuclei RNAseq data from adult mouse cortex and
cerebellum, respectively. Transcriptionally distinct clusters were coloured based on cell
type. (B,G) UMAP plot illustrating the expression level of *Nf1* (red) across different
cortical and cerebellar clusters, respectively. (C-E) *Nf1* expression is higher in cortical
interneurons, differentiating OPCs, and excitatory neurons of adult murine cortex
compared to other neural populations. (H-J) In the adult murine cerebellum, *Nf1*
expression is enriched primarily in Purkinje cells, Golgi INs, Bergmann glia, and OPCs.
MGE_INs, medial ganglionic eminence interneurons; dMGE_INs, dorsal medial
ganglionic eminence interneurons; CGE_INs, caudal ganglionic eminence interneurons;
OPCs, oligodendrocyte precursor cells; MLI, molecular layer interneurons; PLI, Purkinje
layer interneurons; INs, interneurons; UBC, unipolar brush cells.
747

FIGURES

Figure 1

755
756
757

Figure 2

758
759
760
761

762
763 **Figure 3**
764
765
766

767
768
769
Figure 4
771

Figure 5

Figure 6

Figure 7

787
788 **Figure 8**
789

790
791
792   **Figure 9**
793