Mitochondrial protein translation is a heightened dependence and therapeutic vulnerability of chemo-refractory triple negative breast cancer

Mariah J. Berner[1,2,3], Steven W. Wall[1,2,3], Mokryun L. Baek[1,2,3], Audra Lane[1,2,3], Allison S. Greer[1,2,3], Karen Wang[1,2,3], Lacey E. Dobrolecki[1,2], Ivy Strope[1,2,4], Qian Zhu[1,2,4], Bing Zhang[1], Jonathan T. Lei[1,2,5], Michael T. Lewis[1,2,5,6], Gloria V. Echeverria[1,2,3,5,7]

[1]Lester and Sue Smith Breast Cancer, Baylor College of Medicine, Houston, TX, USA
[2]Dan L Duncan Comprehensive Cancer Center, Baylor College of Medicine, Houston, TX, USA
[3]Department of Medicine, Baylor College of Medicine, Houston, TX, USA
[4]Department of Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, USA
[5]Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX, USA
[6]Department of Radiology, Baylor College of Medicine, Houston, TX, USA
[7]Department of Radiation Oncology, Baylor College of Medicine, Houston, TX, USA

Correspondence: Gloria Echeverria (gloria.echeverria@bcm.edu)

ABSTRACT

Triple negative breast cancer (TNBC) patients harboring residual cancer burden following completion of conventional neoadjuvant chemo-immunotherapy regimens have poor relapse-free and overall survival rates despite recent advances in immunotherapies and antibody drug conjugates. We and others have demonstrated the requirement of mitochondrial function for survival of chemo-refractory TNBC, as well as its pervasive association with chemoresistance in human and patient-derived xenograft (PDX) cohorts. We sought to gain new mechanistic insights into the mitochondrial vulnerability of TNBC. Analyses of human and PDX mass spectrometry proteomics datasets revealed that mitochondrial protein translation-related signatures were among the top significantly associated with chemoresistance. Those signatures encompassed many core mitoribosome components as well as the mitoribosome accessory protein, Oxidase (Cytochrome C) Assembly 1-Like (OXA1L), which was consistently enriched in chemoresistant versus chemosensitive TNBCs across datasets. OXA1L, while not yet characterized in cancer, has been reported to be crucial for the termination of translation of the 13 mtDNA-encoded electron transport chain (ETC) proteins and for the insertion of those proteins, as well as nDNA-encoded ETC proteins, into the inner mitochondrial membrane. Together, those functions are crucial for the proper formation and function of the ETC. Therefore, we hypothesized that mitochondrial translation supported by OXA1L supports mitochondrial dependence and chemoresistance in TNBC. Knockdown (KD) of OXA1L in human TNBC cells reduced ETC protein levels, mitochondrial 'respirasome' supercomplex levels, ATP production, and oxidative phosphorylation (oxphos). Of note, OXA1L was required for the characteristic oxphos elevation induced by carboplatin (CRB), and KD significantly enhanced CRB sensitivity. To explore the translational potential of targeting the mitoribosome in TNBC, we leveraged the bacterial ancestry of mitochondria to repurpose the FDA-approved antibiotic tigecycline (TIG) as a chemo-sensitizing drug based on its mitoribosome inhibitory function. Direct measurement of mitochondrial nascent peptide levels revealed that, while CRB elevated mitochondrial translation, TIG potently diminished mitochondrial translation as monotherapy and when combined with CRB or docetaxel (DTX). Further, TIG abolished CRB-induced oxphos, decreased oxphos in combination with DTX, and significantly improved sensitivity to chemotherapies in human TNBC cell lines, PDX-derived spheroids, and in an *in vivo* PDX trial. These findings identify OXA1L-dependent mitochondrial translation and ETC formation as critical determinants of mitochondrial function that support TNBC chemoresistance, justifying further exploration of the clinical potential of repurposed antibiotics for TNBC.

DISCLOSURES

GVE is co-founder, Chief Scientific Officer, and an equity stakeholder of Nemea Therapeutics. G.V.E. formerly received sponsored research funding from Chimerix Inc. G.V.E. receives experimental compounds from the Lead Discovery Center of Germany and from Jazz Pharmaceuticals. MLB is a co-inventor at Nemea Therapeutics. MTL is a founder and limited partner in StemMed Ltd. and a manager in StemMed Holdings, its general partner. He is a founder and equity stakeholder in Tvardi Therapeutics Inc. Some PDX models, including BCM-4272 and BCM-7649, are exclusively licensed to StemMed Ltd., resulting in royalty income to MTL when used for commercial purposes. LED is a compensated employee of StemMed Ltd. Some PDX models, none of which are included in this study, are exclusively licensed to StemMed Ltd., resulting in royalty income to LED. All

58
59

INTRODUCTION

61
Triple negative breast cancer (TNBC) is an aggressive subtype lacking molecularly targeted therapy
options for the 75-85% of patients lacking deleterious BRCA1/2 aberrations[1,2]. TNBC is defined by the molecular
features it lacks, namely expression of the estrogen receptor, progesterone receptor, and overexpression or
amplification of human epidermal growth factor receptor 2. The mainstay treatment for TNBC is a three to six-
month course of neoadjuvant chemotherapy (NACT) consisting of several chemotherapeutic agents
administered simultaneously and/or sequentially prior to surgery[1,3]. There is no one universally accepted NACT
regimen[4]. Standard of care chemotherapies include agents that induce various forms of DNA damage (*e.g.*
carboplatin (CRB), Adriamycin (aka doxorubicin), and cyclophosphamide) and microtubule-perturbing taxanes
(*e.g.* docetaxel (DTX) and paclitaxel). NACT regimens were recently updated to include the immunotherapy
agent Programmed Cell Death 1 (PD-1) inhibitor, with a modest yet statistically significant improvement in
response rate[5]. Despite these many, lengthy treatments, residual tumors persist in approximately 45% of
patients[6,7]. Presence of residual tumor burden following NACT is a strong and significant correlate of poor
relapse-free and overall survival[6]. The five-year survival rate of metastatic TNBC is an unacceptable 12%[8]. Thus,
identifying the molecular dependencies of tumor cells that evade NACT is of utmost importance to improve
outcomes for women living with TNBC.
Mitochondrial metabolism has been reported to support chemotherapy resistance in hematologic
malignancies and solid tumors of breast, colon, prostate, pancreas, and other sites[9-15]. We and others have
established the importance of mitochondrial oxidative phosphorylation (oxphos) in TNBC[9,10,16-19]. Multi-omic
analyses of cohorts of TNBC patients[19,20] and orthotopic patient-derived xenograft (PDX) models[18] have revealed
that not only is oxphos the most upregulated metabolic pathway in TNBC relative to normal mammary tissue[20],
but also that oxphos is among the pathways most significantly and reproducibly associated with
chemoresistance. We demonstrated that inhibition of mitochondrial electron transport chain (ETC) complex I with
IACS-010759[9,21] or of mitochondrial fusion protein optic atrophy 1 with MYLS22[10,22] were both efficacious
preclinical approaches to delay TNBC relapse, although the mechanistic underpinnings of mitochondrial
dependence in TNBC remain poorly understood.
The mitochondrial genome encodes 37 genes, including 2 rRNAs, 22 tRNAs, and 13 mRNAs.
Mitochondria have their own replication, transcription, and translation machinery. The mitoribosome translates
the 13 mtDNA-encoded proteins, all components of ETC complexes I, III, IV, and V[23,24]. The remaining 79 ETC
proteins are encoded in the nucleus (nDNA). Oxidase (Cytochrome C) Assembly 1-Like (OXA1L) is a
transmembrane mitoribosome accessory protein insertase that has been shown in yeast and mammalian cells
to interact with the mitoribosome via its C-terminal tail, promoting translation termination and insertion of nascent
proteins into the inner mitochondrial membrane (IMM), where the ETC resides[24-26]. Furthermore, OXA1L is
responsible for the insertion of 81% of the nuclear-encoded ETC subunits, initially transported into the matrix,
into the IMM for the ETC to properly form[27]. Previous studies have demonstrated the importance of OXA1L for
the formation and function of Complexes I, IV, and V[27-29] in yeast and several human cell types, including
skeletal muscle, dermal fibroblasts, and induced pluripotent stem cells. Its potential role in the function of
Complexes II and III has not been established. OXA1L variants and polymorphisms have been linked with several
mitochondrial disorders[29], including childhood-onset mitochondrial encephalopathy[27], asthma[30,31], and
mitochondrial myopathy[32]. Disruption of OXA1L function contributes to mitochondrial dysfunction in Parkinson's
Disease[33] and in mitochondrial myopathy[32]. However, the functional relevance, regulation, and activity of OXA1L
in cancer has not been reported.
Herein, we identified mitoribosome function as a promising new, targetable metabolic vulnerability of
chemoresistant TNBC, supported by correlative analyses of patient and PDX cohorts and functional studies in
laboratory models. Analyses of multi-omic datasets[19,34,35] revealed a pervasive, positive association of levels of
mitochondrial ribosome proteins, as well as the accessory protein OXA1L, with TNBC chemoresistance. We
provide novel evidence that OXA1L supports protein levels spanning all five ETC complexes. Further, we find
mitochondrial translation and OXA1L are functional dependencies of chemoresistant TNBC. We went on to
demonstrate that repurposing conventional antibiotics to inhibit mitoribosome function may be a promising

approach to overcome the mitochondrial adaptations and survival of chemo-refractory TNBC cells.
111

METHODS
Analysis of publicly available datasets for molecular profiles of TNBC PDX and TNBC tumors
Quantitative chemotherapy responses measured as $\log_2$ fold change in tumor volume, and qualitative mRECIST
categorization after four weekly cycles of single-agent carboplatin or docetaxel, along with $\log_{10}$ transformed
mass spectrometry-based proteomic data from 50 TNBC PDXs, was downloaded from supplemental tables in
the Lei et al, 2025 manuscript[18]. Associations between proteomics data and quantitative responses were
assessed by Spearman correlation and Student's t-test was used for proteomics data and qualitative responses.
$\log_2$ transformed Tandem Mass Tag (TMT) mass spectrometry-based proteomics data for the CPTAC-TNBC
study was downloaded from supplemental tables in the Anurag et al, 2022 manuscript[19]. Baseline proteomic data
was available for 48 patients treated with neoadjuvant carboplatin plus docetaxel (17 pCR, 31 non-pCR),
including 14 paired samples collected after 48-72 hours of chemotherapy. For baseline analysis, Wilcoxon -rank-
sum test was used to compare proteins of non-pCR samples to pCR samples. Gene Set Enrichment analysis
with Reactome and MitoCarta geneset collections was performed using WebGestalt[36] with signed -log10 p-
values as input (positive sign if higher in non-pCR vs pCR, negative sign if lower). For paired on-treatment vs
baseline analysis, samples were analyzed using a moderated t-test with limma[37] using a paired design.
127

For breast cancer cell lines, $\log_2$-transformed TPM (Transcripts Per Million) and shRNA dependency scores
(D2_combined_gene_dep_scores.csv) were downloaded from the DepMap resource (https://depmap.org).
TNBC subtype information for cell lines was retrieved from Lehmann et al[38] and metabolic subtypes were
retrieved from Gong et al[20]. Associations between *OXA1L* RNA with TNBC subtypes and metabolic subtypes
were assessed by ANOVA and associations between *OXA1L* dependency scores with *OXA1L* RNA levels were
assessed by Pearson correlation.
134

MRP signature generation
The MRP signature comprises all MRPL and MRPS genes quantified in TNBC PDXs at the protein level.
Signature scores for individual PDX and clinical samples were calculated using single sample Gene Set
Enrichment Analysis (ssGSEA)[39] implemented through the ssGSEA2.0 tool
(https://github.com/broadinstitute/ssGSEA2.0) and reported as normalized enrichment scores (NES). NES
values were then used to assess associations with therapeutic response.
141

Cell Culture
The MDA-MB-231 and MCF10a cell lines were initially purchased from the American Type Culture Collection,
while the SUM159pt cell line was initially purchased from BioIVT. MDA-MB-231 cells were cultured in RPMI-
1640 glucose-free media (Gibco, 11879020) supplemented with 5 mM glucose (Gibco, A2494001), 10% heat-
inactivated FBS (R&D, S11550), and 1% antibiotic-antimycotic solution (Corning, 30-004-CI). SUM159pt cells
were cultured in the same base media as MDA-MB-231 cells and supplemented with the same glucose and
Antibiotic-Antimycotic concentrations. SUM159pt cells were also supplemented with 5% heat-inactivated FBS
(R&D, S11550), 5 mg/ml insulin (Santa Cruz Biotechnology, sc-360248), and 1 mg/ml hydrocortisone (Sigma-
Aldrich, H4001). MCF10a cells were cultured in MEGM Mammary Epithelial Cell Growth Medium BulletKit
(Lonza, CC-3150) except for the substitution of the GA-1000 with 1% antibiotic-antimycotic Solution (Corning,
30-004-CI). All cells used in the experiments were maintained at fewer than 20 passages.
153

Measurement of cell viability and confluency
Cells were counted using ViaStain AOPI staining solution (Revvity Health Sciences Inc. CS2-0106), which was
added to cells immediately prior to counting with an Nexcelom Cellometer (Revvity Health Sciences, Inc.). For
CellTiter-Glo assays, cells were measured using the Promega CellTiter-Glo™ Luminescent Cell Viability Assay
(PR-G7572) according to the manufacturer's instructions. Cell confluency was assessed longitudinally using an
Incucyte S3-C2 (Sartorius). Cells were seeded in a 96-well plate (Corning #35072) and five live-cell images per
well were taken with the 10x objective every four hours for the duration of the experiment. Cell confluency was
analyzed using the Incucyte image analysis software (Sartorius).
162

Authentication of cell lines and PDX models
Cell lines and all PDX models were tested quarterly for mycoplasma using the Universal Mycoplasma Detection
Kit (ATCC, 30-1012K). Identity of cell lines and PDX models was validated by short-tandem repeat (STR) DNA

fingerprinting at the M.D. Anderson Cancer Center Cytogenetics and Cell Authentication Core. For STR DNA
fingerprinting analysis, the Promega 16 High Sensitivity Kit (Catalog #DC2100) was used, and the results were
compared with online databases (DSMZ/ATCC/JCRB/RIKEN).
169
Gene silencing
Transient siRNA-mediated gene knockdowns were conducted with 5 nM siRNA targeting human *OXA1L* (5'-
CAGAAAAAACAUGGUAUUAAA-3', 5'-CUGGAAUUUAUGCAUGUUGAU-3', 5'-
AAGUUUUCCAGUCGAAUCAGA-3') and Mission siRNA universal negative control #1 (Sigma, SIC001-
10NMOL). Knockdown experiments were conducted using 10cm and 15cm plates, depending on the amount of
material required for the downstream analyses. For 10cm plates and 48-hour cell pellet harvests, cells were
seeded in 10cm at 300K, while for 120-hour cell pellet harvests, cells were seeded at 25K. For 15 cm plates and
48-hour harvests, cells were seeded at 750K, and for 120-hour harvests, cells were seeded at 75K.
178
For 10cm plates 48hr harvests, 7.5 µl of 10µM siRNA was combined with 30 µL of Lipofectamine RNAiMAX
(Invitrogen, 13778150) with 3 mL OPTI-MEM media (Gibco, 11058021) was prepared. Following 15 minutes of
incubation, complexes were added to cells that had been seeded 48-hour before at densities above. For 10 cm
plates with 120-hour harvests, the amount of Lipofectamine RNAiMAX was adjusted to account for the smaller
number of cells at the time of transfection, instead of 30 µL, 7.5 µL of Lipofectamine was used for cells that would
be harvested 120 hours post-transfection. For the 15 cm dish, lipofection reagents were scaled up 2.5 times for
the respective endpoint harvest time. Cells were harvested 48-120 hours post-transfection either on ice using a
cell scraper or with trypsin if cell counting was needed for downstream analyses.
187
In vitro **drug treatments**
Chemotherapies were prepared as 10 mM stock solutions, aliquoted and stored at -80°C. Docetaxel (Selleck
Chemicals S1148) was dissolved in DMSO and Carboplatin (Selleck Chemicals S1215) was dissolved in
complete media. Aliquots of these stock solutions were thawed at time of use, then discarded. Cells were seeded
two days prior to treatments such that, at the time of treatment, cells were ~30% confluent. Docetaxel was
administered at 1 - 5 nM and carboplatin 25 – 100 µM. Cells were collected 48-120 hours following treatment.
Tigecycline (Selleck Chemicals S1403) and chloramphenicol (Selleck Chemicals S1677) were prepared fresh in
DMSO. Tigecycline and chloramphenicol doses are indicated within each figure legend. For combination
treatments, drugs were administered simultaneously, and doses are indicated within each figure legend.
197
DNA extraction and relative mtDNA to nDNA quantification
Cell pellets were snap-frozen in liquid nitrogen and stored at -80°C. DNA extraction was conducted using the
Mag-Bind® Blood and Tissue DNA HDQ 96 Kit (OmegaTek M6399-00) as per the manufacturer's protocol. The
optional steps in the manufacturer's protocol: 10 µL of RNase A was added per sample, and the elution buffer
was heated to 70 °C prior to elution, were taken. The extracted DNA concentration was determined using the
NanoDrop 2000 (Thermo Scientific).
204
To quantify the relative amount of mtNDA per nDNA, qPCR primers for mtDNA[10] were used, while nuclear DNA
primers were used to quantify nDNA. The mtDNA primers were MT-ND1 (forward: 5′-
ATGGCCAACCTCCTACTCCT-3′, reverse: 5′-TAGATGTGGCGGGTTTTAGG-3'), and MT-ND6 (forward: 5′-
TGGGGTTAGCGATGGAGGTAGG, reverse: 5'-AATAGGATCCTCCCGAATCAAC-3'). The nDNA primers were
RGPD1 (forward 5′- GTGGAGCCACTGAGAATGGT-3′, reverse: 5′-GCATGCCTGGCTGATTTTAT-3'), and
FUNDC2P2 (forward 5′- TGAGTCAGTGGACCTTGCAG-3′, reverse: 5′-CAGAATGGTTTGCAAGCTGA-3').
qPCR was conducted using Universal SYBR Green Supermix (Bio-Rad, 1725121), 2-4 ng of extracted DNA,
and 1 µM of primers. The qPCR conditions are as follows: 50C for 2min, then 95 °C hot start for 10min, followed
by 40 cycles of 95 °C for 15 seconds and 60 °C for 1 minute. Relative copy number was then computed using
the relative $2^{-\Delta\Delta Cq}$ method.
215
RNA extraction and reverse transcription quantitative PCR (RT-qPCR)
Total RNA was extracted using TRIzol reagent (Invitrogen™, 15-596-018) according to the manufacturer's
instructions. Briefly, approximately 0.5×10^5 cells were collected in TRIzol reagent. RNA was precipitated with
isopropanol and washed with 75% ethanol. After resuspension in nuclease-free water at 55 °C for 10 min, RNA
was treated with DNase using the TURBO DNA-free™ Kit (Invitrogen™, AM1907) according to the
manufacturer's instructions. RNA was further purified using the RNA Clean & Concentrator kit (Zymo Research,

R1017). cDNA was synthesized using the SuperScript™ III First-Strand Synthesis System (Invitrogen™,
18080051). Quantitative PCR was performed using TaqMan™ Gene Expression Assays (Applied Biosystems™,
4369016) with FAM- or VIC-labeled probes for individual genes (Applied Biosystems™, 4448490). Real-time
qPCR was carried out on a CFX Opus Real-Time PCR System (Bio-Rad) using CFX Maestro software. Relative
gene expression was calculated using the $2^{-\Delta\Delta Cq}$ method. ΔCq was defined as Cq(target) − Cq($TUBB$). $\Delta\Delta Cq$
was calculated by subtracting the mean ΔCq of the control group (vehicle or siNT) from the ΔCq of each
experimental sample (Tigecycline treatment or siOXA1L).
229
Western blotting
Cell pellets were lysed in RIPA buffer (Thermo Scientific, 89901) containing protease inhibitor cocktail (Roche,
11836153001) and phosphatase inhibitor cocktail (PhosSTOP, Roche, 4906837001). Proteins were quantified
with Pierce™ BCA Protein Assay Kits (ThermoFisher Scientific 23225). Criterion TGX precast Midi Protein Gels
(4-20%, Bio-Rad, 5671095) were used for SDS-PAGE. Gels were transferred to nitrocellulose membranes (Bio-
Rad 1620112) using a Trans-Bolt Turbo Transfer System set at 2.5A, 25V, 7min (Bio-Rad, 1704150). EveryBlot
Blocking Buffer (Bio-Rad, 12010020) was used to block membranes for 10 minutes at room temperature and for
the dilution of antibodies. The following primary antibodies were used: anti-OXA1L (1:1,000, Proteintech, 21055-
1-AP), anti-beta-tubulin (1:1,000, Cell Signaling Technology, CST2146), anti-beta-actin (1:1,000, Cell Signaling
Technology, CST3700), anti-Phospho Histone H2AX (Ser139) (1:1,000, Cell Signaling Technology, CST2577),
anti-Tom70 (1:3,000, Proteintech, 14528-1-AP), anti-Mito Profile total OXPHOS Human WB antibody cocktail
(1:500, abcam, ab110411), anti-Complex I NDUFB8 (1:1,000, abcam, ab110242), anti-Complex I MT-ND3
(1:500, abcam, ab192306), anti-Complex II SDHB (1:1,000, abcam, ab14714), anti-Complex III UQCRC2
(1:4,000, abcam, ab14745), anti-Complex IV MT-COXI (1:2,000, Abcam, ab14705), anti-Complex IV MT-COXII
(1:1,000, abcam, ab110258), anti-Complex IV MT-COXIV (1:1,000, Cell Signaling Technology, CST4844), anti-
Complex V ATP5A (1:1,000, abcam, ab14748), anti-Complex V MT-ATP8 (1:1,000, Cell Signaling Technology,
CST96857), anti-caspase-3 (1:1,000, Cell Signaling Technology), and anti-cleaved caspase 3 (Asp175) (5A1E)
(1:1,000, Cell Signaling Technology).
Secondary antibodies were used at a dilution of 1:10,000 in blocking buffer: horseradish peroxidase-
conjugated anti-rabbit IgG H&L (Abcam, ab205718) and horseradish peroxidase-conjugated anti-mouse IgG
H&L (Abcam, ab205719). Membranes were incubated in secondary antibodies for one hour, with rocking, at
room temperature. Membranes were then washed with PBST three times each for 10 minutes. For detection,
Clarity Western ECL substrate (Bio-Rad, 1705060) was used, and images were captured on a ChemiDoc Touch
imaging system (Bio-Rad).
254
Mitochondrial fractionation
Mitochondrial fractions were extracted by differential centrifugation, as previously described[40]. Briefly, cell pellets
containing >5 million cells were resuspended in RSB hypo buffer (10 mM NaCl, 1.5 mM MgCl$_2$, 10 mM Tris-HCl,
pH 7.5) on ice to allow for swelling of the cells. Cell lysis was facilitated on ice by Dounce homogenization via
50 strokes with the small clearance pestle (DWK, 885302-0002) in a 2mL tissue grinding tube (DWK, 885303-
0002). Lysis was halted by the addition of MSH buffer (210 mM mannitol; 70 mM sucrose; 5 mM Tris-HCl; and
1 mM EDTA, pH 7.5). RSB and MSH buffers were supplemented with cOmplete Protease Inhibitor Cocktail
(Sigma, 11836170001) and PhosStop-phosphatase inhibitor tablets (Roche, 04906837001) to inhibit
phosphatase and protease activity, respectively, and all centrifugation steps were performed in a prechilled (4°C)
centrifuge. Cell suspensions were centrifuged at 600 × g for 5 minutes, and the supernatant was transferred to a
new tube by pipetting. Supernatants were then centrifuged at 1,200 × g for 5 minutes to pellet any remaining
nuclei from the sample. The supernatants were again collected, then centrifuged at 7,000 × g for 10 minutes to
pellet the mitochondria-enriched fraction. Supernatants containing cytosolic fractions were collected in fresh
tubes. Cytoplasmic fractions were further processed by centrifuging at 16,000 × g for 20 minutes to pellet cellular
debris. The supernatant (cytosolic protein) was then transferred to a new tube and stored at -80°C.
270
271 The mitochondria-enriched pellet was resuspended in MSH buffer and spun at 7,000 × g for 10 minutes to wash
272 mitochondria. After discarding the wash supernatant, the pellet was resuspended for its final wash in MSH buffer
273 and spun at 10,000 × g for 10 minutes. The supernatant was discarded, and the crude mitochondrial pellet was
274 resuspended in 50-100 µl MSH buffer and proteins were quantified using the Pierce BCA Protein Assay Kit
275 (Thermo-Fisher, 23223), then stored at -80°C.
276
Blue native polyacrylamide gel electrophoresis (BN-PAGE)

25-60 µg of mitochondria in MSH suspension from the fractionation were aliquoted and pelleted at 10,000 × *g* for
10 minutes at 4°C. After removing any remaining supernatant and the MSH buffer, mitochondrial pellets were
resuspended in 1X NativePAGE Sample Buffer (Invitrogen) with 1% digitonin (8 g/g; Invitrogen) by gentle
pipetting (~20 times). Samples were incubated on ice for 15 minutes to solubilize the mitochondrial proteins. The
samples were then centrifuged in a precooled centrifuge (4°C) for 30 minutes at 18,000 × *g* and supernatants
were transferred to new tubes. Next, 1-2 µL of 0.5% G-250 Coomassie dye (Invitrogen) was added to the
samples. Samples were then loaded into pre-cast NativePAGE Novex 3-12% Bis–Tris gels in the XCell SureLock
Mini-Cell system (Invitrogen) according to the manufacturer's protocol. Briefly, the entire apparatus was
assembled and placed in an ice bath. The wells of the gel were filled with dark blue cathode buffer (1X
NativePAGE running buffer and 1X cathode additive), then samples and NativeMark unstained protein standard
(Invitrogen) were loaded. The inner chamber was filled with dark blue Cathode buffer, and the outer chamber
was filled with anode buffer (1X NativePAGE running buffer) until it was 1/3 full. The gel was run at 100V for 30
minutes followed by 150V for another 30 minutes. The dark blue cathode buffer was then replaced with the light
blue cathode buffer (1X NativePAGE running buffer and 0.1X Cathode Additive), and the gel was run for an
additional 90-120 minutes at 250 V.
293
Imaging Coomassie-stained BN-PAGE gels
The gel was transferred into a container with 4°C DI water, quickly rinsed with fresh 4°C DI water, then incubated
in Fix/Destain solution (40% methanol + 7% acetic acid) for 1 hour on an orbital shaker at room temperature.
Halfway through the incubation, the solution was refreshed. After incubation, Fix/Destain solution was removed
and replaced with DI water. The gel was incubated in DI water overnight on an orbital shaker at room
temperature. White-light images were taken on ChemiDoc Touch imaging system (Bio-Rad).
300
Immunoblotting of BN-PAGE gels
Proteins from BN-PAGE gels were transferred to a PVDF membrane using 1X NuPAGE Transfer Buffer
(Invitrogen) and the Trans-blot Turbo transfer system (Bio-Rad) run at 2.5 amps for 18 minutes. Membranes
were rinsed in DI water then incubated in 8% acetic acid for 15 minutes to fix the proteins. Membranes were
then washed with DI water and allowed to dry air for ~30 minutes or until dry. Once dry, membranes were
rehydrated with 100% methanol, which also removed the blue Coomassie staining, then washed in DI water
before blocking with Every-blot blocking buffer (Bio-Rad) for 10 min. Membranes were then incubated at 4°C
overnight in primary antibody. After washing with PBS including 0.1% Tween-20 (PBST) three times for ten
minutes, HRP-linked secondary antibody was added at room temperature for one hour. Preparation of primary
and secondary antibodies and their dilutions were done the same way as for western Blotting. Membranes were
subjected to three more ten-minute PBST washes, then developed using Clarity Western ECL substrate (Bio-
Rad). Images were taken on ChemiDoc Touch imaging system (Bio-Rad).
313
Seahorse analysis of mitochondrial function
Mitochondrial function was assessed using the Agilent Technologies XF Cell Mito Stress Test (Agilent
Technologies, 103015-100) and the Seahorse XFe96 extracellular flux analyzer. Briefly, on the day of the assay,
viable cells were counted by ViaStain AOPI staining solution (Revvity Health Sciences Inc. CS2-0106) and
seeded at a density of 25,000 to 50,000 per well in 180 µl of RPMI seahorse assay media (Agilent Technologies
103576-100) on XFe96 cell culture microplates (Agilent Technologies, 103794-100) that had been previously
treated with Poly-D-lysine hydrobromide (MedChem Express Cat # HY-139201A). Cells were then incubated in
a CO2-free incubator at 37°C for one hour, and the calibrant cartridge was put into the Seahorse XFe96
extracellular flux analyzer for calibration. Next, the cell plate was put into the analyzer to run the assay. For the
assay, mitochondrial inhibitory drugs were prepared as per the manufacturer's guidelines in RPMI Seahorse
Assay Media; oligomycin (1.5 µM), FCCP (1 µM), and rotenone A/antimycin (0.5 µM).
325
Clonogenicity measurement
Cells were treated in 10 cm dishes with tigecycline at 50 mM or at the IC_{50} for each respective cell line (MDA-
MB-231 8 mM or SUM159 3 mM). Forty-eight hours after treatment, cells were counted using ViaStain AOPI
staining solution (Revvity Health Sciences Inc. CS2-0106). Then a 10,000 cells per 10 mL cell dilution was made
and either 100 or 50 cells per condition were plated in 6-well plates in triplicate. The media was refreshed every
two days. Seven days after the initial replating, cells were stained with crystal violet (Sigma-Aldrich HT90132),
washed, and left to dry. Colonies were manually counted the following day.
333

Mitochondrial fluorescent non-canonical amino acid tagging (Mito-FUNCAT)

The method was adapted from Yousefi et al, 2021[41] and Malik et al, 2023[42]. Cells were plated on coverslips (Thorlabs CG15) placed in a 6-well dish (Corning 3516) that had been previously sterilized with 70% ethanol and treated with 1X Poly-D-lysine hydrobromide prepared by the company manual (MedChem Express HY-139201A). Chemotherapy treatments were conducted two days after seeding, then two days after treatment, cells were labeled, stained, and mounted.

To label the cells, the media was aspirated, and the cells were washed with PBS. Cells were then given customized methionine-free complete media (Thermo Fisher Scientific, removed methionine from RPMI A1451701) and respective translation inhibitors for one hour. To inhibit cytosolic translation, 100 mg/ml cycloheximide (Research Products International C81040-1.0) was used, and to inhibit mitochondrial translation, 50 mM tigecycline (Selleck Chemicals S1403) was used. For the negative control, both inhibitors were administered at the previously listed concentrations. Next, 500 mM HPG (click-iT metabolic labeling reagent C10186, Invitrogen) was added to the cells for 30 minutes. After labeling, cells were incubated on ice for two minutes in buffer A (dH$_2$0 base, 10 mM HEPES, 10 nM NaCl, 5 mM KCl, 300 mM sucrose, and 0.015% (w/v) digitonin). Buffer A was then removed, and cells were incubated in buffer A.1 (dH$_2$0 base, 10 mM HEPES, 10 nM NaCl, 5 mM KCl, and 300 mM sucrose) for 15 seconds. Cells were then fixed with 4% paraformaldehyde (Alfa Aesar 47392-9M) for 30 minutes, with gentle rocking at room temperature. After fixation, cells were washed with PBS for five minutes, followed by incubation in 100 mM NH$_4$Cl in PBS for 15 minutes. This was followed by three washes each for five minutes in staining solution (PBS base, 5% BSA, 5% tryptone, and 0.1% Triton X-100). Cells were washed briefly with 3% BSA in PBS. Then, the incorporated methionine analog, HPG, was "clicked" to a fluorescent tag using the Click-iT cell reaction buffer kit (Thermo Scientific C10269) based on the manufacturer's instructions including 3 µM Azide (AlexaFluor 594 Azide, Thermo Scientific A10270) for 30 minutes at room temperature. Afterwards, cells were washed with 3% BSA in PBS.

To stain the mitochondria, cells were incubated in primary CoxIV antibody (Abcam Cat# ab33985) for one hour at room temperature (1:200 in 3% BSA). After incubation, cells were washed three times in 3% BSA for five minutes each. Then the cells were incubated in secondary antibody, goat anti-mouse Alexa Fluor 488 (Jackson ImmunoResearch Laboratory, Cat# 115-547-003, 1:500 dilution in 3% BSA). After incubation, cells were washed with 3% BSA and stained with DAPI for five minutes at room temperature. Following this, the cells were washed with 3% BSA and then incubated for five minutes in PBS including 500 mM NaCl. Last, cells were washed with PBS three times, then mounted on slides (Fisher Scientific 12-550-15) with mounting media (Invitrogen Prolong Glass Antifade Mountant P36982). Slides were stored at room temperature and protected from light until imaging, which was completed within 14 days.

Imaging was conducted with a Zeiss LSM 880 with Airyscan confocal microscope using a 63X oil objective. The channels and powers of the lasers used for imaging were as follows: channel 405 at 2.0 W, channel 488 at 4.5 W, and channel 561 at 15.0 W. The acquisition mode frame size was 1912x1912, and the zoom was set at 2. Z-stacks were taken with slices at 0.16 µM intervals. Images were then processed using Fiji ImageJ.

Mito-FUNCAT image analysis

Fluorescence image analysis was performed using skimage, numpy, and scipy libraries to quantify colocalization of mitochondrial translation. Multi-channel TIFF images were processed to extract HPG-Azide 494 (Channel 1, marking nascent protein synthesis), COXIV 488 (Channel 2, labeling mitochondria), and DAPI (Channel 3, identifying nuclear regions). To ensure consistent background removal, thresholding was first applied on raw images using an Otsu-derived intensity cutoff from the HPG-only control (positive signal). Pixels below this threshold were masked before subsequent processing. After thresholding, fluorescence intensities were normalized using the maximum signal intensity of the retained pixels to prevent artificial enhancement of low signal. Nuclear and cytosolic compartments were segmented using thresholding on the DAPI channel, defining the nucleus as DAPI-positive regions and the cytoplasm as the inverse. Colocalization analysis was conducted separately for nuclear and cytoplasmic compartments. Pearson's correlation coefficient was calculated between Channels 1 and 2 to assess global signal correlation. Percent colocalization in the nucleus was defined as the proportion of overlapping positive pixels within the DAPI mask, normalized to the total DAPI-positive area. Percent colocalization in the cytoplasm was calculated as the proportion of overlapping signal outside the DAPI mask, normalized to the total cytoplasmic area where either Channel 1 or Channel 2 exhibited positive signal. Mitochondrial translation per cell was determined by combining the positive pixels from Channels 1 and 2 that

390 overlap within the cytoplasm and then was normalized to cell count. The proportion of actively translating
391 mitochondria was determined by the difference between the overlapping positive pixels from channel 2 positive
392 pixels (COXIV) divided by channel 2 positive pixels in the cytoplasm.
393
PDX *ex vivo* spheroid studies
Tumor chunks were digested in Dulbecco's modified Eagle's medium (DMEM):F12 (Cytiva HyClone,
SH30023.01) supplemented with 5% FBS, 0.45% collagenase A (Roche, 1088793), 0.086% Hyaluronidase
(Sigma-Aldrich, H3506), and 2% BSA (Sigma-Aldrich, A9418) for 4 hours at 37°C while rotating. After the initial
digestion period, cells were centrifuged at 170 × g for 8 minutes at room temperature, resuspended in red blood
cell (RBC) lysis buffer (Invitrogen 00-433-57), and incubated at room temperature for 3 minutes with periodic
inversion of the tube. At the end of the incubation, the RBC buffer was inactivated with a matched volume of
PDX epithelial cell growth medium (DMEM:F12 (Cytiva Hyclone, SH30023.01) supplemented with 5% FBS and
1X antibiotics). Cells were centrifuged at 120 × g for 5 minutes at room temperature, and the supernatant was
removed. Cells were then resuspended in trypsin (Corning 25-053-CI) for 3 minutes at 37°C, inverted periodically.
The trypsin was quenched with 2X volume of PDX media and spun down at 170xg for 5 minutes at room
temperature. The supernatant was aspirated and cells were resuspended in Dispase solution (Stemcell
Technologies, NC9886504) supplemented with DNase I (1mg/ml, Stemcell Technologies, NC9007308) for 5
minutes at 37°C with periodic inverting. The dispase was inactivated with an equal volume of PDX media, and
cells were collected by centrifugation at 170xg for 5 minutes at room temperature. The supernatant was aspirated
and the cell pellet was resuspended in PDX media and passed through a 70 μm filter followed by a 40 μm filter.
The initial cell count was determined, and MACS separation was initiated to enrich for human PDX tumor cells
by depleting mouse stromal cells. MACS separation (Miltenyi Biotec 130-104-694) was performed according to
the manufacturer's protocol. The final cell count of the enriched human cells was obtained, and 5,000 cells were
plated per 50 μl in growth medium in 96-well low-attachment plates (Corning 3474). Cells were treated with TIG
and CRB 24 hours post-seeding and CTG assays were conducted 48 hours post-treatment.
415
416 **Animal Studies**
417 Animal studies were conducted under approved BCM IACUC protocol AN-8243, which was in accordance with
418 the National Institutes of Health *Guide for the Care and Use of Laboratory Animals*. Following the Association
419 for Assessment and Accreditation of Laboratory Animal Care guidelines, mice were ethically euthanized.
420
421 For spheroid cultures, TNBC PDX tumor pieces were orthotopically implanted into Scid/Bg mice and left to grow
422 until 800-1200 mm^3. Tumors were collected, finely minced with scalpels (VWR 100499-580 and 100499-578),
423 and processed for spheroid culture.
424
The PIM001-P PDX model was established and characterized at the University of Texas MD Anderson Cancer
Center[9,43]. Briefly, cryo-preserved cell suspensions were thawed, AO/PI counted using a Cellometer K2
(Nexcelom Bioscience), and viable cells were resuspended in a 1:1 volume mixture of Matrigel (Corning 354234)
and complete cell growth medium (Dulbecco's modified Eagle's medium: F12, Cytiva Hyclone, SH30023.01)
supplemented with 5% FBS. A total volume of 20 μl of 400,000 cells were unilaterally injected into the fourth
mammary fat pad of each female Scid/Bg [C.B-17/IcrHsd-PrkdcscidLystbg J] mouse (Envigo).
431
The chemotherapeutic docetaxel (DTX; Hospira, 0409-0201-10) was supplied as a solution containing
polysorbate 80 NF, 4 mg anhydrous citric acid USP, 23% v/v dehydrated alcohol USP, q.s., with polyethylene
glycol 300 NF at 20 mg/kg. DTX was administered by i.p. injection at a dose volume of 2 mL/kg. The
chemotherapeutic carboplatin (CRB; TEVA Pharmaceuticals, 0703-4248-01) was supplied at 10 mg/ml in water
for i.p. injection at 50 mg/kg at a dose volume of 5 mL/kg. The antibiotic tigecycline (TIG; Selleck Chemicals,
S1403) was supplied as 200 mg aliquots, which were then dissolved at 25 mg/mL in 10% DMSO + 90% sterile
water and sonicated. The i.p. injection was 100 mg/kg at a dose volume of 4 mL/kg. Three cycles of TIG were
given, consisting of daily administration for 10 days, followed by a 3-day break.
440
Immunohistochemistry (IHC)
Harvested tumors were immediately sliced with scalpels and fixed in 10% formalin (Sigma HT501128-4L) at
room temperature for 48 hours, followed by three 10-minute washes with PBS. Tissues were stored at 4°C in
70% ethanol until processing the fixed tumors into paraffin blocks, from which 3 μm sections were cut. IHC
staining included routine hematoxylin and eosin (H&E; Epredia, 72711 and 71311) staining. Additional IHC was

conducted with antibodies against Ki67 (Dako Cat# M70240 at 1:200) and CC3 (Cell Signaling Technologies,
CST9661 at 1:50) following antigen retrieval with 0.1M Tris-HCL 15 min at full pressure (above 90°C) with a
pressure cooker.
449

Statistical analysis
Data shown are representative of at least three biological experiments. The bar graphs indicate the mean, and
error bars represent the standard error of the mean (SEM). GraphPad Prism and R software, as indicated within
the figure legends, were used for some statistical testing. Outliers were identified by the ROUT method using
Prism and removed. For multiple comparisons, the post-hoc Dunnett's test was used after the one-way ANOVA.
When data were not normally distributed, Mann-Whitney tests were used.
456

458 **RESULTS**
459

Mitochondrial translation and OXA1L are associated with TNBC chemotherapy resistance in human and
PDX cohorts.
Publicly available proteomic data from a Clinical Proteomic Tumor Analysis Consortium (CPTAC) TNBC
study demonstrated oxphos was among the most significant Hallarmks50 GSEA pathways associated with
resistance to DTX+CRB in tumors prior to therapeutic exposure[19]. In light of our prior findings that mitochondrial
function is a heightened vulnerability of chemo-refractory TNBC[9,10], we sought to extend analyses of those data
to gain more in-depth insights into correlates of TNBC chemoresistance (Fig S1A-C). We found mitochondrial
translation-related pathways were significantly enriched in pre-treatment tumors of patients who went on to
harbor residual cancer burden (RCB) compared to those with a complete pathologic response (pCR; Fig 1A,
S1A-B). A subset of those patients provided on-treatment biopsies three days after initiation of DTX+CRB. Again,
mitochondrial translation-related pathways were among the significantly enriched pathways in matched on-
treatment relative to pre-treatment tumors (Fig 1A, S1C).
Complementing this, proteomic profiling of orthotopic TNBC patient-derived xenograft (PDX) tumors from
a large-scale preclinical chemotherapy trial[18] corroborated that prior to therapeutic exposure, the levels of
proteins related to mitochondrial translation and ETC assembly were significantly higher in PDX models that
were refractory to DTX and CRB than in those that were sensitive (Fig 1B). In fact, the Reactome 'mitochondrial
translation termination' pathway was the top pathway associated with chemotherapy resistance among the 40
significantly upregulated pathways[18] and was also enriched in the CPTAC dataset. Among the clinical and PDX
datasets, there were 63 total leading-edge proteins in this pathway upregulated in chemoresistant samples (data
not shown). Most proteins were mitochondrial ribosomal proteins (MRPs): 31 large subunit proteins (MRPLs)
and 24 small subunit proteins (MRPSs), as well as mitoribosome recycling factor (MRRF), another component
of the core mitoribosome machinery. The mitoribosome accessory protein OXA1L was, aside from core
mitoribosome components, the protein most strongly associated with chemotherapy resistance that overlapped
between the patient and PDX datasets (Fig. 1C-D). Additionally, OXA1L protein level significantly increased in
the subset of the TNBC patient tumors biopsied on-chemotherapy treatment compared to matched pre-treatment
tumors (Fig. 1D). We generated an MRP signature but did not observe a significant association with
chemoresistance (Fig S1D). Given the dual functions of OXA1L in mitochondrial translation and insertion of ETC
proteins into the IMM for ETC formation, these findings led us to hypothesize that mitoribosome function, largely
supported by OXA1L, may be a major contributor to the metabolic rewiring[9,10] that supports TNBC
chemoresistance.
490

OXA1L is required for ETC formation and mitochondrial function in human TNBC cells.
Transient siRNA-mediated knockdown (KD) of *OXA1L* in MDA-MB-231 and SUM159pt human TNBC
cells decreased the abundance of mtDNA and nDNA-encoded ETC proteins comprising all five complexes, while
not affecting cell confluency over time (Fig. 2A, S2A, & S3A). Notably, the reduction in protein levels became
apparent five days after transfection, but not as early as two days, likely because the pool of existing ETC
proteins had not yet turned over. The effect of *OXA1L* KD was specifically at the protein level, with no significant
reduction in mtDNA or mtRNA levels observed in either cell line, except for siRNA #1 that modestly decreased
mtDNA content in SUM159pt cells (Fig. S3B-C). Concomitantly, oxygen consumption rate (OCR, *i.e.,* oxphos)
significantly decreased, and extracellular acidification rate (ECAR, *i.e.* glycolysis) tended to non-significantly
increase five days following *OXA1L* KD (Fig. 2B & S2B). ATP levels also significantly decreased upon KD (Fig
2C & S2C). Together, these findings indicate that *OXA1L* supports oxphos through protein-level regulation of

mtDNA- and nDNA-encoded ETC proteins in human TNBC cells.

Although *OXA1L* KD potently reduced the abundance of several key ETC proteins, some were only
modestly decreased (Fig. 2A and S2A), possibly due to their inherent stability and turnover rates. This led us to
ask how the formation of ETC complexes and supercomplexes (SCs) was affected by *OXA1L* loss. Blue native
PAGE of proteins extracted from purified mitochondria, followed by Coomassie staining, showed an overall
disruption of SC formation (Fig. 2D & S2D). Immunoblotting for UQCRC2 (CIII) revealed substantial disruption
of partitioning into SCs, including respiring SCs (aka respirasomes), which have been reported to support oxphos
efficiency[44,45] (Fig. 2E & S2E). Furthermore, *OXA1L* KD dramatically impaired full assembly and dimerization of
ATP synthase (Fig. 2F & S2F). Together, these findings reveal OXA1L is required for proper ETC protein levels,
SC formation, and oxphos in human TNBC cells.

We used the Characterized Cell Line Encyclopedia gene dependency database results from RNAi
screens[46] to compare *OXA1L* dependence among breast cancer cell lines. Within TNBC, there was no significant
correlation between *OXA1L* mRNA levels and the dependency score (Fig S4A-B), suggesting that mitochondrial
translation dependence is influenced by additional factors. Further, there was no significant association between
OXA1L RNA level or *OXA1L* dependency score with transcriptomic[38] (Fig. S4C-D) or metabolic[20] (Fig. S4E)
subtypes of TNBC (Fig S4A-E). Importantly, non-tumor immortalized mammary epithelial MCF10a cells had a
lower OXA1L dependency score than all the TNBC cell lines included, except for the CAL148 cell line (Fig. S4A).
KD of *OXA1L* in MCF10a cells had no significant effects on cell growth, ETC protein levels, or OCR[10] (Fig. S4F-
H). Together, these data suggest a selective dependency on OXA1L in TNBC, supporting a potential therapeutic
window.
522
OXA1L promotes oxphos adaptation and survival of chemotherapy refractory TNBC cells.

OCR was significantly elevated by high-dose CRB (Fig 3A, S5A, S6A), consistent with our previous report
of oxphos elevation by several DNA-damaging chemotherapies[10]. *OXA1L* KD abolished both the high- and low-
dose CRB-induced OCR increases (Fig. 3A, S5A, S6A). Further, *OXA1L* KD potently reduced the abundance of
mtDNA- or nDNA-encoded ETC proteins in the presence and absence of CRB (Fig. 3B & S5B). Moreover,
OXA1L KD significantly increased sensitivity to CRB treatments (Fig 3C & S5C).

Platinum chemotherapies are often combined with taxanes in the neoadjuvant setting for TNBC. Because
OXA1L was identified as a leading edge protein associated with resistance to CRB as well as DTX[18] in the PDX
cohort, we also sought to test the contribution of OXA1L to DTX response. DTX caused a slight elevation of OCR
in SUM159pt cells and no change in OCR in MDA-MB-231 cells. In both cases, *OXA1L* KD reduced oxphos
when combined with DTX (Fig 3A, S5A, S6A). OXA1L significantly enhanced DTX sensitivity in both cell lines
(Fig 3C & S5C). It is important to note that when DTX and CRB were combined, as is commonly used in the
clinic, OCR was elevated, albeit to a lesser extent than with CRB alone (Fig. 3A, S5A, S6A). Then with *OXA1L*
KD, we observed a diminished OCR increase when DTX was combined with CRB. Together, these findings
demonstrate the requirement for OXA1L for both oxphos and chemoresistance in chemo-refractory TNBC cells.
538
Repurposed antibiotics disrupt mitochondrial translation and metabolic adaptations in TNBC cells.

Several conventional antibiotics inhibit the bacterial ribosome, which shares high structural homology
with the mammalian mitoribosome[47] based on the theory of endosymbiosis[47]. Previous research suggests
inhibiting mitochondrial translation using Tigecycline (TIG), an FDA-approved antibiotic used for treatment of
sepsis, has anticancer activity in preclinical models of liver, breast, and ovarian cancer[47-51], although the
mechanistic relationship with mitochondrial translation remains unexplored. We hypothesized that combining
TIG with conventional chemotherapies would disrupt the pro-survival mitochondrial adaptations elicited by
chemotherapies. It is important to point out that, unlike OXA1L, TIG is only expected to disrupt mitoribosome
activity and does not have a documented role in the insertion of nDNA-encoded ETC proteins into the IMM. We
found the IC$_{50}$ of TIG in human TNBC cells to be 3-8 μM (Fig. 4A & S7A), a six to 16-fold lower dose than used
previously in breast cancer *in vitro* models to perturb cancer stem-like cell features[47]. We were able to
recapitulate the previously observed potent inhibition of tumor initiation capacity with 50 µM TIG, but not with the
IC$_{50}$ doses (Fig. S8A-D). Interestingly, TIG treatment did not appear to increase apoptosis, with a notable
reduction in caspase 3 cleavage compared to vehicle treated cells (Fig. 4B, S7B), consistent with its seemingly
cytostatic effects on TNBC cells quantified longitudinally (Fig. S8E).

The abundance of ETC proteins (both mtDNA- and nDNA-encoded, comprising all five ETC complexes),
but not mtDNA or mtRNA levels (except for *MT-ND1* in MDA-MB-231 cells), decreased with only 48 hours of TIG
treatment (Fig. 4B-D & S7B-D) in both cell lines. Interestingly, we observed significant increases in several

mtRNA transcript levels in SUM159pt cells upon TIG treatment, perhaps reflecting an adaptive response to
translation inhibition. ETC respirasome formation was significantly decreased by TIG (Fig 4E-F & S7E-F), and
we hypothesize this may have contributed to the reduction of nDNA-encoded ETC proteins induced by TIG.
Indeed, there is evidence that SC assembly supports the stability of proteins within them[44]. Concordantly, TIG
significantly reduced OCR (Fig. 4G, S7G).
To directly test the ability of TIG to inhibit mitochondrial translation, we conducted Mito-FUNCAT[41,42],
enabling quantification of nascent peptide levels based on labeling of the methionine analog HPG. Mitochondrial
translation was measured by localizing HPG within mitochondria while inhibiting cytosolic translation with
cycloheximide (CHX). As expected, CHX inhibited cytosolic translation while high-dose TIG did not (Fig S9A).
Furthermore, one hour of treatment with high-dose TIG reduced mean mitochondrial translation per cell (number
of overlapping pixels of HPG-azide with COXIV) by approximately 86% (Fig. S9A), while two days of TIG at its
IC_{50} reduced mean mitochondrial translation per cell by approximately 36% (Fig. 5A). Two days of CRB
significantly increased the proportion of mitochondrial area harboring active translation, whereas TIG significantly
reduced this proportion as monotherapy or combined with CRB. Further, TIG significantly reduced total
mitochondrial translation per cell when combined with either chemotherapy relative to the chemotherapy agent
alone.
This imaging approach afforded us the ability to appreciate heterogeneity of mitochondrial translation
within and between cells. In the case of CRB, the total amount of mitochondrial translation per cell was highly
heterogeneous, with some cells actively producing high levels of nascent mitochondrial peptides while other cells
were seemingly silent. In converse, DTX treatment resulted in a relatively homogeneous amount of mitochondrial
translation per cell, but highly heterogeneous proportions of actively translating mitochondria. In other words,
while most DTX-treated cells had similar total amounts of mitochondrial translation, some cells had larger
proportions of non-translating mitochondria than did others. These patterns may reflect differences in how
mitochondrial translation is regulated across mitochondria within individual cells
Concordantly, TIG combined with chemotherapy decreased ETC protein abundance and disrupted CRB-
induced OCR elevation (Fig. 5B-C, S9B, S10A-B). TIG, combined with DTX, reduced OCR and ETC protein
levels (Fig. 5D, S9C, S10C). Together, these data support the notions that human TNBC cells surviving CRB
harbor increased mitochondrial translation and mitochondrial function and that TIG rapidly and potently disrupts
this effect.
586
Mitoribosome inhibition enhances chemotherapy response in TNBC models.
TIG significantly increased the sensitivity of MDA-MB-231 and SUM159pt cells to CRB and DTX (Fig. 6-
A-D). An additional FDA-approved antibiotic, chloramphenicol, could similarly disrupt oxphos and
chemoresistance in the context of CRB (Fig. S11A-B). Furthermore, as observed with *OXA1L* KD in the TNBC
cells, TIG reduced ETC protein levels but did not inhibit OCR of MCF10a cells (Fig. S12A-B). Additionally, only
the highest dose of TIG tested moderately sensitized MCF10a cells to CRB, while all other doses left CRB
sensitivity unaltered or even slightly reduced (Fig. S12C).
Next, we tested the efficacy of TIG monotherapy and in combination with CRB in spheroid cultures from
freshly digested tumors that had been depleted of mouse stroma from three orthotopic PDX models, representing
three TNBC patients: PIM001-P[9,43], BCM-4272, and BCM-7649[52]. TIG and CRB, as monotherapies, were
efficacious across all three models to varying degrees. TIG maintained potency when combined with CRB in
BCM-4272 and BCM-7649 spheroids, and the combination produced an additive effect in PIM001-P spheroids
(Fig. 7A).
We then sought to test whether TIG could improve control of residual tumors persisting following 'NACT'
in vivo in our previously characterized PDX model of residual TNBC, PIM001-P, in which tumor regrowth
invariably follows initial tumor regressions induced by a variety of conventional chemotherapies[9,53]. Once
mammary tumors reached 150 mm^3, animals began treatment with CRB (50 mg/kg, i.p, administered on D0),
DTX (20 mg/kg, i.p. administered on D0), or TIG (100 mg/kg, i.p. q.d. for 10 days followed by a 3-day holiday).
Chemotherapies were administered for one cycle, and TIG was administered for three cycles. As a single agent,
TIG significantly slowed tumor growth (Fig. 7B-C). Treatment of residual PIM001-P tumors, defined as such when
they reached their volume nadir after a single dose of CRB or DTX, significantly delayed the regrowth of residual
tumor (Fig. 7B-C).
Using tumors collected after the second cycle of TIG (considered on-treatment), we affirmed target
engagement by western blotting for mtDNA-encoded ETC proteins (Fig. 7D). Immunohistochemistry revealed
that TIG moderately reduced Ki67 both as a monotherapy and when administered following chemotherapy.
Additionally, TIG increased cleaved caspase-3 staining in combination-treated tumors (Fig. 7E). Typically, during

a TIG cycle, the mice's body weight dropped ~5-10% but then recovered during the drug holiday (Fig. S13A).
Although TIG is safely used in routine clinical care, we found there are limitations in administering it to mice.
While gross pathology analysis did not reveal overt abnormalities in major organs, peritonitis of abdominal
tissues was consistently visually apparent. These studies provide proof of concept evidence that combining TIG
with chemotherapy is a potential therapeutic approach to further explore for chemo-refractory TNBC.
618
619
DISCUSSION
In humans, the mitoribosome translates the 13 mtDNA-encoded transcripts in the mitochondrial matrix,
all of which are components of ETC Complexes I, III, IV, and V. The remaining ETC components are encoded in
nDNA. OXA1L has been studied most thoroughly in yeast, and its function in mammalian cells has been recently
established. Its two reported roles are: 1) orchestrating translation termination of the 13 mtDNA-encoded ETC
proteins[25,26], and 2) co-translational insertion of mtDNA-encoded, as well as nDNA-encoded, ETC proteins into
the IMM[24-27]. Additionally, OXA1L has been linked to mitochondrial diseases[27,29,32], asthma[31,54], osteoporosis[55,56],
pain[57], diabetes complications[58], liver disease[59], Parkinson's disease[33], and cancer[60-63]. Despite these
associations, there are limited functional studies of OXA1L in any disease, and no functional studies of OXA1L
in the context of cancer have been reported. Our study provides evidence that OXA1L is crucial for mitochondrial
translation, ETC formation and function, oxphos, and chemoresistance in TNBC. Its appearance across multiple
external TNBC datasets suggests it's potentially generalizable role in human TNBC.
OXA1L KD disrupted mitochondrial protein translation, ETC formation, and mitochondrial function. Those
phenotypes became apparent five days, but not two days, after KD, likely because the existing pool of ETC
proteins had yet to be turned over to uncover the effect[64]. Furthermore, we observed a decrease in both mtDNA-
encoded and nDNA-encoded ETC proteins following KD, supporting the notion that OXA1L has dual functions
in mitochondrial translation and the insertion of ETC proteins into the IMM. ETC complexes can stand alone or
be in SCs with other complexes. In addition to other functions, SC formation has been shown to enhance electron
flow efficiency and stabilize the individual ETC proteins comprising the complexes[44,45,65]. We observed a
significant decrease in SC formation with *OXA1L* KD, which could explain the decrease in nDNA-encoded ETC
protein levels.
It is important to note that we did not observe changes in mtDNA-encoded proteins or OCR in the
immortalized, non-transformed mammary epithelial MCF10a cells with *OXA1L* KD. We previously reported that
MCF10a has a higher mitochondrial copy number[10], which could explain why these cells were less affected by
the *OXA1L* perturbation than the TNBC cell lines. Further, chemotherapy treatments did not elicit mitochondrial
rewiring in MCF10a cells[10], suggesting nontumor cells may be more 'resistant' to mitochondrial perturbations
than are tumor cells. This finding supports the notion that there may be a therapeutic window for translation of
mitochondria-targeting therapies to the clinic.
OXA1L was required for the CRB-induced OCR boost in residual cells and increased the sensitivity of
TNBC cells to both CRB and DTX *in vitro*. In our previous studies, we demonstrated that reducing mitochondrial
function, induced by chemotherapy, made cells more sensitive to DNA-damaging agents[9,10] so the increase in
sensitivity to CRB upon *OXA1L* KD was somewhat expected. We speculate that the increased sensitivity to DTX
upon *OXA1L* KD could be due to the cell's limited "reservoir" of oxphos capacity, with KD diminishing it below a
threshold required for cells to respond to the stress of chemotherapy. Another possibility is that DTX treatment
induced greater ROS[10], and when combined with OXA1L reduction, previously shown to induce ROS[32], ROS
accumulated to a lethal level. Future molecular studies will shed important light on these potentially divergent
mechanisms.
Since there are currently no inhibitors of OXA1L, we focused on targeting mitochondrial translation using
FDA-approved antibiotics, given the mitochondria's bacterial ancestry. Treating TNBC cells at an IC_{50} dose of
TIG disrupted mitochondrial protein translation, ETC formation, and mitochondrial function. In both MDA-MB-
231 and SUM159pt cells, we observed an additive effect of combining TIG with chemotherapies, and we were
able to recapitulate this finding with an additional antibiotic, chloramphenicol. Importantly, this additive effect was
specific to the cancer cell lines tested, as it was not observed in MCF10a cells.
Administration of TIG to CRB- or DTX-refractory residual tumors significantly delayed the regrowth of
orthotopic PIM001-P PDX tumors. A limitation in the *in vivo* evaluation of TIG was its adverse effects on the
abdominal cavity in mice, which had not been reported in previous mouse studies in the literature[48-51,66-71]. The
dosage of TIG we administered to mice was 100 mg/kg, corresponding to a human equivalent dose[72] of 8.13
mg/kg, that is about 12-fold less than human doses typically used to treat bacterial infections and about 37-fold

lower than the clinical MTD determined in an AML cancer phase I clinical trial of TIG monotherapy
(NCT01332786)[73]. Thus, clinically relevant doses of TIG were not achievable in the strain of mice we used with
intraperitoneal administration. Additional animal strains and administration routes would be worth testing for
further investigating the translational potential of TIG in TNBC. Our findings provide a foundation for future
expanded studies to identify putative biomarkers of TIG sensitivity for clinical translation. Further, PDX mice lack
a complete immune system, so future studies in immune competent TNBC models should be prioritized. It will
also be important to consider the effects of antibiotics on the gut and breast microbiomes in the context of human
TNBC, as microbiomes can influence tumor behaviors[74,75].

FIGURE LEGENDS

**Fig 1. Mitochondrial translation-related pathways were enriched in chemoresistant tumors at baseline
and elevated on-treatment with chemotherapy.** (A) Mitochondrial-related pathways that were significantly
enriched (FDR < 0.05) in non-pCR vs pCR CPTAC TNBC patient biopsies at baseline and on-treatment are
shown. (B) GSEA Reactome pathway analysis of TNBC PDX models[18], with only significantly altered pathways
displayed. Pathways including the term 'mitochondria' are shown in red. (C) Leading edge proteins from the top
pathway, 'Reactome mitochondrial protein translation termination,' upregulated in chemoresistant TNBC PDX
models are shown[18]. Plotted are the $-\log_{10}$ p-value from Spearman correlations. The proteins above the dotted
line are significantly enriched with p<0.05. (D) OXA1L protein abundance in TNBC patient[19] and PDX[18] cohorts
comparing responsive and non-responsive tumors. P-values were derived from one-tailed t-tests and from limma
moderated t-test with a paired design, respectively. Cartoon images were adapted from BioRender.

**Fig 2. The mitoribosome accessory protein OXA1L is required for protein production, formation, and
function of the ETC.** MDA-MB-231 cells were transfected with siRNAs targeting *OXA1L* (si1, si2, si3) or a
negative control sequence (NC). (A) western blots of whole cell lysates probing for ETC proteins at day 2 (D2)
and day 5 (D5) post-transfection are shown. (B) Seahorse MitoStress Tests of cells at D2 and D5 post-
transfection. A representative biological replicate of the D5 timepoint is shown as the tombstone plot. All D2 and
D5 biological replicates are represented in the energy plot. All biological replicates from D5 are shown in bar
plots. OCR and ECAR levels were normalized to cell counts. (C) Relative ATP at D5 post-transfection. (D) Native
Coomassie stain to assess SC formation with SDS-PAGE western blotting of mitochondrial and cytoplasmic
fractions. (E) Blue native-PAGE immunoblotting for complex III: UQCRC2 and measurement of SC bands
compared to total UQCRC2 quantified by densitometry. (F) Blue native-PAGE immunoblotting for complex V:
ATP5A and measurement of monomer and dimer bands compared to total ATP5A quantified by densitometry.
Blue native-PAGE immunoblotting for complex III using UQCRC2 and quantification of SC bands compared to
total UQCRC2 with densitometry. Data are shown as mean ±SEM, and individual data points represent biological
replicates. Significant comparisons are indicated; *p < 0.05, **p < 0.01, ***p < 0.001, ****p<0.0001 by one-way
ANOVA.

**Fig 3. OXA1L KD diminished chemotherapy-induced metabolic adaptation and sensitized TNBC cells to
chemotherapy.** MDA-MB-231 cells were transfected with siRNAs targeting *OXA1L* (si1, si2, si3) or the negative
control (NC) followed by treatment with chemotherapy. Chemotherapy treatments were vehicle (VEH), CRB_{hi}
(100 µM), CRB_{low} (25 µM), DTX (5nM), and DTX+CRB_{low}. Readouts were conducted at D5 post-transfection and
D2 post-chemotherapy treatment. (A) Seahorse MitoStress Tests of cells at D5 post-transfection. A
representative biological replicate is shown in the tombstone plot, while all biological replicates are represented
in the energy plot and bar graphs. OCR and ECAR values were normalized to viable cell counts. (B) Western
blotting of whole cell lysates probing for ETC proteins. (C) IC_{50} values were measured in eight-point dose curves
and relative values are displayed. Data are shown as mean ±SEM, and individual data points represent biological
replicates. Significant comparisons are indicated; *p < 0.05, **p < 0.01, ***p < 0.001, ****p<0.0001 by one-way
ANOVA.

**Fig 4. The bacterial ancestry of mitochondria allows for repurposing of FDA-approved antibiotics to
inhibit mitochondrial translation.** MDA-MB-231 cells were treated with vehicle (VEH) or tigecycline (TIG 8 µM)
for two days. (A) IC_{50} dilution curve, (B) western blotting of whole cell lysates probing for ETC proteins, (C)
mtDNA:nDNA ratio, and (D) qRT-PCR analysis of mitochondrial gene expression. Expression of mtDNA-encoded
genes was normalized to *TUBB*. (E) Coomassie staining of a native gel to assess SC formation, and SDS-PAGE

western blot of mitochondrial and cytoplasm fractions. (F) Blue native-PAGE immunoblotting for complex III: UQCRC2 and measurement of SC bands compared to total UQCRC2 by densitometry. (G) Seahorse MitoStress Tests showing a representative biological replicate for the tombstone plot and all biological replicates in the bar graphs. OCR and ECAR values were normalized to viable cell counts. Data are shown as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; *p < 0.05, **p < 0.01, ***p < 0.001, ****p<0.0001 by unpaired t-test.

Fig 5. TIG diminishes chemotherapy-induced mitochondrial adaptations. MDA-MB-231 cells were treated with VEH, TIG (8 µM), CRB (100 µM), and DTX (5 nM) for two days. (A) Representative images from Mito-FUNCAT are shown with nuclei masked (scale bar = 10 µmicron). The total colocalization of positive pixels for HPG-Azide and COXIV after nuclear masking per cell is shown in the first plot. In the second plot, the proportion of mitochondria actively translating is shown. Each point represents the quantification of an image. 25-30 cells were imaged per condition. Three independent biological replicates were conducted. The solid lines within the truncated violin plot are at the quartiles, and the dotted line is at the median. Normality and lognormality statistical tests revealed data were non-normal, so Mann-Whitney U Tests were performed between treatment pairs of interest. Significant comparisons are indicated; *p < 0.05, **p < 0.01, ***p < 0.001, ****p<0.0001). (B) Western blotting of whole cell lysates probing for ETC proteins. (C&D) Seahorse MitoStress Tests showing representative tombstone plots from one biological replicate. The energy plot includes all the biological replicates conducted. Data are shown as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; *p < 0.05, **p < 0.01, ***p < 0.001, ****p<0.0001 by one-way ANOVA.

Fig 6. TIG increases the sensitivity of TNBC cells to chemotherapy. (A) Incucyte analysis of MDA-MB-231 cells treated with dilutions of TIG alone or simultaneously with CRB_{hi} (100µM). The confluency of the cells readout at 240hrs (D10) was used for the one-way ANOVA followed by the post-hoc Dunnett's test. (B) Incucyte analysis of SUM159pt cells treated with dilutions of TIG alone or simultaneously with CRB_{hi} (100µM). The confluency of the cells readout at 240hrs (D10) was used for the one-way ANOVA followed by the post-hoc Dunnett's test. (C) Hoechst measurement of relative cell counts on D7 of MDA-MB-231 cells after treatment with 2.5 nM of DTX followed by of TIG. (D) Hoechst measurement of relative cell counts on D7 of SUM159pt cells after treatment with DTX (5 nM) followed by TIG. Data are shown as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; *p < 0.05, **p < 0.01, ***p < 0.001, ****p<0.0001 by one-way ANOVA.

Fig 7. TIG delays the regrowth of an orthotopic TNBC PDX models at the residual state. (A) Three PDX models (PIM001-P, BCM-4272, and BCM-7649) were used to generate spheroid cultures for *ex-vivo* treatments with TIG and CRB. (B) The PIM001-P TNBC PDX model was orthotopically implanted into female SCID/bg mice to assess TIG given as a monotherapy or in at the residual state following either DTX or CRB treatment. There were 8 mice per treatment group. Normalized tumor volumes are shown. Mice were treated upfront with one dose of CRB (dark green arrow), one dose of DTX (dark blue arrow), or a cycle of TIG (red brackets). In the CRB and DTX treatment arms, TIG treatment cycles (light green or blue brackets) began at the respective residual timepoint (the approximate tumor volume nadir) for each chemotherapy agent. Asterisks represent when mice on TIG treatment were collected. (C) Kaplan-Meier plot of days to reach 800 mm3 for each tumor volume. The log-rank p-values are displayed. (D) Western blotting of tumors collected on treatment probing for ETC proteins. (EEE) Slides stained with H&E or antibodies against Ki67 and CC3 (scale bar = 50 µm) are shown from PIM001-P tumors collected the last treatment day of the second cycle. Cartoon images adapted from BioRender.

SUPPLEMENTAL FIGURE LEGENDS

Fig S1. Mitoribosome protein signature tends to be enriched in non-pCR clinical and PDX samples. (A&B) Significant pathways enriched in the non-pCR vs pCR CPTAC TNBC patient biopsies at baseline from Gene Set Enrichment analysis with Reactome and MitoCarta geneset collections that was performed using WebGestalt[36]. Significance set at FDR < 0.05. (C) Significant pathways enriched in the on-treatment vs. baseline CPTAC TNBC patient biopsies from the Gene Set Enrichment analysis with MitoCarta geneset collections that was performed using WebGestalt[36]. Significance set at FDR < 0.05. (D) MRP protein signature comparing CPTAC-TNBC patient tumors[19] and TNBC PDXs[18] cohorts comparing responsive vs. non-responsive tumors. P-values derived from limma moderated t-test with a paired design and from t-test, respectively.

Fig S2. Mitoribosome accessory protein OXA1L is required for ETC protein levels and mitochondrial function in SUM159pt cells. SUM159pt cells were transfected with siRNAs targeting *OXA1L* (si1, si2, si3) or the negative control (NC). (A) Western blotting of whole cell lysates probing for ETC proteins at Day 2 (D2) and Day 5 (D5) post-transfection. (B) Seahorse MitoStress Tests of cells at D2 and D5 post-transfection. A representative biological replicate of D5 is shown as the tombstone plot. All D2 and D5 biological replicates are represented in the energy plot. All biological replicates from D5 are shown in the bar graphs. OCR and ECAR values were normalized to viable cell counts. (C) Relative ATP at D5 post-transfection. (D) Native Coomassie stain to assess SC formation with SDS-PAGE western blotting of mitochondrial and cytoplasmic fractions. (E) Blue native-PAGE immunoblotting for complex III: UQCRC2 and measurement of SC bands compared to total UQCRC2 quantified by densitometry. (F) Blue native-PAGE immunoblotting for complex V: ATP5A and measurement of monomer and dimer bands compared to total ATP5A quantified by densitometry. Data are shown as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p<0.0001$ by one-way ANOVA.

Fig S3. *OXA1L* KD does not affect TNBC cell confluency over time or mtDNA ratio, but does have some effect on mt-mRNA levels. MDA-MB-231 and SUM159pt cells were transfected with siRNA targeting *OXA1L* (si1, si2, si3) or the negative control (NC). (A) Incucyte analysis of cellular confluence, (B) mtDNA:nDNA ratio quantification, and (C) qRT-PCR analysis of mitochondrial RNA levels. Expression of mtDNA-encoded genes was normalized to *TUBB*. Data are shown as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p<0.0001$ by one-way ANOVA.

Fig S4. MCF10a OCR is not significantly affected by OXA1L KD. (A) *OXA1L* mRNA expression aligned with TNBC subtype, metabolic subtype (MPS), mRNA expression, and dependency score. The more negative the score is, the greater the gene dependency. (B) *OXA1L* dependency score plotted against *OXA1L* mRNA expression. P-values derived from Pearson Correlation. (C) TNBC subtype plotted against mRNA expression. (D) *OXA1L* dependency score plotted against TNBC subtype. (E) OXA1L dependency score plotted against MPS. (F) MCF10a cells were transfected with siRNAs targeting *OXA1L* (si1, si2, si3) or the negative control (NC). Incucyte analysis of cellular confluence with *OXA1L* KD. (G) Western blotting of MCF10a whole cell lysates on days D2 and D5 after transfection, probing for mtDNA-encoded proteins. (H) Seahorse MitoStress Tests of MCF10a cells D2 and D5 post-transfection. A representative biological replicate from D5 is shown in the tombstone plot. D2 and D5 timepoints from all the biological replicates are represented in the energy plot. Only D5 are shown in bar graphs. OCR and ECAR values were normalized to viable cell counts. Data are shown as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p<0.0001$ by one-way ANOVA.

Fig S5. OXA1L diminishes chemotherapy-induced metabolic adaptations and sensitizes SUM159pt cells to chemotherapy. SUM159pt cells were transfected with siRNAs targeting *OXA1L* (si1, si2, si3) or the negative control (NC), followed by treatment with chemotherapy. Chemotherapy treatments were CRB$_{hi}$ (100 µM), CRB$_{low}$ (25 µM), DTX (5nM), and DTX+CRB$_{low}$. Readouts were conducted at D5 post-transfection and D2 post-chemotherapy treatment. (A) Seahorse MitoStress Tests of cells at D5 post-transfection. A representative biological replicate is shown in the tombstone plot, while all biological replicates are represented in the energy plot and bar graphs. OCR and ECAR values were normalized to viable cell counts. (B) Western blotting of whole cell lysates, probing for ETC proteins. (C) IC$_{50}$ values were measured in eight-point dose curves, and relative values are displayed. Data are shown as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p<0.0001$ by one-way ANOVA.

Fig S6. *OXA1L* KD diminishes CRB-induced OCR boost. MDA-MB-231 and SUM159pt cells were transfected with siRNAs targeting *OXA1L* (si1, si2, si3) or the negative control (NC). Following transfection, cells were treated with chemotherapy. OCR and ECAR values were normalized to viable cell counts. (A) Data from Seahorse MitoStress Tests are shown as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p<0.0001$ by one-way ANOVA.

Fig S7. TIG targets mitochondrial translation in SUM159 cells. SUM159pt cells were treated with vehicle (VEH) or tigecycline (TIG 3 µM) for two days. (A) IC$_{50}$ dilution curve, (B) western blotting of whole cell lysates

probing for ETC proteins, (C) mtDNA:nDNA ratio, and (D) qRT-PCR analysis of mitochondrial gene expression.
Expression of mtDNA-encoded genes was normalized to *TUBB*. (E) Coomassie staining of a native gel to assess
SC formation, and SDS-PAGE western blot of mitochondrial and cytoplasm fractions. (F) Blue native-PAGE
immunoblotting for complex III: UQCRC2 and measurement of SC bands compared to total UQCRC2 by
densitometry. (G) Seahorse MitoStress Tests showing a representative biological replicate for the tombstone plot
and all biological replicates in the bar graphs. OCR and ECAR values were normalized to viable cell counts.
Data are shown as mean ±SEM from a minimum of three independent biological replicates. Significant
comparisons are indicated; $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p<0.0001$ by unpaired t-test.
844
Fig S8. TNBC cells respond to TIG in a dose-dependent manner. MDA-MB-231 cells were treated with TIG
at a high dose (50 µM) and the IC_{50} dose (8 µM). (A) Relative viable cells from AOPI counts on D2 after treatment.
(B) Colony counts on D7 after seeding treated cells at 100 cells per well. SUM159 cells treated with TIG at a
high dose (50 µM) and the IC_{50} dose (3 µM). (C) Relative viable cells from AO/PI counting on D2 after treatment.
(D) Colony counts on D7 after seeding treated cells at 100 cells per well. (E) Incucyte analysis of MDA-MB-231
and SUM159pt cells treated with TIG dilutions. The confluency of the cells readout at the end of the experiment
(120hrs) was used for the one-way ANOVA followed by the post-hoc Dunnett's test. Data are shown as mean
±SEM, and individual data points represent biological replicates. Significant comparisons are indicated; $*p <$
0.05, $**p < 0.01$, $***p < 0.001$, $****p<0.0001$ by one-way ANOVA.
854
Fig S9. TIG diminishes OCR in MDA-MB-231 cells treated with chemotherapy. (A) Mito-FUNCAT analysis
of control samples of MDA-MB-231 cells. Cells were given HPG, a methionine analog, to be incorporated into
nascent polypeptides, which were then fluorescently detected via click chemistry. Additionally, cells were treated
with inhibitors to block either cytosolic translation (cycloheximide, CHX) or mitochondrial translation (TIG, high
dose, 50 µM). Representative images from each condition are shown (scale bar = 10 µm), and the quantification
of HPG-Azide and COXIV total colocalization of positive pixels for HPG-Azide and COXIV per cell after nuclear
masking is plotted in the first plot. In the second plot, the proportion of mitochondria actively translating is shown.
Each dot represents quantification of an image. 25-30 cells were imaged per condition. Three independent
biological replicates were conducted. The solid lines within the truncated violin plot are at the quartiles, and the
dotted line is at the median. (B) Seahorse MitoStress Tests of MDA-MB-231 cells treated with TIG (8 µM) as a
monotherapy or in combination with CRB_{hi} (100 µM) or CRB_{low} (25 µM). (C) Seahorse MitoStress Tests of MDA-
MB-231 cells treated with TIG (8 µM) as a monotherapy or with docetaxel (DTX 5 nM). Data are shown as mean
±SEM from a minimum of three independent biological replicates. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$,
$****p<0.0001$ by one-way ANOVA.
869
Fig S10. TIG diminishes chemotherapy-induced metabolic adaptations in SUM159 cells. SUM159pt cells
were treated with VEH, TIG (3 µM), CRB (100 µM), or DTX (5 nM) for two days. (A) Western blotting of whole
cell lysates probing for ETC proteins. (B&C) Seahorse MitoStress Tests showing representative tombstone plots
from one biological replicate. The energy plot includes all the biological replicates conducted. Data are shown
as mean ±SEM, and individual data points represent biological replicates. Significant comparisons are indicated;
$*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p<0.0001$ by one-way ANOVA.
876
Fig S11. Chloramphenicol sensitizes MDA-MB-231 cells to CRB treatment. MDA-MB-231 cells were treated
with chloramphenicol. (A) Incucyte analysis of cells treated at different dilutions of chloramphenicol alone or with
CRB_{hi} (100 µM). The confluency of the cells readout at the end of the experiment (164hrs) was used for the one-
way ANOVA followed by the post-hoc Dunnett's test. (B) Seahorse MitoStress Tests showing representative
tombstone plots from one biological replicate. The energy plot includes all the biological replicates conducted.
Data are shown as mean ±SEM, and individual data points represent biological replicates. Significant
comparisons are indicated; $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p<0.0001$ by one-way ANOVA.
884
Fig S12. TIG does not affect mitochondrial function or chemotherapy sensitivity in MCF10a cells. MCF10a
cells were treated with VEH, TIG (8 µM), or CRB (10 µM) for two days unless noted otherwise. (A) Western
blotting of whole cell lysates probing for ETC proteins. (B) Seahorse MitoStress Tests showing representative
tombstone plots from one biological replicate. The energy plot includes all the biological replicates conducted.
(C) Incucyte analysis of cells treated with different dilutions of TIG alone or with CRB_{hi} (100 µM). The confluency
of the cells readout at the end of the experiment (312hrs) was used for the one-way ANOVA followed by the post-
hoc Dunnett's test. Data are shown as mean ±SEM, and individual data points represent biological replicates.

Significant comparisons are indicated; *p < 0.05, **p < 0.01, ***p < 0.001, ****p<0.0001 by one-way ANOVA.

Fig S13. TIG does not significantly impact body weight in mice. (A) Female Scid/bg mice bearing orthotopic PIM001-P tumors were treated with TIG and CRB. Normalized body weights to the start of treatment are plotted. Mice were treated upfront with one dose of CRB (dark green arrow), one dose of DTX (dark blue arrow), or a cycle of TIG (red brackets). In the CRB and DTX treatment arms, TIG treatment cycles (light green or blue brackets) began at the respective residual timepoint for each chemotherapy agent. Asterisks represent when mice on TIG treatment were collected. Cartoon images adapted from BioRender.

ACKNOWLEDGEMENTS

We are grateful to the breast cancer patients who donated their biopsies for cell line and PDX model generation. We are grateful to Janice Cowden and Amy Beumer for providing patient advocacy support for this research. Drs. Eric Jaehnig, PhD and Meenakshi Anurag, PhD aided in on-treatment CPTAC patient proteomic signature analysis. Dr. Junegoo Lee and Ms. Basmala Essam assisted with mouse experiments. Dr. Bora Lim provided translational guidance. Dr. Helen Piwnica-Worms provided the PIM001-P PDX through a materials transfer agreement with the University of Texas MD Anderson Cancer Center, and model generation was supported by a generous gift from the Cazalot Foundation through the MD Anderson Women's Cancer Moonshot Program.

BCM core facilities that supported this work included: The Optical Imaging & Vital Microscopy Core for Mito-FUNCAT imaging; The Mouse Metabolism and Phenotyping Core (NIH fund P30DK144025 & UM1HG006348) for Seahorse Bioanalyzer analysis; Patient-derived Xenograft Core (P30 Cancer Center Support Grant NCI CA125123, CPRIT Core Facilities Support Grant RP220646); Breast Center Path Core (supported by the Breast Center and a variety of research grants awarded to its faculty, including one of nine Specialized Programs of Research Excellence (SPORE) in Breast Cancer granted by the National Institute of Health. S10 grant (1S10OD028671-01) for digital imaging of the IHC slides) for processing tumor tissue into FFPE blocks and IHC staining. STR DNA fingerprinting was conducted by the Cytogenetics and Cell Authentication Core at M.D. Anderson Cancer Center.

AUTHOR CONTRIBUTIONS

MJB and GVE were responsible for study conceptualization, carrying out the study, data presentation, and manuscript writing.

SWW conducted ETC complex analyses and provided experimental guidance under the supervision of GVE.

MLB conducted qRT-PCR and provided experimental guidance under the supervision of GVE.

AL conducted mammary epithelial cell experiments and assisted with computational analyses under the supervision of GVE.

ASG conducted mouse experiments under the supervision of GVE.

KW conducted qPCR assays under the supervision of GVE.

LED aided in PDX experiments under the supervision of MTL.

IS conducted fluorescence colocalization quantitative analyses under the supervision of QZ.

QZ oversaw fluorescence colocalization quantitative analyses.

BZ aided in oversight of proteogenomic analyses.

JTL conducted computational analyses of transcriptomic and proteomic datasets under the supervision of BZ and MTL.

948
949 MTL aided in oversight of PDX studies and proteogenomic analyses.
950
951

FUNDING
G.V.E. is a Cancer Prevention Research Institute of Texas (CPRIT) Scholar in Cancer Research. The
authors were supported by CPRIT RR200009 to G.V.E.; National Institutes of Health (NIH) K22-CA241113 to
GVE, R37CA269783-01A1 to G.V.E., and T32 predoctoral training grant T32GM136560-02 to MJB; an American
Cancer Society Research Scholar Grant RSG-22-093-01-CCB to GVE; a Mary Kay Ash Foundation Cancer
Research Grant 02-24 to GVE; a Breast Cancer Alliance Young Investigator Award to GVE; a National Science
Foundation (NSF) Graduate Research Fellowship 2140736 to MJB; a Myra Branum Wilson Baylor Research
Advocates for Student Scientists (BRASS) Scholarship to MJB; an American Cancer Society Postdoctoral
Fellowship PF-24-1293970-01-TBI to SW; a Frank & Sandra Kimmel Endowment Postdoc Fellowship to MLB.
The content is solely the responsibility of the authors and does not necessarily represent the official views of the
NIH, NSF, CPRIT, ACS, or BRASS.

963
964

965 **REFERENCES CITED**
966
1　　Harbeck, N. *et al.* Breast cancer. *Nat Rev Dis Primers* **5**, 66 (2019). https://doi.org:10.1038/s41572-019-
0111-2
2　　Pavese, F. *et al.* BRCA Mutation Status in Triple-Negative Breast Cancer Patients Treated with
Neoadjuvant Chemotherapy: A Pivotal Role for Treatment Decision-Making. *Cancers (Basel)* **14** (2022).
https://doi.org:10.3390/cancers14194571
3　　Bianchini, G., De Angelis, C., Licata, L. & Gianni, L. Treatment landscape of triple-negative breast
cancer - expanded options, evolving needs. *Nat Rev Clin Oncol* **19**, 91-113 (2022).
https://doi.org:10.1038/s41571-021-00565-2
4　　Vagia, E., Mahalingam, D. & Cristofanilli, M. The Landscape of Targeted Therapies in TNBC. *Cancers*
(Basel) **12** (2020). https://doi.org:10.3390/cancers12040916
5　　Schmid, P. *et al.* Pembrolizumab for Early Triple-Negative Breast Cancer. *N Engl J Med* **382**, 810-821
(2020). https://doi.org:10.1056/NEJMoa1910549
6　　Symmans, W. F. *et al.* Long-Term Prognostic Risk After Neoadjuvant Chemotherapy Associated With
Residual Cancer Burden and Breast Cancer Subtype. *J Clin Oncol* **35**, 1049-1060 (2017).
https://doi.org:10.1200/JCO.2015.63.1010
7　　Yau, C. *et al.* Residual cancer burden after neoadjuvant chemotherapy and long-term survival
outcomes in breast cancer: a multicentre pooled analysis of 5161 patients. *Lancet Oncol* **23**, 149-160
(2022). https://doi.org:10.1016/S1470-2045(21)00589-1
8　　*SEER*Explorer: An interactive website for SEER cancer statistics*, <https://seer.cancer.gov/statistics-
network/explorer/application.html?site=622&data_type=1&graph_type=2&compareBy=race&chk_race_
6=6&chk_race_5=5&chk_race_4=4&chk_race_9=9&chk_race_8=8&rate_type=2&hdn_sex=3&age_ran
ge=1&stage=101&advopt_precision=1&advopt_show_ci=on#resultsRegion0> (2026).
9　　Echeverria, G. V. *et al.* Resistance to neoadjuvant chemotherapy in triple-negative breast cancer
mediated by a reversible drug-tolerant state. *Sci Transl Med* **11** (2019).
https://doi.org:10.1126/scitranslmed.aav0936
10　　Baek, M. L. *et al.* Mitochondrial structure and function adaptation in residual triple negative breast
cancer cells surviving chemotherapy treatment. *Oncogene* **42**, 1117-1131 (2023).
https://doi.org:10.1038/s41388-023-02596-8
11　　Vasan, K., Werner, M. & Chandel, N. S. Mitochondrial Metabolism as a Target for Cancer Therapy. *Cell*
Metab **32**, 341-352 (2020). https://doi.org:10.1016/j.cmet.2020.06.019
12　　Decker, C. W. *et al.* Mitofusin-2 mediates doxorubicin sensitivity and acute resistance in Jurkat
leukemia cells. *Biochem Biophys Rep* **24**, 100824 (2020). https://doi.org:10.1016/j.bbrep.2020.100824
13　　Vellinga, T. T. *et al.* SIRT1/PGC1alpha-Dependent Increase in Oxidative Phosphorylation Supports
Chemotherapy Resistance of Colon Cancer. *Clin Cancer Res* **21**, 2870-2879 (2015).
https://doi.org:10.1158/1078-0432.CCR-14-2290

Avolio, R., Matassa, D. S., Criscuolo, D., Landriscina, M. & Esposito, F. Modulation of Mitochondrial Metabolic Reprogramming and Oxidative Stress to Overcome Chemoresistance in Cancer. *Biomolecules* **10** (2020). https://doi.org:10.3390/biom10010135

Pendleton, K. E., Wang, K. & Echeverria, G. V. Rewiring of mitochondrial metabolism in therapy-resistant cancers: permanent and plastic adaptations. *Front Cell Dev Biol* **11**, 1254313 (2023). https://doi.org:10.3389/fcell.2023.1254313

Lee, K. M. *et al.* MYC and MCL1 Cooperatively Promote Chemotherapy-Resistant Breast Cancer Stem Cells via Regulation of Mitochondrial Oxidative Phosphorylation. *Cell Metab* **26**, 633-647 e637 (2017). https://doi.org:10.1016/j.cmet.2017.09.009

Evans, K. W. *et al.* Oxidative Phosphorylation Is a Metabolic Vulnerability in Chemotherapy-Resistant Triple-Negative Breast Cancer. *Cancer Res* **81**, 5572-5581 (2021). https://doi.org:10.1158/0008-5472.CAN-20-3242

Lei, J. T. *et al.* Patient-Derived Xenografts of Triple-Negative Breast Cancer Enable Deconvolution and Prediction of Chemotherapy Responses. *bioRxiv* (2025). https://doi.org:10.1101/2024.12.09.627518

Anurag, M. *et al.* Proteogenomic Markers of Chemotherapy Resistance and Response in Triple-Negative Breast Cancer. *Cancer Discov* **12**, 2586-2605 (2022). https://doi.org:10.1158/2159-8290.CD-22-0200

Gong, Y. *et al.* Metabolic-Pathway-Based Subtyping of Triple-Negative Breast Cancer Reveals Potential Therapeutic Targets. *Cell Metab* **33**, 51-64 e59 (2021). https://doi.org:10.1016/j.cmet.2020.10.012

Molina, J. R. *et al.* An inhibitor of oxidative phosphorylation exploits cancer vulnerability. *Nat Med* **24**, 1036-1046 (2018). https://doi.org:10.1038/s41591-018-0052-4

Herkenne, S. *et al.* Developmental and Tumor Angiogenesis Requires the Mitochondria-Shaping Protein Opa1. *Cell Metab* **31**, 987-1003 e1008 (2020). https://doi.org:10.1016/j.cmet.2020.04.007

Chinnery, P. F. & Hudson, G. Mitochondrial genetics. *Br Med Bull* **106**, 135-159 (2013). https://doi.org:10.1093/bmb/ldt017

Christian, B. E. & Spremulli, L. L. Mechanism of protein biosynthesis in mammalian mitochondria. *Biochim Biophys Acta* **1819**, 1035-1054 (2012). https://doi.org:10.1016/j.bbagrm.2011.11.009

Itoh, Y. *et al.* Mechanism of membrane-tethered mitochondrial protein synthesis. *Science* **371**, 846-849 (2021). https://doi.org:10.1126/science.abe0763

Haque, M. E. *et al.* Properties of the C-terminal tail of human mitochondrial inner membrane protein Oxa1L and its interactions with mammalian mitochondrial ribosomes. *J Biol Chem* **285**, 28353-28362 (2010). https://doi.org:10.1074/jbc.M110.148262

Thompson, K. *et al.* OXA1L mutations cause mitochondrial encephalopathy and a combined oxidative phosphorylation defect. *EMBO Mol Med* **10** (2018). https://doi.org:10.15252/emmm.201809060

Stiburek, L. *et al.* Knockdown of human Oxa1l impairs the biogenesis of F1Fo-ATP synthase and NADH:ubiquinone oxidoreductase. *J Mol Biol* **374**, 506-516 (2007). https://doi.org:10.1016/j.jmb.2007.09.044

Coenen, M. J., Smeitink, J. A., Smeets, R., Trijbels, F. J. & van den Heuvel, L. P. Mutation detection in four candidate genes (OXA1L, MRS2L, YME1L and MIPEP) for combined deficiencies in the oxidative phosphorylation system. *J Inherit Metab Dis* **28**, 1091-1097 (2005). https://doi.org:10.1007/s10545-005-4483-y

Pires, A. O. *et al.* Polymorphisms in the DAD1 and OXA1L genes are associated with asthma and atopy in a South American population. *Mol Immunol* **101**, 294-302 (2018). https://doi.org:10.1016/j.molimm.2018.07.014

Pires, A. O. *et al.* New variants of the DAD1 and OXA1L genes are associated with asthma and atopy in an adult population. *Gene* **937**, 149124 (2025). https://doi.org:10.1016/j.gene.2024.149124

Zhan, Y. *et al.* OXA1L deficiency causes mitochondrial myopathy via reactive oxygen species regulated nuclear factor kappa B signalling pathway. *Clin Transl Med* **15**, e70385 (2025). https://doi.org:10.1002/ctm2.70385

Xurui Shen, R. H., Huiwei Zhong, Peixin Sun, Jinlun Kylian Zhang, Yu Zhang, Hanting Yang. *Oxa1L buffers mitochondrial vulnerability by coupling translation to membrane insertion*, 2026).

Petrosyan, V. *et al.* Identifying biomarkers of differential chemotherapy response in TNBC patient-derived xenografts with a CTD/WGCNA approach. *iScience* **26**, 105799 (2023). https://doi.org:10.1016/j.isci.2022.105799

Lei, J. T. *et al.* Patient-Derived Xenografts of Triple-Negative Breast Cancer Enable Deconvolution and Prediction of Chemotherapy Responses. *bioRxiv* (2024). https://doi.org:10.1101/2024.12.09.627518

058 36 Elizarraras, J. M. *et al.* WebGestalt 2024: faster gene set analysis and new support for metabolomics
059  and multi-omics. *Nucleic Acids Res* **52**, W415-W421 (2024). https://doi.org:10.1093/nar/gkae456

060 37 Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing and microarray
061  studies. *Nucleic Acids Res* **43**, e47 (2015). https://doi.org:10.1093/nar/gkv007

062 38 Lehmann, B. D. *et al.* Identification of human triple-negative breast cancer subtypes and preclinical
063  models for selection of targeted therapies. *J Clin Invest* **121**, 2750-2767 (2011).
064  https://doi.org:10.1172/JCI45014

065 39 Barbie, D. A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven cancers require
066  TBK1. *Nature* **462**, 108-112 (2009). https://doi.org:10.1038/nature08460

067 40 Clayton, D. A. & Shadel, G. S. Isolation of mitochondria from cells and tissues. *Cold Spring Harb Protoc*
068  **2014**, pdb top074542 (2014). https://doi.org:10.1101/pdb.top074542

069 41 Yousefi, R. *et al.* Monitoring mitochondrial translation in living cells. *EMBO Rep* **22**, e51635 (2021).
070  https://doi.org:10.15252/embr.202051635

071 42 Malik, N. *et al.* Dysregulation of Mitochondrial Translation Caused by CBFB Deficiency Cooperates with
072  Mutant PIK3CA and Is a Vulnerability in Breast Cancer. *Cancer Res* **83**, 1280-1298 (2023).
073  https://doi.org:10.1158/0008-5472.CAN-22-2525

074 43 Echeverria, G. V. *et al.* High-resolution clonal mapping of multi-organ metastasis in triple negative
075  breast cancer. *Nat Commun* **9**, 5079 (2018). https://doi.org:10.1038/s41467-018-07406-4

076 44 Javadov, S., Jang, S., Chapa-Dubocq, X. R., Khuchua, Z. & Camara, A. K. Mitochondrial respiratory
077  supercomplexes in mammalian cells: structural versus functional role. *J Mol Med (Berl)* **99**, 57-73
078  (2021). https://doi.org:10.1007/s00109-020-02004-8

079 45 Berndtsson, J. *et al.* Respiratory supercomplexes enhance electron transport by decreasing
080  cytochrome c diffusion distance. *EMBO Rep* **21**, e51015 (2020).
081  https://doi.org:10.15252/embr.202051015

082 46 McFarland, J. M. *et al.* Improved estimation of cancer dependencies from large-scale RNAi screens
083  using model-based normalization and data integration. *Nat Commun* **9**, 4610 (2018).
084  https://doi.org:10.1038/s41467-018-06916-5

085 47 Lamb, R. *et al.* Antibiotics that target mitochondria effectively eradicate cancer stem cells, across
086  multiple tumor types: treating cancer like an infectious disease. *Oncotarget* **6**, 4569-4584 (2015).
087  https://doi.org:10.18632/oncotarget.3174

088 48 Fu, X. *et al.* Targeting mitochondrial respiration selectively sensitizes pediatric acute lymphoblastic
089  leukemia cell lines and patient samples to standard chemotherapy. *Am J Cancer Res* **7**, 2395-2405
090  (2017).

091 49 Tan, J., Song, M., Zhou, M. & Hu, Y. Antibiotic tigecycline enhances cisplatin activity against human
092  hepatocellular carcinoma through inducing mitochondrial dysfunction and oxidative damage. *Biochem*
093  *Biophys Res Commun* **483**, 17-23 (2017). https://doi.org:10.1016/j.bbrc.2017.01.021

094 50 Hu, B. & Guo, Y. Inhibition of mitochondrial translation as a therapeutic strategy for human ovarian
095  cancer to overcome chemoresistance. *Biochem Biophys Res Commun* **509**, 373-378 (2019).
096  https://doi.org:10.1016/j.bbrc.2018.12.127

097 51 Jones, R. A. *et al.* RB1 deficiency in triple-negative breast cancer induces mitochondrial protein
098  translation. *J Clin Invest* **126**, 3739-3757 (2016). https://doi.org:10.1172/JCI81568

099 52 Zhang, X. & Lewis, M. T. Establishment of Patient-Derived Xenograft (PDX) Models of Human Breast
100  Cancer. *Curr Protoc Mouse Biol* **3**, 21-29 (2013). https://doi.org:10.1002/9780470942390.mo120140

101 53 Berner, M. J. *et al.* Three-dimensional analysis of mitochondria in a patient-derived xenograft model of
102  triple negative breast cancer reveals mitochondrial network remodeling following chemotherapy
103  treatments. *bioRxiv* (2024). https://doi.org:10.1101/2024.09.09.611245

104 54 Costa, G. N. *et al.* A genome-wide association study of asthma symptoms in Latin American children.
105  *BMC Genet* **16**, 141 (2015). https://doi.org:10.1186/s12863-015-0296-7

106 55 Zhang, L., Yang, Y., Geng, D. & Wu, Y. Identification of Potential Therapeutic Targets and Molecular
107  Regulatory Mechanisms for Osteoporosis by Bioinformatics Methods. *Biomed Res Int* **2021**, 8851421
108  (2021). https://doi.org:10.1155/2021/8851421

109 56 Vrscaj, L. A., Marc, J. & Ostanek, B. Differential expression of lncRNAs and mRNAs in bone marrow-
110  derived mesenchymal stem cells under continuous and intermittent teriparatide treatment. *Biomed*
111  *Pharmacother* **189**, 118293 (2025). https://doi.org:10.1016/j.biopha.2025.118293

112 57 Wotton, J. M. *et al.* Identifying genetic determinants of inflammatory pain in mice using a large-scale
113  gene-targeted screen. *Pain* **163**, 1139-1157 (2022). https://doi.org:10.1097/j.pain.0000000000002481

58   Xu, W., Liang, Y., Zhuang, Y. & Yuan, Z. Identification of miRNA-mRNA Regulatory Networks
Associated with Diabetic Retinopathy using Bioinformatics Analysis. *Endocr Metab Immune Disord*
Drug Targets **23**, 1628-1636 (2023). https://doi.org:10.2174/1871530323666230419081351
59   Meroni, M. *et al.* A novel gene signature to diagnose MASLD in metabolically unhealthy obese
individuals. *Biochem Pharmacol* **218**, 115925 (2023). https://doi.org:10.1016/j.bcp.2023.115925
60   Helbig, D., Mauch, C., Buettner, R. & Quaas, A. Immunohistochemical expression of melanocytic and
myofibroblastic markers and their molecular correlation in atypical fibroxanthomas and pleomorphic
dermal sarcomas. *J Cutan Pathol* **45**, 880-885 (2018). https://doi.org:10.1111/cup.13346
61   Zheng, X. *et al.* An original cuproptosis-related genes signature effectively influences the prognosis and
immune status of head and neck squamous cell carcinoma. *Front Genet* **13**, 1084206 (2022).
https://doi.org:10.3389/fgene.2022.1084206
62   Wu, F., Ning, H., Sun, Y., Wu, H. & Lyu, J. Integrative exploration of the mutual gene signatures and
immune microenvironment between benign prostate hyperplasia and castration-resistant prostate
cancer. *Aging Male* **26**, 2183947 (2023). https://doi.org:10.1080/13685538.2023.2183947
63   Goze, O. F., Buettner, R., Odenthal, M. & Heydt, C. Genetic diversity in melanomagenesis: a
comprehensive analysis of BRAF, NRAS, KIT and novel mutations across melanoma subtypes. *BMC*
Cancer **26**, 205 (2025). https://doi.org:10.1186/s12885-025-15458-1
64   Morgenstern, M. *et al.* Quantitative high-confidence human mitochondrial proteome and its dynamics in
cellular context. *Cell Metab* **33**, 2464-2483 e2418 (2021). https://doi.org:10.1016/j.cmet.2021.11.001
65   Kohler, A., Barrientos, A., Fontanesi, F. & Ott, M. The functional significance of mitochondrial respiratory
chain supercomplexes. *EMBO Rep* **24**, e57092 (2023). https://doi.org:10.15252/embr.202357092
66   Jia, X., Gu, Z., Chen, W. & Jiao, J. Tigecycline targets nonsmall cell lung cancer through inhibition of
mitochondrial function. *Fundam Clin Pharmacol* **30**, 297-306 (2016). https://doi.org:10.1111/fcp.12199
67   Ren, A., Qiu, Y., Cui, H. & Fu, G. Tigecycline exerts an antitumoral effect in oral squamous cell
carcinoma. *Oral Dis* **21**, 558-564 (2015). https://doi.org:10.1111/odi.12311
68   Skrtic, M. *et al.* Inhibition of mitochondrial translation as a therapeutic strategy for human acute myeloid
leukemia. *Cancer Cell* **20**, 674-688 (2011). https://doi.org:10.1016/j.ccr.2011.10.015
69   Tang, C. *et al.* Antibiotic drug tigecycline inhibited cell proliferation and induced autophagy in gastric
cancer cells. *Biochem Biophys Res Commun* **446**, 105-112 (2014).
https://doi.org:10.1016/j.bbrc.2014.02.043
70   Yang, R. *et al.* Tigecycline Inhibits Glioma Growth by Regulating miRNA-199b-5p-HES1-AKT Pathway.
Mol Cancer Ther **15**, 421-429 (2016). https://doi.org:10.1158/1535-7163.MCT-15-0709
71   Zhong, X. *et al.* Antibiotic drug tigecycline reduces neuroblastoma cells proliferation by inhibiting Akt
activation in vitro and in vivo. *Tumour Biol* **37**, 7615-7623 (2016). https://doi.org:10.1007/s13277-015-
4613-6
72   Nair, A. B. & Jacob, S. A simple practice guide for dose conversion between animals and human. *J*
Basic Clin Pharm **7**, 27-31 (2016). https://doi.org:10.4103/0976-0105.177703
73   Reed, G. A. *et al.* A Phase 1 study of intravenous infusions of tigecycline in patients with acute myeloid
leukemia. *Cancer Med* **5**, 3031-3040 (2016). https://doi.org:10.1002/cam4.845
74   Helmink, B. A., Khan, M. A. W., Hermann, A., Gopalakrishnan, V. & Wargo, J. A. The microbiome,
cancer, and cancer therapy. *Nat Med* **25**, 377-388 (2019). https://doi.org:10.1038/s41591-019-0377-7
75   Nobels, A., van Marcke, C., Jordan, B. F., Van Hul, M. & Cani, P. D. The gut microbiome and cancer:
from tumorigenesis to therapy. *Nat Metab* **7**, 895-917 (2025). https://doi.org:10.1038/s42255-025-
01287-w

Figure 1.

A. NACT: DTX + CRB (CPTAC, PMID: 36001024) → Surgery: pCR or RCB
n=48

Baseline

n=14 On-treatment (Day 3)

Baseline nonpCR vs. pCR
- Reactome: Mitochondrial translation
- Reactome: Mitochondrial translation initiation
- Reactome: Mitochondrial translation elongation
- Reactome: Mitochondrial translation termination
- Mitocarta: Translation
- Mitocarta: Mitochondrial central dogma
- Mitocarta: Mitochondrial ribosome
- Mitocarta: mtRNA metabolism
- Mitocarta: mt-tRNA synthetases

0.0 0.5 1.0 1.5 2.0 2.5
NES

On-treatment vs. baseline
- Mitocarta: Translation
- Mitocarta: Mitochondrial central dogma
- Mitocarta: mtRNA metabolism
- Mitocarta: mt-tRNA synthetases

0.0 0.5 1.0 1.5 2.0 2.5
NES

B. Reactome geneset (FDR < 0.05)

$-\log_{10}$(FDR)

Mitochondrial translation
Mitochondrial translation initiation
Mitochondrial translation elongation
Mitochondrial translation termination

10.0
7.5
5.0
2.5
0.0

Lower in chemoresistant
Higher in chemoresistant

-2 -1 0 1 2
Normalized Enrichment Score

C. 4 x q7d DTX or CRB (PMID: 39713418) → Relative TV
n=55

DTX

$-\log_{10}$ (p-value)

- OXA1L
- MRPS
- MRPL
- Other
p<0.05

3
2
1
0

MRPS11 OXA1L
MRPL21
MRPL43 MRPS22
MRPL47 MRPS40
MRPS9 MRPS5
MRPL15 MRPL54
MRPS26 MRPS33
MRPS31 MRPS18C
MRPS24

Down in resistant
Up in resistant

CRB

MRPL49
MRPL52
MRPL14 MRRF
MRPL15 MRPL30
MRPL13 MRPL23
MRPL19 OXA1L
MRPL50 MRPS30
MRPL28

Down in resistant
Up in resistant

-0.6 -0.3 0.0 0.3 0.6 -0.6 -0.3 0.0 0.3 0.6
Spearman correlation coefficient

D.

PDXs

OXA1L protein abundance ($\log_{10}$ intensity)

p=0.04

CR   Non-CR

CPTAC-TNBC

pCR response ⊞ Yes ⊞ No
Paired

OXA1L protein abundance ($\log_{10}$ intensity)

1
0
-1
-2

p = 0.01

Baseline   On-TX

Figure 2.

A. MDA-MB-231

B.

C.

D.

SDS-PAGE

Native Coomassie

E. Native WB: UQCRC2 (CIII)

F. Native WB: ATP5A (CV)

Figure 3.
A. MDA-MB-231

Figure 4.

A. MDA-MB-231

Relative viable cells / Log Concentration (nM)

$IC_{50} = 8 \mu M$

B.

VEH / TIG

CI — NDUFB8
MT-ND3
CII — SDHB
CIII — UQCRC2
COX4
CIV — MT-COX1
MT-COX2
CV — ATP5A
MT-ATP8
Total C3
CC3
β-actin
TOM70
β-tubulin

C.

% mtDNA/nucDNA copy number

ns

VEH / TIG

D.

● Vehicle ● Tig

Relative mRNA level

****

*

OXA1L MT-ND1 MT-ND2 MT-COX2 MT-COX3 MT-ATP8

E.

VEH / TIG
M C M C

UQCRC2
TOM70
β-tubulin

SDS-PAGE

Ladder VEH TIG

(kDa)

Native Coomassie

F.

VEH / TIG

SC: I+III₂+IVₙ (Respirasomes) — 1236 / 1048
SC: III₂+IV₂ — 720
SC: III₂+IV — 720
III₂¹ —
480
242

Extended Exposure

SC: I+III₂+IVₙ (Respirasomes) — 1236 / 1048
SC: III₂+IV₂ — (kDa)

Native WB: UQCRC2 (CIII)

% of UQCRC2 in SC (vs total)

**

VEH / TIG

G.

● VEH ● TIG

Normalized OCR (pmol/min/Norm. Unit) / Time (minutes)

FCCP
Oligo
Rot/AA

Normalized basal OCR (pmol/min) ** VEH / TIG

Normalized maximal OCR (pmol/min) * VEH / TIG

Normalized basal ECAR (pmol/min) * VEH / TIG

Normalized basal OCR/ECAR (pmol/min) ns VEH / TIG

Figure 5.

A. MDA-MB-231

Figure 6.

A. MDA-MB-231

B. SUM159pt

C. MDA-MB-231

D. SUM159pt

Figure 7.