A metabolic model based on a pangenome core unveils new biochemical

features of the phytopathogen *Xylella fastidiosa*

Corbín-Agustí, Paola[1*]; Álvarez-Herrera, Miguel[1]; Román-Écija, Miguel[2]; Álvarez, Patricia[1] ;

Tortajada, Marta[3]; Landa, Blanca B.[2*]; Peretó, Juli[1,4]

[1]Institute for Integrative Systems Biology, I2SysBio (CSIC-Universitat de València), Paterna, Spain.

[2]Institute for Sustainable Agriculture (IAS), Spanish National Research Council (CSIC), Córdoba,

Spain.

[3]Department of Biomedical Sciences, Faculty of Health Sciences, Universidad Cardenal Herrera,

Centro de Estudios Universitarios, UCH-CEU, Alfara del Patriarca, Spain.

[4]Department of Biochemistry and Molecular Biology, University of Valencia, Burjassot, Spain.

*Corresponding authors: Blanca B. Landa blanca.landa@ias.csic.es, Paola Corbín Agustí

paola.corbin@uv.es

15   **Abstract**

Xylella fastidiosa is a xylem-limited phytopathogen bacterium responsible for severe diseases

in numerous plant species of major agricultural importance. Despite its economic impact, its

metabolism remains poorly characterized due to the bacterium's fastidious growth and the

limited availability of defined culture media. In this work, we reconstructed the first

pangenome-based genome-scale metabolic model for *X. fastidiosa*, integrating the conserved

metabolic capabilities of 18 strains representing five described subspecies. The resulting core

metabolic model, Xfcore, was manually curated and used to investigate the metabolic potential

of the species. Model simulations predict minimal nutritional requirements that guide the

formulation of defined media capable of supporting biofilm formation in vitro. Analysis of the

metabolic network also suggests an undescribed metabolic pathway that enables growth on

acetate as a sole carbon source. Furthermore, the model predicts that *X. fastidiosa* could

overproduce polyamines, compounds previously associated with virulence in other

phytopathogens. Experimental analyses confirm the production and secretion of polyamines

in several *X. fastidiosa* strains, providing the first in vitro evidence of polyamine production in

this pathogen. These findings suggest that polyamine biosynthesis may represent an

uncharacterized virulence factor in *X. fastidiosa*, potentially contributing to bacterial protection

against host-induced oxidative stress. Overall, the Xfcore model provides a systems-level

framework to explore the metabolism of *X. fastidiosa*, generate testable hypotheses about its

physiology and virulence, and establish a basis for future strain-specific reconstructions and

host-pathogen metabolic interaction studies.

Keywords: *Xylella fastidiosa* pangenome, metabolic model, polyamines, acetate metabolism,

defined medium

Introduction

Xylella fastidiosa is a fastidious, Gram-negative, xylem-limited phytopathogenic bacterium that infects more than 700 plant species worldwide (EFSA et al., 2026), including several crops of major agro-economic relevance. In some of these hosts, the bacterium causes devastating diseases such as Pierce's disease (PD) in grapevine, citrus variegated chlorosis (CVC), almond leaf scorch (ALS), and olive quick decline syndrome (OQDS). The bacterium colonizes the xylem vessels of its plant hosts, and symptoms development depends on several bacterial virulence factors, particularly the formation of biofilm and host defense responses (Landa et al., 2022; De La Fuente et al., 2024). Biofilms also develop in the anterior foregut of sap-feeding insect vectors, which mediate bacterial transmission (Chatterjee et al., 2008).

Xylella fastidiosa is also characterized by substantial intraspecific diversity. Recent computational analyses based on k-mer signatures support a revised taxonomy that groups the species into three major clades: clade I (corresponding to subsp. *pauca*), clade II (subsp. *multiplex*) and clade III (including three subsp.: *fastidiosa*, *morus* and *sandyi*) (Denancé et al., 2019). In parallel, multilocus sequence typing (MLST) has been widely used for the epidemiological characterization of isolates (Scally et al., 2005) and 90 sequence types (STs) have been described to date (Jolley et al., 2018; EPPO 2023). Although STs do not provide genome-wide resolution and therefore cannot capture the full extent of gene-content variation, they remain useful for rapid strain typing, subspecies assignment, and initial epidemiological interpretation. However, the high recombination rates and the existence of some non-monophyletic STs raise concerns about the phylogenetic resolution and the robustness of current classification schemes based solely on MLST (Hanage et al., 2005; Vanhove et al., 2019; Landa et al., 2020). This genomic complexity suggests that marker-based classification alone is insufficient to infer functional traits, including metabolic capabilities that may influence host association, niche adaptation, and pathogenicity (Landa et al., 2022).

Despite decades of research, key aspects of *X. fastidiosa* physiology –particularly those linked to virulence, host adaptation, and nutritional requirements– remain poorly understood. This knowledge gap is especially relevant because the bacterium inhabits the xylem, a nutritionally restricted environment in which successful colonization, multiplication, and biofilm formation are expected to depend strongly on metabolic adaptation. Its inability to grow efficiently on standard bacteriological media, reflected in the species epithet "*fastidiosa*", continues to be a defining feature of the species (Davis et al., 1981; Wells et al., 1987; Chatterjee et al., 2008). Empirical efforts have produced improved culture media that support faster growth (Davis et al., 1980; Wells et al., 1981; Davis et al., 1981), but none fully overcome its fastidiousness

(Campanharo et al., 2003). Moreover, growth in currently available defined media remains
slow and biomass production is limited. As a result, the metabolic basis of nutrient use,
physiological specialization, and virulence-associated traits in *X. fastidiosa* is still only partially
resolved (De La Fuente et al., 2024).

Genome-scale metabolic models (GEMs) are structured knowledge bases that integrate
biochemical, genetic and genomic data into a system-level representation of the metabolic
capabilities of an organism (O'Brien et al., 2015). Over the past two decades, GEMs have
been applied in systems biology, enabling in silico prediction of genotype-phenotype
relationships, reverse ecology paradigm to infer ecological features from genomic data, such
as nutrient requirements prediction and growth medium design, and the discovery of metabolic
network properties or pan-reactome comparisons (Yus et al., 2009; Monk et al., 2014; Gu et
al., 2019; Ye et al., 2022; Qian & Ye, 2024). Constraint-based reconstruction and analysis
(COBRA) is one of the foundational and most widely used approaches for working with GEMs.
It achieves high predictive power by imposing physicochemical and environmental constraints
(such as stoichiometry, thermodynamics and nutrient availability) that define a solution space
encompassing all feasible metabolic states of the cell under different conditions (Ibarra et al.,
2002; Palsson, 2015).

Here, we present an in silico approach to investigate key aspects of *X. fastidiosa* physiology
through the reconstruction of a GEM. Although two metabolic models have previously been
reconstructed for this phytopathogen, both are strain-specific (Gerlin et al., 2020; Oliveira et
al., 2023). In this study, to account for the genetic diversity within the species while focusing
on its conserved metabolic functions, a pangenome was assembled and used it as the
template for a core metabolic reconstruction. The resulting model represents a species-level
GEM that reflects metabolic functions conserved across *X. fastidiosa,* rather than traits
restricted to particular subspecies or strains. Then, the resulting reconstruction was
experimentally validated by growth-phenotype profiling of a selection of *X. fastidiosa* strains
via Biolog Phenotyping assays. We also used the model to design a defined minimal growth
medium based on predicted nutritional requirements –an aspect that has received limited
attention despite *X. fastidiosa* being the first phytopathogenic bacterium with a fully sequenced
genome (de Macedo Lemos et al., 2003). Beyond this, exploration of the metabolic network
highlighted two notable features of the bacterial physiology. First, given the absence in *X.*
fastidiosa of previously described acetate assimilation pathways, we sought a metabolic
solution that would explain growth on acetate as the sole carbon source. The resulting
pathway emerges from the combination of submodules from other metabolic routes, providing
genomic support for acetate utilization suggested by model-based analyses. Second, the

model predicts a remarkable capacity for polyamine biosynthesis. Polyamines are metabolites
that are not only produced by plants as part of their oxidative stress and defense responses,
but have also been previously linked to virulence in other phytopathogens such as *Ralstonia*
solanacearum (Lowe-Power et al., 2018) and *Pseudomonas syringae* (Solmi et al., 2025). We
experimentally confirmed polyamine production in *X. fastidiosa*, suggesting a possible role in
host-pathogen interactions. Together, this work delivers the first species-level GEM of *X.*
fastidiosa and establishes a systems biology framework to investigate its physiology,
nutritional requirements, and metabolic determinants of pathogenicity.

Results

Metabolic characteristics of the *Xylella fastidiosa* pangenome and the core metabolic model

A curated genome-scale metabolic reconstruction was generated based on the core genes of
18 *X. fastidiosa* strains representing the subspecies *fastidiosa*, *pauca*, *multiplex*, *morus* and
sandyi (Table 1), and including the strains previously used for metabolic modelling (Gerlin et
al., 2020; Oliveira et al., 2023). These strains were isolated from a diverse set of host plants,
including grapevine, almond, sweet orange, olive, coffee and several wild species. This host
diversity reflects the broad ecological range of the species and provides a representative basis
for assessing its pangenomic and metabolic variability. Moreover, this pangenome also
incorporates recently emerged European strains, with six isolates from Spain, Italy and
France, in order to account for the lineages involved in the outbreaks starting to be detected
in 2013, which have resulted in substantial agricultural and economic losses in Europe
(Sanchez et al., 2019; De la Fuente, 2024).

Pangenome analysis revealed a total of 2660 ortholog groups (OGs), comprising 40828
genes, of which 1372 OGs included genes found in the 18 strains (Fig. 1A). The core genome
consists of 26831 genes, constituting 65.72% of the genes analyzed. The accessory genome
includes 13421 genes, corresponding to 32.87% of the total pangenome. Among the analyzed
strains, XYL468 exhibited the highest number of accessory genes (890), whereas Salento-1
displayed the lowest (595). A total of 576 genes were classified as unique, accounting for
1.41% of all genes. The strain with the largest number of unique genes was XYL1961/18, with
141 unique genes, while U24D contained only 2 unique genes. Only a limited number of OGs
showed strict subspecies-specific distribution.

Functional annotation of the pangenome revealed clear differences between the core,
accessory, and unique gene sets (Fig. S1). Core ortholog groups were predominantly enriched
in metabolic functions, particularly carbohydrate, amino acid, nucleotide, energy and glycan
metabolism, as well as metabolism of cofactors and vitamins. These were followed in
prevalence by genes associated with genetic information processing and with signalling and
cellular processes. In contrast, accessory genes showed a stronger representation of the latter
two categories, including a higher proportion of functions associated with prokaryotic defense
systems, secretion and transport systems, and quorum-sensing pathways, alongside a
substantial fraction of genes with unknown function. Metabolic genes were also present within
the accessory set, although to a lesser extent. For the unique gene set, only 6.2% (36 out of
576) of the genes could be functionally annotated, and none were linked to core metabolic
pathways. Instead, the annotated genes mapped to categories such as prokaryotic defense
systems, quorum sensing, or DNA replication and repair.

The core ortholog set identified in the pangenome analysis was used to generate a draft
genome-scale metabolic reconstruction using the automated reconstruction software
ModelSEED (Henry et al., 2010). Curation was performed iteratively. Initial efforts focused on
central metabolism and biomass production. Subsequently, complete biosynthetic routes for
amino acids, lipids, nucleotides, energy metabolism, and selected transport systems were
manually reviewed. Pathways associated with redox balance, cofactor biosynthesis, and host-
associated growth were carefully curated, including quinone and shikimate biosynthesis,
polyamine metabolism, the S-adenosylmethionine (SAM) cycle, sulfur and nitrogen
metabolism, and the methionine salvage pathway. Vitamin and cofactor biosynthetic pathways
were curated in detail, encompassing pantothenate, folate, riboflavin, thiamine, and porphyrin
metabolism. Finally, pathways associated with cell envelope biogenesis and host interaction
were refined, including exopolysaccharide (EPS) and lipopolysaccharide (LPS) biosynthesis,
as well as fatty acid metabolism, and the biosynthesis and export of the LesA protein, following
previous strain-specific metabolic reconstructions (Gerlin et al., 2020; Oliveira et al., 2023).

During this iterative process, a set of accessory genes were identified as present in the
majority of the strains, and considered relevant for modelling the general metabolic behavior
of *X. fastidiosa* (Table S1). These genes were not included into the core set of sequences due
to pseudogenization events in specific strains. In several cases, this was the result from
frameshift mutations caused by single-nucleotide insertions or deletions within repetitive
nucleotide regions, which may allow some enzymatic activity or reflect sequencing artefacts
(Cooley & Wright, 2024). For instance, the accessory gene XF_RS10205, encoding 5'-
methylthioadenosine phosphorylase, is annotated as a pseudogene in strain 9a5c despite

being reviewed in UniProt (Q9PAZ2) as a functional enzyme. In total, 43 accessory genes were incorporated into the metabolic reconstruction.

The final genome-scale metabolic reconstruction (Xfcore; Supplementary Material S1) included 1034 reactions, 1075 metabolites (corresponding to 1013 unique metabolites), 451 genes, and two cellular compartments: cytoplasm and extracellular space (Fig. 1B). Of the metabolites, there were 63 extracellular and 1012 cytosolic. The model includes 69 boundary reactions (exchange, sink, and demand reactions), and 76 transport reactions enabling metabolite exchange between compartments. Among the 451 genes, 407 belong to the core pangenome, while 43 are accessory genes (Table S1). Additionally, there is a gene called "Unknown", indicating a functional requirement in the network despite the absence of genetic evidence. A total of 83% of reactions (859 out of 1034) are associated with gene-protein-reaction (GPR) rules. Of the 175 reactions lacking GPR associations, 99 correspond to boundary or transport reactions, the majority of which do not require gene assignment. Model quality and consistency evaluated using the Memote testing suite (Lieven et al., 2020), yielded a total score of 85% and a consistency score of 97% (Supplementary Material S2). Annotation completeness scores were 66% for metabolites, 73% for reactions, and 53% for genes. The coverage of SBO terms reached 83%, indicating strong semantic and structural annotation.

The core gene set of *X. fastidiosa* encoded nearly all central metabolic functions, although several specific gaps and distinctive features were identified. The Embden-Meyerhof-Parnas (EMP) and the Entner-Doudoroff (ED) glycolytic pathways were fully represented in the model. However, one enzyme from the gluconeogenic pathway –namely, fructose-1,6-bisphosphatase (FBPase)– was not identified in any gene set of the pangenome. This absence would suggest limitations in the performance of gluconeogenesis. Nevertheless, extensive experimental evidence indicates that *X. fastidiosa* is capable of converting pyruvate into sugars (Davis et al., 1980, 1981; Almeida et al., 2004; Gerlin et al., 2020). As previously proposed by Oliveira et al. (2023), FBPase activity (EC 3.1.3.11) can be functionally replaced by a reversible pyrophosphate-dependent phosphofructokinase (PPi-PFK; EC 2.7.1.90), encoded by gene XF_RS01160. This enzyme, homologous to the PPi-PFK described in *Xanthomonas campestris* (UniProt: B0RP51), is reversible and can therefore catalyze both the forward glycolytic reaction and the reverse gluconeogenic conversion. Accordingly, this reaction was implemented in the model to account for both activities.

The oxidative and non-oxidative branches of the pentose phosphate pathway (PPP) were evaluated in the core gene set. Glucose-6-phosphate dehydrogenase (EC 1.1.1.49) –an enzyme shared with the ED pathway– was present in all strains except Pr8x, where it appears

as a pseudogene; nevertheless, this activity was retained in the model to reflect its functional relevance. Two additional activities were absent from the core pangenome: 6-phosphogluconate dehydrogenase (PPP oxidative phase) and transaldolase (TALA, PPP non-oxidative phase). Although the absence of the oxidative branch of the pentose phosphate pathway prevents NADPH generation through this route, reducing power may be supplied by the ED pathway and additional reactions such as the malic enzyme. No transhydrogenase activity was identified in the model. Flux redistribution through the reversible reactions of the PPP non-oxidative branch allows synthesis of essential precursors such as phosphoribosyl pyrophosphate (PRPP). In contrast, the absence of TALA is expected to impair growth on pentoses as sole carbon sources, as it disrupts sedoheptulose-7-phosphate (S7P) metabolism. In previously published models (Oliveira et al., 2023), growth on ribose was enabled by incorporating a sedoheptulose-1,7-bisphosphate (SBP) bypass reaction that converts S7P into SBP using PPi instead of ATP, and subsequently cleaved into dihydroxyacetone phosphate (DHAP) and erythrose-4-phosphate (E4P). Experimental and computational studies have shown that bacteria lacking TALA can metabolize S7P through an SBP aldolase pathway (Nakahigashi et al., 2009; Koendjbiharie et al., 2020; Garschagen et al., 2021). For the phosphorylation step, we assigned this activity to the pyrophosphate-dependent phosphofructokinase (PPi-PFK; XF_RS01160), which has been shown in other bacteria to catalyze reversible PPi-dependent phosphoryl transfer reactions (Koendjbiharie et al., 2020). For the aldolase reaction, which converts SBP to DHAP and E4P, we hypothesized that the fructose-bisphosphate aldolase encoded by XF_RS03490 may exhibit substrate promiscuity and catalyze both FBP and SBP cleavage. Such dual specificity has been described in other bacteria, including *Synechocystis* sp. PCC 6803, where class I and II aldolases can participate in SBP metabolism in MetaCyc.

Every enzymatic step of the tricarboxylic acid (TCA) cycle was represented within the core pangenome. The oxoglutarate dehydrogenase complex (ODC) was supported by all its subunits, although the E1 component (XF_RS06555) was absent in strain 3124. The E2 component (XF_RS06550) was conserved in the core genome. The pyruvate dehydrogenase complex (PDC), which catalyzes an analogous reaction and shares a similar multienzyme architecture with ODC, showed strain-specific variation in the E2 subunit. The gene encoding the E2 component (XF_RS03665; dihydrolipoyllysine-residue acetyltransferase) was annotated as a pseudogene in strains XYL468 and CFBP8418, however, inconsistencies between genome assemblies in NCBI suggest potential sequencing or annotation artefacts. So, this complex was conserved in the metabolic model. Two candidate genes encoding the E3 subunit (dihydrolipoyl dehydrogenase) were identified in the pangenome (XF_RS06540

and XF_RS03660). Only XF_RS06540 was part of the core gene set and was therefore
assigned as the shared E3 component for both ODC and PDC in the model. Importantly, no
enzymes associated with the glyoxylate shunt were identified in any analyzed strain.

Remarkably, pyruvate carboxylase (EC 6.4.1.1), which catalyzes the ATP-dependent
conversion of pyruvate to oxaloacetate (OAA), was not identified in the *X. fastidiosa*
pangenome and was therefore not included in the model. No alternative reaction directly
linking pyruvate to oxaloacetate was detected. The absence of this anaplerotic step limits the
replenishment of tricarboxylic acid (TCA) cycle intermediates when sugars or organic acid
compounds are supplied as sole carbon sources. Under these conditions, growth requires
TCA intermediates to sustain biosynthetic demands. The reconstructed network lacks
sufficient anaplerotic reactions to compensate for carbon withdrawal from the cycle, resulting
in growth impairment in silico. To evaluate whether growth could be sustained using plant
metabolites, we considered incorporating TCA cycle intermediates or their precursors –e.g.,
glutamine or glutamate– which could be supplied by the host, to simulate bacterial growth.

As previously described, the core metabolic network of *X. fastidiosa* contained complete
biosynthetic pathways for all amino acids. Lipid biosynthesis pathways were also identified
and included in the reconstruction, whereas no evidence of lipid degradation pathways –e.g.,
fatty acids β-oxidation– was detected in the pangenome. Additionally, no auxotrophies were
identified for essential vitamins or cofactors.

As for quinone-dependent electron transport, only the ubiquinone biosynthetic pathway was
identified in the pangenome, although it appeared incomplete. No candidate gene was
detected for two intermediate reactions corresponding to the decarboxylation and
hydroxylation of 3-polyprenyl-4-hydroxybenzoate. Nevertheless, the upstream and
downstream steps of the pathway were conserved, and no alternative quinone biosynthetic
route was identified. Given the apparent continuity of the pathway and the absence of
pseudogenization in adjacent enzymatic steps, it is likely that these missing reactions are
catalyzed either by highly divergent enzymes not detected by sequence homology, by
promiscuous enzymatic activities, or through non-canonical mechanisms that remain to be
characterized.

Finally, the biomass reaction was formulated to include all known cellular components and
their respective contributions per gram of biomass. Following extensive curation, including
comparative evaluation against the biomass formulations of the *Escherichia coli* model
iJO1366 (Orth et al., 2011) and the *X. fastidiosa* model Xfm1158 (Gerlin et al., 2020), the
biomass equation included 78 metabolites.

Flux balance analysis (FBA) predicted a maximum specific growth rate of 0.413 h⁻¹ when

glutamine, one of the major components of xylem sap, was supplied as the sole carbon source

at an uptake rate of 7.25 mmol·gDW⁻¹·h⁻¹ (Supplementary Material S3). The corresponding

optimal flux distribution is shown in Fig. 1C. Under these conditions, most of the imported

glutamine is first converted into glutamate and subsequently into α-ketoglutarate, which enters

the tricarboxylic acid (TCA) cycle. Carbon flux proceeds through the cycle to generate malate

and oxaloacetate. Malate is partially converted to pyruvate via the malic enzyme, generating

NADPH and contributing to cellular reducing power. Pyruvate is then redirected toward

gluconeogenesis to supply precursors required for lipid, cell wall, and other biomass

components. In parallel, a fraction of pyruvate is converted into acetyl-CoA, which supports

fatty acid biosynthesis and citrate formation, ultimately reconnecting with α-ketoglutarate

within the TCA cycle. The Xfcore model further predicted uptake of inorganic ions, phosphate,

and sulfate under these growth conditions, while CO_2 and NH_3 were secreted as metabolic

by-products. Oxygen was also required, as the model did not predict growth under anaerobic

conditions.

Experimental validation of the metabolic reconstruction

Phenotypic characterization of *Xylella fastidiosa* strains

The phenotypic characterization of strains ESVL, De Donno, IVIA5901, Temecula1 and

XYL1961/18 using the phenotype microarray system of Biolog (Supplementary Material S4,

Figs. S2 and S3) showed partially overlapping metabolic fingerprints. Among strains, De

Donno showed the highest proportion of assimilated carbon sources (31.6% and 38.9% for

PM1 and PM2, respectively), whereas Temecula1 showed the lowest (26.3% and 21.1%,

respectively) (Table S2). Overall, the five strains consistently metabolized 34 out of the 190

compounds tested (17.9%): 17 compounds in PM1 and 17 compounds in PM2 plates (Table

2; Fig. S4 and S5). Additionally, five compounds were classified as exhibiting "Undetermined"

growth across all strains, due to positive growth in only two of the four biological replicates. In

contrast, 50 compounds were not assimilated by any of the strains (Supplementary Material

S4; Figs. S4 and S5).

The main groups of compounds consistently utilized as carbon sources by all *X. fastidiosa*

strains in the PM1 and PM2 plates included i) carbohydrates and their derivatives, ii) organic

acids, iii) lipids and fatty acids, and iv) amino acids derivatives and amines. These in vitro

results were compared with in silico growth predictions for each compound, when possible (Table 2, Supplementary Material S5).

In silico predictions were consistent with in vitro results for 7 out of the 34 metabolites that supported growth in vitro (D-ribose, D-xylose, D-fructose-6-phosphate, D-glucosamine, α-ketoglutaric acid, pyruvic acid, and acetic acid) (Table 2, Supplementary Material S5). Nineteen out of the 34 metabolites could not be evaluated using the Xfcore model because they were not included in the reconstructed metabolic network or their metabolism could not be represented. These included all lipids and fatty acids, as well as amino acid derivatives and amines. For the remaining eight metabolites, growth was observed in vitro but could not be supported in silico (L-arabinose, L-lyxose, 2-deoxy-D-ribose, D-psicose, D-tagatose, α-ketobutyric acid, acetoacetic acid, and dihydroxyacetone).

Additionally, several compounds that did not support growth for all strains in the PM plates were evaluated in silico due to their reported abundance in the xylem sap of olive, almond, and citrus trees (Table 2, Supplementary Material S5) (Youssefi et al., 2000; Purcino et al., 2007; Anguita-Maeso et al., 2021; Fausto et al., 2021; Surano et al., 2024). D-glucose, one of the major sugars detected in olive xylem sap, was predicted to sustain growth in silico, consistent with previous strain-specific models (Gerlin et al., 2020; Oliveira et al., 2023). However, this prediction was not supported experimentally, as only one strain showed undetermined growth while the remaining four exhibited no growth in the PM assays (Fig. S4). Citric acid, also reported as a common organic acid in xylem sap, was predicted to sustain growth in silico. The corresponding transporter was present in the core pangenome, and growth has been supported in other reconstructions. Experimentally, citrate supported growth in one strain, showed indeterminate growth in one strain, and resulted in no growth in the remaining three strains.

Regarding amino acids commonly reported in xylem sap (glutamine, glutamate, aspartate, arginine, asparagine, alanine, serine, and proline), in silico simulations predicted growth on L-glutamine, L-glutamate, L-aspartate, and L-alanine (Table 2, Supplementary Material S5). However, experimental validation showed partial or limited agreement (Fig. S4 and S5). L-aspartate supported growth in vitro in two strains, while one strain showed undetermined growth and two showed no growth. In contrast, L-glutamate, L-glutamine, and L-alanine predominantly resulted in undetermined or no-growth phenotypes across strains despite being predicted to sustain growth in silico. L-arginine, L-asparagine, L-serine, and L-proline were not predicted to support growth in silico. Consistently, these amino acids did not sustain robust growth in vitro, with most strains showing either no growth or undetermined phenotypes.

Growing *Xylella fastidiosa* in minimal media predicted by the model

Empirical efforts have led to the development of several culture media for *X. fastidiosa*, including PD2 (Davis et al., 1980), BCYE (Wells et al., 1981) and PW (Davis et al., 1981). However, these formulations do not fully overcome the fastidious growth requirements of the bacterium. In this work, a reverse ecology approach based on constraint-based metabolic modeling was applied to predict a synthetic defined minimal medium capable of sustaining biomass production in the Xfcore model. Under a fixed L-glutamine uptake constraint (7.25 mmol·gDW^{-1}·h^{-1}), the minimal medium prediction identified oxygen, phosphate, sulfate, glutamine (as both carbon and nitrogen source), and inorganic ions as required components (see Supplementary Material S6 for the complete list of metabolites and corresponding uptake fluxes).

Based on these predicted requirements, six synthetic defined minimal media were formulated (mXC1–mXC6; Table S3). Glutamine was used as the sole carbon and nitrogen source in mXC1 and mXC2, given its abundance in xylem sap, despite not supporting growth in the phenotypic characterization plates. Two different iron sources were tested in these media, ferric pyrophosphate and hemin chloride, to assess iron bioavailability. Acetate was used as the carbon source in mXC3 and mXC4, supplemented with ammonium sulfate as a nitrogen source. To validate model predictions regarding biotin dependency under acetate growth conditions (see below Acetate metabolism section), mXC4 was additionally supplemented with biotin. Finally, ribose was tested as the sole carbon source in mXC5 and mXC6. In mXC6, cysteine was evaluated as an alternative nitrogen source, as it showed strong utilization in nitrogen assimilation assays (Biolog PM3 plate; data not shown).

All minimal media supported, to some extent, the growth and biofilm formation in all *X. fastidiosa* strains studied (Fig. 2; Supplementary Material S6). Interestingly, all strains were able to grow with glutamine media, despite glutamine not supporting growth in the previous phenotypic characterization of Biolog PM plates. The reference medium PD3 significantly supported higher sAUGPC values than all minimal media across strains (Fig. 2A), except for XYL1961/18, which showed higher growth in the Glutamine+Fe formulation (mXC1) and no significant differences were detected for the rest of the media. In strain XYL468, no significant differences were observed between PD3 and Glutamine+Hemin or Ribose+Cysteine media. sAUGPC values in PD3 ranged from 0.002±0.0011 to 0.072±0.1344, being between 3.3 to 58.4-fold higher than those obtained in the minimal media.

On the other hand, biofilm formation (Fig. 2B) showed comparable levels across most minimal
media relative to PD3. The exception was strain De Donno, which produced significantly
higher biofilm in the control medium (PD3). Strain IVA5901 exhibited enhanced biofilm
formation in Ribose+NH_4 compared to PD3. Overall, a general trend was observed in which
synthetic defined media containing ribose supported higher biofilm production across most
strains, followed by acetate-based media.

Pairwise comparisons between media formulations sharing the same carbon source revealed
no significant differences for growth or biofilm formation associated with iron source, biotin
supplementation in acetate media, or nitrogen source substitution in ribose-based
formulations (data not shown).

Acetate metabolism in *Xylella fastidiosa*

Experimental evidence supports the ability of *X. fastidiosa* to grow on acetate as the sole
carbon source in several strains (Gerlin et al., 2020; this work). In this work, all previously
described metabolic solutions to produce biomass from acetate were evaluated (see Material
and Methods). These included: i) the glyoxylate cycle (Kornberg & Krebs, 1957); ii) a
ferredoxin-dependent pyruvate synthase (Segura et al., 2008), typically associated with
anaerobic metabolism; iii) the ethylmalonyl-CoA (EMC) pathway (Erb et al., 2007); iv) the
methylaspartate cycle (MaC) characteristic of haloarchaea (Khomyakova et al., 2011); and v)
part of the 3-hydroxypropionate (3HP) autotrophic bi-cycle (Herter et al., 2002). Across the
18 strains comprising the pangenome, none of these previously described pathways were fully
supported by genomic evidence, lacking all or some of the necessary enzymes for these
routes. Alternative glyoxylate and propionyl-CoA assimilation modules, including the
methylcitrate cycle (Brämer & Steinbüchel, 2001; Dolan et al., 2018), were also examined.

The Xfcore metabolic model predicted growth on acetate through an alternative, yet
undescribed metabolic solution (Fig. 3). This consisted of a modular pathway with 11 steps
(Table S4), connecting acetate to central carbon metabolism, generating pyruvate and
succinate and enabling net carbon assimilation into biomass. This pathway integrates two
enzymatic subsets present in previously described metabolic routes. The first one corresponds
to the initial half of the 3HP bi-cycle, linking acetyl CoA to the synthesis of propionyl-CoA (the
remaining reactions of the cycle are absent from the pangenome). The second subset
corresponds to the methylcitrate pathway, which enables the assimilation of propionyl-CoA
into pyruvate and succinate. During the reconstruction process, minor gaps were identified in

the draft network. These gaps were resolved through manual genome evaluation and literature

reviewing. All reactions included in the final pathway are supported by genes present in the

pangenome. Below we describe the route in more detail, indicating the specific steps shown

in Fig. 3.

First, acetate enters the cell through a sodium-dependent acetate symporter (XF_RS09795),

which is present in all strains analyzed. Subsequently, acetate is converted into acetyl-CoA

by acetate-CoA ligase (step 1; XF_RS09805). This gene is absent in the J1A12 strain due to

a single nucleotide deletion (loss of an adenine at position 1120) within a repetitive region of

adenine. Apart from this frameshift mutation, the sequence is identical to that found in the

remaining strains. In step 2, acetyl-CoA carboxylase catalyzes the ATP-dependent

carboxylation of acetyl-CoA to produce malonyl-CoA. This enzyme requires biotin as a

cofactor for its catalytic activity. Malonyl-CoA is then reduced to malonate semialdehyde (3-

oxopropanoate; step 3) by a NADP-dependent malonyl-CoA reductase. Although the

associated gene (XF_RS05765) is annotated as aspartate semialdehyde dehydrogenase,

BLAST analysis and Pfam domain architecture match the reviewed UniProt entry A4YEN2

from *Metallosphaera sedula*, which performs this specific reaction. Step 4 involves the

conversion of malonate semialdehyde to 3-hydroxypropionate, catalyzed by 3-

hydroxypropionate dehydrogenase. This reaction was incorporated into the model and

assigned to gene XF_RS00605, annotated as NADP-dependent 3-hydroxy acid

dehydrogenase. In step 5, 3-hydroxypropionate is converted into its corresponding CoA-

activated form by an acyl-CoA synthetase, again encoded by XF_RS09805. In this case,

BLAST analysis and Pfam domain architecture matched the reviewed UniProt entry A4YGR1

from *M. sedula*, which catalyzes this specific reaction. This enzyme, annotated as "acs" in

several strains, would exhibit substrate flexibility, being also responsible for acetate activation.

Step 6 involves the dehydration of 3-hydroxypropionyl-CoA to acryloyl-CoA, catalyzed by an

enoyl-CoA hydratase, operating in the reverse direction. Although the candidate gene

XF_RS04720 identified did not yield a strong BLAST match to experimentally characterized

acryloyl-CoA hydratases, it is annotated as an enoyl-CoA hydratase, consistent with the

proposed reaction. Furthermore, Pfam domain analysis revealed the presence of the

conserved ECH_1 domain, which is characteristic of enoyl-CoA hydratase family enzymes

and is found in all reference proteins associated with this activity. Step 7 corresponds to the

final reaction of the 3HP bi-cycle module, involving the conversion of acryloyl-CoA into

propionyl-CoA. This reaction is catalyzed by an NADPH-dependent acryloyl-CoA reductase.

Two candidate genes belonging to the same orthologous group were identified as potentially

responsible for this activity (XF_RS07485 and XF_RS10335), both annotated as NAD(P)-

dependent alcohol dehydrogenases. BLAST analysis showed that XF_RS07485 matched the reviewed UniProt entry A4YGN2 from *M. sedula*.

Propionyl-CoA is the metabolic junction between the module of part of 3HP bi-cycle and the putative second module of the acetate assimilation, namely, the methylcitrate cycle. Step 8 corresponds to the condensation of propionyl-CoA with oxaloacetate to form 2-methylcitrate. This reaction is classically catalyzed by methylcitrate synthase (PrpC). No canonical *prpC* gene was identified in the core genome. However, the gene XF_RS06495, which is conserved across all strains, annotated as citrate synthase (GltA), was identified as a potential candidate for this step. Given this structural homology and the absence of a dedicated *prpC* gene, GltA was considered the most parsimonious candidate to catalyze the condensation of propionyl-CoA with oxaloacetate in the reconstructed pathway. Such substrate flexibility has been reported for related enzymes within this family (for instance, *Bacillus subtilis*, UniProt entry P45858, as reported in BRENDA database). Step 9 corresponds to the conversion of 2-methylcitrate into 2-methylaconitate, which proceeds through two consecutive activities: 2-methylcitrate dehydratase (EC 4.2.1.117; AcnD) followed by 2-methylaconitate *cis-trans* isomerase (EC 5.3.3.7; PrpF). These transformations were included in the model as a combined step. The *acnD* locus (XF_RS05260) was annotated as a pseudogene in a subset of genomes (Temecula1, M23, 9a5c, U24D and Ann1). Among the strains used experimentally, only Temecula1 carried this pseudogene annotation, whereas the remaining experimental strains did not show this disruption. How this strain solves the absence of a functional AcnD remains unsolved, and it would be worth studying the substrate promiscuity of other aconitases. The gene *prpF* was identified in the core gene set of the pangenome (XF_RS05265). Step 10 corresponds to aconitase AcnB, present in all strains and encoded by XF_RS01240, and annotated as an aconitate hydratase 2 / 2-methylisocitrate dehydratase. Finally, step 11 corresponds to methylisocitrate lyase (PrpB), which belongs to the core gene set and is encoded by XF_RS05225, which produces pyruvate and succinate.

Compared with the flux distribution obtained when glutamine was used as the sole carbon source (Fig. 1C), in the flux distribution corresponding to acetate assimilation (Fig. 3; Supplementary Material S7) pyruvate is no longer converted into acetyl-CoA, as this metabolite is directly generated from acetate (Fig. 3, step 1). The pyruvate generated in the proposed pathway is redirected toward gluconeogenesis, similar to the glutamine-based simulation. Under acetate conditions, the malic enzyme also contributes to NADPH generation, although at 42% of the flux of the one observed with glutamine. This reduction may seem a paradox since the acetate assimilation pathway proposed here shows an increased demand of reducing power. A higher proportion of flux is directed through

oxaloacetate (95% vs 27.5% with glutamine). This increased TCA cycle flux implies a higher flux through isocitrate dehydrogenase (IDH). The Xfcore model contains both the NAD- and NADP-dependent IDH isozymes, and when the model was evaluated with pFBA under the same conditions, with acetate 76% of the flux through IDH was generating NADPH, whereas with glutamine as the only C source, all the flux went through the NAD-dependent IDH isoenzyme.

Polyamines production by *Xylella fastidiosa* strains

As previously described for other phytopathogens, such as *R. solanacearum* and *P. syringae* polyamine production could represent a potential virulence factor (Lowe-Power et al., 2018; Solmi et al., 2025) that has not yet been characterized in *X. fastidiosa*. In the Xfcore metabolic model, the biosynthesis pathways for the polyamines for putrescine, spermidine and spermine were well represented, including the S-adenosylmethionine (SAM) regeneration cycle required for the formation of spermidine and spermine biosynthesis (Fig. S6).

To evaluate the potential for polyamine overproduction, the presence of polyamine export systems in *X. fastidiosa* was evaluated. Polyamines are positively charged at physiological pH and therefore require active transport mechanisms to cross biological membranes. In *R. solanacearum*, putrescine secretion has been experimentally reported (Lowe-Power et al., 2018). However, no homologs of known polyamine exporters have been identified in its genome (Igarashi & Kashiwagi, 2010; Sugiyama et al., 2017), suggesting the existence of a yet uncharacterized transport system. In *X. fastidiosa*, no clear genomic evidence supporting a dedicated polyamine export system was identified. Therefore, instead of assuming extracellular exchange reactions, cytosolic demand reactions were introduced in the Xfcore model to simulate polyamine overproduction.

When polyamine demand reactions were enforced or used as the objective function (see Material and Methods), the model predicted feasible production of putrescine, spermidine, and spermine, beyond biomass requirements (Fig. 4A). However, increasing polyamine production resulted in a progressive reduction in biomass flux, revealing a trade-off between growth and polyamine production. The extent of this trade-off varied among the three polyamines, with differences in maximum achievable fluxes at minimum biomass levels. Under these conditions, the model predicted maximum fluxes of 7.09 mmol·gDW^{-1}·h^{-1} for putrescine, 1.57 mmol·gDW^{-1}·h^{-1} for spermidine, and 0.87 mmol·gDW^{-1}·h^{-1} for spermine, by individually maximizing each polyamine demand reaction.

When putrescine overproduction was constrained to zero, spermidine and spermine exhibited a linear inverse relationship, with limited flexibility concerning the production of biomass and combined polyamine (Fig. 4A). In contrast, when putrescine overproduction was allowed together with either spermidine or spermine, a broader feasible solution for similar biomass levels space was observed. In these cases, similar biomass levels could be obtained with different combinations of polyamine production, as reflected by the expansion of the biomass contour levels.

The experimental results on polyamine production by *X. fastidiosa* strains corroborated the findings of the Xfcore metabolic model (Fig 4B and Fig. S7; Supplementary Material S8). In a first experiment (Fig. S7) six *X. fastidiosa* strains (De Donno, ESVL, IVIA5770, Temecula1, XYL1961/18 and XYL468) produced and secreted polyamines in vitro. Putrescine, spermidine and spermine were detected in cell pellets from both biofilm and planktonic growth (CP), as well as in the culture supernatant (SP). Spermidine was produced at higher concentrations than putrescine and spermine.

In a second experiment (Fig. 4B), polyamine levels were analyzed in three distinct sample types: planktonic cell pellets (CP-P), biofilm-forming cell pellets (CP-B), and the culture supernatant (SP). Again, spermidine was the most abundant polyamine produced. Mean spermidine production across strains and time points ranged from 7.85 ± 8.74 to 38.00 ± 40.81 ng·mL^{-1} in CP-P and from 3.43 ± 2.51 to 442.67 ± 162.38 ng·mL^{-1} in CP-B. In SP, spermidine levels ranged from 0 to 180.32 ± 57.91 ng·mL^{-1}. In contrast, mean putrescine values ranged from 1.95 ± 0.32 to 2.85 ± 1.60 ng/ml in CP-P and from 1.67 ± 0.31 to 5.18 ± 1.01 ng·mL^{-1} in CP-B, while SP values ranged from 0.24 ± 0.27 to 5.65 ± 3.40 ng·mL^{-1}. Spermine production ranged from 2.04 ± 1.45 to 11.15 ± 14.39 ng·mL^{-1} in CP-P, from 2.7 ± 0.86 to 7.14 ± 2.83 ng·mL^{-1} in CP-B, and from 4.75 ± 0.91 to 22.41 ± 12.46 ng·mL^{-1} in SP.

There were no significant differences observed between time points within each sample type, except for the four comparisons marked with an asterisk in Fig. 4B. Nevertheless, CP-B samples showed an overall tendency for putrescine and spermidine to decrease between 5 and 12 days, a trend that was weaker and less consistent in CP-P. In the supernatant (SP), spermidine showed the clearest evidence of extracellular accumulation, whereas putrescine and spermine were also detected extracellularly but at lower levels. In some strains, supernatant polyamine levels showed a tendency to increase at 12 days compared to 5 days. Overall, these results indicate that spermidine is the most prevalent polyamine produced by *X. fastidiosa* strains, suggesting that putrescine is converted into spermidine, which is then further converted into spermine, but at a lower rate.

Trade-off between virulence factors and biomass

Previous reconstructions of *X. fastidiosa* metabolism (Gerlin et al., 2020; Oliveira et al., 2023) have considered the secretion of two known virulence factors: the LesA protein (Nascimento et al., 2016) and exopolysaccharides (EPS) (da Silva et al., 2001). The metabolic model of the phytopathogen *R. solanacearum* (Peyraud et al., 2016), also incorporated polyamine secretion due to its experimental confirmation.

To account for these virulence-associated processes, the Xfcore metabolic model incorporated the reactions required for LesA biosynthesis. The gene responsible for LesA production (XF_RS01515 or XF_RS01510) was present in the core gene set, and the ATP-dependent transporter required for its export was also part of the core pangenome, with the exception of one gene involved in the secretion cluster of strain J1A12 (XF_RS06440, *gspK*). However, as described in other cases, the annotation of this gene as a pseudogene was due to a deletion of a G at position 786 within a repetitive G-rich region.

For EPS biosynthesis, the *gum* operon responsible for EPS synthesis and export was found in the core pangenome, with the exception of the *gumF* gene, which encodes an acetyltransferase responsible for the final step of the biosynthetic pathway. This gene was annotated as a pseudogene in strains U24D and Salento 2; however, since the rest of the operon was present and the pseudogene annotation results from a single point mutation that could represent a sequencing error, EPS production was incorporated into the Xfcore model.

In this work, several exploration scenarios were analyzed by simulating the combined production of LesA protein, exopolysaccharides (EPS), and one of the polyamines: putrescine (Fig. 5A), spermidine (Fig. 5B), or spermine (Fig. 5C). Under these conditions, the model predicted maximum fluxes of 0.02 mmol·gDW^{-1}·h^{-1} for LesA protein, 1.45 mmol·gDW^{-1}·h^{-1} for EPS, while the maximum achievable fluxes for putrescine, spermidine and spermine correspond to those previously described.

In these simulations, a clear trade-off between virulence factor secretion and biomass production was observed (Fig. 5). The combined production of LesA with EPS showed a linear inverse relationship. For the production of putrescine together with the known virulence factors (Fig. 5A), a nearly linear inverse relationship was observed between the combined production of these compounds and biomass formation. Interestingly, in the case of spermidine (Fig. 5B) and spermine (Fig. 5C), the feasible solution space showed greater compatibility between LesA production and polyamine secretion. In these scenarios, different rates of LesA

production remain feasible even when spermidine or spermine are produced at high levels,
indicating greater metabolic flexibility for LesA secretion while maintaining the same biomass
flux. This behavior is reflected by the broader feasible region observed in the phase planes.
In particular, a change in slope appears when LesA production reaches approximately 0.02
mmol·gDW⁻¹·h⁻¹. A similar but less pronounced pattern is observed for EPS production.
Overall, EPS secretion appears to be more compatible with spermidine or spermine production
than with putrescine production, particularly near the upper limit of EPS flux (around 1.3
mmol·gDW⁻¹·h⁻¹).

Discussion

In this work, we reconstructed the core metabolic model of *X. fastidiosa* based on the
pangenome of 18 strains, representing five described subspecies of the phytopathogen. This
approach accounts for the high genetic diversity within the species while allowing the
identification of conserved metabolic capabilities to gain insight into the common traits of this
bacterium. Previous metabolic reconstructions of *X. fastidiosa* were strain-specific, including
the model for *X. fastidiosa* subsp. *multiplex* CFBP 8418 (Gerlin et al., 2020) and the model for
X. fastidiosa subsp. *pauca* De Donno (Oliveira et al., 2023). In the first study, genome
comparisons were performed with the strains *X. fastidiosa* subsp. *pauca* 9a5c and subsp.
fastidiosa Temecula1, suggesting that the results could be extrapolated to these subspecies.
In contrast, our reconstruction explicitly incorporates these strains within a broader
pangenome framework, providing a more robust representation of the conserved metabolic
potential of *X. fastidiosa*.

Moreover, the Xfcore metabolic model and the pangenome assembly can serve as a
foundation for the reconstruction of strain-specific models, following the protocol described in
Norsigian et al. (2020). This framework highlights how the availability of a curated base model
can significantly reduce the effort required for the generation of multi-strain metabolic models.
Such approaches have already been successfully applied to reconstruct strain-specific models
for the *Lactobacillaceae* family and for *Aspergillus fumigatus* species (Mirhakkak et al., 2023;
Ardalani et al., 2024). Additionally, our reconstructed model incorporates the ortholog group
associated with each gene in the network annotation, enabling the extraction of strain-specific
genetic information when required.

The Xfcore metabolic model contains 42 genes annotated as pseudogenes in some of the

strains. However, several cases suggest that these annotations may not reflect true gene loss

but sequencing mistakes, as some genes appear intact in alternative genome assemblies of

the same strain or contain frameshift mutations in homopolymeric regions prone to sequencing

errors (Cooley & Wright, 2024). Moreover, some of these genes belong to otherwise complete

metabolic pathways, making the loss of a single intermediate reaction metabolically unlikely.

For these reasons, pseudogene annotations were carefully evaluated during model curation.

Some peculiarities were identified when analyzing the characteristics of the metabolic network.

For instance, early studies on *X. fastidiosa* metabolism reported the absence of fructose-1,6-

bisphosphatase (FBPase) activity and suggested that gluconeogenesis could not occur in this

bacterium (Facincani et al., 2003). Consistent with these observations, no FBPase was found

in the pangenome. However, gluconeogenesis could be supported by a reversible

pyrophosphate-dependent phosphofructokinase (EC 2.7.1.90), which is encoded in the core

genome.

In addition, the bacterium appears to have a limited set of reactions for reducing power

production, as both 6-phosphogluconate dehydrogenase and transhydrogenase are absent

from the Xfcore network. In minimal medium with glutamine, pyruvate can be generated

through the malic enzyme, which contributes to NADPH production and represents the main

connection between the TCA cycle and gluconeogenesis. We did not find any evidence for

reactions directly linking pyruvate or phosphoenolpyruvate with the TCA cycle. Therefore, *X.*

fastidiosa appears to lack typical anaplerotic reactions to replenish the cycle, suggesting that

its metabolism operates under relatively constrained network configurations, fully dependent

of the environmental supply of precursors of the cycle intermediates, such as the amino acids

glutamine or glutamate present in the xylem. Consistent with this observation, simulations in

the Xfcore model indicate that growth on hexoses or pentoses, or acetate, requires the

presence of a TCA cycle intermediate to initiate the cycle. In the absence of such

intermediates, the model cannot sustain growth, reflecting that limited capacity to replenish

TCA metabolites. In previous reconstructions, different assumptions allowed growth under

these conditions: in the model by Gerlin et al. (2020), growth was supported by the baseless

inclusion of the glyoxylate cycle, whereas in the reconstruction by Oliveira et al. (2023) growth

was enabled by allowing the malic enzyme to operate reversibly, effectively generating malate

from   pyruvate.   When   the   malic   enzyme   is   constrained   to   operate   in   the

oxidative/decarboxylative direction, growth decreases and requires the presence of a TCA

intermediate.

The phenotypic characterization performed in this work provides a robust dataset, including four independent experiments and five strains belonging to the three main *X. fastidiosa* subspecies (*fastidiosa*, *multiplex* and *pauca*). Previous studies have reported Biolog results for *X. fastidiosa*, such as the work of Gerlin et al. (2020), although in that case the analysis was limited to the strain CFBP8418 (subsp. *multiplex*) used for metabolic modelling. For sugars, the in silico simulations indicated that the Entner–Doudoroff (ED) pathway represents the main glycolytic route through which the metabolic network can metabolize these compounds. This result is consistent with the model proposed by Oliveira et al. (2023), which reached the same conclusion and suggested that this could help protect the bacterium against reactive oxygen species (ROS) generated by the plant defense system (Wang et al., 2017). As previously reported in *P. putida*, the metabolism through the ED pathway has been associated with increased tolerance to oxidative stress due to enhanced NADPH generation (Chavarría et al., 2013).

In axenic cultures, *X. fastidiosa* can grow with acetate as a sole carbon source. Remarkably, acetate is a major organic acid present in olive xylem sap (Anguita-Maeso et al., 2021). Intriguingly, GEMs for this organism lack any canonical pathway for acetate assimilation, and previous metabolic reconstructions have addressed this issue using different strategies. In the Xfm1158 model (Gerlin et al., 2020), growth on acetate was enabled by incorporating the glyoxylate cycle, despite the absence of genomic evidence supporting the presence of its key enzymes. In contrast, the iMS50 model (Oliveira et al., 2023), adopted a more conservative position and did not include the glyoxylate cycle, resulting in the inability to simulate growth on acetate. In our reconstruction, the Xfcore model suggested a previously undescribed metabolic solution for growth on acetate. In brief, we propose the first part of the 3HP bi-cycle –an autotrophic pathway described in *Chloroflexus auranticus* by Herter et al. (2002) (Supplementary Figure S8, panel A)–combined with the conversion of propionyl CoA into a biomass precursor. The assimilation of propionyl CoA in *X. fastidiosa* would use a pathway different to the one proposed by Zarzycki and Fuchs (2011) for the co-assimilation of acetate in *C. auranticus* (Supplementary Figure S8, panel B). Thus, in the Xfcore model propionyl CoA can be assimilated into pyruvate through the methylcitrate pathway firstly described in yeasts by Tabuchi et al. (1974) and which is also operative in bacteria like *E. coli* grown in propionate (Brock et al., 2002) (Supplementary Figure S8, panel C).

We can compare this new pathway for acetate assimilation with the ones previously described in the literature (Supplementary Table S6). The putative acetate assimilation pathway in *X. fastidiosa* (Xf pathway) represents a slight variation of the 3HP pathway described in *C. auranticus*, in terms of energy balance and carbon efficiency; in both cases, there is a net gain

of carbon. The main difference is that the Xf pathway involves two more enzymatic steps than

the half 3HP bi-cycle. Both pathways are more demanding of reducing power than the rest of

known routes and, as discussed before, this may explain the increased flux through the TCA

cycle in Xfcore under acetate conditions. It is remarkable that *X. fastidiosa* harbors two

structurally different IDH isoenzymes, specific for NAD+ or NADP+, respectively (Lv et al.,

2018). Interestingly, the evolutionary origin of a NADP-dependent IDH has been correlated

with an increased demand of reducing power associated with the use of acetate as carbon

source (Zhu et al., 2005). Comparing all the pathways, the shortest solutions are represented

by the anaerobic assimilation of acetate by ferredoxin-dependent pyruvate synthase and the

widely distributed glyoxylate cycle (Supplementary Table S6).

Our proposal of a combination of part of the 3HP bi-cycle with the methylcitrate pathway to

explain acetate assimilation by *X. fastidiosa* highlights the role of functional modularity in

metabolic adaptation and evolution. Metabolic modules can reorganize or combine to explore

biochemical diversity (Schada von Borzyskowski et al., 2020). Functional degeneracy

between diverse metabolic solutions also can reflect different dynamics of adaptation to

environmental challenges, as described for *Paracoccus denitrificans* (Kremer et al., 2019). On

the other hand, in our hypothetical solution for acetate assimilation in *X. fastidiosa* we have

proposed several steps catalyzed by enzymes with relaxed substrate specificity. Thus,

referring to the pathway in Fig. 3, an acyl-CoA synthetase operates in steps 1 and 5, citrate

synthase is also operative as methylcitrate synthase (step 8), and finally aconitase AcnD will

act in step 10. Those cases would represent additional examples of the well known metabolic

economy observed in symbionts (either pathogens or mutualists) and sustained through

reactions operated by promiscuous or multifunctional enzymes (Yus et al., 2009; Kim et al.,

2010; Gil & Peretó, 2015). Gerlin et al. (2020) have correlated the lack of enzymatic

redundancy in the *X. fastidiosa* metabolic network with a reduced robustness and the

characteristic fastidious growth. In any case, we are aware that our in silico hypothesis

requires experimental validation, through omics approaches, biochemical characterization of

enzyme activities, and ^{13}C-fluxomics experiments. Furthermore, phylogenetic analysis of the

involved enzymes also could shed light on the origin and evolutionary tinkering of the pathway,

as well as its distribution in the tree of life.

Our results revealed some discrepancies between in vitro phenotyping, published

observations, and model predictions. Consistent growth on glutamine or glutamate was not

detected in the Biolog PM assays for the strains analyzed, whereas growth was sustained in

silico. However, glutamine was selected as the main carbon source for the minimal media

experiments and simulations because its utilization has been widely reported in the literature

for *X. fastidiosa* (Almeida et al., 2004; de Macedo Lemos et al., 2003; Gerlin et al., 2020). In addition, in the minimal media experiments performed in this work, glutamine sustained both bacterial growth and biofilm formation. As per the same reason, glutamic acid was used in the phenotypic simulations because it provided a direct entry point into the TCA cycle, representing a known nitrogen source present in xylem sap (Surano et al., 2024). This glutamate supplementation was consistent with the medium composition used in the Biolog phenotypic assays (available via manufacturer). Moreover, its assimilation is consistent with the predictions of the Xfcore model, which requires substrates capable of replenishing TCA cycle intermediates.

Some carbohydrates and derivatives, as well as some organic acids, supported growth in vitro but not in silico, suggesting that certain metabolites and metabolic reactions may be missing from the model, possibly because the corresponding genes are not present in all strains of the pangenome. In the case of lipid substrates, in silico validation could not be performed because the β-oxidation pathway has not been described in *X. fastidiosa*. However, growth on lipid derivatives has been previously reported (Gerlin et al., 2020), suggesting that some alternative, yet undescribed, metabolic pathways may exist. As shown for acetate assimilation, the metabolic model may help guide the search for such alternative pathways by identifying potential metabolic gaps.

Finally, several compounds commonly present in xylem sap were also evaluated. Although no growth was detected in vitro for some of them (D-glucose, citrate, L-aspartate and L-alanine), the metabolic model predicted that these substrates could theoretically support growth. These discrepancies may be explained by differences in experimental conditions. In this work, growth was evaluated using OD_{600} measurements, whereas Biolog PM assays typically rely on respiration-based detection. Additionally, factors such as substrate concentration, medium composition, or transport limitations may influence the ability of cells to utilize certain compounds in vitro. For instance, a low initial cell density could limit the activity of the glutamate transporter, preventing the flux required to sustain growth, unlike other carbon sources such as pyruvate. Moreover, in xylem sap *X. fastidiosa* is likely exposed to combinations of carbon sources, which could compensate for metabolic limitations that are not captured when substrates are tested individually in Biolog PM assays.

Metabolic reconstructions have been previously used to design minimal growth media for fastidious microorganisms. A notable example is the work for *Mycoplasma pneumoniae* to define a synthetic minimal medium (Yus et al., 2009). This strategy of reverse ecology implicitly validates that genomic information and reconstructed networks can guide the

formulation of culture media for other fastidious or uncultured bacteria (Cross et al., 2019;
Lewis et al., 2021). Under strictly defined conditions, the use of sugars or other substrates that
do not directly feed the TCA cycle would theoretically require the addition of a TCA
intermediate. However, in empirical in vitro conditions, intracellular metabolite pools may
partially compensate for this requirement, as cells are not metabolically empty at the beginning
of the experiment. In natural environments, on the other hand, the chemical diversity
accessible to the bacterium would compensate for this metabolic obstacle. Interestingly,
although none of the minimal medium tested here supported higher growth, with few
exceptions, than the control medium, biofilm formation was observed at comparable levels
under these restricted conditions. *X. fastidiosa* is known to colonize xylem vessels primarily in
the form of biofilms, which play a key role in its pathogenicity (Marques et al., 2002). This
approach aimed to explore whether minimal environmental conditions, such as in the xylem
sap, could trigger biofilm development independently of optimal planktonic growth, which was
confirmed with the results shown.

Another important trait explored in this work was polyamine production in *X. fastidiosa*.
Polyamines are well-established modulators of biofilm formation and adhesion in many
bacterial pathogens. They influence biofilm development through effects on gene regulation,
signaling pathways, and extracellular matrix production. However, there was currently no
direct experimental evidence that polyamines are produced by *X. fastidiosa*, modulate its
biofilm formation or adhesion. A previous study analyzing nitrogen metabolism in citrus trees
infected with *X. fastidiosa* reported high levels of putrescine exclusively in infected plants
(Purcino et al., 2007), although this observation was interpreted as associated with plant
production, as polyamines are also known to be produced by plants as part of their stress
signalling responses (Blázquez, 2024). In other phytopathogens, such as *R. solanacearum*
and *P. syringae* polyamine production has been described as a virulence factor (Lowe-Power
et al., 2018; Solmi et al., 2025), but this trait has not yet been characterized in *X. fastidiosa*. In
these pathogens, polyamines have been proposed to protect bacteria against oxidative stress
generated by plant defense responses and may contribute to symptom development during
infection.

Metabolic modelling has previously been used to explore production of virulence-associated
processes in plant pathogens. Other reconstructions of *X. fastidiosa* metabolism (Gerlin et al.,
2020; Oliveira et al., 2023) have considered the secretion of two known virulence factors: the
LesA protein (Nascimento et al., 2016) and exopolysaccharides (EPS) (da Silva et al., 2001).
The metabolic model of the phytopathogen *R. solanacearum* (Peyraud et al., 2016), also
incorporated polyamine secretion due to its experimental confirmation. We extended this

concept by exploring the relationship between polyamine production and the virulence-
associated factors in *X. fastidiosa*, including the secretion of the LesA protein and
exopolysaccharides (EPS).

In this work, both metabolic modelling and experimental analyses indicated that *X. fastidiosa*
is capable of producing and secreting polyamines. Metabolic simulations revealed a clear
trade-off between biomass production, polyamine biosynthesis, and other known virulence
factors, suggesting that the synthesis of these compounds may occur at a metabolic cost for
the bacterium. Differences in the maximum achievable fluxes for the three polyamines at
minimum biomass levels reflect their distinct biosynthetic costs and precursor requirements
within the metabolic network. In particular, spermidine and spermine production appear to be
metabolically coupled due to their dependence on shared precursors. Additionally, the
simulations revealed a clear trade-off between biomass production and the secretion of
virulence-related compounds. Interestingly, spermidine and spermine production appeared to
be more compatible with the secretion of LesA and EPS across a broader range of similar
biomass levels.

Our experimental data, despite the variability typically observed when working with fastidious
bacteria in vitro, showed a higher accumulation of spermidine in the culture supernatant.
Additionally, polyamine levels tended to be higher in biofilm-associated cell pellets (CP-B) than
in planktonic cells, suggesting a link between polyamine production and the biofilm state of
this phytopathogen. While polyamines are established regulators of bacterial biofilms–often
acting through EPS synthesis or second messenger pathways (Karatan & Michael, 2013)–
their role in *X. fastidiosa* remains untested experimentally. All current works for *X. fastidiosa*
focus on EPS composition or processing (Janissen et al., 2015; Castro et al., 2023), the role
of surface adhesins and pili (Scala et al., 2025), or of DSF-family quorum sensing signals
(Horgan et al., 2024), without an explicit link to polyamine metabolism or signaling. Our results
open the possibility that polyamine production could represent an additional virulence-related
trait in *X. fastidiosa*. Further in vivo studies will be necessary to determine whether the
polyamines detected in infected plants originate from bacterial metabolism and to clarify their
role during host colonization by this species.

Overall, the Xfcore metabolic model provides a systems-level framework to investigate the
metabolism of *X. fastidiosa* and to generate new hypotheses about its physiology and
virulence strategies. In future work, this reconstruction could be integrated with plant metabolic
models to explore host–pathogen interactions within the xylem environment.

Conclusions

This work presents the first pangenome-based metabolic reconstruction for *X. fastidiosa*, integrating the conserved metabolism across five subspecies of this phytopathogen. After experimental validation, the Xfcore model enabled the prediction of minimal nutritional requirements, which guided the formulation of defined media capable of supporting biofilm formation in vitro. This success served as further implicit validation of the predictive power of the model. Furthermore, the model suggested a previously undescribed metabolic solution for acetate assimilation into biomass precursors. The simulations also predicted the capacity of *X. fastidiosa* to overproduce polyamines, a result that was confirmed experimentally, representing the first in vitro evidence of polyamine production in this pathogen. Beyond these insights, the Xfcore model provides a robust framework for the reconstruction of strain-specific metabolic models and offers a valuable platform for future studies integrating bacterial and plant metabolism to investigate host–pathogen interactions.

Materials and Methods

Genome data preparation and pangenome reconstruction

A total of 18 *X. fastidiosa* strains were considered for the pangenome assembly (Table 1). Complete, chromosomal, whole-genome data was retrieved from the NCBI Genome database (Sayers et al., 2024). For each strain, the associated plasmid sequences, when available, were also inspected using the BlastKOALA annotation server (Kanehisa et al., 2016) and the NCBI annotation. As none of the plasmid-encoded genes were associated with relevant metabolic functions, plasmid sequences were excluded from the pangenome analysis. Coding sequences, excluding those annotated as pseudogenes, were clustered into ortholog groups (OGs) using InParanoid (Remm et al., 2001), applying a minimum of 80% sequence identity and at least 50% pairwise aligned sequence length, using the BLOSUM 80 substitution matrix. Pairwise ortholog clusters generated by InParanoid were merged into multi-strain ortholog groups using QuickParanoid (http://pl.postech.ac.kr/QuickParanoid/). Genes without a best bidirectional hit in any other genome were classified as "unique" while "accessory" genes were defined as those present at least in two strains. Genes belonging to OGs shared across all strains were considered "core". Core, accessory and unique gene sets were annotated with BlastKOALA automatic annotated server to explore the functional traits.

Metabolic model reconstruction and refinement

For the constraint-based metabolic model reconstruction and analysis (COBRA), established protocols were followed (Thiele & Palsson, 2010; Nogales, 2017). An initial draft reconstruction of the metabolic network was generated based on the core gene set of the *X. fastidiosa* 9a5c strain using the automatic workflow provided by ModelSEED (Overbeek et al., 2005; Henry et al., 2010). In addition, draft reconstructions were generated for all remaining strains to support the manual curation of the core draft model. Curated genome-scale metabolic models of *Escherichia coli* iJO1366 (Orth et al., 2011) obtained from BiGG Models (King et al., 2016), *Xylella fastidiosa* Xfm1158 (Gerlin et al., 2020) and iMS508 (Oliveira et al., 2023) from BioModels (Malik-Sheriff et al., 2020), and *Xanthomonas campestris* pv. *campestris* str. B100 (Schatschneider et al., 2013) were used as references during the curation process.

Subsequently, the model was iteratively refined to ensure consistency with available physiological and biochemical data. To facilitate thorough curation of the draft reconstruction, reactions, metabolites, and genes were annotated using identifiers from multiple biochemical databases, including KEGG (Kanehisa et al., 2025), MetaCyc (Caspi et al., 2020), BioCyc (Karp et al., 2019) BiGG Models (King et al., 2016), ModelSEED (Overbeek et al., 2005; Henry et al., 2010), and NCBI databases. SBOannotator (Leonidou et al., 2023) was used to assign SBO terms. BioCyc (Karp et al., 2019), KEGG (Kanehisa et al., 2025), BRENDA (Chang et al., 2021) and Pfam databases (Mistry et al., 2021) were consulted to support manual curation. Consistency analysis of the topological network was conducted to identify gap metabolites and blocked reactions (Ponce-de-Leon et al., 2015). During this step, reaction stoichiometries were revised, including mass and charge balances, reaction directionality and cofactor usage. Similarity-based searches were used to infer missing functions in the model to sequences from the *X. fastidiosa* pangenome, based on sequence and functional homology with other organisms. These analyses were performed using BLASTp (Camacho et al., 2009) and HMMER (Mistry et al., 2013) toolboxes against local sequence databases built from the *X. fastidiosa* pangenome whole-genome sequences. Within the HMMER framework, hmmscan and hmmsearch and pfamscan (Madeira et al., 2024) were used to identify conserved protein domains and domain architectures, providing additional support for functional inference based on profile hidden Markov models. Although all pseudogenes were initially excluded from the pangenome analysis–defined as genes predicted to have lost functionality due to frameshift mutations, partial or incomplete sequences, internal codon stops or a high proportion of ambiguous nucleotides–this criterion was later revised in order to avoid overlooking any essential metabolic activity, during the manual model curation. To that end, every missing function was also searched within the accessory gene sets. Accessory functions that were

found in the majority of strains and pseudogenized only in a limited number of strains, typically due to point mutations visualized with the MEGAX software (Kumar et al., 2018), were treated as core functions to better represent the general metabolic phenotype of the species (Table S1). Reactions were assigned a confidence score ranging from 0 to 4 according to the level of supporting evidence, following the criteria defined by Thiele and Palsson (2010). Curation efforts focused on pathways central to the conserved physiology of *X. fastidiosa*, including central carbon metabolism, complete biosynthetic routes for amino acids, lipid and nucleotide metabolism, energy metabolism, and selected transport systems (Simpson et al., 2000; Meidanis et al., 2002). In addition, pathways related to redox balance, cofactor biosynthesis and host-associated growth were carefully curated. These included quinone and shikimate biosynthesis, polyamine metabolism, the S-adenosylmethionine (SAM) cycle, sulfur and nitrogen metabolism, and the methionine salvage pathway. Vitamin and cofactor biosynthetic pathways were curated in detail, encompassing pantothenate, folate, riboflavin, thiamine, and porphyrin metabolism. Finally, pathways associated with cell envelope biogenesis and host interaction were refined. This included the biosynthesis of exopolysaccharides (EPS) and lipopolysaccharides (LPS), as well as other fatty acid metabolism. Furthermore, the biosynthesis and export reactions of the LesA protein were incorporated to the model, following previous strain-specific metabolic reconstructions (Gerlin et al., 2020; Oliveira et al., 2023).

In parallel with the rest of the network, the biomass reaction was also curated. The biomass reaction generated by ModelSEED was used as a template. Comparison of the biomass reactions of the reference models and the pangenome core one was done. The suitability of each metabolite in the template and potential biomass components were evaluated, including the molar contribution of each one, based on biochemical data and the biomass reactions from the reference models.

Manipulation of metabolic reconstructions was carried out with the COBRApy library in Python (Ebrahim et al., 2013). NetworkX v2.4 (Hagberg et al., 2008), Escher v1.7.3 (King et al., 2015), yED Graph Editor v3.18.2 (yWorks GmbH, 2020) and Cytoscape v3.7.1 (Shannon et al., 2003) were used for metabolic network visualization during the curation process. The quality of the final model was evaluated using the MEMOTE v.0.13.0 tool (Lieven et al., 2020). The final GEM is provided in JSON, MAT-File and Systems Biology Markup Language (SBML) formats (Supplementary Material S1).

In silico constraints and growth conditions

For reversible reactions, upper and lower bounds of 1000 and -1000 mmol·gDW^{-1}·h^{-1}, were respectively applied. For irreversible reactions, the lower bound–or in specific cases, the upper bound–was fixed at 0 mmol·gDW^{-1}·h^{-1}. Maximization of growth rate was set as the objective function. Oxygen exchange was left unconstrained. The growth-associated maintenance energy (GAM) was fixed at 50.12 mmol·gDW^{-1}·h^{-1}, while the non-growth-associated maintenance energy (NGAM) was set to 3.073 mmol·gDW^{-1}·h^{-1}, according to Gerlin et al. (2020).

For simulation of growth conditions, the lower bounds of exchange reactions were set to zero except for the inorganic ions included in the predicted minimal medium.

For minimal medium simulations, glutamine uptake flux was constrained to -7.25 mmol gDW^{-1} h^{-1} as the carbon and nitrogen source, based on environmental, biochemical and previous metabolic modeling data (Davis, 1992; Gerlin et al., 2020; Orth et al., 2011). For trade-off simulations, glutamine uptake was fixed at -15.665 mmol gDW^{-1} h^{-1} (see *Polyamine and virulence factor trade-off Prediction section* in Materials and Methods).

To evaluate phenotypic substrate utilization data in silico, including acetate growth conditions, carbon substrate uptakes were normalized by the number of carbon per molecule, ensuring an equivalent maximum carbon availability across metabolites. A carbon uptake rate of 100 C-mmol·gDW^{-1}·h^{-1} was imposed, following the methodology of Gerlin and coworkers (2020). Ammonia uptake was left unconstrained as the nitrogen source. Additionally, L-glutamate uptake was set to -6 mmol gDW^{-1} h^{-1} to evaluate differences in growth between substrates without limitations imposed by ammonium uptake and anaplerotic reactions of the network.

The complete set of exchange flux constraints in each simulation is provided in Table S5.

Flux Balance and Flux Variability Analyses

Flux Balance Analysis (FBA) and parsimonious Flux Balance Analysis (pFBA) were performed using the COBRApy software (Ebrahim et al., 2013). The optimization problems were solved with the GNU Linear Programming Kit (GLPK) v.5.0 or the IBM ILOG CPLEX Optimizer v.22.1.1. FBA and pFBA were used to compute the maximum value of the objective function and the corresponding flux distribution (Orth et al., 2010; Thiele & Palsson, 2010). Maximization of the growth rate was set as the default cellular objective function. For modelling

949 and simulating growth specific conditions, bounds of some reactions were modified as
950 previously specified in the *In silico constraints and growth conditions* section.

Design of synthetic defined minimal media

To design a model-driven minimal medium sustaining the FBA-predicted growth rate under a
fixed L-glutamine uptake constraint (7.25 mmol·gDW^{-1}·h^{-1}), the number of active exchange
reactions was minimized. This was formulated as a mixed-integer linear programming (MILP)
problem and solved using the GLPK solver (Free Software Foundation, 2018), employing the
minimal_medium implementation from the COBRApy medium module (Ebrahim et al., 2013).
Based on the predicted metabolite requirements, six synthetic defined minimal media were
designed (mXC1-mXC6, Table S3). The initial formulations (mXC1 and mXC2) were derived
from the in silico minimal medium, using glutamine as carbon and nitrogen source, and
different iron sources. Concentrations of all components were adapted from existing media
known to support *X. fastidiosa* growth. Concentrations of glutamine, hemin chloride and
inorganic salts were adapted from PW synthetic medium (Davis et al., 1981), while ferric
pyrophosphate concentrations were derived from BCYE (Wells et al., 1981) and 3G10R media
(Leite et al., 2004). For subsequent formulations (mXC3 to mXC6), glutamine was replaced
with equivalent quantities of alternative carbon sources (sodium acetate or ribose),
supplementing with nitrogen sources. Generally, ammonium sulfate was used as an inorganic
nitrogen source, whereas cysteine was used as an alternative nitrogen source in mXC6
formulation. To validate the biotin-related predictions under growth on acetate, mXC4 medium
was supplemented with biotin. Concentration of $(NH_4)_2SO_4$ was obtained from XVM2
(*Xanthomonas* apoplast medium) medium (Hiery et al., 2013), cysteine from BCYE and XF-
26 (Chang, 1993), and biotin from SXM (Xylem fluid medium) (Neumann, 2003). All media
were adjusted to a pH of 6.9 and brought to a final volume of 1 L with sterile distilled water.
PD3 broth (Davis et al., 1981) was used as a positive growth control.

Acetate assimilation pathways research

To investigate the metabolic strategy employed by *X. fastidiosa* for the conversion of acetate
into biomass, the diverse metabolic assimilation pathways described in the literature were
searched against the *X. fastidiosa* genome. Reference protein sequences corresponding to
the enzymatic steps involved in each pathway were retrieved from the BRENDA database
based on their associated EC numbers. Reciprocal BLASTP (v2.9.0+) (Camacho et al., 2009)

searches were conducted between these reference sequences and the annotated amino acid

sequences of *X. fastidiosa*. Substantial outcomes were obtained by setting specific filtering

parameters (coverage greater than 80 %, e-value ≤ 1e−6). Additionally, hmmscan, hmmsearch

and pfamscan (Madeira et al., 2024) were used to identify conserved protein domains and

domain architectures, providing additional support for functional inference. Finally, the results

were manually revised and some selected candidates and enzymatic activities were reviewed

using the UniProt database (UniProt Consortium, 2025) to validate results.

Polyamine and virulence factors trade-off prediction

Polyamine biosynthesis and degradation pathways were manually curated and revised as

described in the metabolic reconstruction section. Special attention was paid to S-

adenosylmethionine (SAM) regeneration and the L-methionine salvage pathway. As no

experimentally validated polyamine export mechanism has been described in *X. fastidiosa*,

demand reactions for putrescine, spermidine, and spermine were incorporated to simulate

intracellular production and secretion potential.

To investigate potential trade-offs between biomass production, polyamine secretion and

virulence factors synthesis (EPS and LesA protein), a two-step optimization strategy was

applied. First, an experimental-like scenario was reproduced constraining biomass production

(0.1608 h^{-1}), EPS biosynthesis (1.296 mmol gDW^{-1} h^{-1}) and LesA production (0.0024 mmol

gDW^{-1} h^{-1}) as described in Gerlin et al. (2020). Under these constraints, glutamine uptake was

minimized, resulting in a flux of -15.665 mmol gDW^{-1} h^{-1}. This value was used as the carbon

and nitrogen source uptake for all trade-off simulations. In the second step, biomass was

constrained to a minimum of 0.1608 h^{-1}, and the feasible production space of three target

metabolites was explored. For each metabolic product, the maximum achievable flux at the

minimum fixed biomass level was determined by FBA by maximizing the corresponding

demand or export reaction. Uniform grids (30 intervals per axis) ranging from zero to the

maximum achievable flux were generated for each metabolite. For every combination of target

fluxes (p1, p2, p3), biomass was constrained to a minimum value of 0.1608 h^{-1} and the model

was re-optimized maximizing the growth rate as the objective function using the CPLEX

Optimizer. Only feasible solutions were retained. This sampling enabled the reconstruction of

two-dimensional phase planes approximating the surface of the Pareto front, representing the

trade-off between metabolite production and biomass growth rate. Phase planes were

obtained by fixing one product to zero and projecting the feasible region of the remaining two

metabolites. The resulting solution space was visualized using tricontour and tricontourf functions from the Matplotlib library. Biomass flux was represented as a continuous color gradient.

Bacterial strains and culture conditions

Seven *X. fastidiosa* strains used in the in vitro studies including two from subspecies *fastidiosa* isolated from grapevine in California (Temecula1) and Balearic island (IVIA5770), three from subspecies *multiplex* isolated from almond trees in Alicante (Spain) (ESVL and IVIA5901) and wild olive in Balearic island (XYL468) and two from subspecies *pauca* isolated from olive trees (De Donno and XYL1961/18) in Apulia (Italy) and Ibiza (Spain), respectively. Stocks of *X. fastidiosa* strains were stored in phosphate buffered saline (PBS) plus 50 % glycerol at -80 °C. For all the experiments, strains were recovered from glycerol stocks on plates of PD3 (Davis et al., 1981) or BCYE (Wells et al., 1981) media incubated at 28 °C in the dark. After 6 to 8 days, cells were re-streaked onto new plates and cultured for 5 to 7 days before using them in each experiment.

Phenotypic characterization of *Xylella fastidiosa* strains

Phenotypic characterization of five *X. fastidiosa* strains (ESVL, De Donno, IVIA5901, Temecula1, and XYL1961/18) was performed using the Biolog phenotype microarray (PM) system (Biolog, Hayward, California). The potential utilization of 190 carbon sources was determined using the PM1 and PM2 plates. For PM plate inoculation, bacterial cells were scraped from solid culture plates and resuspended in the IF-0a GN/GP base inoculating fluid (Biolog, Hayward, California) and the bacterial suspension was mixed with the corresponding additive solutions for each PM plate and the Dye Mix G, to reach a final optical density at 600 nm (OD_{600}) of 0.08 according to the manufacturer's protocol. Subsequently, 100 µL of a cell suspension of each *X. fastidiosa* strain was added to each well of the PM microplates and incubated at 28 °C during 9 days without agitation. Bacterial growth was assessed by measuring OD_{600} using a microplate absorbance reader i-Mark TM spectrophotometer (Bio-Rad, Hercules, CA, USA). Four independent experiments were conducted for each strain and PM substrate-type microplate.

Growth of *Xylella fastidiosa* strains in designed synthetic defined minimal media

In vitro culture experiments using the six minimal media (Table S3) were conducted with seven *X. fastidiosa* strains (ESVL, De Donno, IVIA5770, IVIA5901, Temecula1, XYL1961/18, and XYL468) as described by Román-Écija and coworkers (2023) to assess the ability of the designed minimal media to support total and planktonic growth, as well as biofilm formation of the seven *X. fastidiosa* strains. Briefly, a cell suspension of each strain was diluted in each minimal medium to reach at OD_{600} = 0.08. Cell suspensions (200 μL) were inoculated in polystyrene 96-well microplates (Sarstedt, Nümbrecht, Germany) and incubated at 28 °C during 7 days without agitation. At least four independent experiments were conducted, and each experiment included 5 replicated wells per treatment. Non-inoculated medium served as negative control. Bacterial growth was measured daily by measuring OD_{600}. After the 7-day incubation, planktonic growth was assessed by transferring 150 μL cell suspension to a new plate and measuring the OD_{600}. Biofilm formation was estimated using the crystal violet assay as described (Román-Écija et al. 2023).

Polyamines production by *Xylella fastidiosa* strains in liquid medium

To validate the metabolic model predictions for the ability of *X. fastidiosa* to produce polyamines (putrescine, spermidine and spermine) during in vitro culture two experiments were performed. An initial experiment was carried out to determine the production of polyamines by *X. fastidiosa* during *in vitro* culture. For that, 10-mL cell suspensions at OD_{600} = 0.05 of six *X. fastidiosa* strains (ESVL, De Donno, IVIA5770, Temecula1, XYL1961/18 and XYL468) were used to inoculate 50-mL glass flasks and incubated under agitation at 28 °C. After 9 days of growth, two types of samples were collected for polyamines extraction. One type of sample included *X. fastidiosa* cells from planktonic growth plus those present in the biofilms formed on the glass surface at the air-liquid interface. For that, cells were scratched from the attachment ring formed on the glass wall and vortexed. Then, the bacterial cell pellet (CP) was obtained after centrifugation at 14,000 rpm for 10 min of the entire culture. The second type of sample included the culture supernatant (SP). At least two biological replicates for each strain were performed. In the second experiment, three *X. fastidiosa* strains were used (ESVL, De Donno and Temecula1 strains). In this experiment, three types of samples were obtained after 5 and 12 days of growth. For each strain and sampling point, culture cells in planktonic growth were collected by transferring the entire culture to a new falcon tube. Then, a cell pellet from planktonic growth (CP-P) and a supernatant (SP) were obtained after

centrifugation as described above. To collect cells growing in biofilms, the biofilm formed on
the glass surface was scraped with a loop and resuspended with a pipette using 1 mL of sterile
distilled water. Then, a cell pellet from biofilms (CP-B) was collected by the centrifugation and
removal of the supernatant. In this second experiment, four biological replicates for each strain
and condition were obtained.

The polyamine extraction and quantification of the bacterial cell pellets (CP-P and CP-B) and
supernatant (SP) samples were carried out according to Solé-Gil and coworkers (2019) with
small modifications. Briefly, bacterial cell pellets were resuspended in 1 mL sterile water,
treated with lysozyme (10 mg·mL^{-1}), and kept on ice for 10 minutes. Subsequently, bacterial
cells were disrupted by sonication (10 cycles of 30 seconds each). After centrifugation at
10,000 rpm for 10 min, 500 µL of the supernatants were treated with 2.5 mL of perchloric acid
5 % (w/v). Culture supernatants (2.5 mL) were also treated with 2.5 mL of perchloric acid. At
this point, pre-treated samples were stored at -20 °C and sent for analysis of presence of
polyamines at the Metabolomics Service of the Institute of Molecular and Cellular Biology of
Plants (IBMCP, Valencia). Before analysis, 500 µL of the internal standard (1, 6-
Hexanediamine 1 mM, HD) were added to each sample. Polyamines were quantified by HPLC
analysis as described by Santos and coworkers (2021). Measurements were done in triplicate.

1097

Data processing and statistical analysis

Growth curves of the different *X. fastidiosa* strains growing in each PM compound or minimal
media were obtained by accumulating OD$_{600}$ values over time from the date of inoculation. To
evaluate growth in minimal media, the OD$_{600}$ value of the control wells was subtracted from
each time point to obtain the net absorbance. Cumulative growth was then calculated by
subtracting the initial net absorbance at the beginning of the experiment from the net
absorbance on each subsequent day. For both experiments, the standardized area under the
growth progress curve (sAUGPC) was calculated using the trapezoidal integration method
standardized by duration of the experiment in days (Simko & Piepho, 2012).

In the PM study, metabolic assimilation by *X. fastidiosa* was determined by normalizing the
sAUGPC of each compound against the strain- and plate-specific negative control (NC).
Following the criteria of Gerlin and coworkers (2020), a compound was considered as positive
carbon source ('Growth') if its sAUGPC was at least 1.14 times higher than the NC.
Specifically, a normalized sAUGPC was calculated as 100 x [sAUGPC - (1.14 * sAUGPC$_{NC}$)].
Negative values after normalization were set to zero. Growth was categorized based on

consistency: a compound was considered assimilated only if the normalized sAUGPC remained positive in at least three of the four biological replicates. Conversely, it was classified as 'No growth' if assimilation occurred in zero or one replicates; otherwise, growth was assessed to be 'Undetermined'.

Polyamine concentrations were quantified in both cell pellets (CP-P and CP-B) and supernatants (SP). For SP samples, net polyamine content was calculated by subtracting the mean concentration for each polyamine measured of the non-inoculated negative control medium (PD3). This correction was not applied to the pellet samples, as centrifugation removes the influence of the initial medium. Any negative values after normalization were set to zero.

Statistical analyses and data visualization were conducted using R software. Given the variability of bacterial growth and polyamines production, Wilcoxon rank-sum tests were used for all pairwise comparisons ($P \leq 0.05$). For minimal media and biofilm experiments, each tested medium was compared against the PD3 control, and between media pairs sharing the same carbon source. For polyamine data, differences between time points within each sample type and strain were assessed.

Data were visualized using violin plots for growth and biofilm formation in minimal media, and polyamine data. For polyamine results, a pseudo-logarithmic scale was used to facilitate comparison across several orders of magnitude. To visualize the metabolic profiles across PM plates, two heatmap approaches were used. First, a quantitative heatmap was generated to represent the normalized sAUGPC values, with a pseudo-logarithmic transformation for the color scale. Values of zero (no growth) were set as transparent. Second, discrete heatmaps were obtained to summarize growth categories across the four biological replicates. All visualizations were produced using ggplot2, ggpubr and ggh4x packages in R.

Author contributions

Paola Corbín Agustí: Conceptualization, Methodology, Formal analysis, Visualization, Investigation, Software, Data curation, Writing– Original draft, Reviewing and Editing. **Miguel Álvarez Herrera**: Methodology, Formal analysis, Visualization, Investigation, Software, Writing– Reviewing and editing. **Miguel Román Écija**: Methodology, Formal analysis, Investigation, Writing– Reviewing and editing. **Patricia Álvarez**: Methodology, Investigation, Software, Writing– Reviewing and editing. **Marta Tortajada**: Conceptualization, Writing–

Reviewing and editing. **Juli Peretó**: Conceptualization, Supervision, Funding acquisition, Writing– Original draft, Reviewing and Editing. **Blanca B. Landa**: Conceptualization, Supervision, Funding acquisition, Writing– Original draft, Reviewing and Editing.

Funding

P.C.-A. was recipient of a predoctoral fellowship from the Generalitat Valenciana "Subvenciones para la contratación de personal investigador de carácter predoctoral (ACIF)", with reference ACIF/2021/110. M.Á.-H. was supported by the Generalitat Valenciana and the European Social Fund "ESF Investing in your future" through grant CIACIF/2022/333. J.P. acknowledges the generic support of CYTED (Iber-Xyfas network "Red Iberoamericana para la vigilancia de *Xylella fastidiosa*", ref. 119RT0569) as well as of his host institutions Universitat de València and CSIC. This work was financed by the HORIZON-CL6-2021-FARM2FORK-01 project BeXyl (Beyond *Xylella*, Integrated Management Strategies for Mitigating *Xylella fastidiosa* impact in Europe; Grant ID No 101060593). The funders had no role in study design, data collection and analysis, decision to publish, or presentation of the manuscript.

Conflicts of Interest

M. Tortajada is a former employee of Biopolis-Archer Daniels Midland and contributed to this study in their personal capacity. The authors declare no conflicts of interest.

Data Availability Statement

All relevant data are within the manuscript and its additional supporting information. Detailed supplementary files and the code will be made publicly available upon peer-reviewed publication and are available from the corresponding author on request.

Acknowledgements

The authors thank the expert support of Miguel Ángel Blázquez and Francisco Vera-Sirera (IBMCP, UPV-CSIC) and the IBMCP Metabolomics Facility for the polyamine analyses. The authors thank Gonzalo de Oyanguren for the initial steps into the pangenome analysis. Figures 1C and 3 were created in BioRender. Smith, J. (2025).

References

Almeida, R. P. P., Mann, R., & Purcell, A. H. (2004). *Xylella fastidiosa* cultivation on a minimal solid defined medium. *Current Microbiology*, *48*(5), 368–372. https://doi.org/10.1007/s00284-003-4219-x

Anguita-Maeso, M., Haro, C., Montes-Borrego, M., Fuente, L. D. L., Navas-Cortés, J. A., & Landa, B. B. (2021). Metabolomic, ionic and microbial characterization of olive xylem sap reveals differences according to plant age and genotype. *Agronomy*, *11*(6). https://doi.org/10.3390/agronomy11061179

Ardalani, O., Phaneuf, P. V., Mohite, O. S., Nielsen, L. K., & Palsson, B. O. (2024). Pangenome reconstruction of *Lactobacillaceae* metabolism predicts species-specific metabolic traits. *mSystems*, *9*(7), e00156-24. https://doi.org/10.1128/msystems.00156-24

Blázquez, M. A. (2024). Polyamines: their role in plant development and stress. *Annual Review of Plant Biology*, *75*(Volume 75, 2024), 95–117. https://doi.org/10.1146/annurev-arplant-070623-110056

Brämer, C. O., & Steinbüchel, A. (2001). The methylcitric acid pathway in *Ralstonia eutropha*: New genes identified involved in propionate metabolism. *Microbiology*, *147*(8), 2203–2214. https://doi.org/10.1099/00221287-147-8-2203

Brock, M., Maerker, C., Schütz, A., Völker, U., & Buckel, W. (2002). Oxidation of propionate to pyruvate in *Escherichia coli*: Involvement of methylcitrate dehydratase and aconitase. *European Journal of Biochemistry*, *269*(24), 6184–6194. https://doi.org/10.1046/j.1432-1033.2002.03336.x

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1), 421. https://doi.org/10.1186/1471-2105-10-421

Campanharo, J. C., Lemos, M. V. F., & Lemos, E. G. de M. (2003). Growth optimization procedures for the phytopathogen *Xylella fastidiosa*. *Current Microbiology*, *46*(2), 0099–0102. https://doi.org/10.1007/s00284-002-3829-z

Caspi, R., Billington, R., Keseler, I. M., Kothari, A., Krummenacker, M., Midford, P. E., Ong, W. K., Paley, S., Subhraveti, P., & Karp, P. D. (2020). The MetaCyc database of metabolic pathways and enzymes—A 2019 update. *Nucleic Acids Research*, *48*(D1), D445–D453. https://doi.org/10.1093/nar/gkz862

Castro, C., Ndukwe, I., Heiss, C., Black, I., Ingel, B. M., Guevara, M., Sun, Y., Azadi, P., Sun, Q., & Roper, M. C. (2023). *Xylella fastidiosa* modulates exopolysaccharide polymer length and the dynamics of biofilm development with a β-1,4-endoglucanase. *mBio*,

14(5), e01395-23. https://doi.org/10.1128/mbio.01395-23

Chang, A., Jeske, L., Ulbrich, S., Hofmann, J., Koblitz, J., Schomburg, I., Neumann-Schaal,
M., Jahn, D., & Schomburg, D. (2021). BRENDA, the ELIXIR core data resource in
2021: New developments and updates. *Nucleic Acids Research*, *49*(D1), D498–D508.
https://doi.org/10.1093/nar/gkaa1025

Chang, C. J. (1993). *Xylella fastidiosa*: Cultivation in chemically defined medium.
Phytopathology, *83*(2), 192. https://doi.org/10.1094/Phyto-83-192

Chatterjee, S., Almeida, R. P. P., & Lindow, S. (2008). Living in two worlds: the plant and
insect lifestyles of *Xylella fastidiosa. Annual Review of Phytopathology*, *46*(1), 243–
271. https://doi.org/10.1146/annurev.phyto.45.062806.094342

Chavarría, M., Nikel, P. I., Pérez-Pantoja, D., & de Lorenzo, V. (2013). The Entner-Doudoroff
pathway empowers *Pseudomonas putida* KT2440 with a high tolerance to oxidative
stress. *Environmental Microbiology*, *15*(6), 1772–1785. https://doi.org/10.1111/1462-
2920.12069

Cooley, N. P., & Wright, E. S. (2024). Many purported pseudogenes in bacterial genomes are
bona fide genes. *BMC Genomics*, *25*(1), 365. https://doi.org/10.1186/s12864-024-
10137-0

Cross, K. L., Campbell, J. H., Balachandran, M., Campbell, A. G., Cooper, C. J., Griffen, A.,
Heaton, M., Joshi, S., Klingeman, D., Leys, E., Yang, Z., Parks, J. M., & Podar, M.
(2019). Targeted isolation and cultivation of uncultivated bacteria by reverse genomics.
Nature Biotechnology, *37*(11), 1314–1321. https://doi.org/10.1038/s41587-019-0260-
6

da Silva, F. R., Vettore, A. L., Kemper, E. L., Leite, A., & Arruda, P. (2001). Fastidian gum:
The *Xylella fastidiosa* exopolysaccharide possibly involved in bacterial pathogenicity.
FEMS Microbiology Letters, *203*(2), 165–171. https://doi.org/10.1111/j.1574-
6968.2001.tb10836.x

Davis, M. J. (1992). Fastidious bacteria of plant vascular tissues and their invertebrate vectors.
In *The Prokaryotes* (pp. 4030–4049). *Springer*. https://doi.org/10.1007/978-1-4757-
2191-1_66

Davis, M. J., French, W. J., & Schaad, N. W. (1981). Axenic culture of the bacteria associated
with phony disease of peach and plum leaf scald. *Current Microbiology*, *6*(5), 309–314.
https://doi.org/10.1007/BF01566883

Davis, M. J., Gillaspie, A. G., Harris, R. W., & Lawson, R. H. (1980). Ratoon Stunting Disease
of Sugarcane: Isolation of the Causal Bacterium. *Science*, *210*(4476), 1365–1367.
https://doi.org/10.1126/science.210.4476.1365

De La Fuente, L., Navas-Cortés, J. A., & Landa, B. B. (2024). Ten challenges to understanding

and managing the insect-transmitted, xylem-limited bacterial pathogen *Xylella fastidiosa*. *Phytopathology®, 114*(5), 869–884. https://doi.org/10.1094/PHYTO-12-23-0476-KC

de Macedo Lemos, E. G., Alves, L. M. C., & Campanharo, J. C. (2003). Genomics-based design of defined growth media for the plant pathogen *Xylella fastidiosa*. *FEMS Microbiology Letters, 219*(1), 39–45. https://doi.org/10.1016/S0378-1097(02)01189-8

Denancé, N., Briand, M., Gaborieau, R., Gaillard, S., & Jacques, M.-A. (2019). Identification of genetic relationships and subspecies signatures in *Xylella fastidiosa*. *BMC Genomics, 20*(1), 239. https://doi.org/10.1186/s12864-019-5565-9

Dolan, S. K., Wijaya, A., Geddis, S. M., Spring, D. R., Silva-Rocha, R., & Welch, M. (2018). Loving the poison: The methylcitrate cycle and bacterial pathogenesis. *Microbiology, 164*(3), 251–259. https://doi.org/10.1099/mic.0.000604

Ebrahim, A., Lerman, J. A., Palsson, B. O., & Hyduke, D. R. (2013). COBRApy: COnstraints-Based Reconstruction and Analysis for Python. *BMC Systems Biology, 7*(1), 74. https://doi.org/10.1186/1752-0509-7-74

EFSA (European Food Safety Authority), E. F. S., Cavalieri, V., Fasanelli, E., Furnari, G., Gibin, D., Gutierrez Linares, A., La Notte, P., Pasinato, L., & Stancanelli, G. (2026). Update of the *Xylella* spp. Host plant database – Systematic literature search up to 30 June 2025. *EFSA Journal, 24*(1), e9840. https://doi.org/10.2903/j.efsa.2026.9840

Erb, T. J., Berg, I. A., Brecht, V., Müller, M., Fuchs, G., & Alber, B. E. (2007). Synthesis of C_5-dicarboxylic acids from C_2-units involving crotonyl-CoA carboxylase/reductase: The ethylmalonyl-CoA pathway. *Proceedings of the National Academy of Sciences, 104*(25), 10631–10636. https://doi.org/10.1073/pnas.0702791104

Facincani, A. P., Ferro, J. A., Pizauro Jr., J. M., Pereira Jr., H. A., Lemos, E. G. de M., Prado, A. L. do, & Ferro, M. I. T. (2003). Carbohydrate metabolism of *Xylella fastidiosa*: Detection of glycolytic and pentose phosphate pathway enzymes and cloning and expression of the enolase gene. *Genetics and Molecular Biology, 26*, 203–211. https://doi.org/https://doi.org/10.1590/S1415-47572003000200015

Fausto, C., Araniti, F., Mininni, A. N., Crecchio, C., Scagliola, M., Bleve, G., Dichio, B., & Sofo, A. (2021). Differential olive grove management regulates the levels of primary metabolites in xylem sap. *Plant and Soil, 460*(1–2), 281–296. https://doi.org/10.1007/s11104-020-04800-0

Garschagen, L. S., Franke, T., & Deppenmeier, U. (2021). An alternative pentose phosphate pathway in human gut bacteria for the degradation of C5 sugars in dietary fibers. *The FEBS Journal, 288*(6), 1839–1858. https://doi.org/10.1111/febs.15511

Gerlin, L., Cottret, L., Cesbron, S., Taghouti, G., Jacques, M.-A., Genin, S., & Baroukh, C.

(2020). Genome-scale investigation of the metabolic determinants generating bacterial fastidious growth. *mSystems*, *5*(2), 10.1128/msystems.00698-19. https://doi.org/10.1128/msystems.00698-19

Gil, R., & Peretó, J. (2015). Small genomes and the difficulty to define minimal translation and metabolic machineries. *Frontiers in Ecology and Evolution*, *3*. https://doi.org/10.3389/fevo.2015.00123

Gu, C., Kim, G. B., Kim, W. J., Kim, H. U., & Lee, S. Y. (2019). Current status and applications of genome-scale metabolic models. *Genome Biology*, *20*(1), 121. https://doi.org/10.1186/s13059-019-1730-3

Hagberg, A., Swart, P. J., & Schult, D. A. (2008). Exploring network structure, dynamics, and function using NetworkX. In G. Varoquaux, T. Vaught, & J. Millman (Eds.), *Proceedings of the 7th Python in Science Conference (SciPy2008)* (pp. 11–15).

Hanage, W. P., Fraser, C., & Spratt, B. G. (2005). Fuzzy species among recombinogenic bacteria. *BMC Biology*, *3*(1), 6. https://doi.org/10.1186/1741-7007-3-6

Henry, C. S., DeJongh, M., Best, A. A., Frybarger, P. M., Linsay, B., & Stevens, R. L. (2010). High-throughput generation, optimization and analysis of genome-scale metabolic models. *Nature Biotechnology*, *28*(9), 977–982. https://doi.org/10.1038/nbt.1672

Herter, S., Fuchs, G., Bacher, A., & Eisenreich, W. (2002). A bicyclic autotrophic CO2 fixation pathway in *Chloroflexus aurantiacus*. *Journal of Biological Chemistry*, *277*(23), 20277–20283. https://doi.org/10.1074/jbc.M201030200

Hiery, E., Adam, S., Reid, S., Hofmann, J., Sonnewald, S., & Burkovski, A. (2013). Genome-wide transcriptome analysis of *Clavibacter michiganensis* subsp. *michiganensis* grown in xylem mimicking medium. *Journal of Biotechnology*, *168*(4), 348–354. https://doi.org/10.1016/j.jbiotec.2013.09.006

Horgan, C., Baccari, C., O'Driscoll, M., Lindow, S. E., & O'Sullivan, T. P. (2024). BDSF analogues inhibit quorum sensing-regulated biofilm production in *Xylella fastidiosa*. *Microorganisms*, *12*(12), 2496. https://doi.org/10.3390/microorganisms12122496

Ibarra, R. U., Edwards, J. S., & Palsson, B. O. (2002). *Escherichia coli* K-12 undergoes adaptive evolution to achieve in silico predicted optimal growth. *Nature*, *420*(6912), 186–189. https://doi.org/10.1038/nature01149

Igarashi, K., & Kashiwagi, K. (2010). Characteristics of cellular polyamine transport in prokaryotes and eukaryotes. *Plant Physiology and Biochemistry: PPB*, *48*(7), 506–512. https://doi.org/10.1016/j.plaphy.2010.01.017

Janissen, R., Murillo, D. M., Niza, B., Sahoo, P. K., Nobrega, M. M., Cesar, C. L., Temperini, M. L. A., Carvalho, H. F., De Souza, A. A., & Cotta, M. A. (2015). Spatiotemporal distribution of different extracellular polymeric substances and filamentation mediate

Xylella fastidiosa adhesion and biofilm formation. *Scientific Reports*, *5*(1), 9856. https://doi.org/10.1038/srep09856

Jolley, K. A., Bray, J. E., & Maiden, M. C. J. (2018). Open-access bacterial population genomics: BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Research*, *3*, 124. https://doi.org/10.12688/wellcomeopenres.14826.1

Kanehisa, M., Furumichi, M., Sato, Y., Matsuura, Y., & Ishiguro-Watanabe, M. (2025). KEGG: Biological systems database as a model of the real world. *Nucleic Acids Research*, *53*(D1), D672–D677. https://doi.org/10.1093/nar/gkae909

Kanehisa, M., Sato, Y., & Morishima, K. (2016). BlastKOALA and GhostKOALA: KEGG Tools for functional characterization of genome and metagenome sequences. *Journal of Molecular Biology*, *428*(4), 726–731. https://doi.org/10.1016/j.jmb.2015.11.006

Karatan, E., & Michael, A. J. (2013). A wider role for polyamines in biofilm formation. *Biotechnology Letters*, *35*(11), 1715–1717. https://doi.org/10.1007/s10529-013-1286-3

Karp, P. D., Billington, R., Caspi, R., Fulcher, C. A., Latendresse, M., Kothari, A., Keseler, I. M., Krummenacker, M., Midford, P. E., Ong, Q., Ong, W. K., Paley, S. M., & Subhraveti, P. (2019). The BioCyc collection of microbial genomes and metabolic pathways. *Briefings in Bioinformatics*, *20*(4), 1085–1093. https://doi.org/10.1093/bib/bbx085

Khomyakova, M., Bükmez, Ö., Thomas, L. K., Erb, T. J., & Berg, I. A. (2011). A methylaspartate cycle in Haloarchaea. *Science*, *331*(6015), 334–337. https://doi.org/10.1126/science.1196544

Kim, J., Kershner, J. P., Novikov, Y., Shoemaker, R. K., & Copley, S. D. (2010). Three serendipitous pathways in *E. coli* can bypass a block in pyridoxal-5'-phosphate synthesis. *Molecular Systems Biology*, *6*, 436. https://doi.org/10.1038/msb.2010.88

King, Z. A., Dräger, A., Ebrahim, A., Sonnenschein, N., Lewis, N. E., & Palsson, B. O. (2015). Escher: A web application for building, sharing, and embedding data-rich visualizations of biological pathways. *PLOS Computational Biology*, *11*(8), e1004321. https://doi.org/10.1371/journal.pcbi.1004321

King, Z. A., Lu, J., Dräger, A., Miller, P., Federowicz, S., Lerman, J. A., Ebrahim, A., Palsson, B. O., & Lewis, N. E. (2016). BiGG Models: A platform for integrating, standardizing and sharing genome-scale models. *Nucleic Acids Research*, *44*(D1), D515–D522. https://doi.org/10.1093/nar/gkv1049

Koendjbiharie, J. G., Hon, S., Pabst, M., Hooftman, R., Stevenson, D. M., Cui, J., Amador-Noguez, D., Lynd, L. R., Olson, D. G., & van Kranenburg, R. (2020). The pentose phosphate pathway of cellulolytic clostridia relies on 6-phosphofructokinase instead of transaldolase. *The Journal of Biological Chemistry*, *295*(7), 1867–1878.

https://doi.org/10.1074/jbc.RA119.011239

Kornberg, H. L., & Krebs, H. A. (1957). Synthesis of cell constituents from C2-units by a modified tricarboxylic acid cycle. *Nature*, *179*(4568), 988–991. https://doi.org/10.1038/179988a0

Kremer, K., Van Teeseling, M. C. F., Schada Von Borzyskowski, L., Bernhardsgrütter, I., Van Spanning, R. J. M., Gates, A. J., Remus-Emsermann, M. N. P., Thanbichler, M., & Erb, T. J. (2019). Dynamic metabolic rewiring enables efficient acetyl coenzyme a assimilation in *Paracoccus denitrificans*. *mBio*, *10*(4), e00805-19. https://doi.org/10.1128/mBio.00805-19

Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Molecular Biology and Evolution*, *35*(6), 1547–1549. https://doi.org/10.1093/molbev/msy096

Landa, B. B., Castillo, A. I., Giampetruzzi, A., Kahn, A., Román-Écija, M., Velasco-Amo, M. P., Navas-Cortés, J. A., Marco-Noales, E., Barbé, S., Moralejo, E., Coletta-Filho, H. D., Saldarelli, P., Saponari, M., & Almeida, R. P. P. (2020). Emergence of a plant pathogen in Europe associated with multiple intercontinental introductions. *Applied and Environmental Microbiology*, *86*(3), e01521-19. https://doi.org/10.1128/AEM.01521-19

Landa, B. B., Saponari, M., Feitosa-Junior, O. R., Giampetruzzi, A., Vieira, F. J. D., Mor, E., & Robatzek, S. (2022). *Xylella fastidiosa*'s relationships: The bacterium, the host plants, and the plant microbiome. *New Phytologist*, *234*(5), 1598–1605. https://doi.org/10.1111/nph.18089

Leite, B., Andersen, P. C., & Ishida, M. L. (2004). Colony aggregation and biofilm formation in xylem chemistry-based media for *Xylella fastidiosa*. *FEMS Microbiology Letters*, *230*(2), 283–290. https://doi.org/10.1016/S0378-1097(03)00917-0

Leonidou, N., Fritze, E., Renz, A., & Dräger, A. (2023). SBOannotator: A Python tool for the automated assignment of systems biology ontology terms. *Bioinformatics*, *39*(7). https://doi.org/10.1093/bioinformatics/btad437

Lewis, W. H., Tahon, G., Geesink, P., Sousa, D. Z., & Ettema, T. J. G. (2021). Innovations to culturing the uncultured microbial majority. *Nature Reviews Microbiology*, *19*(4), 225–240. https://doi.org/10.1038/s41579-020-00458-8

Lieven, C., Beber, M. E., Olivier, B. G., Bergmann, F. T., Ataman, M., Babaei, P., Bartell, J. A., Blank, L. M., Chauhan, S., Correia, K., Diener, C., Dräger, A., Ebert, B. E., Edirisinghe, J. N., Faria, J. P., Feist, A. M., Fengos, G., Fleming, R. M. T., García-Jiménez, B., … Zhang, C. (2020). MEMOTE for standardized genome-scale metabolic model testing. *Nature Biotechnology*, *38*(3), 272–276. https://doi.org/10.1038/s41587-

020-0446-y

Lowe-Power, T. M., Hendrich, C. G., Von Roepenack-Lahaye, E., Li, B., Wu, D., Mitra, R., Dalsing, B. L., Ricca, P., Naidoo, J., Cook, D., Jancewicz, A., Masson, P., Thomma, B., Lahaye, T., Michael, A. J., & Allen, C. (2018). Metabolomics of tomato xylem sap during bacterial wilt reveals *Ralstonia solanacearum* produces abundant putrescine, a metabolite that accelerates wilt disease. *Environmental Microbiology*, *20*(4), 1330–1349. https://doi.org/10.1111/1462-2920.14020

Lv, P., Tang, W., Wang, P., Cao, Z., & Zhu, G. (2018). Enzymatic characterization and functional implication of two structurally different isocitrate dehydrogenases from *Xylella fastidiosa*. *Biotechnology and Applied Biochemistry*, *65*(2), 230–237. https://doi.org/10.1002/bab.1560

Madeira, F., Madhusoodanan, N., Lee, J., Eusebi, A., Niewielska, A., Tivey, A. R. N., Lopez, R., & Butcher, S. (2024). The EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024. *Nucleic Acids Research*, *52*(W1), W521–W525. https://doi.org/10.1093/nar/gkae241

Malik-Sheriff, R. S., Glont, M., Nguyen, T. V. N., Tiwari, K., Roberts, M. G., Xavier, A., Vu, M. T., Men, J., Maire, M., Kananathan, S., Fairbanks, E. L., Meyer, J. P., Arankalle, C., Varusai, T. M., Knight-Schrijver, V., Li, L., Dueñas-Roca, C., Dass, G., Keating, S. M., … Hermjakob, H. (2020). BioModels—15 years of sharing computational models in life science. *Nucleic Acids Research*, *48*(D1), D407–D415. https://doi.org/10.1093/nar/gkz1055

Marques, L. L. R., Ceri, H., Manfio, G. P., Reid, D. M., & Olson, M. E. (2002). Characterization of biofilm formation by *Xylella fastidiosa* in vitro. *Plant Disease*, *86*(6), 633–638. https://doi.org/10.1094/PDIS.2002.86.6.633

Meidanis, J., Braga, M. D. V., & Verjovski-Almeida, S. (2002). Whole-genome analysis of transporters in the plant pathogen *Xylella fastidiosa*. *Microbiology and Molecular Biology Reviews*, *66*(2), 272–299. https://doi.org/10.1128/MMBR.66.2.272-299.2002

Mirhakkak, M. H., Chen, X., Ni, Y., Heinekamp, T., Sae-Ong, T., Xu, L.-L., Kurzai, O., Barber, A. E., Brakhage, A. A., Boutin, S., Schäuble, S., & Panagiotou, G. (2023). Genome-scale metabolic modeling of *Aspergillus fumigatus* strains reveals growth dependencies on the lung microbiome. *Nature Communications*, *14*(1), 4369. https://doi.org/10.1038/s41467-023-39982-5

Mistry, J., Chuguransky, S., Williams, L., Qureshi, M., Salazar, G. A., Sonnhammer, E. L. L., Tosatto, S. C. E., Paladin, L., Raj, S., Richardson, L. J., Finn, R. D., & Bateman, A. (2021). Pfam: The protein families database in 2021. *Nucleic Acids Research*, *49*(D1), D412–D419. https://doi.org/10.1093/nar/gkaa913

Mistry, J., Finn, R. D., Eddy, S. R., Bateman, A., & Punta, M. (2013). Challenges in homology
search: HMMER3 and convergent evolution of coiled-coil regions. *Nucleic Acids*
Research, *41*(12), e121. https://doi.org/10.1093/nar/gkt263

Monk, J., Nogales, J., & Palsson, B. O. (2014). Optimizing genome-scale network
reconstructions. *Nature Biotechnology*, *32*(5), 447–452.
https://doi.org/10.1038/nbt.2870

Nakahigashi, K., Toya, Y., Ishii, N., Soga, T., Hasegawa, M., Watanabe, H., Takai, Y., Honma,
M., Mori, H., & Tomita, M. (2009). Systematic phenome analysis of *Escherichia coli*
multiple-knockout mutants reveals hidden reactions in central carbon metabolism.
Molecular Systems Biology, *5*(1), MSB200965. https://doi.org/10.1038/msb.2009.65

Nascimento, R., Gouran, H., Chakraborty, S., Gillespie, H. W., Almeida-Souza, H. O., Tu, A.,
Rao, B. J., Feldstein, P. A., Bruening, G., Goulart, L. R., & Dandekar, A. M. (2016).
The type II secreted lipase/esterase LesA is a key virulence factor required for *Xylella*
fastidiosa pathogenesis in grapevines. *Scientific Reports*, *6*(1), 18598.
https://doi.org/10.1038/srep18598

Neumann, M. (2003). Sequence tag analysis of gene expression during pathogenic growth
and microsclerotia development in the vascular wilt pathogen *Verticillium dahliae*.
Fungal Genetics and Biology, *38*(1), 54–62. https://doi.org/10.1016/S1087-
1845(02)00507-8

Nogales, J. (2017). A practical protocol for genome-scale metabolic reconstructions. In T. J.
McGenity, K. N. Timmis, & B. Nogales (Eds), *Hydrocarbon and lipid microbiology*
protocols: genetic, genomic and system analyses of pure cultures (pp. 197–221).
Springer. https://doi.org/10.1007/8623_2014_12

Norsigian, C. J., Fang, X., Seif, Y., Monk, J. M., & Palsson, B. O. (2020). A workflow for
generating multi-strain genome-scale metabolic models of prokaryotes. *Nature*
Protocols, *15*(1), 1–14. https://doi.org/10.1038/s41596-019-0254-3

O'Brien, E. J., Monk, J. M., & Palsson, B. O. (2015). Using genome-scale models to predict
biological capabilities. *Cell*, *161*(5), 971–987.
https://doi.org/10.1016/j.cell.2015.05.019

Oliveira, A., Cunha, E., Silva, M., Faria, C., & Dias, O. (2023). Exploring *Xylella fastidiosa*'s
metabolic traits using a GSM model of the phytopathogenic bacterium. In F. Fdez-
Riverola, M. Rocha, M. S. Mohamad, S. Caraiman, & A. B. Gil-González (Eds),
Practical Applications of Computational Biology and Bioinformatics, 16th International
Conference (PACBB 2022) (Vol. 553, pp. 79–88). Springer International Publishing.
https://doi.org/10.1007/978-3-031-17024-9_8

Orth, J. D., Conrad, T. M., Na, J., Lerman, J. A., Nam, H., Feist, A. M., & Palsson, B. Ø. (2011).

A comprehensive genome-scale reconstruction of *Escherichia coli* metabolism—2011. *Molecular Systems Biology*, *7*(1), 535. https://doi.org/10.1038/msb.2011.65

Orth, J. D., Thiele, I., & Palsson, B. Ø. (2010). What is flux balance analysis? *Nature Biotechnology*, *28*(3), 245–248. https://doi.org/10.1038/nbt.1614

Overbeek, R., Begley, T., Butler, R. M., Choudhuri, J. V., Chuang, H.-Y., Cohoon, M., de Crécy-Lagard, V., Diaz, N., Disz, T., Edwards, R., Fonstein, M., Frank, E. D., Gerdes, S., Glass, E. M., Goesmann, A., Hanson, A., Iwata-Reuyl, D., Jensen, R., Jamshidi, N., … Vonstein, V. (2005). The subsystems approach to genome annotation and its use in the project to annotate 1000 genomes. *Nucleic Acids Research*, *33*(17), 5691–5702. https://doi.org/10.1093/nar/gki866

Palsson, B. Ø. (2015). *Systems Biology: Constraint-based reconstruction and analysis*. Cambridge University Press. https://doi.org/10.1017/CBO9781139854610

Peyraud, R., Cottret, L., Marmiesse, L., Gouzy, J., & Genin, S. (2016). A Resource Allocation Trade-Off between Virulence and Proliferation Drives Metabolic Versatility in the Plant Pathogen Ralstonia solanacearum. *PLOS Pathogens*, *12*(10), e1005939. https://doi.org/10.1371/journal.ppat.1005939

PM 7/24 (5) Xylella fastidiosa. (2023). *EPPO Bulletin*, *53*(2), 205–276. https://doi.org/10.1111/epp.12923

Ponce-de-Leon, M., Calle-Espinosa, J., Peretó, J., & Montero, F. (2015). Consistency analysis of genome-scale models of bacterial metabolism: A metamodel approach. *PLOS ONE*, *10*(12), e0143626. https://doi.org/10.1371/journal.pone.0143626

Purcino, R. P., Medina, C. L., Martins de Souza, D., Winck, F. V., Machado, E. C., Novello, J. C., Machado, M. A., & Mazzafera, P. (2007). *Xylella fastidiosa* disturbs nitrogen metabolism and causes a stress response in sweet orange *Citrus sinensis* cv. Pera. *Journal of Experimental Botany*, *58*(11), 2733–2744. https://doi.org/10.1093/jxb/erm138

Qian, J., & Ye, C. (2024). Development and applications of genome-scale metabolic network models. In *Advances in Applied Microbiology* (Vol. 126, pp. 1–26). Academic Press. https://doi.org/10.1016/bs.aambs.2024.02.002

Remm, M., Storm, C. E. V., & Sonnhammer, E. L. L. (2001). Automatic clustering of orthologs and in-paralogs from pairwise species comparisons. *Journal of Molecular Biology*, *314*(5), 1041–1052. https://doi.org/10.1006/jmbi.2000.5197

Román-Écija, M., Navas-Cortés, J. A., Velasco-Amo, M. P., Arias-Giraldo, L. F., Gómez, L. M., Fuente, L. D. L., & Landa, B. B. (2023). Two *Xylella fastidiosa* subsp. *multiplex* strains isolated from almond in spain differ in plasmid content and virulence traits. *Phytopathology®*, *113*(6), 960–974. https://doi.org/10.1094/PHYTO-06-22-0234-R

Sanchez, F. B., Barreiro, H. J., Soto, E. I., & Rodriguez, C. E. (2019). *The Impact Indicator for Priority Pests (I2P2): A tool for ranking pests according to Regulation (EU) 2016/2031*. Publications Office of the European Union. https://doi.org/10.2760/585182

Santos, A. P., Muratore, L. N., Solé-Gil, A., Farías, M. E., Ferrando, A., Blázquez, M. A., & Belfiore, C. (2021). Extremophilic bacteria restrict the growth of *Macrophomina phaseolina* by combined secretion of polyamines and lytic enzymes. *Biotechnology Reports*, *32*, e00674. https://doi.org/10.1016/j.btre.2021.e00674

Sayers, E. W., Beck, J., Bolton, E. E., Brister, J. R., Chan, J., Connor, R., Feldgarden, M., Fine, A. M., Funk, K., Hoffman, J., Kannan, S., Kelly, C., Klimke, W., Kim, S., Lathrop, S., Marchler-Bauer, A., Murphy, T. D., O'Sullivan, C., Schmieder, E., … Pruitt, K. D. (2024). Database resources of the National Center for Biotechnology Information in 2025. *Nucleic Acids Research*, *53*(D1), D20–D29. https://doi.org/10.1093/nar/gkae979

Scala, V., Salustri, M., Merfa, M. V., Beccaccioli, M., Lascala, L., De La Fuente, L., & Reverberi, M. (2025). XadA-like adhesin XADA2 regulates biofilm formation in *X. fastidiosa* subsp. *fastidiosa* putatively by engaging oleic-acid derived oxylipins. *Molecular Biology Reports*, *52*(1), 263. https://doi.org/10.1007/s11033-025-10259-y

Scally, M., Schuenzel, E. L., Stouthamer, R., & Nunney, L. (2005). Multilocus sequence type system for the plant pathogen *Xylella fastidiosa* and relative contributions of recombination and point mutation to clonal diversity. *Applied and Environmental Microbiology*, *71*(12), 8491–8499. https://doi.org/10.1128/AEM.71.12.8491-8499.2005

Schada von Borzyskowski, L., Bernhardsgrütter, I., & Erb, T. J. (2020). Biochemical unity revisited: Microbial central carbon metabolism holds new discoveries, multi-tasking pathways, and redundancies with a reason. *Biological Chemistry*, *401*(12), 1429–1441. https://doi.org/10.1515/hsz-2020-0214

Schatschneider, S., Persicke, M., Watt, S. A., Hublik, G., Pühler, A., Niehaus, K., & Vorhölter, F.-J. (2013). Establishment, in silico analysis, and experimental verification of a large-scale metabolic network of the xanthan producing *Xanthomonas campestris* pv. campestris strain B100. *Journal of Biotechnology*, *167*(2), 123–134. https://doi.org/10.1016/j.jbiotec.2013.01.023

Segura, D., Mahadevan, R., Juárez, K., & Lovley, D. R. (2008). Computational and Experimental analysis of redundancy in the central metabolism of *Geobacter sulfurreducens*. *PLOS Computational Biology*, *4*(2), e36. https://doi.org/10.1371/journal.pcbi.0040036

Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D., Amin, N., Schwikowski, B., & Ideker, T. (2003). Cytoscape: A software environment for integrated models of biomolecular interaction networks. *Genome Research*, *13*(11),

2498–2504. https://doi.org/10.1101/gr.1239303

Simko, I., & Piepho, H.-P. (2012). The area under the disease progress stairs: Calculation, advantage, and application. *Phytopathology®*, *102*(4), 381–389. https://doi.org/10.1094/PHYTO-07-11-0216

Simpson, A. J., Reinach, F. C., Arruda, P., Abreu, F. A., Acencio, M., Alvarenga, R., Alves, L. M., Araya, J. E., Baia, G. S., Baptista, C. S., Barros, M. H., Bonaccorsi, E. D., Bordin, S., Bové, J. M., Briones, M. R., Bueno, M. R., Camargo, A. A., Camargo, L. E., Carraro, D. M., … Setubal, J. C. (2000). The genome sequence of the plant pathogen *Xylella fastidiosa*. *Nature*, *406*(6792), 151–157. https://doi.org/10.1038/35018003

Solé-Gil, A., Hernández-García, J., López-Gresa, M. P., Blázquez, M. A., & Agustí, J. (2019). Conservation of thermospermine synthase activity in vascular and non-vascular plants. *Frontiers in Plant Science*, *10*. https://doi.org/10.3389/fpls.2019.00663

Solmi, L., Rossi, F. R., Romero, F. M., Bach-Pages, M., Preston, G. M., & Gárriz, A. (2025). The influence of the polyamine synthesis pathways on *Pseudomonas syringae* virulence and plant interaction. *Microbiology*, *171*(6), 001569. https://doi.org/10.1099/mic.0.001569

Sugiyama, Y., Nara, M., Sakanaka, M., Gotoh, A., Kitakata, A., Okuda, S., & Kurihara, S. (2017). Comprehensive analysis of polyamine transport and biosynthesis in the dominant human gut bacteria: Potential presence of novel polyamine metabolism and transport genes. *The International Journal of Biochemistry & Cell Biology*, *93*, 52–61. https://doi.org/10.1016/j.biocel.2017.10.015

Surano, A., del Grosso, C., Musio, B., Todisco, S., Giampetruzzi, A., Altamura, G., Saponari, M., Gallo, V., Mastrorilli, P., Boscia, D., & Saldarelli, P. (2024). Exploring the xylem-sap to unravel biological features of *Xylella fastidiosa* subspecies *pauca* ST53 in immune, resistant and susceptible crop species through metabolomics and in vitro studies. *Frontiers in Plant Science*, *14*. https://doi.org/10.3389/fpls.2023.1343876

Tabuchi, T., Serizawa, N., & Uchiyama, H. (1974). A novel pathway for the partial oxidation of propionyl-coa to pyruvate *via* seven-carbon tricarboxylic acids in yeasts. *Agricultural and Biological Chemistry*, *38*(12), 2571–2572. https://doi.org/10.1080/00021369.1974.10858334

Thiele, I., & Palsson, B. Ø. (2010). A protocol for generating a high-quality genome-scale metabolic reconstruction. *Nature Protocols*, *5*(1), 93–121. https://doi.org/10.1038/nprot.2009.203

UniProt Consortium. (2025). UniProt: The Universal Protein Knowledgebase in 2025. *Nucleic Acids Research*, *53*(D1), D609–D617. https://doi.org/10.1093/nar/gkae1010

Vanhove, M., Retchless, A. C., Sicard, A., Rieux, A., Coletta-Filho, H. D., De La Fuente, L.,

Stenger, D. C., & Almeida, R. P. P. (2019). Genomic diversity and recombination among *Xylella fastidiosa* subspecies. *Applied and Environmental Microbiology*, *85*(13), e02972-18. https://doi.org/10.1128/AEM.02972-18

Wang, P., Lee, Y., Igo, M. M., & Roper, M. C. (2017). Tolerance to oxidative stress is required for maximal xylem colonization by the xylem-limited bacterial phytopathogen, *Xylella fastidiosa*. *Molecular Plant Pathology*, *18*(7), 990–1000. https://doi.org/10.1111/mpp.12456

Wells, J. M., Raju, B. C., Hung, H.-Y., Weisburg, W. G., Mandelco-Paul, L., & Brenner, D. J. (1987). *Xylella fastidiosa* gen. nov., sp. nov: Gram-negative, xylem-limited, fastidious plant bacteria related to *Xanthomonas* spp. *International Journal of Systematic and Evolutionary Microbiology*, *37*(2), 136–143. https://doi.org/10.1099/00207713-37-2-136

Wells, J. M., Raju, B. C., Nyland, G., & Lowe, S. K. (1981). Medium for isolation and growth of bacteria associated with plum leaf scald and phony peach diseases. *Applied and Environmental Microbiology*, *42*(2), 357–363. https://doi.org/10.1128/aem.42.2.357-363.1981

Ye, C., Wei, X., Shi, T., Sun, X., Xu, N., Gao, C., & Zou, W. (2022). Genome-scale metabolic network models: From first-generation to next-generation. *Applied Microbiology and Biotechnology*, *106*(13), 4907–4920. https://doi.org/10.1007/s00253-022-12066-y

Youssefi, F., Brown, P., & Weinbaum, S. (2000). Relationship between tree nitrogen status, xylem and phloem sap amino acid concentrations, and apparent soil nitrogen uptake by almond trees *(Prunus dulcis)*. *The Journal of Horticultural Science and Biotechnology*, *75*(1), 62–68. https://doi.org/10.1080/14620316.2000.11511201

Yus, E., Maier, T., Michalodimitrakis, K., van Noort, V., Yamada, T., Chen, W.-H., Wodke, J. A. H., Güell, M., Martínez, S., Bourgeois, R., Kühner, S., Raineri, E., Letunic, I., Kalinina, O. V., Rode, M., Herrmann, R., Gutiérrez-Gallego, R., Russell, R. B., Gavin, A.-C., … Serrano, L. (2009). Impact of genome reduction on bacterial metabolism and its regulation. *Science*, *326*(5957), 1263–1268. https://doi.org/10.1126/science.1177263

yWorks GmbH (2020). *yEd Graph Editor* (Version 3.18.2) [Computer software]. https://www.yworks.com/products/yed

Zarzycki, J., & Fuchs, G. (2011). Coassimilation of organic substrates via the autotrophic 3-hydroxypropionate bi-cycle in *Chloroflexus aurantiacus*. *Applied and Environmental Microbiology*, *77*(17), 6181–6188. https://doi.org/10.1128/AEM.00705-11

Zhu, G., Golding, G. B., & Dean, A. M. (2005). The selective cause of an ancient adaptation. *Science*, *307*(5713), 1279–1282. https://doi.org/10.1126/science.1106974

1602

Figure Legends

1604

Figure 1. Genome-scale metabolic model reconstruction of *Xylella fastidiosa* through a pangenome based on the subspecies *pauca*, *multiplex*, *fastidiosa*, *morus* and *sandyi*. A) Pangenome analysis illustrating the distribution of core genes, genes shared between subspecies and unique genes. Each bar in the vertical bar chart represents the number of ortholog groups shared by the strains indicated below the chart. B) Summary of the features (reactions, metabolites and genes) from the *X. fastidiosa* metabolic core (Xfcore) reconstructed in this work. C) Predicted flux distribution through the core metabolism, using FBA, in a minimal medium with glutamine as the sole carbon and nitrogen source, achieving a specific growth rate of 0.41 h^{-1}. Metabolic fluxes (mmol·gDW^{-1}·h^{-1}) are portrayed numerically.

1615

Figure 2. Evaluation of six synthetic defined growth media based on Xfcore model predictions. Total bacterial growth measured by the standardized area under the growth progress curve (sAUGPC) (A) and biofilm formation (B) of seven *X. fastidiosa* strains grow at 28 °C for 6 days. PD3 medium is included as a positive control. Statistical differences between each synthetic defined medium and the PD3 control were assessed using Wilcoxon rank-sum tests. Significance levels of comparisons are displayed above each line connecting group pairs (ns: $P>0.05$, *: $P≤0.05$, **: $P≤0.01$, ***: $P≤0.001$, ****: $P≤0.0001$).

1623

Figure 3. Predicted acetate assimilation by the Xfcore model. Flux distribution through the *X. fastidiosa* core metabolism, using FBA, in medium with acetate as the sole carbon and glutamic acid and ammonium as nitrogen source, achieving a specific growth rate of 1.14 h^{-1}. Metabolic fluxes (mmol·gDW^{-1}·h^{-1}) are portrayed numerically. The alternative metabolic strategy for converting acetate into biomass combines two previously described modules: half of the 3-hydroxypropionate (3HP) bicycle (Herter et al., 2002) and the methylcitrate pathway (Brämer & Steinbüchel, 2001; Dolan et al., 2018) for propionyl-CoA assimilation. The enzymes and genes involved in the metabolic steps (1-11) of this pathway are in Supplementary Table S4. Abbreviations for metabolites participating in the 3HP pathway and the methylcitrate pathway: MCOA (malonyl CoA), MSA (malonate semialdehyde or 3-oxopropanoate), 3HP (3-

hydroxypropionate), 3HPCOA (3-hrydroxypropionyl CoA), PP2COA (acryloyl CoA), PPCOA (propionyl CoA), 2MCIT (2-methylcitrate), 2MCACN (2-methylaconitate), MICT (2-methylisocitrate).

Figure 4. Polyamine production by *Xylella fastidiosa.* A) Trade-off between biomass and polyamine production using the Xcore model. This figure illustrates the inverse relationship between growth (biomass accumulation) and the production of polyamines (putrescine, spermidine, and spermine) when one of the polyamines is completely absent. Simulations were performed using a minimal medium where glutamine served as the sole carbon source (with an uptake flux of 15.665 mmol·gDW^{-1}·h^{-1}). For each data point, the biomass reaction was maximized while the production fluxes for the polyamines (putrescine, spermidine, and spermine) were incrementally forced to the values represented by axis in flux units, mmol·gDW^{-1}·h^{-1}. A minimal threshold of 0.1608 mmol·gDW^{-1}·h^{-1} was established for the consideration of growth. B) In vitro production of putrescine, spermidine and spermine across different sample types (cell pellet from biofilm, CP-B; cell pellet from planktonic growth, CP-P; and culture supernatant, SP). Samples were obtained from three *Xylella fastidiosa* strains (De Donno, ESVL and Temecula1) grown at 28 °C in PD3 medium for 5 and 12 days, with four replicates for each sample and time point. Concentrations (in ng·mL^{-1}) are visualized with a pseudo-logarithmic scale. Statistical differences between time points within each sample type and strain were assessed using Wilcoxon rank-sum tests. Only significant results are indicated (∗: $P \leq 0.05$).

Figure 5. Trade-off between biomass and virulence factor production in *X. fastidiosa.* This figure illustrates the inverse relationship between growth (biomass accumulation) and the production of: LesA protein, fastidian gum EPS, and putrescine (A), spermidine (B) or spermine (C) using the Xfcore metabolic mode, when one of the three compounds is completely absent. Simulations were performed using a minimal medium where glutamine served as the sole carbon source (with an uptake flux of 15.664 mmol·gDW^{-1}·h^{-1}). For each data point, the biomass reaction was maximized while the production fluxes for the virulence factors (LesA, EPS, polyamine) were incrementally forced to the values represented by the axis in flux units, mmol·gDW^{-1}·h^{-1}. A minimal threshold of 0.1608 mmol·gDW^{-1}·h^{-1} was established for the consideration of growth.

Table legends

Table 1. Summary of *X. fastidiosa* strains used for the pangenome reconstruction and genome-scale metabolic modeling.

Table 2. Comparison between in silico growth predictions and in vitro phenotypic profiles of *X. fastidiosa*. Experimental growth across various substrates, including carbohydrates, organic acids, lipids, and amino acids, was obtained from Biolog PM plates. In silico growth was predicted through FBA using the Xfcore model as described in Material and Methods.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Table 1.

Identifier	Subspecies	Host	Location	Disease[a]	ID RefSeq
Temecula1	*fastidiosa*	*Vitis vinifera*	California, USA	PD	NC_004556.1
M23	*fastidiosa*	*Prunus dulcis*	California, USA	PD, ALS	NC_010577.1
MUL0034	*morus*	*Morus alba*	California, USA	MLS	NZ_CP006740.1
M12	*multiplex*	*Prunus dulcis*	California, USA	ALS	NC_010513.1
9a5c	*pauca*	*Citrus sinensis*	São Paulo, Brazil	CVC	NC_002488.3
U24d	*pauca*	*Citrus sinensis*	São Paulo, Brazil	CVC	NZ_CP009790.1
J1a12	*pauca*	*Citrus sinensis*	São Paulo, Brazil	Non-pathogenic	NZ_CP009823.1
Pr8x	*pauca*	*Prunus domestica*	São Paulo, Brazil	LS	NZ_CP009826.1
3124	*pauca*	*Coffee arabica*	São Paulo, Brazil	CLS	NZ_CP009829.1
Hib4	*pauca*	*Hibiscus fragilis*	Distrito Federal, Brazil	LS	NZ_CP009885.1
Salento-2	*pauca*	*Olea europaea*	Puglia, Italy	OQDS	NZ_CP016610.1
De Donno	*pauca*	*Olea europaea*	Puglia, Italy	OQDS	NZ_CP020870.1
Ann-1	*sandyi*	*Nerium oleander*	California, USA	OLS	NZ_CP006696.1
ESVL	*multiplex*	*Prunus dulcis*	Alicante, Spain	ALS	NZ_CP099516.1
IVIA5901	*multiplex*	*Prunus dulcis*	Alicante, Spain	ALS	NZ_CP047134.1
XYL1961/18	*pauca*	*Olea europaea var. europaea*	Ibiza, Spain	OQDS	CP171775.1
XYL468	*multiplex*	*Olea europaea var. sylvestris*	Mallorca, Spain	OQDS	NZ_CP136966.1
CFBP8418	*multiplex*	*Spartium junceum*	Corse, France	LS	NZ_CP136979.1

[a] PD: Pierce's Disease; ALS: Almond Leaf Scorch; MLS: Mulberry Leaf Scorch; CVC: Citrus Variegated Chlorosis; CLS: Coffee Leaf Scorch; OQDS: Olive Quick Decline Syndrome; OLS: Oleander Leaf Scorch; LS: Leaf Scorch.

Table 2.

Substrate	Growth assessment	
	In silico[a]	Experimental
Carbohydrates and derivatives		
D-Ribose	+	+
D-Xylose	+	+
L-Arabinose	-	+
L-Lyxose	-	+
2-Deoxy-D-Ribose	-	+
D-Arabinose	n.a.	+
D-Fructose-6-Phosphate	+	+
D-Psicose	-	+
D-Tagatose	-	+
L-Sorbose	n.a.	+
L-Rhamnose	n.a.	+
Inulin	n.a.	+
Pectin	n.a.	+
D-Glucosamine	+	+
3-0-β-D-Galactopyranosyl-D-Arabinose	n.a.	+
Glucuronamide	n.a.	+
L-Galactonic Acid-γ-Lactone	n.a.	+
5-Keto-D-Gluconic Acid	n.a.	+
Organic acids		
∝-Keto-Glutaric Acid	+	+
∝-Keto-Butyric Acid	-	+
∝-Keto-Valeric Acid	n.a.	+
Pyruvic acid	+	+
Acetoacetic Acid	-	+

Acetic Acid	+	+
Oxalic Acid	n.a.	+
Oxalomalic Acid	n.a.	+
Dihydroxy Acetone	-	+
Sorbic Acid	n.a.	+
Lipids and fatty acids		
Capric acid	n.a.	+
Tween 20	n.a.	+
Tween 40	n.a.	+
D,L-Carnitine	n.a.	+
Amino acids and amines		
Tyramine	n.a.	+
Sec-Butylamine	n.a.	+

[a] (+) and (-) indicate, respectively, growth/no growth in the corresponding simulation or experimental result. (n.a.) indicates metabolites not represented in the metabolic reconstruction or not available to model.