Leishmania major co-opts IL-7 feedback in monocytes to suppress CD4+ T-cell immunity

Yan Fu[1,2], Christophe Toussaint[3], Ina Sauerland[1,2], Elisabeth Reiß[1,2], Lars Philipsen[4], Patricia

Gintschel[1,2], Laura Knop[2,5], Andreas Braun[2,6], Bastian Kruse[2,6], Leon-Alexander Dewitz[1,2], Jonas

Amore[1,2], Vladyslava Dovhan[2,5], Sönke Weinert[4], Nouria Jantz-Naeem[1,2], Iris Baars[1,2], Juliane Mohr[1,2],

Julia Prein[1,2], Cora Schwendele[1,2], Philipp Bruno[2,5], Martin Böttcher[7], Nicole Gröger[8], Hannah Van

Hove[9], Melanie Greter[9], Burkhard Becher[9], Sascha Kahlfuß[2,5], Burkhart Schraven[2,5], Robert Geffers[10],

Thomas Tüting[2,6], Nicole Joller[11], Stephan Fricke[2,5,12], Antoine-Emmanuel Saliba[3,13,14]*, Thomas

Schüler[2,5,14]*, Andreas J. Müller[1,2,14,15]*

[1] Experimental Immunodynamics, Institute for Clinical Immunology and Cell Therapeutics, Medical Faculty,
Otto-von-Guericke University Magdeburg, Magdeburg, Germany.

[2] Health Campus Immunology, Infectiology and Inflammation (GCI3) and Center for Health and Medical
Prevention (CHaMP), Otto-von-Guericke-University, Magdeburg, Germany

[3] Helmholtz Institute for RNA-based Infection Research, Helmholtz Centre for Infection Research, Würzburg,
Germany.

[4] Platform for Multiparametric Bioimaging and Cytometry (MPBIC), Medical Faculty, Otto-von-Guericke-
University, Magdeburg, Germany.

[5] Institute for Clinical Immunology and Cell Therapeutics, Medical Faculty, Otto-von-Guericke University
Magdeburg, Magdeburg, Germany

[6] Department of Dermatology, Medical Faculty, Otto-von-Guericke University Magdeburg, Magdeburg,
Germany.

[7] Department of Hematology, Oncology, and Cell Therapy, Medical Faculty, Otto-von-Guericke University
Magdeburg, Magdeburg, Germany.

[8] Department of Zoology/Developmental Neurobiology, Institute of Biology, Otto von Guericke University
Magdeburg, Magdeburg, Germany.

[9] Institute of Experimental Immunology, University of Zurich, Switzerland

[10] Helmholtz Centre for Infection Research, Braunschweig, Germany.

[11] Department of Quantitative Biomedicine, University of Zurich, Zurich, Switzerland.

[12] Fraunhofer Institute for Cell Therapy and Immunology, Leipzig, Germany.

[13] University of Würzburg, Faculty of Medicine, Institute of Molecular Infection Biology, Würzburg, Germany.

[14] These authors contributed equally

[15] Lead contact

* Correspondence: emmanuel.saliba@helmholtz-hiri.de (A.E.S.); thomas.schueler@med.ovgu.de (T.S.);
andreas.mueller@med.ovgu.de (A.J.M.)

Summary

Effective immunity against infections requires a tight balance between pro- and anti-inflammatory

pathways to ensure pathogen clearance while preserving tissue integrity. Intracellular pathogens such as

Leishmania major can exploit host-cell immunosuppressive functions, yet the underlying molecular and

cellular interactions remain incompletely understood. Here we report a monocyte-derived,

hypo-inflammatory PDPN$^+$IL-7R$^+$ subset that harbours low-proliferating parasites, inhibits effector

CD4$^+$ T-cell control of intracellular pathogens, and therefore perpetuates infection. This

immunosuppression is mediated by fibroblast-derived IL-7, which is up-regulated in response to IFNγ.

Thus, our data suggest a cytokine-dependent feedback switch through which IL-7 activates

immunosuppressive monocyte-derived cells when local IFNγ production exceeds a critical level. This

dampens effector CD4$^+$ T cell functions at the expense of pathogen persistence. Targeting this feedback

loop may yield novel immunotherapeutic strategies that enhance immune responses while maintaining

tissue integrity.

Introduction

Inflammation induces the recruitment of large numbers of monocyte-derived myeloid cells which phagocytose invading pathogens, deliver antimicrobial effector molecules and thus limit pathogen load. However, such cells also exert regulatory functions required for the resolution of inflammation and tissue repair (Guilliams et al., 2018). This is of critical importance for host survival, since unrestrained immune responses endanger tissue integrity and organ function (Ortega-Gomez et al., 2013; Postat et al., 2018; Sugimoto et al., 2016). Intracellular pathogens infecting monocyte-derived cells can interfere with anti-microbial effector functions of their host, thereby establishing long-term persistence and chronic disease (Carvalho et al., 2023; Collins-McMillen et al., 2017; Drewry et al., 2019; Mariotti et al., 2004; Stapels et al., 2018). Furthermore, monocyte-derived cells can acquire an immunosuppressive phenotype that counter-regulates effector functions of other immune cells, or shifts immune responses towards a less inflammatory phenotype (Gundra et al., 2017; Zhang et al., 2022). Elucidating and manipulating these diverse functions holds the potential to improve treatment for infectious diseases (Channappanavar et al., 2016; Dorhoi et al., 2014; Wallis et al., 2023) and cancer (Kruse et al., 2023; Mantovani et al., 2022) but the mechanisms by which such functions are induced and deployed remain incompletely understood.

Upon arrival in the tissue, monocytes adapt their functional properties in response to complex environmental cues, including inflammation- and pathogen-derived signals. This context-dependent adaptation is crucial for a balanced response ensuring pathogen elimination and tissue protection at the same time (Veglia et al., 2021). Thus, within a population of activated monocyte-derived cells, a mixture of pathways triggering inflammatory, antimicrobial, and tissue remodeling functions must be activated (Sanin et al., 2022). Single-cell RNA sequencing (scRNAseq) has been instrumental in dissecting monocytes and their derivatives into phenotypically and functionally distinct subpopulations during infection by intracellular bacterial (Bryson et al., 2019; Hoffman et al., 2021), viral (Baasch et al., 2021) or parasitic (Patir et al., 2020) pathogens, as well as in the tumor microenvironment (Cheng et al., 2021; Mulder et al., 2021). Some of these subpopulations have the capacity to critically divert the course of the disease, even if present only at a small number compared to the total monocyte population

(Hoffmann et al., 2022; Zhang et al., 2022). In the course of an infection, the level of complexity is

further increased by a significant diversity of virulence, growth rate and invasiveness of individual

pathogens, thereby additionally modulating the functional properties of the infected cell type (Avraham

et al., 2015; Pisu et al., 2021; Saliba et al., 2016).

To address the differential monocyte-derived cell functions in the context of the diverse pathogen states,

we employed *Leishmania major* (*L. major*) infection as a physiological model for infection-induced

recruitment and activation of monocytes (Heyde et al., 2018; Romano et al., 2017). In this model system,

monocyte-derived cells represent the preferential host cell type for intracellular pathogen proliferation

and, at the same time, are critical for controlling the parasite through expression of the inducible nitric

oxide synthase (iNOS) in response to T cell-derived Interferon-γ (IFNγ) (De Trez et al., 2009; Leon et

al., 2007; Muller et al., 2012; Olekhnovitch et al., 2014). Permanent monocyte recruitment to and

maturation at the site of infection results in a large phenotypic and functional diversity. Importantly, the

proliferation state of the pathogen correlates with the phenotype and function of the infected monocyte-

derived cells (Formaglio et al., 2021; Heyde et al., 2018; Mandell and Beverley, 2017). Also, the

composition and polarization of these cell populations has been shown to critically impact on the

deployment of the anti-*Leishmania* response, e.g. by entertaining type II immune cell circuitries (Lee et

al., 2023), or by modulating the reservoir of monocyte-derived cells permissive for pathogen

proliferation (Carneiro et al., 2020). However, the cellular and molecular interactions which enable

monocyte-derived populations to impact prolongation versus control of the infection remain to be fully

elucidated.

Here, we combined an *in vivo* pathogen proliferation reporter system with scRNAseq to characterize the

function of distinct monocyte-derived subpopulations in the context of their interactions with

intracellular pathogens. With the help of this approach, we identified a subset of Podoplanin (PDPN)-

and Interleukin-7 receptor (IL-7R) expressing cells which are derived from monocytes and preferentially

harbor low-proliferating *L. major*. This PDPN[+]IL-7R[+] population exhibited a hypo-inflammatory

phenotype and suppressed effector functions of *L. major*-specific CD4[+] T cells in response to fibroblast-

derived IL-7. IL-7 production was dependent on IFNγ release in the course of the adaptive immune

response against *L. major*. Interference with IL-7R signaling or depletion of these cells, as well as *Il7*

gene inactivation in fibroblasts, resulted in improved effector T cell responses and subsequent pathogen

clearance.

In summary, we provide evidence for an as-yet unknown feedback loop which dampens T cell responses

as a result of the inflammation-associated up-regulation of fibroblast-derived IL-7, and the subsequent

immunosuppressive activity of IL-7R$^+$ monocyte-derived cells. This mechanism can favour the

pathogen to establish persistence and could therefore represent a new target for immunotherapeutic

approaches.

Results

PDPN⁺IL-7R⁺ phagocytes infected with low proliferating *L. major* exhibit a hypo-inflammatory

phenotype

During local immune responses, tissue- and inflammation-derived signals modulate the phenotype and

function of recruited immune cells (Romano et al., 2017; Saliba et al., 2016; Szabo et al., 2021; Yamada

et al., 2020). However, dissecting these environmental cues has remained challenging. We and others

have shown previously that intracellular *L. major* proliferation differs considerably between infected

monocyte-derived cells recruited to the infection site (Heyde et al., 2018; Mandell and Beverley, 2017).

We therefore utilized pathogen proliferation rate as a proxy for host-pathogen interaction cues which

define distinct monocyte-derived cell functions *in vivo*. For this, we employed an *in vivo* proliferation

assay, which relies on genetically modified *L. major* parasites expressing the photoconvertible green

fluorescence protein mKikume (Lm^{SWITCH}). Upon illumination, mKikume molecules convert from green

to red. Consequently, red molecules are preserved in low-proliferating parasites (Lm^{lo}), but are diluted

by newly synthesized green molecules in high-proliferating *L. major* (Lm^{hi}) (Fig. 1A) (Formaglio et al.,

2021).

To characterize the transcriptome of cells which harbour Lm^{lo} and Lm^{hi}, we infected C57BL/6 (B6) mice

with 2×10^6 Lm^{SWITCH} in the ear. At 19 days post infection (p.i.), the parasites were photoconverted and

CD45⁺ leukocytes were MACS purified 2 days later from the infected ear. Subsequently, single-cell

sorting of uninfected, Lm^{lo}- or Lm^{hi}-containing monocyte-derived cells was performed (Fig. 1B). Both

CD11c^{int-hi} and Ly6c⁺ were selected to cover a broad spectrum of recruited monocyte-derived cells (Fig.

S1A) (Heyde et al., 2018; Olekhnovitch et al., 2014). Lm^{lo}- or Lm^{hi}-infected (termed Lm^{lo}Mo or Lm^{hi}Mo,

respectively) as well as uninfected (termed UMo) single monocyte-derived cells were separated using

index-sorting into individual wells (Fig. 1C and S1B). ScRNAseq using the SMARTseq2 protocol

(Picelli et al., 2013) revealed specifically and consistently high *L. major* transcript counts in cells sorted

as infected, but not uninfected (Fig. 1D). In total, 88% of the sorted cells had passed quality control

based on mitochondria read counts and percentage of ERCC (Fig. S1C, and see details in Methods).

Mapping of the sequenced cells sorted as Lm^{hi}Mo (green), UMo (black) and Lm^{lo}Mo (red) to a batch-

corrected Uniform Manifold Approximation and Projection (UMAP) representation (Fig. 1E, S1D, and

S1E) revealed that cells infected with Lm^{lo} exhibited a unique gene expression profile. In contrast,

uninfected cells and cells infected by Lm^{hi} were not distinguishable on the UMAP representation.

Unsupervised clustering using the Leiden algorithm identified three distinct clusters of cells referred to

as Cluster 1, 2 and 3, with Cluster 3 consisting mainly of Lm^{lo}Mo, (Fig. 1F; Table S1).

Gene ontology (GO) function terms revealed a differential regulation of mainly immune system-related

processes for Cluster 3, but not Cluster 1 and 2. In particular, biological process enrichment in Cluster

3 included immunosuppressive signaling pathways, but lowered immune-activating functions and

antigen processing as compared to Cluster 1 and 2 (Fig. S1F-G; Table S2-3). Regarding differentially

expressed genes (DEG) specific for Cluster 3, we found a strong reduction of the proinflammatory

cytokine gene $Il1b$, chemokines and class II MHC-related genes as compared to Cluster 1 and 2. In

contrast, cathepsins and immune checkpoint ligands were expressed specifically in Cluster 3, in line

with a hypo-inflammatory phenotype suggested by GO terms. Moreover, we found a specific expression

of $Pdpn$ in Cluster 3, which is reported to be involved in immunomodulatory functions of myeloid cells

(Hovd et al., 2024; Rayes et al., 2017). Additionally, $Il7r$ was strongly upregulated in Cluster 3. In

contrast, $Ccr2$ expression was virtually absent in this population (Fig. S1H).

As Cluster 3 consisted mainly of Lm^{lo}Mo, while Lm^{hi}Mo and UMo were evenly dispersed across Cluster

1 and 2 (see Fig. 1E), we defined the five most prominent populations for further analysis, i.e. Lm^{hi}

infected and uninfected cells in Cluster 1 (termed Lm^{hi}Mo1 and UMo1) and Cluster 2 (termed Lm^{hi}Mo2

and UMo2), as well as cells infected with Lm^{lo} in Cluster 3 (termed Lm^{lo}Mo3) (Fig. 1G). The gene

expression pattern specific for Cluster 3 (see Fig. S1H) was consistent with the comparison of Lm^{lo}Mo3

versus Lm^{hi}Mo1, UMo1, Lm^{hi}Mo2 and UMo2 (Fig. 1H). This notably included the significantly higher

expression of $Pdpn$ and $Il7r$, suggesting that PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3 constitute a hypo-inflammatory

cellular subset. These observations were validated by cells sorted according to their differential

expression of CD11c (high in Cluster 1), Ly6c (high in cluster 2, intermediate in infected cluster 1)

observed during index sorting, and CCR2 (high in cluster 2) observed in scRNAseq (Fig. S1I, and S1J).

Quantitative PCR (qPCR) of cells sorted based according to this strategy (Fig. S1K) reproduced the

differential gene expression of Lm^{lo}Mo3 observed in scRNAseq, including the specific expression of

Pdpn and *Il7r*, and the hypo-inflammatory phenotype indicated by lower expression of *Il1b*, *Il1r2* and

Ccl5 (Fig. S1I). Also, Lm^{lo}Mo3 exhibited the lowest levels of pro-IL-1β, CXCL9 protein among all five,

and the lowest IL-6 and IL-12 p35 protein among the infected cell subsets (Fig. 1J). Irrespective of

whether we gated Lm^{lo}Mo3 cells from the infection gated according to CD11c, Ly6C and CCR2 (see

Fig. S1K), or as CD11b⁺Ly6G⁻SiglecF⁻ monocytes gated for PDPN and IL-7R expression, Lm^{lo}Mo3,

PDPN⁺IL-7R⁺ monocytes, respectively, exhibited a low intracellular pathogen proliferation rate (Fig.

S1L-N). These data altogether suggested a PDPN⁺IL-7R⁺ phagocyte population which is infected with

low proliferating *L. major* and which exhibits a hypo-inflammatory phenotype.

IL-7R⁺PDPN⁺ Lm^{lo}Mo3 develop from bone-marrow derived CCR2⁺ monocytes and acquire their

phenotype in the infected tissue

To further characterize the IL-7R⁺PDPN⁺ Lm^{lo}Mo3, we analyzed mannose receptor (CD206) and FcγRI

(CD64) expression, which distinguished tissue-resident macrophages, dendritic cells as well as different

stages of inflammatory monocytes and monocyte-derived macrophages (Carneiro et al., 2020; Lee et

al., 2018; Lee et al., 2023) (Fig. 2A and S2A). Infected IL-7R⁺PDPN⁺ phagocytes consisted mainly of

Ly6C^int monocyte-derived macrophages, but hardly any tissue-resident and monocyte-derived dendritic

cells (TR/MoDC), nor tissue-resident macrophages (TRM) (Fig. 2B and S2B). The same was observed

when analyzing phagocytes infected with low proliferating *L. major* (Fig. 2B and S2C-D). Thus

independently of the gating strategy, IL-7R⁺PDPN⁺ Lm^{lo}Mo3 cells seemed to consist mainly of Ly6C^int-

^lo, monocyte-derived macrophages.

Having shown a monocyte-derived phenotype, we sought to determine the dynamics of IL-7R⁺PDPN⁺

Lm^{lo}Mo3 recruitment and turnover. For this, we employed optical time-stamping with mKikume-

transgenic mice, in which photoconversion allows the identification of cells recruited before and after a

distinct time point (Formaglio et al., 2021; Heyde et al., 2018). The mice were infected with *L. major*

expressing blue-fluorescent mTurquoise ($Lm^{Turquoise}$), and mKikume-expressing host cells in the ear were

photoconverted at 16 dpi, and analyzed by flow cytometry 5 days later (Fig. 2C). This allowed

determining IL-7R and PDPN expression in infected cells and to discriminate photoconverted, long

resident (recruited > 5 days before analysis) cells from short resident cells which had been recruited

after photoconversion (Fig. 2D and S2E). Importantly, PDPN$^+$IL-7R$^+$ cells were found nearly

exclusively in the long resident cell fraction, while all other monocyte-derived cells exhibited a higher

turnover rate and were recruited mainly after the photoconversion (Fig. 2D and E).

Because of their long tissue residence time, we aimed to formally demonstrate whether or not Lm^{lo}Mo3

derive from CCR2$^+$ monocytes. For this, we employed lethally irradiated mice reconstituted with Ccr2-

creERT2 (Croxford et al., 2015) x Ai14 (Rosa-CAG-LSL-tdTomato-WPRE) BM, in which $Ccr2$-

expressing cells permanently express red fluorescent tdTomato upon Tamoxifen treatment. Tamoxifen

was applied either before the infection in order to label cells recruited during the innate wave of

monocyte recruitment, or from day 11 p.i. in order to label monocytes recruited during the established

adaptive immune response (Carneiro et al., 2020) (Fig. 2F). Tamoxifen treatment induced robust

labeling of >75% of blood monocytes for several days (Fig. 2G and S2F). In the infected ear, all

monocyte-derived populations, including PDPN$^+$IL-7R$^+$, showed a large proportion of CCR2-dependent

tdTomato expressing cells if Tamoxifen was applied during the adaptive wave of monocyte recruitment,

but not if the innate wave of monocytes was labeled (Fig. 2H, S2G-H). Therefore, we concluded that

despite its low turnover rate as compared to other monocyte-derived cells, the PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3

population was derived from CCR2$^+$ inflammatory monocytes recruited to the site of infection.

PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3 dampen T cell responses against $L. major$ infection $in vivo$

Having shown IL-7R and PDPN expression for monocyte-derived cells in the infected ear, we assessed

whether other immune cell types at the infection site, in particular non-myeloid cells, might also include

PDPN$^+$IL-7R$^+$ double-positive cells (Fig. S3A-C). While as expected, IL-7R was expressed on several

leukocyte subsets, PDPN expression, and, consequently, PDPN and IL-7R double-expression, was

limited to monocyte-derived, in particular infected, cells (Fig. 3A).

Since all IL-7R$^+$ monocyte-derived cells co-expressed PDPN, we next employed a $Pdpn$-cre based BM

reconstitution approach to specifically abolish IL-7R in these cells (Fig. 3B). For this, we lethally

irradiated WT recipient mice, reconstituted them with either $Pdpn$-cre $\times$ $Il7r^{fl/fl}$ or littermate BM, and

infected them with Lm^{DsRed} 10 weeks after reconstitution. Pathogen control and effector T cell activation

were analyzed at week 3 p.i. We found that IL-7R was significantly diminished in Lm^{lo}Mo3, but not in

lymph node T cells, confirming monocyte-specific gene deletion of *Il7r* (Fig. 3C and S3D). Strikingly,

the *Pdpn-cre* dependent *Il7r* knockout on BM-derived cells significantly reduced *L. major* infected

monocytes and pathogen burden (Fig. 3D and S3E). Furthermore, IFNγ production in effector CD4$^+$ T

cells within the infected ear tissue was elevated (Fig. 3E and S3F), suggesting that disruption of IL-7R

signaling in monocyte-derived cells at the site of infection enhances pathogen control by the adaptive

immune response. We were able to reproduce the enhanced pathogen control in a different experimental

system by reconstituting irradiated recipients with $Rag1^{-/-} \times Il7r^{-/-}$ BM and adoptive transfer of WT CD4$^+$

(Fig. S3G). The resulting animals, in which IL-7R was exclusively present on lymphocytes, also

exhibited enhanced control of *L. major,* confirming the role of IL-7R signaling in non-lymphocyte

immune cells (Fig. S3H-I).

Enhanced pathogen control in the absence of IL-7R signaling in monocyte-derived cells could be due to

cell-intrinsic enhancement of intracellular pathogen control in individual cells devoid of IL-7R, or due

to cell-extrinsic effects. In order to address these possibilities, we generated mixed BM chimeras

reconstituted with 50% CD45.1 WT and 50% $Il7r^{-/-}$ (CD45.2) BM. Mice reconstituted with mixtures of

WT CD45.1 and WT CD45.2 BM served as controls (Fig. 3F). We found that infection rates (Fig. 3G),

the relative abundance, (Fig. 3H), and the parasite proliferation within (Fig. 3I) the different monocyte-

derived cell subsets were independent of the cellular *Il7r* genotype in the mixed BM chimeras.

Therefore, we concluded that IL-7R signaling in the IL-7R$^+$PDPN$^+$ Lm^{lo}Mo3 does not affect their

development, abundance, or cell-intrinsic defense function, but dampens pathogen control in a cell-

extrinsic manner.

Next, we aimed to assess the effect of the depletion of PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3 on the control of *L.*

major. For this, we crossed *Pdpn-cre* transgenic mice to iDTR mice expressing the Diphtheria Toxin

receptor (DTR) after Cre-mediated elimination of a DNA-STOP cassette (Buch et al., 2005). Using these

mice as donors, we created BM chimeras reconstituted with 100% *Pdpn-cre* × iDTR, as well as controls

reconstituted with mixed 50% WT (CD45.1) and 50% *Pdpn-cre* × iDTR (CD45.2) BM (Fig. 3J).

Diphtheria Toxin (DTX) treatment of mice reconstituted with *Pdpn-cre* × iDTR BM reduced

significantly the percentage PDPN$^+$ cells among infected phagocytes (Fig. 3K), indicating successful

depletion at the site of infection. Importantly, the percentage of infected cells, as well as *L. major* tissue

burden was significantly reduced in these animals as compared to non-depleted of only partially

depletable BM chimera reconstituted with 50% WT and 50% *Pdpn-cre* × iDTR BM (Fig. 3L and S3J).

Of note, this depletion was also associated with an elevated IFNγ production by CD4$^+$ effector T cells

(Fig. 3M and S3K). These results underlined an immunosuppressive function of PDPN$^+$IL-7R$^+$ *Lm*^{lo}Mo3

in promoting parasite persistence via the inhibition of CD4$^+$ T effector functions. Thus, we concluded

that IL-7R signaling in *Lm*^{lo}Mo3 ultimately resulted in a dampened T cell response against *L. major* and

impaired pathogen control.

IL-7 from IFNγ-induced fibroblasts establishes a feedback loop that limits the effector T cell

response in the skin

Stromal cells, particularly fibroblasts, have been shown to produce IL-7 in response to *L. major* infection

(Peduto et al., 2009). In line with this, we found that fibroblasts from infected *Pdpn-cre* × *Il7r*$^{fl/fl}$ BM

reconstituted tissues expressed more MHC-I and PD-L1 than controls, indicating their additional

activation due to the enhanced immune response (Figure 4A and S4A). The same was observable in

Pdpn-cre × iDTR BM chimeric mice in which PDPN$^+$ leukocytes were depleted (Fig. 4A and S4B). Of

note, we found that *Il7* gene expression was upregulated in the tissue upon *L. major* infection (Fig. 4B

and S4C). We therefore sought to elucidate if skin fibroblasts could represent the source of the IL-7

modulating the immunosuppressive function of PDPN$^+$IL-7R$^+$ *Lm*^{lo}Mo3. For this, we employed *Prrx-*

cre × *Il7*$^{fl/fl}$ mice, which lack *Il7* gene activity in fibroblasts (Knop et al., 2020). Quantitative PCR of

stromal cells sorted from the ear confirmed that *Prrx* expression was limited to fibroblasts (Fig. 4C),

which were also more abundant in the infected skin than vascular endothelial cells (Fig. 4D and S4D).

Additionally, *Il7* expression was significantly reduced in fibroblasts from infected *Prrx-cre* × *Il7*$^{fl/fl}$

mouse ear tissue, but not in any other stromal cell population, further confirming a fibroblast-specific

Il7 gene deficiency (Fig. 4E-F). This suggested that *Prrx-cre* × *Il7*$^{fl/fl}$ mice indeed lacked *Il7* primarily

in fibroblasts in the infected tissue, and that fibroblasts are present in high numbers as compared to

vascular endothelial cells as alternative producers of IL-7 (Hara et al., 2012). Importantly, *L. major*

clearance was enhanced in *Prrx-cre* × *Il7*$^{fl/fl}$ mice (Fig. 4G) and IFNγ production by CD4$^+$ T cells was

increased (Fig. 4H and S4E). This suggested that fibroblast-derived IL-7 was involved in the suppression

of the immune response against *L. major*.

IFNγ has been reported to induce IL-7 expression in skin fibroblasts (Dai et al., 2021). Therefore, we

hypothesized that enhanced IFNγ production by T cells may contribute to a fibroblast-associated

feedback loop, limiting the effectiveness of an initially productive T cell response. This may be a result

of IL-7 induction by fibroblasts finally leading to immunosuppression of T cells by PDPN+IL-7R+

*Lm*loMo3 and subsequent pathogen persistence. In line with this hypothesis, we had shown previously

that IFNγ from T cells alone is sufficient for *L. major* containment (Muller et al., 2012), and found T

cells represented the main IFNγ producing cell population (Fig. 5A and S5A-B). Also, fibroblasts

cultured in the presence of recombinant IFNγ exhibited increased *Il7* cytokine gene expression (Fig.

5B). This suggested that T cell-derived IFNγ could stimulate fibroblasts in the infectious environment

to produce IL-7.

In order to demonstrate the impact of IFNγ signaling on IL-7 production by fibroblasts in vivo, we

treated *L. major*-infected mice with IFNγ-specific blocking or isotype-matched control antibodies (Fig.

5C). In fibroblasts from the infected tissue of control mice, we observed a strong correlation between

Il7 gene expression and the cellular infection rate. In contrast, this correlation was lost after anti-IFNγ

treatment (Fig. 5D). Consequently, anti-IFNγ treatment reduced fibroblasts *Il7* expression in strongly

infected tissues (Fig. 5E), and decreased fibroblast activation (Fig. 5F and S5C). Therefore, we

concluded that the IFNγ produced during the T cell response against *L. major* induces IL-7 production

in fibroblasts, thereby activating immunosuppressive PDPN+IL-7R+ *Lm*loMo3 which subsequently blunt

pathogen clearance.

Anti-IL-7/IL-7R treatment results in an enhanced immune response against *L. major*

Our data suggested that targeting IL-7R signaling might offer a possibility to prevent

immunosuppression by PDPN+IL-7R+ *Lm*loMo3, thus enhancing effector T cell responses and reducing

pathogen burden. Sustained long-term treatment with anti-IL-7R antibody has been shown to impair T

cell responses (Colpitts et al., 2009). However, IL-7R has been reported to be critical for maintaining

mainly naïve and memory T cells, but less important for effector T cells (Capitini et al., 2009; Fry and

Mackall, 2002; Huster et al., 2004). Thus, IL-7/IL-7R-blockade could suspend immunosuppression by

PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3 when applied temporarily during the acute infection.

To test if such transient anti-IL-7/IL-7R treatment improves the control of $L.$ $major$ infections, we

injected $L.$ $major^{DsRed}$-infected mice with an anti-IL-7/IL-7R antibody cocktail after the establishment

of infection (Fig. 6A). As expected for IL-7-depleting conditions (Grabstein et al., 1993), the treatment

resulted in a decline of immature B cells in the BM and spleen, but not in mature B cells (Fig. S6A-B)

three days after antibody injection. We also observed a slight reduction in total CD4$^+$ T cells, while the

numbers of central memory, effector or regulatory T cells were unchanged in the draining lymph node

(Fig. S6C). This indicated that the short-term treatment did not profoundly affect the peripheral

lymphocyte homeostasis, or T cell effector populations. Importantly, anti-IL-7/IL-7R treatment resulted

within 3-5 days in strongly reduced $L.$ $major$ infection (Fig. 6B-C), enhanced IFNγ production by

effector T cells (Fig. 6D and S6D), and increased fibroblast activation (Fig.6E and S6E) in the skin. To

explore whether anti-IL-7/IL-7R blockade also increased the induction of IFNγ-dependent antimicrobial

effector functions (Muller et al., 2012), we determined iNOS expression in monocyte-derived cells

isolated from the site of infection and found that anti-IL-7/IL-7R treatment resulted in higher iNOS

expression in these cells (Fig.6F and S6F). Collectively, these data indicated that IL-7/IL-7R blockade

efficiently increased the adaptive immune response against established skin infections with $L.$ $major$.

In order to address whether impaired IL-7R signaling had a long-term effect on pathogen control, and

immune activation, we analysed the impact of anti-IL-7/IL-7R treatment several weeks after antibody

treatment. For this, $L.$ $major^{DsRed}$ infected mice were injected with anti-IL-7/IL-7R antibody or isotype

control at day 16 and 20 p.i., and analyzed at week 7 p.i. for pathogen control and immune activation

(Fig. 6G). Weekly measurement of the infected ear thickness indicated a transiently higher inflammation

in anti-IL-7/IL-7R injected versus control animals, suggesting that the treatment increased the amplitude

of the immune response (Fig. 6F). At week 7 p.i., $L.$ $major$ tissue burden was reduced compared to

isotype control-treated animals (Fig. 6I), indicating that although the local pathology seemed

comparable at this late time point, the transient anti-IL-7/IL-7R had a lasting effect on pathogen control.

Furthermore, the fraction of IFNγ-producing effector T cells was strongly increased in the draining

lymph nodes of anti-IL-7/IL-7R treated animals (Fig. 6J and S6G). These data suggested that the

transient blockade of IL-7/IL-7R and suspension of the immunosuppressive function of Lm^{lo}Mo3 in the

early phase of the response supported a long-lived CD4$^+$ T cell response and pathogen control.

Altogether, our data therefore indicate that the IL-7 produced by fibroblasts in response to IFNγ drives

PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3 cells blunting CD4$^+$ T-cell effector functions, and targeted manipulation of this

pathway can facilitate robust anti-*Leishmania major* immunity.

Discussion

Mechanisms that restrict overshooting effector functions are an important component of immune

responses to prevent immunopathological tissue damage (Formaglio et al., 2021; Mishra et al., 2013;

Postat et al., 2018; Sugimoto et al., 2016). As such, the effects of IFNγ, a main mediator of cellular

immune responses, are controlled on different levels, including the restriction of IFNγ-producing cells

(Badovinac et al., 2003) and inhibition of IFNγ signaling (Alexander et al., 1999; Marine et al., 1999).

Pathogens can benefit from such regulatory mechanisms to establish efficient infection (Collins-

McMillen et al., 2017; Dolgachev et al., 2014; Drewry et al., 2019; Mariotti et al., 2004; Navarro et al.,

2011). A better understanding of the cellular and molecular interactions underlying these

immunosuppressive functions would greatly facilitate the development of novel immunotherapeutic

approaches (Gocher et al., 2022; Wykes and Lewin, 2018). Such approaches have so far mainly focused

on T cell functions directly (Ellis et al., 2021), while the potential of monocyte-derived functions is only

beginning to be acknowledged (Kruse et al., 2023; Mantovani et al., 2022). We describe here a new self-

limiting mechanism by which increasing IFNγ levels initiate a feedback loop that ultimately prompts

IL-7R expressing monocyte-derived cells to prevent effector T cells from producing more IFNγ.

Interruption of this feedback loop increased effector T cell function and promoted pathogen clearance.

Our unbiased identification of a PDPN⁺IL-7R⁺ monocyte-derived subset that harbors low-proliferating

L. major and suppresses CD4⁺ T-cell effector function adds a new layer to the picture of myeloid cell

heterogeneity during chronic infection. The plasticity of myeloid cells at infection sites has been

intensely studied, with ample evidence for immunomodulatory functions, for example PD-L2⁺CD206⁺

monocyte-derived macrophages in the context of helminth infection (Gundra et al., 2017; Gundra et al.,

2014; Oyesola et al., 2023), or CD206⁺ tissue resident macrophages which promote type 2 immunity in

non-healing *L. major* Seidman strain infection (Lee et al., 2018; Lee et al., 2020; Lee et al., 2023).

Furthermore, CCR2⁺ monocytes have been shown adopt a dual role in the establishment of parasite-

permissive conditions and the activation of parasite control depending on their time point of recruitment

and exposure to the local cytokine milieu (Carneiro et al., 2020). Our data extend these concepts by

demonstrating that, downstream of IFNγ, stromal fibroblasts secrete IL-7 that engages IL-7R on a

PDPN$^+$ monocyte-derived population, driving these cells towards a hypo-inflammatory phenotype

which dampens CD4$^+$ T cell responses.

We clearly show that PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3 cells originate from CCR2$^+$ monocytes and have a

turnover rate of far more than five days, and do not express TRM markers such as CD206. Thus, we

conclude that the functional overlap with alternatively activated CD206$^+$ tissue-resident macrophages

described previously is at most partial, and that the PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3 adopt their phenotype upon

exposure to the infection site. In line with this, previous reports indicate that both macrophages and

monocytes can upregulate cell surface expression of PDPN and IL-7R in response to inflammatory

stimuli (Hou et al., 2010; Kerrigan et al., 2012; Yazdani et al., 2024).

IL-7R expression by myeloid cells has been shown to promote the development of tissue-resident

macrophages in early ontogeny in mice (Cool et al., 2020; Leung et al., 2019). In humans, monocyte-

derived lung macrophages have been reported to express IL-7R during their development under non-

inflammatory conditions (Evren et al., 2021), and upregulate IL-7R expression in response to

inflammation (Al-Mossawi et al., 2019; Hu et al., 2008). Consequently, IL-7R expressing myeloid cells

were observed in a variety of inflammatory and infectious diseases. For example, serum from patients

infected with *Mycobacterium tuberculosis* (Mtb) as well as Mtb products induces IL-7R on monocytes

(Harelimana et al., 2022). Furthermore, a role of IL-7R in monocytes in modulating the immune

response has been suggested in human patients of SARS COV2 infection, where their presence in the

blood was correlated with lower disease pathology (Zhang et al., 2022). Moreover, *PDPN* and *IL7R*

expression was found in monocyte-derived subpopulations of human head and neck carcinoma

(Hoffmann et al., 2022), and *IL7R* was found to be a signature gene for tumor-associated monocyte-

derived cell subsets in a variety of malignancies (Cheng et al., 2021; Mulder et al., 2021; Nalio Ramos

et al., 2022). Altogether, these findings suggest a more general role for IL-7R expressing monocyte-

derived cells in dampening immune responses in a variety of infectious diseases, or even beyond. With

the identification of a feedback loop fueled by IFNγ, which induces IL-7 production by activated

fibroblasts, we provide a multicellular interaction network for the immunosuppressive function of

PDPN$^+$IL-7R$^+$ Lm^{lo}Mo3. Of note, we also observed PDPN$^+$IL-7R$^-$ single-positive monocyte-derived

cells. These cells might represent an intermediate state, but, alternatively, might deploy

immunoregulatory functions independently of the described IL-7 dependent feedback switch, since not

only IL-7R, but also PDPN has been shown to be involved in immunoregulatory mechanisms (Xie et

al., 2020).

The enhancement of pathogen clearance in response to i) *Il7r* deficiency in or depletion of PDPN[+]IL-

7R[+] *Lm*[lo]Mo3, ii) fibroblast-specific *Il7* gene inactivation, and iii) IL-7/IL-7R blockade, strongly

suggests that IL-7 can mediate an immunosuppressive function. While other cell types could also

produce IFNγ to enhance pathogen control, T cells seem to be a main and sufficient IFNγ producer for

iNOS induction (Muller et al., 2012), and thereby a very likely target of the described feedback

mechanism. At first sight this appears counter-intuitive, since IL-7 stimulates T cell immunity at

multiple levels, such as thymic T cell development and the survival of naïve and memory T cells in the

periphery. However, this is not the case for effector T cells, which down-regulate IL-7R expression and

are thus insensitive to direct IL-7 effects (Bachmann et al., 2005; Huster et al., 2004; Kaech et al., 2003).

Importantly, endogenous IL-7 had previously been shown to suppress IFNγ production and to exacerbate

disease in a skin infection model of mice with *Schistosoma mansoni* (Wolowczuk et al., 1997).

Moreover, treatment with recombinant IL-7 can aggravate *L. major* infection in susceptible mice

(Gessner et al., 1995). In line with these reports, we found that short-term blockade of IL-7/IL-7R affects

the number of naïve T and B cells in secondary lymphoid organs, but not of effector T cells, whose IFNγ

production is increased, resulting in enhanced control of the pathogen. We have shown previously that

even slight manipulation of pathogen permissiveness at week 3 p.i. results in enhanced control later on

(Formaglio et al., 2021). In line with this, we observed an altered course of pathology, with stronger ear

swelling 2-3 weeks after anti-IL-7/IL-7R treatment. The treatment also resulted in more efficient

pathogen clearance on the long-term perspective, including stronger IFNγ production by T cells, which

indicates a long-lasting effect of the transient anti-IL-7/IL-7R administration at the peak of infection.

To specify the critical source of IL-7 mediating an immunosuppressing effect, we employed a

conditional *Il7* knockout mediated by *Prrx-cre*, which is active in the majority of skin fibroblasts around

birth (Leavitt et al., 2020; Liu et al., 2022), and is upregulated during wound healing (Lee et al., 2022).

Importantly, we show that *Il7* expression is affected by the conditional knockout only in fibroblasts, but

not any other stromal cell population sorted from the skin. The finding that *Prrx-cre* mediated *Il7* was

sufficient to enhance *L. major* clearance and effector CD4$^+$ T cell function therefore suggests that

fibroblasts are critical producers of IL-7 in the mechanism. Skin fibroblasts have been shown to increase

Il7 expression in response to inflammation in a variety of models, including the infection of the ear

dermis with *L. major* (Peduto et al., 2009), which we could confirm. IFNγ, which we could show to be

mainly produced by T cells in our system, is sufficient to induce *Il7* expression upon injection in the

skin of uninfected mice. Conversely, anti-IFNγ injection decreases *Il7* expression in a model of

autoimmune alopecia (Dai et al., 2021). Therefore, it is conceivable that IFNγ produced by *L. major*-

specific T cells (Muller et al., 2012) initiates an IL-7 mediated feedback from fibroblasts which

ultimately dampens pathogen clearance. This mechanism might constitute a feedback switch that

activates immunosuppressive monocyte-derived cells as soon as local IFNγ production in the course of

the T cell response exceeds a certain threshold.

Targeting immunoregulatory mechanisms carries the risk of inducing immune-related adverse events

(Yan et al., 2024), including pathological effects of unrestricted T cell functions (Byrne and Fisher,

2017; Johnson et al., 2016; Murakami et al., 2016). Of note, we did only observe a slight, transient

increase of inflammation, while pathogen control was long-lasting. Two explanations are likely for this

finding: First, efficient pathogen control could decrease pathogen burden-dependent inflammation

(Formaglio et al., 2021), thereby masking the increased immune activation related to impaired IL-7

signaling. Second, the effector molecule nitric oxide, which is likely produced at higher amounts in

response to the increased IFNγ, can restrict leukocyte recruitment (Mishra et al., 2013; Postat et al.,

2018), thus counteracting an even more severe pathology in the anti-IL-7/IL-7R treated animals.

Therefore, targeting IL-7 mediated immunosuppression might be especially well-suited for enhancing

immune reactions involving nitric oxide-mediated killing by monocyte-derived cells, as nitric oxide

seems to provide an additional layer of immunoregulation. Such monocyte-mediated killing has been

demonstrated not only in immunity against intracellular pathogens (Olekhnovitch et al., 2014), but also

during T cell therapy against cancer (Kruse et al., 2023).

Taken together, we describe a cytokine-dependent feedback loop that can restrict T cell activation in

response to excessive IFNγ production, a mechanism from which the pathogen can benefit to perpetuate

the infection. Targeting this mechanism to adjust the activation of effector T cell functions opens new

avenues for future immunotherapeutic approaches.

Methods

Ethics Statement

All animal experiments were reviewed and approved by the Ethics Committee of the Office for

Veterinary Affairs of the State of Saxony-Anhalt, Germany (permit license numbers 42502-2-1314 UNI

MD, 42502-2-1393 Uni MD, 42502-2-1575 Uni MD, 42502-2-1586 Uni MD, 42502-2-1615 Uni MD,

42502-2-1672 Uni MD, and 42502-2-1735 Uni MD) in accordance with the legislation of both the

European Union (Council Directive 499 2010/63/EU) and the Federal Republic of Germany (according

to § 8, Section 1 TierSchG, and TierSchVersV).

Parasites, tumor and mouse infections

L. major LRC-L137 V121 wild-type, DsRed-, and mKikume-expressing (*Lm*[SWITCH]) parasites were

described previously (Muller et al., 2013; Sorensen et al., 2003). *L. major* expressing the cyan

fluorescent protein mTurquoise (*Lm*[Turquoise]) was generated by targeted integration of the mTurquoise

gene into the recombinant RNA locus of L. major LRC-L137 V121. For this, the mTurquoise gene

(Goedhart et al., 2010) sequence was synthesized *de novo* (Eurofins Scientific) and cloned via BamHI-

BglII and NotI (New England Biolabs) into a pLEXSY-Hyg2 vector (Jena Bioscience). The resulting

plasmid was linearized usin SwaI (New England Biolabs) and electroporated into *L. major*. Stable

transfectants were selected with 30 mg/ml hygromycin B (Invitrogen). Single clones were obtained by

limiting dilution. All parasites were grown in M119 medium completed with 10% heat-inactivated fetal

calf serum, 0.1 mM adenine, 1 mg/ml biotin, 5 mg/ml hemin, and 2 mg/ml biopterin (all from Sigma)

and kept for a maximum of six passages.

Wild-type C57BL/6J and CD45.1 (B6.SJL-*Ptprc*[a]*Pepc*[b]/BoyJ, JAX stock #002014) mice, *Il7r*[-/-]

(B6.129S7-*Il7r*[tm1Imx]/J, JAX stock #002295,) (Peschon et al., 1994), and *Rag1*[-/-] (B6.129S7-*Rag1*[tm1Mom]/J,

JAX stock #002096) (Mombaerts et al., 1992) mice were purchased from The Jackson Laboratory (Bar

Harbor, ME) via Charles River (Sulzfeld, Germany). Bone marrow from Ccr2-creERT2

(C57BL/6NTac-*Ccr2*[tm2982(T2A-Cre7ESR1-T2A-mKate2)BB]) (Croxford et al., 2015) x Ai14(B6.Cg-

Gt(ROSA)26Sor[tm14(CAG-tdTomato)Hze]) (Madisen et al., 2010) mice was provided by Melanie Greter and

Burkhard Becher (University of Zürich, Switzerland). To obtain *Prrx-cre × Il7^{fl/fl}* animals, *Prrx-cre*

(B6.Cg-Tg(Prrx1-cre)1Cjt/J, JAX stock #005584) (Logan et al., 2002) mice were purchased from The

Jackson Laboratory and crossed to mice harboring floxed *Il7* alleles (*Il7^{fl}Il7^{fl}*), which had been

previously generated by removing a FRT flanked part of a "knockout-first" construct *Il7^{tm1a(EUCOMM)Wtsi}*

(provided by The European Conditional Mouse Mutagenesis Program (EUCOMM)) using a Flpo-

transgenic cross. To obtain *Pdpn-cre × Il7r^{fl/fl}* and *Pdpn-cre ×* iDTR animals, the sperm of Pdpn-cre

mice (C57BL/6N-Tg(Pdpn-icre)14Biat/Biat, EM#11673) (Onder et al., 2011) were purchased from

European Mouse Mutant Archive (EMMA) and used in artificial insemination procedures with *Il7r^{fl/fl}*

(Il7rtm1.1Asin/J, JAX stock #022143) (McCaughtry et al., 2012) females and ROSA26iDTR (CBy.B6-

Gt(ROSA)26Sortm1(HBEGF)Awai/J) (Buch et al., 2005) females, respectively. The animals were

housed and bred under specific-pathogen-free conditions in the central animal facility (ZTL) of the

Medical Faculty at Otto-von-Guericke-Magdeburg, and were used for the experiments in age-and-sex-

matched groups at an age of 8-14 weeks at the beginning of the experiments. Males were used for

experiments involving RNA sequencing, while both males and females were included in the other

experiments. No differences in the reported effects were observed between male and female animals.

For infection, $2x10^6$ stationary phase promastigotes were washed and resuspended in PBS and

subsequently injected in 10 µl into the ear dermis. Where indicated, Tamoxifen (Sigma-Aldrich) was

dissolved in olive oil (Sigma-Aldrich) at a concentration of 20 mg/ml and administered by i.p. injection

(100 mg/kg) for five consecutive days, either five days before the infection, or between days 11 through

15 p.i.

Antibody treatment

For anti-IL-7/IL-7R application, mice were intraperitoneally (i.p.) injected with 500 µg of anti-mouse

IL-7Rα (clone A7R34, Bio X Cell), which is non-depleting (Mai et al., 2014; Seddon et al., 2003), and

500 µg of anti-mouse IL7 (clone M25, Bio X Cell) per treatment (Table S4). Control mice were i.p.

injected with 500 µg of rat IgG2a isotype control (clone 2A3, Bio X Cell) and 500 µg of mouse IgG2b

isotype control (clone MPC-11, Bio X Cell) per treatment. For anti-IFNγ application, mice were i.p.

injected with 500 µg of anti-mouse IFNγ (clone R4-6A2, Bio X Cell) per treatment, while control mice

received 500 µg of rat IgG1 isotype control (clone HRPN, Bio X Cell) per treatment.

Adoptive T cell and BM transfer, irradiation and BM chimera generation

For adoptive transfer, CD4$^+$ T cells were isolated from axillary, brachial, cervical, and inguinal lymph

nodes as well as from the spleen using the CD4 isolation kit and negative selection MACS purification

(Miltenyi Biotech, Bergisch Gladbach, Germany) according to the manufacturer's instructions. 1×10^7

T cells per recipient were injected intravenously. For BM isolation, BM cells were flushed from tibia

and femur with ice-cold non-supplemented RPMI 1640 (PAN Biotec) and filtered through 100 micron

cell strainers. Cells were washed with non-supplemented PBS, and 4×10^7 cells (for BM reconstitution)

per recipient were resuspended in PBS and injected intravenously into the tail vein. For generating BM

chimeras or fully BM reconstituted mice, recipients were γ-irradiated with 9.5 gray before reconstitution

and infected 8 weeks later.

Photoconversion for proliferation measurement and optical time-stamping

Photoconversion of *Lm*^{SWITCH} parasites and cells in the mouse ear was performed with a 405 nm

wavelength, 665 mW/cm2 collimated high-power LED (Thorlabs). The ears of anesthetized mice were

fixed and illuminated from each side for 30 seconds at a distance of 20 cm and were analyzed after 48h

by flow cytometry.

Flow cytometry

Cells were harvested from digested ears as previously described (Heyde et al., 2018). Draining lymph

nodes or spleen were dissociated on a 70 µm cell strainer, and cells were washed with PBS. Surface

staining of cells was done by using the antibodies specified in table S4. For intracellular staining of IL-

1 beta (Pro-form), IL-6, IL-12 p35, CXCL9, Eomes, Foxp3, GATA3, RORgt, T-bet and IFNγ, the cell

suspension was fixed for 1hr at 4°C with fixation buffer (Biolegend), following permeabilization with

Perm/Wash solution (Biolegend) according to the manufacturer's instructions and stained with the

antibodies liested in table S4 prior to staining and analyzed on a Fortessa flow cytometer (BD

Biosciences). Samples with a content of less than 40% of CD11c-expressing cells among CD45[+] were

excluded from quantitative analysis. The mKikume fluorescence with/without photoconversion was

read out at 561 nm excitation and 610/20 nm emission, or 488 nm excitation and 530/30 nm emission,

respectively. An autofluorescence signal was recorded at 488 excitation and 695/40 nm emission. Data

were analyzed using the FlowJo software (FlowJo, LLC). qPCR of sorted cells was performed as

described in the supplemental materials and methods.

Single-cell index sorting and SmartSeq2 sequencing library preparation

For scRNAseq, mouse CD45 MicroBeads (Miltenyi Biotec) were used to isolate leukocytes prior to cell

surface staining according to manufacturer's instructions. In order to review the complete phenotype of

every single cell and associate every phenotype with each sorted single cell, index sorting was performed

on FACS ARIA III sorter (BD Biosciences). Single cells were sorted into different wells from a 96-well

plate (Costar) filled with 4.65 µl of Buffer RLT Plus (Qiagen), 0.25 µl of SUPERase-In RNase-Inhibitor

(Invitrogen), and 0.1 µl of ERCC spike-Mix1 (Invitrogen) pre-diluted 1:500000 in RNAse-free water.

For cell-population RNAseq, 50-100 cells per subpopulation per sample were sorted into the wells.

Sorted cells were immediately frozen and stored at -80 °C. Single-cell libraries were prepared using

Smart-seq2 protocol as previously described (Picelli et al., 2014) with minor modifications: Following

cDNA amplification, ACTB expression was measured in each cDNA library by qPCR. All cDNA

libraries were diluted aiming to an equivalent ct value for ACTB (average = 21.11, SD = 1.05) and were

subsequently used as input for a tagmentation-based protocol (Nextera XT, Illumina) using one quarter

of the recommended volumes, 10 min for tagmentation at 55°C (Saliba et al., 2016). Libraries were

quantified using Qubit dsDNA Hs Assay (Life Technologies) and fragment size of libraries was

analyzed using QIAxcel DNA high resolution Kit (Qiagen) and Bioanalyzer High Sensitivity DNA kit

(Agilent). Libraries were normalized to 2 nM and pooled. Particularly, 218 of single-cell libraries or 87

of cell-population libraries were sequenced for HiSeq PE200 (Illumina). Sequencing data was analysed

as described in the supplemental materials and methods.

Processing of RNA sequencing data

Raw fastq files were processed with Cutadapt (v3.7) (Martin, 2011) to remove poly-A tails and adapter

sequences from the SMART-seq and Nextera protocols. Low-quality bases (Q-score < 20) and reads left

with less than 25 bases after trimming were removed. Sequencing data were mapped to a reference

genome using STAR with default parameters (Dobin et al., 2013). The reference genome was built by

combining the mouse (GRCm38) and *Leishmania major* (ASM272v2) genomes as well as spike-ins

(ERCC92) and the Kikume Green-Red transcript sequences. Raw count matrices were generated from

the mapping results after read count quantification with featureCounts (Subread, v1.6.3) (Liao et al.,

2014). Two count matrices for 218 single cells and 87 cell-population libraries were created.

Quality control and normalization of scRNAseq data

Low-quality cells with more than 2% mitochondrial read counts, more than 20% ERCC spike-ins read

counts or less than 1000 detected genes were discarded. Only mouse genes with at least 5 read counts

detected in at least 10 cells were retained. Read counts for the 194 retained single-cell transcriptomes

were normalized with scNorm (Bacher et al., 2017). Information about the plate for technical biases

(*Condition*) and gene length (*WithinSample*) was considered during the normalization procedure.

Dimensional reduction and clustering of scRNAseq data

Downstream analysis of the single-cell dataset has been performed with the Seurat R package (v4.0)

(Hao et al., 2021). The normalized read count matrix was natural log-transformed and the top 1000

highly variables genes were identified (*FindVariableFeatures*). For dimensional reduction, principal

component analysis was performed (*RunPCA*) to capture the principal components (PCs) explaining the

highest amount of variation in the dataset. Batch effects between mice were corrected with Harmony

(v0.1.1). The top 8 PCs from the corrected PCA were projected into a 2D space by Uniform Manifold

Approximation Projection (*RunUMAP*) for data visualization. In the uncorrected data, we found that

cells from three infected ears were strongly separated from the rest of the cells by in UMAP 1 versus

UMAP 2 (see Fig. S1E). MACS-purified cells from these infected ears revealed a substantially lower

percentage of CD11c$^+$ cells than the remaining samples in other ears. While this war corrected for

scRNAseq, we excluded cells from samples with a CD11c$^+$ cell content of less than 40% from

experiments with functional readout.

A graph-based unsupervised machine learning procedure was used to identify groups of single cells with

similar transcriptomes, i.e. clusters. The top 8 PCs were used to build a KNN graph (*FindNeighbors*;

k = 10) followed by partition into clusters with the Leiden algorithm (Traag et al., 2019). We used the C

implementation for R of the Leiden algorithm (leidenbase v0.1.3, Trapnell Lab) with a resolution set at

0.08. Clustering results were used to label cells on the UMAP.

Differential expression testing and pathway analysis for single-cell data

Gene markers of each cluster were identified by a Wilcoxon Rank-Sum Test with Seurat

(*FindAllMarkers*). Only genes that were expressed in at least 10% of the cells from a given cluster, with

an absolute log2 fold-change > 0.25 and a Bonferroni adjusted p-value < 0.05 were considered as

differentially expressed.

Gene Set Enrichment Analysis (v4.3.2) was used for pathway analysis (Subramanian et al., 2005). DEG

testing was repeated over all genes by a Wilcoxon Rank-Sum Test for each cluster versus all others

through Seurat (*FindMarkers*). Gene-ranked lists were generated after converting all uncorrected p-

values to -log10(p-value) * fold-change direction. Ranked lists were used as input for GSEA preranked

with default parameters and Gene Ontology Biological Process gene sets. Pathways up- or down-

regulated with FDR < 0.05 were considered significant.

Quantitative PCR analysis

For validation of DEG identified in the scRNAseq, 50 cells per sample were sorted from each

subpopulation according to the gating strategy shown in Fig. S2C and S2D and lysed in 10µl lysis buffer

(RNeasy Plus lysis buffer, QIAGEN) containing 5% SUPERase-In RNase-Inhibitor 20U/µl

(Invitrogen). RNA extraction and reverse transcription were performed using the Smart-seq2 protocol

as previously described (Picelli et al., 2014). Primers used are specified in table S4.

For measuring *Il7* gene expression, total RNA was isolated from *L. major*-infected ear tissue using

TRIzol reagent (Invitrogen) and then was reverse-transcribed using the High-Capacity cDNA Reverse

Transcription Kit (Applied Biosystems) following the manufacturer's instructions. In order to correct

for the immune cell infiltrate which was expected to contain leukocytes expressing *Hprt*, but not *Il7*)

hprt CT values were corrected according to the mean difference in RNA extracted.

For measuring the *Il7* and *Prrx1* gene expression in fibroblasts, LEC and BEC, three replicates of 200

cells each were sorted from each mouse ear. RNA extraction and reverse transcription was performed

using the Smart-Seq2 protocol as described above. Quantitative PCR analysis was done using a SYBR

Green Master Mix (Applied Biosystems).

To run the quantitative PCR, 40 cycles of 95 °C (15 s), and 60 °C (60 s) followed by a dissociation

protocol 95 °C (15 s), 60 °C (20 s), and 95 °C (15 s) were performed on an ABI Prism 7000 Sequence

detection system (Applied Biosystems). Technical triplicates were performed for each sample.

Limiting dilution assay

L. major burden within ear tissue was measured using a limiting dilution assay. Quadruplicate samples

of each ear homogenate were serially diluted in 1:2 steps in *Leishmania* culture medium in 96-well

plates and incubated at 26 °C for 14 days. The initial parasite burden was calculated from the median of

the highest dilutions of the quadruplicates which exhibited parasite growth. In order to compensate

experiment-to-experiment variation, the values were normalized to the control groups for pooled data.

Proliferation analysis

The proliferation index of *L. major* for flow cytometry was calculated based on the MFI of

mKikumeGreen and mKikumeRed as described previously (Heyde et al., 2018). In brief, for visualizing

qualitative comparisons within the same sample using the FlowJo software, values were plotted as

647
$$C - 100 * \left(\frac{mKikume\ Red}{mKikume\ Green}\right)_{\text{cell}} \Big/ \left(\frac{mKikume\ Red}{mKikume\ Green}\right)_{\text{mean (all infected cells)}}$$

648   with C chosen between 100 and 250 and kept constant within the same sample for which the comparison

649   was made, and the factor 100 was introduced to analyze integer fluorescence values in FlowJo.

650   For inter-sample comparison, the proliferation index was calculated as

$$651 \qquad 1 - \left(\frac{mKikume\ Red}{mKikume\ Green} \right)_{\text{cell}} / \left(\frac{mKikume\ Red}{mKikume\ Green} \right)_{\text{mean (all infected cells)}}$$

and represented as a percent deviation from the total infected cell population within one sample or

imaged infection site.

Cell culture and stimulation

To generate a fibroblast culture, lymphoid stromal cells were isolated from peripheral lymph nodes of

C57BL/6J mice as described previously (Knop et al., 2020). Afterwards, cells from four mice were

pooled and seeded in a T25 flask in full medium (RPMI 1640 supplemented with 10% (v/v) FCS (FBS

Superior; Sigma) /1% (v/v) P/S (Gibco)/2 mM L-glutamine (Gibco)/1 mM sodium pyruvate (Gibco)/0.1

mM HEPES (Gibco)/50 μM 2-mercaptoethanol (Sigma). Cells were cultured for 24h at 37°C and 5%

CO_2 and non-attached cells were removed. When reaching confluence, cells were detached using

Trypsin/EDTA (Merck Biochrom) and seeded onto T75, and afterwards onto T175. Cells were detached

using Trypsin/EDTA and treated with biotinylated anti-CD45 (30-F11, Biolegend) and anti-CD31

(MEC13.3, Biolegend)-specific antibodies in degassed staining buffer (PBS/ 2 mM EDTA (Carl Roth)/

0,5% (w/v) BSA (AppliChem)) for 15min at 4°C. Cells were washed and resuspended in degassed PBS/

2 mM EDTA and CD45+ and CD31+ cells were depleted using Streptavidin MicroBeads (Miltenyi

Biotec), LS columns (Miltenyi Biotec) and MidiMACS separator (Miltenyi Biotec) according to

manufacturer's instructions. Cells were seeded in a T175 flask in full medium. When reaching

confluence, cells were detached using Accutase (Stempro, Gibco) and stained with anti-CD45.2-APC

(104, Biolegend), gp38-A488 (8.1.1, Biolegend) and CD31-PE/Cy7 (390, eBioscience)-specific

antibodies for 30min at 4°C in staining buffer containing purified anti-CD16/32 (2.4G2 ATCC® HB-

197TM). After washing with PBS/2 mM EDTA, cells were resuspended in full medium containing 2

μg/ml Ciprobay (Bayer) and CD45.2[-] gp38[+]CD31[-] cells were sorted using FACSARIA III (Becton

Dickinson). The sorted cells were seeded at a density of 7400 cells/cm^2 in full medium containing 2

μg/ml Ciprobay. After 48h, Ciprobay was removed and the fibroblasts were expanded until reaching

confluence in a T175 flask.

For IFN-γ stimulation, cells were detached using Accutase and were seeded at 40000 cells/24-Well in

500 μl full medium without or with 50 ng/ml recombinant mouse IFN-γ (R&D Systems). After 24h, the

cells were detached using Accutase and resuspended in TRIzol reagent (Invitrogen). RNA isolation,

reverse transcription and RT-qPCR were performed as described previously (Knop et al., 2020). The

following TaqMan® Gene Expression Assays (Thermo Fisher Scientific) were used according to the

manufacturer's instructions: Il7 (FAM-MGB probe Mm01295804 m1) and Hprt (FAM-MGB probe

Mm00446968 m1). Each sample underwent analysis in triplicates on qTOWER3 G (Analytic Jena).

Subsequently, relative quantification was performed according to the ΔCT method.

Statistical analysis

Statistical analysis was carried out with GraphPad Prism 8 (GraphPad Software, San Diego, CA, USA).

To compare multiple samples in pairwise analysis within datasets with more than two experimental

groups, one-way analysis of variance (ANOVA) was done for datasets that had passed a Shapiro-Wilk

normal distribution test, Kruskal-Wallis tests were performed for datasets with non-normal distribution.

Appropriate multiple comparison post-tests (Bonferroni's for multiple comparisons, Dunnet's for

multiple pairwise comparisons or multiple comparisons with a control group in ANOVA, Dunn's test

for Kruskal-Wallis analyses) were employed as indicated in the respective Figure legends. Two-group

comparisons were analyzed using two-sided, unpaired or paired t-tests for data with normal distribution

and Mann-Whitney tests for datasets for which a Shapiro-Wilk normal distribution suggested non-

normal distribution. Representation of the median or mean, with standard deviation in cases where not

all data points are shown individually, are indicated together with sample size in the Figure legends.

References

Al-Mossawi, H., Yager, N., Taylor, C.A., Lau, E., Danielli, S., de Wit, J., Gilchrist, J., Nassiri, I., Mahe,
E.A., Lee, W.*, et al.* (2019). Context-specific regulation of surface and soluble IL7R expression by an
autoimmune risk allele. Nat Commun *10*, 4575.

Alexander, W.S., Starr, R., Fenner, J.E., Scott, C.L., Handman, E., Sprigg, N.S., Corbin, J.E., Cornish,
A.L., Darwiche, R., Owczarek, C.M.*, et al.* (1999). SOCS1 is a critical inhibitor of interferon gamma
signaling and prevents the potentially fatal neonatal actions of this cytokine. Cell *98*, 597-608.

Avraham, R., Haseley, N., Brown, D., Penaranda, C., Jijon, H.B., Trombetta, J.J., Satija, R., Shalek,
A.K., Xavier, R.J., Regev, A., and Hung, D.T. (2015). Pathogen Cell-to-Cell Variability Drives
Heterogeneity in Host Immune Responses. Cell *162*, 1309-1321.

Baasch, S., Giansanti, P., Kolter, J., Riedl, A., Forde, A.J., Runge, S., Zenke, S., Elling, R., Halenius,
A., Brabletz, S.*, et al.* (2021). Cytomegalovirus subverts macrophage identity. Cell *184*, 3774-3793
e3725.

Bacher, R., Chu, L.F., Leng, N., Gasch, A.P., Thomson, J.A., Stewart, R.M., Newton, M., and
Kendziorski, C. (2017). SCnorm: robust normalization of single-cell RNA-seq data. Nat Methods *14*,
584-586.

Bachmann, M.F., Wolint, P., Schwarz, K., Jager, P., and Oxenius, A. (2005). Functional properties and
lineage relationship of CD8+ T cell subsets identified by expression of IL-7 receptor alpha and CD62L.
J Immunol *175*, 4686-4696.

Badovinac, V.P., Hamilton, S.E., and Harty, J.T. (2003). Viral infection results in massive CD8+ T cell
expansion and mortality in vaccinated perforin-deficient mice. Immunity *18*, 463-474.

Bryson, B.D., Rosebrock, T.R., Tafesse, F.G., Itoh, C.Y., Nibasumba, A., Babunovic, G.H., Corleis, B.,
Martin, C., Keegan, C., Andrade, P.*, et al.* (2019). Heterogeneous GM-CSF signaling in macrophages
is associated with control of Mycobacterium tuberculosis. Nat Commun *10*, 2329.

Buch, T., Heppner, F.L., Tertilt, C., Heinen, T.J., Kremer, M., Wunderlich, F.T., Jung, S., and Waisman,
A. (2005). A Cre-inducible diphtheria toxin receptor mediates cell lineage ablation after toxin
administration. Nat Methods *2*, 419-426.

Byrne, E.H., and Fisher, D.E. (2017). Immune and molecular correlates in melanoma treated with
immune checkpoint blockade. Cancer *123*, 2143-2153.

Capitini, C.M., Chisti, A.A., and Mackall, C.L. (2009). Modulating T-cell homeostasis with IL-7:
preclinical and clinical studies. J Intern Med *266*, 141-153.

Carneiro, M.B., Lopes, M.E., Hohman, L.S., Romano, A., David, B.A., Kratofil, R., Kubes, P.,
Workentine, M.L., Campos, A.C., Vieira, L.Q., and Peters, N.C. (2020). Th1-Th2 Cross-Regulation
Controls Early Leishmania Infection in the Skin by Modulating the Size of the Permissive Monocytic
Host Cell Reservoir. Cell Host Microbe *27*, 752-768 e757.

Carvalho, A., Ferraz, I.A., Hojo-Souza, N.S., Medeiros, F.A.C., Viana, L.A., Bartholomeu, D.C.,
Chaves, A.T., de Souza, T.M., Costa, E.S.M.F., Mendes, T.A.O.*, et al.* (2023). Chagas cardiomyopathy
is associated with a high susceptibility to T. cruzi infection in monocyte-derived macrophages and a
predominance of CD4(+)CD45RO(+) T-cells with immunoregulatory patterns. Acta Trop *237*, 106749.

Channappanavar, R., Fehr, A.R., Vijay, R., Mack, M., Zhao, J., Meyerholz, D.K., and Perlman, S. (2016).
Dysregulated Type I Interferon and Inflammatory Monocyte-Macrophage Responses Cause Lethal
Pneumonia in SARS-CoV-Infected Mice. Cell Host Microbe *19*, 181-193.

Cheng, S., Li, Z., Gao, R., Xing, B., Gao, Y., Yang, Y., Qin, S., Zhang, L., Ouyang, H., Du, P.*, et al.*
(2021). A pan-cancer single-cell transcriptional atlas of tumor infiltrating myeloid cells. Cell *184*, 792-
809 e723.

Collins-McMillen, D., Stevenson, E.V., Kim, J.H., Lee, B.J., Cieply, S.J., Nogalski, M.T., Chan, G.C.,
Frost, R.W., 3rd, Spohn, C.R., and Yurochko, A.D. (2017). Human Cytomegalovirus Utilizes a
Nontraditional Signal Transducer and Activator of Transcription 1 Activation Cascade via Signaling
through Epidermal Growth Factor Receptor and Integrins To Efficiently Promote the Motility,
Differentiation, and Polarization of Infected Monocytes. J Virol *91*.

Colpitts, S.L., Dalton, N.M., and Scott, P. (2009). IL-7 receptor expression provides the potential for
long-term survival of both CD62Lhigh central memory T cells and Th1 effector cells during Leishmania
major infection. J Immunol *182*, 5702-5711.

Cool, T., Worthington, A., Poscablo, D., Hussaini, A., and Forsberg, E.C. (2020). Interleukin 7 receptor
is required for myeloid cell homeostasis and reconstitution by hematopoietic stem cells. Exp Hematol
90, 39-45 e33.

Croxford, A.L., Lanzinger, M., Hartmann, F.J., Schreiner, B., Mair, F., Pelczar, P., Clausen, B.E., Jung,
S., Greter, M., and Becher, B. (2015). The Cytokine GM-CSF Drives the Inflammatory Signature of
CCR2+ Monocytes and Licenses Autoimmunity. Immunity *43*, 502-514.

Dai, Z., Wang, E.H.C., Petukhova, L., Chang, Y., Lee, E.Y., and Christiano, A.M. (2021). Blockade of
IL-7 signaling suppresses inflammatory responses and reverses alopecia areata in C3H/HeJ mice. Sci
Adv *7*.

De Trez, C., Magez, S., Akira, S., Ryffel, B., Carlier, Y., and Muraille, E. (2009). iNOS-producing
inflammatory dendritic cells constitute the major infected cell type during the chronic Leishmania major
infection phase of C57BL/6 resistant mice. PLoS Pathog *5*, e1000494.

Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and
Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.

Dolgachev, V.A., Yu, B., Sun, L., Shanley, T.P., Raghavendran, K., and Hemmila, M.R. (2014).
Interleukin 10 overexpression alters survival in the setting of gram-negative pneumonia following lung
contusion. Shock *41*, 301-310.

Dorhoi, A., Yeremeev, V., Nouailles, G., Weiner, J., 3rd, Jorg, S., Heinemann, E., Oberbeck-Muller, D.,
Knaul, J.K., Vogelzang, A., Reece, S.T., *et al.* (2014). Type I IFN signaling triggers immunopathology
in tuberculosis-susceptible mice by modulating lung phagocyte dynamics. Eur J Immunol *44*, 2380-
2393.

Drewry, L.L., Jones, N.G., Wang, Q., Onken, M.D., Miller, M.J., and Sibley, L.D. (2019). The secreted
kinase ROP17 promotes Toxoplasma gondii dissemination by hijacking monocyte tissue migration. Nat
Microbiol *4*, 1951-1963.

Ellis, G.I., Sheppard, N.C., and Riley, J.L. (2021). Genetic engineering of T cells for immunotherapy.
Nat Rev Genet *22*, 427-447.

Evren, E., Ringqvist, E., Tripathi, K.P., Sleiers, N., Rives, I.C., Alisjahbana, A., Gao, Y., Sarhan, D.,
Halle, T., Sorini, C., *et al.* (2021). Distinct developmental pathways from blood monocytes generate
human lung macrophage diversity. Immunity *54*, 259-275 e257.

Formaglio, P., Alabdullah, M., Siokis, A., Handschuh, J., Sauerland, I., Fu, Y., Krone, A., Gintschel, P.,
Stettin, J., Heyde, S., *et al.* (2021). Nitric oxide controls proliferation of Leishmania major by inhibiting
the recruitment of permissive host cells. Immunity *54*, 2724-2739 e2710.

Fry, T.J., and Mackall, C.L. (2002). Interleukin-7: from bench to clinic. Blood *99*, 3892-3904.

Gessner, A., Will, A., Vieth, M., Schroppel, K., and Rollinghoff, M. (1995). Stimulation of B-cell
lymphopoiesis by interleukin-7 leads to aggravation of murine leishmaniasis. Immunology *84*, 416-422.

Gocher, A.M., Workman, C.J., and Vignali, D.A.A. (2022). Interferon-gamma: teammate or opponent
in the tumour microenvironment? Nat Rev Immunol *22*, 158-172.

Goedhart, J., van Weeren, L., Hink, M.A., Vischer, N.O., Jalink, K., and Gadella, T.W., Jr. (2010).
Bright cyan fluorescent protein variants identified by fluorescence lifetime screening. Nat Methods *7*,
137-139.

Grabstein, K.H., Waldschmidt, T.J., Finkelman, F.D., Hess, B.W., Alpert, A.R., Boiani, N.E., Namen,
A.E., and Morrissey, P.J. (1993). Inhibition of murine B and T lymphopoiesis in vivo by an anti-
interleukin 7 monoclonal antibody. J Exp Med *178*, 257-264.

Guilliams, M., Mildner, A., and Yona, S. (2018). Developmental and Functional Heterogeneity of
Monocytes. Immunity *49*, 595-613.

Gundra, U.M., Girgis, N.M., Gonzalez, M.A., San Tang, M., Van Der Zande, H.J.P., Lin, J.D., Ouimet,
M., Ma, L.J., Poles, J., Vozhilla, N.*, et al.* (2017). Vitamin A mediates conversion of monocyte-derived
macrophages into tissue-resident macrophages during alternative activation. Nat Immunol *18*, 642-653.

Gundra, U.M., Girgis, N.M., Ruckerl, D., Jenkins, S., Ward, L.N., Kurtz, Z.D., Wiens, K.E., Tang, M.S.,
Basu-Roy, U., Mansukhani, A.*, et al.* (2014). Alternatively activated macrophages derived from
monocytes and tissue macrophages are phenotypically and functionally distinct. Blood *123*, e110-122.

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W.M., 3rd, Zheng, S., Butler, A., Lee, M.J., Wilk, A.J.,
Darby, C., Zager, M.*, et al.* (2021). Integrated analysis of multimodal single-cell data. Cell *184*, 3573-
3587 e3529.

Hara, T., Shitara, S., Imai, K., Miyachi, H., Kitano, S., Yao, H., Tani-ichi, S., and Ikuta, K. (2012).
Identification of IL-7-producing cells in primary and secondary lymphoid organs using IL-7-GFP
knock-in mice. J Immunol *189*, 1577-1584.

Harelimana, J.D., Ahor, H.S., Benner, B., Hellmuth, S., Adankwah, E., Minadzi, D., Aniagyei, W.,
Lamptey, M.N.K., Arthur, J., Yeboah, A.*, et al.* (2022). Cytokine-induced transient monocyte IL-7Ra
expression and the serum milieu in tuberculosis. Eur J Immunol *52*, 958-969.

Heyde, S., Philipsen, L., Formaglio, P., Fu, Y., Baars, I., Hobbel, G., Kleinholz, C.L., Seiss, E.A., Stettin,
J., Gintschel, P.*, et al.* (2018). CD11c-expressing Ly6C+CCR2+ monocytes constitute a reservoir for
efficient Leishmania proliferation and cell-to-cell transmission. PLoS Pathog *14*, e1007374.

Hoffman, D., Tevet, Y., Trzebanski, S., Rosenberg, G., Vainman, L., Solomon, A., Hen-Avivi, S., Ben-
Moshe, N.B., and Avraham, R. (2021). A non-classical monocyte-derived macrophage subset provides
a splenic replication niche for intracellular Salmonella. Immunity *54*, 2712-2723 e2716.

Hoffmann, C., Noel, F., Grandclaudon, M., Massenet-Regad, L., Michea, P., Sirven, P., Faucheux, L.,
Surun, A., Lantz, O., Bohec, M.*, et al.* (2022). PD-L1 and ICOSL discriminate human Secretory and
Helper dendritic cells in cancer, allergy and autoimmunity. Nat Commun *13*, 1983.

Hou, T.Z., Bystrom, J., Sherlock, J.P., Qureshi, O., Parnell, S.M., Anderson, G., Gilroy, D.W., and
Buckley, C.D. (2010). A distinct subset of podoplanin (gp38) expressing F4/80+ macrophages mediate
phagocytosis and are induced following zymosan peritonitis. FEBS Lett *584*, 3955-3961.

Hovd, A.K., Nayar, S., Smith, C.G., Kanapathippillai, P., Iannizzotto, V., Barone, F., Fenton, K.A., and
Pedersen, H.L. (2024). Podoplanin expressing macrophages and their involvement in tertiary lymphoid
structures in mouse models of Sjogren's disease. Front Immunol *15*, 1455238.

Hu, X., Chung, A.Y., Wu, I., Foldi, J., Chen, J., Ji, J.D., Tateya, T., Kang, Y.J., Han, J., Gessler, M.*, et
al.* (2008). Integrated regulation of Toll-like receptor responses by Notch and interferon-gamma
pathways. Immunity *29*, 691-703.

Huster, K.M., Busch, V., Schiemann, M., Linkemann, K., Kerksiek, K.M., Wagner, H., and Busch, D.H.
(2004). Selective expression of IL-7 receptor on memory T cells identifies early CD40L-dependent
generation of distinct CD8+ memory T cell subsets. Proc Natl Acad Sci U S A *101*, 5610-5615.

Johnson, D.B., Balko, J.M., Compton, M.L., Chalkias, S., Gorham, J., Xu, Y., Hicks, M., Puzanov, I.,
Alexander, M.R., Bloomer, T.L.*, et al.* (2016). Fulminant Myocarditis with Combination Immune
Checkpoint Blockade. N Engl J Med *375*, 1749-1755.

Kaech, S.M., Tan, J.T., Wherry, E.J., Konieczny, B.T., Surh, C.D., and Ahmed, R. (2003). Selective
expression of the interleukin 7 receptor identifies effector CD8 T cells that give rise to long-lived
memory cells. Nat Immunol *4*, 1191-1198.

Kerrigan, A.M., Navarro-Nunez, L., Pyz, E., Finney, B.A., Willment, J.A., Watson, S.P., and Brown,
G.D. (2012). Podoplanin-expressing inflammatory macrophages activate murine platelets via CLEC-2.
J Thromb Haemost *10*, 484-486.

Knop, L., Deiser, K., Bank, U., Witte, A., Mohr, J., Philipsen, L., Fehling, H.J., Muller, A.J., Kalinke,
U., and Schuler, T. (2020). IL-7 derived from lymph node fibroblastic reticular cells is dispensable for
naive T cell homeostasis but crucial for central memory T cell survival. Eur J Immunol *50*, 846-857.

Kruse, B., Buzzai, A.C., Shridhar, N., Braun, A.D., Gellert, S., Knauth, K., Pozniak, J., Peters, J.,
Dittmann, P., Mengoni, M*., et al.* (2023). CD4(+) T cell-induced inflammatory cell death controls
immune-evasive tumours. Nature *618*, 1033-1040.

Leavitt, T., Hu, M.S., Borrelli, M.R., Januszyk, M., Garcia, J.T., Ransom, R.C., Mascharak, S.,
desJardins-Park, H.E., Litzenburger, U.M., Walmsley, G.G*., et al.* (2020). Prrx1 Fibroblasts Represent
a Pro-fibrotic Lineage in the Mouse Ventral Dermis. Cell Rep *33*, 108356.

Lee, K.W., Yeo, S.Y., Gong, J.R., Koo, O.J., Sohn, I., Lee, W.Y., Kim, H.C., Yun, S.H., Cho, Y.B.,
Choi, M.A*., et al.* (2022). PRRX1 is a master transcription factor of stromal fibroblasts for
myofibroblastic lineage progression. Nat Commun *13*, 2793.

Lee, S.H., Charmoy, M., Romano, A., Paun, A., Chaves, M.M., Cope, F.O., Ralph, D.A., and Sacks,
D.L. (2018). Mannose receptor high, M2 dermal macrophages mediate nonhealing Leishmania major
infection in a Th1 immune environment. J Exp Med *215*, 357-375.

Lee, S.H., Chaves, M.M., Kamenyeva, O., Gazzinelli-Guimaraes, P.H., Kang, B., Pessenda, G., Passelli,
K., Tacchini-Cottier, F., Kabat, J., Jacobsen, E.A*., et al.* (2020). M2-like, dermal macrophages are
maintained via IL-4/CCL24-mediated cooperative interaction with eosinophils in cutaneous
leishmaniasis. Sci Immunol *5*.

Lee, S.H., Kang, B., Kamenyeva, O., Ferreira, T.R., Cho, K., Khillan, J.S., Kabat, J., Kelsall, B.L., and
Sacks, D.L. (2023). Dermis resident macrophages orchestrate localized ILC2 eosinophil circuitries to
promote non-healing cutaneous leishmaniasis. Nat Commun *14*, 7852.

Leon, B., Lopez-Bravo, M., and Ardavin, C. (2007). Monocyte-derived dendritic cells formed at the
infection site control the induction of protective T helper 1 responses against Leishmania. Immunity *26*,
519-531.

Leung, G.A., Cool, T., Valencia, C.H., Worthington, A., Beaudin, A.E., and Forsberg, E.C. (2019). The
lymphoid-associated interleukin 7 receptor (IL7R) regulates tissue-resident macrophage development.
Development *146*.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics *30*, 923-930.

Liu, H., Li, P., Zhang, S., Xiang, J., Yang, R., Liu, J., Shafiquzzaman, M., Biswas, S., Wei, Z., Zhang,
Z*., et al.* (2022). Prrx1 marks stem cells for bone, white adipose tissue and dermis in adult mice. Nat
Genet *54*, 1946-1958.

Logan, M., Martin, J.F., Nagy, A., Lobe, C., Olson, E.N., and Tabin, C.J. (2002). Expression of Cre
Recombinase in the developing mouse limb bud driven by a Prxl enhancer. Genesis *33*, 77-80.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D.,
Hawrylycz, M.J., Jones, A.R*., et al.* (2010). A robust and high-throughput Cre reporting and
characterization system for the whole mouse brain. Nat Neurosci *13*, 133-140.

Mai, H.L., Boeffard, F., Longis, J., Danger, R., Martinet, B., Haspot, F., Vanhove, B., Brouard, S., and
Soulillou, J.P. (2014). IL-7 receptor blockade following T cell depletion promotes long-term allograft
survival. J Clin Invest *124*, 1723-1733.

Mandell, M.A., and Beverley, S.M. (2017). Continual renewal and replication of persistent Leishmania
major parasites in concomitantly immune hosts. Proc Natl Acad Sci U S A *114*, E801-E810.

Mantovani, A., Allavena, P., Marchesi, F., and Garlanda, C. (2022). Macrophages as tools and targets
in cancer therapy. Nat Rev Drug Discov *21*, 799-820.

Marine, J.C., Topham, D.J., McKay, C., Wang, D., Parganas, E., Stravopodis, D., Yoshimura, A., and
Ihle, J.N. (1999). SOCS1 deficiency causes a lymphocyte-dependent perinatal lethality. Cell *98*, 609-
616.

Mariotti, S., Teloni, R., Iona, E., Fattorini, L., Romagnoli, G., Gagliardi, M.C., Orefici, G., and Nisini,
R. (2004). Mycobacterium tuberculosis diverts alpha interferon-induced monocyte differentiation from
dendritic cells into immunoprivileged macrophage-like host cells. Infect Immun *72*, 4385-4392.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet. journal *17*, 10-12.

McCaughtry, T.M., Etzensperger, R., Alag, A., Tai, X., Kurtulus, S., Park, J.H., Grinberg, A., Love, P.,
Feigenbaum, L., Erman, B., and Singer, A. (2012). Conditional deletion of cytokine receptor chains
reveals that IL-7 and IL-15 specify CD8 cytotoxic lineage fate in the thymus. J Exp Med *209*, 2263-
2276.

Mishra, B.B., Rathinam, V.A., Martens, G.W., Martinot, A.J., Kornfeld, H., Fitzgerald, K.A., and
Sassetti, C.M. (2013). Nitric oxide controls the immunopathology of tuberculosis by inhibiting NLRP3
inflammasome-dependent processing of IL-1beta. Nat Immunol *14*, 52-60.

Mombaerts, P., Iacomini, J., Johnson, R.S., Herrup, K., Tonegawa, S., and Papaioannou, V.E. (1992).
RAG-1-deficient mice have no mature B and T lymphocytes. Cell *68*, 869-877.

Mulder, K., Patel, A.A., Kong, W.T., Piot, C., Halitzki, E., Dunsmore, G., Khalilnezhad, S., Irac, S.E.,
Dubuisson, A., Chevrier, M., *et al.* (2021). Cross-tissue single-cell landscape of human monocytes and
macrophages in health and disease. Immunity *54*, 1883-1900 e1885.

Muller, A.J., Aeschlimann, S., Olekhnovitch, R., Dacher, M., Spath, G.F., and Bousso, P. (2013).
Photoconvertible pathogen labeling reveals nitric oxide control of Leishmania major infection in vivo
via dampening of parasite metabolism. Cell Host Microbe *14*, 460-467.

Muller, A.J., Filipe-Santos, O., Eberl, G., Aebischer, T., Spath, G.F., and Bousso, P. (2012). CD4+ T
cells rely on a cytokine gradient to control intracellular pathogens beyond sites of antigen presentation.
Immunity *37*, 147-157.

Murakami, N., Borges, T.J., Yamashita, M., and Riella, L.V. (2016). Severe acute interstitial nephritis
after combination immune-checkpoint inhibitor therapy for metastatic melanoma. Clin Kidney J *9*, 411-
417.

Nalio Ramos, R., Missolo-Koussou, Y., Gerber-Ferder, Y., Bromley, C.P., Bugatti, M., Nunez, N.G.,
Tosello Boari, J., Richer, W., Menger, L., Denizeau, J., *et al.* (2022). Tissue-resident FOLR2(+)
macrophages associate with CD8(+) T cell infiltration in human breast cancer. Cell *185*, 1189-1207
e1125.

Navarro, S., Cossalter, G., Chiavaroli, C., Kanda, A., Fleury, S., Lazzari, A., Cazareth, J., Sparwasser,
T., Dombrowicz, D., Glaichenhaus, N., and Julia, V. (2011). The oral administration of bacterial extracts
prevents asthma via the recruitment of regulatory T cells to the airways. Mucosal Immunol *4*, 53-65.

Olekhnovitch, R., Ryffel, B., Muller, A.J., and Bousso, P. (2014). Collective nitric oxide production
provides tissue-wide immunity during Leishmania infection. J Clin Invest *124*, 1711-1722.

Onder, L., Scandella, E., Chai, Q., Firner, S., Mayer, C.T., Sparwasser, T., Thiel, V., Rulicke, T., and
Ludewig, B. (2011). A novel bacterial artificial chromosome-transgenic podoplanin-cre mouse targets
lymphoid organ stromal cells in vivo. Front Immunol *2*, 50.

Ortega-Gomez, A., Perretti, M., and Soehnlein, O. (2013). Resolution of inflammation: an integrated
view. EMBO Mol Med *5*, 661-674.

Oyesola, O.O., Hilligan, K.L., Namasivayam, S., Howard, N., Clancy, C.S., Zhao, M., Oland, S.D.,
Kiwanuka, K.N., Garza, N.L., Lafont, B.A.P., *et al.* (2023). Exposure to lung-migrating helminth
protects against murine SARS-CoV-2 infection through macrophage-dependent T cell activation. Sci
Immunol *8*, eadf8161.

Patir, A., Gossner, A., Ramachandran, P., Alves, J., Freeman, T.C., Henderson, N.C., Watson, M., and
Hassan, M.A. (2020). Single-cell RNA-seq reveals CD16(-) monocytes as key regulators of human
monocyte transcriptional response to Toxoplasma. Sci Rep *10*, 21047.

Peduto, L., Dulauroy, S., Lochner, M., Spath, G.F., Morales, M.A., Cumano, A., and Eberl, G. (2009).
Inflammation recapitulates the ontogeny of lymphoid stromal cells. J Immunol *182*, 5789-5799.

Peschon, J.J., Morrissey, P.J., Grabstein, K.H., Ramsdell, F.J., Maraskovsky, E., Gliniak, B.C., Park,
L.S., Ziegler, S.F., Williams, D.E., Ware, C.B.*, et al.* (1994). Early lymphocyte expansion is severely
impaired in interleukin 7 receptor-deficient mice. J Exp Med *180*, 1955-1960.

Picelli, S., Bjorklund, A.K., Faridani, O.R., Sagasser, S., Winberg, G., and Sandberg, R. (2013). Smart-
seq2 for sensitive full-length transcriptome profiling in single cells. Nat Methods *10*, 1096-1098.

Picelli, S., Faridani, O.R., Bjorklund, A.K., Winberg, G., Sagasser, S., and Sandberg, R. (2014). Full-
length RNA-seq from single cells using Smart-seq2. Nat Protoc *9*, 171-181.

Pisu, D., Huang, L., Narang, V., Theriault, M., Le-Bury, G., Lee, B., Lakudzala, A.E., Mzinza, D.T.,
Mhango, D.V., Mitini-Nkhoma, S.C.*, et al.* (2021). Single cell analysis of M. tuberculosis phenotype
and macrophage lineages in the infected lung. J Exp Med *218*.

Postat, J., Olekhnovitch, R., Lemaitre, F., and Bousso, P. (2018). A Metabolism-Based Quorum Sensing
Mechanism Contributes to Termination of Inflammatory Responses. Immunity *49*, 654-665 e655.

Rayes, J., Lax, S., Wichaiyo, S., Watson, S.K., Di, Y., Lombard, S., Grygielska, B., Smith, S.W.,
Skordilis, K., and Watson, S.P. (2017). The podoplanin-CLEC-2 axis inhibits inflammation in sepsis.
Nat Commun *8*, 2239.

Romano, A., Carneiro, M.B.H., Doria, N.A., Roma, E.H., Ribeiro-Gomes, F.L., Inbar, E., Lee, S.H.,
Mendez, J., Paun, A., Sacks, D.L., and Peters, N.C. (2017). Divergent roles for Ly6C+CCR2+CX3CR1+
inflammatory monocytes during primary or secondary infection of the skin with the intra-phagosomal
pathogen Leishmania major. PLoS Pathog *13*, e1006479.

Saliba, A.E., Li, L., Westermann, A.J., Appenzeller, S., Stapels, D.A., Schulte, L.N., Helaine, S., and
Vogel, J. (2016). Single-cell RNA-seq ties macrophage polarization to growth rate of intracellular
Salmonella. Nat Microbiol *2*, 16206.

Sanin, D.E., Ge, Y., Marinkovic, E., Kabat, A.M., Castoldi, A., Caputa, G., Grzes, K.M., Curtis, J.D.,
Thompson, E.A., Willenborg, S.*, et al.* (2022). A common framework of monocyte-derived macrophage
activation. Sci Immunol *7*, eabl7482.

Seddon, B., Tomlinson, P., and Zamoyska, R. (2003). Interleukin 7 and T cell receptor signals regulate
homeostasis of CD4 memory cells. Nat Immunol *4*, 680-686.

Sorensen, M., Lippuner, C., Kaiser, T., Misslitz, A., Aebischer, T., and Bumann, D. (2003). Rapidly
maturing red fluorescent protein variants with strongly enhanced brightness in bacteria. FEBS Lett *552*,
110-114.

Stapels, D.A.C., Hill, P.W.S., Westermann, A.J., Fisher, R.A., Thurston, T.L., Saliba, A.E.,
Blommestein, I., Vogel, J., and Helaine, S. (2018). Salmonella persisters undermine host immune
defenses during antibiotic treatment. Science *362*, 1156-1160.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A.,
Pomeroy, S.L., Golub, T.R., Lander, E.S., and Mesirov, J.P. (2005). Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S
A *102*, 15545-15550.

Sugimoto, M.A., Sousa, L.P., Pinho, V., Perretti, M., and Teixeira, M.M. (2016). Resolution of
Inflammation: What Controls Its Onset? Front Immunol *7*, 160.

Szabo, P.A., Dogra, P., Gray, J.I., Wells, S.B., Connors, T.J., Weisberg, S.P., Krupska, I., Matsumoto,
R., Poon, M.M.L., Idzikowski, E.*, et al.* (2021). Longitudinal profiling of respiratory and systemic
immune responses reveals myeloid cell-driven lung inflammation in severe COVID-19. Immunity *54*,
797-814 e796.

Traag, V.A., Waltman, L., and van Eck, N.J. (2019). From Louvain to Leiden: guaranteeing well-
connected communities. Sci Rep *9*, 5233.

Veglia, F., Sanseviero, E., and Gabrilovich, D.I. (2021). Myeloid-derived suppressor cells in the era of
increasing myeloid cell diversity. Nat Rev Immunol *21*, 485-498.

Wallis, R.S., O'Garra, A., Sher, A., and Wack, A. (2023). Host-directed immunotherapy of viral and
bacterial infections: past, present and future. Nat Rev Immunol *23*, 121-133.

Wolowczuk, I., Delacre, M., Roye, O., Giannini, S.L., and Auriault, C. (1997). Interleukin-7 in the skin
of Schistosoma mansoni-infected mice is associated with a decrease in interferon-gamma production
and leads to an aggravation of the disease. Immunology *91*, 35-44.

Wykes, M.N., and Lewin, S.R. (2018). Immune checkpoint blockade in infectious diseases. Nat Rev
Immunol *18*, 91-104.

Xie, Z., Shao, B., Hoover, C., McDaniel, M., Song, J., Jiang, M., Ma, Z., Yang, F., Han, J., Bai, X.*, et*
al. (2020). Monocyte upregulation of podoplanin during early sepsis induces complement inhibitor
release to protect liver function. JCI Insight *5*.

Yamada, K.J., Heim, C.E., Xi, X., Attri, K.S., Wang, D., Zhang, W., Singh, P.K., Bronich, T.K., and
Kielian, T. (2020). Monocyte metabolic reprogramming promotes pro-inflammatory activity and
Staphylococcus aureus biofilm clearance. PLoS Pathog *16*, e1008354.

Yan, T., Yu, L., Zhang, J., Chen, Y., Fu, Y., Tang, J., and Liao, D. (2024). Achilles' Heel of currently
approved immune checkpoint inhibitors: immune related adverse events. Front Immunol *15*, 1292122.

Yazdani, R., Askari, M., Ahmadi, A.M., Azizi, G., Ciric, B., Boehm, A., Zhang, G.X., and Rostami, A.
(2024). Mouse monocytes express CD127 by immune cells, not LPS. Front Immunol *15*, 1356004.

Zhang, B., Zhang, Y., Xiong, L., Li, Y., Zhang, Y., Zhao, J., Jiang, H., Li, C., Liu, Y., Liu, X.*, et al.*
(2022). CD127 imprints functional heterogeneity to diversify monocyte responses in inflammatory
diseases. J Exp Med *219*.

1008

Acknowledgments: The helpful discussions with Tim Lämmermann, Tatjana Ruhl, Gabriel Sollberger,

and all members of the Institute of Molecular and Clinical Immunology are gratefully acknowledged.

We thank Tamas Dolowschiak for invaluable advice for setting up the scRNAseq library construction.

We also thank Roland Hartig from the MPBIC platform and the team of the ZTL of the Medical Faculty

OvGU for excellent technical support, and Ehsan Vafadarnejad for the support with initial scRNAseq

data analysis.

1015

1016   **Funding:** This work was supported by funding from the European Research Council (ERC) under the

1017   European Union's Horizon 2020 research and innovation program (StG ImmProDynamics, grant

1018   agreement 714233 to A.J.M and grant 677200 to N.J.), the German Research Foundation (DFG)

1019   (SFB854-B31, MU3744/6-1, and SPP2225 EXIT (MU3744/5-1)).

1020

1021   **Author contributions:** YF, TS, AES and AJM conceived of and designed the study. YF, IS, ER, LP,

1022   PG, LK, BK, LAD, VD, JA, NJN, MB, IB, JM, JP, CS, BP, and AJM conducted experiments, YF, ER,

1023   CT, LP, JM, SK, LK, BK, SF, AB, RG and AJM analyzed the data. NG, TT, BS, HVH, MG, BB, NJ,

1024   and TS provided critical technology. YF, TS, AES and AJM wrote the manuscript. All authors revised,

1025   commented and approved the manuscript.

1026

1027   **Competing interests:** The authors declare no competing interests.

1028

1029   **Data and materials availability:** RNAseq data supporting the findings of this study have been

1030   deposited at Gene Expression Omnibus under accession no. GSE228455. R scripts used for the analysis

1031   of the scRNAseq data are available on GitHub: https://github.com/saliba-lab/Monocytes_leishmania

1032

1033

 Figure legends

 Figure 1.

 PDPN$^+$IL-7R$^+$ phagocytes infected with low proliferating *L. major* exhibit a hypo-inflammatory

 phenotype.

 A. Experimental setup for *in vivo L. major* proliferation measurement using *Lm*^{SWITCH}. **B.** Experimental

 setup for index single-cell sorting of CD11c^{int-hi} and Ly6c$^+$ monocytes from 10 individual mouse ears

 infected with proliferation reporter parasites at 3 weeks post-infection (p.i.). **C.** Index mKikume red and

 green reporter fluorescence values. **D.** *L. major* reads detected in the sorted monocytes infected with

 high (green) or low (red) proliferating pathogen, or uninfected (black). **E.** UMAP representation and

 mapping of cells sorted as high-proliferator infected (green), uninfected (black) and low-proliferator

 infected (red). **F.** Leiden clustering of the scRNAseq data on the UMAP representation. **G.** Fraction of

 monocytes sorted as infected with *Lm*^{hi} (green), uninfected (black) or infected with *Lm*^{lo} (red) among

 the three clusters identified by the Leiden algorithm. The five highlighted most prominent populations

 (*Lm*^{hi}Mo1, UMo1, Cluster 1; *Lm*^{hi}Mo2, Umo2, Cluster 2; *Lm*^{lo}Mo3, Cluster 3) were selected for

 differentially expressed gene (DEG) analysis. **H.** Normalized expression (color-code) and cell fraction

 per population with detected expression (circle size) for selected DEG differentially expressed in

 Lm^{lo}Mo3 versus all other populations. **I.** Quantitative PCR (qPCR) analysis of selected genes identified

 by scRNAseq. Expression levels were determined for replicates of 50-100 cells sorted according to the

 gating strategy outlined in Fig. S1K. Expression levels relative to *Gapdh* and normalized to *Lm*^{lo}Mo3

 are shown. Each symbol represents 50-100 cell replicates; data was generated from two independent

 experiments. Horizontal lines denote the median. ****, $p<0.0001$;***, $p<0.001$;**, $p<0.01$;*, $p<0.05$

 compared to the *Lm*^{lo}Mo3 population according to one-way ANOVA with Dunnett's post-test. **J.** Mean

 fluorescence intensity (MFI) of pro-IL-1β, CXCL9, IL-6, and IL-12 p35 among the *Lm*^{hi}Mo1, Umo1,

 Lm^{hi}Mo2, Umo2, and *Lm*^{lo}Mo3 (according to the gating strategy shown in Fig. S1K) from *L. major*

 infected C56BL/6J mouse ear skin at 3 weeks p.i. Each dot represents one infected ear analyzed, data

 pooled from two independent experiments. Horizontal lines denote mean. ****, $p<0.0001$; **, $p<0.01$;

 *, $p<0.05$ according to one-way ANOVA multiple comparison with Dunnett's post-test.

1062

1063

Figure 2. IL-7R⁺PDPN⁺ *Lm*ˡᵒMo3 develop from bone-marrow derived CCR2⁺ monocytes and acquire their phenotype in the infected tissue.

A. Gating to determine tissue-resident and monocyte-derived dendritic cells (TR/MoDC), tissue-resident macrophages (TRM) and different monocyte-derived populations (left) and proportion of these populations among CD11b⁺Ly6G⁻SiglecF⁻ phagocytes (right). **B.** Proportion of the cell populations shown in (A) among *Lm*ʰⁱMo1, Umo1, *Lm*ʰⁱMo2, Umo2, and *Lm*ˡᵒMo3 (H) and among infected CD11b⁺Ly6G⁻SiglecF⁻ cells with differential PDPN and IL-7R expression and different intracellular pathogen proliferation rates. Horizontal lines denote the median. Each dot represents one infected ear analyzed, data pooled from two independent experiments. **C.** Experimental approach to identify newly recruited monocytes by photoconversion of mKikume transgenic mice (left). **D.** Upper left: Infected CD11b⁺Ly6G⁻SiglecF⁻ cells from *Lm*ᵀᵘʳᑫᵘᵒⁱˢᵉ-infected ear tissue at 3 weeks p.i. from mKikume transgenic mice before (green curve) and 0 h after photoconversion (red curve). Lower left: Identification of short resident mKikume green (recruited <5 days after photoconversion) and long resident mKikume red (recruited >5 days after photoconversion) cells among infected CD11b⁺Ly6G⁻SiglecF⁻ cells. Right: Examples of the proportion long resident cells (mKikume red⁺) among infected CD11b⁺Ly6G⁻SiglecF⁻ cells with differential PDPN and IL-7R expression. **E.** Quantification of short and long resident cells among infected CD11b⁺Ly6G⁻SiglecF⁻ cells with differential PDPN and IL-7R expression. Horizontal lines denote the median. Each dot represents one infected analyzed ear, data pooled from two independent experiments. **F.** Experimental approach to demonstrate the origin of *Lm*ˡᵒMo3 from bone marrow-derived, CCR2⁺ monocytes recruited during the innate and adaptive phase of the anti-*Leishmania* response. **G.** CCR2-creᴱᴿ-dependent TdTomato expression in different blood leukocytes cells before, and at day 2, 6 and 23 after the beginning of five daily Tamoxifen injections. Horizontal lines denote the median. Each dot represents one animal, data pooled from two independent experiments. **H.** Fraction of CCR2-creᴱᴿ-dependent tdTomato expressing cells among PDPN and IL-7R expressing monocyte-derived cells. The cre recombinase activity was induced in CCR2⁺ monocytes for the innate phase of the immune response by injecting Tamoxifen before infection (left), or for the adaptive phase

1090   by injection at 11-15 days p.i. (right). Horizontal lines denote the median. Each symbol represents one

1091   infection site per condition obtained from two independent experiments.

1092

Figure 3. PDPN+IL-7R+ *Lm*loMo3 dampen T cell responses against *L. major* infection *in vivo*.

A. Proportion of PDPN-IL-7R+ (left), PDPN+IL-7R- (middle) and PDPN+IL-7R+ (right) fractions among

different leukocytes at the infection site at week 3 p.i. Bars denote the median. Each symbol represents

one infection site per condition obtained from two independent experiments. **B.** Experimental setup for

cell-specific *Il7r*-deficiency in PDPN+ leukocytes. **C.** Validation of cell-specific IL-7R deficiency.

Upper panels: Example and MFI analysis for IL-7R surface expression on *Lm*loMo3 (gated according to

figure S1K) in the infected ear tissue at 3 weeks p.i. Lower panels: Example and MFI analysis of IL-7R

expression on CD4+ T cells in the draining lymph node (dLN) at week 3 p.i. Values are normalized to

control animals. Each dot represents one infected ear or dLN analyzed. Isotype control-stained samples

are shown as a grey histogram and dashed line in the data plot, respectively. Horizontal lines denote

mean, data pooled from 2 independent experiments. **, $p < 0.01$; ns, not significant according to unpaired

t-test. **D.** *L. major*-infected cell fraction among recruited monocytes in the ear tissue (left) and limiting

dilution assay (LDA) of *L. major* tissue burden normalized to control animals (right) at week 3 p.i. of

lethally irriatiated mice either reconstituted with littermate control (black symbols) or *Pdpn-cre × Il7r*fl/fl

BM (blue symbols). **E.** IFNγ+ fraction and IFNγ MFI among effector CD4+ T cells within *L. major*-

infected ear at week 3 p.i.for mice reconstituted with either littermate control (black symbols) or *Pdpn-*

*cre × Il7r*fl/fl BM (blue symbols). Each symbol in (D-E) represents one individual mouse ear. Horizontal

lines denote the mean. ***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$ according to unpaired t-test. Data

representative of two independent experiments. **F.** Experimental setup for mixed BM chimera

reconstituted with 50% CD45.1 WT and 50% *Il7r*-/- (CD45.2) BM. Mice reconstituted with mixtures of

WT CD45.1 and WT CD45.2 BM served as controls. **G.** *L. major* infected cell fraction among recruited

monocytes in the ear tissue at week 3 p.i., either WT BM chimera compartment (black symbols), or *Il7r*-

/- BM chimera compartment (blue symbols). **H.** Fraction of *Lm*hiMo1, *Lm*hiMo2, and *Lm*loMo3 (gated

according to figure S1K) among monocyte-derived cells of each BM chimera compartment, either WT

BM chimera compartment (closed symbols), or *Il7r*-/- BM chimera compartment (open symbols) within

L. major infected ear tissue at 3 week p.i. Each symbol represents one individual mouse ear, data pooled

from two independent experiments. Horizontal lines denote the median. No significance detected

according to one-way ANOVA multiple comparison with Dunn's post-test. **I**. Intracellular pathogen

proliferation rates among Lm^{hi}Mo1, Lm^{hi}Mo2, and Lm^{lo}Mo3 (gated according to figure S1K), either WT

BM chimera compartment (closed symbols), or $Il7r^{-/-}$ BM chimera compartment (open symbols). Each

symbol represents one individual mouse ear, data pooled from two independent experiments. Horizontal

lines denote the median. ****, $p < 0.0001$ according to one-way ANOVA multiple comparison with

Dunnett post-test. **J.** Experimental setup for depletion of *Pdpn*-expressing leukocytes of lethally

irradiated mice reconstituted with either *Pdpn-cre* × iDTR BM cells, or 50% of *Pdpn-cre* × iDTR and

50% of CD45.1 WT BM cells. **K.** Validation of depletion of PDPN$^+$ cells among infected leukocytes.

Upper panel: Example of PDPN staining of infected cells from ear tissue at 3 weeks p.i. in depleted

(blue curve) or non-depleted control animals (black curve). Lower panel: Percentage of PDPN$^+$ cells

among infected leukocytes from ear tissue at 3 weeks p.i. in depleted (blue symbols) or non-depleted

control animals (black symbols). Horizontal lines denote the mean. Each dot represents one infected ear

analyzed. *, $p < 0.05$; ns, not significant according to unpaired t-test. **L.** *L. major* infected phagocyte

fraction (left) and *L. major* burden (right, normalized to non-depleted control) in the ear at week 3 p.i.

in non-depleted (black symbols) or partially depleted (grey symbols) controls versus 100% *Pdpn*-

dependent depleted BM chimeras (blue symbols). **M.** Fraction (left) and MFI (right) of IFNγ$^+$ expressing

CD4$^+$CD44$^+$ effector T cells from ear tissue (left) in the ear at week 3 p.i. in non-depleted (black

symbols) or partially depleted (grey symbols) controls versus 100% *Pdpn*-dependent depleted BM

chimeras (blue symbols). Each symbol in (L) and (M) represents one individual mouse ear, data

representative of two independent experiments. Horizontal lines denote the median. **, $p < 0.01$

according to Mann-Whitney test.

1141

Figure 4. Fibroblast-produced IL-7 contributes to pathogen persistence and suppression of IFNγ

production in effector T cells.

A. Upper panels: H2KB$^+$ or PD-L1$^+$ fraction and H2KB or PD-L1 MFI of among fibroblasts from mice

reconstituted with *Pdpn-cre* × $Il7r^{WT/WT}$ littermate control BM (black symbols) or *Pdpn-cre* × $Il7r^{fl/fl}$ BM

(blue symbols). Data normalized to mice reconstituted with littermate control BM. Lower panels:

H2KB$^+$ or PD-L1$^+$ fraction and H2KB or PD-L1 MFI of among fibroblasts from non-depleted control

(black symbols) or *Pdpn*-dependent 100% depleted (blue symbols) ear infections. Each symbol

represents one individual mouse ear. Data normalized to non-depleted control animals. Horizontal lines

denote the mean. ***, p<0.001; **, p<0.01; *, p<0.05 according to unpaired t-test. Data shown

correspond to the experiments shown in fig. 3C-E, , and K, respectively. **B.** Expression of *Il7* within *L.*

major-infected (week 3 p.i.) and non-infected ear tissue. Expression as determined by ΔΔCT between

Il7 and *Hprt* qPCR. CT values are corrected according to the mean difference in RNA extracted (see fig.

S4C). Each symbol represents one individual mouse ear. Horizontal lines denote the mean. *, p<0.05

according to unpaired t-test. **C.** Expression level of *Prrx* among fibroblasts, LEC, BEC and PDPN$^-$CD31$^-$

non-immune cells within *L. major*-infected ear tissue at week 3 p.i.. Expression as determined by ΔΔCT

between *Prrx* and *Hprt* qPCR followed by normalization to PDPN$^-$CD31$^-$ cells. Each symbol represents

one individual mouse ear. Horizontal lines denote the mean. ****, p<0.0001 according to one-way

ANOVA multiple comparison with Dunnett's post-test. Data in (A-C) pooled from two independent

experiments. **D.** Proportion of fibroblasts, LEC, and BEC among CD45$^-$TER119$^-$7AAD$^-$ cells within the

L. major infected ear at week 3 p.i.. Each symbol represents one individual mouse ear. Horizontal lines

denote the mean. ****, p<0.0001 according to one-way ANOVA multiple comparison with Dunnett's

post-test. **E.** Experimental setup to test for *Prrx*-expressing fibroblasts as a necessary source of IL-7 to

promote *Lm*^{lo}Mo3 mediated suppression of CD4$^+$ T cell responses in the *L. major* infected ear tissue. **F.**

Expression level of *Il7* among fibroblasts, LEC, and BEC within *L. major*-infected ear tissue from

littermate control (*Prrx-Cre* × *Il7*$^{wt/wt}$) (black symbols) and *Prrx-cre* × *Il7*$^{fl/fl}$ (blue symbols). Expression

as determined by ΔCT between *Il7* and *Hprt* qPCR. **G.** Left: the fraction of *L. major* infected cells

among recruited monocytes in the ear tissue at week 3 p.i., littermate control (*Prrx-Cre* × *Il7*$^{wt/wt}$) (filled

symbols) and *Prrx-cre* × *Il7*$^{fl/fl}$ (open symbols). Differently shaded symbols represent two independent

experiments. Each symbol represents one individual mouse ear. Horizontal lines denote the median.

***, p<0.001 according to two-way ANOVA comparing genotypes. Right: Limiting dilution assay

(LDA) of *L. major* tissue burden in the ear. Data normalized to littermate control animals. **H.** IFNγ$^+$

fraction and MFI of IFNγ signal among effector CD4$^+$ T cells within *L. major*-infected ear at week 3 p.i.

of either littermate control (black symbols) or *Prrx-cre × Il7^{fl/fl}* animals (blue symbols). Data normalized

to littermate control animals. Each symbol in (G-I) represents one individual mouse ear, data are from

two independent experiments. Horizontal lines denote the mean. *, p<0.05 according to unpaired t-test.

1177

Figure 5. IFNγ activates fibroblasts to produce IL-7 depending of the infection amplitude.

A. Proportion of IFNγ$^+$ produced by T, NK, and dendritic cells at week 3 p.i. Each symbol represents

one individual mouse ear. Horizontal lines denote the mean. ****, p<0.0001 according to unpaired t-

test. **B.** Expression of the *Il7* cytokine gene in ex vivo cultivated fibroblasts without (black symbols) and

with (red symbols) addition of IFNγ$^+$ for 24h. Horizontal lines denote the median. **, p<0.01 according

to Mann-Whitney test. **C.** Experimental setup to test the impact of anti-IFNγ treatment in fibroblasts

within an ongoing *L. major* infection. **D.** Correlation (r^2, with p value indicated) and linear regression

(red line) between *L. major* infection rate (determined by flow cytometry) and *Il7* expression level

(qPCR) in fibroblasts within the infected ear (week 3 p.i.), in either isotype control (black symbols,

upper graph) or anti-IFNγ treated (orange symbols, lower graph) animals. Each symbol represents one

replicate of flow cytometry analysis from six individual infected ears per condition with qPCR of up to

3 pools of 200 fibroblasts sorted from the same infected ear. The vertical dashed line shows the threshold

defined for analyzing highly infected tissues. **E.** Comparison of *Il7* expression in fibroblasts sorted from

tissues with an infection rate above the threshold (dashed line) shown in (D). **F.** H2KB$^+$ or PD-L1$^+$

fraction and H2KB or PD-L1 MFI among fibroblasts isolated from week 3 p.i. infected ears of animals

treated with either isotype control (black symbols) or anti-IFNγ antibody (orange symbols). Each symbol

in (E-G) represents one individual mouse ear, horizontal lines denote the mean. ***, p<0.001; **,

p<0.01; *, p<0.05 according to unpaired t-test. Data were collected in two independent experiments.

1196

1197   **Figure 6. Anti-IL-7/IL-7R treatment results in an enhanced immune response against *L. major*.**

1198   **A.** Hypothesis (box) and experimental setup for testing the effects of anti-IL-7/IL-7R treatment during

1199   *L. major* infection. Treatment was applied twice (day 16, 20, black arrows, data shown in panel B-F) or

1200   once (day 18, grey arrow, data shown in panel C). **B-C.** Limiting dilution assay (LDA) of *L. major* tissue

1201   burden in the ear following anti-IL-7/IL-7R treatment normalized to isotype control treated animals (B)

and *L. major* infected cell fraction among recruited monocytes with two-dosage (left panel) or single-dosage anti-IL-7/IL-7R treatment (right panel) (C). **D.** IFNγ$^+$ fraction (left) and IFNγ MFI (normalized to isotype control treatment, right) among effector CD4$^+$ T cells isolated from the ears at week 3 p.i. of either isotype antibody (black symbols) or anti-IL-7R/IL-7 treated animals (blue symbols). **E.** H2KB$^+$ and PD-L1$^+$ fraction among fibroblasts, and the MFI of H2KB or PD-L1 among fibroblasts (normalized to isotype control animals), either isotype control (black symbols) or anti-IL-7R/IL-7 treated (blue symbols). **F.** iNOS$^+$ fraction and iNOS MFI (normalized to isotype control animals) among total recruited monocyte-derived cells (left panel) and infected monocyte-derived cells (right panel), either isotype control (black symbols) or anti-IL-7R/IL-7 treated samples (blue symbols). Each symbol in (B-F) represents one individual mouse ear, data pooled from two independent experiments. Horizontal lines denote the mean. . **, $p < 0.01$; *, $p < 0.05$ according to unpaired t-test. **G.** Experimental setup to determine the long-term effects of anti-IL-7/IL-7R treatment. **H.** Infection-associated immune cell recruitment determined by ear diameter of isotype control-treated (grey) and anti-IL-7/IL-7R treated (blue) animals. Data were normalized to the mean of the corresponding values of the same ear measured before the treatment. Bars denote the mean+SD. Each symbol represents one individual ear repeatedly measured throughout the infection time course. Data obtained from two independent experiments. ****, $p < 0.0001$ *, $p < 0.05$ according to 2-way ANOVA (treatment, experiment) with Bonferroni post-test. **I.** Limiting dilution assay (LDA) of *L. major* tissue burden in the ear at week 7 p.i. of animals previously treated (week 3 p.i.) with isotype control-treated (black symbols) or anti-IL-7/IL-7R treated (blue symbols). **J.** IFNγ$^+$ fraction among CD4$^+$ T cells isolated from the draining lymph node at week 7 p.i. of animals previously treated (week 3 p.i.) with either isotype control (black symbols) or anti-IL-7R/IL-7 antibody (blue symbols). Each symbol in (I-J) represents one individual ear. Data pooled from two independent experiments and normalized to control animals of each experiment for limiting dilution. Horizontal lines denote the median. *, $p < 0.05$ according unpaired Mann-Whitney test. Each symbol represents one individual mouse ear. Horizontal lines denote the mean. ***, $p < 0.001$; according to unpaired Mann-Whitney test.

Figure 1

Figure 2

Figure 4

Figure 5

A IFNγ-producing leukocytes (ear)

% of IFNγ⁺

T cells NK XCR1^hi XCR1^int XCR1^lo

****

B *In vitro* fibroblast *Il7* expression

Fold *hprt*

**

● Control
○ 24h IFNγ

C WT mice Lm^{DsRed} Anti-IFNγ

Day 0 15,18 21 Fibro-blast IL-7?

CD4⁺ T_eff: IFNγ
Fibroblasts: IL-7
Lm^{lo}Mo3: IL-7R

D Isotype

Il7 expression (fold *hprt*)

r² = 0.46
p < 0.01

Anti-IFNγ

r² = 0.02
p = 0.59

% infected CD45⁺

E Fibroblasts in highly infected tissues

Il7 expression (fold *hprt*)

*

● Isotype control
○ Anti-IFNγ

F Fibroblast activation

H2-KB

% Fibroblasts Norm. MFI

* *

PD-L1

% Fibroblasts Norm. MFI

** *

● Isotype control
○ Anti-IFNγ

Figure 6