MLL3/4 methyltransferases regulate the differentiation of pluripotent stem cells through coordinating glycolysis and mitochondrial respiration

Suza Mohammad Nur[1,3], Yunbo Jia[1], Muyi Ye[1], Caylin A. Lepak[1], Issam Ben-Sahra[4,5], and Kaixiang Cao[1,2,3,*]

[1]Department of Biochemistry, Case Western Reserve University

[2]Department of Genetics and Genome Sciences, Case Western Reserve University

[3]Case Comprehensive Cancer Center, Case Western Reserve University, Cleveland, OH 44106, USA

[4]Department of Biochemistry and Molecular Genetics, Northwestern University

[5]Robert H. Lurie Comprehensive Cancer Center, Northwestern University, Chicago, IL 60611, USA

Running title: MLL3/4 regulate cellular respiration to facilitate pluripotent stem cell differentiation

Keywords: MLL3, MLL4, stem cells, differentiation, cellular metabolism, glycolysis, TCA cycle

*Correspondence should be sent to the following address:

Kaixiang Cao, PhD

Department of Biochemistry, School of Medicine, Case Western Reserve University

10900 Euclid Avenue, Cleveland, OH 44106, USA

Office: (216) 368-1218

Email: kxc725@case.edu

Abstract

Enhancer-regulating epigenetic modifiers play critical roles in normal physiological processes and human pathogenesis. The major enhancer regulator paralogs MLL3 and MLL4 (MLL3/4) belong to the lysine methyltransferase 2 (KMT2) family, which catalyzes the methylation of lysine 4 on histone H3 (H3K4me). MLL3/4 are required for enhancer activation and are essential for mammalian development and stem cell differentiation. Although recent studies have linked MLL3/4 with different metabolic pathways in the regulation of stem cell self-renewal and cancer cell growth, the mechanisms connecting enhancer function to metabolic control remain elusive. Here, using respiration flux assays, stable isotope tracing, transcriptomics, and stem cell biology techniques, we show that the loss of MLL3/4 impairs glycolysis and mitochondrial respiration in murine embryonic stem cells. Mechanistically, MLL3/4 deficiency suppresses the expression of the rate-limiting glycolytic enzyme hexokinase 2 (HK2) and compromises the function of the oxoglutarate dehydrogenase (OGDH) complex, thereby coordinately impairing central carbon metabolism. Remarkably, combined restoration of HK2 and OGDH rescues the metabolic defects caused by MLL3/4 loss and reinstates differentiation capacity. Taken together, our study identifies a direct link between enhancer-regulating epigenetic machineries and metabolic control of cell fate transition, providing a mechanistic framework for understanding how enhancer malfunction contributes to developmental abnormalities and human diseases.

Introduction

Enhancer deregulation is a major driver of human congenital disorders (*1-4*). In mammals, nucleosomes at enhancers are mono-methylated on lysine 4 of histone H3 (H3K4) by the paralogous enzymes Mixed-lineage Leukemia 3 (MLL3, also known as KMT2C) and Mixed-lineage Leukemia 4 (MLL4, also known as KMT2D) (*5-9*). Mice null for MLL4 die shortly after gastrulation, while MLL3 loss leads to neonatal lethality (*10, 11*). Heterozygous loss-of-function mutations in MLL4 lead to Kabuki syndrome (*1*) and those in MLL3 cause Kleefstra syndrome 2 (*4, 12, 13*). Both disorders are characterized by craniofacial dysmorphism and developmental delay, highlighting the instructive role of MLL3/4 in normal development and human pathogenesis. However, molecular mechanisms by which MLL3/4 function in biologically normal and pathological processes remain poorly understood.

Recent results have revealed a functional link between MLL3/4 and cellular metabolism. MLL3/4 and nucleotide synthesis pathways co-regulate the self-renewal of pluripotent stem cells (PSCs) (*14*). Moreover, steady-state abundance of glycolytic metabolites such as glucose-6-phosphate (G6P) and fructose-6-phosphate (F6P), and tricarboxylic acid (TCA) cycle metabolites including succinate, fumarate, and malate, is reduced in PSCs lacking MLL3/4 (*14*). In contrast, MLL4-deficient lung cancer and melanoma cells adopt upregulated glycolysis (*15, 16*), suggesting a context-dependent role of MLL3/4 in regulating cellular metabolism. How MLL3/4 modulate metabolic pathways and whether they govern cell fate via these pathways are largely unknown.

Glycolysis and mitochondrial respiration are tightly regulated in PSCs. Naive PSCs, such as murine embryonic stem cells (ESCs), exhibit metabolic bivalency, utilizing both glycolysis and mitochondrial respiration for energy production, whereas primed PSCs are predominantly glycolytic (*17*). On the other hand, somatic cells are more dependent on mitochondrial respiration

(*18, 19*). Inhibition of glycolysis drives the exit from pluripotency, while stimulation of glycolysis

enhances somatic cell reprogramming to induced PSCs (iPSCs) (*18, 20*), suggesting that the

balance between glycolysis and mitochondrial respiration instructs cell fate transition. To date,

how metabolic states are established during development and whether they instruct cell identity

remain enigmatic.

To investigate the functional relationship of MLL3/4, cellular metabolism, and PSC

differentiation, we examined the impact of MLL3/4 loss on glycolysis and mitochondrial

respiration in ESCs. Both extracellular acidification rate (ECAR) and oxygen consumption rate

(OCR) are reduced in MLL3/4-null ESCs. Stable isotope tracing reveals a reduction in the labeling

of specific metabolites in both glycolysis and the TCA cycle upon MLL3/4 loss. MLL3/4 are

required for the expression of hexokinase 2 (HK2) and for proper lipoylation of dihydrolipoamide

S-succinyltransferase (DLST), the E2 subunit of the oxoglutarate dehydrogenase complex

(OGDHc). Moreover, combined overexpression of HK2 and OGDH partially rescues the

metabolic and differentiation defects caused by MLL3/4 loss. Our results provide novel insight

into understanding the crosstalk between epigenetic machineries and cellular metabolism.

Importantly, they may pave the way for developing novel strategies to target human diseases driven

by enhancer deregulation.

Results

MLL3/4 loss dampens glycolytic activity in ground-state ESCs

We previously found that the loss of MLL3/4 in ESCs reduced the steady-state abundance of glycolytic intermediates, including G6P and F6P (*14*), suggesting that MLL3/4 regulate glycolysis in naive PSCs. To directly test this idea, we measured the ECAR of wildtype (WT) and MLL3/4 double knockout (DKO) ESCs (*14*) using a glycolysis stress assay. MLL3/4 deficiency markedly reduced both basal glycolysis and glycolytic capacity (Fig. 1A-C), suggesting that MLL3/4 are required to maintain the glycolytic activity in naive PSCs. To determine if the decreased steady-state abundance of glycolytic metabolites reflects impaired glucose metabolism, we next performed stable isotope tracing with $[^{13}C_6]$-glucose in WT and MLL3/4 DKO ESCs, followed by liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis (Fig. 1D). Consistent with results in steady-state metabolomics, the incorporation of ^{13}C into F6P/G1P (glucose-1-phosphate)/G6P (m+6) was significantly reduced by MLL3/4 loss (Fig. 1E and Fig. S1A). Moreover, MLL3/4 loss reduced labeling of lactate (m+3) from $[^{13}C_6]$-glucose (Fig. 1F and Fig. S1B), corresponding to the decreased glycolytic activity in MLL3/4 DKO ESCs. To determine whether this defect was caused by reduced glucose import, we measured glucose uptake and found no significant difference between WT and MLL3/4 DKO ESCs (Fig. 1G). These findings indicate that MLL3/4 loss does not impair glucose entry into cells but instead disrupts early rate-limiting steps in glycolysis, thereby suppressing glycolysis in ESCs.

To elucidate the mechanism underlying this defect, we examined RNA-seq data (*14*) for genes encoding glycolytic enzymes involved in the production and utilization of G6P and F6P. Among these, *Hk2*, which encodes hexokinase 2, the enzyme that catalyzes the first committed step of glycolysis by converting glucose to G6P, was significantly downregulated in MLL3/4 DKO

ESCs (Fig. S1C). On the other hand, the expression levels of genes encoding glucose-6-phosphate

isomerase (*Gpi1*), phosphofructokinase 1 (*Pfkl* and *Pfkp*), and phosphofructokinase 2 (*Pfkfb1* and

Pfkfb2) were largely unchanged (Fig. S1C). Consistent with the transcriptomic data, western

blotting analysis showed that HK2 protein levels were markedly reduced by MLL3/4 loss, whereas

levels of HK1 (hexokinase 1), the paralog of HK2, remained largely unaffected (Fig. 1H). To

investigate how MLL3/4 regulate *Hk2* expression, we analyzed ChIP-seq datasets (*7, 14*) and

found at least six putative enhancers near the *Hk2* locus, which were bound by MLL4 and exhibited

reduced H3K27ac levels in MLL3/4 DKO ESCs (Fig. S1D). These data suggest that MLL3/4 could

promote *Hk2* expression through these putative enhancers. Taken together, our results indicate that

glycolysis in naive PSCs requires MLL3/4 and that the decreased HK2 levels may be responsible

for the defective glycolysis caused by MLL3/4 loss.

MLL3/4 are required for mitochondrial respiration in ground-state ESCs

Metabolic bivalency, defined by the concurrent use of glycolysis and mitochondrial respiration, is

one of the hallmarks of naive PSCs (*21, 22*). To examine the role of MLL3/4 in regulating

mitochondrial respiration in naive PSCs, we measured the OCR of WT and MLL3/4 DKO ESCs

using a mitochondrial stress assay. Interestingly, MLL3/4 deficiency significantly reduced both

basal and maximal respiration (Fig. 2A-C), suggesting that MLL3/4 are required to maintain the

bivalent metabolic program of naive PSCs. To define where mitochondrial metabolism was

disrupted, we performed stable isotope tracing with $[^{13}C_6]$-glucose labeling (Fig. 2D). MLL3/4

loss did not have a significant impact on the labeling of citrate/isocitrate (m+2) and alpha-

ketoglutarate (α-KG; m+2) (Fig. 2E-F and Fig. S2A-B). In contrast, labeling of succinate (m+2),

fumarate (m+2), and malate (m+2) was significantly reduced in MLL3/4 DKO ESCs (Fig. 2G-I

and Fig. S2C-E). These findings suggest that MLL3/4 loss disrupts TCA cycle activity downstream

of α-KG, thereby contributing to the impaired mitochondrial respiration.

To further test this model, we traced glutamine carbon entry into the TCA cycle using

$[^{13}C_5]$-glutamine (Fig. 2J). Although MLL3/4 loss did not significantly affect labeling of glutamate

(m+5) (Fig. 2K and Fig. S2F), that of α-KG (m+5) was significantly reduced in MLL3/4 DKO

ESCs (Fig. 2L and Fig. S2G). This reduction may reflect impaired conversion of glutamate to α-

KG, although the expression levels of *Glud1*, which encodes glutamate dehydrogenase (GDH), the

enzyme catalyzing the conversion of glutamate to α-KG, were comparable between WT and

MLL3/4 DKO ESCs (Fig. S2L). Consistent with results from $[^{13}C_6]$-glucose labeling, labeling of

succinate (m+4), fumarate (m+4), and malate (m+4) from $[^{13}C_5]$-glutamine was significantly

reduced in MLL3/4 DKO ESCs, while that of citrate/isocitrate (m+4) was largely unchanged (Fig.

2M-P and Fig. S2H-K). Furthermore, expression levels of multiple genes encoding enzymes

involved in α-KG and downstream TCA cycle metabolism were comparable between MLL3/4

DKO and WT ESCs (Fig. S2L). In addition, protein levels of OGDH, SUCLA2, SUCLG1,

SUCLG2, and SDHA remained largely unperturbed upon MLL3/4 loss (Fig. 2Q and Fig. S2M-N),

suggesting that MLL3/4 regulate TCA cycle metabolites via mechanisms other than transcriptional

control of metabolic enzymes.

To further elucidate the molecular mechanisms by which MLL3/4 regulate the TCA cycle,

we examined the protein levels of DLST and dihydrolipoamide dehydrogenase (DLD), E2 and E3

components of the OGDHc. MLL3/4 loss had little impact on their levels (Fig. 2R-S), suggesting

that MLL3/4 are unlikely to regulate mitochondrial respiration via modulating the abundance of

OGDHc components. Lipoylated DLST is essential for OGDHc activity and for the conversion of

α-KG to succinyl-CoA (*23-25*). To understand if MLL3/4 loss impairs the activity of DLST, we

examined the levels of lipoylated proteins in WT and MLL3/4 DKO ESCs using a specific anti-

lipoate antibody (*26*). MLL3/4 loss led to a reproducible decrease of the lipoylated protein species

with lower molecular weight, attributed to lipoylated DLST based on its size (Fig. 2T). The

decrease in lipoylation was specific to DLST as levels of other abundantly lipoylated proteins in

the mitochondria, such as dihydrolipoamide acetyltransferase (DLAT) in the pyruvate

dehydrogenase complex and dihydrolipoamide branched-chain transacylase (DBT) in the

branched-chain alpha-keto acid dehydrogenase complex (*24, 25*), were comparable in MLL3/4

DKO and WT ESCs (Fig. 2T). Taken together, these results suggest that MLL3/4 loss impairs

mitochondrial respiration and disrupts metabolic flux through reducing DLST lipoylation in naive

PSCs.

Combined overexpression of HK2 and OGDH rescues metabolic defects caused by MLL3/4

loss

To determine if MLL3/4 regulate glycolysis and mitochondrial respiration in ESCs through

activating HK2, we overexpressed 3×HA-tagged HK2 in MLL3/4 DKO ESCs with a doxycycline

(DOX)-inducible system (*27, 28*) (Fig. S3A). HK2 overexpression increased basal glycolysis and

glycolytic capacity in MLL3/4 DKO cells (Fig. S3B-D); however, OCR during basal respiration

was reduced upon HK2 expression (Fig. S3E-G), suggesting that HK2 overexpression renders

MLL3/4 DKO ESCs more dependent on glycolysis for energy production.

Since HK2 overexpression led to an expected increase in ECAR but a reduction in OCR,

we sought to restore both glycolysis and mitochondrial respiration in MLL3/4 DKO ESCs to study

the functional relationship of MLL3/4, metabolic state, and the differentiation capability of PSCs.

We hypothesized that the decrease of ^{13}C incorporation in succinate, fumarate, and malate in both

$[^{13}C_6]$-glucose and $[^{13}C_5]$-glutamine labelled MLL3/4-null ESCs results from the defective

OGDHc. Although OGDH levels were comparable between MLL3/4 DKO and WT ESCs, OGDH

overexpression could potentially compensate for the functionally impaired OGDHc and reduced

TCA cycle metabolites in MLL3/4-null ESCs. We therefore overexpressed 3×HA-tagged HK2 and

3×FLAG-tagged OGDH together in MLL3/4 DKO ESCs using DOX-inducible expression vectors

carrying distinct drug resistance genes (Fig. 3A). Increased levels of HK2 and OGDH led to

elevated rates of basal glycolysis, glycolytic capacity, and both basal and maximal respiration (Fig.

3B-G). Furthermore, stable isotope tracing by labeling MLL3/4 DKO and HK2/OGDH double-

overexpressing (DOE) MLL3/4 DKO cells with $[^{13}C_6]$-glucose indicated that labeling of

metabolites reduced by MLL3/4 loss such as F6P/G1P/G6P (m+6), lactate (m+3), succinate (m+2),

and fumarate (m+2) was partially restored by HK2/OGDH DOE (Fig. 3H-M, Fig. S4A-B, and Fig.

S4E-G). As expected, we did not observe a significant difference in the labeling of citrate/isocitrate

(m+2) or α-KG (m+2) (Fig. 3J and Fig. S4C-E). Notably, malate (m+2) labeling was not

significantly increased upon HK2/OGDH DOE in MLL3/4 DKO cells (Fig. 3M and Fig. S4H),

suggesting that there are additional regulatory layers on the levels of malate besides HK2 and

OGDHc. Taken together, these results suggest that MLL3/4 regulate glycolysis and mitochondrial

respiration through activating *Hk2* expression and maintaining the normal function of the OGDHc.

Combined overexpression of HK2 and OGDH rescues differentiation defects caused by

MLL3/4 loss

To assess the impact of MLL3/4 loss on cell fate transition, we generated embryoid bodies (EBs)

from WT and MLL3/4 DKO ESCs. As expected, MLL3/4 loss led to a decrease in the size of day

6 EBs (Fig. 4A-B), indicating that MLL3/4 are required for PSC differentiation. RNA-seq analysis

showed that MLL3/4 loss caused the significant deregulation of 4,547 genes in day 6 EBs (Fig.

4C). Among genes significantly downregulated in MLL3/4 DKO EBs, 52.3% (1,192 of 2,280)

were significantly upregulated during WT EB differentiation (Fig. 4D), suggesting that these genes

are dependent on MLL3/4 to be activated during differentiation. We thus defined them as MLL3/4-

dependent differentiation genes. Gene ontology (GO) analysis showed that the top 300 upregulated

genes of the 1,192 MLL3/4-dependent differentiation genes were enriched with embryogenesis-

related genes and known MLL4 targets essential for actin cytoskeleton organization (*7*) (Fig. 4E),

further indicating that MLL3/4 loss impairs PSC differentiation. Indeed, pluripotency genes were

significantly upregulated in MLL3/4 DKO EBs and marker genes of meso- and endo-derm

lineages were downregulated in EBs by MLL3/4 loss (Fig. S5A-B). These data demonstrate that

MLL3/4 are required for PSC differentiation and related transcriptional reconfiguration.

To examine if the partial restoration of glycolysis and mitochondrial respiration had any

impact on the differentiation capacity of ESCs lacking MLL3/4, we treated MLL3/4 DKO and

HK2/OGDH DOE MLL3/4 DKO cells with DOX and differentiated them toward EBs. EB

differentiation revealed that HK2/OGDH DOE led to a significant increase in the size of day 6

EBs (Fig. 4F-G), suggesting that restoring glycolysis and mitochondrial respiration rescues

differentiation defects caused by MLL3/4 loss. Moreover, RNA-seq analysis identified 1,778

significantly deregulated genes by HK2/OGDH DOE compared with MLL3/4 DKO ESCs treated

with DOX (Fig. 4H). Gene set enrichment analysis (GSEA) and hierarchical clustering analysis of

MLL3/4-dependent differentiation genes (Fig. 4D) revealed a significant upregulation of them

upon HK2/OGDH DOE (Fig. 4I-J). Indeed, 34.3% of genes upregulated by HK2/OGDH DOE

(300 of 875) were dependent on MLL3/4 to be activated during differentiation (Fig. 4K),

suggesting that the impaired metabolic state contributes to differentiation defects caused by

MLL3/4 loss. GO analysis of the 300 significantly rescued genes showed enrichment of genes

related to embryo morphogenesis and mesodermal development such as the development of renal

and skeletal systems (Fig. 4L). Consistently, HK2/OGDH DOE led to the significant

downregulation of pluripotency genes and the significant upregulation of differentiation marker

genes (Fig. S5C-D). Taken together, these results suggest that defects in glycolysis and

mitochondrial respiration contribute to differentiation failures of naive PSCs lacking MLL3/4.

Discussion

Discussion

Epigenetic regulators, including histone modifiers, play central roles in regulating cell fate

transition during PSC differentiation, somatic cell reprogramming, and normal development (*29-*

35); however, the underlying mechanisms remain incompletely understood, in part because many

epigenetic regulators function independently from their catalytic activity (*6, 7, 28, 36-43*). Besides

being essential for cellular differentiation and mammalian embryogenesis (*11, 44*), loss-of-

function mutations in MLL3 and MLL4 drive congenital disorders such as Kleefstra syndrome 2

and Kabuki syndrome, as well as multiple types of cancer (*1, 4, 13, 45*). Recent advances reveal a

novel link between MLL3/4 and cellular metabolism; however, whether and how the roles of

MLL3/4 in regulating cell fate and metabolic pathways intersect are unclear. Here, we unveil a

previously unrecognized function of MLL3/4 in regulating glycolysis and mitochondrial

respiration in murine ESCs. Our findings suggest that MLL3/4 govern cell fate transition during

the differentiation of naive PSCs through regulating the expression levels of *Hk2* and the function

of OGDHc, two rate-limiting nodes in glycolysis and the TCA cycle, respectively. Importantly,

restoring defective glycolysis and mitochondrial respiration may be an effective strategy to treat

diseases driven by MLL3/MLL4 loss-of-function.

We find that MLL3/4 loss causes decreased glycolysis and mitochondrial respiration in

ESCs. Metabolic bivalency is a defining feature of naive PSCs such as murine ESCs (*22*). The

decrease of glycolysis in MLL3/4-null cells may hinder the cells from exiting naive pluripotency,

as formative PSCs and primed PSCs are highly glycolytic (*17*). On the other hand, inhibiting

glycolysis promotes differentiation as somatic cells have much elevated levels of oxidative

phosphorylation compared with PSCs (*18-20*). Therefore, the dampened mitochondrial respiration

caused by MLL3/4 loss may impair the differentiation of PSCs into germ layers and/or specific

lineages. Our data provide a mechanistic explanation for the failure of MLL4-null ESCs to exit

naive pluripotency and for the severe gastrulation defects observed in MLL3/4-null embryos (*7,*

10, 46).

255   Our data support that MLL3/4 facilitate glycolysis in naive ESCs through maintaining the

expression of *Hk2*, which encodes the rate-limiting enzyme hexokinase 2 in glycolysis. It is

possible that MLL3/4 regulate *Hk2* expression through enhancers as we identified multiple

putative *cis*-regulatory regions that may be directly regulated by MLL3/4. Understanding whether

and how MLL3/4 function through these putative enhancers to control the expression levels of

Hk2 is an interesting topic. Shadow enhancers have been reported in different organisms including

Drosophila, mouse, and human (*47-49*). Generating individual and compound enhancer mutations

will elucidate how the putative enhancers regulate *Hk2*. Combination of MLL3/4 loss-of-function

and enhancer mutations will further reveal the functional relationship between MLL3/4 and

putative enhancers in regulating *Hk2*. These future studies are essential for understanding the

mechanisms by which MLL3/4 regulate glycolysis in naive PSCs.

266   Interestingly, overexpressing HK2 alone only partially restores glycolysis in ESCs lacking

MLL3/4, suggesting that MLL3/4 support glycolysis in naive PSCs through additional

mechanisms. Determining whether other critical regulators of glycolysis such as

phosphofructokinase, pyruvate kinase, and lactate dehydrogenase play a role in dampening

glycolysis in the context of MLL3/4 loss-of-function is an important future study. Genome-wide

and targeted genetic screens may also identify additional factors that rescue glycolysis in MLL3/4-

null ESCs. It is noteworthy that HK2 overexpression leads to a decrease of OCR in MLL3/4 DKO

ESCs, suggesting that these cells are more dependent on glycolysis as an energy source in

comparison to oxidative phosphorylation under high levels of HK2. Since steady-state acetyl-CoA

abundance was reduced in MLL3/4 DKO ESCs (*14*) and our stable isotope tracing assays were not

able to detect labeled acetyl-CoA, it is interesting to examine if and how MLL3/4 loss affects the

mitochondrial transport of pyruvate in ESCs in the future. Defining the role of MLL3/4 in other

pathways leading to the synthesis of acetyl-CoA such as fatty acid metabolism may also facilitate

understanding how MLL3/4 regulate the conjunction of glycolysis and the TCA cycle.

The reduction in ^{13}C incorporation into succinate, fumarate, and malate in both [^{13}C$_6$]-

glucose and [^{13}C$_5$]-glutamine labeling of MLL3/4 DKO ESCs suggests a functionally defective

OGDHc caused by MLL3/4 loss. Indeed, the levels of lipoylated DLST were decreased in

MLL3/4-null cells, suggesting a reduced activity of the OGDHc. It is likely that the decrease of

lipoylation is specific to DLST as MLL3/4 loss has little impact on the levels of lipoylated DLAT.

It was previously shown that the α/β hydrolase ABHD11 binds OGDHc and maintains the activity

of lipoylated DLST by preventing the formation of lipoyl adducts (*26*). It will be interesting to

investigate the functional relationship of MLL3/4, ABHD11, and OGDHc in the future to

understand how MLL3/4 regulate the activity of OGDHc. It is also noteworthy that ^{13}C

incorporation into α-KG (m+5) is impaired in MLL3/4 DKO ESCs labelled with [^{13}C$_5$]-glutamine,

although α-KG (m+2) levels through [^{13}C$_5$]-glucose labeling and steady-state levels of α-KG were

largely unchanged by MLL3/4 loss (*14*). Besides glutamate dehydrogenase, glutamate pyruvate

transaminase and glutamate oxaloacetate transaminase also can convert glutamate to α-KG.

MLL3/4 may also regulate the transport of cytosolic glutamate into mitochondria. Future studies

on the role of MLL3/4 in regulating glutamate transport and the conversion of glutamate to α-KG

would be important to understand the crosstalk between MLL3/4 and glutaminolysis.

α-KG is an essential co-substrate of Fe(II)/α-KG-dependent dioxygenases including

Jumonji C (JmjC) domain-containing histone demethylases and the ten-eleven translocation (TET)

family of DNA hydroxylases (*50, 51*), playing a role in regulating the levels of methylation on

different lysine residues of histones and 5-methyl Cytosine (5mC) on DNA in mammalian cells.

Due to their structural similarity to α-KG, 2-hydroxyglutarate, succinate, and fumarate inhibit α-

KG-dependent dioxygenases by acting as competitive inhibitors of α-KG (*52, 53*). Our results that

MLL3/4 loss in ESCs leads to decreased levels of succinate and fumarate, through steady-state

metabolomics (*14*) and stable isotope tracing with $[^{13}C_6]$-glucose and $[^{13}C_5]$-glutamine, suggest a

possibility that activities of multiple histone demethylases and TETs are increased in cells with

MLL3/4 loss-of-function. Such a possibility provides a mechanistic explanation for the DNA

hypomethylation observed in MLL4-depleted B cells in mouse and MLL4 mutant prostate cancers

in human (*54*). Understanding the impact of the potential alteration in levels of histone methylation

and DNA methylation on the differentiation capability of MLL3/4 mutant PSCs is very interesting

and may provide insight into understanding the molecular mechanisms underlying the functional

relationship of metabolic state, epigenetic machineries, and cell fate transition.

In summary, our study reveals a previously unappreciated crosstalk between epigenetic

modifiers exemplified by MLL3/4 and glycolysis and mitochondrial respiration in regulating cell

fate. Our findings on MLL3/4 regulating the levels of HK2 and lipoylated DLST could pave the

way for designing therapeutic strategies to target human diseases driven by loss-of-function

mutations of MLL3/4.

Methods

Antibodies

The following primary antibodies were used in this study: anti-HK2 (Cell Signaling Technology 2867), anti-HK1 (Cell Signaling Technology 2024), anti-OGDH (Proteintech 17110-1-AP), anti-SDHA (Abcam 14715), anti-SUCLA2 (Proteintech 12627-1-AP), anti-SUCLG1 (Proteintech 14923-1-AP), anti-SUCLG2 (Proteintech NBP1-32521), anti-FLAG (MiliporeSigma F3165), anti-HA (MiliporeSigma H3663), anti-DLST (Cell Signaling Technology 5556), anti-DLD (Santa Cruz 365977), anti-Lipoic acid (Calbiochem 437695), anti-Tubulin (Developmental Studies Hybridoma Bank E7), and anti-HSP90 (Santa Cruz 13119).

ESC Culture and EB differentiation

ESCs were cultured in serum-free medium supplemented with 2i/LIF as previously described (*55*). For the generation of HK2 overexpressing and HK2/OGDH DOE MLL3/4 DKO ESCs, cells were transfected with DOX-inducible PiggyBac vectors (*27*) as previously described (*28*) using an Amaxa Nucleofector (Lonza). EB differentiation was performed using the hanging-drop method as previously described (*28*). For DOX-induced overexpression, ESCs were treated with 100 ng/ml doxycycline for 48 hours prior to downstream assays. For differentiation experiments, doxycycline (100 ng/ml) were added 24 hours prior to EB formation and maintained throughout the 6-day differentiation period.

Seahorse extracellular flux assays

ESCs were seeded in Seahorse XF microplates (Agilent) and cultured overnight with 2i/LIF serum-free medium. Cells were then incubated in Seahorse XF assay medium (Agilent) at 37°C in a non-

CO_2 incubator for 1 hour. For the Glycolysis Stress Test (Agilent 103020-100), glucose (10 mM),

oligomycin (2 µM), and 2-DG (50 mM) were sequentially injected. For the Mitochondrial Stress

Test (Agilent 103015-100), oligomycin (1.5 µM), FCCP (0.5 µM), and Rot/AA (0.5 µM) were

sequentially injected. Cell Titer Glo assay (Promega) was performed in parallel on replica plates

to determine the cell number of each well. Individual raw ECAR and OCR values recorded by the

Seahorse XF Analyzer were normalized against the average Cell Titer Glo values of control wells

to derive normalized ECAR and normalized OCR values. Wave software (Agilent) was used to

extract individual ECAR and OCR measurements, which were subsequently used to generate the

bar plots.

Stable isotope tracing and metabolomics

Stable isotope labeling and metabolite extraction were performed as previously described (*14*).

ESCs were incubated with media containing 25 mM [$^{13}C_6$]-glucose or 4 mM [$^{13}C_5$]-glutamine for

2 hours before methanol extraction. Metabolomic analyses were performed at the Northwestern

Metabolomics Core Facility.

Glucose uptake assay

Glucose update was measured using the Glucose Uptake-Glo Assay Kit (Promega J1342)

according to the manufacturer's instructions. In brief, 2×10^4 cells were seeded on each well of a

96-well flat bottom plate with opaque walls and clear bottoms (Corning 7200587). After 16 hours

of culture, cells were incubated in glucose-free medium for 1 hour at 37°C in a CO_2 incubator.

Cells were then treated with 1 mM 2-DG for 10 minutes at room temperature, followed by the

treatment of stop buffer, neutralization buffer, and detection reagent following the instructions of

the manufacturer. After 2.5 hours of dark-conditioned incubation, the luminescence was measured

using a GloMax luminometer (Promega). The luminescence data were normalized by cell numbers

measured with the Cell Titer Glo assay (Promega) on replica plates in parallel.

RNA-seq and analysis

Total RNA was extracted using TRIzol reagent (Thermo Fisher) following the manufacturer's

instructions. Each cell line was analyzed in at least two biological replicates. Extracted RNA were

treated with DNase I (Sigma) and purified with the RNeasy Plus Mini Kit (Qiagen). Ribosomal

RNA was removed from the purified RNA with the NEBNext rRNA Depletion Kit (New England

BioLabs). RNA libraries were then prepared with the NEBNext Ultra II Directional RNA Library

Prep Kit (New England BioLabs). The pooled libraries were sequenced with 150 bp paired-end

reads on the NovaSeq platfrom (Illumina).

Raw reads were processed with Trimmomatic v0.39(*56*) to remove adaptors and low-

quality reads with the parameter "-q 30". The cleaned reads were aligned to the mm9 genome

assembly using STAR v2.5.1 with default parameter (*57*). Reads were normalized to total read

counts per million (cpm) and visualized as bigwig-formatted coverage tracks using deepTools

v3.5.6 (*58*). Gene expression was quantified using HTseq V2.0.2 (*59*), and differential gene

expression analysis was performed with EdgeR v4.0.16 (*60*). Significantly differentially expressed

genes were identified with Benjamini-Hochberg-adjusted p value cutoff < 0.01 and a log fold

change threshold greater than |1|. hierarchical clustering heatmaps were generated using the ggpubr

v0.6.0 (https://rpkgs.datanovia.com/ggpubr/). GSEA plots were generated with clusterProfiler

v4.10.1 (*61*). Gene ontology (GO) analysis was performed using Metascape v3.5.20260201 (*62*).

Acknowledgements

We thank the Cao lab members for their helpful suggestions and discussions. We thank Dr. Peng Gao and the Northwestern Metabolomics Core Facility for performing the metabolomics study. We thank Emmalee W. Cooke for technical support. We thank Yalu Zhou for bioinformatic analysis support. This study was supported in part by grants R35GM150668, R00HD094906, and P30CA043703 from the National Institutes of Health to K.C.

393 **Author contributions**

394 K.C. conceived the study. S.M.N., I.B., and K.C. designed the research. S.M.N., Y.J., and M.Y.

395 performed experiments. S.M.N., Y.J., M.Y., and K.C. analyzed the data. S.M.N. and K.C. wrote

396 the manuscript. C.A.L. and I.B. critically read the manuscript.

Figure legends

Figure 1. The loss of MLL3/4 leads to impaired glycolytic activity in ground-state ESCs.

(A) Representative ECAR traces of WT and MLL3/4 DKO ESCs measured by the Seahorse XF glycolysis stress test assay. Data were normalized by the number of cells at the time of the assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment was repeated three times independently. 2-DG: 2-Deoxy-D-glucose.

(B-C) Bar plots of ECAR showing the levels of glycolysis (B) and glycolytic capacity (C) in WT and MLL3/4 DKO ESCs. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(D) Schematic of glycolysis illustrating the fate of ^{13}C carbons upon the incubation with $[^{13}C_6]$-glucose.

(E-F) Bar plots showing the levels of F6P/G1P/G6P (m+6, E) and lactate (m+3, F) in WT and MLL3/4 DKO ESCs labeled with $[^{13}C_6]$-glucose. n=3 biological replicates per sample. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(G) Bar plots showing the levels of glucose uptake in WT and MLL3/4 DKO ESCs. n=6 replicates per sample. Experiments were repeated three times independently and representative results are shown. Data were normalized by the number of cells at the time of the assay and are presented as mean ± S.D. p-values were calculated using Student's t-test. RLU: relative light unit.

(H) Western blots showing the levels of HK1 and HK2 in WT and MLL3/4 DKO ESCs. 10 and 20 µg of cell lysates were loaded for each sample. Experiments were repeated three times independently and representative results are shown. Protein levels in each lane were quantified by normalizing with Tubulin signals of the corresponding lane.

Figure 2. The loss of MLL3/4 leads to impaired mitochondrial respiration in ground-state ESCs.

(A) Representative OCR traces of WT and MLL3/4 DKO ESCs measured by the Seahorse XF mitochondrial stress test assay. Data were normalized by the number of cells at the time of the assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment was repeated three times independently. FCCP: Carbonyl Cyanide-4-(trifluoromethoxy) phenylhydrazone. ROT/AA: Rotenone and Antimycin A.

(B-C) Bar plots showing the levels of OCR during basal (B) and maximal (C) respiration in WT and MLL3/4 DKO ESCs. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(D) Schematic of the TCA cycle illustrating the fate of ^{13}C carbons upon the incubation with [$^{13}C_6$]-glucose.

(E-I) Bar plots showing levels of citrate/isocitrate (m+2, E), α-KG (m+2, F), succinate (m+2, G), fumarate (m+2, H), and malate (m+2, I) in WT and MLL3/4 DKO ESCs labeled with [$^{13}C_6$]-glucose. n=3 biological replicates per sample. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(J) Schematic of the TCA cycle illustrating the fate of ^{13}C carbons upon the incubation with [$^{13}C_5$]-glutamine.

(K-P) Bar plots showing levels of glutamate (m+5, K), α-KG (m+5, L), succinate (m+4, M), fumarate (m+4, N), malate (m+4, O), and citrate/isocitrate (m+4, P) in WT and MLL3/4 DKO ESCs labeled with [$^{13}C_5$]-glutamine. n=3 biological replicates per sample. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(Q-T) Western blots showing the levels of OGDH (Q), DLST (R), DLD (S), and lipoylated proteins (T) in WT and MLL3/4 DKO ESCs. 10 and 20 μg of cell lysates were loaded for each sample. Experiments were repeated three times independently and representative results are shown. Protein levels of OGDH (in Q) , DLST (in R), DLD (in S), and lipoylated DLST (in T) were quantified by normalizing with Tubulin or HSP90 signals of the corresponding lane.

Figure 3. Combined overexpression of HK2 and OGDH rescues metabolic defects of MLL3/4 DKO ESCs.

(A) Western blots showing levels of HK2, 3×HA-tagged HK2, OGDH, and 3×FLAG-tagged OGDH in MLL3/4 DKO ESCs and MLL3/4 DKO ESCs overexpressing HK2/OGDH treated with or without DOX. 10 and 20 μg of cell lysates were loaded for each sample. Experiments were repeated three times independently and representative results are shown. Protein levels in each lane of the anti-HK2 and anti-OGDH blots were quantified by normalizing with Tubulin signals of the corresponding lane.

(B) Representative ECAR traces of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX) measured by the Seahorse XF glycolysis stress test assay. Data were normalized by the number of cells at the time of the assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment was repeated three times independently.

(C-D) Bar plots of ECAR showing the level of glycolysis (C) and glycolytic capacity (D) in MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX). Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(E) Representative OCR traces of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and

MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX) measured by the Seahorse XF

mitochondrial stress test assay. Data were normalized by the number of cells at the time of the

assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment

was repeated three times independently.

(F-G) Bar plots showing the levels of OCR during basal (F) and maximal (G) respiration in

MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing

HK2/OGDH (DOE+DOX). Data are presented as mean ± S.D. p-values were calculated using

Student's t-test.

(H-M) Bar plots showing levels of F6P/G1P/G6P (m+6, H), lactate (m+3, I), α-KG (m+2, J),

succinate (m+2, K), fumarate (m+2, L), and malate (m+2, M) in MLL3/4 DKO ESCs treated with

DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX) labeled

with [$^{13}C_6$]-glucose. n=3 biological replicates per sample. Data are presented as mean ± S.D. p-

values were calculated using Student's t-test.

Figure 4. Combined overexpression of HK2 and OGDH rescues differentiation defects

caused by MLL3/4 loss.

(A) Representative images of day 6 WT and MLL3/4 DKO EBs. Scale bar: 250μm.

(B) Quantification of EB sizes in (A). n=50 for each genotype. Experiments were repeated three

times independently and representative results are shown.

(C) Correlation plots of RNA-seq data between day 6 MLL3/4 DKO and WT EBs. Data were

derived from two biological replicates. Statistical significance was determined by two-sided Wald

test and p-values were corrected for multiple testing using the Benjamini–Hochberg method.

Significantly up- and downregulated genes (adjusted p<0.01, $|$logFC$|$>1) were labeled in red and

blue with numbers of genes noted, respectively.

(D) Venn diagram identifying 1,192 MLL3/4-dependent differentiation genes.

(E) GO analysis of top 300 MLL3/4-dependent differentiation genes.

(F) Representative images of day 6 MLL3/4 DKO EBs treated with DOX (DKO+DOX) and

MLL3/4 DKO EBs overexpressing HK2/OGDH (DOE+DOX). Scale bar: 250μm.

(G) Quantitation of EB sizes in (F). n=50 for each genotype. Experiments were repeated three

times independently and representative results are shown.

(H) Correlation plots of RNA-seq data between MLL3/4 DKO EBs treated with DOX

(DKO+DOX) and MLL3/4 DKO EBs overexpressing HK2/OGDH (DOE+DOX).

(I) GSEA analysis of MLL3/4-dependent differentiation genes (1,192 in D) comparing MLL3/4

DKO EBs treated with DOX (DKO+DOX) and MLL3/4 DKO EBs overexpressing HK2/OGDH

(DOE+DOX). RES: running enrichment score; RLM: ranked list metric; NES: normalized

enrichment score; FDR: false discovery rate.

(J) Hierarchical clustering analysis of the expression levels of 1,192 MLL3/4-dependent

differentiation genes in MLL3/4 DKO EBs treated with DOX (DKO+DOX) and MLL3/4 DKO

EBs overexpressing HK2/OGDH (DOE+DOX). FPKM: Fragments per kilobase of transcript per

million mapped reads.

(K) Venn diagram showing the overlap of MLL3/4-dependent differentiation genes and

significantly upregulated genes in MLL3/4 DKO EBs overexpressing HK2/OGDH (red in H).

(L) GO analysis of 300 DOE-rescued differentiation genes (overlapping genes in K).

Figure S1. The impact of MLL3/4 loss on glycolytic metabolites, glycolytic gene expression, and putative enhancers at the *Hk2* locus.

(A-B) Stable isotope tracing of WT and MLL3/4 DKO ESCs labeled with $[^{13}C_6]$-glucose. The levels of isotopologues (m+0 to m+6) of F6P/G6P/G1P (A) and lactate (B) were shown as stacked bar plots.

(C) Relative expression levels of genes coding key glycolytic enzymes involved in the production and utilization of G6P, F6P, and lactate. CPM (counts per million reads) from RNA-seq data (*14*) of MLL3/4 DKO ESCs were normalized to those of WT ESCs. p-values were determined by two-sided Wald test and corrected using the Benjamini–Hochberg method.

(D) Genome browser view of MLL4 ChIP-seq signals in WT ESCs, and that of H3K27ac and H3K4me3 ChIP-seq signals in WT and MLL3/4 DKO ESCs at the *Hk2* locus. Putative enhancers bound by MLL4 around *Hk2* gene were highlighted.

Figure S2. The impact of MLL3/4 loss on TCA cycle metabolites and the expression of TCA cycle-related genes.

(A-E) Stable isotope tracing of WT and MLL3/4 DKO ESCs labeled with $[^{13}C_6]$-glucose. The levels of isotopologues (m+0 to m+6) of citrate/isocitrate (A), α-KG (B), succinate (C), fumarate (D), and malate (E) were shown as stacked bar plots.

(F-K) Stable isotope tracing of WT and MLL3/4 DKO ESCs labeled with $[^{13}C_5]$-glutamine. The levels of isotopologues (m+0 to m+6) of glutamate (F), α-KG (G), succinate (H), fumarate (I), malate (J), and citrate/isocitrate (K) were shown as stacked bar plots.

(L) Relative expression levels of genes coding TCA-cycle enzymes involved in the production and utilization of α-KG, succinate, fumarate, and malate. CPM (counts per million reads) from RNA-

seq data (*14*) of MLL3/4 DKO ESCs were normalized to those of WT ESCs. p-values were

determined by two-sided Wald test and corrected using the Benjamini–Hochberg method.

(M-N) Western blots showing levels of Succinyl-CoA Synthetase subunits (M), SDHA (N, top

blot), and FH (N, middle blot). 10 and 20 μg of cell lysates were loaded for each sample.

Experiments were repeated three times independently and representative results are shown. Protein

levels in each lane were quantified by normalizing with HSP90 signals of the corresponding lane.

Figure S3. HK2 overexpression leads to an increase in glycolysis but a decrease in

mitochondrial respiration in MLL3/4 DKO ESCs.

(A) Western blots showing the induction of 3×HA-tagged HK2 in HK2-overexpressing MLL3/4

DKO ESCs by DOX. 10 and 20 μg of cell lysates were loaded for each sample. Experiments were

repeated three times independently and representative results are shown. Protein levels of HK2 in

each lane were quantified by normalizing with Tubulin signals of the corresponding lane.

(B) Representative ECAR traces of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and

MLL3/4 DKO ESCs overexpressing HK2 (3×HA-HK2+DOX) measured by the Seahorse XF

glycolysis stress test assay. Data were normalized by the number of cells at the time of the assay

and are represented as mean ± S.D. Each data point had six replicates, and the experiment was

repeated three times independently.

(C-D) Bar plots of ECAR showing the levels of glycolysis (C) and glycolytic capacity (D) in

MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing

HK2 (3×HA-HK2+DOX). Data are presented as mean ± S.D. p-values were calculated using

Student's t-test.

(E) Representative OCR traces of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and

MLL3/4 DKO ESCs overexpressing HK2 (3×HA-HK2+DOX) measured by the Seahorse XF

mitochondrial stress test assay. Data were normalized by the number of cells at the time of the

assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment

was repeated three times independently.

(F-G) Bar plots showing the OCR levels in MLL3/4 DKO ESCs treated with DOX (DKO+DOX)

and MLL3/4 DKO ESCs overexpressing HK2 (3×HA-HK2+DOX) during basal (F) and maximal

(G) respiration. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

Figure S4. Stable isotope tracing of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and

MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX) labeled with [^{13}C$_6$]-glucose.

The levels of isotopologues of F6P/G1P/G6P (A, m+0 to m+6), lactate (B m+0 to m+3),

citrate/isocitrate (m+2 in C and m+0 to m+6 in D), α-KG (E, m+0 to m+5), succinate (F, m+0 to

m+4), fumarate (G, m+0 to m+4), and malate (H, m+0 to m+4) were shown.

Figure S5. The impact of MLL3/4 loss and HK2/OGDH overexpression on the expression

levels of marker genes of pluripotency and differentiation in EBs.

(A-B) Relative expression levels of pluripotency genes (A) and differentiation markers (B) in

MLL3/4 DKO EBs compared with WT EBs. Log2 fold change of RNA-seq signals (CPM) were

shown. Data are presented as mean values ± SD. n = 2 biologically independent experiments. All

p-values were smaller than 0.01 compared with WT EBs.

(C-D) Relative expression levels of pluripotency genes (C) and differentiation markers (D) in

MLL3/4 DKO EBs overexpressing HK2/OGDH (DOE+DOX) compared with MLL3/4 DKO EBs

treated with DOX (DKO+DOX). Log2 fold change of RNA-seq signals (CPM) were shown. Data

are presented as mean values ± SD. n = 2 biologically independent experiments. All p-values were

smaller than 0.01 compared with DKO+DOX EBs.

References

1. S. B. Ng, A. W. Bigham, K. J. Buckingham, M. C. Hannibal, M. J. McMillin, H. I. Gildersleeve, A. E. Beck, H. K. Tabor, G. M. Cooper, H. C. Mefford, C. Lee, E. H. Turner, J. D. Smith, M. J. Rieder, K. Yoshiura, N. Matsumoto, T. Ohta, N. Niikawa, D. A. Nickerson, M. J. Bamshad, J. Shendure, Exome sequencing identifies MLL2 mutations as a cause of Kabuki syndrome. *Nat Genet* **42**, 790-793 (2010).

2. I. D. Krantz, J. McCallum, C. DeScipio, M. Kaur, L. A. Gillis, D. Yaeger, L. Jukofsky, N. Wasserman, A. Bottani, C. A. Morris, M. J. Nowaczyk, H. Toriello, M. J. Bamshad, J. C. Carey, E. Rappaport, S. Kawauchi, A. D. Lander, A. L. Calof, H. H. Li, M. Devoto, L. G. Jackson, Cornelia de Lange syndrome is caused by mutations in NIPBL, the human homolog of Drosophila melanogaster Nipped-B. *Nat Genet* **36**, 631-635 (2004).

3. E. K. Schorry, M. Keddache, N. Lanphear, J. H. Rubinstein, S. Srodulski, D. Fletcher, R. I. Blough-Pfau, G. A. Grabowski, Genotype-phenotype correlations in Rubinstein-Taybi syndrome. *Am J Med Genet A* **146A**, 2512-2519 (2008).

4. T. S. Koemans, T. Kleefstra, M. C. Chubak, M. H. Stone, M. R. F. Reijnders, S. de Munnik, M. H. Willemsen, M. Fenckova, C. Stumpel, L. A. Bok, M. Sifuentes Saenz, K. A. Byerly, L. B. Baughn, A. P. A. Stegmann, R. Pfundt, H. Zhou, H. van Bokhoven, A. Schenck, J. M. Kramer, Functional convergence of histone methyltransferases EHMT1 and KMT2C involved in intellectual disability and autism spectrum disorder. *PLoS Genet* **13**, e1006864 (2017).

5. D. Hu, X. Gao, M. A. Morgan, H. M. Herz, E. R. Smith, A. Shilatifard, The MLL3/MLL4 branches of the COMPASS family function as major histone H3K4 monomethylases at enhancers. *Mol Cell Biol* **33**, 4745-4754 (2013).

6. R. Rickels, H. M. Herz, C. C. Sze, K. Cao, M. A. Morgan, C. K. Collings, M. Gause, Y. H. Takahashi, L. Wang, E. J. Rendleman, S. A. Marshall, A. Krueger, E. T. Bartom, A. Piunti, E. R. Smith, N. A. Abshiru, N. L. Kelleher, D. Dorsett, A. Shilatifard, Histone H3K4 monomethylation catalyzed by Trr and mammalian COMPASS-like proteins at enhancers is dispensable for development and viability. *Nat Genet* **49**, 1647-1653 (2017).

7. K. Cao, C. K. Collings, M. A. Morgan, S. A. Marshall, E. J. Rendleman, P. A. Ozark, E. R. Smith, A. Shilatifard, An Mll4/COMPASS-Lsd1 epigenetic axis governs enhancer function and pluripotency transition in embryonic stem cells. *Sci Adv* **4**, eaap8747 (2018).

8. A. Shilatifard, The COMPASS family of histone H3K4 methylases: mechanisms of regulation in development and disease pathogenesis. *Annu Rev Biochem* **81**, 65-95 (2012).

9. N. D. Heintzman, R. K. Stuart, G. Hon, Y. Fu, C. W. Ching, R. D. Hawkins, L. O. Barrera, S. Van Calcar, C. Qu, K. A. Ching, W. Wang, Z. Weng, R. D. Green, G. E. Crawford, B. Ren, Distinct and predictive chromatin signatures of transcriptional promoters and enhancers in the human genome. *Nat Genet* **39**, 311-318 (2007).

10. J. E. Lee, C. Wang, S. Xu, Y. W. Cho, L. Wang, X. Feng, A. Baldridge, V. Sartorelli, L. Zhuang, W. Peng, K. Ge, H3K4 mono- and di-methyltransferase MLL4 is required for enhancer activation during cell differentiation. *Elife* **2**, e01503 (2013).

11. D. Ashokkumar, Q. Zhang, C. Much, A. S. Bledau, R. Naumann, D. Alexopoulou, A. Dahl, N. Goveas, J. Fu, K. Anastassiadis, A. F. Stewart, A. Kranz, MLL4 is required after implantation, whereas MLL3 becomes essential during late gestation. *Development* **147**, (2020).

12. V. Faundes, W. G. Newman, L. Bernardini, N. Canham, J. Clayton-Smith, B. Dallapiccola, S. J. Davies, M. K. Demos, A. Goldman, H. Gill, R. Horton, B. Kerr, D. Kumar, A. Lehman, S. McKee, J. Morton, M. J. Parker, J. Rankin, L. Robertson, I. K. Temple, S. Clinical Assessment of the Utility of, S. Evaluation as a Service, S. Deciphering Developmental Disorders, S. Banka, Histone Lysine Methylases and Demethylases in the Landscape of Human Developmental Disorders. *Am J Hum Genet* **102**, 175-187 (2018).

13. D. Rots, S. Choufani, V. Faundes, A. J. M. Dingemans, S. Joss, N. Foulds, E. A. Jones, S. Stewart, P. Vasudevan, T. Dabir, S. M. Park, R. Jewell, N. Brown, L. Pais, S. Jacquemont, K. Jizi, C. Ravenswaaij-Arts, H. Y. Kroes, C. Stumpel, C. W. Ockeloen, I. J. Diets, M. Nizon, M. Vincent, B. Cogne, T. Besnard, M. Kambouris, E. Anderson, E. H. Zackai, C. McDougall, S. Donoghue, A. O'Donnell-Luria, Z. Valivullah, M. O'Leary, S. Srivastava, H. Byers, N. Leslie, S. Mazzola, G. E. Tiller, M. Vera, J. J. Shen, R. Boles, V. Jain, E. Brischoux-Boucher, E. Kinning, B. N. Simpson, J. C. Giltay, J. Harris, B. Keren, A. Guimier, P. Marijon, B. B. A. Vries, C. S. Motter, B. A. Mendelsohn, S. Coffino, E. H. Gerkes, A. Afenjar, P. Visconti, E. Bacchelli, E. Maestrini, A. Delahaye-Duriez, C. Gooch, Y. Hendriks, H. Adams, C. Thauvin-Robinet, S. Josephi-Taylor, M. Bertoli, M. J. Parker, J. W. Rutten, O. Caluseriu, H. J. Vernon, J. Kaziyev, J. Zhu, J. Kremen, Z. Frazier, H. Osika, D. Breault, S. Nair, S. M. E. Lewis, F. Ceroni, M. Viggiano, A. Posar, H. Brittain, T. Giovanna, G. Giulia, L. Quteineh, R. Ha-Vinh Leuchter, E. Zonneveld-Huijssoon, C. Mellado, I. Marey, A. Coudert, M. I. Aracena Alvarez, M. G. P. Kennis, A. Bouman, M. Roifman, M. I. Amoros Rodriguez, J. D. Ortigoza-Escobar, V. Vernimmen, M. Sinnema, R. Pfundt, H. G. Brunner, L. Vissers, T. Kleefstra, R. Weksberg, S. Banka, Pathogenic variants in KMT2C result in a neurodevelopmental disorder distinct from Kleefstra and Kabuki syndromes. *Am J Hum Genet* **111**, 1626-1642 (2024).

14. Z. Zhao, K. Cao, J. Watanabe, C. N. Philips, J. M. Zeidner, Y. Ishi, Q. Wang, S. R. Gold, K. Junkins, E. T. Bartom, F. Yue, N. S. Chandel, R. Hashizume, I. Ben-Sahra, A. Shilatifard, Therapeutic targeting of metabolic vulnerabilities in cancers with MLL3/4-COMPASS epigenetic regulator mutations. *J Clin Invest* **133**, (2023).

15. H. Alam, M. Tang, M. Maitituoheti, S. S. Dhar, M. Kumar, C. Y. Han, C. R. Ambati, S. B. Amin, B. Gu, T. Y. Chen, Y. H. Lin, J. Chen, F. L. Muller, N. Putluri, E. R. Flores, F. J. DeMayo, L. Baseler, K. Rai, M. G. Lee, KMT2D Deficiency Impairs Super-Enhancers to Confer a Glycolytic Vulnerability in Lung Cancer. *Cancer Cell* **37**, 599-617 e597 (2020).

16. M. Maitituoheti, E. Z. Keung, M. Tang, L. Yan, H. Alam, G. Han, A. K. Singh, A. T. Raman, C. Terranova, S. Sarkar, E. Orouji, S. B. Amin, S. Sharma, M. Williams, N. S. Samant, M. Dhamdhere, N. Zheng, T. Shah, A. Shah, J. B. Axelrad, N. E. Anvar, Y. H. Lin, S. Jiang, E. Q. Chang, D. R. Ingram, W. L. Wang, A. Lazar, M. G. Lee, F. Muller, L. Wang, H. Ying, K. Rai, Enhancer Reprogramming Confers Dependence on Glycolysis and IGF Signaling in KMT2D Mutant Melanoma. *Cell Rep* **33**, 108293 (2020).

17. W. Zhou, M. Choi, D. Margineantu, L. Margaretha, J. Hesson, C. Cavanaugh, C. A. Blau,
M. S. Horwitz, D. Hockenbery, C. Ware, H. Ruohola-Baker, HIF1alpha induced switch
from bivalent to exclusively glycolytic metabolism during ESC-to-EpiSC/hESC
transition. *EMBO J* **31**, 2103-2116 (2012).

18. C. D. Folmes, T. J. Nelson, A. Martinez-Fernandez, D. K. Arrell, J. Z. Lindor, P. P. Dzeja,
Y. Ikeda, C. Perez-Terzic, A. Terzic, Somatic oxidative bioenergetics transitions into
pluripotency-dependent glycolysis to facilitate nuclear reprogramming. *Cell Metab*
14, 264-271 (2011).

19. A. D. Panopoulos, O. Yanes, S. Ruiz, Y. S. Kida, D. Diep, R. Tautenhahn, A. Herrerias,
E. M. Batchelder, N. Plongthongkum, M. Lutz, W. T. Berggren, K. Zhang, R. M. Evans,
G. Siuzdak, J. C. Izpisua Belmonte, The metabolome of induced pluripotent stem cells
reveals metabolic changes occurring in somatic cell reprogramming. *Cell Res* **22**,
168-177 (2012).

20. A. Moussaieff, M. Rouleau, D. Kitsberg, M. Cohen, G. Levy, D. Barasch, A. Nemirovski,
S. Shen-Orr, I. Laevsky, M. Amit, D. Bomze, B. Elena-Herrmann, T. Scherf, M. Nissim-
Rafinia, S. Kempa, J. Itskovitz-Eldor, E. Meshorer, D. Aberdam, Y. Nahmias, Glycolysis-
mediated changes in acetyl-CoA and histone acetylation control the early
differentiation of embryonic stem cells. *Cell Metab* **21**, 392-402 (2015).

21. A. De Los Angeles, F. Ferrari, R. Xi, Y. Fujiwara, N. Benvenisty, H. Deng, K.
Hochedlinger, R. Jaenisch, S. Lee, H. G. Leitch, M. W. Lensch, E. Lujan, D. Pei, J.
Rossant, M. Wernig, P. J. Park, G. Q. Daley, Hallmarks of pluripotency. *Nature* **525**,
469-478 (2015).

22. P. Du, J. Wu, Hallmarks of totipotent and pluripotent stem cell states. *Cell Stem Cell*
31, 312-333 (2024).

23. E. A. Rowland, T. M. Greco, C. K. Snowden, A. L. McCabe, T. J. Silhavy, I. M. Cristea,
Sirtuin Lipoamidase Activity Is Conserved in Bacteria as a Regulator of Metabolic
Enzyme Complexes. *mBio* **8**, (2017).

24. E. A. Rowland, C. K. Snowden, I. M. Cristea, Protein lipoylation: an evolutionarily
conserved metabolic regulator of health and disease. *Curr Opin Chem Biol* **42**, 76-85
(2018).

25. J. E. Cronan, Assembly of Lipoic Acid on Its Cognate Enzymes: an Extraordinary and
Essential Biosynthetic Pathway. *Microbiol Mol Biol Rev* **80**, 429-450 (2016).

26. P. S. J. Bailey, B. M. Ortmann, A. W. Martinelli, J. W. Houghton, A. S. H. Costa, S. P. Burr,
R. Antrobus, C. Frezza, J. A. Nathan, ABHD11 maintains 2-oxoglutarate metabolism
by preserving functional lipoylation of the 2-oxoglutarate dehydrogenase complex.
Nat Commun **11**, 4046 (2020).

27. C. Buecker, R. Srinivasan, Z. Wu, E. Calo, D. Acampora, T. Faial, A. Simeone, M. Tan,
T. Swigut, J. Wysocka, Reorganization of enhancer patterns in transition from naive to
primed pluripotency. *Cell Stem Cell* **14**, 838-853 (2014).

28. C. Zeng, J. Chen, E. W. Cooke, A. Subuddhi, E. T. Roodman, F. X. Chen, K. Cao,
Demethylase-independent roles of LSD1 in regulating enhancers and cell fate
transition. *Nat Commun* **14**, 4944 (2023).

29. J. A. Lessard, G. R. Crabtree, Chromatin regulatory mechanisms in pluripotency. *Annu*
Rev Cell Dev Biol **26**, 503-532 (2010).

30. S. Feng, S. E. Jacobsen, W. Reik, Epigenetic reprogramming in plant and animal
development. *Science* **330**, 622-627 (2010).

31. B. Papp, K. Plath, Epigenetics of reprogramming to induced pluripotency. *Cell* **152**,
1324-1343 (2013).

32. W. W. Tee, D. Reinberg, Chromatin features and the epigenetic regulation of
pluripotency states in ESCs. *Development* **141**, 2376-2390 (2014).

33. S. Schlesinger, E. Meshorer, Open Chromatin, Epigenetic Plasticity, and Nuclear
Organization in Pluripotency. *Dev Cell* **48**, 135-150 (2019).

34. A. Burton, M. E. Torres-Padilla, Epigenome dynamics in early mammalian
embryogenesis. *Nat Rev Genet* **26**, 587-603 (2025).

35. X. Wan, S. Zhang, J. Li, D. Kong, M. Chen, Epigenetic remodeling during early
embryonic development. *Front Cell Dev Biol* **14**, 1750381 (2026).

36. R. Eskeland, M. Leeb, G. R. Grimes, C. Kress, S. Boyle, D. Sproul, N. Gilbert, Y. Fan, A.
I. Skoultchi, A. Wutz, W. A. Bickmore, Ring1B compacts chromatin structure and
represses gene expression independent of histone ubiquitination. *Mol Cell* **38**, 452-
464 (2010).

37. R. S. Illingworth, M. Moffat, A. R. Mann, D. Read, C. J. Hunter, M. M. Pradeepa, I. R.
Adams, W. A. Bickmore, The E3 ubiquitin ligase activity of RING1B is not essential for
early mouse development. *Genes Dev* **29**, 1897-1902 (2015).

38. C. C. Sze, K. Cao, C. K. Collings, S. A. Marshall, E. J. Rendleman, P. A. Ozark, F. X. Chen,
M. A. Morgan, L. Wang, A. Shilatifard, Histone H3K4 methylation-dependent and -
independent functions of Set1A/COMPASS in embryonic stem cell self-renewal and
differentiation. *Genes Dev* **31**, 1732-1737 (2017).

39. L. Wang, C. K. Collings, Z. Zhao, K. A. Cozzolino, Q. Ma, K. Liang, S. A. Marshall, C. C.
Sze, R. Hashizume, J. N. Savas, A. Shilatifard, A cytoplasmic COMPASS is necessary
for cell survival and triple-negative breast cancer pathogenesis by regulating
metabolism. *Genes Dev* **31**, 2056-2066 (2017).

40. K. M. Dorighi, T. Swigut, T. Henriques, N. V. Bhanu, B. S. Scruggs, N. Nady, C. D. Still,
2nd, B. A. Garcia, K. Adelman, J. Wysocka, Mll3 and Mll4 Facilitate Enhancer RNA
Synthesis and Transcription from Promoters Independently of H3K4
Monomethylation. *Mol Cell* **66**, 568-576 e564 (2017).

41. K. Cao, M. Ugarenko, P. A. Ozark, J. Wang, S. A. Marshall, E. J. Rendleman, K. Liang, L.
Wang, L. Zou, E. R. Smith, F. Yue, A. Shilatifard, DOT1L-controlled cell-fate
determination and transcription elongation are independent of H3K79 methylation.
Proc Natl Acad Sci U S A **117**, 27365-27373 (2020).

42. M. A. J. Morgan, A. Shilatifard, Epigenetic moonlighting: Catalytic-independent
functions of histone modifiers in regulating transcription. *Sci Adv* **9**, eadg6593 (2023).

43. A. Khan, C. Zhang, P. G. Nguyen, J. M. Metts, L. C. Collins, K. Jain, C. A. Mills, L. Vlach,
K. Li, A. L. Brademeyer, B. M. Bowman, M. B. Major, J. Aube, L. E. Herring, W. K.
Rathmell, F. M. Mason, I. J. Davis, Q. Zhang, B. D. Strahl, A SETD2-CDK1-lamin axis
maintains nuclear morphology and genome stability. *Nat Cell Biol* **27**, 1327-1341
(2025).

44. C. Wang, J. E. Lee, B. Lai, T. S. Macfarlan, S. Xu, L. Zhuang, C. Liu, W. Peng, K. Ge,
Enhancer priming by H3K4 methyltransferase MLL4 controls cell fate transition. *Proc*
Natl Acad Sci U S A **113**, 11871-11876 (2016).

45. C. C. Sze, A. Shilatifard, MLL3/MLL4/COMPASS Family on Epigenetic Regulation of
Enhancer Function and Cancer. *Cold Spring Harb Perspect Med* **6**, (2016).

46. G. Xie, J. E. Lee, A. D. Senft, Y. K. Park, Y. Jang, S. Chakraborty, J. J. Thompson, K.
McKernan, C. Liu, T. S. Macfarlan, P. P. Rocha, W. Peng, K. Ge, MLL3/MLL4
methyltransferase activities control early embryonic development and embryonic
stem cell differentiation in a lineage-selective manner. *Nat Genet* **55**, 693-705 (2023).

47. J. W. Hong, D. A. Hendrix, M. S. Levine, Shadow enhancers as a source of evolutionary
novelty. *Science* **321**, 1314 (2008).

48. K. Cao, C. K. Collings, S. A. Marshall, M. A. Morgan, E. J. Rendleman, L. Wang, C. C.
Sze, T. Sun, E. T. Bartom, A. Shilatifard, SET1A/COMPASS and shadow enhancers in
the regulation of homeotic gene expression. *Genes Dev* **31**, 787-801 (2017).

49. E. Z. Kvon, R. Waymack, M. Gad, Z. Wunderlich, Enhancer redundancy in
development and disease. *Nat Rev Genet* **22**, 324-336 (2021).

50. Y. Tsukada, J. Fang, H. Erdjument-Bromage, M. E. Warren, C. H. Borchers, P. Tempst,
Y. Zhang, Histone demethylation by a family of JmjC domain-containing proteins.
Nature **439**, 811-816 (2006).

51. M. Tahiliani, K. P. Koh, Y. Shen, W. A. Pastor, H. Bandukwala, Y. Brudno, S. Agarwal, L.
M. Iyer, D. R. Liu, L. Aravind, A. Rao, Conversion of 5-methylcytosine to 5-
hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* **324**, 930-
935 (2009).

52. W. Xu, H. Yang, Y. Liu, Y. Yang, P. Wang, S. H. Kim, S. Ito, C. Yang, P. Wang, M. T. Xiao,
L. X. Liu, W. Q. Jiang, J. Liu, J. Y. Zhang, B. Wang, S. Frye, Y. Zhang, Y. H. Xu, Q. Y. Lei, K.
L. Guan, S. M. Zhao, Y. Xiong, Oncometabolite 2-hydroxyglutarate is a competitive
inhibitor of alpha-ketoglutarate-dependent dioxygenases. *Cancer Cell* **19**, 17-30
(2011).

53. M. Xiao, H. Yang, W. Xu, S. Ma, H. Lin, H. Zhu, L. Liu, Y. Liu, C. Yang, Y. Xu, S. Zhao, D.
Ye, Y. Xiong, K. L. Guan, Inhibition of alpha-KG-dependent histone and DNA
demethylases by fumarate and succinate that are accumulated in mutations of FH
and SDH tumor suppressors. *Genes Dev* **26**, 1326-1338 (2012).

54. K. Skvortsova, E. Masle-Farquhar, P. L. Luu, J. Z. Song, W. Qu, E. Zotenko, C. M. Gould,
Q. Du, T. J. Peters, Y. Colino-Sanguino, R. Pidsley, S. S. Nair, A. Khoury, G. C. Smith, L.
A. Miosge, J. H. Reed, J. G. Kench, M. A. Rubin, L. Horvath, O. Bogdanovic, S. M. Lim,
J. M. Polo, C. C. Goodnow, C. Stirzaker, S. J. Clark, DNA Hypermethylation
Encroachment at CpG Island Borders in Cancer Is Predisposed by H3K4
Monomethylation Patterns. *Cancer Cell* **35**, 297-314 e298 (2019).

55. E. W. Cooke, C. Zeng, S. M. Nur, Y. Jia, A. Huang, J. Chen, P. Gao, F. X. Chen, F. Jin, K.
Cao, H3K79 methylation and H3K36 trimethylation synergistically regulate gene
expression in pluripotent stem cells. *Sci Adv* **11**, eadt8765 (2025).

56. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

57. A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M.
Chaisson, T. R. Gingeras, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,
15-21 (2013).
58. F. Ramirez, D. P. Ryan, B. Gruning, V. Bhardwaj, F. Kilpert, A. S. Richter, S. Heyne, F.
Dundar, T. Manke, deepTools2: a next generation web server for deep-sequencing
data analysis. *Nucleic Acids Res* **44**, W160-165 (2016).
59. S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).
60. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
139-140 (2010).
61. T. Wu, E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan, X. Fu,
S. Liu, X. Bo, G. Yu, clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. *Innovation (Camb)* **2**, 100141 (2021).
62. Y. Zhou, B. Zhou, L. Pache, M. Chang, A. H. Khodabakhshi, O. Tanaseichuk, C. Benner,
S. K. Chanda, Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nat Commun* **10**, 1523 (2019).

A Day 6 EBs
WT
DKO

B EB size

C DKO *vs.* WT day 6 EBs

D Upregulated genes during EB differentiation
1,633
1,192
1,088
Downregulated genes in DKO EBs

E
Actin cytoskeleton organization
Tube morphogenesis
Embryonic morphogenesis
Tissue morphogenesis
Positive regulation of locomotion

F Day 6 EBs
DKO+DOX
DOE+DOX

G EB size

H DOE *vs.* DKO day 6 EBs

I 1,192 MLL3/4-dependent differentiation genes
NES=2.5692
FDR=0

J

K MLL3/4-dependant differentiation genes
892
300
575
Upregulated genes in DOE EBs

L
Renal system development
Embryonic morphogenesis
Skeletal system development
Brain development
Mesenchyme development