MLL3/4 methyltransferases regulate the differentiation of pluripotent stem cells via cellular

respiration

Suza Mohammad Nur[1,3], Yunbo Jia[1], Muyi Ye[1], Caylin A. Lepak[1], Issam Ben-Sahra[4,5], and

Kaixiang Cao[1,2,3,*]

[1]Department of Biochemistry, Case Western Reserve University

[2]Department of Genetics and Genome Sciences, Case Western Reserve University

[3]Case Comprehensive Cancer Center, Case Western Reserve University, Cleveland, OH 44106,

USA

[4]Department of Biochemistry and Molecular Genetics, Northwestern University

[5]Robert H. Lurie Comprehensive Cancer Center, Northwestern University, Chicago, IL 60611,

USA

Running title: MLL3/4 regulate cellular respiration to facilitate pluripotent stem cell differentiation

Keywords: MLL3, MLL4, stem cells, differentiation, cellular metabolism, glycolysis, TCA cycle

*Correspondence should be sent to the following address:

Kaixiang Cao, PhD

Department of Biochemistry, School of Medicine, Case Western Reserve University

10900 Euclid Avenue, Cleveland, OH 44106, USA

Office: (216) 368-1218

Email: kxc725@case.edu

Abstract

Enhancer-regulating epigenetic modifiers play critical roles in normal physiological processes and human pathogenesis. The major enhancer regulator paralogs MLL3 and MLL4 (MLL3/4) belong to the lysine methyltransferase 2 (KMT2) family, which catalyzes the methylation of lysine 4 on histone H3 (H3K4me). MLL3/4 are required for enhancer activation and are essential for mammalian development and stem cell differentiation. Recent studies have linked MLL3/4 with different metabolic pathways in the context of stem cell self-renewal and cancer cell growth; however, the underlying mechanisms remain elusive. Here, we utilize Seahorse extracellular flux analysis, stable isotope tracing, stem cell biology techniques, and transcriptomic analysis to investigate the functional relationship of MLL3/4, cellular respiration, and stem cell differentiation. Our results indicate that the loss of MLL3/4 impairs glycolytic activity and mitochondrial respiration in murine embryonic stem cells by downregulating the rate-limiting glycolytic enzyme Hexokinase 2 (HK2) and impairing the function of the Alpha-ketoglutarate dehydrogenase (OGDH) complex. Furthermore, simultaneously overexpression of HK2 and OGDH rescues defects in both cellular respiration and differentiation caused by MLL3/4 loss. Taken together, our study reveals a novel mechanism by which epigenetic machineries such as MLL3/4 govern the differentiation of pluripotent stem cells and facilitates the understanding of disease pathogenesis driven by enhancer malfunction.

Introduction

Enhancer deregulation is a major driver of human congenital disorders (*1-4*). In mammals, nucleosomes at enhancers are mono-methylated on lysine 4 of histone H3 (H3K4) by the paralogous enzymes Mixed-lineage Leukemia 3 (MLL3, also known as KMT2C) and Mixed-lineage Leukemia 4 (MLL4, also known as KMT2D) (*5-9*). Mice null for MLL4 die shortly after gastrulation, while MLL3 loss leads to neonatal lethality (*10, 11*). Heterozygous loss-of-function mutations in MLL4 lead to Kabuki syndrome (*1*) and those in MLL3 cause Kleefstra syndrome 2 (*4, 12, 13*). Both disorders are characterized by craniofacial dysmorphism and developmental delay, highlighting the instructive role of MLL3/4 in normal development and human pathogenesis. However, the underlying mechanisms remain poorly understood.

Recent results have revealed a functional link between MLL3/4 and cellular metabolism. MLL3/4 and nucleotide synthesis pathways co-regulate the self-renewal of pluripotent stem cells (PSCs) (*14*). Moreover, steady-state levels of glycolytic metabolites such as glucose-6-phosphate (G6P) and fructose-6-phosphate (F6P), and tricarboxylic acid (TCA) cycle metabolites including succinate, fumarate, and malate, are reduced in PSCs lacking MLL3/4 (*14*). In contrast, MLL4-deficient lung cancer and melanoma cells adopt upregulated glycolysis (*15, 16*), suggesting a context-dependent role of MLL3/4 in regulating cellular metabolism. How MLL3/4 modulate metabolic pathways and whether they govern cell fate via these pathways are largely unknown.

Cellular respiration is tightly regulated in PSCs. Naive PSCs, such as murine embryonic stem cells (ESCs), exhibit metabolic bivalency, utilizing both glycolysis and mitochondrial respiration for energy production, whereas primed PSCs are predominantly glycolytic (*17*). On the other hand, somatic cells are more dependent on mitochondrial respiration (*18, 19*). Inhibition of glycolysis drives the exit from pluripotency, while stimulation of glycolysis enhances somatic cell

reprogramming to induced PSCs (iPSCs) (*18, 20*), suggesting that the balance between glycolysis

and mitochondrial respiration instructs cell fate transition. To date, how metabolic states are

established during development and whether they instruct cell identity remain enigmatic.

To investigate the functional relationship of MLL3/4, cellular metabolism, and PSC

differentiation, we examined the impact of MLL3/4 loss on cellular respiration in ESCs. Both

extracellular acidification rate (ECAR) and oxygen consumption rate (OCR) are reduced in

MLL3/4-null ESCs. Stable isotope tracing reveals a reduction of metabolite levels in both

glycolysis and the TCA cycle upon MLL3/4 loss. MLL3/4 are required for the expression of

Hexokinase 2 (HK2) and for proper lipoylation of Dihydrolipoamide S-Succinyltransferase

(DLST), the E2 subunit of the Alpha-ketoglutarate Dehydrogenase complex (OGDHc). Moreover,

simultaneous overexpression of HK2 and OGDH partially rescues defects in cellular respiration

and differentiation caused by MLL3/4 loss. Our results provide novel insight into understanding

the crosstalk between epigenetic machineries and cellular metabolism. Importantly, they may pave

the way for developing novel strategies to target human diseases driven by enhancer deregulation.

Results

MLL3/4 loss dampens glycolytic activity in ground-state ESCs

We previously showed that steady-state levels of glycolysis-related metabolites, such as G6P and F6P, were reduced by the loss of MLL3/4 in ESCs (*14*), suggesting that MLL3/4 may regulate glycolysis in naive PSCs. To determine the impact of MLL3/4 loss on the rate of glycolysis, we measured the ECAR of wildtype (WT) and MLL3/4 double knockout (DKO) ESCs (*14*) with the glycolysis stress test using the Seahorse XF Analyzer. The levels of glycolysis and glycolytic capacity were significantly reduced by MLL3/4 loss (Fig. 1A-C), suggesting that MLL3/4 are required to maintain glycolytic activity in naive PSCs. To understand if the decrease in the steady-state levels of metabolites reflects altered glycolytic activity, we performed stable isotope tracing with $[^{13}C_6]$-glucose in WT and MLL3/4 DKO ESCs, followed by liquid chromatography-tandem mass spectrometry (LC-MS/MS) analysis (Fig. 1D). Consistent with results in steady-state metabolomics, the incorporation of ^{13}C into F6P/G1P (glucose-1-phosphate)/G6P (m+6) was significantly reduced by MLL3/4 loss (Fig. 1E and Fig. S1A). Moreover, levels of lactate (m+3) were reduced by MLL3/4 loss (Fig. 1F and Fig. S1B), corresponding to the decreased glycolytic activity in MLL3/4 DKO ESCs. To determine if the decreased levels of glycolytic metabolites resulted from defective glucose import, we measured the rate of glucose uptake and found that MLL3/4 loss did not have a significant impact on cellular import of glucose in ESCs (Fig. 1G), suggesting that MLL3/4 loss impairs early rate-limiting steps in glycolysis, leading to reduced glycolytic activity in ESCs.

To elucidate the mechanism underlying reduced rates of glycolysis in MLL3/4 DKO ESCs, we analyzed RNA-seq data (*14*) to examine the expression levels of genes encoding glycolytic enzymes involved in G6P and F6P metabolism. We found that *Hk2*, the gene encoding the rate-

limiting enzyme Hexokinase 2 that catalyzes the phosphorylation of glucose to G6P, was

significantly downregulated by MLL3/4 loss (Fig. S1C). On the other hand, the expression levels

of genes encoding Glucose-6-phosphate isomerase (*Gpi1*), Phosphofructokinase 1 (*Pfkl* and *Pfkp*),

and Phosphofructokinase 2 (*Pfkfb1* and *Pfkfb2*) were comparable between WT and MLL3/4 DKO

ESCs (Fig. S1C). We then measured the levels of Hexokinases in WT and MLL3/4 DKO ESCs by

western blotting. HK2 levels were reduced by the loss of MLL3/4, while the levels of HK1

(Hexokinase 1), the paralog of HK2, remained largely unchanged (Fig. 1H). To investigate how

MLL3/4 regulate the expression levels of *Hk2*, we analyzed ChIP-seq datasets (*7, 14*) and found

at least six putative enhancers near the *Hk2* locus that are bound by MLL4 and exhibit reduced

H3K27ac levels upon MLL3/4 loss (Fig. S1D), suggesting that MLL3/4 regulate *Hk2* through

these putative enhancers. Taken together, these data indicate that the glycolytic activity of naive

PSCs requires MLL3/4 and that the decreased HK2 levels may be responsible for the defective

glycolysis caused by MLL3/4 loss.

MLL3/4 are required for mitochondrial respiration in ground-state ESCs

The metabolic bivalency is one of the hallmarks of naive PSCs (*21, 22*). To examine the role of

MLL3/4 in regulating mitochondrial respiration in naive PSCs, we measured the OCR of WT and

MLL3/4 DKO ESCs with the mitochondrial stress assay using the Seahorse XF Analyzer.

Interestingly, the levels of basal and maximal respiration were significantly reduced by MLL3/4

loss in ESCs (Fig. 2A-C), suggesting that MLL3/4 are required to maintain the metabolic bivalency

in naive PSCs. Analysis of ^{13}C incorporation into TCA-cycle metabolites via stable isotope tracing

using [^{13}C$_6$]-glucose labeling (Fig. 2D) showed that MLL3/4 loss did not have a significant impact

on the levels of citrate/isocitrate (m+2) and alpha-ketoglutarate (α-KG) (m+2) (Fig. 2E-F and Fig.

S2A-B). In contrast, the levels of succinate (m+2), fumarate (m+2), and malate (m+2) were significantly reduced in MLL3/4 DKO ESCs (Fig. 2G-I and Fig. S2C-E), suggesting that potential disruptions in TCA cycle steps downstream of α-KG contribute to the impaired mitochondrial respiration driven by MLL3/4 loss.

To further examine this possibility, we performed stable isotope tracing with $[^{13}C_5]$-glutamine labeling (Fig. 2J). Although MLL3/4 loss did not have a significant impact on the levels of glutamate (m+5) (Fig. 2K and Fig. S2F), the levels of α-KG (m+5) were significantly reduced in MLL3/4 DKO ESCs (Fig. 2L and Fig. S2G). This reduction may reflect impaired conversion of glutamate to α-KG, although the expression levels of *Glud1*, which encodes Glutamate Dehydrogenase (GDH), the enzyme catalyzing the conversion of glutamate to α-KG, were comparable between WT and MLL3/4 DKO ESCs (Fig. S2L). Consistent with results from $[^{13}C_6]$-glucose labeling, the levels of succinate (m+4), fumarate (m+4), and malate (m+4) were significantly reduced in MLL3/4 DKO ESCs, while those of citrate/isocitrate (m+4) were largely unchanged after $[^{13}C_5]$-glutamine labeling (Fig. 2M-P and Fig. S2H-K). Furthermore, expression levels of multiple genes encoding enzymes involved in α-KG and downstream TCA cycle metabolism were comparable between MLL3/4 DKO and WT ESCs (Fig. S2L). In addition, protein levels of OGDH, SUCLA2, SUCLG1, SUCLG2, and SDHA remained largely unperturbed upon MLL3/4 loss (Fig. S2M-O), suggesting that MLL3/4 regulate TCA cycle metabolites via mechanisms other than transcriptional control of metabolic enzymes. Taken together, these results demonstrate that MLL3/4 loss impairs mitochondrial respiration and disrupts metabolic flux at specific steps within the TCA cycle in naive PSCs.

Simultaneous overexpression of HK2 and OGDH rescues defects in cellular respiration caused by MLL3/4 loss

To determine if MLL3/4 regulate cellular respiration in ESCs through activating HK2, we overexpressed 3×HA-tagged HK2 in MLL3/4 DKO ESCs with a doxycycline (DOX)-inducible system (*23, 24*) (Fig. S3A). HK2 overexpression increased glycolytic activity in MLL3/4 DKO cells (Fig. S3B-D); however, OCR during basal respiration was reduced upon HK2 expression (Fig. S3E-G), suggesting that HK2 overexpression renders MLL3/4 DKO ESCs more dependent on glycolysis for energy production.

Since HK2 overexpression led to an expected increase in ECAR but a reduction in OCR, we sought to restore both glycolysis and mitochondrial respiration in MLL3/4 DKO ESCs to study the functional relationship of MLL3/4, cellular respiration, and the differentiation capability of PSCs. We hypothesized that the decrease of ^{13}C incorporation in succinate, fumarate, and malate in both $[^{13}C_6]$-glucose and $[^{13}C_5]$-glutamine labelled MLL3/4-null ESCs results from the potentially defective OGDHc caused by the loss of MLL3/4. Although OGDH levels were comparable between MLL3/4 DKO and WT ESCs, OGDH overexpression could potentially compensate for reduced levels of succinate, fumarate, and malate in MLL3/4-null ESCs. We therefore simultaneously overexpressed 3×HA-tagged HK2 and 3×FLAG-tagged OGDH in MLL3/4 DKO ESCs using DOX-inducible expression vectors carrying distinct drug resistance genes (Fig. 3A). Increased levels of HK2 and OGDH led to elevated rates of glycolysis, glycolytic capacity, and both basal and maximal respiration (Fig. 3B-G). Furthermore, stable isotope tracing by labeling MLL3/4 DKO and HK2/OGDH double-overexpressing (DOE) MLL3/4 DKO cells with $[^{13}C_6]$-glucose indicated that the levels of metabolites reduced by MLL3/4 loss such as F6P/G1P/G6P (m+6), lactate (m+3), succinate (m+2), and fumarate (m+2) were partially restored

by HK2/OGDH DOE (Fig. 3H-M, Fig. S4A-B, and Fig. S4E-G). As expected, we did not observe

a significant difference in the levels of citrate/isocitrate (m+2) or α-KG (m+2) (Fig. 3J and Fig.

S4C-E). Notably, malate (m+2) levels were not significantly increased upon HK2/OGDH DOE in

MLL3/4 DKO cells (Fig. 3M and Fig. S4H), suggesting that there are additional regulatory layers

on the levels of malate besides HK2 and OGDHc.

Our results that HK2/OGDH overexpression rescues defects in cellular respiration and

levels of glycolytic and TCA cycle metabolites suggest that MLL3/4 may regulate glycolysis and

mitochondrial respiration through activating *Hk2* expression and maintaining the normal function

of the OGDHc. To further elucidate the molecular mechanisms by which MLL3/4 regulate the

TCA cycle, we examined the levels of DLST and Dihydrolipoamide Dehydrogenase (DLD), E2

and E3 components of the OGDHc. MLL3/4 loss had little impact on their levels (Fig. 3N-O),

suggesting that MLL3/4 are unlikely to regulate mitochondrial respiration via modulating the

abundance of OGDHc components. Lipoylated DLST is essential for the conversion of α-KG to

succinyl-CoA (*25-27*). To understand if MLL3/4 loss impairs the activity of DLST, we examined

the levels of lipoylated proteins in WT and MLL3/4 DKO ESCs using a specific anti-lipoate

antibody (*28*). MLL3/4 loss led to a reproducible decrease of the lipoylated protein species with

lower molecular weight, attributed to lipoylated DLST based on its size (Fig. 3P). The decrease in

lipoylation was specific to DLST as levels of other abundantly lipoylated proteins in the

mitochondria, such as Dihydrolipoamide Acetyltransferase (DLAT) in the Pyruvate

Dehydrogenase complex and Dihydrolipoamide Branched-chain Transacylase (DBT) in the

Branched-chain Alpha-keto Acid Dehydrogenase complex (*26, 27*), were comparable in MLL3/4

DKO and WT ESCs (Fig. 3P). Taken together, these results suggest that MLL3/4 regulate cellular

respiration by maintaining the levels of HK2 and the activity of OGDHc in naive PSCs, and that

simultaneous overexpression of HK2 and OGDH partially rescues defects in cellular respiration

caused by MLL3/4 loss.

Simultaneous overexpression of HK2 and OGDH rescues differentiation defects caused by

MLL3/4 loss

To assess the impact of MLL3/4 loss on cell fate transition, we generated embryoid bodies (EBs)

from WT and MLL3/4 DKO ESCs. As expected, MLL3/4 loss led to a decrease in the size of day

6 EBs (Fig. 4A-B), indicating that MLL3/4 are required for PSC differentiation. RNA-seq analysis

showed that MLL3/4 loss caused the significant deregulation of 4,547 genes in day 6 EBs (Fig.

4C). Among genes significantly downregulated in MLL3/4 DKO EBs, 52.3% (1,192 of 2,280)

were significantly upregulated during WT EB differentiation (Fig. 4D), suggesting that these genes

are dependent on MLL3/4 to be activated during differentiation. We thus defined them as MLL3/4-

dependent differentiation genes. Gene ontology (GO) analysis showed that the top 300 upregulated

genes of the 1,192 MLL3/4-dependent differentiation genes were enriched for embryogenesis-

related genes and known MLL4 targets essential for actin cytoskeleton organization (*7*) (Fig. 4E),

further indicating that MLL3/4 loss impairs PSC differentiation. Indeed, pluripotency genes were

significantly upregulated in MLL3/4 DKO EBs and marker genes of meso- and endo-derm

lineages were downregulated in EBs by MLL3/4 loss (Fig. S5A-B). These data demonstrate that

MLL3/4 are required for PSC differentiation and related transcriptional reconfiguration.

To examine if the partial restoration of cellular respiration had any impact on the

differentiation capacity of ESCs lacking MLL3/4, we treated MLL3/4 DKO and HK2/OGDH DOE

MLL3/4 DKO cells with DOX and differentiated them toward EBs. EB differentiation revealed

that HK2/OGDH DOE led to a significant increase in the size of day 6 EBs (Fig. 4F-G), suggesting

that restoring cellular respiration rescues differentiation defects caused by MLL3/4 loss. Moreover,

RNA-seq analysis identified 1,778 significantly deregulated genes by HK2/OGDH DOE

compared with MLL3/4 DKO ESCs treated with DOX (Fig. 4H). Gene set enrichment analysis

(GSEA) and hierarchical clustering analysis of MLL3/4-dependent differentiation genes (Fig. 4D)

revealed a significant upregulation of them upon HK2/OGDH DOE (Fig. 4I-J). Indeed, 34.3% of

genes upregulated by HK2/OGDH DOE (300 of 875) were dependent on MLL3/4 to be activated

during differentiation (Fig. 4K), suggesting that the impaired cellular respiration contributes to

differentiation defects caused by MLL3/4 loss. GO analysis of the 300 significantly rescued genes

showed enrichment of genes related to embryo morphogenesis and mesodermal development such

as the development of renal and skeletal systems (Fig. 4L). Consistently, HK2/OGDH DOE led

to the significant downregulation of pluripotency genes and the significant upregulation of

differentiation marker genes (Fig. S5C-D). Taken together, these results suggest that defects in

cellular respiration contribute to differentiation failures of naive PSCs lacking MLL3/4.

Discussion

Epigenetic regulators, including histone modifiers, play central roles in regulating cell fate

transition during PSC differentiation, somatic cell reprogramming, and normal development (*29-*

35); however, the underlying mechanisms remain incompletely understood, in part because many

epigenetic regulators function independently of their catalytic activity (*6, 7, 24, 36-43*). Besides

being essential for cellular differentiation and mammalian embryogenesis (*11, 44*), loss-of-

function mutations in MLL3 and MLL4 drive congenital disorders such as Kleefstra syndrome 2

and Kabuki syndrome, as well as multiple types of cancer (*1, 4, 13, 45*). Recent advances reveal a

novel link between MLL3/4 and cellular metabolism; however, whether and how the roles of

MLL3/4 in regulating cell fate and metabolic pathways intersect are unclear. Here, we unveil a

previously unrecognized function of MLL3/4 in regulating glycolysis and mitochondrial

respiration in murine ESCs. Our findings suggest that MLL3/4 govern cell fate transition during

the differentiation of naive PSCs through regulating the expression levels of *Hk2* and the function

of OGDHc, two rate-limiting nodes in glycolysis and the TCA cycle, respectively. Importantly,

restoring defective cellular respiration may be an effective strategy to treat diseases driven by

MLL3/MLL4 loss-of-function.

We find that MLL3/4 loss causes decreased glycolytic activity and mitochondrial

respiration in ESCs. Metabolic bivalency is a defining feature of naive PSCs such as murine ESCs

(*22*). The decrease of glycolytic activity in MLL3/4 null cells may hinder the cells from exiting

naive pluripotency, as formative PSCs and primed PSCs are highly glycolytic (*17*). On the other

hand, inhibiting glycolysis promotes differentiation as somatic cells have much elevated levels of

oxidative phosphorylation compared with PSCs (*18-20*). Therefore, the dampened mitochondrial

respiration caused by MLL3/4 loss may impair the differentiation of PSCs into germ layers and/or

specific lineages. Our data provide a mechanistic explanation for the failure of MLL4-null ESCs

to exit naive pluripotency and for the severe gastrulation defects observed in MLL3/4-null embryos

(*7, 10, 46*).

Our data support that MLL3/4 facilitate glycolysis in naive ESCs through maintaining the

expression of *Hk2*, which encodes the rate-limiting enzyme Hexokinase 2 in glycolysis. It is likely

that MLL3/4 regulate *Hk2* expression through enhancers as we identified multiple putative *cis-*

regulatory regions that may be directly regulated by MLL3/4. Understanding whether and how

MLL3/4 function through these enhancers to control the expression levels of *Hk2* is an interesting

topic. Shadow enhancers have been reported in different organisms including *Drosophila*, mouse,

and human (*47-49*). Generating individual and compound enhancer mutations will elucidate how

the putative enhancers regulate *Hk2*. Combination of MLL3/4 loss-of-function mutations and

enhancer mutants will further reveal the functional relationship between MLL3/4 and putative

enhancers in regulating *Hk2*. These future studies are essential for understanding the mechanisms

by which MLL3/4 regulate glycolysis in naive PSCs.

Interestingly, overexpressing HK2 alone only partially restores glycolytic activity in ESCs

lacking MLL3/4, suggesting that MLL3/4 support glycolysis in naive PSCs through additional

mechanisms. Determining whether other critical regulators of glycolysis such as

Phosphofructokinase, Pyruvate Kinase, and Lactate Dehydrogenase play a role in dampening

glycolysis in the context of MLL3/4 loss-of-function is an important future study. Genome-wide

and targeted genetic screens may also identify additional factors that rescue glycolysis in MLL3/4-

null ESCs. It is noteworthy that HK2 overexpression leads to a decrease of OCR in MLL3/4 DKO

ESCs, suggesting that these cells are more dependent on glycolysis as an energy source in

comparison to oxidative phosphorylation under high levels of HK2. Since steady-state acetyl-CoA

levels were reduced in MLL3/4 DKO ESCs (*14*) and our stable isotope tracing assays were not

able to detect acetyl-CoA, it is interesting to examine if and how MLL3/4 loss affects the

mitochondrial transport of pyruvate in ESCs in the future. Defining the role of MLL3/4 in other

pathways leading to the synthesis of acetyl-CoA such as fatty acid metabolism may also facilitate

understanding how MLL3/4 regulate the conjunction of glycolysis and the TCA cycle.

The reduction in the ^{13}C incorporation into succinate, fumarate, and malate in both $[^{13}C_6]$-

glucose and $[^{13}C_5]$-glutamine labeling of MLL3/4 DKO ESCs suggests a functionally defective

OGDHc caused by MLL3/4 loss. Indeed, the levels of lipoylated DLST were decreased in

MLL3/4-null cells, suggesting a reduced activity of the OGDHc. It is likely that the decrease of

lipoylation is specific to DLST as MLL3/4 loss has little impact on the levels of lipoylated DLAT.

It was previously shown that the α/β hydrolase ABHD11 binds OGDHc and maintains the activity

of lipoylated DLST by preventing the formation of lipoyl adducts (*28*). It will be interesting to

investigate the functional relationship between MLL3/4, ABHD11, and OGDHc in the future to

understand how MLL3/4 regulate the activity of OGDHc. It is also noteworthy that the levels of

^{13}C incorporation into α-KG (m+5) is impaired in MLL3/4 DKO ESCs in the $[^{13}C_5]$-glutamine

labeling experiments, although α-KG (m+2) levels through $[^{13}C_5]$-glucose labeling and steady-

state levels of α-KG were largely unchanged by MLL3/4 loss (*14*). Besides Glutamate

Dehydrogenase, Glutamate Pyruvate Transaminase and Glutamate Oxaloacetate Transaminase

also can convert glutamate to α-KG. Future studies on the role of MLL3/4 in regulating the

conversion of glutamate to α-KG would be important to understand the crosstalk between MLL3/4

and glutaminolysis.

α-KG is an essential co-substrate of Fe(II)/α-KG-dependent dioxygenases including

Jumonji C (JmjC) domain-containing histone demethylases and the ten-eleven translocation (TET)

family of DNA hydroxylases (*50, 51*), playing a role in regulating the levels of methylation on

different lysine residues of histones and 5-methyl Cytosine (5mC) on DNA in mammalian cells.

Due to their structural similarity to α-KG, 2-hydroxyglutarate, succinate, and fumarate inhibit α-

KG-dependent dioxygenases by acting as a competitive inhibitor of α-KG (*52, 53*). Our results

that MLL3/4 loss in ESCs lead to a decreased levels of succinate and fumarate, through steady-

state metabolomics and stable isotope tracing with $[^{13}C_6]$-glucose and $[^{13}C_5]$-glutamine, suggest a

possibility that the activities of multiple histone demethylases and TETs are increased in cells with

MLL3/4 loss-of-function. Such a possibility provides a mechanistic explanation for the DNA

hypomethylation observed in MLL4-depleted B cells in mouse and MLL4 mutant prostate cancers

in human (*54*). Understanding the impact of the potential alterations in the levels of histone

methylation and DNA methylation on the differentiation capability of MLL3/4-mutant PSCs is

very interesting and may provide insight into understanding the molecular mechanisms underlying

the functional relationship of cellular respiration, epigenetic machineries, and cell fate transition.

In summary, our study reveals a previously unappreciated crosstalk between epigenetic

modifiers exemplified by MLL3/4 and cellular respiration pathways glycolysis and the TCA cycle

in regulating cell fate. Our findings on MLL3/4 regulating the levels of HK2 and lipoylated DLST

could pave the way for designing therapeutic strategies to target human diseases driven by loss-

of-function mutations of MLL3/4.

Methods

Antibodies

The following primary antibodies were used in this study: anti-HK2 (Cell Signaling Technology

2867), anti-HK1 (Cell Signaling Technology 2024), anti-OGDH (Proteintech 17110-1-AP), anti-

SDHA (Abcam 14715), anti-SUCLA2 (Proteintech 12627-1-AP), anti-SUCLG1 (Proteintech

14923-1-AP), anti-SUCLG2 (Proteintech NBP1-32521), anti-FLAG (MiliporeSigma F3165),

anti-HA (MiliporeSigma H3663), anti-DLST (Cell Signaling Technology 5556), anti-DLD (Santa

Cruz 365977), anti-Lipoic acid (Calbiochem 437695), anti-Tubulin (Developmental Studies

Hybridoma Bank E7), and anti-HSP90 (Santa Cruz 13119).

ESC Culture and EB differentiation

ESCs were cultured in serum-free medium supplemented with 2i/LIF as previously described (*55*).

For the generation of HK2 overexpressing and HK2/OGDH DOE MLL3/4 DKO ESCs, cells were

transfected with DOX-inducible PiggyBac vectors (*23*) as previously described (*24*) using an

Amaxa Nucleofector (Lonza). EB differentiation was performed using the hanging-drop method

as previously described (*24*). For DOX-induced overexpression, ESCs were treated with 100 ng/ml

doxycycline for 48 hours prior to downstream assays. For differentiation experiments, doxycycline

(100 ng/ml) were added 24 hours prior to EB formation and maintained throughout the 6-day

differentiation period.

Seahorse extracellular flux assays

ESCs were seeded in Seahorse XF microplates (Agilent) and cultured overnight with 2i/LIF serum-

free medium. Cells were then incubated in Seahorse XF assay medium (Agilent) at 37°C in a non-

CO_2 incubator for 1 hour. For the Glycolysis Stress Test (Agilent 103020-100), glucose (10mM),

oligomycin (2 µM), and 2-DG (50mM) were sequentially injected. For the Mitochondrial Stress

Test (Agilent 103015-100), oligomycin (1.5 µM), FCCP (0.5 µM), and Rot/AA (0.5 µM) were

sequentially injected. Cell Titer Glo assay (Promega) was performed in parallel on replica plates

to determine the cell number of each well. Individual raw ECAR and OCR values recorded by the

Seahorse XF Analyzer were normalized against the average Cell Titer Glo values of control wells

to derive normalized ECAR and normalized OCR values. Wave software (Agilent) was used to

extract individual ECAR and OCR measurements, which were subsequently used to generate the

bar plots.

Stable isotope tracing and metabolomics

Stable isotope labeling and metabolite extraction were performed as previously described (*14*).

ESCs were incubated with media containing 25 mM $[^{13}C_6]$-glucose or 4mM $[^{13}C_5]$-glutamine for

2 hours before methanol extraction. Metabolomic analyses were performed at the Northwestern

Metabolomics Core Facility.

Glucose uptake assay

Glucose update was measured using the Glucose Uptake-Glo Assay Kit (Promega J1342)

according to the manufacturer's instructions. In brief, 2×10^4 cells were seeded on each well of a

96-well flat bottom plate with opaque walls and clear bottoms (Corning 7200587). After 16 hours

of culture, cells were incubated in glucose-free medium for 1 hour at 37°C in the tissue-culture

incubator. Cells were then treated with 1mM 2-DG for 10 minutes at room temperature, followed

by the treatment of stop buffer, neutralization buffer, and detection reagent following the

instructions of the manufacturer. After 2.5 hours of dark-conditioned incubation, the luminescence

was measured using a GloMax luminometer (Promega). The luminescence data were normalized

by cell numbers measured with the Cell Titer Glo assay (Promega) on replica plates in parallel.

RNA-seq and analysis

Total RNA was extracted using TRIzol reagent (Thermo Fisher) following the manufacturer's

instructions. Each cell line was analyzed in at least two biological replicates. Extracted RNA were

treated with DNase I (Sigma) and purified with the RNeasy Plus Mini Kit (Qiagen). Ribosomal

RNA was removed from the purified RNA with the NEBNext rRNA Depletion Kit (New England

BioLabs). RNA libraries were then prepared with the NEBNext Ultra II Directional RNA Library

Prep Kit (New England BioLabs). The pooled libraries were sequenced with 150 bp paired-end

reads on the NovaSeq platfrom (Illumina).

Raw reads were processed with Trimmomatic v0.39(*56*) to remove adaptors and low-

quality reads with the parameter "-q 30". The cleaned reads were aligned to the mm9 genome

assembly using STAR v2.5.1 with default parameter (*57*). Reads were normalized to total read

counts per million (cpm) and visualized as bigwig-formatted coverage tracks using deepTools

v3.5.6 (*58*). Gene expression was quantified using HTseq V2.0.2 (*59*), and differential gene

expression analysis was performed with EdgeR v4.0.16 (*60*). Significantly differentially expressed

genes were identified with Benjamini-Hochberg-adjusted p value cutoff < 0.01 and a log fold

change threshold greater than |1|. hierarchical clustering heatmaps were generated using the ggpubr

v0.6.0 (https://rpkgs.datanovia.com/ggpubr/). GSEA plots were generated with clusterProfiler

v4.10.1 (*61*). Gene ontology (GO) analysis was performed using Metascape v3.5.20260201 (*62*).

Acknowledgements

We thank the Cao lab members for their helpful suggestions and discussions. We thank Dr. Peng Gao and the Northwestern Metabolomics Core Facility for performing the metabolomics study. We thank Emmalee W. Cooke for technical support. We thank Yalu Zhou for bioinformatic analysis support. This study was supported in part by grants R35GM150668, R00HD094906, and P30CA043703 from the National Institutes of Health to K.C.

396 **Author contributions**

397 K.C. conceived the study. S.M.N., I.B., and K.C. designed the research. S.M.N., Y.J., and M.Y.

398 performed experiments. S.M.N., Y.J., M.Y., and K.C. analyzed the data. S.M.N. and K.C. wrote

399 the manuscript. C.L. and I.B. critically read the manuscript.

    Figure legends

Figure 1. The loss of MLL3/4 leads to impaired glycolytic activity in ground-state ESCs.

(A) Representative ECAR traces of WT and MLL3/4 DKO ESCs measured by the Seahorse XF

glycolysis stress test assay. Data were normalized by the number of cells at the time of the assay

and are represented as mean ± S.D. Each data point had six replicates, and the experiment was

repeated three times independently. 2-DG: 2-Deoxy-D-glucose.

(B-C) Bar plots of ECAR showing the levels of glycolysis (B) and glycolytic capacity (C) in WT

and MLL3/4 DKO ESCs. Data are presented as mean ± S.D. p-values were calculated using

Student's t-test.

(D) Schematic of glycolysis illustrating the fate of ^{13}C carbons upon the incubation with $[^{13}C_6]$-

glucose.

(E-F) Bar plots showing the levels of F6P/G1P/G6P (m+6, E) and lactate (m+3, F) in WT and

MLL3/4 DKO ESCs labeled with $[^{13}C_6]$-glucose. n=3 biological replicates per sample. Data are

presented as mean ± S.D. p-values were calculated using Student's t-test.

(G) Bar plots showing the levels of glucose uptake in WT and MLL3/4 DKO ESCs. n=6 replicates

per sample. Experiments were repeated three times independently and representative results were

shown. Data were normalized by the number of cells at the time of the assay and are presented as

mean ± S.D. p-values were calculated using Student's t-test. RLU: relative light unit.

(H) Western blots showing the levels of HK1 and HK2 in WT and MLL3/4 DKO ESCs. 10 and

20µg of cell lysates were loaded for each sample. Experiments were repeated three times

independently and representative results were shown. Protein levels in each lane were quantified

by normalizing with Tubulin signals of the corresponding lane.

Figure 2. The loss of MLL3/4 leads to impaired mitochondrial respiration in ground-state ESCs.

(A) Representative OCR traces of WT and MLL3/4 DKO ESCs measured by the Seahorse XF mitochondrial stress test assay. Data were normalized by the number of cells at the time of the assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment was repeated three times independently. FCCP: Carbonyl Cyanide-4-(trifluoromethoxy) phenylhydrazone. ROT/AA: Rotenone and Antimycin A.

(B-C) Bar plots showing the levels of OCR during basal (B) and maximal (C) respiration in WT and MLL3/4 DKO ESCs. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(D) Schematic of the TCA cycle illustrating the fate of ^{13}C carbons upon the incubation with $[^{13}C_6]$-glucose.

(E-I) Bar plots showing levels of citrate/isocitrate (m+2, E), α-KG (m+2, F), succinate (m+2, G), fumarate (m+2, H), and malate (m+2, I) in WT and MLL3/4 DKO ESCs labeled with $[^{13}C_6]$-glucose. n=3 biological replicates per sample. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(J) Schematic of the TCA cycle illustrating the fate of ^{13}C carbons upon the incubation with $[^{13}C_5]$-glutamine.

(K-P) Bar plots showing levels of glutamate (m+5, K), α-KG (m+5, L), succinate (m+4, M), fumarate (m+4, N), malate (m+4, O), and citrate/isocitrate (m+4, P) in WT and MLL3/4 DKO ESCs labeled with $[^{13}C_5]$-glutamine. n=3 biological replicates per sample. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

Figure 3. Simultaneous overexpression of HK2 and OGDH rescues metabolic defects of MLL3/4 DKO ESCs.

(A) Western blots showing levels of HK2, 3×HA-tagged HK2, OGDH, and 3×FLAG-tagged OGDH in MLL3/4 DKO ESCs and MLL3/4 DKO ESCs overexpressing HK2/OGDH treated with or without DOX. 10 and 20μg of cell lysates were loaded for each sample. Experiments were repeated three times independently and representative results were shown. Protein levels in each lane of the anti-HK2 and anti-OGDH blots were quantified by normalizing with Tubulin signals of the corresponding lane.

(B) Representative ECAR traces of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX) measured by the Seahorse XF glycolysis stress test assay. Data were normalized by the number of cells at the time of the assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment was repeated three times independently.

(C-D) Bar plots of ECAR showing the level of glycolysis (C) and glycolytic capacity (D) in MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX). Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(E) Representative OCR traces of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX) measured by the Seahorse XF mitochondrial stress test assay. Data were normalized by the number of cells at the time of the assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment was repeated three times independently.

(F-G) Bar plots showing the levels of OCR during basal (F) and maximal (G) respiration in MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX). Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(H-M) Bar plots showing levels of F6P/G1P/G6P (m+6, H), lactate (m+3, I), α-KG (m+2, J), succinate (m+2, K), fumarate (m+2, L), and malate (m+2, M) in MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX) labeled with $[^{13}C_6]$-glucose. n=3 biological replicates per sample. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

(N-P) Western blots showing the levels of DLST (N), DLD (O), and lipoylated proteins (P) in WT and MLL3/4 DKO ESCs. 10 and 20μg of cell lysates were loaded for each sample. Experiments were repeated three times independently and representative results were shown. Protein levels of DLST (in N), DLD (in O), and lipoylated DLST (in P) were quantified by normalizing with HSP90 signals of the corresponding lane.

Figure 4. Simultaneous overexpression of HK2 and OGDH rescues differentiation defects caused by MLL3/4 loss.

(A) Representative images of day 6 WT and MLL3/4 DKO EBs. Scale bar: 250μm.

(B) Quantification of EB sizes in (A). n=50 for each genotype. Experiments were repeated three times independently and representative results were shown.

(C) Correlation plots of RNA-seq data between day 6 MLL3/4 DKO and WT EBs. Data were derived from two biological replicates. Statistical significance was determined by two-sided Wald test and p-values were corrected for multiple testing using the Benjamini–Hochberg method.

Significantly up- and downregulated genes (adjusted p<0.01, │logFC│>1) were labeled in red and

blue with numbers of genes noted, respectively.

(D) Venn diagram identifying 1,192 MLL3/4-dependent differentiation genes.

(E) GO analysis of top 300 MLL3/4-dependent differentiation genes.

(F) Representative images of day 6 MLL3/4 DKO EBs treated with DOX (DKO+DOX) and

MLL3/4 DKO EBs overexpressing HK2/OGDH (DOE+DOX). Scale bar: 250μm.

(G) Quantitation of EB sizes in (F). n=50 for each genotype. Experiments were repeated three

times independently and representative results were shown.

(H) Correlation plots of RNA-seq data between MLL3/4 DKO EBs treated with DOX

(DKO+DOX) and MLL3/4 DKO EBs overexpressing HK2/OGDH (DOE+DOX).

(I) GSEA analysis of MLL3/4-dependent differentiation genes (1,192 in D) comparing MLL3/4

DKO EBs treated with DOX (DKO+DOX) and MLL3/4 DKO EBs overexpressing HK2/OGDH

(DOE+DOX). RES: running enrichment score; RLM: ranked list metric; NES: normalized

enrichment score; FDR: false discovery rate.

(J) Hierarchical clustering analysis of the expression levels of 1,192 MLL3/4-dependent

differentiation genes in MLL3/4 DKO EBs treated with DOX (DKO+DOX) and MLL3/4 DKO

EBs overexpressing HK2/OGDH (DOE+DOX). FPKM: Fragments per kilobase of transcript per

million mapped reads.

(K) Venn diagram showing the overlap of MLL3/4-dependent differentiation genes and

significantly upregulated genes in MLL3/4 DKO EBs overexpressing HK2/OGDH (red in H).

(L) GO analysis of 300 DOE-rescued differentiation genes (overlapping genes in K).

Figure S1. The impact of MLL3/4 loss on glycolytic metabolites, glycolytic gene expression,

and putative enhancers at the *Hk2* locus.

(A-B) Stable isotope tracing of WT and MLL3/4 DKO ESCs labeled with $[^{13}C_6]$-glucose. The

levels of isotopologues (m+0 to m+6) of F6P/G6P/G1P (A) and lactate (B) were shown as stacked

bar plots.

(C) Relative expression levels of genes coding key glycolytic enzymes involved in the anabolism

and catabolism of G6P, F6P, and lactate. CPM (counts per million reads) from RNA-seq data (*14*)

of MLL3/4 DKO ESCs were normalized to those of WT ESCs. p-values were determined by two-

sided Wald test and corrected using the Benjamini–Hochberg method.

(D) Genome browser view of MLL4 ChIP-seq signals in WT ESCs, and that of H3K27ac and

H3K4me3 ChIP-seq signals in WT and MLL3/4 DKO ESCs at the *Hk2* locus. Putative enhancers

bound by MLL4 around *Hk2* gene were highlighted.

Figure S2. The impact of MLL3/4 loss on TCA-cycle metabolites and the expression of TCA-

cycle-related genes.

(A-E) Stable isotope tracing of WT and MLL3/4 DKO ESCs labeled with $[^{13}C_6]$-glucose. The

levels of isotopologues (m+0 to m+6) of citrate/isocitrate (A), α-KG (B), succinate (C), fumarate

(D), and malate (E) were shown as stacked bar plots.

(F-K) Stable isotope tracing of WT and MLL3/4 DKO ESCs labeled with $[^{13}C_5]$-glutamine. The

levels of isotopologues (m+0 to m+6) of glutamate (F), α-KG (G), succinate (H), fumarate (I),

malate (J), and citrate/isocitrate (K) were shown as stacked bar plots.

(L) Relative expression levels of genes coding TCA-cycle enzymes involved in the anabolism and

catabolism of α-KG, succinate, fumarate, and malate. CPM (counts per million reads) from RNA-

seq data (*14*) of MLL3/4 DKO ESCs were normalized to those of WT ESCs. p-values were

determined by two-sided Wald test and corrected using the Benjamini–Hochberg method.

(M-O) Western blots showing levels of OGDH (M), Succinyl-CoA Synthetase subunits (N), SDHA

(O, top blot), and FH (O, middle blot). 10 and 20μg of cell lysates were loaded for each sample.

Experiments were repeated three times independently and representative results were shown.

Protein levels in each lane were quantified by normalizing with Tubulin or HSP90 signals of the

corresponding lane.

Figure S3. HK2 overexpression leads to an increase in the glycolytic activity but a decrease

in mitochondrial respiration in MLL3/4 DKO ESCs.

(A) Western blots showing the induction of 3×HA-tagged HK2 in HK2-overexpressing MLL3/4

DKO ESCs by DOX. 10 and 20μg of cell lysates were loaded for each sample. Experiments were

repeated three times independently and representative results were shown. Protein levels of HK2

in each lane were quantified by normalizing with Tubulin signals of the corresponding lane.

(B) Representative ECAR traces of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and

MLL3/4 DKO ESCs overexpressing HK2 (3×HA-HK2+DOX) measured by the Seahorse XF

glycolysis stress test assay. Data were normalized by the number of cells at the time of the assay

and are represented as mean ± S.D. Each data point had six replicates, and the experiment was

repeated three times independently.

(C-D) Bar plots of ECAR showing the levels of glycolysis (C) and glycolytic capacity (D) in

MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and MLL3/4 DKO ESCs overexpressing

HK2 (3×HA-HK2+DOX). Data are presented as mean ± S.D. p-values were calculated using

Student's t-test.

(E) Representative OCR traces of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and

MLL3/4 DKO ESCs overexpressing HK2 (3×HA-HK2+DOX) measured by the Seahorse XF

mitochondrial stress test assay. Data were normalized by the number of cells at the time of the

assay and are represented as mean ± S.D. Each data point had six replicates, and the experiment

was repeated three times independently.

(F-G) Bar plots showing the OCR levels in MLL3/4 DKO ESCs treated with DOX (DKO+DOX)

and MLL3/4 DKO ESCs overexpressing HK2 (3×HA-HK2+DOX) during basal (F) and maximal

(G) respiration. Data are presented as mean ± S.D. p-values were calculated using Student's t-test.

Figure S4. Stable isotope tracing of MLL3/4 DKO ESCs treated with DOX (DKO+DOX) and

MLL3/4 DKO ESCs overexpressing HK2/OGDH (DOE+DOX) labeled with [^{13}C$_6$]-glucose.

The levels of isotopologues of F6P/G1P/G6P (A, m+0 to m+6), lactate (B m+0 to m+3),

citrate/isocitrate (m+2 in C and m+0 to m+6 in D), α-KG (E, m+0 to m+5), succinate (F, m+0 to

m+4), fumarate (G, m+0 to m+4), and malate (H, m+0 to m+4) were shown.

Figure S5. The impact of MLL3/4 loss and HK2/OGDH overexpression on the expression

levels of marker genes of pluripotency and differentiation in EBs.

(A-B) Relative expression levels of pluripotency genes (A) and differentiation markers (B) in

MLL3/4 DKO EBs compared with WT EBs. Log2 fold change of RNA-seq signals (CPM) were

shown. Data are presented as mean values ± SD. n = 2 biologically independent experiments. All

p-values were smaller than 0.01 compared with WT EBs.

(C-D) Relative expression levels of pluripotency genes (C) and differentiation markers (D) in

MLL3/4 DKO EBs overexpressing HK2/OGDH (DOE+DOX) compared with MLL3/4 DKO EBs

treated with DOX (DKO+DOX). Log2 fold change of RNA-seq signals (CPM) were shown. Data

are presented as mean values ± SD. n = 2 biologically independent experiments. All p-values were

smaller than 0.01 compared with DKO+DOX EBs.

References

1.   S. B. Ng, A. W. Bigham, K. J. Buckingham, M. C. Hannibal, M. J. McMillin, H. I.
590   Gildersleeve, A. E. Beck, H. K. Tabor, G. M. Cooper, H. C. Mefford, C. Lee, E. H. Turner,
591   J. D. Smith, M. J. Rieder, K. Yoshiura, N. Matsumoto, T. Ohta, N. Niikawa, D. A.
592   Nickerson, M. J. Bamshad, J. Shendure, Exome sequencing identifies MLL2 mutations
593   as a cause of Kabuki syndrome. *Nat Genet* **42**, 790-793 (2010).
594   2.   I. D. Krantz, J. McCallum, C. DeScipio, M. Kaur, L. A. Gillis, D. Yaeger, L. Jukofsky, N.
595   Wasserman, A. Bottani, C. A. Morris, M. J. Nowaczyk, H. Toriello, M. J. Bamshad, J. C.
596   Carey, E. Rappaport, S. Kawauchi, A. D. Lander, A. L. Calof, H. H. Li, M. Devoto, L. G.
597   Jackson, Cornelia de Lange syndrome is caused by mutations in NIPBL, the human
598   homolog of Drosophila melanogaster Nipped-B. *Nat Genet* **36**, 631-635 (2004).
599   3.   E. K. Schorry, M. Keddache, N. Lanphear, J. H. Rubinstein, S. Srodulski, D. Fletcher, R.
600   I. Blough-Pfau, G. A. Grabowski, Genotype-phenotype correlations in Rubinstein-
601   Taybi syndrome. *Am J Med Genet A* **146A**, 2512-2519 (2008).
602   4.   T. S. Koemans, T. Kleefstra, M. C. Chubak, M. H. Stone, M. R. F. Reijnders, S. de Munnik,
603   M. H. Willemsen, M. Fenckova, C. Stumpel, L. A. Bok, M. Sifuentes Saenz, K. A. Byerly,
604   L. B. Baughn, A. P. A. Stegmann, R. Pfundt, H. Zhou, H. van Bokhoven, A. Schenck, J.
605   M. Kramer, Functional convergence of histone methyltransferases EHMT1 and
606   KMT2C involved in intellectual disability and autism spectrum disorder. *PLoS Genet*
607   **13**, e1006864 (2017).
608   5.   D. Hu, X. Gao, M. A. Morgan, H. M. Herz, E. R. Smith, A. Shilatifard, The MLL3/MLL4
609   branches of the COMPASS family function as major histone H3K4 monomethylases
610   at enhancers. *Mol Cell Biol* **33**, 4745-4754 (2013).
611   6.   R. Rickels, H. M. Herz, C. C. Sze, K. Cao, M. A. Morgan, C. K. Collings, M. Gause, Y. H.
612   Takahashi, L. Wang, E. J. Rendleman, S. A. Marshall, A. Krueger, E. T. Bartom, A. Piunti,
613   E. R. Smith, N. A. Abshiru, N. L. Kelleher, D. Dorsett, A. Shilatifard, Histone H3K4
614   monomethylation catalyzed by Trr and mammalian COMPASS-like proteins at
615   enhancers is dispensable for development and viability. *Nat Genet* **49**, 1647-1653
616   (2017).
617   7.   K. Cao, C. K. Collings, M. A. Morgan, S. A. Marshall, E. J. Rendleman, P. A. Ozark, E. R.
618   Smith, A. Shilatifard, An Mll4/COMPASS-Lsd1 epigenetic axis governs enhancer
619   function and pluripotency transition in embryonic stem cells. *Sci Adv* **4**, eaap8747
620   (2018).
621   8.   A. Shilatifard, The COMPASS family of histone H3K4 methylases: mechanisms of
622   regulation in development and disease pathogenesis. *Annu Rev Biochem* **81**, 65-95
623   (2012).
624   9.   N. D. Heintzman, R. K. Stuart, G. Hon, Y. Fu, C. W. Ching, R. D. Hawkins, L. O. Barrera,
625   S. Van Calcar, C. Qu, K. A. Ching, W. Wang, Z. Weng, R. D. Green, G. E. Crawford, B.
626   Ren, Distinct and predictive chromatin signatures of transcriptional promoters and
627   enhancers in the human genome. *Nat Genet* **39**, 311-318 (2007).

10. J. E. Lee, C. Wang, S. Xu, Y. W. Cho, L. Wang, X. Feng, A. Baldridge, V. Sartorelli, L.
Zhuang, W. Peng, K. Ge, H3K4 mono- and di-methyltransferase MLL4 is required for
enhancer activation during cell differentiation. *Elife* **2**, e01503 (2013).

11. D. Ashokkumar, Q. Zhang, C. Much, A. S. Bledau, R. Naumann, D. Alexopoulou, A.
Dahl, N. Goveas, J. Fu, K. Anastassiadis, A. F. Stewart, A. Kranz, MLL4 is required after
implantation, whereas MLL3 becomes essential during late gestation. *Development*
147, (2020).

12. V. Faundes, W. G. Newman, L. Bernardini, N. Canham, J. Clayton-Smith, B.
Dallapiccola, S. J. Davies, M. K. Demos, A. Goldman, H. Gill, R. Horton, B. Kerr, D.
Kumar, A. Lehman, S. McKee, J. Morton, M. J. Parker, J. Rankin, L. Robertson, I. K.
Temple, S. Clinical Assessment of the Utility of, S. Evaluation as a Service, S.
Deciphering Developmental Disorders, S. Banka, Histone Lysine Methylases and
Demethylases in the Landscape of Human Developmental Disorders. *Am J Hum*
Genet **102**, 175-187 (2018).

13. D. Rots, S. Choufani, V. Faundes, A. J. M. Dingemans, S. Joss, N. Foulds, E. A. Jones,
S. Stewart, P. Vasudevan, T. Dabir, S. M. Park, R. Jewell, N. Brown, L. Pais, S.
Jacquemont, K. Jizi, C. Ravenswaaij-Arts, H. Y. Kroes, C. Stumpel, C. W. Ockeloen, I.
J. Diets, M. Nizon, M. Vincent, B. Cogne, T. Besnard, M. Kambouris, E. Anderson, E. H.
Zackai, C. McDougall, S. Donoghue, A. O'Donnell-Luria, Z. Valivullah, M. O'Leary, S.
Srivastava, H. Byers, N. Leslie, S. Mazzola, G. E. Tiller, M. Vera, J. J. Shen, R. Boles, V.
Jain, E. Brischoux-Boucher, E. Kinning, B. N. Simpson, J. C. Giltay, J. Harris, B. Keren,
A. Guimier, P. Marijon, B. B. A. Vries, C. S. Motter, B. A. Mendelsohn, S. Coffino, E. H.
Gerkes, A. Afenjar, P. Visconti, E. Bacchelli, E. Maestrini, A. Delahaye-Duriez, C.
Gooch, Y. Hendriks, H. Adams, C. Thauvin-Robinet, S. Josephi-Taylor, M. Bertoli, M. J.
Parker, J. W. Rutten, O. Caluseriu, H. J. Vernon, J. Kaziyev, J. Zhu, J. Kremen, Z. Frazier,
H. Osika, D. Breault, S. Nair, S. M. E. Lewis, F. Ceroni, M. Viggiano, A. Posar, H. Brittain,
T. Giovanna, G. Giulia, L. Quteineh, R. Ha-Vinh Leuchter, E. Zonneveld-Huijssoon, C.
Mellado, I. Marey, A. Coudert, M. I. Aracena Alvarez, M. G. P. Kennis, A. Bouman, M.
Roifman, M. I. Amoros Rodriguez, J. D. Ortigoza-Escobar, V. Vernimmen, M. Sinnema,
R. Pfundt, H. G. Brunner, L. Vissers, T. Kleefstra, R. Weksberg, S. Banka, Pathogenic
variants in KMT2C result in a neurodevelopmental disorder distinct from Kleefstra
and Kabuki syndromes. *Am J Hum Genet* **111**, 1626-1642 (2024).

14. Z. Zhao, K. Cao, J. Watanabe, C. N. Philips, J. M. Zeidner, Y. Ishi, Q. Wang, S. R. Gold,
K. Junkins, E. T. Bartom, F. Yue, N. S. Chandel, R. Hashizume, I. Ben-Sahra, A.
Shilatifard, Therapeutic targeting of metabolic vulnerabilities in cancers with MLL3/4-
COMPASS epigenetic regulator mutations. *J Clin Invest* **133**, (2023).

15. H. Alam, M. Tang, M. Maitituoheti, S. S. Dhar, M. Kumar, C. Y. Han, C. R. Ambati, S. B.
Amin, B. Gu, T. Y. Chen, Y. H. Lin, J. Chen, F. L. Muller, N. Putluri, E. R. Flores, F. J.
DeMayo, L. Baseler, K. Rai, M. G. Lee, KMT2D Deficiency Impairs Super-Enhancers to
Confer a Glycolytic Vulnerability in Lung Cancer. *Cancer Cell* **37**, 599-617 e597 (2020).

16. M. Maitituoheti, E. Z. Keung, M. Tang, L. Yan, H. Alam, G. Han, A. K. Singh, A. T. Raman,
C. Terranova, S. Sarkar, E. Orouji, S. B. Amin, S. Sharma, M. Williams, N. S. Samant,
M. Dhamdhere, N. Zheng, T. Shah, A. Shah, J. B. Axelrad, N. E. Anvar, Y. H. Lin, S. Jiang,
E. Q. Chang, D. R. Ingram, W. L. Wang, A. Lazar, M. G. Lee, F. Muller, L. Wang, H. Ying,

K. Rai, Enhancer Reprogramming Confers Dependence on Glycolysis and IGF Signaling in KMT2D Mutant Melanoma. *Cell Rep* **33**, 108293 (2020).

17. W. Zhou, M. Choi, D. Margineantu, L. Margaretha, J. Hesson, C. Cavanaugh, C. A. Blau, M. S. Horwitz, D. Hockenbery, C. Ware, H. Ruohola-Baker, HIF1alpha induced switch from bivalent to exclusively glycolytic metabolism during ESC-to-EpiSC/hESC transition. *EMBO J* **31**, 2103-2116 (2012).

18. C. D. Folmes, T. J. Nelson, A. Martinez-Fernandez, D. K. Arrell, J. Z. Lindor, P. P. Dzeja, Y. Ikeda, C. Perez-Terzic, A. Terzic, Somatic oxidative bioenergetics transitions into pluripotency-dependent glycolysis to facilitate nuclear reprogramming. *Cell Metab* **14**, 264-271 (2011).

19. A. D. Panopoulos, O. Yanes, S. Ruiz, Y. S. Kida, D. Diep, R. Tautenhahn, A. Herrerias, E. M. Batchelder, N. Plongthongkum, M. Lutz, W. T. Berggren, K. Zhang, R. M. Evans, G. Siuzdak, J. C. Izpisua Belmonte, The metabolome of induced pluripotent stem cells reveals metabolic changes occurring in somatic cell reprogramming. *Cell Res* **22**, 168-177 (2012).

20. A. Moussaieff, M. Rouleau, D. Kitsberg, M. Cohen, G. Levy, D. Barasch, A. Nemirovski, S. Shen-Orr, I. Laevsky, M. Amit, D. Bomze, B. Elena-Herrmann, T. Scherf, M. Nissim-Rafinia, S. Kempa, J. Itskovitz-Eldor, E. Meshorer, D. Aberdam, Y. Nahmias, Glycolysis-mediated changes in acetyl-CoA and histone acetylation control the early differentiation of embryonic stem cells. *Cell Metab* **21**, 392-402 (2015).

21. A. De Los Angeles, F. Ferrari, R. Xi, Y. Fujiwara, N. Benvenisty, H. Deng, K. Hochedlinger, R. Jaenisch, S. Lee, H. G. Leitch, M. W. Lensch, E. Lujan, D. Pei, J. Rossant, M. Wernig, P. J. Park, G. Q. Daley, Hallmarks of pluripotency. *Nature* **525**, 469-478 (2015).

22. P. Du, J. Wu, Hallmarks of totipotent and pluripotent stem cell states. *Cell Stem Cell* **31**, 312-333 (2024).

23. C. Buecker, R. Srinivasan, Z. Wu, E. Calo, D. Acampora, T. Faial, A. Simeone, M. Tan, T. Swigut, J. Wysocka, Reorganization of enhancer patterns in transition from naive to primed pluripotency. *Cell Stem Cell* **14**, 838-853 (2014).

24. C. Zeng, J. Chen, E. W. Cooke, A. Subuddhi, E. T. Roodman, F. X. Chen, K. Cao, Demethylase-independent roles of LSD1 in regulating enhancers and cell fate transition. *Nat Commun* **14**, 4944 (2023).

25. E. A. Rowland, T. M. Greco, C. K. Snowden, A. L. McCabe, T. J. Silhavy, I. M. Cristea, Sirtuin Lipoamidase Activity Is Conserved in Bacteria as a Regulator of Metabolic Enzyme Complexes. *mBio* **8**, (2017).

26. E. A. Rowland, C. K. Snowden, I. M. Cristea, Protein lipoylation: an evolutionarily conserved metabolic regulator of health and disease. *Curr Opin Chem Biol* **42**, 76-85 (2018).

27. J. E. Cronan, Assembly of Lipoic Acid on Its Cognate Enzymes: an Extraordinary and Essential Biosynthetic Pathway. *Microbiol Mol Biol Rev* **80**, 429-450 (2016).

28. P. S. J. Bailey, B. M. Ortmann, A. W. Martinelli, J. W. Houghton, A. S. H. Costa, S. P. Burr, R. Antrobus, C. Frezza, J. A. Nathan, ABHD11 maintains 2-oxoglutarate metabolism by preserving functional lipoylation of the 2-oxoglutarate dehydrogenase complex. *Nat Commun* **11**, 4046 (2020).

29. J. A. Lessard, G. R. Crabtree, Chromatin regulatory mechanisms in pluripotency. *Annu*
Rev Cell Dev Biol **26**, 503-532 (2010).
30. S. Feng, S. E. Jacobsen, W. Reik, Epigenetic reprogramming in plant and animal
development. *Science* **330**, 622-627 (2010).
31. B. Papp, K. Plath, Epigenetics of reprogramming to induced pluripotency. *Cell* **152**,
1324-1343 (2013).
32. W. W. Tee, D. Reinberg, Chromatin features and the epigenetic regulation of
pluripotency states in ESCs. *Development* **141**, 2376-2390 (2014).
33. S. Schlesinger, E. Meshorer, Open Chromatin, Epigenetic Plasticity, and Nuclear
Organization in Pluripotency. *Dev Cell* **48**, 135-150 (2019).
34. A. Burton, M. E. Torres-Padilla, Epigenome dynamics in early mammalian
embryogenesis. *Nat Rev Genet* **26**, 587-603 (2025).
35. X. Wan, S. Zhang, J. Li, D. Kong, M. Chen, Epigenetic remodeling during early
embryonic development. *Front Cell Dev Biol* **14**, 1750381 (2026).
36. R. Eskeland, M. Leeb, G. R. Grimes, C. Kress, S. Boyle, D. Sproul, N. Gilbert, Y. Fan, A.
I. Skoultchi, A. Wutz, W. A. Bickmore, Ring1B compacts chromatin structure and
represses gene expression independent of histone ubiquitination. *Mol Cell* **38**, 452-
464 (2010).
37. R. S. Illingworth, M. Moffat, A. R. Mann, D. Read, C. J. Hunter, M. M. Pradeepa, I. R.
Adams, W. A. Bickmore, The E3 ubiquitin ligase activity of RING1B is not essential for
early mouse development. *Genes Dev* **29**, 1897-1902 (2015).
38. C. C. Sze, K. Cao, C. K. Collings, S. A. Marshall, E. J. Rendleman, P. A. Ozark, F. X. Chen,
M. A. Morgan, L. Wang, A. Shilatifard, Histone H3K4 methylation-dependent and -
independent functions of Set1A/COMPASS in embryonic stem cell self-renewal and
differentiation. *Genes Dev* **31**, 1732-1737 (2017).
39. L. Wang, C. K. Collings, Z. Zhao, K. A. Cozzolino, Q. Ma, K. Liang, S. A. Marshall, C. C.
Sze, R. Hashizume, J. N. Savas, A. Shilatifard, A cytoplasmic COMPASS is necessary
for cell survival and triple-negative breast cancer pathogenesis by regulating
metabolism. *Genes Dev* **31**, 2056-2066 (2017).
40. K. M. Dorighi, T. Swigut, T. Henriques, N. V. Bhanu, B. S. Scruggs, N. Nady, C. D. Still,
2nd, B. A. Garcia, K. Adelman, J. Wysocka, Mll3 and Mll4 Facilitate Enhancer RNA
Synthesis and Transcription from Promoters Independently of H3K4
Monomethylation. *Mol Cell* **66**, 568-576 e564 (2017).
41. K. Cao, M. Ugarenko, P. A. Ozark, J. Wang, S. A. Marshall, E. J. Rendleman, K. Liang, L.
Wang, L. Zou, E. R. Smith, F. Yue, A. Shilatifard, DOT1L-controlled cell-fate
determination and transcription elongation are independent of H3K79 methylation.
Proc Natl Acad Sci U S A **117**, 27365-27373 (2020).
42. M. A. J. Morgan, A. Shilatifard, Epigenetic moonlighting: Catalytic-independent
functions of histone modifiers in regulating transcription. *Sci Adv* **9**, eadg6593 (2023).
43. A. Khan, C. Zhang, P. G. Nguyen, J. M. Metts, L. C. Collins, K. Jain, C. A. Mills, L. Vlach,
K. Li, A. L. Brademeyer, B. M. Bowman, M. B. Major, J. Aube, L. E. Herring, W. K.
Rathmell, F. M. Mason, I. J. Davis, Q. Zhang, B. D. Strahl, A SETD2-CDK1-lamin axis
maintains nuclear morphology and genome stability. *Nat Cell Biol* **27**, 1327-1341
(2025).

44.   C. Wang, J. E. Lee, B. Lai, T. S. Macfarlan, S. Xu, L. Zhuang, C. Liu, W. Peng, K. Ge,
761   Enhancer priming by H3K4 methyltransferase MLL4 controls cell fate transition. *Proc*
762   *Natl Acad Sci U S A* **113**, 11871-11876 (2016).
763   45.   C. C. Sze, A. Shilatifard, MLL3/MLL4/COMPASS Family on Epigenetic Regulation of
764   Enhancer Function and Cancer. *Cold Spring Harb Perspect Med* **6**, (2016).
765   46.   G. Xie, J. E. Lee, A. D. Senft, Y. K. Park, Y. Jang, S. Chakraborty, J. J. Thompson, K.
766   McKernan, C. Liu, T. S. Macfarlan, P. P. Rocha, W. Peng, K. Ge, MLL3/MLL4
767   methyltransferase activities control early embryonic development and embryonic
768   stem cell differentiation in a lineage-selective manner. *Nat Genet* **55**, 693-705 (2023).
769   47.   J. W. Hong, D. A. Hendrix, M. S. Levine, Shadow enhancers as a source of evolutionary
770   novelty. *Science* **321**, 1314 (2008).
771   48.   K. Cao, C. K. Collings, S. A. Marshall, M. A. Morgan, E. J. Rendleman, L. Wang, C. C.
772   Sze, T. Sun, E. T. Bartom, A. Shilatifard, SET1A/COMPASS and shadow enhancers in
773   the regulation of homeotic gene expression. *Genes Dev* **31**, 787-801 (2017).
774   49.   E. Z. Kvon, R. Waymack, M. Gad, Z. Wunderlich, Enhancer redundancy in
775   development and disease. *Nat Rev Genet* **22**, 324-336 (2021).
776   50.   Y. Tsukada, J. Fang, H. Erdjument-Bromage, M. E. Warren, C. H. Borchers, P. Tempst,
777   Y. Zhang, Histone demethylation by a family of JmjC domain-containing proteins.
778   *Nature* **439**, 811-816 (2006).
779   51.   M. Tahiliani, K. P. Koh, Y. Shen, W. A. Pastor, H. Bandukwala, Y. Brudno, S. Agarwal, L.
780   M. Iyer, D. R. Liu, L. Aravind, A. Rao, Conversion of 5-methylcytosine to 5-
781   hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* **324**, 930-
782   935 (2009).
783   52.   W. Xu, H. Yang, Y. Liu, Y. Yang, P. Wang, S. H. Kim, S. Ito, C. Yang, P. Wang, M. T. Xiao,
784   L. X. Liu, W. Q. Jiang, J. Liu, J. Y. Zhang, B. Wang, S. Frye, Y. Zhang, Y. H. Xu, Q. Y. Lei, K.
785   L. Guan, S. M. Zhao, Y. Xiong, Oncometabolite 2-hydroxyglutarate is a competitive
786   inhibitor of alpha-ketoglutarate-dependent dioxygenases. *Cancer Cell* **19**, 17-30
787   (2011).
788   53.   M. Xiao, H. Yang, W. Xu, S. Ma, H. Lin, H. Zhu, L. Liu, Y. Liu, C. Yang, Y. Xu, S. Zhao, D.
789   Ye, Y. Xiong, K. L. Guan, Inhibition of alpha-KG-dependent histone and DNA
790   demethylases by fumarate and succinate that are accumulated in mutations of FH
791   and SDH tumor suppressors. *Genes Dev* **26**, 1326-1338 (2012).
792   54.   K. Skvortsova, E. Masle-Farquhar, P. L. Luu, J. Z. Song, W. Qu, E. Zotenko, C. M. Gould,
793   Q. Du, T. J. Peters, Y. Colino-Sanguino, R. Pidsley, S. S. Nair, A. Khoury, G. C. Smith, L.
794   A. Miosge, J. H. Reed, J. G. Kench, M. A. Rubin, L. Horvath, O. Bogdanovic, S. M. Lim,
795   J. M. Polo, C. C. Goodnow, C. Stirzaker, S. J. Clark, DNA Hypermethylation
796   Encroachment at CpG Island Borders in Cancer Is Predisposed by H3K4
797   Monomethylation Patterns. *Cancer Cell* **35**, 297-314 e298 (2019).
798   55.   E. W. Cooke, C. Zeng, S. M. Nur, Y. Jia, A. Huang, J. Chen, P. Gao, F. X. Chen, F. Jin, K.
799   Cao, H3K79 methylation and H3K36 trimethylation synergistically regulate gene
800   expression in pluripotent stem cells. *Sci Adv* **11**, eadt8765 (2025).
801   56.   A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina
802   sequence data. *Bioinformatics* **30**, 2114-2120 (2014).

57.   A. Dobin, C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M.
Chaisson, T. R. Gingeras, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,
15-21 (2013).
58.   F. Ramirez, D. P. Ryan, B. Gruning, V. Bhardwaj, F. Kilpert, A. S. Richter, S. Heyne, F.
Dundar, T. Manke, deepTools2: a next generation web server for deep-sequencing
data analysis. *Nucleic Acids Res* **44**, W160-165 (2016).
59.   S. Anders, P. T. Pyl, W. Huber, HTSeq--a Python framework to work with high-
throughput sequencing data. *Bioinformatics* **31**, 166-169 (2015).
60.   M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* **26**,
139-140 (2010).
61.   T. Wu, E. Hu, S. Xu, M. Chen, P. Guo, Z. Dai, T. Feng, L. Zhou, W. Tang, L. Zhan, X. Fu,
S. Liu, X. Bo, G. Yu, clusterProfiler 4.0: A universal enrichment tool for interpreting
omics data. *Innovation (Camb)* **2**, 100141 (2021).
62.   Y. Zhou, B. Zhou, L. Pache, M. Chang, A. H. Khodabakhshi, O. Tanaseichuk, C. Benner,
S. K. Chanda, Metascape provides a biologist-oriented resource for the analysis of
systems-level datasets. *Nat Commun* **10**, 1523 (2019).