# Expression levels of α5 subunit-containing GABA-A receptors in the prelimbic cortex are associated with visual perceptual learning

Matthew C. D. Bailey[1], Emilie Preisler[1], Clara Valazquez Sanchez[1], Lucia Marti Prats[1], Olivia Stupart[1], Livia J.F. Wilod Versprille[1], Johann F. du Hoffmann[2], Zoe Kourtzi[1], Jeffrey W. Dalley[1,3,#]

1 Department of Psychology, University of Cambridge, Downing St, Cambridge CB2 3EB, UK; 2 Boehringer Ingelheim Pharma GmbH & Co. KG, Div. Research Germany, Biberach an der Riβ, Germany; 3 Department of Psychiatry, Herschel Smith Building for Brain and Mind Sciences, University of Cambridge CB2 0SZ, UK

Short title: Inhibitory modulation of visual perceptual learning

Word count: 5,701
Abstract: 219
Introduction: 583
Discussion: 1,853
Figures: 5

# **Author for correspondence**: Professor Jeffrey W. Dalley, Department of Psychology,
University of Cambridge, Downing St, Cambridge CB2 3EB, UK. Tel. +44(0)1223 765 291;
Email: jwd20@cam.ac.uk

**Abstract**

Perceptual learning is a temporally dynamic process involving the acquisition and
integration of sensory information necessary for adaptive decision making. Resolving the
neural basis of perceptual learning could uncover new therapeutic targets for schizophrenia
and other neurodevelopmental disorders that implicate impaired perceptual acuity. In the
present study, we developed a novel touchscreen task which utilizes orientation
discrimination to assess visual perceptual learning (VPL) in male and female rats. Based on
previous evidence we hypothesised that VPL would depend on inhibitory neurotransmission
mediated by γ-amino butyric acid (GABA). Segregating subjects based on 'poor learning'
(lower tertile) and 'good learning' (upper tertile) revealed dose-dependent improvements in
VPL in poor learners following the administration of a GABA-B agonist (R-baclofen) and an
α5 subunit specific GABA-A (GABRA5) positive allosteric modulator (alogabat)
administered early in learning. Poor VPL performance was associated with a significant
reduction in GABRA5 expression in dorsal regions of the prefrontal cortex (PFC), most
notably the prelimbic cortex. Reduced GABRA5 expression in this region was co-localized
to somatostatin- and parvalbumin-expressing interneurons. These findings indicate that
inter-individual variation in the expression of GABRA5 in selective PFC populations of
inhibitory interneurons may determine the speed and acuity of VPL. Based on these
findings, interventions that restore GABRA5 signalling in the PFC may hold therapeutic
relevance for neuropsychiatric disorders involving deficits in perceptual learning.

**Key words:** GABA; adaptive decision making; spatial transcriptomics; primary visual
cortex; prefrontal cortex; parvalbumin interneurons

## 1.0 Introduction

Visual perceptual learning (VPL) refers to long-term improvements in visual discrimination that emerge after repeated training. Previous studies have revealed finely tuned stimulus- and context-specific cortical responses during VPL[1–4]. Such responses are linked to synaptic plasticity in primary visual cortex (V1) with intrinsic neurons exhibiting small receptive fields and narrow feature selectivity[5–7]. Higher visual processing areas including parietal and frontal cortices involved in decision-making and attention are progressively also recruited alongside classical visual processing areas such as V1 and superior colliculus to refine behavioral output and improve visual discrimination[8–10]. However, the relative contributions of bottom-up sensory *versus* top-down decisional and attentional control in VPL remain poorly understood.

Inhibitory control is fundamental to reshaping cortical circuits throughout development and adulthood, coordinating plasticity during sensitive periods[11,12] and VPL[13–16]. γ-Aminobutyric acid (GABA), the primary inhibitory neurotransmitter in the brain acts in concert with glutamate, the primary excitatory neurotransmitter, to regulate local circuit and global network excitability *via* the excitatory-inhibitory balance (E/I). Dysregulated E/I balance is associated with perceptual deficits in neurodevelopmental disorders such as autism [17,18] and schizophrenia[19–21], with aberrant inhibition driving network abnormalities[22–27]. Resolving the mechanism by which inhibition contributes to VPL may thus inform the development of new interventions to improve perceptual learning in a variety of disorders.

At the cortical circuit level, distinct populations of inhibitory interneurons contribute to experience-dependent remodelling of sensory, associative and executive networks recruited during learning and attentional tasks[28–31]. These include parvalbumin (Pvalb)-, somatostatin (Sst)-, and vasoactive intestinal peptide (VIP)- expressing neurons in V1, which dynamically remap their activity as VPL proceeds[31–35]. During late stages of learning, increased attentional engagement recruits higher-order inhibitory microcircuits to sharpen stimulus selectivity and enhance network fidelity[36–39]. However, within these microcircuits, GABAergic interneurons exhibit significant receptor and receptor composition heterogeneity and are recruited in a context dependent manner to modify network excitability[40–43].

Given the established role of the α5 subunit containing GABA-A receptor (GABRA5) in hippocampal-dependent learning and memory[44–47], the present study investigated the potential of GABRA5 to enhance VPL. Although GABRA5 represent less than 5% of

cortically expressed GABA-A receptors, they are expressed both intra- and extra-

synaptically and thus contribute respectively to phasic and tonic inhibition[45,48,49]. GABRA5

receptors also show dynamic activity-dependent synaptic re-localisation, a process

hypothesised to facilitate fine tuning of tonic and phasic inhibitory tone to preserve learned

associations in a context dependent manner[50]. Thus, modulation of GABRA5 receptors may

offer a promising strategy to rescue or enhance visual perceptual learning in autism,

schizophrenia and other neurodevelopmental disorders.

To investigate the role of GABRA5 in VPL we established a novel touchscreen task for rats

involving the discrimination of visual gratings with distinct orientations. We compared sex-

differences in task acquisition and baseline performance before testing the sensitivity of the

task to a range of GABAergic compounds that affect distinct aspects of GABAergic synaptic

transmission and stratified these effects according to poor and good learning performance

on the task. Given that the rate of learning in visual touchscreen discrimination tasks is

linked to thresholds of visual acuity[51] we also investigated the dimensional relationship of

VPL in individual animals and expression levels of the GABRA5 mRNA transcript and

protein product in V1 and frontal cortex. We hypothesised that enhancing GABAergic

inhibition with selective GABA reuptake inhibition (tiagabine) and a positive allosteric

modulator of GABRA5 (alogabat)[52] would enhance VPL. We further hypothesised that inter-

individual variation in VPL would relate to GABRA5 expression with faster visual perceptual

learning in animals exhibiting increased GABRA5 expression.

**2.0 Methods**

**2.1 Subjects**

Thirty-two Lister-hooded rats (16 male, 16 female) were raised from time mated dams

(Envigo, Bicester, UK), weaned at post-natal-day (PND) 21, and housed in groups of four

with environmental enrichment under a 12/12hr reverse light/dark cycle (lights on 6am).

From weaning, animals had *ad libitum* access to food (Safe A40, Germany) and water. After

PND26, animals were handled 3 times per week to habituate them to human contact. On

PND56 animals were food restricted to no less than 85% free feeding weight for the

remainder of the experiment. Animals were weighed at least once per week. Animals were

sacrificed at the end of the experiment by schedule 1 methods for tissue collection. All

procedures were carried out in accordance with the UK (1986) Animal (Scientific

Procedures) Act (PPL PP9536688) and were approved by the University of Cambridge

Animal Welfare and Ethical Review Body (AWERB).

**2.2 Visual perceptual learning task**

Operant chambers running K-Limbic software (Med Associates, UK) were integrated with a

1024x768 resolution portrait touchscreen (Nexio, UK) on one wall, and a food magazine

with light, pellet dispenser, house light, and tone generator on the opposite wall (see Figure

1A). A camera was installed overhead to record behavior.

To investigate VPL, a novel 3-stimuli orientation-based visual discrimination paradigm

adapted from Horner and colleagues[53] was developed using sinusoidal stimuli presented in

three equally spaced response windows (200x200 pixels), located 150 pixels from the

bottom of the touchscreen. Target stimuli were assigned at the beginning of training,

counterbalanced within the cohort (16 animals 0°, 16 animals 90°) and spatially randomised

on a trial-by-trial basis to reduce procedural learning and location bias. Pretraining

proceeded with chamber acclimatisation with reward pellets (TestDiet AIN-76, 5TUL, Cat

No.1811155) (PT1), appetitive autoshaping to the touchscreen by association of target

grating with reward (PT2), timeout for background touches (PT3), and the introduction of

distractors in the form of two blank stimuli (PT4). Criteria for progression was >80% correct

on 2 sessions (see Figure S1).

Animals then completed a series of visual discrimination training sessions to assess the

suitability of chosen angular separations from target (ang_sep) before moving to the final

task. Beginning with a 90° (VD1) separation from the target stimuli, training progressed by

decreasing the separation angle through 45°, 30°, 15° and finally 10° (VD5). Animals were

advanced to the next training stage depending on difficulty after 2 consecutive sessions at

criteria: >80% 90°→45°, >70% 45°→30°, >60% 30°→15°, >50% 15°→10°, >50% 10°→final

task. Animals that reached VD5 but failed to reach criteria after 5 sessions were deemed to

have reached their perceptual limit. All animals were moved to the final task for

pharmacological interventions.

The final task involved 150 trials split into five, 30-trial blocks of distractor orientation which

converged on the target stimulus orientation (Easy - 90°, 45°, 30°, 15° and 10° - Hard – in

degrees from target angle). On any given trial, three stimuli were presented, one target

grating, one distractor grating, and one white stimulus randomised on a trial-by-trial basis.

Here, target distractor distance (TDD) was classified as 'close' when target grating and

distractor grating were presented in adjacent windows, and 'far' when target distractor

gratings were separated by the third, white stimuli. TDD was employed to investigate the

role of attentional load and simultaneous suppression of multiple stimuli on perceptual

decision making. A correct response was rewarded with a sucrose pellet and auditory tone

whereas incorrect responses at either the distractor stimuli (distractor grating or white) or

background were signalled with a 10s house light pulse timeout and no food reward.

Sessions lasted on average 1 hour with an inter-trial-interval (ITI) of 10s (see Figure 1B, C).

For further details of the training stages and task contingencies refer to Figure S1.

**Figure 1 (overleaf): visual discrimination task and experimental timeline**
**A** Operant chamber apparatus. Med Associates chambers showing the position of the house-light, tone generator and food magazine with a light connected to a pellet dispenser on one side of the chamber and a 1024 x 768 resolution touchscreen on the other. A CCTV camera was mounted overhead to record behavior. **B** Trial wise decomposition of final task contingencies. A nose poke in the illuminated food magazine initiated a trial. Stimuli were then presented on the touchscreen (1 target grating, 2 distractors (1 blank, 1 grating) with animals making a choice between the 3 presented stimuli. A correct response was reinforced with a sugar pellet paired with tone (magazine light on during collection). Incorrect responses at either the distractor or background stimuli received a 10 second timeout with no reward and house light illumination. **C** Baseline task perceptual challenge and distractor presentation sequence. In total, the task consisted of 150 trials in blocks of 30 at each distractor frequency, starting at 90° from target. Stimuli were presented randomly on a trial-by-trial basis. **D** Experimental timeline showing behavioral stratification, pharmacological challenges and molecular analyses. Figure created using https://app.biorender.com/

## 2.3 Subject stratification

To investigate individual differences in visual perceptual learning, animals were stratified into 'learning type' groups based on their task acquisition rate. For males (n=16), animals that completed pretraining and VD training during the thirty-three-session training period were classified as "good learners" (n=5), whereas animals that failed to reach visual discrimination training were classified as "poor learners" (n=6). Animals that completed pretraining but did not finish VD training were classed as "intermediate learners" (n=5) and represented the mean task acquisition rate. Only good and poor learners were considered for final analysis to investigate what drives good vs poor VPL.

Although females were included in the study, it was not possible to use the same stratification procedure as males because considerably more sessions were required for females to acquire the task (mean 52 sessions) and baseline task response rates were variable. Moreover, females trained on the basic discrimination task failed to complete the more difficult training stages of the VPL task, as shown in Figure S2, meaning that no animals were classed as 'good learners' despite the extended training period. Instead, most females were classed as 'poor' (n=8), with the remainder which either reached criterion on the PT4, or begun VD training were classed as 'intermediate' (n=8). These groups were used in comparing within-sex performance, pharmacology and molecular variations, and can be found in the supplementary material section.

**2.4 Pharmacological interventions**

Selective GABAergic compounds were administered according to a Latin square design. The compounds (and doses) were tested in with a 3-day dosing schedule (day 1 (dose level 1), day 2 (no drug), day 3 (baseline) and at least 3 day washout period between each compound: (1) alogabat (3, 10, 30 mg/kg) (2) tiagabine hydrochloride (0.5, 2.5, 7.5 mg/kg); (3) R-baclofen (0.05, 0.5, 1.5 mg/kg). Alogabat (provided by Boehringer Ingelheim Pharma, Germany) was administered in 5% HCl (1M) and 95% 2-(hydroxypropyl)-β-cyclodextrin (20%) at 2 ml/kg (pH 7-7.5). Tiagabine-HCl (Insight Biotechnology, UK-CAS 145821-59-6) was administered in deionised water pH 7-7.5 at 1 ml/kg. R-Baclofen (Insight Biotechnology, UK-CAS 69308-37-8) was administered in 0.9% saline (pH 7-7.5) at 1 ml/kg. Tiagabine and R-baclofen were administered simultaneously to male and female animals, whilst alogabat was administered first to males, and as the final compound to females, accounting for the time difference in pertaining. All drugs were administered *via* intraperitoneal (i.p.) injection with 30-minute pretreatment time (See Figure 1D and Figure S10 for further details).

**2.5 Tissue collection**

Rats were anesthetised with isoflurane (Sigma-Aldrich, UK) *via* inhalation before decapitation 5 days after the final VPL session. Brains were collected and snap-frozen with dry-ice pre-cooled isopentane (Sigma-Aldrich, UK) and stored at -80°C. Brains were sectioned using a cryostat (Leica, UK) to obtain coronal slices of the medial prefrontal cortex (mPFC) (5.16 to 2.28 mm from Bregma) and V1 (-5.6 to -9.1 mm from Bregma). Each hemisphere was prepared for Western blotting (WB) with sections cut at 200 μm or RNAscope *in situ* hybridisation (RNAScope) with sections cut at 12 μm. Hemispheres were counterbalanced between the two assays, and slices were mounted on Menzel gläzer superfrost® slides (Thermo Fisher Scientific, UK) for WB or Superfrost® plus slides (Thermo Fisher Scientific, UK) for RNAscope. Sections were stored at -80°C prior to processing.

**2.6 Western immunoblot analysis**

WB were conducted using a WB kit according to manufacturer guidelines (Invitrogen-Life Technologies, UK). Four regions-of-interest (ROI)s were extracted from the slices of each rat: V1, Cg1, PrL, and IL. After immunoblotting, GABRA5 expression was developed using a primary antibody for GABRA5 (Biorbyt, UK – 1:1000 Cat No. orb374299) incubated

overnight at 4°C on a shaker. The primary antibody for the housekeeping protein actin

(Abcam, UK – 1:10,000 Cat No. ab8226) was incubated for 1h at room temperature.

Secondary antibodies, IRDye680RD and IRDye800CW (LI-COR, UK – 1:15,000 Cat No:

680, 926-68071; 800, 926-32210) were incubated at room temperature for 1h alongside a

chameleon duo ladder (LI-COR, UK – Cat No. 928-60000) before imaging (Odyssey

Imaging System, LI-COR, UK).

**2.7 RNAscope fluorescence in-situ hybridisation**

The spatial transcriptomic profile of GABRA5 in the mPFC and V1 was assayed in replicate

samples to investigate cell-type specific GABRA5 mRNA expression. The assay and target

probes were performed[54] according to the fresh frozen tissue protocol provided by the

manufacturer (RNAscope fluorescent multiplex reagent kit, ACD, UK) with specific details

given in the supplementary section (Table 1).

Tissue was imaged using a Zeiss Axio Imager M2 microscope with an AxioCam MRm

microscope camera (Oberkochen, Germany) running Visiopharm software (Medicon Valley,

Denmark). Approximately 15 images were each taken of V1 and mPFC subregions

(cingulate cortex - Cg1; prelimbic cortex – PrL; infralimbic cortex - IL) with 20X

magnification. Signals were quantified using an RNAscope signal analysis pipeline

developed by Velazquez-Sanchez et al.[55], which utilised a MATLAB script (The MathWorks

Inc) combined with a FIJI/ImageJ[56] machine learning algorithm to count mRNA molecules

within DAPI labelled cells.

**2.8 Data analysis**

Behavioral data were collected on a trial-by-trial basis and pre-processed to remove

outliers. Baseline outliers were removed based on the following criteria: <96/150 trials

completed in each session – only reaching the third block of distractor, and response

latencies (<100ms) that were too fast to be instrumental in nature (e.g. arising from

unintentional tail contacts with the touchscreen). For the pharmacological experiments,

outliers were removed for failed/incomplete dosing or a failure to reach the trial completion

criteria on <2/4 doses of each compound (n=1 removed per compound), final n: n=5 poor

learners, n=5 good learners. Reaction time responses were inverse transformed (-100/RT)

to normalise distribution of the data and facilitate more robust modelling.

Statistical comparisons were made in R (Version 4.4.2)[57] using within-subjects mixed-

effects modelling, a generalised logit link function to quantify trial wise accuracy, and a

linear model to analyse inverse transformed mean response latencies (inverseRT). Fixed-

effects relationships between angular separation from target (ang_sep), sex, learning type

(LT) and target distractor distance (TDD) were investigated. Animal ID was added as a

random effect with a random intercept to account for within group individual variation (see

Figure S3). Pharmacological models were similar, although model selection showed TDD

did not have significant explanatory power and was therefore not included in the final

model. Ang_sep was also only included as a fixed effect rather than interaction due to

model convergence violations. Where appropriate, estimated-marginal-means pairwise

comparisons with Tukey HSD correction were performed and reported as log-odds ratios for

generalised models.

The analysis of Western blot data was carried out using Welch's two sample t-test to

compare learning type means within cortical region. A within-subjects linear mixed-effects

model was applied to molecular data to investigate the relationships between learning-type

and GABRA5 mRNA expression within specified cell types in the various brain regions of

interest. Each animal was added as a random effect with a random intercept to account for

within group individual variation. An ANOVA was performed to identify main effects, and

where appropriate, estimated-marginal-means pairwise comparisons with Tukey HSD

correction were performed between groups within region, within cell type, and adjusted p

values ≤ 0.05 reported (see Figure S4). All variables and models were performed and

evaluated for analysis suitability using normality and model evaluation metrics available in

the Performance package in R (see Figure S5). Full summaries of statistical models can be

found in supplementary table 2.

**3.0 Results**

**3.1 Modelling VPL with a novel orientation discrimination task**

Assessment of performance on the baseline version of the VPL task revealed that

narrowing the angular separation from the target stimulus decreased discriminative

accuracy (Figure 2A) (Compared to ang_sep90: ang_sep45: $Pr(>|z|)$ = 0.970, ang_sep30:

$Pr(>|z|)$ = $2.14e^{-05}$, ang_sep15: $Pr(>|z|)$ < $2e^{-16}$, ang_sep10, $Pr(>|z|)$ < $2e^{-16}$). Here there

was also a within-block sex difference in trial wise mean accuracy where females displayed

lower accuracy from block 2 onwards (ang_sep90, F-M: adj.p value = 0.418, ang_sep45, F-

M: adj.p value = 0.0471 ang_sep30, F-M: adj.p value = 0.0531 ang_sep15, F-M: adj.p value

= 0.0275, ang_sep10, F-M: adj.p value = 0.0247). However, the accuracy of males and

females across all blocks was not significantly different (Compared to F: M: $Pr(>|z|)$ =

0.732) (Figure 2B). When the relative location of the target and distractor gratings was

considered (Target Distractor Distance – TDD), both sexes were impaired by a greater

distance between stimuli (F, TDDclose – TDDfar: adj.p value = <.0001. M, TDDclose –

TDDfar: adj.p value = 0.0002), with females showing a greater reduction in accuracy than

males (TDDfar, F-M: adj.p value = 0.02) (Figure 2C). There was no such effect with close

TDD (TDDclose, F-M: adj.p value = 0.163). Analysis of concordant inverse transformed

response latencies (inverseRT) revealed a slowing of responses with narrowing angular

separation (Figure 2D) ($F_{(4,269)}$ = 123, $Pr(>|z|)$ < $2.2e^{-16}$) and a modest blockwise sex effect

(ang_sep90, F-M: adj.p value = 0.848, ang_sep45, F-M: adj.p value = 0.363, ang_sep30, F-

M: adj.p value = 0.204, ang_sep15, F-M: adj.p value = 0.0696, ang_sep10, F-M: adj.p value

= 0.0781 (Figure 2E). However, there was no significant difference between males and

females on inverseRT across the full task ($F_{(1, 29)}$ = 1.54, $Pr(>F)$ 0.225) or when analysed

by TDD (Sex:TDD interaction, $F_{(1, 269)}$ = 0.224, $Pr(>F)$ = 0.637 (Figure 2F).

**Figure 2 (overleaf): Baseline performance measures on the VPL task**
**A** Trial-wise mean accuracy by sex presented as mean ± SEM. Dashed lines denote distractor frequency blocks. Annotations denote between-sex comparisons within each distractor frequency block. **B** Distribution of baseline task accuracy segregated by sex. Each data point denotes an individual animal. **C** Distribution of mean task accuracy for each target distractor distance (TDD). Comparisons made within TDD-between sex and within sex-between TDD. **D** Trial-wise mean inverseRT by sex presented as mean ± SEM. Dashed lines mark distractor frequency blocks. Annotations denote between-sex comparisons, within distractor frequency block. **E** Distribution of mean inverseRT segregated by sex. Each data point denotes an individual animal **F** Distribution of mean inverseRT by TDD. Comparisons made within TDD-between sex and within sex-between TDD.
Baseline sessions displayed for each sex with statistical comparisons made by generalized linear mixed-effects model with binomial link for accuracy **(A-C).** Linear mixed-effects model for inverseRT **(D-F)**. Boxplots denote median, interquartile range with whiskers extending to 1.5x IQR. Outliers denoted by solid points. Pairwise comparisons made via estimated-marginal-means to "Female" condition in each case, within block **(A,D)** and "close" **(C,F)**. Effect sizes: "p value" = p<0.1, "*" = p<0.05, "***" = p<0.001 **(D-F)** as log-odds ratios **(A-C)**.

## 3.2 Segregation of good *versus* poor learners on the VPL task

After validating the perceptual challenge of our paradigm, subjects were stratified based on their VPL performance. Given that, for difficult learning paradigms, rodents require extensive training and are often over trained by the time they are evaluated at baseline, we segregated animals during the task acquisition stage. To achieve this, we grouped subjects by their final training stage before the drug challenges. Those which completed all pretraining, reaching the baseline task were classed as "Good Learners" (n = 5), those which did not complete pretraining were "Poor Learners" (n = 6) (Figure 3A). Unfortunately, due to differing pre-training exposures, we could not directly compare sexes with this stratification approach, instead we completed a concordant stratification analysis for female animals available in the supplementary materials (Figures S7, S8, S9).

In males, groupwise comparisons revealed within-block learning differences in accuracy (Figure 3B) where good learners were more accurate than poor learners in all but the final block (ang_sep90, poor-good: adj.p value = 0.0166, ang_sep45, poor-good: adj.p value =

0.0057, ang_sep30, poor-good: adj.p value = 0.0083, ang_sep15, poor-good: adj.p value =

0.0095, ang_sep10, poor-good: adj.p value = 0.169).This was associated with good

learners displaying a greater full task accuracy (Figure 3C) (poor – good: Pr(>|z|) = 0.0253)

and both TDD close and far accuracies than poor learners (Figure 3D) (TDDclose, Poor-

Good: adj.p value = 0.0258, TDDfar, Poor-Good adj.p value = 0.0114). Additionally, whilst

poor learner accuracy was negatively impacted by increased TDD, good learners displayed

only a marginal reduction in performance between TDDclose and TDDfar trials (Poor,

TDDclose – TDDfar: adj.p value <.0001, Good, TDDclose – TDDfar: adj.p value = 0.0599).

Analysis of inverseRT highlighted no blockwise (Figure 3E) (ang_sep90, Poor-Good: adj.p

value = 0.263, ang_sep45, Poor-Good:  adj.p value = 0.203 ang_sep30, Poor-Good: adj.p

value = 0.17, ang_sep15, Poor-Good: adj.p value = 0.204, ang_sep10, Poor-Good: adj.p

value = 0.254) or taskwise (Figure 3F) differences between learning types (poor – good: $F_{(1, 9)}$ = 1.81, Pr(>F) = 0.212. There were also no within or between group effects of TDD on

inverse RT (Figure 3G) ($F_{(1, 89)}$ = 0.558, Pr(>F) = 0.457).

**Figure 3 (overleaf) Summary measures of perceptual performance on the VPL task**
**A** Summary of stratification procedure based on training stage reached at the end of 33 sessions of pretraining (males only). **B** Trial-wise mean accuracy by learning type presented as mean ± SEM. Dashed lines mark distractor frequency blocks. Annotations denote between-learning type comparisons, within distractor frequency block. **C** Distribution of baseline accuracy by learning type. **D** Distribution of mean accuracy segregated by target distractor distance. Points denote individual animals per TDD. Comparisons made within TDD-between learning type and within learning type-between TDD. **E** Trial-wise mean inverseRT by learning type presented as mean ± SEM. Dashed lines mark distractor frequency blocks. Annotations denote between-learning type comparisons, within distractor frequency block. **F** Distribution of mean inverseRT by learning type.
**G** Distribution of mean inverseRT by target distractor distance. Points denote individual animals per TDD. Comparisons made within TDD-between learning type and within learning type-between TDD. Baseline sessions are displayed for each learning type with statistical comparisons made by generalized linear mixed-effects model with binomial link for accuracy (**B-D**). Linear mixed-effects model for inverseRT (**E-G**). Boxplots denote median, interquartile range with whiskers extending to 1.5x IQR. Outliers denoted by solid points. Pairwise comparisons made via estimated-marginal-means to "Poor learner" condition in each case, within block **(B,E)** and "close" **(D,G)**. Effect sizes: "*" = p<0.05, "**" = p<0.01**(E-G)** as log-odds ratios **(B-D)**.

Bailey et al. 2026

| Learning Type | Criteria | Total Animals |
|---|---|---|
| Poor Learner | Failed complete Pre-training | 6 |
| Good Learner | Reached final task | 5 |

### 3.3 GABAergic pharmacology

To investigate the contribution of GABRA5 to VPL we administered the GABRA5 positive allosteric modulator (PAM) alogabat during task performance. To interrogate the specificity of the behavioral effects of this GABRA5-based compound we also administered two clinically available compounds: tiagabine, a GABA reuptake inhibitor and baclofen, a GABA B agonist.

As shown in Figure 4A, tiagabine had no significant effect on accuracy (compared to Veh: 0.5 mg/kg, $Pr(>|z|) = 0.831$, 2.5 mg/kg, $Pr(>|z|) = 0.461$, 7.5 mg/kg, $Pr(>|z|) = 0.961$). However, in good learners, tiagabine had a dose dependent effect on response latencies (main effect of dose: $F_{(3,161)} = 17.3$, $Pr(>F) = 8.88e^{-10}$). Here, low (Veh-0.5 mg/kg: adj.p value = 0.0365) and middle doses (Veh-2.5 mg/kg: adj.p value = 0.0001) increased response speed whereas the highest dose slowed responding compared to all other doses (Veh-7.5 mg/kg: adj.p value = 0.031, 0.5 mg/kg-7.5 mg/kg: adj.p value < 0.0001, 2.5 mg/kg-7.5 mg/kg: adj.p value < 0.0001) (Figure 4B).

Administration of R-baclofen impacted accuracy in both learning groups (Figure 4C) (compared to Veh: 0.05 mg/kg, $Pr(>|z|) = 0.862$, 0.5 mg/kg, $Pr(>|z|) = 2.89e^{-05}$, 1.5 mg/kg, $Pr(>|z|) = 0.00182$). Here, middle-dose R-baclofen improved "Good" learner accuracy (Good: Veh-0.5 mg/kg: adj.p value = 0.048), and had a positive effect on accuracy in "Poor" learners (Poor: Veh - 0.5 mg/kg: adj.p value = 0.0002, 0.05 mg/kg – 0.5 mg/kg: adj.p value = 0.0001). High-dose baclofen improved the accuracy of "Poor" leaners (Poor: Veh-1.5 mg/kg: adj.p value = 0.0099, Veh-0.5 mg/kg: adj.p value = 0.008) but this comparison was relatively low powered due to subject exclusion (Figure S6). Low dose R-baclofen had no impact on "Poor" learner accuracy (Poor: Veh-0.05 mg/kg: adj.p value = 0.998), but there was a positive trend associated with "Good" learner performance (Good: Veh-0.05 mg/kg: adj.p value = 0.0868). Whilst there was a main effect of R-baclofen dose on inverseRT ($F_{(3,139)} = 6.05$, $Pr(>F) = 6.68e^{-04}$), this was limited to "Poor" learners only. Here, dose wise comparisons revealed faster responses at the intermediate dose (Poor: Veh-0.5 mg/kg: adj.p value = 0.0026), and slowing at high doses (Poor: Veh-1.5 mg/kg: adj.p value = 0.0014, 0.05-1.5 mg/kg: adj.p value < 0.0001, 0.5-1.5 mg/kg: adj.p value < 0.0001) compared to vehicle and low dose conditions (Figure 4D).

Like baclofen, alogabat administration modified accuracy in both learning groups (Veh-3 mg/kg, $Pr(>|z|) = 1.30e^{-05}$, Veh-10 mg/kg, $Pr(>|z|) = 9.87e^{-12}$, Veh-30 mg/kg, $Pr(>|z|) =$

0.479). Within groups, "Poor" learner accuracy demonstrated an 'inverted-U' dose-response

with stepwise improvements in accuracy from vehicle at 3 mg/kg (Poor:Veh-3 mg/kg: adj.p

value = 0.0001), and 10 mg/kg doses (Poor:Veh-10 mg/kg: adj.p value < 0.0001) whereas

high dose exhibited no difference compared with the vehicle control group (Poor:Veh-30

mg/kg: adj.p value = 0.894). In contrast, a dose of 30 mg/kg alogabat improved "Good"

learner accuracy (adj.p value = 0.0001) with lower doses showing comparable accuracy

with the vehicle group (Figure 4E) (Veh-3 mg/kg: adj.p value = 0.701, Veh-10 mg/kg: adj.p

value = 0.986). Like tiagabine, alogabat had significant effects on response latencies in

"Good" learners only. Thus, a dose of 10 mg/kg resulted in faster responses compared with

vehicle (Veh-10 mg/kg: adj.p value = 0.0275) and 3 mg/kg (3 -10 mg/kg: adj.p value =

0.0193), whilst 30 mg/kg slowed responding compared to the 10 mg/kg dose level (10-30

mg/kg: adj.p value < 0.0001) (Figure 4F).

**Figure 4 Baseline-dependent effects of GABAergic interventions on the VPL task**
**A** Distribution of full task mean accuracy following the administration of the GAT-1 inhibitor tiagabine separated by learning group. **B** Distribution of full task mean inverseRT following the administration of the GAT-1 inhibitor tiagabine separated by learning group. **C** Distribution of full task mean accuracy following the administration of the GABA B agonist R-baclofen separated by learning group. **D** Distribution of full task mean inverseRT following the administration of GABA B agonist R-baclofen separated by learning group. **E** Distribution of full task mean accuracy following the administration of the GABRA5 positive allosteric modulator alogabat separated by learning group. **F** Distribution of full task mean inverseRT following the administration of the GABRA5 positive allosteric modulator alogabat separated by learning group.

Generalized linear mixed-effects model with binomial link **(A,C,E)**. Linear mixed-effects model **(B,D,F)**. Boxplots denote median, interquartile range with whiskers extending to 1.5x IQR. Outliers denoted by solid points. Pairwise comparisons made via estimated marginal means to Vehicle condition in each case, within "Learning type". Effect sizes: "p value" = p<0.1, "*" = p<0.05, "**" = p<0.01, "***" = p<0.001**(B,D,F)** and as log-odds ratios **(A,C,E)**.

**3.4 Relating perceptual performance to GABRA5 expression**

To identify the putative substrates underlying the variability in alogabat's drug effects we next investigated GABRA5 expression in primary visual cortex (V1) and subregions of the mPFC (Cg1, PrL, IL – see Figure 5A) using Western immunoblot and RNA scope fluorescence in-situ-hybridisation (RNA Scope) (Figure 5B-D).

Western blot protein analysis highlighted an increased level of GABRA5 protein expression in the PrL of good learners (Poor - Good: p-value = 0.0393) (Figure 5E). We then assessed GABRA5 mRNA in selective subpopulations of inhibitory interneurons, specifically: parvalbumin (Pvalb)-, somatostatin (Sst)-, vasoactive intestinal peptide (VIP)- and excitatory pyramidal cells (slc17A7) in each target cortical region of interest. Statistical comparison between learning types revealed a greater number of GABRA5 and Sst co-expressing cells in the PrL of good learners (Poor - Good: adj.p value = 0.0006) (Figure 5F), an increased number of GABRA5 and Sst overlapping mRNA copies in the Cg1 and PrL of good learners (Cg1:Poor - Good: adj.p value = 0.0187, PrL: Poor - Good: adj.p value = 0.0038), and a greater number of GABRA5 and Pvalb overlapping mRNA copies in the PrL of good learners (Poor - Good: adj.p value = 0.0031) (Figure 5G). There was no difference in within-region total-DAPI derived learning-type cell counts (LT:Region - $F_{(3,22.4)}$ = 0.102, Pr(>F) = 0.958). All other comparisons were non-significant. A summary of the molecular quantification data is given in supplementary Table 2.

**Figure 5 (overleaf) Relationship of GABRA5 expression perceptual acuity and learning rate**
**A** Schematic illustrations showing target regions of interest, left = primary visual cortex (V1) -5.6mm – -8.16 mm from bregma, right = prefrontal cortex (PFC) 4.6 mm – 2.52 mm from bregma. Sub-regions: anterior cingulate cortex (Cg1), prelimbic cortex (PrL), infralimbic cortex (IL). **B** Representative section of PFC showing fluorescent signal aligned with ROIs. **C** RNAscope assays from representative good versus poor learners on the VPL task. Scale bar 100 µm. Assay 1 GABRA5 (red); Pvalb (green); Sst (yellow): Assay 2 GABRA5 (red); slc17A7 (green); VIP (yellow). Arrows highlight cells expressing target probes in each case. **D** Representative Western-immunoblot gel for GABRA5 protein expression in good versus poor learning groups with actin as the housekeeper control for each ROI. **E** Statistical comparison of Western-immunoblot derived GABRA5 protein expression in cortical ROIs between stratified "learning types". **F** Cell-wise quantification of GABRA5 co-expression from RNAscope assays within inhibitory interneuron subtypes: parvalbumin (Pvalb), somatostatin (Sst), vasoactive intestinal peptide (VIP) and excitatory pyramidal cells (slc17A7) by "Learning type", per target cortical region. **G** Puncta-wise quantification of GABRA5 co-expression from RNAscope assays within inhibitory interneuron subtypes: parvalbumin (Pvalb), somatostatin (Sst), vasoactive intestinal peptide (VIP) and excitatory pyramidal cells (slc17A7) by "learning type" in target cortical regions. A modulus transformation (p=0.3) was applied to scale the RNAscope y-axes to enable better interpretability between high and low expression values on the same plot. mRNA coexpression values are displayed for each "learning type" as mean ± SEM per subject. RNAscope was performed using 2 assays to assess GABRA5 within (1: Pvalb, Sst, 2: VIP, slc17A7) highlighted with dashed line. Pairwise comparisons made via linear mixed-effects model to "poor learner" condition each case via estimated-marginal-means, within region and cell type: "p value" = p<0.1, "*" = p<0.05, "**" = p<0.01, "***" = p<0.001.

**4.0 Discussion**

In this study we investigated the role of inhibitory neurotransmission in VPL and specifically an underlying involvement of α5 subunit containing GABA-A receptors. Using a novel orientation discrimination task, we assessed the association between GABRA5 expression and task acquisition and validated our findings with selective GABAergic pharmacological agents in both male and female animals. We report that superior learning on a visual discrimination perceptual task is accompanied by increased GABRA5 expression in Sst and Pvalb interneuronal populations in the prelimbic subregion of the medial prefrontal cortex. Since VPL acuity in poor learners was improved following positive allosteric GABRA5 modulation during the early task sessions, our findings indicate that VPL may depend in part on optimal GABRA5 mediated inhibition in the prelimbic cortex early in the learning process.

**4.1 Modelling perceptual learning and attentional load**

Orientation discrimination is a commonly employed behavioral paradigm to investigate perceptual acuity in rodents[28,38] and humans[58,59]. However, cross-species translation is often challenged by differences in task configuration and decoding precisely how rodents solve tasks of this nature. To address these issues, we developed a novel progressive orientation discrimination paradigm to test rodent visual perceptual learning in a manner as close to human perceptual tasks as possible. Like virtual-reality or head-fixed running wheel approaches[60,61], we used progressive orientation discrimination as the perceptual challenge. However, unlike 2-choice perceptual tasks in rodents[62], we added a third stimulus. By simultaneously presenting target and distractor gratings close or far from one another, we not only reduced procedural location-based learning but also widened the window for pharmacological effects (from chance performance of 50% to 33%). In addition, the configuration of our novel task increased attentional demands[63], thus mimicking the high attentional load of equivalent VPL tasks in humans[64]. Crucially, our approach also preserved fine discrimination without restraint, facilitating naturalistic behavior and negating the limitations of head fixed approaches commonly used in go/no-go-based perceptual tasks. With these features, our task permitted the investigation and decoding of multiple components of visual perceptual learning whilst also enabling future temporally resolved neural recording techniques such as fiber photometry[65] and electrocorticography (ECoG)[66,67].

**4.2 VPL variability is associated with GABRA5 expression in the prelimbic cortex**

To investigate intrinsic differences in perceptual learning in our task, we stratified behavior

at the conclusion of the pretraining stage. At this point in acquisition, stimulus reward

associations (PT2+3), distractor interference tolerance (PT4, VD1-5), and spatial flexibility

(PT2-VD5) could be assessed. Our approach aimed to separate the upper and lower

tertiles of learning rate to determine whether GABRA5 expression in key components of the

visual processing pathway were associated with task acquisition and could function as

predictors of subsequent baseline performance.

VPL involves the integration of bottom-up sensory processing and coordinated responses

from top-down attentional control mechanisms[68–73]. Our finding that higher GABRA5 protein

expression in a subregion of the PFC – the PrL – is associated with faster perceptual

learning indicates that top-down inhibitory control may be an important determinant of early

contingency learning in this task. Supporting this notion, animals showing faster rates of

learning were also more tolerant of a wider separation between the target and distractor

stimuli, consistent with a role of the PrL in maintaining attentional focus in the face of

distraction[74,75]. Indeed, higher cortical regions including the PFC and parietal association

cortex (PC) are implicated in the control of interference suppression[76] in response to

multiple simultaneously presented visual stimuli on such tasks as multiple-object-tracking

(MOT)[77] and visual discrimination[78]. Such observations imply that top-down control

processes may be necessary for acquiring attentional selection strategies to enable

discriminative acuity in the present VPL task.

Within each session we found block-wise differences in VPL performance between fast and

slower learning male and female rats, which diminished toward the most challenging

discrimination block (10° from target). This finding suggests that whilst faster contingency

learning during pretraining is beneficial in overall task performance, when the task reaches

fine levels of discrimination, even robust interference suppression strategies are ineffective.

Since the PrL contributes to the formation of stimulus reward action-associations and

dynamic cue-outcome predictions when the stimulus is either 'rewarded' or 'not rewarded',

rather than changes to its 'value'[79,80] we suggest that when target and distractor stimuli

converge in orientation, so too do their perceived values. Here, groupwise perceptual

strategies recruited early in learning and putatively modulated by GABRA5 in the PrL may

break down due to high levels of uncertainty between the perceived values of target and

distractor stimuli. Instead, processing may depend more on bottom-up sensory integration

or other stimulus-value associated regions such as the orbitofrontal cortex[81,82]. However, it

could also be argued that 10-degree discrimination represents a perceptual limit for rats in

this task. Taken together, these findings suggest that PrL GABRA5 expression may enable

faster acquisition of top-down stimulus-reward association weights and interference

suppression strategies during pretraining. However, during late VPL, a limit is reached for

fine discriminations that may be independent of GABRA5 in the PrL.

**4.3 GABRA5 selective agents may alter performance by modifying tonic inhibitory**

**activity**

The pharmacological interventions in this study aimed to validate the observed differential

expression of GABRA5 in the PrL of good and poor learners. To achieve this objective we

administered a specific GABRA5 PAM, alogabat, and compared its effects to clinical

compounds that potentiate metabolic (tiagabine) and post-synaptic tonic inhibitory activity

(R-baclofen), both of which have demonstrated pro-cognitive effects in different

contexts[83,84]. Here, we found a lack of efficacy in metabolic GABA potentiation with

tiagabine, but a positive dose-dependent effect of baclofen on discriminative accuracy in

both poor and good learning male rats. In contrast, alogabat elicited a dose-dependent

enhancement in VPL performance that varied according to learning group and timing of

administration.

Our findings with tiagabine and baclofen suggest that extracellular GABA levels do not

impact task performance at nonsedative doses, but that moderate post-synaptic

potentiation via the GABA B receptor improves perceptual acuity regardless of task

acquisition rate whilst sparing side effects. However, the effects of alogabat are more

challenging to interpret. It should be noted that administration timing of alogabat was

variable, firstly between male and female animals (sessions 1-13 male, 16-23 female), and

secondly, based on relative task experience between good and poor learning males

(sessions: 1-8 poor, ~5-13 good) (Figure S10). We acknowledge this variation as a

limitation but suggest the accompanied behavioural effects shed light on the time

dependent mechanisms by which GABRA5 acts in this task.

Alogabat administration resulted in an apparent inverted-U dose response effect on

accuracy in poor learning males, and a positive log-linear dose response in good learning

males. Considering poor learning males were yet to asymptote during administration we

suggest that potentiating GABRA5 in this window enables faster learning of the top-down

stimulus reward associations. Whereas, in good learning males which did not display

asymptotic performance, refinements could only be made at the highest dose (details of

session-wise baseline performance found in Figure S11). Whilst these responses relate to

groupwise PrL GABRA5 expression, the similar performance effects of baclofen and

alogabat in male rats suggests that alogabat may be modifying cortical tonic inhibition.

Tonic GABA currents are typically viewed as constraining learning and memory while

sharpening sensory processing (reviewed by Koh., et al. 2023)[85]. Although GABA B

receptors contribute substantially to tonic inhibition, several extrasynaptic GABA A subtypes

including GABRA5 also mediate shunting inhibition[45,86,87]. This GABA A-based mechanism

has been proposed to narrow the temporal integration window of incoming signals, thereby

increasing the signal-to-noise ratio and improving the fidelity of sensory representations[88,89].

In this context, pharmacological potentiation of GABRA5 would be expected to bias

microcircuitry toward stability and noise-suppression, potentially enhancing distractor

resistance. Indeed, consistent with the clinical indications of alogabat [52] and similar pro-

cognitive GABRA5 PAMs [90,91] we suggest that modulating GABRA5 in this way may benefit

refinements in early plastic learning periods that appear to rely on receptor expression

presumably related to dynamic synaptic localisation.

Our pharmacological interventions support the notion that extrasynaptic GABRA5 receptors

regulate a balance between stability and plasticity favouring stability for faster task

acquisition. However, tuning by GABRA5 pharmacology during early exposure to the final

task may enable long-term refinements of stimuli representations, which depend on intrinsic

receptor expression. Moreover, once these representations are well trained, ceiling effects

of increased tonic GABA inhibition is evident.

**4.4 Sst and pvalb interneuron GABRA5 mRNA expression underlies behavioral and**

**pharmacological variation**

Our RNAScope findings indicate that higher GABRA5 expression in mPFC pvalb and Sst

interneuron populations benefit early task learning and long-term baseline performance.

Based on the established connections between GABRA5 and Sst interneurons with

dendritic input gating refinement to pyramidal cells (PYR) [92–96], we posit that sensory signal-

to-noise ratio and input integration mediated by these cells may be an important

determinant of distractor suppression mediated by the PrL during VPL acquisition. A

putative involvement of Pvalb interneurons also suggests GABRA5+ cells contribute to PrL

recurrent inhibitory activity to fine tune the local E/I balance, thereby enhancing task

acquisition via trial-by-trial prediction error updates[97,98]. The absence of GABRA5

expression differences in VIP and slc17A7 expressing PYR cells potentially highlights

canonical microcircuit mechanisms where VIP interneurons disinhibit PYR excitability

through inhibition of Sst suppression of PYR cells and would be bypassed by increased

tonic inhibition from GABRA5 expressing pvalb and Sst populations[16,92]. These findings

suggest that GABRA5 clusters in Pvalb and Sst interneurons of the PrL function to bias the

microcircuitry towards greater tonic inhibitory tuning of sensory input fidelity. This

mechanism would account for faster learning in animals exhibiting a higher intrinsic

expression of GABRA5.

The profound differences observed in the acquisition of the present VPL task in male and

female rats precludes a meaningful comparison to be made in terms of drug effects and

GABRA5 dependent mechanisms. The longer acquisition observed in female rats suggests

females may be more susceptible to distraction interference than males. Indeed, a previous

study also reported longer training periods of female rats in similar 3-stimuli operant task

compared to males[99]. Additionally, we acknowledge that the timing of Alogabat

administration renders stable performance vs learning effects inseparable. Although, we

argue that based on the consistency of RNAScope findings across the cohort, the

similarities in experienced animal responses to alogabat and baclofen, and the dynamic

synaptic localisation of GABRA5 receptors, our results display useful learning-based

findings on the contributions of GABRA5. Finally, our results present longitudinal behaviour,

but static molecular readouts.

**4.5 Conclusions**

Overall, we report a convergent set of findings implicating GABRA5 mediated inhibitory

processes during adaptive visually guided behaviors. We report a putative involvement of

GABRA5 expressed Sst and pvalb interneurons of the PrL in a visual orientation

discrimination task that involves set-shifting and simultaneous suppression of distractor

stimuli in the visual field. We validated GABRA5's role in our task via comparing selective

GABAergic agents between groups of animals stratified according to poor and good

learning during early acquisition of the task. Our findings indicate that suppression of

interference by the PrL may facilitate faster learning in animals exhibiting high expression of

GABRA5 in PrL Sst and Pvalb expressing interneurons. Thus, in conclusion, GABRA5

expressed in the PrL may mediate tonic inhibitory control of excitatory sensory input to

facilitate top-down learning of perceptual task contingencies.

**Acknowledgements**

This work was supported by the Wellcome Trust (223131/Z/21/Z) and a studentship from

Boehringer Ingelheim Pharma. The experimental work was carried out under a Home Office

Project Licence held by Prof. Dr. A. L. Milton.

**Author Contributions**

Conceptualization: MCDB. Experimental design: MCDB. Data analysis: MCDB, OSRPS,

LVW. Investigation: MCDB, EP, CVS, LMP. Visualization: MCDB, EP, CVS, LMP.

Supervision: JFdH, JWD. Writing–original draft: MCDB JWD. Writing–review & editing: all

authors

**Conflicts of Interest**

The authors declare no competing interests

## 5.0 References

1.   Ahissar, M. & Hochstein, S. Task difficulty and the specificity of perceptual learning. *Nature* **387**, 401–6
(1997).
2.   Ball, K. & Sekuler, R. Direction-specific improvement in motion discrimination. *Vision Res.* **27**, 953–965
(1987).
3.   Karni, A. & Sagi, D. Where practice makes perfect in texture discrimination: evidence for primary visual
cortex plasticity. *Proceedings of the National Academy of Sciences* **88**, 4966–4970 (1991).
4.   Shiu, L. P. & Pashler, H. Improvement in line orientation discrimination is retinally local but dependent
on cognitive set. *Percept. Psychophys.* **52**, 582–8 (1992).
5.   Hubel, D. H. & Wiesel, T. N. Receptive fields, binocular interaction and functional architecture in the
cat's visual cortex. *J. Physiol.* **160**, 106–154 (1962).
6.   Schoups, A., Vogels, R., Qian, N. & Orban, G. Practising orientation identification improves orientation
coding in V1 neurons. *Nature* **412**, 549–553 (2001).
7.   Hubel, D. H. & Wiesel, T. N. Receptive fields and functional architecture of monkey striate cortex. *J.
Physiol.* **195**, 215–243 (1968).
8.   Mukai, I. *et al.* Activations in Visual and Attention-Related Areas Predict and Correlate with the Degree
of Perceptual Learning. *The Journal of Neuroscience* **27**, 11401–11411 (2007).
9.   Kahnt, T., Grueschow, M., Speck, O. & Haynes, J.-D. Perceptual Learning and Decision-Making in
Human Medial Frontal Cortex. *Neuron* **70**, 549–559 (2011).
10.  Jing, R., Yang, C., Huang, X. & Li, W. Perceptual learning as a result of concerted changes in
prefrontal and visual cortex. *Current Biology* **31**, 4521-4533.e3 (2021).
11.  Ganguly, K., Schinder, A. F., Wong, S. T. & Poo, M. ming. GABA itself promotes the developmental
switch of neuronal GABAergic responses from excitation to inhibition. *Cell* **105**, 521–532 (2001).
12.  Kirmse, K. *et al.* GABA depolarizes immature neurons and inhibits network activity in the neonatal
neocortex in vivo. *Nat. Commun.* **6**, (2015).
13.  Froemke, R. C. Plasticity of cortical excitatory-inhibitory balance. *Annu. Rev. Neurosci.* **38**, 195–219
(2015).
14.  van Versendaal, D. & Levelt, C. N. Inhibitory interneurons in visual cortical plasticity. *Cellular and
Molecular Life Sciences* **73**, 3677–3691 (2016).
15.  Desrosiers, J. & Basha, D. Cortical Inhibition, Plasticity, and Sleep. *The Journal of Neuroscience* **43**,
523–525 (2023).
16.  Tremblay, R., Lee, S. & Rudy, B. GABAergic Interneurons in the Neocortex: From Cellular Properties to
Circuits. *Neuron* **91**, 260–292 (2016).
17.  Hollestein, V. *et al.* Excitatory/inhibitory imbalance in autism: the role of glutamate and GABA gene-
sets in symptoms and cortical brain structure. *Transl. Psychiatry* **13**, 1–9 (2023).
18.  Cooper, R. A. *et al.* Reduced hippocampal functional connectivity during episodic memory retrieval in
autism. *Cerebral Cortex* **27**, 888–902 (2017).
19.  Lesh, T. A., Niendam, T. A., Minzenberg, M. J. & Carter, C. S. Cognitive control deficits in
schizophrenia: Mechanisms and meaning. *Neuropsychopharmacology* **36**, 316–338 (2011).
20.  RA McCutcheon, R. K. P. M. Cognitive impairment in schizophrenia: aetiology, pathophysiology, and
treatment. *Mol. Psychiatry* **28**, 1902–1918 (2023).
21.  Howes, O. D., Bukala, B. R. & Beck, K. Schizophrenia: from neurochemistry to circuits, symptoms and
treatments. *Nature Reviews Neurology 2023 20:1* **20**, 22–35 (2023).
22.  van Hugte, E. J. H., Schubert, D. & Kasri, N. N. E/I balance in epilepsies and neurodevelopmental
disorders: depolarizing GABA as a common mechanism. *Epilepsia* 1–15 (2023) doi:10.1111/epi.17651.
23.  Ferguson, B. R. & Gao, W. J. Pv interneurons: critical regulators of E/I balance for prefrontal cortex-
dependent behavior and psychiatric disorders. *Front. Neural Circuits* **12**, (2018).
24.  Mueller-Buehl, C., Wegrzyn, D., Bauch, J. & Faissner, A. Regulation of the E/I-balance by the neural
matrisome. *Front. Mol. Neurosci.* **16**, (2023).

25. Kaar, S. J., Nottage, J. F., Angelescu, I., Marques, T. R. & Howes, O. D. Gamma Oscillations and Potassium Channel Modulation in Schizophrenia: Targeting GABAergic Dysfunction. *Clin. EEG Neurosci.* **55**, 203–213 (2024).

26. Hoftman, G. D. *et al.* Altered cortical expression of GABA-related genes in schizophrenia: Illness progression vs developmental disturbance. *Schizophr. Bull.* **41**, 180–191 (2015).

27. Marín, O. Parvalbumin interneuron deficits in schizophrenia. *Eur Neuropsychopharmacol* **82**, 44–52 (2024).

28. Poort, J. *et al.* Learning and attention increase visual response selectivity through distinct mechanisms. *Neuron* **110**, 686–697 (2022).

29. Yazaki-Sugiyama, Y., Kang, S., Câteau, H., Fukai, T. & Hensch, T. K. Bidirectional plasticity in fast-spiking GABA circuits by visual experience. *Nature* **462**, 218–21 (2009).

30. Kuchibhotla, K. V *et al.* Parallel processing by cortical inhibition enables context-dependent behavior. *Nat. Neurosci.* **20**, 62–71 (2017).

31. Kato, H. K., Gillet, S. N. & Isaacson, J. S. Flexible Sensory Representations in Auditory Cortex Driven by Behavioral Relevance. *Neuron* **88**, 1027–1039 (2015).

32. Cheng, R. *et al.* A disinhibitory microcircuit in the temporal association cortex for fear retrieval to pure tones. *Sci. Rep.* **15**, 1–16 (2025).

33. Khan, A. G. *et al.* Distinct learning-induced changes in stimulus selectivity and interactions of GABAergic interneuron classes in visual cortex. *Nat. Neurosci.* **21**, 851–859 (2018).

34. Myers-Joseph, D., Wilmes, K. A., Fernandez-Otero, M., Clopath, C. & Khan, A. G. Disinhibition by VIP interneurons is orthogonal to cross-modal attentional modulation in primary visual cortex. *Neuron* **112**, 628-645.e7 (2024).

35. Aime, M. *et al.* Paradoxical somatodendritic decoupling supports cortical plasticity during REM sleep. *Science (1979).* **376**, 724–730 (2022).

36. Snyder, A. C., Morais, M. J. & Smith, M. A. Dynamics of excitatory and inhibitory networks are differentially altered by selective attention. *J. Neurophysiol.* **116**, 1807–1820 (2016).

37. Snyder, A. C., Yu, B. M. & Smith, M. A. A Stable Population Code for Attention in Prefrontal Cortex Leads a Dynamic Attention Code in Visual Cortex. *Journal of Neuroscience* **41**, 9163–9176 (2021).

38. Speed, A., Del Rosario, J., Mikail, N. & Haider, B. Spatial attention enhances network, cellular and subthreshold responses in mouse visual cortex. *Nat. Commun.* **11**, 505 (2020).

39. Raven, F. & Aton, S. J. The Engram's Dark Horse: How Interneurons Regulate State-Dependent Memory Processing and Plasticity. *Front. Neural Circuits* **15**, (2021).

40. Olsen, R. W. & Sieghart, W. GABAA receptors: Subtypes provide diversity of function and pharmacology. *Neuropharmacology* **56**, 141–148 (2009).

41. Weir, C. J. Ion channels, receptors, agonists and antagonists. *Anaesthesia & Intensive Care Medicine* **17**, 645–651 (2016).

42. Pinard, A., Seddik, R. & Bettler, B. GABA B receptors. Physiological functions and mechanisms of diversity. in *Advances in Pharmacology* vol. 58 231–255 (2010).

43. Kohl, M. M. & Paulsen, O. The roles of GABA B receptors in cortical network activity. in *Advances in Pharmacology* vol. 58 205–229 (Academic Press Inc., 2010).

44. Prut, L. *et al.* A reduction in hippocampal GABA $_A$ receptor α5 subunits disrupts the memory for location of objects in mice. *Genes Brain Behav.* **9**, 478–488 (2010).

45. Caraiscos, V. B. *et al.* Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by α5 subunit-containing γ-aminobutyric acid type A receptors. *Proceedings of the National Academy of Sciences* **101**, 3662–3667 (2004).

46. Zarnowska, E. D., Keist, R., Rudolph, U. & Pearce, R. A. GABAA receptor α5 subunits contribute to GABA A,slow synaptic inhibition in mouse hippocampus. *J. Neurophysiol.* **101**, 1179–1191 (2009).

47. Engin, E. *et al.* Tonic Inhibitory Control of Dentate Gyrus Granule Cells by α5-Containing GABA $_A$ Receptors Reduces Memory Interference. *The Journal of Neuroscience* **35**, 13698–13712 (2015).

48. Serwanski, D. R. *et al.* Synaptic and nonsynaptic localization of GABAA receptors containing the alpha5 subunit in the rat brain. *J. Comp. Neurol.* **499**, 458–70 (2006).

49.    Panzanelli, P. *et al.* Distinct mechanisms regulate GABA A receptor and gephyrin clustering at
perisomatic and axo-axonic synapses on CA1 pyramidal cells. *J. Physiol.* **589**, 4959–4980 (2011).

50.    Davenport, C. M. *et al.* Relocation of an Extrasynaptic GABAA Receptor to Inhibitory Synapses
Freezes Excitatory Synaptic Strength and Preserves Memory. *Neuron* **109**, 123-134.e4 (2021).

51.    Crijns, E. & Op de Beeck, H. The Visual Acuity of Rats in Touchscreen Setups. *Vision* **4**, 4 (2019).

52.    Cecere, G. *et al.* Preclinical pharmacology of alogabat: a novel GABAA-α5 positive allosteric modulator
targeting neurodevelopmental disorders with impaired GABAA signaling. *Front. Pharmacol.* **16**,
1626078 (2025).

53.    Horner, A. E. *et al.* The touchscreen operant platform for testing learning and memory in rats and mice.
*Nat. Protoc.* **8**, 1961–1984 (2013).

54.    Wang, F. *et al.* RNAscope. *The Journal of Molecular Diagnostics* **14**, 22–29 (2012).

55.    Velazquez-Sanchez, C., Muresan, L., Marti-Prats, L. & Belin, D. The development of compulsive
coping behaviour is associated with a downregulation of Arc in a Locus Coeruleus neuronal ensemble.
*Neuropsychopharmacology* **48**, 653–663 (2023).

56.    Schindelin, J. *et al.* Fiji: An open-source platform for biological-image analysis. *Nature Methods* vol. 9
676–682 Preprint at https://doi.org/10.1038/nmeth.2019 (2012).

57.    R: The R Project for Statistical Computing. https://www.r-project.org/.

58.    Jia, K. *et al.* Visual perceptual learning modulates decision network in the human brain: The evidence
from psychophysics, modeling, and functional magnetic resonance imaging. *J. Vis.* **18**, 9 (2018).

59.    Edden, R. A. E., Muthukumaraswamy, S. D., Freeman, T. C. A. & Singh, K. D. Orientation
Discrimination Performance Is Predicted by GABA Concentration and Gamma Oscillation Frequency in
Human Primary Visual Cortex. *The Journal of Neuroscience* **29**, 15721–15726 (2009).

60.    Poort, J. *et al.* Learning Enhances Sensory and Multiple Non-sensory Representations in Primary
Visual Cortex. *Neuron* **86**, 1478–1490 (2015).

61.    Wal, A., Klein, F. J., Born, G., Busse, L. & Katzner, S. Evaluating Visual Cues Modulates Their
Representation in Mouse Visual and Cingulate Cortex. *The Journal of Neuroscience* **41**, 3531–3544
(2021).

62.    Sale, A. *et al.* Visual perceptual learning induces long-term potentiation in the visual cortex.
*Neuroscience* **172**, 219–225 (2011).

63.    Yoo, S. A., Martinez-Trujillo, J. C., Treue, S., Tsotsos, J. K. & Fallah, M. Attention to visual motion
suppresses neuronal and behavioral sensitivity in nearby feature space. *BMC Biol.* **20**, 1–19 (2022).

64.    Edden, R. A. E., Muthukumaraswamy, S. D., Freeman, T. C. A. & Singh, K. D. Orientation
discrimination performance is predicted by GABA concentration and gamma oscillation frequency in
human primary visual cortex. *Journal of Neuroscience* **29**, 15721–15726 (2009).

65.    Simpson, E. H. *et al.* Lights, fiber, action! A primer on in vivo fiber photometry. *Neuron* **0**, (2023).

66.    Xie, K. *et al.* Portable wireless electrocorticography system with a flexible microelectrodes array for
epilepsy treatment. *Sci. Rep.* **7**, 1–8 (2017).

67.    Proctor, C. M. *et al.* An Electrocorticography Device with an Integrated Microfluidic Ion Pump for
Simultaneous Neural Recording and Electrophoretic Drug Delivery In Vivo. *Adv. Biosyst.* **3**, e1800270–
e1800270 (2018).

68.    Barbera, D., Priebe, N. J. & Glickfeld, L. L. Feedforward mechanisms of cross-orientation interactions
in mouse V1. *Neuron* **110**, 297-311.e4 (2022).

69.    Bastos, A. M. *et al.* Visual areas exert feedforward and feedback influences through distinct frequency
channels. *Neuron* **85**, 390–401 (2015).

70.    Kok, P., Bains, L. J., Van Mourik, T., Norris, D. G. & De Lange, F. P. Selective Activation of the Deep
Layers of the Human Primary Visual Cortex by Top-Down Feedback. *Current Biology* **26**, 371–376
(2016).

71.    Huh, C. Y. L., Peach, J. P., Bennett, C., Vega, R. M. & Hestrin, S. Feature-Specific Organization of
Feedback Pathways in Mouse Visual Cortex. *Current Biology* **28**, 114-120.e5 (2018).

72.    Schäfer, R., Vasilaki, E. & Senn, W. Perceptual Learning via Modification of Cortical Top-Down Signals.
*PLoS Comput. Biol.* **3**, e165 (2007).

73. Zhang, S. *et al.* Selective attention. Long-range and local circuits for top-down modulation of visual
cortex processing. *Science (1979).* **345**, 660–665 (2014).
74. Sharpe, M. J. & Killcross, S. The Prelimbic Cortex Contributes to the Down-Regulation of Attention
Toward Redundant Cues. *Cerebral Cortex* **24**, 1066–1074 (2014).
75. Auger, M. L., Meccia, J. & Floresco, S. B. Regulation of sustained attention, false alarm responding
and implementation of conditional rules by prefrontal GABAA transmission: comparison with NMDA
transmission. *Psychopharmacology (Berl).* **234**, 2777–2792 (2017).
76. Wöstmann, M. *et al.* Ten simple rules to study distractor suppression. *Prog. Neurobiol.* **213**, 102269
(2022).
77. Meyerhoff, H. S., Papenmeier, F. & Huff, M. Studying visual attention using the multiple object tracking
paradigm: A tutorial review. *Atten. Percept. Psychophys.* **79**, 1255–1274 (2017).
78. Frangou, P. *et al.* Learning to optimize perceptual decisions through suppressive interactions in the
human brain. *Nat. Commun.* **10**, 1–12 (2019).
79. Ostlund, S. B. & Balleine, B. W. Lesions of Medial Prefrontal Cortex Disrupt the Acquisition But Not the
Expression of Goal-Directed Learning. *The Journal of Neuroscience* **25**, 7763–7770 (2005).
80. Niedringhaus, M. & West, E. A. Prelimbic cortex neural encoding dynamically tracks expected outcome
value. *Physiol. Behav.* **256**, 113938 (2022).
81. Gourley, S. L., Zimmermann, K. S., Allen, A. G. & Taylor, J. R. The Medial Orbitofrontal Cortex
Regulates Sensitivity to Outcome Value. *The Journal of Neuroscience* **36**, 4600–4613 (2016).
82. Marquardt, K., Sigdel, R. & Brigman, J. L. Touch-screen visual reversal learning is mediated by value
encoding and signal propagation in the orbitofrontal cortex. *Neurobiol. Learn. Mem.* **139**, 179–188
(2017).
83. Sałat, K. *et al.* The effect of GABA transporter 1 (GAT1) inhibitor, tiagabine, on scopolamine-induced
memory impairments in mice. *Pharmacological Reports* **67**, 1155–1162 (2015).
84. Stoppel, L. J. *et al.* R-Baclofen Reverses Cognitive Deficits and Improves Social Interactions in Two
Lines of 16p11.2 Deletion Mice. *Neuropsychopharmacology* **43**, 513–524 (2018).
85. Koh, W., Kwak, H., Cheong, E. & Lee, C. J. GABA tone regulation and its cognitive functions in the
brain. *Nat Rev Neurosci* **24**, 523–39 (2023).
86. Farrant, M. & Nusser, Z. Variations on an inhibitory theme: phasic and tonic activation of GABA(A)
receptors. *Nat. Rev. Neurosci.* **6**, 215–29 (2005).
87. Perez-Sanchez, J. *et al.* α5GABAA Receptors Mediate Tonic Inhibition in the Spinal Cord Dorsal Horn
and Contribute to the Resolution Of Hyperalgesia. *J. Neurosci. Res.* **95**, 1307–1318 (2017).
88. Pavlov, I., Savtchenko, L. P., Kullmann, D. M., Semyanov, A. & Walker, M. C. Outwardly rectifying
tonically active GABAA receptors in pyramidal cells modulate neuronal offset, not gain. *J. Neurosci.* **29**,
15341–15350 (2009).
89. Brickley, S. G. & Mody, I. Extrasynaptic GABA A Receptors: Their Function in the CNS and Implications
for Disease. *Neuron* **73**, 23–34 (2012).
90. Bernardo, A. M. *et al.* Procognitive and neurotrophic benefits of α5-GABA-A receptor positive allosteric
modulation in a β-amyloid deposition mouse model of Alzheimer's disease pathology. *Neurobiol. Aging*
**147**, 49–59 (2025).
91. Koh, M. T., Rosenzweig-Lipson, S. & Gallagher, M. Selective GABAA α5 positive allosteric modulators
improve cognitive function in aged rats with memory impairment. *Neuropharmacology* **64**, 145–152
(2013).
92. Pfeffer, C. K., Xue, M., He, M., Huang, Z. J. & Scanziani, M. Inhibition of inhibition in visual cortex: the
logic of connections between molecularly distinct interneurons. *Nature Neurosci.* **16**, 1068–1076
(2013).
93. Cao, J.-W., Guan, W., Yu, Y.-C. & Fu, Y. Synaptic Transmission from Somatostatin-expressing
Interneurons to Excitatory Neurons Mediated by α5-subunit-containing GABAA Receptors in the
Developing Visual Cortex. *Neuroscience* **449**, 147–156 (2020).
94. Huang, Y. Feature selective inhibitory mechanisms enable expectation suppression in cortical
microcircuits. *Scientific Reports 2025 15:1* **15**, 44597- (2025).

95.   Fee, C. *et al.* Behavioral Deficits Induced by Somatostatin-Positive GABA Neuron Silencing Are
Rescued by Alpha 5 GABA-A Receptor Potentiation. *International Journal of*
*Neuropsychopharmacology* **24**, 505–518 (2021).
96.   Schulz, J. M., Knoflach, F., Hernandez, M.-C. & Bischofberger, J. Dendrite-targeting interneurons
control synaptic NMDA-receptor activation via nonlinear α5-GABAA receptors. *Nat. Commun.* **9**, 3576
(2018).
97.   Znamenskiy, P. *et al.* Functional specificity of recurrent inhibition in visual cortex. *Neuron* **112**, 991-
1000.e8 (2024).
98.   Jia, K. *et al.* Recurrent inhibition refines mental templates to optimize perceptual decisions. *Sci. Adv.*
**10**, 31 (2024).
99.   Kutash, L. A. *et al.* Sustained attention performance deficits in the three-choice serial reaction time task
in male and female rats after experimental brain trauma. *Brain Res.* **1808**, 148336 (2023).
803
804