Apical Localization of RNA Polymerases Modulate Transcription Dynamics and Supercoiling Domains Revealed by Cryo-ET

Meng Zhang[1,2,7]*, Cristhian Cañari-Chumpitaz[1,3,4,7]*, Jianfang Liu[2], Bibiana Onoa[1,5], Sinead de Cleir[6], Enze Cheng[8], Katherinne I. Requejo[1,7], Carlos Bustamante[1,6,7,8,9,10,11,12, †]

[1] California Institute for Quantitative Biosciences, University of California, Berkeley, USA
[2] The Molecular Foundry, Lawrence Berkeley National Laboratory, Berkeley, USA
[3] Department of Chemistry, University of California Berkeley, Berkeley, CA, USA
[4] Department of Biology, Stanford University, Stanford, CA USA
[5] Innovative Genomics Institute, University of California Berkeley, Berkeley, CA, USA
[6] Department of Molecular and Cell Biology, University of California, Berkeley, USA
[7] Howard Hughes Medical Institute, University of California, Berkeley, USA
[8] Department of Physics, University of California, Berkeley, USA
[9] Department of Chemistry, University of California, Berkeley, USA
[10] Molecular Biophysics and Integrative Bioimaging Division, Lawrence Berkeley National Laboratory, USA
[11] Kavli Energy Nanoscience Institute, University of California, Berkeley, USA
[12] Jason Choy laboratory of single molecule biophysics, University of California, Berkeley, USA
* These authors contributed equally
† Correspondence should be addressed to: C. B. (carlosb@berkeley.edu)

SUMMARY

Protein interactions with canonical B-form DNA are well-characterized, yet the effect on these interactions of torsionally constrained DNA—ubiquitous in cells—remains underexplored. Using cryo-electron tomography (cryo-ET), we 3D-reconstructed entire negatively supercoiled DNA substrates bound to active RNA polymerase (RNAP), revealing diverse DNA supercoiling conformations and their interplay with transcription. RNAP preferentially localizes at plectoneme apices in a swiveled, pause-prone state. RNAP, along with other DNA-melting proteins like dCas9, can act as torsional roadblocks that segregate "twin-supercoiling domains" during active transcription, independent of external DNA/RNAP tethering. Co-transcribing RNAPs further intensify this domain separation: tandem oriented RNAPs relieve negative supercoiling more effectively than opposing ones, promote greater RNAP accumulation and enhanced elongation, both in vitro and in vivo. Topoisomerase I relieves torsional stress and facilitates RNAP escape from apical stalls, thereby supporting apical transcription regulation. Together, these findings support a load-and-release mechanism at plectoneme apices that may underlie supercoiling-dependent transcriptional bursting.

INTRODUCTION

The influence of DNA supercoiling on both prokaryotic and eukaryotic DNA transactions[1]—including transcription, replication, and chromatin segregation,[2,3] underscores the role played by deviations from canonical B-form DNA in cellular metabolism. In particular, prokaryotic genomes, inherently accumulate negative torsional stress with an average supercoiling density of ~ –0.06. Accordingly, the interplay between DNA supercoiling and the activities of DNA-binding proteins, such as the collective transcriptional behavior of RNA polymerases (RNAPs) on supercoiled templates, or the torsional regulation by DNA gyrase and topoisomerase, remain an area of active research.[4-8]

The need to visualize DNA supercoiling dynamics has spurred the development of diverse biophysical approaches. In vitro single-molecule fluorescence microscopy enables real-time tracking of DNA plectoneme formation.[9] Atomic Force Microscopy (AFM) has further allowed the detection of kinks generated in supercoiled DNA.[10] However, these approaches typically require partial confinement or surface immobilization of DNA. Cryo-electron tomography (cryo-ET) has revisited this topic using small minicircles in a more native state,[11-13] but the ~300 bp minicircles were not of sufficient length to capture the complexity of the plectoneme structures.[14] In structural studies, DNA transcription has primarily focused on elucidating RNA polymerase (RNAP) states and cofactor-mediated regulation on linear templates,[15-19] while the supercoiling aspect of the template is often overlooked. The "twin-supercoiling domain model",[20] that describes overwinding (positive supercoils) ahead of the RNAP and transcription-induced DNA unwinding (negative supercoils) at its wake, still lacks detailed 3D structural characterization despite strong biochemical support.[21-24] Consequently, questions persist such as: What are the structures naturally adopted by supercoiled DNA in 3D? What is the spatial relationship between a transcribing RNAP and its supercoiled substrate? And how do supercoiling and transcription mutually affect each other?

In this cryo-ET study, we achieved direct 3D visualization of individual plasmid particles (~2 kbp) under their naturally occurring state of negative supercoiling, allowing precise quantification of their 3D conformational dynamics. Using this plasmid, we stalled and initiated *E.coli* RNAP to systematically investigate the mutual influence between supercoiling and transcription. We determined the orientation of RNAPs on their templates and discovered that their preferred apical binding on negatively supercoiled DNA plectonemes persists during active transcription. RNAPs were found swiveled in this apical configuration, functionally facilitating initiation but hindering elongation—a previously unrecognized mode. Interestingly, we discover that this apical binding also applies to dCas9, turning it into a "soft" torsional barrier that hinders free DNA rotation. The simultaneous apical binding of dCas9 and RNAP on opposite apices of a plasmid leads to the first structural visualization of supercoiling-organized transcription domains, inducing non-plectonemic DNA loops and promoting multi-RNAP slow co-transcriptional events. This novel apical-torsional regulatory mechanism provides a direct structural basis for understanding and experimentally probing how co-transcribing RNAPs impose mutual torsional constraints under tandem and opposing gene contexts, yielding contrasting in vivo transcriptional outcomes. Additionally,

introducing *E.coli* Topoisomerase I (TopI) in the presence of RNAP only partially releases the torsion within the plasmid, which, however, is sufficient to disrupt RNAP's apical positioning and increase transcription elongation activity. Therefore, we propose an 'on-off' switch model of apical constraint that plays a critical role in linking torsional stress to transcription regulation.

RESULTS

Cryo-ET per-particle analysis enables full 3D DNA tracing of negatively supercoiled plasmids

To explore RNAP's interactions with its naturally occurring substrate, we first examined the structure and dynamics of a ~2 kbp modified pUC19 -T7A1U circular plasmid (**Figure 1A)** isolated from early stationary-phase *E. coli* cells. 2D gel analysis revealed that the two DNA strands of the plasmid loop over each other 15 times fewer (ΔLk= −15) compared to their relaxed state (Lk_0 =186), presenting a physiologically relevant supercoiling density ($\sigma=\Delta Lk/Lk_0$) of −0.08 (**Figure 1B**). The size of pUC19 permits multiple plectoneme formation, as observed by AFM, leading to significant conformational deviation among ΔLk variants. (**Figure 1C**). To capture the native topological diversity of DNA free from surface constraints, we leveraged recent advances in cryo-electron tomography (cryo-ET)[25-27] to resolve the 3D conformation of large plasmid molecules.

We initiated cryo-EM imaging under low-salt conditions (5 mM KCl) to capture loosely coiled plasmids,[14] revealing predominantly long plectonemes (**Figure 1D, orange arrows**) and occasional sharp kinks (**Figure 1D, red circles**) resembling those seen by high-resolution AFM.[10] Following a deep learning-based segmentation,[28] local "per-particle" 3D reconstruction[26], and IsoNet missing wedge correction[29] (**Video S1 and Supplementary Data 1**), we were able to obtain the non-averaged 3D maps of individual plasmids (**Figure 1E**) at resolution of ~30-40 Å (**Supplementary Data, Particle gallery #006**). Using a DNA detection algorithm with manual curation (**Supplementary Data 2**), we traced full 3D plasmid contours, enabling precise supercoiling quantification. For the example in **Figure 1E**, spatial quantification[30] of right-handed crossings in 3D gave a writhe (Wr) of −6.2. Accurate structural analysis is difficult to achieve from 2D projections alone due to projection artifacts—e.g., DNA crossing number varies (**Figure 1F, orange arrows**), smooth curves may appear as sharp kinks, and true bends can be obscured in certain views. (**Figure 1G, panels I-II)**. To overcome these limitations, we measured 3D curvature (reciprocal of osculating sphere radius; **Figure 1H**) and inter-helix spacing in plectonemes (**Figure 1G, panel III**). The curvature distribution showed two peaks (**Figure 1I, red arrows**), corresponding to plectoneme apices.

We then compared plasmids under low salt (5 mM K⁺; **Figure S1A**) and high salt (40 mM K⁺, 5 mM Mg²⁺, near-physiological ionic strength; **Figure S1B**) conditions, (see full dataset in **Supplementary Data Particle Gallery**). Under higher salt, plasmids exhibited increased elongation, indicated by a larger radius of gyration (Rg), and greater DNA intertwining with mean Wr shifting from −5.7 to −10.2; *P* <0.0001 (**Figure S1C-D**). This result aligns with theoretical predictions that electrostatic screening favors DNA

crossing,[14] promoting the conversion of negative supercoiling into writhe at the expense
of twist (ΔTw). As a result, plectoneme width decreased with high salt from 13.2 to
8.4 nm; *P* <0.001 (**Figure S1E**), accompanied by increased mean curvature,
particularly at distal apices (from 0.19 to 0.23 nm⁻¹; *P* <0.001, **Figure S1F-G**). Given
that some proteins recognize highly curved DNA region via indirect readout[31], we
assessed apex number distributions under both conditions and observed comparable
results: ~60% of plasmids remaining unbranched (two apices; **Figure S1H**). Clearly,
generation of new apices is energetically unfavorable. These analyses provide a
structural framework for understanding supercoiling dynamics (see below).
143
RNAP apical binding favors promoter escape and affects the configuration of the
negatively supercoiled template
An early study in the 1990s revealed unexpected apical localization of RNAP on
supercoiled DNA,[32] a finding recently revisited by single-molecule experiments.[9] To gain
structural insights, we incubated RNAP with pUC19 -T7A1U plasmids (bearing a single
T7A1-driven gene; **Figure 2A, top circle**) at a 3:1 ratio under high-salt conditions and
obtained stalled transcription elongation complexes (sTECs) via UTP starvation, halting
RNAP 19 nt downstream of the transcription start site (TSS). 3D reconstruction revealed
that ~90% of RNAPs localized at plectoneme apices (**Figure 2A, lower panels**). Sub-
tomogram averaging (STA) resolved RNAP orientation at the apex.[28] Averaging all
RNAP particles (bound and unbound) yielded a 14 Å map with blurred DNA entry/exit
sites (**Figure S2A-B, top**), whereas averaging only plasmid-bound RNAPs (17 Å
resolution) revealed a DNA density protrusion indicating the downstream direction
(**Figure S2A, bottom, black arrow**). This feature is absent in RNAP structures resolved
with linear DNA templates (4YLN, 6CA0, 6JBQ, 6N60,[33-36] which show only upstream
DNA bound to the σ factor (**Figure S2C**). Mapping STA-determined RNAP orientations
onto their respective plasmid models (**Figure 2A, top right**) recovered DNA information
otherwise averaged out. Superimposing all RNAPs at apical regions highlighted DNA's
conformational flexibility (**Figure 2B**), while superimposing DNA at the same region
revealed relatively restricted RNAP apical localization (**Figure 2C**).
Negative DNA supercoiling is believed to facilitate transcription initiation by
enhancing the kinetics of open complex formation through DNA unwinding.[37,38] However,
its impact on other initiation events, such as promoter search and escape, remains
unclear. Interestingly, bulk transcription assays on nicked and negatively supercoiled (–
sc) templates revealed more efficient promoter escape on –sc DNA, evidenced by a
reduced rate of abortive initiation before RNAP reached the 19 nt stall site (**Figure 2D**).
Structural superposition of the apically stalled TECs (sTEC) in –sc plasmids onto
transcription initiation complexes (TICs) assembled on linear DNA (such as 6CA0)
revealed a significant deviation in the consensus upstream DNA trajectory—
approximately 170° in bending angle in the former (**Figure S2C**). This substantial
reorientation of the upstream DNA in –sc complexes would disrupt most of DNA-σ-
factor interaction, if present (**Figure 2E**). As weakening of DNA-σ contacts is a
necessary step for RNAP release from the promoter and transition into elongation[39], this
apical configuration on –sc DNA provides a possible explanation for the increased
promotor clearance observed in the bulk assay (**Figure 2E**).

Having characterized RNAP's apical binding, we asked whether this feature is
unique to RNAP or shared by other DNA-binding proteins. With the same imaging setup,
we replaced RNAP with catalytically inactive dCas9 guided by an sgRNA targeting the
plasmid's transcription termination site (TTS) (**Figure 2F, top circle**). We chose dCas9
for its RNAP-like DNA binding (i.e. forming a bubble via DNA-RNA hybrid) and its
visibility (~10 nm) in cryo-ET. Notably, dCas9 also exhibited apical binding on –sc
plasmids (**Figure 2F, bottom**) with STA analysis revealing the consensus entry-exit
DNA orientation **(Figure S2D, arrows**; 17Å resolution, **Figure S2E**). While both proteins
share apical binding, quantitative 3D analysis enables the identification of structural
differences between dCas9- and RNAP-plasmid complexes (P.Cas vs. sTEC), with the
former displaying more elongated, tightly wound plectonemes, as indicated by
increased Rg and a more negative Wr (–9.5 to –11.8; P <0.0001, **Figure S2F, top**).
Enlarged distal loops were also observed at dCas9-bound apices, showing reduced
curvature (from 0.25 to 0.22 nm^{-1}; P <0.05) and increased DNA spacing (from 7.7 to
10.3 nm; P <0.05) compared to RNAP (**Figure S2F, bottom**). Mapping dCas9
orientation highlighted these distinctions (**Figure 2G-H**).
Why do two DNA-binding proteins with similar mechanisms reshape plectonemes
differently? Structural comparison of RNAP[40] and dCas9[41] binding pockets revealed that,
despite its larger size (15 vs. 10 nm), RNAP features a more acute binding pocket with
a sharper curvature than dCas9 (**Figure 2I**), favoring DNA melting over negative writhe.
Coarse-grained molecular dynamics (CGMD) simulations using oxDNA[42] further support
this conclusion, showing that tighter apex bending with strand separations of 4.8 and
6.4 nm (**Figure S3A**) yield sharper apices and reduced negative writhe compared to
looser apex bending (8 nm separation) (**Figure S3B, left and middle**). A positive
correlation (r =0.51) was observed between apex curvature and writhe number, based
on data pooled from all trajectories (**Figure S3B, right**). Taken together, these findings
suggest that apex localization is not unique to RNAP, but reflects a general structural
adaptation of supercoiled DNA to the geometry of bound proteins that minimizes the
energy of the overall configuration. Significantly, we found that DNA sequence at the
promoter and dCas9 binding site played a minor effect on proteins apical localization[43]
(**Figure S3C**).
210
Transcription induces branched supercoiling while preserving RNAP's apical
localization
RNAP's translocation along the DNA helix necessitates rotational movement by either
the DNA or RNAP to keep the DNA template in register within the RNAP active site.
Simultaneous rotational constraints on both RNAP and DNA, such as RNAP tethering to
cellular structures[44,45] and DNA confinement within dense genome architecture,[46-48] can
lead to the formation of twin supercoiling domains.[20] However, such external constraints
might not always be present in various cellular contexts. It is therefore intriguing to
investigate whether intrinsic constraints in TECs resulting from RNAP's apical
localization and a negative supercoiling background[48-50] can sustain transcription and
contribute to topological-twin-domain delineation.
To investigate the above scenario, we added the full set of NTPs (100 µM each)
to the stalled TECs (sTECs) and imaged the sample 10 minutes after resuming

transcription. Remarkably, the 3D reconstruction reveals the continued apical positioning of RNAP in TECs (**Figure 3A-B, purple arrows**), with nascent RNA density intermittently observed around RNAP in both 3D maps (**Figure 3A, right**) and 2D slices (**Figure 3C top, red contours**), but absent in sTECs (**Figure 3C bottom**). However, the limited resolution of cryo-ET, along with conformational heterogeneity of RNA[51] prevented the tracing of complete RNA transcripts, permitting only the mapping of the RNA protrusion direction relative to RNAP (**Figure S3D, arrows**). Quantitative analysis revealed that, during active transcription, plasmids predominantly adopt a multi-branched configuration for the first time (>78% of structures, **Figure 3D**), resulting in a population with reduced spatial extension and smaller Rg (**Figure S3E, top, red arrow**). Additionally, we observed occasional emergence of blunt apices in TECs (hollow purple arrows in both **Figure 3A and Figure S3E**), a feature rarely seen in sTECs (**Figure 2A**). While the overall plasmid writhe was similar between TECs and sTECs (–9.5 vs. –9.8) (**Figure S3E, top**), segmentation of individual branches (**Figure S3F, I-III**) followed by analysis of writhe density (branch Wr/branch length) showed that TECs exhibit a broader distribution with a slightly increases (–0.13 vs. –0.15 nm⁻¹; P <0.05; **Figure S3F, bottom**). These observations suggest that intrinsic –sc constraints, coupled with active DNA translocation relative to RNAP, are sufficient to generate a mild degree of torsional imbalance in the plasmid, promoting the formation of a new supercoiling domains (plectoneme branching). Branching appears to preferentially occur in the region of the plasmid proximal to RNAP (**Figure S3G**), consistent with its role as the source of torsional stress.

DNA torsional stress resulting from transcription of torsionally constrained templates is known to impede transcription elongation, causing RNAP stalling and increased backtracking as shown in single-molecule assays.[21] Given the persistent apical localization of RNAP observed in our experiments during active transcription elongation, we wondered if transcription rate was also affected in these conditions. Using radiolabeled α-^{32}P-ATP, single-round in vitro transcription assays on –sc DNA displayed significantly slower elongation kinetics (**Figure 3E**), as well as increased pause frequency and duration (**Figure 3F**), than those performed on nicked DNA. We note that, in our bulk transcription assay, the topological constraint arises from the natural steady-state degree of negative supercoiling of the plasmids extracted from the cell. The slower elongation kinetics observed supports the notion that the apical localization of RNAP hinders its rotation around DNA, and that the inefficient rotation of the larger DNA molecule becomes rate-limiting to transcription.[32]

Alternatively, the topological stress under which the apical RNAP operates could induce a conformational change in the enzyme which could, in turn, affect its dynamics. To test this hypothesis, we performed single-particle analysis (SPA) of TECs on –sc templates. 3D classification yielded three classes, of which Class 1 and Class 2 (TEC1–SC and TEC2–SC), comprise 34% and 40% of particles, respectively (**Figure S4A-C**) attaining in both cases 2.9 Å resolution (**Figure S4D-E**) and clearly showing melted DNA and RNA transcripts at the active site. Minor Class 3 (26%) exhibited faint DNA density and was excluded from further analysis. Close inspection revealed that TEC1–SC and TEC2–SC are both in a post-translocated state, evidenced by an unfolded trigger helix and absence of the incoming NTP (**Figure S4F**). Using the NTP-bound

RNAP structure (PDB: 6RH3) as a reference, we measured the swivel angle following
Kang et al.[52] and found that TEC1–SC exhibits significant swiveling (~2.7°; **Figure 3G**),
comparable to the swivel state displayed by the His-elemental paused elongation
complex (His-ePEC, **Figure 3H**), that results from a pause stabilized by an RNA hairpin
formation. Interestingly, we notice that the TEC1–SC DNA orientation (**Figure 3I,**
dashed line) and the DNA confinement geometry in apically located RNAP (**Figure 3I,**
black arrow) align with the orientation of the swivel module (**Figure 3I, red arrow**).
Because the downstream DNA contacts the swivel module, these observations suggest
that bending of the constrained apical DNA drives the swivel conformation, thereby
slowing transcription elongation.
Building on these findings, we propose a model for how transcription takes place
in intrinsically –sc DNA (**Figure 3J**): The persistent apical positioning of RNAP requires
DNA to rotate during transcription as it is thread through the enzyme (**Figure 3J, I**).
However, the propagation of DNA rotation throughout the entire plasmid structure is
slow,[53,54] so that transcription induces transient (+) torsional stress ahead of the enzyme
and (–) torsional stress at its wake (**Figure 3J, II**). In topologically constrained DNA, the
(+) torsional stress is rapidly absorbed by the –sc background, and therefore, the
cancellation of (+) and (–) supercoils through diffusion across the plasmid does not
occur (**Figure 3J, III**). Meanwhile, since high salt conditions favor partitioning of ΔLk into
writhe over twist (**Figure S1D**), the extra (–) torsional stress generated at the wake of
the polymerase promotes the formation of a new negative plectoneme branch.
290
Dual apical protein binding promotes the formation of topological domains
The results above highlighted RNAP's role in generating torsional stress during non-
equilibrium transcription of supercoiled DNA. Torsional stress during transcription can
also accumulate due to the presence of torsional roadblocks, such as DNA-binding
proteins and transcription factors,[55,56] leading to the formation of topological domains
that can affect RNAP molecules distant kilobases away.[57] Nonetheless, the mechanism
through which small DNA-binding proteins impose rotational constraints and favor the
generation of topological domains remains elusive.
To gain insight into this process, we employed dCas9 again but in this case for
its capability of delaying DNA rotation, as reported in magnetic tweezers assays.[58] To
confirm that dCas9 can serve as a programmable torsional roadblock, we performed
DNA relaxation experiments using topoisomerase I (TopI) in the absence and presence
of dCas9 bound to the transcriptional termination site (TTS) (**Figure 4A, top circle**).
TopI relaxes torsion in –sc DNA by rotating the helix until equilibrium (ΔLk $=$ 0) is
achieved. Our results demonstrate that dCas9 delays the relaxation kinetics of pUC19
plasmids, and topoisomers with ΔLk from –3 to –1 still persist after a 30-minute reaction
(**Figure S5A**). We repeated the experiment using human Topoisomerase IB (hTopIB),
which relaxes supercoiling via a direct DNA rotation mechanism rather than strand
passage (as in TopI), and similarly observed that dCas9 delays DNA relaxation (**Figure**
S5B). Because dCas9 primarily delays, rather than halts, –sc relaxation, we term it a
"soft" torsional block. To understand this mechanism, we performed CGMD simulations
to investigate how a –sc plasmid relaxes after nicking, with or without an apical
constraint. Results reveal that imposing an apical constraint on one of the two apices

slows supercoiling torsional relaxation dynamics when the nick is positioned between the apices (**Video S2 and Figure S5C, Model II-III vs. I**). However, when the nick is placed directly on the apex, relaxation becomes fastest, even with the opposite apex constrained (**Figure S5C, Model IV**). Based on the observed relaxation speeds (IV > I > II and III), and given that model IV is the only scenario where torsional propagation encounters no apices (with the nick created at the very end), it suggests that the apex, particularly the confined apex (mimicking the effect of dCas9 binding), hinders free DNA rotation.

To experimentally test the role of torsional roadblocks on transcription, we first initiated transcription on pUC19 plasmids with dCas9 bound at the TTS (**Figure 4A, top circle**). Imaging of stalled TECs (sTEC-Cas) harboring a ~20 nt transcript revealed that RNAP and dCas9 predominantly localize to opposite plectoneme ends (**Figure 4A top left, and bottom columns 1-3**). We note that on unbranched plasmids, their apical localization is mutually reinforcing, as both favor and induce bending at their target sites (**Figure 2A,F**). Even under this mild transcription and stalling scenario, sTEC-Cas exhibited noticeably less negative writhe domains compared to the dCas9-bound-only plasmid (P.Cas), resembling instead those of active TECs (**Figure S5D, red arrow**). Notably, mild transcription conditions also led to a significant increase in non-apical binding cases (**Figure 4A, blue label** and **Figure S5E**) for both RNAP and dCas9 (from 18% to 41% and from 17% to 47%, respectively, **Figure 4B**). Given that the increased off-apex dCas9 ratio was not due to additional dCas9 binding events (both 6% for P.Cas and sTEC-Cas; **Figure 4C**), it suggests that transcription-induced accumulation of (+) torsional stress between the front of RNAP and dCas9, and (−) stress between dCas9 and the back of RNAP, promotes partial escape of the dCas9 from the apices as the system seeks to minimize its energy.

Torsional block enhances cooperative RNAP transcription

As shown above, mild (+) and (−) torsional stresses differentially accumulate in each half of the plasmid, delineated by RNAP and dCas9, resulting in more frequent off-apical dCas9 events, suggesting the "soft" nature of dCas9 when acting as a torsional block. Accordingly, we anticipated increased off-apex localization of dCas9 after 10 minutes of active RNAP transcription with the full set of NTPs at 100 µM (TEC-Cas; **Figure 4D**). Indeed, the off-apex ratios of both RNAP and dCas9 increased, with dCas9 reaching 77% (**Figure 4B**). Given a −sc background ($\Delta Lk = −15$), the "twin-domain" model should result in the DNA segment in front of the RNAP to be less −sc and the domain at its back more so. Consistent with this expectation, we observed the coexistence of large, non-plectonemic DNA loops and tightly wound DNA regions within the same TEC-Cas particle (**Figure 4D, top left**). Quantitative analysis confirmed that, although overall plasmid morphology remained similar (**Figure S6A, top**), TEC-Cas exhibited a broader writhe density distribution across plectonemic branches, with two peaks above and one slightly below the mean of the plasmid-only control (**Figure S6B, red arrows**). This shift toward less negative writhe was accompanied by blunter apices, characterized by reduced apex curvature (from 2.5 to 2.3 nm^{-1}; $P < 0.05$) and increased apical-DNA spacing (from 8.5 to 11.1 nm; $P < 0.001$, **Figure S6A, bottom**). Notably, less-negative domains were more prominent, which seems contrary to the expected

symmetry of (+) and (−) torsion generated by RNAP (**Figure 4E, I**). This asymmetry can be rationalized by the marked increase in multi-RNAP binding (from 29% to 50%; **Figure 4C**) observed in TEC-Cas. Accordingly, as each bound RNAP unwinds ~10 bp of DNA, they absorb excess negative supercoiling (**Figure 4E, II**). Moreover, the increase in the proportion of multi-branch form plasmids in TEC-Cas (from 32% to 50%, **Figure 4F**), providing an additional mechanism to buffer excess negative supercoiling (**Figure 4F, III**).

Since structural studies provide only static snapshots of domain separation, we employed CGMD to better visualize how torsional domains dynamically emerge as a consequence of apical constraint in the absence of external tethering. We treated the domains as harmonic (linear torsional springs)[59] to mimic the (+) and (−) torsion generated by RNAP. The simulations compare a free versus constrained apex in managing torsional stress (**Video S3**). Notably, the constrained apex successfully recapitulated the formation of large loops and branching structures observed experimentally.

Leveraging dCas9's gRNA-directed DNA binding as a fiducial marker, we tentatively mapped RNAP positions and orientations on the plasmid construct (**Figure S6C**). Some RNAPs were found outside the transcriptional region or oriented upstream of the TSS (brown and blue arrows), which we interpret as non-specific binding enhanced by transcription-induced −sc domain. In contrast, most RNAPs were located within the transcriptional region and oriented downstream (pink arrows), with some plasmids displaying multiple transcribing RNAPs. The direct visualization of two RNAPs on the same template, both exhibiting nascent RNA protrusions (**Figure 4D, green labels**), supports this interpretation. These findings suggest that dCas9-induced topological imbalance in TECs promotes cooperative RNAP transcription, potentially enhancing transcriptional kinetics. To corroborate this hypothesis and link structure to function, we performed a bulk transcription assay in the presence of dCas9 (**Figure S6D**). The results show that while pause frequency remains unchanged, pause escape is faster in the TEC-Cas—particularly at the 45 nt and 55 nt pause sites (**Figure 4G, blue vs. green curve**). This enhanced pause escape suggests that the accumulation of (+) torsion ahead of RNAP, induced by the dCas9 torsional block, relieves the enzyme's apical constraint on a −sc template, thereby facilitating elongation—though not to the level observed with nicked templates (**Figure 4G, orange curve**). Together, these results suggest that, in the context of −sc, twin-domain formation facilitates both RNAP initiation and elongation, with a stronger effect on the former.

TopI removes the apical constraint of RNAP facilitating DNA transcription
Although the presence of a torsional roadblock like dCas9 promotes transcription by multiple RNAPs, it could not fully replicate the elongation dynamics seen on nicked circular plasmids. Accordingly, we sought to determine if TopI[8,60,61] (**Figure 5A**) could alleviate the apical constraints on RNAP by relaxing the DNA plectoneme and thereby relieve RNAP from its pauses.

Bulk transcription assays on −sc plasmid with TopI confirmed markedly improved transcription (**Figure 5B, yellow vs. green, and Figure S6E**), exhibiting kinetics similar to those of nicked templates without torsional stress (**Figure 5B, orange**). Based on

these results, we conducted a cryo-ET imaging study of RNAP transcription in the presence of TopI (RNAP:TopI:plasmid=3:1:1). 10 minutes after the re-start of the sTEC, TEC-TopI sample revealed the presence of large RNA transcripts (**Figure 5C**). Moreover, TopI activity alone is sufficient to displace 50% of RNAPs from their apical localization, even in the absence of a torsional block (**Figure 5D, green arrows, and Figure 5E**), yielding the highest off-apex ratio across all tested conditions (**Figure 5F**), and without inducing plasmid branching (**Figure 5G**). Quantitative analysis showed that blunt apices predominated in TopI-relaxed plasmids, in contrast to unrelaxed ones, with a marked reduction in curvature (from 0.25 to 0.18 nm^{-1}; P <0.0001) and an increase in DNA spacing (from 7.3 to 15.2 nm; P <0.0001). (**Figure S6F**). We attribute this prominent apex structural change to the relaxation of the plectoneme (mean Wr =−6.4; **Figure S6F**). Analysis of RNAP orientations in TEC-TopI particles reveals highly dynamic entry–exit DNA (**Figure 5H**), in contrast to the apically constrained configuration observed in TECs (**Figure 3B**). These results suggest that not only torsional constraint, but also a high degree of –sc, is required for RNAP apical localization and the longer pauses observed during single-round transcription.

Interestingly, while TopI facilitates RNAP transcription, RNAP also modulates TopI activity. Previous bulk assays showed that TopI alone (1:1 with plasmid) rapidly relaxed supercoils (ΔLk from −15 to −1) within minutes (**Figure S5A**). However, the addition of RNAP (plasmid:RNAP:TopI = 1:3:1) notably slowed relaxation kinetics, with topoisomer bands clustering at ΔLk < −4 even after 30 minutes (**Figure 5I**). 2D gel further resolved these bands, revealing accumulation of topoisomers at ΔLk = −8 (**Figure S7A**). Given the reported coupling between RNAP and TopI[8] (**Figure 5A, dashed arrow**), which could potentially impair TopI function, we performed a control experiment with excess TopI (plasmid:RNAP:TopI = 1:3:9; **Figure S7B**). The relaxation kinetics were similar to those observed at a 1:3:1 ratio, indicating that the slower relaxation is not due to TopI inactivation upon interaction with RNAP, but rather reflects RNAP's torsional block function—similar to the dCas9-mediated TopI downregulation (**Figure S5A**). Moreover, hTopIB, known not to interact with RNAP, showed a similarly slowed relaxation pattern on RNAP-bound plasmids, further supporting the above interpretation (**Figure 5J**). Interestingly, the torsional block effect of RNAP appears to be more pronounced than that of dCas9 (dCas9-topoisomers clustered at ΔLk = −1 to −3), likely due to its larger size and its ability to induce sharper kinks when it melts the DNA duplex. Indeed, supporting the requirement of DNA melting for DNA-binding protein to act as a roadblock, use of RNAP core—which binds DNA but cannot form stable transcription bubbles—resulted in faster relaxation than the holo-enzyme and ended in fully relaxed plasmids. (**Figure S7C**). Consistently, dead EcoRI mutant (dEcoRI), which bends but does not melt DNA, failed to act as a torsional barrier and permitted rapid, full relaxation (**Figure S7D**). These results suggest that both DNA bending and duplex melting are required for a protein to function effectively as a torsional block.

Tandem promoters facilitate supercoiling domain formation and favors transcription over opposing promoters

We learned from the above experiments that the restricted rotational freedom of RNAP
at apices confers dual functions: (1) generation of torsional strain during transcription, a
prerequisite for the twin-domain model, and (2) serve as a torsional roadblock for a
second transcribing RNAP as is also observed with dCas9. This dual function parallels
the cellular scenario where multiple RNAPs co-transcribe the same genomic DNA,
enabling transcriptional coupling between neighboring genes through the formation of
topological domains.[62] To gain insight into this system, we designed dual-promoter
circular pUC19 plasmids (2.8 kb) containing two identical T7A1 promoters, each with a
U-less stalling sequence, driving transcription of AmpR and mEGFP, either in opposite
(topologically mixed convergent and divergent; **Figure 6A, left**) or tandem (i.e., same-
direction; **Figure 6B, left**) orientations. After UTP starvation–induced stalling, TECs
were imaged after 10 minutes of transcription. To facilitate visualization of these more
complex structures, we projected plasmids onto a circular layout (**Figure 6A-B,
middle**), with the rim color-coded to match the 3D molecular model (**Figure 6A-B,
right**). Arrowheads indicate the transcription direction of bound RNAPs, and circles
mark apical sites. Inner circular sectors represent individual plectonemes (e.g., I, II, and
III), color-coded by its writhe density (blue–pink–red, –0.3 to 0.1) (**Figure 6A, middle**).
Significantly, as RNAP loading increased, plasmids underwent major
reconfiguration from elongated plectonemes to expanded, globular structures
(**Figure 6C, opposing, and Figure 6D, tandem**). Under identical reaction conditions,
we observed a higher number of RNAPs in tandem TECs (T-TECs) compared to
opposing TECs (O-TECs). T-TEC particles exhibited up to 10 bound RNAPs and 9
apices, with 60% of plasmids bound by more than 3 RNAPs and 77% forming more
than 3 branches (**Figure 6E**). In contrast, multi-RNAP loading and multi-apices were
down to 16% and 35% in O-TECs, and dropped further to 0% and 4%, respectively in
single-promoter TECs, respectively. As a result, the highly branched T-TECs exhibited
reduced plectonemic extension, with a significantly smaller Rg compared to O-TECs
(64 nm vs. 79 nm; P <0.001, **Figure 6F**). Statistical analysis confirmed a positive
correlation between RNAP occupancy and plasmid branching (r = 0.61, **Figure 6G**),
suggesting that RNAP number potentially reflects the generation of domain separation.
Notably, the higher number of RNAPs was not necessarily localized at apices. In both
O-TECs and T-TECs, increased multi-RNAP binding led to elevated off-apex RNAP
localization, reaching 44% for O-TECs and 54% for T-TECs (**Figure 6H**), in contrast to
off-apex localization of single TECs (8%), but comparable to levels observed in TEC-
Cas and TEC-TopI (45% and 50%, respectively).
The overall phenomena of multi-RNAP binding and multi-branching are not
fundamentally distinct from those observed in TEC-Cas, but rather represent more
extreme cases, especially in T-TECs. This result supports the continued validity of the
TEC-Cas model (**Figure 4E**), with the RNAP version of "dCas9" reinforcing domain
separation through its dual functions. In this context, we compare O-TECs and T-TECs
in their supercoiling domain formation. Visually, as more RNAPs were loaded, a shift
toward non-plectonemic DNA loops (lower negative writhe density; pink circle segments,
Figure S7E) was more frequently observed in T-TECs than in O-TECs. This trend was
further confirmed as increased number of RNAPs led to a broader writhe density
distribution of T-TECs, displaying enriched hypo and hyper –sc domains compared to

O-TECs (**Figure 6I, red arrows**). In contrast, a subset of O-TECs exhibited plectoneme
writhe densities extending into the far-positive regime (**Figure 6I, blue arrow**).
Examination of these particles revealed, for the first time, TECs with long positive
plectonemes (**Figure 6C and S7E, red label**), supporting that opposing RNAPs,
transcribing toward each other mutually act as torsional blocks that accumulate positive
supercoiling. One T-TEC also showed short positive supercoiling (**Figure S7E**), but
likely due to DNA entanglement between plectonemes. Nevertheless, this observation
also suggests the possibility of mutual torsional interference between tandem RNAPs
due to differential translocation speeds, consistent with the on–off apex model observed
in TEC-Cas particles.
Interestingly, we observed a higher correlation between the writhe density and
RNAP number in the O-TECs than T-TECs (r =0.46 vs. r =0,14, **Figure 6I**). We interpret
it as the accumulation of torsional stress in the opposing geometry leading to higher
changes in writhe compared to the cancellation of torsional stress when RNAP
transcribes in tandem (i.e., further multi-RNAP loading has limited impact on torsional
accumulation). Furthermore, by identifying all possible RNAP pairs on the plasmid and
measuring their angular separation in polar plots, we found that opposing RNAPs were
more likely to be separated by 2.7 radians (~1.2 kbp) (**Figure 6J**), in contrast to the
uniform distribution observed for tandem RNAPs—further supporting a model of
transcriptional hindrance in opposing configurations.
To confirm the functional consequences of the observed structural differences
between tandem and opposing gene arrangements, we measured gene expression
both in vitro and in vivo. RNA molecular beacon assays[63] revealed comparable in vitro
transcript levels of AmpR and mEGFP within the same construct (**Figure 6K**), as
expected since both genes are driven by identical promoters (T7A1). However, higher
RNA levels were observed with the tandem arrangement compared to the opposing one
for both genes (**Figure 6K**). We also compared in vivo gene expression of mEGFP by
measuring fluorescence emission in live *E. coli* cells harboring either construct, grown in
M9GluCAAT medium. Consistent with the in vitro results, tandem constructs showed
nearly a ten-fold higher fluorescence than their opposing counterparts (**Figure 6L-M**).
These results are highly consistent with our cryo-ET observations: the tandem gene
arrangement promotes increased transcriptional activity, including higher RNAP loading,
increased numbers of off-apex RNAPs less prone to pausing, and lower accumulated
torsional stress due to the cancellation of neighboring RNAPs.
527
DISCUSSION
We demonstrate that cryo-ET, combined with modern image processing, enables
detailed 3D visualization of highly dynamic DNA/protein complexes previously
underexplored. By reconstructing the full DNA backbone of plasmids extracted from
cells and bearing natural –sc densities, we precisely quantify plectoneme
conformational dynamics. This marks a major advance in understanding DNA
supercoiling, previously limited to surface-level AFM,[10,64] negative-stain electron
microscopy,[65] or cryo-ET imaging of ~300 bp minicircles.[12] By directly imaging individual
~2-3 kb plasmids, we quantify writhe and show that under near-physiological ionic
strength (40 mM KCl, 5 mM MgCl$_2$), torsional stress is stored primarily as writhe (Wr ≈

−10 for ∆Lk ~ −15; **Figure S1B**, **S1D**). We find that lowering the ionic strength reduces
Wr to −5, increases DNA strand spacing of plectonemic DNA and lowers apex curvature.
Previously, such 3D conformational changes were accessible only through coarse-
grained Brownian dynamics simulations.[66,67] Our findings provide a basis for refining
theoretical models to better reflect DNA's typical topological state in cells.
Our full DNA tracing in 3D reveals that DNA-melting proteins like RNAP and
dCas9 preferentially localize at plectoneme apices when bound to −sc DNA (**Figures**
2A, 2F). We find that this apical localization, first noted by ten Heggeler-Bordier et al.[32]
and revisited by Janissen et al.,[9] is sequence-independent and persists as RNAP
translocates along DNA (**Figure 3A**). Direct observation of RNA transcripts shows that
this geometry helps segregate nascent RNA from the DNA body (**Figure 3A-B**),
potentially aiding co-transcriptional events such as RNA folding, RNA processing,
ribosome binding, while preventing RNA-DNA hybrid formation. Notably, RNAP
localizes apically only at σ ≈ −0.08 (∆Lk = −15); relaxation by TopI to σ ≈ −0.04 (∆Lk =
−8) reduces this localization (**Figure 5F**), suggesting a possible role of −sc on gene
expression.
Beyond stabilizing open complex formation,[37] we show that −sc promotes
transcription by facilitating promoter escape and reducing abortive initiation (**Figure 2D**).
The ~170° bending induced on the upstream DNA in the apically positioned TECs
disrupts the majority of σ-factor:DNA contacts, a requirement for promoter clearance,
offering an additional layer of regulation and helping rationalize the supercoiling
sensitivity of certain bacterial promoters.[68] Negative supercoiling also enhances RNAP
binding following the formation of hyper-negative domains (**Figure S6B**), likely
accelerating promoter search and triggering transcriptional cascades.
However, supercoiling can also impede transcription by promoting RNAP
pausing (**Figure 3F**), in line with single-molecule studies.[21] In contrast to those studies,
which used surface fixed DNA, our work shows that under no additional torsional barrier
other than the inherit-torsional stress of plectonemic DNA, the apical localization of
RNAP can induce a swiveled state in RNAP with increased propensity to pausing
(**Figure 3G**). While nascent RNA structures, transcription factors, and DNA sequence
are known to promote pausing by stabilizing the swiveled state, our results reveal that
DNA topology, through apical localization, can similarly regulate this conformational
transition. This supercoiling-induced pausing is reversed by TopI (**Figure 5B**) and also
by the action of co-transcribing RNAPs in tandem, which reduces apical localization
(**Figure 6H**), restores elongation, and promote a more "fluid" transcriptional mode.[24]
The twin-supercoiled domain model has long explained transcription-induced
supercoiling in prokaryotes and eukaryotes,[5,24,69,70] but direct visualization of the resulting
supercoils (**Figure 7A, left**) has remained elusive. Early models suggested that long
RNA transcripts or anchoring to cellular structures were required to prevent RNAP
rotation and allow torsional stress to build. Here, we show direct structural evidence that
rotationally constrained, apically localized RNAP—with even short (~20 nt) transcripts—
can induce new plectoneme formation. This suggests long transcripts are not necessary
to constrain RNAP, aligning with recent RNase-treatment studies.[9] A single apically
positioned RNAP can generate plectoneme branching, as observed in our experiments
and predicted by other simulation work,[71] though the latter required unrealistically high

elongation rates and forces. Incorporating our 3D model of the apical RNAP configuration may help reconcile this discrepancy. Under physiological conditions with initial negative supercoiling, positive supercoil formation was rare, contrary to textbook models. Instead, we observed writhe density changes upstream and downstream of RNAP (**Figure 7A, right**), especially when torsional barriers such as neighboring RNAPs were present. This role of RNAP and DNA-melting proteins like dCas9 may extend to similar enzymes in both domains of life. DNA-bending proteins like dEcoRI did not show this effect, but other factors such as nucleoid-associated proteins[72] or nucleosomes—which store ~1.7 DNA turns—may also act as torsional barriers.[71,73] Together, these observations define a revised twin-supercoil model that updates a canonical textbook framework under physiologically relevant, tether-free negative supercoiling.

Negative supercoiling promotes RNAP initiation but hinders elongation, supporting a pulse-like transcription model reminiscent of in vivo "transcriptional bursting".[74,75] Single-molecule studies have linked supercoiling to this behavior.[22] Based on our findings, we propose an updated model for –sc DNA (**Figure 7B, I–III**). Initially (**Figure 7B, I–III**), physiological DNA stores negative torsion that facilitates transcription initiation, promoter escape, and apical RNAP localization. In the absence of additional torsional roadblocks besides an apically located RNAP, transcription-induced supercoiling may lead to the formation of additional plectonemes (**Figure 7B, IV, RNAP alone**). In contrast, additional torsional blocks segment topological domains—reducing the negative writhe in front of the RNAP while increasing it behind, leading to enhanced RNAP binding and transcriptional burst initiation (**Figure 7B, IV +additional torsional block**). When RNAPs bind in tandem, they can cancel each other's torsional stress, promoting transcriptional synergy (**Figure 7B, V, Tandem**); however, opposing RNAPs may stall due to accumulated stress (**Figure 7B, V, Opposing**). To complete transcription, topoisomerase relieves –sc, displacing RNAPs from the apex and facilitating elongation (**Figure 7B, VI**). The requirement for topoisomerase in active transcription is supported by the direct interaction between *E. coli* TopI and RNAP[8,60] and by the observed recruitment of topoisomerase at transcription start sites.[61] The resulting relief of negative torsion reduces the likelihood of new initiation, effectively acting as an OFF switch. The cycle can restart via ATP-dependent DNA gyrase activity, which restores negative supercoiling.

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Carlos Bustamante (carlosb@berkeley.edu).

Materials availability
All unique reagents generated in this study are available from the lead contact with a completed Materials Transfer Agreement.

DATA AVAILABILITY

The cryo-ET maps of ~2 kb negatively supercoiled pUC19-T7A1U plasmids under various conditions were montaged and deposited in the Electron Microscopy Data Bank (EMDB). These include maps of plasmids under low salt conditions (EMD-47843), high salt conditions without additional proteins (EMD-47844), with stalled RNA polymerase (EMD-47847), with dCas9 (EMD-47849), with RNA polymerase during active transcription (EMD-47850), with both stalled RNA polymerase and dCas9 (EMD-47851), with transcribed RNA polymerase and dCas9 (EMD-47853), and with transcribed RNA polymerase and Topoisomerase I (EMD-47855). The cryo-ET maps of ~2.8 kb negatively supercoiled pUC19-T7A1U plasmids containing opposing promoters and tandem promoters were montaged and deposited in the EMDB under accession numbers EMD-71622 and EMD-71618, respectively. The single-particle analysis maps of TEC1–SC and TEC2–SC were deposited in the EMDB under accession numbers EMD-71675 and EMD-71676, with their corresponding atomic models available in the PDB under accession codes 9PIP and 9PIQ, respectively. The individual particle 3D reconstructed maps, models, original gel images, fluorescence images, the script calculate the TECs swivel angle, and data used to generate the statistical analysis presented in the figures have been deposited on GitHub Zenodo at (https://doi.org/10.5281/zenodo.16423113).

ACKNOWLEDGMENTS

We thank Dr. Liang Meng Wee for the expression and purification of dCas9. We thank Rodrigo Fregoso, Matthew Lo, Allison Cordova and Jimena Pavlovitch for assistance in the early stages of the project. We also thank Dr. Gang Ren for technical support and valuable feedback, and Professor Zev Bryant and Dr. Noor Sayyad for their insightful comments on the work. Data were collected at the Cal-Cryo facility, Berkeley QB3 Institute. The work at the molecular foundry, LBNL, was supported by the Office of Science, Office of Basic Energy Sciences of the United States Department of Energy (contract no. DE-AC02-05CH11231). This research was supported by grants from the US National Institutes of Health R01GM032543 (C.B). C.B. is Howard Hughes Medical Institute investigator.

AUTHOR CONTRIBUTIONS

M.Z., C.C.-C., and C.B. conceived the study and designed the research. C.C.-C. prepared the oligonucleotide and plasmid. C.C.-C. performed the bulk transcription assay. C.C.-C., S.D.C., and K.I.R. performed the relaxation assays. C.C.-C. and K.I.R. performed in vitro and in vivo measurements of gene expression for the opposing and tandem constructs. B.O. conducted the AFM imaging. M.Z. prepared the EM specimens, and M.Z. and J.L. collected the EM data. M.Z. managed the 3D reconstruction workflow. M.Z., S.D.C., and E.C. contributed to the modeling. M.Z. performed single particle cryo-EM analysis. M.Z. interpreted the data and prepared the figures. J.F and M.Z. performed the molecular dynamic simulation. M.Z. and C.C.-C. wrote the original draft, with revisions by B.O., J.L., S.D.C., K.I.R., E.C. and C.B. supervised the work.

DECLARATION OF INTERESTS

672   The authors declare no competing interests.
673
674   **MATERIALS AND METHODS**
675
676   **Oligonucleotides and RNA**
677   All DNA oligonucleotides listed below were obtained from Integrated DNA Technologies
678   (IDT). Except for PCR primers, all oligonucleotides were purified in-house using
679   denaturing urea-polyacrylamide gels (PAGE) prepared from SequaGel UreaGel 29:1
680   Concentrate (National Diagnostics).
681
Preparation of single guide RNAs (sgRNAs)
sgRNA specific oligonucleotides and scaffold oligonucleotides (CBD03 and CBD04)
were ordered from IDT and purified in-house by polyacrylamide-gel electrophoresis
(PAGE). In a 30 µL reaction, oligonucleotides (20 µL, 50 µM) were annealed and
extended by T4 DNA polymerase in CutSmart 1X buffer to produce the dsDNA template.
After heat inactivation, 10 µL of this reaction (~33.3 µM dsDNA template) were used for
a 20 µL in vitro transcription reaction using HiScribe T7 In vitro Transcription Kit (New
England Biolabs). In vitro transcription reactions were performed at 37°C for 6 hours,
after which Dnase I was added to degrade the DNA template. sgRNAs were purified by
urea PAGE. Bands corresponding to the correct RNA size were cut out as gel slices,
eluted overnight at room temperature in 2X PK buffer (200 mM Tris-HCl, pH 7.5, 25 mM
EDTA, pH 8.0, 300 mM NaCl and 2% SDS (w/v)), phenol-chloroform extracted and
precipitated with 2X volume of 200-proof 100% ethanol (Koptec). Then, samples were
air dried and suspended in UltraPure DNase/RNase-free distilled water (Invitrogen,
Thermo Fisher Scientific).
697
Cloning of Single promoter plasmid pUC19-T7A1U, pUC19-T7A1-AmpR-mEGFP-
Tandem and pUC19-T7A1-AmpR-mEGFP-Opposing constructs
Single promoter plasmid pUC19-T7A1-U was generated using around-the-horn PCR
from template plasmid pUC19 using primers (CBD01/CBD02) with extensions
containing the T7A1 promoter sequence as well mutations in the initially transcribed
region to form a U-less cassette. These primers also contained EcoRI recognition sites
for subsequent digestion and circularization by ligation. PCR products were digested
with EcoRI and DpnI and were further purified by agarose-gel extraction. Purified PCR
products were ligated using T4 DNA Ligase and transformed into DH5α cells. The
tandem construct (pUC19-T7A1-AmpR-mEGFP-Tandem) was generated using Gibson
Assembly of three PCR fragment assembly reaction: fragment 1 (PCR with template
pUC19-T7A1-U using primers CBD05/06, fragment (PCR with template pUC19-T7A1-U
using primers CBD07/08), and fragment 3 (PCR from plasmid containing mEGFP using
primers CBD09/10). Each PCR product was purified by gel-extraction. Gibson Assembly
was carried using NEB Gibson Assembly Master Mix (NEB, #E2611L) following
manufacturer instructions and then assembly reaction transformed into DH5α cells. The
opposing construct (pUC19-T7A1-AmpR-mEGFP-Opposing) was obtained using
Golden Gate Assembly from two PCR fragments: fragment 1 (PCR with template
pUC19-T7A1-AmpR-mEGFP-Tandem using primers CBD11/12) and fragment 2 (PCR

with template pUC19-T7A1-U using primers CBD13/14). The PCR fragments were treated with BsaI (NEB) and BsmbI (NEB), respectively, and then ligated using T4 DNA ligase (NEB). Given the instability of this plasmid in DH5α cells, NEB® Stable Competent *E. coli* cells were used for transformation instead. In all cases, plasmids from positive clones were extracted using QIAGEN miniprep kit and their sequences were confirmed by DNA sequencing.

For large scale production of plasmids, plasmids were retransformed into DH5α or NEB Stable competent cells, accordingly. Cells were grown in 1 L of LB media and extracted using QIAGEN Maxiprep kit following manufacturer instructions. For bulk biochemical as well as cryo-ET structural studies, supercoiled plasmid pUC19-T7A1-U was further purified using low-melting agarose (SeaPlaque Agarose, Lonza) from 1% agarose gel (1X TAE with 1X SYBR Safe DNA Staining). Under these electrophoretic conditions nicked DNA was well separated from supercoiled DNA. The band corresponding to supercoiled DNA was excised and extracted using β-Agarase (New England Biolabs), followed by phenol-chloroform extraction and ethanol precipitation. To desalt and concentrate the samples, the purified plasmids were further purified with Monarch PCR & DNA Cleanup Kit (New England Biolabs) using manufacturer instructions. The isolated purified plasmids were stored at –20 °C until further use.

Generation of different DNA topologies

To generate different DNA topologies (nicked, $\Delta Lk=0$, $\Delta Lk < 0$, and $\Delta Lk>0$) for AFM, pUC19-T7A1-U plasmids were treated with different enzymes to obtain the desired topology and they were further purified by agarose gel extraction using the β-Agarase described above. To generate nicked topology, plasmids (20 µg) were incubated in a 100 µL reaction with nicking endonuclease Nt.BspQI (20 units) in NEB 3.1 Buffer for 6 hours at 50 °C. For $\Delta Lk = 0$ topology, nicked plasmids (20 µg) were re-ligated by incubation in a 50 µL mixture with T4 DNA ligase (800 units) in T4 DNA ligase 1X Buffer at room temperature. For $\Delta Lk < 0$ topology, we used isolated plasmids with their native supercoiling degree as extracted from the stationary phase cells. For $\Delta Lk > 0$ topology, plasmids (600 fmol) were incubated in a 20 µL reaction with reverse gyrase TopR2 (0.1 µM), and ATP (1 mM) in Reverse Gyrase 1X Buffer (50 mM Tris–HCl pH 8.0, 20 mM MgCl$_2$, 100 mM NaCl, 0.5 mM DTT, and 0.5 mM EDTA) for 1 hour at 75 °C. Once plasmid topologies were generated, plasmids were purified using the Monarch PCR DNA Clean-up Kit.

Expression and purification of *Streptococcus pyogenes* catalytically inactive dCas9 and nickase Cas9 D10A (SpCas9 D10A)

Purification of SpCas9 D10A was performed using a reported protocol.[76] Briefly, plasmid MJ825 (Addgene, #39315) or MJ841 (Addgene, #39318) was transformed into BL21(DE3) cells. One liter of Terrific Broth culture containing 50 µg/mL kanamycin was grown at 37°C. Upon reaching OD 0.6, cells were induced with IPTG to a final concentration of 0.5 mM at 20°C and were grown overnight (~16 hours). Cells were harvested by spinning at 5000 rpm for 10 minutes and the cell pellet was stored at –80 °C. Cells were resuspended in 50 mL Lysis Buffer (50 mM Tris-HCl pH 7.5, NaCl 500 mM, 5% (v/v) glycerol, 1 mM DTT, supplemented with 4 tablets of mini-EDTA free

protease inhibitor) and lysed using a sonicator. The lysate was centrifuged for 30
minutes at 4°C at 20,000 g. The clarified supernatant was passed through a HisTrap 5
mL column, washed with buffer NiA (50 mM Tris-HCl pH 7.5, NaCl 500 mM, 5% (v/v)
glycerol, 1 mM DTT, 10 mM imidazole) and eluted on a linear gradient with buffer NiB
(50 mM Tris-HCl pH 7.5, NaCl 500 mM, 5% (v/v) glycerol, 1 mM DTT, 300 mM
imidazole). Positive fractions containing SpCas9 D10A (verified by SDS-PAGE) were
combined, and 2 mL of TEV protease were added and dialyzed overnight on Dialysis
buffer (50 mM Tris-HCl pH 7.5, NaCl 500 mM, 5% (v/v) glycerol, 1 mM DTT). The
samples were further purified on HiTrap Heparin column using a linear gradient of KCl
from 200 mM to 1000 mM (50 mM Tris-HCl pH 7.5, 5% (v/v) glycerol, 1 mM DTT),
followed by size-exclusion chromatography on a Superdex S300 column using Gel
Filtration buffer (20 mM Tris-HCl pH 7.5, KCl 200 mM, 5% (v/v) glycerol, 1 mM DTT).
774
Expression and purification of *S. solfataricus* reverse gyrase (TopR2)
The gene fragment expressing the reverse gyrase (TopR2) was obtained by PCR from
Sulfolobus solfataricus genomic DNA (ATCC, 35092D-5). This fragment was inserted
into an expression plasmid with His6 N-terminal tag with a TEV protease site (2Bc-T,
Addgene #37236) using ligation independent cloning. Protein expression was carried
with this plasmid (2Bc-T-TopR2) following a reported protocol.[77] Briefly, plasmid 2Bc-T-
TopR2 was transformed into Rosetta (DE3) cells. One liter of 2xYT culture containing
100 µg/mL ampicillin, 34 µg/mL chloramphenicol, and 1% glucose was grown at 37°C.
Upon reaching OD 0.5, cells were induced with IPTG to a final concentration of 1 mM at
37°C and were grown for 6 hours. Cells were harvested by spinning at 5000 rpm for 10
min and the cell pellet was stored at –80°C. The pellets of induced cells were
resuspended in Lysis buffer (40 mM Tris–HCl pH 8.0, 100 mM NaCl, 1 mM DTT, 0.1
mM EDTA). After sonication on ice, the sample was centrifuged at 30,000 g for 30
minutes. The resulting supernatant underwent heat treatment at 75°C for 13 minutes
and was clarified by centrifugation at 30,000 g for 20 minutes. Polyethylenimine was
then added to achieve a final concentration of 0.3%, and the mixture was stirred for 1
hour before being centrifuged at 40,000 g for 30 minutes. The supernatant was adjusted
to 70% $(NH_4)_2SO_4$ saturation, stirred for 30 minutes at 4°C, and centrifuged at 20,000 g
for 30 minutes. The proteins were dissolved in buffer B (Lysis Buffer with 200 mM NaCl
and 10% (v/v) ethylene glycol). After dialysis against buffer B, the proteins were loaded
onto a 5 mL HiTrap Heparin HP column pre-equilibrated with buffer B using an FPLC
AKTA system (GE Healthcare). The column was washed with 50 mL of buffer B, and
bound proteins were eluted with a NaCl gradient ranging from 0.2 to 1.2 M NaCl in
buffer B.
799
SpCas9 D10A nicking assay
To test the activity of the sgRNA and nickase dCas9, a nicking assay was developed to
monitor activity by electrophoretic mobility of DNA in agarose gel. sgRNA (90 fmol) was
incubated at room temperature for 5 min with SpCas9 D10A (90 fmol) in Cas9 Reaction
Buffer (20 mM HEPES pH 6.5, 100 mM NaCl, 5 mM $MgCl_2$, 0.1 mM EDTA). Supercoiled
plasmid (9 fmol) was added, and the reaction was kept at 37°C for one hour. The

samples were treated with Proteinase K and then loaded into an agarose gel (1%
agarose, 1X TAE 2.5 µg/mL chloroquine).
808

DNA topoisomerase relaxation assay

The DNA relaxation activity of *E. coli* DNA topoisomerase I on supercoiled pUC19-
T7A1U was tested under different reaction conditions. In all reaction conditions, a total
reaction volume of ~40 µL containing ~200 ng (~ 0.157 pmol) of negatively supercoiled
plasmid was used and relaxation was carried out in TB40 buffer (20 mM Tris pH 8.0, 40
mM KCl, 5 mM $MgCl_2$, 0.02 mg/mL BSA, and 1 mM DTT). An incubation at 37°C for 20
minutes was done for all conditions prior to the addition of *E. coli* Topoisomerase I (New
England Biolabs, M0301S, 0.720 pmol). (1) For Topoisomerase I only condition,
topoisomerase was added after an incubation of 37°C for 20 minutes. (2) To test the
effect of a stalled transcription complex, *E.coli* RNA Polymerase Holoenzyme (New
England Biolabs, M0551S, 1.700 pmol) was supplemented to the previous reaction
condition with 10 µM of ATP, GTP, CTP (Thermo Scientific, R0481) and 50 µM of GpA
dinucleotide (TriLink Biotechnologies) and incubated during 20 min at 37°C. (3) To test
for the effect of a dCas9: sgRNA complex, a sgRNA:dCas9 complex at a concentration
of 1.55 µM was first formed by incubating 15.5 pmoles of sgRNA and 15.5 pmoles of
dCas9 in TB40 buffer (10 µL) at 25°C for 10 minutes. Afterward, the sgRNA: dCas9
complex was incubated with the supercoiled plasmids for 20 minutes at 37 °C. (4) To
test the effect of *E.coli* RNAP without DNA-melting, *E.coli* RNAP core (NEB, 1.7 pmol)
was supplemented to the plasmid-only condition and incubated at 37°C incubation prior
to the addition of topoisomerase. (5) To test the effect DNA binding protein that does
not melt DNA, dEcoRI (EcoRI Q111, 2 pmol, a gift from ModrichLab) was supplemented
to the plasmid-only condition and incubated at the 37°C incubation prior to the addition
of topoisomerase.
In each assay, after 20 minutes of incubation, *E. coli* Topoisomerase I (NEB ,
0.72 pmol) or human topoisomerase (hTop1B, 4U,Topogen) was added to a total
reaction volume of 40 µL and time-points were taken at 0, 1, 2, 3, 5, 10, 15, and 30
minutes and quenched in a 100 µL of a Stop Buffer containing 8 U Proteinase K (NEB),
4 µL Glycogen (5 mg/mL, Invitrogen), and 86 µL of 2X PK buffer (Tris 200 mM pH 7.5,
EDTA 25 mM, NaCl 300 mM, SDS 2%). The timepoints were extracted with phenol (pH >
7.5) and phenol/chloroform/isoamyl alcohol (25:24:1 Mixture, pH 6.7/8.0,
FisherBioReagents™) and then precipitated with 70% ethanol and redissolved in EB
buffer. The time points were electrophoresed using a 1.5% (w/v) agarose gel that was
run at 80V for 3 h with 1X TAE buffer (40 mM Tris-acetate, pH 8.1, 2 mM EDTA) in ice
and post-stained in a solution of SYBR Safe in water before being visualized using
Typhoon imager.
844

2D agarose gel electrophoresis

846 *DNA topoisomer ladder generation*
847 Plasmid pUC19-T7A1U was nicked with Nt.BspQI (New England Biolabs). Nicked
848 pUC19-T7A1U plasmids (400 ng) were religated in 1X T4 DNA ligase buffer with
849 increasing amounts of ethidium bromide (0, 40 ,80, 120, 160, 200, 400, and 800 ng)
850 using 100 U of T4 DNA ligase (New England Biolabs) in 40 µL reactions. The reaction

proceeded overnight at room temperature and was quenched with Proteinase K (NEB), followed by extraction using 2-butanol, phenol-chloroform extraction, and ethanol precipitation. A mixture of the different ligation reactions was made to generate the 2D topoisomer ladder.

Separation of DNA topoisomers in 2D agarose gel

A large 2% agarose slab was cast in 1X TAE containing 2.5 µg/mL chloroquine. 100 ng of plasmid was loaded on a narrow lane in the 2D gel and electrophoresis was carried out with the following conditions: 1) For the first dimension: 40V (22 hours) in 1X TAE 2.5 µg/mL chloroquine and 2) For the second dimension: 60V (8 hours) in 1X TAE 25 µg/mL chloroquine. The gel was then stained using SYBR Safe before imaging on Typhoon FLA 9500.

In vitro Transcription assay

To test the effect of DNA supercoiling, soft-torsional block by dCas9 and Topoisomerase I in single-round transcription elongation, we performed experiments using radiolabeled RNA. Stalled elongation complexes labeled with radiolabeled nascent RNA were formed by assembling *E.coli* RNA polymerase Holoenzyme (10 pmol, New England Biolabs) in pUC19-T7A1U (1 pmol) TB40 buffer (20 mM Tris pH 8.0, 40 mM KCl, 5 mM MgCl$_2$, 0.02 mg/mL BSA, and 1 mM DTT) in the presence of 10 µM of ATP, GTP, CTP (Thermo Scientific, R0481) and 50 µM of GpA dinucleotide (TriLink Biotechnologies) and α-P32-ATP (Perkin Elmer). The reactions were incubated for 20 min at 37 °C and then placed on ice. The excess of radiolabeled nucleotides was removed using a size-exclusion column Microspin G-25 (GE Healthcare). The stalled complexes were reinitiated by adding NTPs at 10 µM and reaction allowed to proceed at room temperature. Time points were quenched using 2X RNA formamide loading dye and taken at 0, 20, 40, 60, 80, 100, 120, 180, 240, 300, 450, 600, 750, 900, 1050, 1200, 1350, 1500, 1650, 1800 s. Quenched samples were ran in 10% denaturing urea gel 19:1 acrylamide/bisacrylamide, the resolved RNA were imaged on Typhoon FLA 9500 and the bands were quantified with ImageQuant TL 8.2.

The reactions were performed using nicked and supercoiled DNA templates of single promoter pUC19-T7A1U. For conditions where dCas9 was used, dCas9:sgRNA targeting TTS (CBR01) were incubated at 10 min after addition of *E. coli* RNAP Holoenzyme. When Topoisomerase I was used, *E. coli* Topoisomerase I was added 10 min after addition of *E coli* RNAP and allowed to relax DNA before isolating stalled elongation complexes. The abortive initiation measurements in **Figure 2D** were based on autoradiographed RNA intensities at time point 0 for both nicked and –sc pUC19-T7A1U templates. As can be seen from these experiments, more efficient loading and stalling of RNAP was observed on –sc DNA templates than nicked after removal of the excess radiolabeled nucleotides (i.e.: higher absolute amount was observed at 19 nt stalled site for the –sc condition than the nicked condition). To account for this, gel intensities were scaled accordingly in **Figure 3E**, and lane-normalization was used for quantification accordingly (See below in Gel quantification).

Gel quantification

The gel images were initially adjusted using the GIMP cage transformation function to correct bending along the horizontal direction, facilitating subsequent auto-processing steps. The quantification of gel intensity was facilitated by using the Python script gel_lane_finder script,[78] which enable the automated annotation of gel lanes. The two-dimensional band was averaged in the horizontal direction, resulting in a one-dimensional array representing the pixel-wise intensity along the annotated gel lane. To standardize the intensity values, which may be influenced by variations in sample loading for each lane, the sum of pixel intensity in the one-dimensional array within each lane was employed to normalize the intensity at each pixel point. Specific representative pausing positions were traced across the temporal evolution of the transcription. For supercoiling relaxation experiments, the presence of a small fraction of pre-existing nicked species in the negatively supercoiled DNA template (i.e., the top band at time zero, $\Delta Lk = 0$) may affect the interpretation of the results. Therefore, the $\Delta Lk = 0$ lane was excluded from the analysis.

Real-time transcription assays using a molecular beacon

To measure RNA synthesis in real time for both AmpR and mEGFP gene from the opposing and tandem constructs, we adapted a fluorescence-based methodology using molecular beacons that possess a 2'-O-methylribonucleotide backbone.[63] Molecular beacons CBMB01 and CBMB02 targeting 3' RNA ends of AmpR and mEGFP transcripts were used in separate reactions. Briefly, each reaction was performed in TB40 buffer (20 mM Tris pH 8.0, 40 mM KCl, 5 mM $MgCl_2$, 0.02 mg/mL BSA, and 1 mM DTT) in the presence of 1 mM of ATP, GTP, CTP, UTP (Thermo Scientific, R0481), 50 µM of GpA dinucleotide (TriLink Biotechnologies), 8 nM of plasmid, 85 nM of *E.coli* RNAP Holoenzyme (NEB), and 500 nM of molecular beacon. To ensure that all of the transcription reactions started simultaneously, the reaction mixtures were prepared on an ice-cold metal block. Then, all transcription reactions were incubated simultaneously in an iQ5 Thermal Cycler for 2 hour at 37°C. During the reaction, the fluorescence intensity FAM of the molecular beacon was recorded every 20 s.

Measurement of GFP fluorescence emission in *E. coli* MG1655 with tandem and opposing constructs

The following experiments were performed using similar experimental conditions as reported by Kim et al.[24] Briefly, plasmids for the tandem and opposing constructs (pUC19-T7A1-AmpR-mEGFP-Tandem and pUC19-T7A1-AmpR-mEGFP-Tandem) were transformed into *E.coli* MG1655 cells using standard electroporation protocols. Positive clones were selected by their growth on an ampicillin antibiotic plate and by their fluorescence emission in a trans-illuminator under blue light. Three colonies from each construct were selected, mixed and a ~ 5 mL inoculate grown in M9GluCAAT media containing M9 minimal medium (6 g/L Na_2HPO_4, 3 g/L KH_2PO_4, 0.5 g/L NaCl, 1 g/L NH_4Cl, 2 mM $MgSO_4$, 0.1 mM $CaCl_2$) supplemented with 0.2% glucose, 0.1% casamino acids (Difco Laboratories),1 mg/L thiamine and ampicillin 50 µg/mL at 37 °C. The overnight inoculate was then diluted 10,000 fold and then grown to exponential phase in M9GluCAAT with ampicillin 50 µg/mL to an optical density $OD_{600}=0.2$. Then, ~0.5 µL of these cells were spotted into an agarose pad made with 1% agar and

M9GluCAAT covered with a no. 1.5 coverslip, then imaged in the microscope at room
temperature. Imaging by phase contrast and fluorescence microscopy was performed
on a Nikon Eclipse Ti microscope equipped with either a 1.40 NA phase-contrast oil
objective, a Hamamatsu Orca-Flash4.0 V2 CMOS camera, a Sola Light Engine
(Lumencor) and pE-4000 (CoolLED) light source. For GFP excitation/emission the
ET470/40x Chroma dichroic was used. The microscopes were controlled by the Nikon
Elements software.
948
Atomic Force Microscopy
Slow scan Atomic Force Microscopy (AFM)
DNA samples of different topologies of pUC19-T7A1U (1-2 nM) were diluted in 10 mM
HEPES pH 7.0 and 5 mM MgCl$_2$ prior to sample deposition. Two microliters of DNA
were deposited on freshly cleaved mica and incubated at room temperature for 2 or 3
minutes. Mica was rinsed with 50 µL of water five times and dried under N$_2$ gas flow.
AFM measurements were performed with a Multimode AFM Nanoscope 8 (Bruker Co.).
The samples were imaged in tapping mode; the silicon cantilevers (Nanosensors) were
excited at their resonance frequency (280–350 kHz) with free amplitudes of 2-10 nm.
The image amplitude (set point As) and free amplitude (A0) ratio (As/A0) was kept at
0.8. All samples were imaged at room temperature in air, at a relative humidity of 30%.
Raw static AFM images acquired in air were flattened and leveled using Gwyddion 2.5.
A mask of the entire DNA molecule using thresholding was obtained and only pixels
corresponding to DNA regions shown in **Figure 1C**.
963
Cryo-ET sample preparation for transcriptional modulation of DNA supercoiling
Preparation of circular plasmid under low salt and high salt conditions
For the preliminary investigation into the conformational dynamics of DNA supercoiling
in response to varying ionic strength, the above extracted and purified pUC19-T7A1U
plasmid was diluted to a concentration of 100 nM under two distinct buffer conditions:
low salt and high salt buffer (20 mM Tris-Cl, 0.5 mM DTT, and 2% Trehalose; 20 mM
Tris-Cl, 40 mM KCl, 5 mM MgCl$_2$, 0.5 mM DTT, and 2% Trehalose, respectively)
preceding the application onto the EM grids.
972
Formation of RNAP-pUC19-T7A1U stalled complex
To investigate the localization of RNAP on the plasmid, a 10 µL incubation reaction was
prepared to induce RNAP stalling 19 nucleotides downstream from the TSS of the T7A1
promoter. This was achieved by employing UTP starvation in a system composed of 60
nM pUC19-T7A1U plasmid, 180 nM RNAP (3:1 ratio to the plasmid), 50 µM GpA
dinucleotide primer (facilitating RNAP initiation), and 10 µM rNTP mix (excluding UTP).
The reaction buffer consisted of 20 mM Tris-Cl, 40 mM KCl, 5 mM MgCl$_2$, 0.5 mM DTT,
and 2% Trehalose. This reaction was then allowed to incubate for 20 minutes at 37°C to
create the RNAP stall complex before application onto the TEM grids.
982
983 *Formation of dCas9-pUC19-T7A1U complex*
984 Following the same incubation protocol as described above, the RNAP in the system
985 was replaced by dCas9, an alternative DNA-binding protein. To do so, the dCas9

protein and its guiding RNA (sgRNA) were first assembled in a separate vial at a 1:1 ratio, reaching a concentration of 500 nM. This assembly was achieved by incubating the dCas9-sgRNA mixture in a buffer containing 20 mM Tris-Cl, 40 mM KCl, 5 mM $MgCl_2$, and 0.5 mM DTT for 10 minutes at room temperature. Subsequently, the incubated dCas9-sgRNA complex was introduced into the plasmid, yielding a system containing 60 nM pUC19-T7A1U, 120 nM dCas9-sgRNA (2:1 ratio to the plasmid), a GpA dinucleotide primer (50 µM) for initiation, and an rNTP mix (10 µM, excluding UTP). The reaction buffer composition remained consistent with 20 mM Tris-Cl, 40 mM KCl, 5 mM $MgCl_2$, 0.5 mM DTT, and 2% Trehalose. This reaction mixture was further incubated for 20 minutes at 37°C before being applied to the TEM grids for further analysis.

Formation of transcribed (none-equilibrium) RNAP-pUC19-T7A1U complex

To study the activation of RNAP transcription on the negatively supercoiled plasmid, the RNAP stalled complex was first prepared as described above. At the end of its 20-minute incubation at 37°C, rNTP (including UTP) was added to the solution, adjusting the concentration of each nucleotide to reach 100 µM. The transcription reaction continued for 10 minutes at room temperature before being transferred onto ice prior to application onto the TEM grids. For single-particle analysis, the transcription elongation complex concentration (both RNAP and plasmid) was scaled up 4-fold without changing the buffer conditions.

Formation of stalled and transcribed RNAP-dCas9-pUC19-T7A1U complexes

To study active RNAP transcription on the negatively supercoiled plasmid in the presence of dCas9 torsional block, the previous dCas9-pUC19-T7A1U complex was first prepared. Subsequently, RNAP was introduced into the incubated solution at a 3:1:1 ratio to dCas9 and plasmid, with concentrations of 180 nM, 60 nM, and 60 nM, respectively, maintained under the same physiological salt condition. In this sequence, the addition of rNTP (10 µM final concentration, excluding UTP) allowed the accumulation of mild torsion as RNAP initiated transcription from the TSS to the U-less stalling site, spanning 19 nucleotides. To induce additional torsional stress in the system, rNTP (including UTP) was added to the stalled RNAP-dCas9-pUC19-T7A1U complex at 100 µM final concentration. The sample was subjected to a 10 minutes incubation at room temperature for DNA transcription before being transferred to the EM grids.

Formation of transcribed RNAP-topI-pUC19-T7A1U complexes

To relieve RNAP from its apical constraint, topoisomerase I was introduced into the transcription system. The stalled RNAP-pUC19-T7A1U complex was initially prepared following the same established protocol. After a 20-minute incubation at 37°C, topoisomerase I was added to the solution at a ratio of 3:1:1 with respect to RNAP and plasmid at concentrations of 180 nM, 60 nM, and 60 nM, respectively, under the same physiological salt condition. Subsequently, rNTP (including UTP) was added to the solution mixture, reaching a final concentration of 100 µM, and the transcription persisted for 10 minutes at room temperature before application onto the EM grids.

Formation of transcribed RNAP-pUC19-T7A1U dual-promoter complexes

To study RNAP transcription activation on negatively supercoiled plasmids containing dual promoters (i.e., opposing and tandem constructs), RNAP-stalled complexes were prepared using the same procedure as described above for the single-promoter pUC19-T7A1U template, with the exception that the RNAP concentration was doubled to 360 nM (corresponding to a 6:1 molar ratio of RNAP to plasmid) to accommodate dual-promoter initiation. Following a 20-minute incubation at 37 °C, rNTPs (including UTP) were added to a final concentration of 100 μM for each nucleotide. The transcription reaction was allowed to proceed for 10 minutes at room temperature and then placed on ice prior to application onto TEM grids.

TEM specimen preparation

3 μL of a sample after the above incubation was deposited onto a glow-discharged (PELCO easiGlow™ Glow Discharge Cleaning System) 200 mesh Quantifoil gold grid (hole size ranging from 1 micron to 2 microns, Electron Microscopy Sciences) for 30 seconds. Following a 20-second on-grid incubation, the grid was plunge-frozen in liquid ethane at ~95% humidity and 15°C using a Leica EM GP rapid-plunging device (Leica, Buffalo Grove, IL, USA) after controlled blotting with filter paper (3-5 s). The resulting flash-frozen grids were then transferred into liquid nitrogen for storage.

TEM data acquisition

The Cryo-EM data were acquired using a Titan Krios (FEI) transmission electron microscope operating at 300 kV high tension and equipped with a Gatan energy filter. Imaging was performed with a Gatan K3 Summit direct electron detection camera, employing a magnification of x53,000 (where each pixel of the micrographs corresponds to 1.67 Å in specimens) in super-resolution and correlated double sampling (CDS) mode. Tilt series of the samples were captured from -51° to +51° at 3° step or -55° to +55° at 5° step increments using a dose symmetry scheme. Data collection was automated using SerialEM software[79] to track the specimen and maintain a defocus of ~2.5 μm. The total dose for the tilt image series ranged from ~110-150 e⁻/Å². At each tilt angle, a total of 8 frames were collected, with an exposure time of 0.25 s per frame. Single-particle movie data were collected using a Gatan K3 direct electron detector at a nominal magnification of ×81,000, corresponding to a pixel size of 1.05 Å. Each exposure lasted 7.39 seconds and was recorded over 50 frames, with a total electron dose of 50 e⁻/Å². Micrographs were acquired in non-CDS and super-resolution modes, with a defocus range of 0.6–1.6 μm, using SerialEM.

Cryo-ET tilt series processing and 3D reconstruction

The beam-induced motion of cryo-EM frames was corrected using MotionCor2.[80] To improve the contrast of low-dose cryo-ET tilt series images, a deep learning-based denoising method (NOISE2NOISE[81]) was implemented, involving the division of motion-corrected movie frames into even and odd halves for network training and subsequent particle resolution estimation (**Supplementary Data 2, I**). Predictions for even and odd frames for a single tilt were averaged together. The defocus value of the cryo-ET tilt series was determined using GCTF.[82] A carbon area perpendicular to the tilt axis was

included during data collection to assist in contrast transfer function (CTF) detection and later tilt series alignment. The tilt series were initially aligned using IMOD with patch tracking function[83] and subsequently imported into e2tomo software[28] for 3D reconstruction (**Supplementary Data 2, II**). Approximately 100 subtomogram patches containing DNA features were cropped for manual annotation and subsequently submitted for training. This convolutional neural network (CNN) was then applied across all tomograms for DNA identification (**Supplementary Data 2, III**).

Individual particle 3D reconstruction refinement and modeling
After the deep learning-based DNA annotation, individual plasmid particles were manually labeled in the tomogram (**Supplementary Data 1, III, right**). In brief, the DNA-annotated tomograms were binned by 4 (6.68 Apix) and divided into isolated small surface pieces within Chimera.[84] Density segments from a single plasmid particle were then manually selected and grouped, followed by low-pass filtering to a 6 nm resolution to serve as the particle-shape mask. The criteria for selecting plasmid particles from the annotated tomogram included: 1) the particle should display supercoiling, manifesting as a helical structure (cyan color) without nicking (red color) (**Supplementary Data 1, III, right**), 2) plasmid particles should be isolated and not significantly entangled with others; and 3) particles not partially attached to the carbon area. The center of mass of the particle shape mask was calculated (**Supplementary Data 1, IV, left**), and this information was used to crop 1400x1400 pixel local tilt series within the raw large micrograph before denoising. This cropped tilt series, with a plasmid particle in the center, was binned by 7 (a larger crop area and bin=8 were used for the later opposing and tandem construct plasmid particles for their larger size) and then submitted for local 3D reconstruct with EMAN2.[28] To minimize artifacts caused by the limited tilt angle range, the 3D reconstructed maps were missing-wedge compensated using IsoNet software[29] (**Supplementary Data 1, V**), employing the previously created low-resolution mask. The output map was then low-pass filtered to 2 nm, serving as the final map for the modeling.

Plasmid modeling was facilitated through the following steps: from the 3D reconstructed maps of the plasmid particle, a spine line representing the particle's super-helical axis was initially computed (**Supplementary Data 2, first row, left**). This involved applying a strong Gaussian kernel to the map, followed by skeletonization using the 'lee' method from the scikit-image package. Along the determined spine line, a sampling cylinder with dimensions of 60 nm diameter and 15 nm height was generated, utilizing a 5 nm step. The DNA density within the cylinder was divided into two regions, and the weight centers for each region were recorded (**Supplementary Data 2, first row, right**). Subsequently, the tracing centers were interconnected based on rotation and distance to the previous centers, with a subsequent round of manual correction undertaken to address misconnection cases, particularly in low-quality areas of the EM map (**Supplementary Data 2, second row, left**). The resulting threaded points underwent final processing through smoothing and interpolation with a 2 nm spatial interval, yielding the final model for the 3D map of the supercoiled DNA (**Supplementary Data 2, second row, right**).

Evaluation of the cryo-ET 3D reconstruction resolution

The resolution for the individual particle reconstructions were estimated by two methods. 1) Data-to-Data based analysis: the Fourier Shell Correlation (FSC) was calculated between two independently reconstructed 3D maps, in which each map was based on one-half of the tilt-series (split by even and odd frames for each tilt). The frequencies at which the FSC curve first falls to values of 0.143 were used to represent the reconstruction resolution. 2) Data-to-Model based analysis: the FSC curve between the final 3D reconstruction and the density map converted from the corresponding fitting model was calculated. The frequencies at which the FSC curve fell below 0.5 was used to estimate the resolution. The density map of the fitting model was generated by pdb2mrc in EMAN software.[85]

Sub-tomogram averaging of RNAP and dCas9

The sub-tomogram averaging of RNAP and dCas9 particles was carried out utilizing the EMAN2 e2tomo. Briefly, the CTF-corrected and imod-aligned raw tilt series were imported into the software, followed by e2tomo 3D reconstruction, yielding a 4×-binned tomogram (equivalent to 6.68 Å with an unbinned pixel size of 1.67 Å). Subsequently, approximately 100 particles for each protein type were manually selected, and ab-initio maps were reconstructed for the subsequent round of reference-based boxing. Employing a template matching threshold value (vthr) of 7.5, a total of 1781 and 875 particles were extracted for RNAP and dCas9, respectively. Particle cropping, alignment, and averaging procedures were executed using the e2spt_refine_new.py script. No symmetry was applied during particle alignment. Following 8 rounds of alignment (iters=p,p,p,t,p,p,t,r), resolutions of 14.1 Å and 17.6 Å were determined for RNAP and dCas9, respectively (0.143 cut-off), employing Fourier Shell Correlation (FSC) with odd and even particles from the masked, final average. By mapping the orientation-determined particles back to their tomogram using e2spt_mapptclstotomo.py, DNA-bound particles were selectively chosen if their center of mass was within a distance of <1 nm to any points of the plasmid model. A subset of 232 RNAP and 116 dCas9 particles was then subjected to another round refinement (iters=p,p,p,t,p), resulting in resolutions of 17.7 Å and 18.9 Å, respectively, evaluated under the same standards.

Single-particle analysis of TECs on negatively supercoiled DNA template

Single-particle 3D reconstruction was performed using cryoSPARC.[86] The detailed data processing workflow for TECs on negatively supercoiled DNA templates is shown in **Figure S4A**. A total of 10,476 micrographs underwent patch motion correction, patch CTF estimation, and particle picking, yielding 2,063,152 particles. Initial 2D classification and ab initio reconstruction revealed moderate orientation bias of RNAP particles. To preserve rare views, particle cleanup was performed via three rounds of heterogeneous refinement against 6 models—5 representing noisy junk references and one low-pass-filtered good model from the initial 3D reconstruction—instead of standard 2D classification. This cleanup yielded 611,747 particles (binned 3, in a 108-pixel box), which were used for homogeneous refinement, achieving a 6.5 Å reconstruction. Subsequent 3D classification at 15 Å resolution separated two major classes: TIC (186,816 particles, with σ-factor bound) and TEC (424,931 particles, without σ). The

TEC particles were re-boxed (unbinned, in a 324-pixel box) and subjected to ab initio
reconstruction, non-uniform refinement, and orientation rebalancing using cryoSPARC's
default settings. Further 3D classification at 10 Å resolution of TEC particles yielded
three classes: Class I (TEC1–SC) and Class II (TEC2–SC), both showing DNA in the
RNAP active site and a downstream DNA density protrusion, and Class III, which
showed only faint DNA density and was not analyzed further. Non-uniform and local
refinement of TEC1–SC (46,743 particles) and TEC2–SC (54,869 particles) yielded final
structures at 2.9 Å resolution. Maps were post-processed with DeepEMhancer.[87] Initial
models were built by fitting domains from known TEC structures (e.g., PDB: 6ALH) into
the density maps using ChimeraX.[88] Models were iteratively refined using
phenix.real_space_refine,[89] manually adjusted in ISOLDE,[90] and validated with
MolProbity[91] (**Supplementary Data 4**). The swivel angle calculation for TEC1–SC and
TEC2–SC followed the definition of the core and swivel modules as described by Kang
et al.,[52] and using PDB structure 6RH3 as the reference. Additional structures (1:6RH3,
2:8EG8,4:6ALH, 5:8EHI, 7:8EHF, and 8:7PYK) were also included in the measurement
as control. Measurements were performed in ChimeraX, and the ChimeraX script is
available in the Zenodo repository.

Plectoneme apical site sequence-based prediction
Prediction of DNA plectoneme formation loci along the pUC19-T7A1U sequence was
performed using a computational model developed by Kim et al.[43] To account for the
circular topology of the plasmid while matching the required linear DNA input format, the
pUC19-T7A1U sequence was duplicated three times and joined head-to-tail to simulate
continuity; only the central region corresponding to the original plasmid was plotted to
eliminate edge artifacts. To provide a reference for interpreting the plectoneme
formation scores, a control sequence consisting of polyA–Widom 601–polyA was used.
This construct includes the well-characterized Widom 601 nucleosome positioning
element, flanked by polyadenine tracts, and is known for its high intrinsic bendability,
which provides a comparative baseline for sequence-driven plectoneme localization.

Quantification of supercoiled DNA plasmid morphology
The quantification of the supercoiled DNA plasmid's morphology involved
measurements of radius of gyration, global writhe number, and plectoneme branch
writhe density. To quantify these characteristics, the eigenvalues of the gyration tensor
were calculated and designated as r1, r2, and r3, with the relationship r1 > r2 > r3. The
radius of gyration (R) was determined by the formula $R = r1^2 + r2^2 + r3^2$. The calculation
of the writhe number was facilitated by the E-CAM polymer_data_analysis module,
which implements a double integral computation method proposed by Klenin &
Langowski.[92]

Determination of the curvature and apexes along the supercoiled DNA
1207 To determine the curvature of the plasmid along its DNA trajectory, a loop iterated from
1208 index 1 to N-1 for each point on the DNA plasmid model. The curvature (κ) at each point
1209 (i) was computed with its two neighboring points (i-1 and i+1) using the formula κ = 1/R
1210 = 4S/fgh,[93] where R is the radius of the circle circumscribing a triangle formed by the

three points. Here, S represents the area of the triangle, and f, g, and h are the side lengths opposite the vertices of the triangle. The curvature at each point along the DNA trajectory was color-coded using a gradient, with points exhibiting the largest curvature values near the distal ends of each plectoneme branch utilized as the apexes of the plasmid.

Calculation of the distance between DNA along the plasmid's helical axis

To calculate the distance between DNA along the supercoiling helical axis, the points on the DNA plasmid model were sequentially sorted and segmented into different color groups using the apical points on the plasmid (**Figure S1E**). For each point (i) in a group, its nearest neighboring point in another color group was determined, and the center of mass as well as the distance between the two points were calculated. This calculation was repeated from point 1 to N for each segment, resulting in a list of centers and distances that were used to represent writhe helix axis and the distances between the DNA along the supercoiling helix (**Figure S3F, II**). Given the difficulty of defining the start and end site for a circular plasmid, the apexes were utilized as the zero index to align the calculated distance list.

Determination of the local writhe density of the plasmid

Utilizing the previously calculated writhe helical axis trace of the model (**Figure S3F, II**), the branches of the plasmid can be determined and split at the junction (**Figure S3F, I**). Each helical axis branch was color-coded, and these color codes were used to segment the points on the plasmid model based on their vicinity (**Figure S3F, III**). These branch segmentations were treated as smaller closed plasmids and subjected to writhe number calculation using the previously described method.[92] In the statistical analysis, to account for the greater contribution of longer plectonemes to writhe density, long DNA plectonemes were segmented into 100 nm units. Plectonemes shorter than 100 nm were left unsegmented and included as individual units.

Construction of superimposed models and mapping RNAP position on plasmid

To align all models from the same experimental condition, orientation-determined particles within the same group were back-mapped to their respective tomograms. Concurrently, the plasmid model was also back-mapped to the tomogram, so that it allows the determination of the relative orientation between the plasmid and plasmid-bound particles (RNAP and dCas9). Given the current resolution limitations of cryo-ET in resolving DNA sequence information, the position of dCas9 was utilized as a marker to infer the TSS, assuming specificity in dCas9 binding as per the construct design. The entire length of the plasmid trace was normalized to 1950 base pairs, with the point nearest to the dCas9 particle designated as 1090 bp. As none of the particles are symmetrical, the upstream and downstream positions of the bound RNAP were determined relative to the plasmid.

Coarse-grained simulation of DNA supercoiling relaxation

Coarse-grained molecular dynamics (CGMD) simulations were conducted using the GPU-accelerated, sequence-dependent oxDNA2 model.[42,94] The simulation environment

was set up under the canonical NVT ensemble (constant number of particles, volume, and temperature). To enhance computational efficiency, the 525-bp minicircle DNA system were employed. The in-silico plasmid model was generated using TacoxDNA,[95] with a DNA twist deficit of -5 and a writhe of 0, yielding a supercoiling density of -0.1. The circular DNA plasmid was energy minimized and relaxed using parameters tailored from oxDNA example simulations (at dna.physics.ox.ac.uk). These parameters included the following (thermostat = john, interaction_type = DNA2, newtonian_steps = 103, diff_coeff = 2.5, salt_concentration = 0.5, T = 300K, dt = 0.005, verlet_skin = 0.05, rcut = 2.0).

To study the impact of varying levels of plectoneme apex constraint on DNA supercoiling structural dynamics, simulations were initiated by introducing a small bulge on the minicircle DNA (**Figure S3A, left**), which facilitated apex formation at the target site during relaxation. The DNA base pairs at the two distal ends of the bulge were subjected to different levels of constraint, implemented as a mutual force trap between the two red sites (**Figure S3A, dashed line box**), with spring stiffness set to 1 and spring equilibrium distances (r0) set to 8.0 nm, 6.4 nm (80%), and 4.8 nm (60%). A fully unconstrained condition was also included. For each condition, the plasmid was first energy minimized for 2×10^6 steps, followed by relaxation for 3×10^7 steps (equivalent to 300 μs), with a frame sampling interval of 5×10^4 steps, yielding 600 frames. The final 500 frame structures after plasmid relaxation and apices formation were used to compute the global writhe distribution and plectoneme apex curvature statistics.

To study the effect of plectoneme apex constraint on DNA supercoiling torsional relaxation, simulations were initiated by relaxing the minicircle for 3×10^6 steps (30 μs) until the formation of plectonemes and the appearance of apices. The final frame structure was edited to create a nicked plasmid by introducing a single-strand break at base pair index 940 for models I and II, 828 for model III, and 564 for model IV (**Figure S5C**). This nicked configuration was designed to mimic the activity of Topoisomerase I (TopI). Four DNA supercoiling torsional relaxation simulations were conducted: For model I, the nicked plasmid was allowed to relax without any external constraints for 3×10^7 steps (equivalent to 300 μs). For the model II, III, and IV, a mutual force trap was applied to constrain the sharp apex region, covering approximately 20 base pairs. These traps, functioning as springs, were formed between base pairs at indices 798, 251 on one side and 817 and 232 on the other. The spring stiffness was set to 1, and r0 was defined as the initial distance between the mass centers of each pair of base pairs in the initial structure divided by a factor of 2. The second batch of simulations were also run 3×10^7 steps. All simulations were repeated two times using the same relaxation parameters mentioned above. Movies were generated using the last run of each simulation at a rate of 1×10^5 steps per frame.

The equilibrated final frame structure prior to the nick was also utilized to simulate the formation of DNA twin domains. In brief, two rotating harmonic traps were applied to the base pairs at indices 107 and 942, as well as indices 366 and 683, positioned within the intermediate region of the plectoneme. This approach segments plectoneme two sections: the first section contains an apex with constraints, while the second section features a free apex. This setup allows for a comparison of the effects of (+) torsional propagation on one strand, (-) torsional propagation on the other, and their

mutual cancellation when the rotating trap is applied. The center of rotation was defined
as the center of mass of the selected base pair, with the rotation axis determined as the
unit vector extending from the center of the base pair to the center of mass of another
base pair located 10 base pairs upstream. The stiffness of the trap was set to 10 in
oxDNA, with a rotation rate of $1x10^5$. The confined apex was configured as described
earlier. The simulation was run for $3x10^6$ steps and with one repeat. Movies were
created with frames captured every $1x10^4$ steps.
1308

SUPPLEMENTAL INFORMATION

Video S1. 3D reconstruction workflow of a representative negatively supercoiled 2 kbp
plasmid particle
Video S2. Coarse-grained MD simulation of negatively supercoiled plasmid relaxation
with and without apical constraint
Video S3. Coarse-grained MD simulation of negative DNA supercoiling in managing
positive and negative torsion with and without apical constraints
1316
1317

KEY RESOURCE TABLE

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Bacterial strains		
BL21(DE3) pLysS *E. coli* competent cells	Macrolab, QB3-Berkeley	N/A
DH5α E. coli competent cells	Macrolab, QB3-Berkeley	N/A
NEB® Stable *E. coli* competent cells	New England Biolabs	C3040H
MG1655	Coli Genetic Stock Center at Yale	Cat#CGSC6300
MG1655/pUC19-T7A1U-AmpR-mEGFP-Tandem	This study	N/A
MG1655/pUC19-T7A1U-AmpR-mEGFP-Opposing	This study	N/A
Rosetta(DE3) pLysS	Macrolab, QB3-Berkeley	N/A
Chemicals, peptides, and recombinant proteins		
β-Agarase I	New England Biolabs	Cat#M0392S
Ampicillin	Sigma-Aldrich	Cat#A0166; CAS: 69-53-4
ATP, [α^{32}P]	Perkin Elmer	Cat#BLU003H250UC
Bovine serum albumin (BSA)	New England Biolabs	Cat#B9000
BsaI-HF®v2	New England Biolabs	Cat#R3733
BsmBI-v2	New England Biolabs	Cat#R0739
Calcium chloride dihydrate	Sigma-Aldrich	Cat#C5080; CAS: 10035-04-8
Chloramphenicol	Sigma-Aldrich	Cat#C0378; CAS: 56-75-7
Chloroform	Sigma-Aldrich	Cat#C2432; CAS: 67-66-3
Chloroquine diphosphate salt	Sigma-Aldrich	Cat#C6628; CAS: 50-63-5
cOmpleteTM, EDTA-free protease inhibitor cocktail	Roche	Cat#11873580001
Dithiothreitol (DTT)	Sigma-Aldrich	Cat#43815; CAS: 3483-12-3
EcoRI-HF®	New England Biolabs	Cat: #R3101S
dEcoRI (EcoRI R111Q)	Wright, D. et al[96]	N/A
E. coli RNA Polymerase, Core Enzyme	New England Biolabs	Cat#M0550S
E. coli RNA Polymerase, Holoenzyme	New England Biolabs	Cat#M0551S
E. coli topoisomerase I	New England Biolabs	Cat#M0301S
Ethanol, 200-proof	Koptec	Cat#V1016; CAS: 64-17-5

Ethylenediaminetetraacetic acid, EDTA	Sigma-Aldrich	Cat#E6758; CAS: 60-00-4
Formamide (Deionized)	Sigma-Aldrich	Cat#F9037; CAS: 75-12-7
Glycerol	Sigma-Aldrich	Cat#G5516; CAS: 56-81-5
Glycogen (5 mg/ml)	Invitrogen	Cat#AM9510
GpA RNA Dinucleotid（5'-3'）	TriLink Biotechnologies	Cat#O-31009-05
HEPES	Sigma-Aldrich	Cat# H4034; CAS: 7365-45-9
Human Topoisomerase IB	Topogen	TG2005HRC3
Imidazole CAS: 367-93-1	Sigma-Aldrich	Cat#I5513; CAS: 288-32-4
Isopropyl b-D-1-thiogalactopyranoside, IPTG	Thermo Fisher Scientific	Cat#50-490-794; CAS: 367-93-1
Kanamycin sulfate	Sigma-Aldrich	Cat#60615; CAS: 70560-51-9
LB broth, Miller	Fisher Bioreagents	Cat#BP1426-2
Magnesium chloride, MgCl2	Thermo Fisher Scientific	Cat# BP214; CAS: 7786-30-3
Magnesium sulfate, MgSO4	Sigma-Aldrich	Cat#M7506; CAS: 7487-88-9
Nt.BspQI	New England Biolabs	Cat#R0644S
NTP Set, 100 mM Solution	Thermo-Scientific	Cat#R0481
Phenol:Chloroform:Isoamyl Alcohol 25:24:1	Fisher Scientific	Cat#BP1752I-100
Phusion High-Fidelity DNA Polymerase	New England Biolabs	Cat#M0530S
Potassium chloride, KCl	Sigma-Aldrich	Cat#P9541; CAS: 7447-40-7
Potassium phosphate dibasic, K_2HPO4	Sigma-Aldrich	Cat#60353; CAS: 7758-11-4
Potassium phosphate monobasic, KH_2PO4	Sigma-Aldrich	Cat#P9791; CAS: 7778-77-0
Proteinase K	New England Biolabs	Cat#P8107S
SeaPlaque® Agarose	Lonza	Cat:#50100
SequaGel UreaGel 29:1 Concentrate	National Diagnostics	Cat#EC-828
Sodium chloride, NaCl	Sigma-Aldrich	Cat#S9888; CAS: 7647-14-5
Sodium dodecyl sulfate, SDS	Sigma-Aldrich	Cat#L5750; CAS: 151-21-3
Sodium phosphate dibasic, Na_2HPO4	Sigma-Aldrich	Cat#S3264; CAS: 7558-79-4
S. solfataricus reverse gyrase (TopR2)	This study	N/A
SYBR™ Safe DNA Gel Stain	Invitrogen	Cat:#S33102
Terrific Broth	Thermo Fisher Scientific	Cat#BP2468500
TEV Protease	Macrolab, QB3-Berkeley	N/A

TopVision Agarose	Thermo Fisher Scientific	Cat#R0491
T4 DNA Ligase	New England Biolabs	Cat#M0202
D-(+)-Trehalose dihydrate	Sigma-Aldrich	Cat#T9531
Tris Base	Genesee Scientific	Cat#18-146; CAS: 77-86-1
Urea	Sigma-Aldrich	Cat#U1250; CAS: 57-13-6
Critical commercial assays		
HiScribe® T7 High Yield RNA Synthesis Kit	New England Biolabs	Cat# E2040L
Monarch® Spin PCR & DNA Cleanup Kit (5 µg)	New England Biolabs	Cat# T1130S
QIAprep Spin Miniprep Kit	QIAGEN	Cat# 27104
QIAGEN Plasmid Maxi Kit	QIAGEN	Cat# 12162
Deposited data		
Cryo-ET maps of P. LS particles	Electron Microscopy Data Bank (EMDB)	EMD-47843
Cryo-ET maps of P. HS particles	Electron Microscopy Data Bank (EMDB)	EMD-47844
Cryo-ET maps of sTEC particles	Electron Microscopy Data Bank (EMDB)	EMD-47850
Cryo-ET maps of P. Cas particles	Electron Microscopy Data Bank (EMDB)	EMD-47849
Cryo-ET maps of TEC particles	Electron Microscopy Data Bank (EMDB)	EMD-47850
Cryo-ET maps of sTEC-Cas particles	Electron Microscopy Data Bank (EMDB)	EMD-47851
Cryo-ET maps of TEC-Cas particles	Electron Microscopy Data Bank (EMDB)	EMD-47853
Cryo-ET maps of TEC-TopI particles	Electron Microscopy Data Bank (EMDB)	EMD-47855
Cryo-ET maps of Opp-TEC particles	Electron Microscopy Data Bank (EMDB)	EMD-71622
Cryo-ET maps of Tan-TEC particles	Electron Microscopy Data Bank (EMDB)	EMD-71618
SPA maps of TEC1–SC	Electron Microscopy Data Bank (EMDB)	EMD-71675
SPA maps of TEC2–SC	Electron Microscopy Data Bank (EMDB)	EMD-71676
PDB of TEC1–SC	Protein Data Bank (PDB)	9PIP
PDB of TEC2–SC	Protein Data Bank (PDB)	9PIQ
Oligonucleotides		
CBD01: AATACTAGAATTCTTATCAAAAAGAGTATTGACTTAAAGTCTAACCTATAGGATACTTACAGCCGAAAAAAGCAACAAAAAAATTGAAAAAGGAAGAGTATGAGTATTCAAC	IDT	N/A
CBD02: AATACTAGAATTCCTGCATTAATGAATCGGCC	IDT	N/A

CBD03: AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATA ACGGACTAGCCTTATTTTAACTTGCTATTTCTAGCTCTA AAAC	IDT	N/A
CBD04: TTCTAATACGACTCACTATAGTGGTCATGAGATTATCAA AAGTTTTAGAGCTAGAAATAG	IDT	N/A
CBD05 GCGCCCAATACGCAAACCG	IDT	N/A
CBD06 CCAGGAACCGTAAAAAGGCCG	IDT	N/A
CBD07 GGCCTTTTTACGGTTCCTGGTGGCCGATTCATTAATGC AGGAATTCTTATC	IDT	N/A
CBD08 TCCTCGCCCTTGCTCACCATACTCTTCCTTTTTCAATTT TTTTGTTGCTTTTTTCG	IDT	N/A
CBD09: ATGGTGAGCAAGGGCGAGGAG	IDT	N/A
CBD10: GCGGTTTGCGTATTGGGCGCGATTAAAACGAAAGGCC CAGTCTTTCGA	IDT	N/A
CBD11: CCACGTCTCATTGAGATCCTTTTTTTTCTGCGCGTAATC T	IDT	N/A
CBD12: CCACGTCTCAACGCGCGGGGAGAGGCG	IDT	N/A
CBD13: CCACGTCTCAGCGTGAAGATCCTTTGATCTTTTCTACG GGGTCTG	IDT	N/A
CBD14: CCACGTCTCATCAATGGCCGATTCATTAATGCAGGAAT TCTTATC	IDT	N/A
CBR01 (sgRNA targeting TTS): GUGGUCAUGAGAUUAUCAAAAGUUUUAGAGCUAGAA AUAGCAAGUUAAAAUAAGGCUAGUCCGUUAUCAACU UGAAAAAGUGGCACCGAGUCGGUGCUUUU	This study	N/A
CBMB01 (Molecular Beacon for AmpR): /56-FAM/ mCmGmCmUmAmUmCmUmAmAmAmGmUmAmUmAm UmAmUmGmAmGmUmAmAmAmGmCmG/3BHQ_1/	IDT	N/A
CBMB02 (Molecular Beacon for mEGFP): /56-FAM/ mCmGmCmUmAmGmCmGmCmCmUmCmAmGmCmUm AmUmCmAmUmGmUmCmGmAmGmCmG/3BHQ_1/	IDT	N/A
Recombinant DNA		
pET His6 LIC cloning vector (2Bc-T)	Addgene	37236
p2Bc-T-TopR2	This study	N/A
pMJ825	Addgene	#39315
pMJ841	Addgene	#39318
pUC19	New England Biolabs	N3041S
pUC19-T7A1U	This study	N/A

pUC19-T7A1U-AmpR-mEGFP-Tandem	This study	N/A
pUC19-T7A1U-AmpR-mEGFP-Opposing	This study	N/A
Sulfolobus solfataricus genomic DNA	ATCC	35092D-5
Software and algorithms		
IMOD	Kremer et al.[83]	https://bio3d.colorado.edu/imod/
Serial EM	Schorb et al.[79]	https://bio3d.colorado.edu/SerialEM/
MotionCor2	Zheng et al.[80]	https://msg.ucsf.edu/software
UCSF Chimera	Pettersen et al.[84]	https://www.cgl.ucsf.edu/ chimera/
ChimeraX	Meng et al.[88]	
EMAN2	Chen et al.[28]	https://blake.bcm.edu/ emanwiki/EMAN2
CARE	Weigert et al.[97]	https://csbdeep.bioimage computing.com/tools/care/
DNA plectoneme site prediction	Kim et al.[43]	https://github.com/elifesciences-publications/Plectoneme_analysis
Cryo-SPARC	Punjani et al.[86]	https://cryosparc.com/
3D skeletonizing	Lee et al.[98]	https://scikit-image.org/docs/0.13.x/auto_examples/edges/plot_skeleton.html
IsoNet	Liu et al.[29]	https://github.com/IsoNet-cryoET/IsoNet
oxDNA	Suma et al.[95]	https://dna.physics.ox.ac.uk/index.php/Main_Page
DeepEMhancer	Sanchez-Garcia et al.[87]	https://github.com/rsanchezgarc/deepEMhancer
ISOLDE	Croll et al.[90]	https://tristanic.github.io/isolde/
PHENIX	Liebschner et al.[89]	https://phenix-online.org/documentation/overviews/cryo-em_index.html
MolProbity	Chen et al.[91]	https://www.phenix-online.org/documentation/reference/molprobity_tool.html
The E-CAM Software Library	Eastman et al.[30]	https://e-cam.readthedocs.io/en/latest/
Other		

Amicon Ultra-0.5 Centrifugal Filter Unit, 3K MWCO	Thermo Fisher Scientific	Cat#UFC500324
Amicon Ultra-0.5 Centrifugal Filter Unit, 10K MWCO	Thermo Fisher Scientific	Cat#UFC501024
Amicon Ultra-15 Centrifugal Filter Unit, 10K MWCO	Thermo Fisher Scientific	Cat#UFC901024
Amicon Ultra-15 Centrifugal Filter Unit, 30K MWCO	Thermo Fisher Scientific	Cat#UFC903008
HiPrepTM Sephacryl S300 16/60	GE Healthcare	Cat#17-1167-01
HisTrap HP	GE Healthcare	Cat#17-5248-02
HiTrap Heparin HP	GE Healthcare	Cat#17-0407-01
HiTrap Q HP	GE Healthcare	Cat#17-1154-01
UltraPure DNase/RNase-Free Distilled Water	Thermo Fisher Scientific	Cat#10977-023
Quantifoil R 2/1 200 mesh Au	Electron Microscopy Sciences	Cat#661-200-AU
Quantifoil R 3.5/1 200 mesh Au	Electron Microscopy Sciences	Cat#660-200-AU

1320
1321

References

1.   Baranello, L., Levens, D., Gupta, A., and Kouzine, F. (2012). The importance of being
supercoiled: How DNA mechanics regulate dynamic processes. Bba-Gene Regul Mech
1819, 632-638. 10.1016/j.bbagrm.2011.12.007.
2.   Gagua, A.V., Belintsev, B.N., and Lyubchenko Yu, L. (1981). Effect of base-pair stability
on the melting of superhelical DNA. Nature *294*, 662-663. 10.1038/294662a0.
3.   Yan, Y., Leng, F., Finzi, L., and Dunlap, D. (2018). Protein-mediated looping of DNA
under tension requires supercoiling. Nucleic acids research *46*, 2370-2379.
10.1093/nar/gky021.
4.   Tripathi, S., Brahmachari, S., Onuchic, J.N., and Levine, H. (2022). DNA supercoiling-
mediated collective behavior of co-transcribing RNA polymerases. Nucleic acids
research *50*, 1269-1279. 10.1093/nar/gkab1252.
5.   Patel, H.P., Coppola, S., Pomp, W., Aiello, U., Brouwer, I., Libri, D., and Lenstra, T.L.
(2023). DNA supercoiling restricts the transcriptional bursting of neighboring eukaryotic
genes. Molecular cell *83*, 1573-1587 e1578. 10.1016/j.molcel.2023.04.015.
6.   Fu, Z., Guo, M.S., Zhou, W., and Xiao, J. (2024). Differential roles of positive and
negative supercoiling in organizing the E. coli genome. Nucleic acids research *52*, 724-
737. 10.1093/nar/gkad1139.
7.   Vayssieres, M., Marechal, N., Yun, L., Lopez Duran, B., Murugasamy, N.K., Fogg, J.M.,
Zechiedrich, L., Nadal, M., and Lamour, V. (2024). Structural basis of DNA crossover
capture by Escherichia coli DNA gyrase. Science *384*, 227-232.
10.1126/science.adl5899.
8.   Vidmar, V., Borde, C., Bruno, L., Takacs, M., Batisse, C., Saint-Andre, C., Zhu, C.,
Espeli, O., Lamour, V., and Weixlbaumer, A. (2024). DNA topoisomerase I acts as
supercoiling sensor for transcription elongation in E. coli. bioRxiv.
9.   Janissen, R., Barth, R., Polinder, M., van der Torre, J., and Dekker, C. (2024). Single-
molecule visualization of twin-supercoiled domains generated during transcription.
Nucleic acids research *52*, 1677-1687. 10.1093/nar/gkad1181.
10. Pyne, A.L.B., Noy, A., Main, K.H.S., Velasco-Berrelleza, V., Piperakis, M.M., Mitchenall,
L.A., Cugliandolo, F.M., Beton, J.G., Stevenson, C.E.M., Hoogenboom, B.W., et al.

(2021). Base-pair resolution analysis of the effect of supercoiling on DNA flexibility and
major groove recognition by triplex-forming oligonucleotides. Nature communications *12*,
1053. 10.1038/s41467-021-21243-y.
11. Demurtas, D., Amzallag, A., Rawdon, E.J., Maddocks, J.H., Dubochet, J., and Stasiak, A.
(2009). Bending modes of DNA directly addressed by cryo-electron microscopy of DNA
minicircles. Nucleic acids research *37*, 2882-2893. 10.1093/nar/gkp137.
12. Irobalieva, R.N., Fogg, J.M., Catanese, D.J., Jr., Sutthibutpong, T., Chen, M., Barker,
A.K., Ludtke, S.J., Harris, S.A., Schmid, M.F., Chiu, W., and Zechiedrich, L. (2015).
Structural diversity of supercoiled DNA. Nature communications *6*, 8440.
10.1038/ncomms9440.
13. Amzallag, A., Vaillant, C., Jacob, M., Unser, M., Bednar, J., Kahn, J.D., Dubochet, J.,
Stasiak, A., and Maddocks, J.H. (2006). 3D reconstruction and comparison of shapes of
DNA minicircles observed by cryo-electron microscopy. Nucleic acids research *34*, e125.
10.1093/nar/gkl675.
14. Vologodskii, A.V., and Cozzarelli, N.R. (1994). Conformational and Thermodynamic
Properties of Supercoiled DNA. Annu Rev Bioph Biom *23*, 609-643. DOI
10.1146/annurev.bb.23.060194.003141.
15. Abdella, R., Talyzina, A., Chen, S., Inouye, C.J., Tjian, R., and He, Y. (2021). Structure
of the human Mediator-bound transcription preinitiation complex. Science *372*, 52-56.
10.1126/science.abg3074.
16. Aibara, S., Schilbach, S., and Cramer, P. (2021). Structures of mammalian RNA
polymerase II pre-initiation complexes. Nature *594*, 124-128. 10.1038/s41586-021-
03554-8.
17. Chen, X.Z., Qi, Y.L., Wu, Z.H., Wang, X.X., Li, J.B., Zhao, D., Hou, H.F., Li, Y., Yu, Z.S.,
Liu, W.D., et al. (2021). Structural insights into preinitiation complex assembly on core
promoters. Science *372*, 480-+. ARTN eaba8490

10.1126/science.aba8490.
18. Kettenberger, H., Armache, K.J., and Cramer, P. (2004). Complete RNA polymerase II
elongation complex structure and its interactions with NTP and TFIIS. Molecular cell *16*,
955-965. DOI 10.1016/j.molcel.2004.11.040.
19. Gnatt, A.L., Cramer, P., Fu, J.H., Bushnell, D.A., and Kornberg, R.D. (2001). Structural
basis of transcription:: An RNA polymerase II elongation complex at 3.3 Å resolution.
Science *292*, 1876-1882. 10.1126/science.1059495.
20. Liu, L.F., and Wang, J.C. (1987). Supercoiling of the DNA-Template during Transcription.
Proceedings of the National Academy of Sciences of the United States of America *84*,
7024-7027. DOI 10.1073/pnas.84.20.7024.
21. Ma, J., Bai, L., and Wang, M.D. (2013). Transcription Under Torsion. Science *340*, 1580-
1583. 10.1126/science.1235441.
22. Chong, S.S., Chen, C.Y., Ge, H., and Xie, X.S. (2014). Mechanism of Transcriptional
Bursting in Bacteria. Cell *158*, 314-326. 10.1016/j.cell.2014.05.038.
23. Leng, F.F., Amado, L., and McMacken, R. (2004). Coupling DNA supercoiling to
transcription in defined protein systems. Journal of Biological Chemistry *279*, 47564-
47571. 10.1074/jbc.M403798200.
24. Kim, S., Beltran, B., Irnov, I., and Jacobs-Wagner, C. (2019). Long-Distance Cooperative
and Antagonistic RNA Polymerase Dynamics via DNA Supercoiling. Cell *179*, 106-119
e116. 10.1016/j.cell.2019.08.033.
25. Jentink, N., Purnell, C., Kable, B., Swulius, M.T., and Grigoryev, S.A. (2023).
Cryoelectron tomography reveals the multiplex anatomy of condensed native chromatin

and its unfolding by histone citrullination. Molecular cell *83*, 3236-3252 e3237.
10.1016/j.molcel.2023.08.017.

26.   Zhang, M., Díaz-Celis, C., Liu, J., Tao, J., Ashby, P.D., Bustamante, C., and Ren, G.
(2024). Angle between DNA linker and nucleosome core particle regulates array
compaction revealed by individual-particle cryo-electron tomography. Nature
communications *15*, 4395.

27.   Beel, A.J., Azubel, M., Mattei, P.J., and Kornberg, R.D. (2021). Structure of mitotic
chromosomes. Molecular cell *81*, 4369-4376 e4363. 10.1016/j.molcel.2021.08.020.

28.   Chen, M., Bell, J.M., Shi, X., Sun, S.Y., Wang, Z., and Ludtke, S.J. (2019). A complete
data processing workflow for cryo-ET and subtomogram averaging. Nature methods *16*,
1161-1168. 10.1038/s41592-019-0591-8.

29.   Liu, Y.T., Zhang, H., Wang, H., Tao, C.L., Bi, G.Q., and Zhou, Z.H. (2022). Isotropic
reconstruction for electron tomography with deep learning. Nature communications *13*,
6482. 10.1038/s41467-022-33957-8.

30.   Eastman, P., Swails, J., Chodera, J.D., McGibbon, R.T., Zhao, Y., Beauchamp, K.A.,
Wang, L.P., Simmonett, A.C., Harrigan, M.P., Stern, C.D., et al. (2017). OpenMM 7:
Rapid development of high performance algorithms for molecular dynamics. PLoS
Comput Biol *13*, e1005659. 10.1371/journal.pcbi.1005659.

31.   Samee, M.A.H., Bruneau, B.G., and Pollard, K.S. (2019). A De Novo Shape Motif
Discovery Algorithm Reveals Preferences of Transcription Factors for DNA Shape
Beyond Sequence Motifs. Cell Syst *8*, 27-42 e26. 10.1016/j.cels.2018.12.001.

32.   ten Heggeler‐Bordier, B., Wahli, W., Adrian, M., Stasiak, A., and Dubochet, J. (1992).
The apical localization of transcribing RNA polymerases on supercoiled DNA prevents
their rotation around the template. The EMBO journal *11*, 667-672.

33.   Zuo, Y., and Steitz, T.A. (2015). Crystal structures of the E. coli transcription initiation
complexes with a complete bubble. Molecular cell *58*, 534-540.
10.1016/j.molcel.2015.03.010.

34.   Narayanan, A., Vago, F.S., Li, K., Qayyum, M.Z., Yernool, D., Jiang, W., and Murakami,
K.S. (2018). Cryo-EM structure of Escherichia coli sigma(70) RNA polymerase and
promoter DNA complex revealed a role of sigma non-conserved region during the open
complex formation. The Journal of biological chemistry *293*, 7367-7375.
10.1074/jbc.RA118.002161.

35.   Fang, C., Li, L., Shen, L., Shi, J., Wang, S., Feng, Y., and Zhang, Y. (2019). Structures
and mechanism of transcription initiation by bacterial ECF factors. Nucleic acids
research *47*, 7094-7104. 10.1093/nar/gkz470.

36.   Braffman, N.R., Piscotta, F.J., Hauver, J., Campbell, E.A., Link, A.J., and Darst, S.A.
(2019). Structural mechanism of transcription inhibition by lasso peptides microcin J25
and capistruin. Proceedings of the National Academy of Sciences of the United States of
America *116*, 1273-1278. 10.1073/pnas.1817352116.

37.   Wood, D.C., and Lebowitz, J. (1984). Effect of supercoiling on the abortive initiation
kinetics of the RNA-I promoter of ColE1 plasmid DNA. The Journal of biological
chemistry *259*, 11184-11187.

38.   Ueshima, R., Fujita, N., and Ishihama, A. (1989). DNA supercoiling and temperature
shift affect the promoter activity of the Escherichia coli rpoH gene encoding the heat-
shock sigma subunit of RNA polymerase. Mol Gen Genet *215*, 185-189.
10.1007/BF00339716.

39.   Henderson, K.L., Felth, L.C., Molzahn, C.M., Shkel, I., Wang, S., Chhabra, M., Ruff, E.F.,
Bieter, L., Kraft, J.E., and Record, M.T., Jr. (2017). Mechanism of transcription initiation
and promoter escape by E. coli RNA polymerase. Proceedings of the National Academy

of Sciences of the United States of America *114*, E3032-E3040.
10.1073/pnas.1618675114.

40.   Finn, R.D., Orlova, E.V., Gowen, B., Buck, M., and van Heel, M. (2000). Escherichia coli
RNA polymerase core and holoenzyme structures. The EMBO journal *19*, 6833-6844.
10.1093/emboj/19.24.6833.

41.   Josephs, E.A., Kocak, D.D., Fitzgibbon, C.J., McMenemy, J., Gersbach, C.A., and
Marszalek, P.E. (2015). Structure and specificity of the RNA-guided endonuclease Cas9
during DNA interrogation, target binding and cleavage. Nucleic acids research *43*, 8924-
8941. 10.1093/nar/gkv892.

42.   Sulc, P., Romano, F., Ouldridge, T.E., Rovigatti, L., Doye, J.P., and Louis, A.A. (2012).
Sequence-dependent thermodynamics of a coarse-grained DNA model. The Journal of
chemical physics *137*, 135101. 10.1063/1.4754132.

43.   Kim, S.H., Ganji, M., Kim, E., van der Torre, J., Abbondanzieri, E., and Dekker, C.
(2018). DNA sequence encodes the position of DNA supercoils. eLife *7*.
10.7554/eLife.36557.

44.   Brogna, S., Sato, T.A., and Rosbash, M. (2002). Ribosome components are associated
with sites of transcription. Molecular cell *10*, 93-104. Doi 10.1016/S1097-2765(02)00565-
8.

45.   Liu, Y., Li, P.Y., Fan, L., and Wu, M.H. (2018). The nuclear transportation routes of
membrane-bound transcription factors. Cell Commun Signal *16*. ARTN 12

10.1186/s12964-018-0224-3.

46.   Gilbert, N., and Allan, J. (2014). Supercoiling in DNA and chromatin. Current Opinion in
Genetics & Development *25*, 15-21. 10.1016/j.gde.2013.10.013.

47.   Worcel, A., Strogatz, S., and Riley, D. (1981). Structure of Chromatin and the Linking
Number of DNA. P Natl Acad Sci-Biol *78*, 1461-1465. DOI 10.1073/pnas.78.3.1461.

48.   Luijsterburg, M.S., White, M.F., van Driel, R., and Dame, R.T. (2008). The Major
Architects of Chromatin: Architectural Proteins in Bacteria, Archaea and Eukaryotes. Crit
Rev Biochem Mol *43*, 393-418. 10.1080/10409230802528488.

49.   Matsumoto, K., and Hirose, S. (2004). Visualization of unconstrained negative supercoils
of DNA on polytene chromosomes of Drosophila. Journal of cell science *117*, 3797-3805.
10.1242/jcs.01225.

50.   Ljungman, M., and Hanawalt, P.C. (1995). Presence of negative torsional tension in the
promoter region of the transcriptionally poised dihydrofolate reductase gene in vivo.
Nucleic acids research *23*, 1782-1789. 10.1093/nar/23.10.1782.

51.   Ding, J.Y., Lee, Y.T., Bhandari, Y., Schwieters, C.D., Fan, L.X., Yu, P., Tarosov, S.G.,
Stagno, J.R., Ma, B.Y., Nussinov, R., et al. (2023). Visualizing RNA conformational and
architectural heterogeneity in solution. Nature communications *14*. 10.1038/s41467-023-
36184-x.

52.   Kang, J.Y., Mishanina, T.V., Bao, Y., Chen, J., Llewellyn, E., Liu, J., Darst, S.A., and
Landick, R. (2023). An ensemble of interconverting conformations of the elemental
paused transcription complex creates regulatory options. Proceedings of the National
Academy of Sciences of the United States of America *120*, e2215945120.
10.1073/pnas.2215945120.

53.   Thomen, P., Bockelmann, U., and Heslot, F. (2002). Rotational drag on DNA: a single
molecule experiment. Physical review letters *88*, 248102.
10.1103/PhysRevLett.88.248102.

54.   Nelson, P. (1999). Transport of torsional stress in DNA. Proceedings of the National
Academy of Sciences of the United States of America *96*, 14342-14347.
10.1073/pnas.96.25.14342.

55. Chatterjee, P., Goldenfeld, N., and Kim, S. (2021). DNA Supercoiling Drives a Transition
between Collective Modes of Gene Synthesis. Physical review letters *127*. ARTN
218101

10.1103/PhysRevLett.127.218101.
56. Horberg, J., and Reymer, A. (2020). Specifically bound BZIP transcription factors
modulate DNA supercoiling transitions. Scientific reports *10*, 18795. 10.1038/s41598-
020-75711-4.
57. Kouzine, F., Gupta, A., Baranello, L., Wojtowicz, D., Ben-Aissa, K., Liu, J.H., Przytycka,
T.M., and Levens, D. (2013). Transcription-dependent dynamic supercoiling is a short-
range genomic force. Nature structural & molecular biology *20*, 396-403.
10.1038/nsmb.2517.
58. Aldag, P., Welzel, F., Jakob, L., Schmidbauer, A., Rutkauskas, M., Fettes, F., Grohmann,
D., and Seidel, R. (2021). Probing the stability of the SpCas9-DNA complex after
cleavage. Nucleic acids research *49*, 12411-12421. 10.1093/nar/gkab1072.
59. Bryant, Z., Stone, M.D., Gore, J., Smith, S.B., Cozzarelli, N.R., and Bustamante, C.
(2003). Structural transitions and elasticity from torque measurements on DNA. Nature
424, 338-341. 10.1038/nature01810.
60. Cheng, B.K., Zhu, C.X., Ji, C.L., Ahumada, A., and Tse-Dinh, Y.C. (2003). Direct
interaction between

RNA polymerase and the zinc ribbon domains of DNA topoisomerase I. Journal of Biological
Chemistry *278*, 30705-30710. 10.1074/jbc.M303403200.
61. Sutormin, D., Galivondzhyan, A., Musharova, O., Travin, D., Rusanova, A., Obraztsova,
K., Borukhov, S., and Severinov, K. (2022). Interaction between transcribing RNA
polymerase and topoisomerase I prevents R-loop formation in E. coli. Nature
communications *13*, 4524. 10.1038/s41467-022-32106-5.
62. Bignaud, A., Cockram, C., Borde, C., Groseille, J., Allemand, E., Thierry, A., Marbouty,
M., Mozziconacci, J., Espéli, O., and Koszul, R. (2024). Transcription-induced domains
form the elementary constraining building blocks of bacterial chromosomes. Nature
structural & molecular biology *31*. 10.1038/s41594-023-01178-2.
63. Marras, S.A., Gold, B., Kramer, F.R., Smith, I., and Tyagi, S. (2004). Real-time
measurement of in vitro transcription. Nucleic acids research *32*, e72.
10.1093/nar/gnh068.
64. Lyubchenko, Y.L., and Shlyakhtenko, L.S. (1997). Visualization of supercoiled DNA with
atomic force microscopy in situ. Proceedings of the National Academy of Sciences of the
United States of America *94*, 496-501. 10.1073/pnas.94.2.496.
65. Boles, T.C., White, J.H., and Cozzarelli, N.R. (1990). Structure of plectonemically
supercoiled DNA. Journal of molecular biology *213*, 931-951. 10.1016/S0022-
2836(05)80272-4.
66. Arman, F., Helmut, S., and Reza, E.M. (2015). Molecular Dynamics Simulation of
Supercoiled DNA Rings.
67. Huang, J., Schlick, T., and Vologodskii, A. (2001). Dynamics of site juxtaposition in
supercoiled DNA. Proceedings of the National Academy of Sciences of the United
States of America *98*, 968-973. 10.1073/pnas.98.3.968.
68. Forquet, R., Nasser, W., Reverchon, S., and Meyer, S. (2022). Quantitative contribution
of the spacer length in the supercoiling-sensitivity of bacterial promoters. Nucleic acids
research *50*, 7287-7297. 10.1093/nar/gkac579.
69. Bentin, T., and Nielsen, P.E. (2002). In vitro transcription of a torsionally constrained
template. Nucleic acids research *30*, 803-809. 10.1093/nar/30.3.803.

70. Teves, S.S., and Henikoff, S. (2014). Transcription-generated torsional stress
destabilizes nucleosomes. Nature structural & molecular biology *21*, 88-94.
10.1038/nsmb.2723.
71. Fosado, Y.A.G., Michieletto, D., Brackley, C.A., and Marenduzzo, D. (2021).
Nonequilibrium dynamics and action at a distance in transcriptionally driven DNA
supercoiling. Proceedings of the National Academy of Sciences of the United States of
America *118*. ARTN e1905215118

10.1073/pnas.1905215118.
72. Leng, F.F., and McMacken, R. (2002). Potent stimulation of transcription-coupled DNA
supercoiling by sequence-specific DNA-binding proteins. Proceedings of the National
Academy of Sciences of the United States of America *99*, 9139-9144.
10.1073/pnas.142002099.
73. Brahmachari, S., Tripathi, S., Onuchic, J.N., and Levine, H. (2024). Nucleosomes play a
dual role in regulating transcription dynamics. Proceedings of the National Academy of
Sciences of the United States of America *121*, e2319772121. 10.1073/pnas.2319772121.
74. Golding, I., Paulsson, J., Zawilski, S.M., and Cox, E.C. (2005). Real-time kinetics of
gene activity in individual bacteria. Cell *123*, 1025-1036. 10.1016/j.cell.2005.09.031.
75. So, L.H., Ghosh, A., Zong, C., Sepulveda, L.A., Segev, R., and Golding, I. (2011).
General properties of transcriptional time series in Escherichia coli. Nature genetics *43*,
554-560. 10.1038/ng.821.
76. Jinek, M., Chylinski, K., Fonfara, I., Hauer, M., Doudna, J.A., and Charpentier, E. (2012).
A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity.
Science *337*, 816-821. 10.1126/science.1225829.
77. Bizard, A., Garnier, F., and Nadal, M. (2011). TopR2, the second reverse gyrase of
Sulfolobus solfataricus, exhibits unusual properties. Journal of molecular biology *408*,
839-849. 10.1016/j.jmb.2011.03.030.
78. Aquilina, M., and Dunn, K.E. (2023). Multiplexed Label-Free Biomarker Detection by
Targeted Disassembly of Variable-Length DNA Payload Chains. Anal Sens *3*. ARTN
e202200082

10.1002/anse.202200082.
79. Schorb, M., Haberbosch, I., Hagen, W.J.H., Schwab, Y., and Mastronarde, D.N. (2019).
Software tools for automated transmission electron microscopy. Nat Methods *16*, 471-
477. 10.1038/s41592-019-0396-9.
80. Zheng, S.Q., Palovcak, E., Armache, J.P., Verba, K.A., Cheng, Y., and Agard, D.A.
(2017). MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
electron microscopy. Nat Methods *14*, 331-332. 10.1038/nmeth.4193.
81. Buchholz, T.O., Krull, A., Shahidi, R., Pigino, G., Jekely, G., and Jug, F. (2019). Content-
aware image restoration for electron microscopy. Methods Cell Biol *152*, 277-289.
10.1016/bs.mcb.2019.05.001.
82. Zhang, K. (2016). Gctf: Real-time CTF determination and correction. J Struct Biol *193*, 1-
12. 10.1016/j.jsb.2015.11.003.
83. Kremer, J.R., Mastronarde, D.N., and McIntosh, J.R. (1996). Computer visualization of
three-dimensional image data using IMOD. J Struct Biol *116*, 71-76.
10.1006/jsbi.1996.0013.
84. Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C.,
and Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory research
and analysis. J Comput Chem *25*, 1605-1612. 10.1002/jcc.20084.

85.   Ludtke, S.J., Baldwin, P.R., and Chiu, W. (1999). EMAN: semiautomated software for
high-resolution single-particle reconstructions. J Struct Biol *128*, 82-97.
10.1006/jsbi.1999.4174.
86.   Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC:
algorithms for rapid unsupervised cryo-EM structure determination. Nat Methods *14*,
290-296. 10.1038/nmeth.4169.
87.   Sanchez-Garcia, R., Gomez-Blanco, J., Cuervo, A., Carazo, J.M., Sorzano, C.O.S., and
Vargas, J. (2021). DeepEMhancer: a deep learning solution for cryo-EM volume post-
processing. Commun Biol *4*, 874. 10.1038/s42003-021-02399-1.
88.   Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H.,
and Ferrin, T.E. (2023). UCSF ChimeraX: Tools for structure building and analysis.
Protein Sci *32*, e4792. 10.1002/pro.4792.
89.   Liebschner, D., Afonine, P.V., Baker, M.L., Bunkoczi, G., Chen, V.B., Croll, T.I., Hintze,
B., Hung, L.W., Jain, S., McCoy, A.J., et al. (2019). Macromolecular structure
determination using X-rays, neutrons and electrons: recent developments in Phenix.
Acta Crystallogr D Struct Biol *75*, 861-877. 10.1107/S2059798319011471.
90.   Croll, T.I. (2018). ISOLDE: a physically realistic environment for model building into low-
resolution electron-density maps. Acta Crystallogr D Struct Biol *74*, 519-530.
10.1107/S2059798318002425.
91.   Chen, V.B., Arendall, W.B., 3rd, Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral,
G.J., Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: all-atom
structure validation for macromolecular crystallography. Acta crystallographica. Section
D, Biological crystallography *66*, 12-21. 10.1107/S0907444909042073.
92.   Klenin, K., and Langowski, J. (2000). Computation of writhe in modeling of supercoiled
DNA. Biopolymers *54*, 307-317. 10.1002/1097-0282(20001015)54:5<307::AID-
BIP20>3.0.CO;2-Y.
93.   Johnson, R.A. (1929). Modern geometry: an elementary treatise on the geometry of the
triangle and the circle (Houghton, Mifflin Company).
94.   Burns, J.R., Seifert, A., Fertig, N., and Howorka, S. (2016). A biomimetic DNA-based
channel for the ligand-controlled transport of charged molecular cargo across a
biological membrane. Nat Nanotechnol *11*, 152-156.
95.   Suma, A., Poppleton, E., Matthies, M., Sulc, P., Romano, F., Louis, A.A., Doye, J.P.K.,
Micheletti, C., and Rovigatti, L. (2019). TacoxDNA: A user-friendly web server for
simulations of complex DNA structures, from single strands to origami. Journal of
computational chemistry *40*, 2586-2595. 10.1002/jcc.26029.
96.   Wright, D.J., King, K., and Modrich, P. (1989). The Negative Charge of Glu-111 Is
Required to Activate the Cleavage Center of Ecori Endonuclease. J Biol Chem *264*,
11816-11821.
97.   Weigert, M., Schmidt, U., Boothe, T., Muller, A., Dibrov, A., Jain, A., Wilhelm, B.,
Schmidt, D., Broaddus, C., Culley, S., et al. (2018). Content-aware image restoration:
pushing the limits of fluorescence microscopy. Nat Methods *15*, 1090-1097.
10.1038/s41592-018-0216-7.
98.   Lee, T.C., Kashyap, R.L., and Chu, C.N. (1994). Building Skeleton Models Via 3-D
Medial Surface Axis Thinning Algorithms. Cvgip-Graph Model Im *56*, 462-478. DOI
10.1006/cgip.1994.1042.

1638

Figure 1: 3D reconstruction of an individual negative-supercoiled (–sc) DNA plasmid
(**A**) pUC19 plasmid construct features a T7A1 promoter (green), a U-less stalling site (black), and a transcriptional region (pink). (**B**) 2D gel electrophoresis of the –sc pUC19-T7A1U plasmid and ΔLk quantification. (**C**) AFM images of pUC19 plasmids showing ΔLk variants. (**D**) Cryo-EM images of –sc plasmid (orange arrowhead) highlighting sharp DNA kinks (red circle). (**E**) Cryo-ET per-particle 3D reconstruction steps of a representative –sc plasmid. (**F**) Zoom-in view of the final map and model from E, showing two intertwined DNA segments (yellow and green) with the DNA crossing marked by arrowheads in the view. (**G**) Different 3D views influence 2D assessments of DNA curvature (panels I, II) and spacing (panel III). (**H**) Color-coded map of the plasmid in F, with high-curvature regions in red and low-curvature regions in blue. (**I**) Tracing of the DNA curvature in H along the circular DNA plasmid.

Figure 2: Conformational dynamics and regulatory effect of apically bound proteins on DNA supercoiling
(**A**) Cryo-ET (3D maps and models) of the −sc plasmids bound with stalled RNAP (sTEC), with plasmid construct design in circle and plectoneme apex modeling steps on top right. RNAPs in sTECs are marked by purple arrowheads in the bottom particle collection (**B**) The superimposition of all bound RNAPs reveals apical DNA dynamics. The dashed vector indicates RNAP orientation. (**C**) The superimposition of all apical DNA segments shows the RNAP orientation dynamics (purple arrowheads). (**D**) Electrophoresis gel shows reduced abortive transcript levels on −sc DNA compared to nicked DNA. (**E**) Schematic highlighting DNA geometry conversion in TEC on −sc DNA (top), and in TIC (bottom left) and TEC (bottom right) on linear DNA templates. (**F**) Cryo-ET of dCas9 (orange) bound to −sc plasmids. (**G and H**) Superimposition of all bound dCas9s and all apical DNA segments, respectively. (**I**) Schematic highlights dCas9's smaller pocket curvature compared to that of RNAP, reshaping bound plasmids with enlarged distal ends and tightly intertwined body.

Figure 3: Interplay between apically bound RNAP and –sc DNA during non-equilibrium transcription
(**A**) Cryo-ET of TEC on –sc templates after 10 minutes of transcription, with RNAP and large DNA loops indicated by solid and hollow purple arrowheads, respectively. (**B**) Alignment and superimposition of all apical DNA segments. (**C**) Z-dimensional slices (10 nm-thickness) of sub-tomograms comparing transcribed (top) and stalled (bottom) RNAPs, overlaid with models. (**D**) Plasmid apex number distribution; N=84. (**E-F**) Single-round *in vitro* transcription of RNAP transcription on nicked and –sc templates over time, with quantification of their pause release kinetics, respectively. (**G**) Single-particle cryo-EM reconstruction of the major RNAP class, TEC1–SC on –sc DNA, overlaid with the non-swiveled 6RH3 model. (**H**) Comparison of TEC1–SC swivel angle to reported RNAP structures from active transcription to various paused states. (**I**) TEC1–SC active-site DNA orientation and subunit swivel direction. (**J**) Schematic illustration of transcription-induced new plectoneme formation on –sc plasmid. DNA translational and rotational motion is indicated by green arrows, with orange color denoting drags.

Figure 4: Plectoneme apical confinement promotes topological domains and multi-RNAP transcription
(**A**) Cryo-ET of –sc plasmids simultaneously bound by stalled RNAP (purple) and dCas9 (orange). Representative particles show RNAP and dCas9 co-localized at the apex (top left) or off-apex (top right), both consistent with the opposing-position construct design (circular plot). (**B**) Off-apex particle ratios for RNAP and dCas9 across conditions; N=301 (**C**) Ratio of plasmids bound by multiple RNAPs or dCas9; N=324. (**D**) Cryo-ET of –sc plasmid resuming transcription for 10 minutes in the presence of torsional roadblock dCas9 (**E**) Schematics illustrating RNAP transcription-induced topological domains and supercoiling rearrangement in the presence of dCas9. (**F**) Quantification of plasmids' apex number; N=162. (**G**) Assessment of RNAP pause release in the presence of dCas9 during transcription via electrophoresis. The RNA band intensities were quantified at three strong pausing sites and run-off site.

Figure 5: TopI releases RNAP from apical constraint during transcription
(**A**) Schematic showing TopI bound to DNA or coupled with RNAP. (**B**) Assessment of RNAP pause release in the presence of TopI from single-round *in vitro* transcription assay. (**C**) Cryo-ET z-dimensional slice (50 nm-thickness) of sample after 10 minutes of transcription in the presence of TopI. (**D**) Cryo-ET of the plasmid particles in the presence of RNAP and TopI after 10 minutes of of transcription (TEC-TopI). Apical and non-apical RNAP binding are indicated by purple and green arrowheads, respectively. (**E**) Representative TEC–TopI complexes showing RNAPs escape from the apices. (**F-G**) Quantification of the non-apical RNAP ratio (N=90) and plasmid's apex number (N=85), respectively. (**H**) Superimposition of all bound RNAPs (top panel) and all apical DNA segments (bottom panel). (**I-J**) DNA supercoiling relaxation assays mediated by TopI and hTopIB in the presence of RNAP, respectively, with ΔLk quantification shown on the right.

Figure 6: Tandem promoters enhance topological domain formation and transcription compared to opposing promoters.
(**A**) Opposing dual-T7A1U promoter construct (left) driving transcription of AmpR and mEGFP. Circular plasmid layout (middle) with rim color-matched to the 3D model (right). Arrowheads indicate RNAP transcription direction; circles mark apical sites. Inner sectors represent individual plectonemes (I–III), color-coded by writhe density (blue–pink–red). (**B**) Tandem promoter construct, identical to A, but with AmpR flipped. (**C-D**) Representative TECs from opposing and tandem constructs, ordered by RNAP count. (**E**) Plasmid distribution by bound RNAP number and apex count; N=90. (**F-H**) Quantification of plasmid Rg (N=67), correlation between RNAP number and apex count (N=206), and ratio of RNAP located off-apex; N=232. (**I**) Plectoneme writhe density distribution vs. bound RNAP number; N=230. (**J**) Distance distribution of opposing RNAPs (left) and tandem RNAPs (right) on the circular plasmid layout. Shaded regions indicate less likely RNAP separations, attributed to the non-specifically bound RNAP population; N=212. (**K**) Quantification of *in vitro* gene expression by molecular beacon assay. (**L-M**) Imaging and quantification of in vivo GFP gene expression in *E. coli* MG1655 carrying either plasmid of tandem or opposing constructs; N=529.

Figure 7. Revisited twin-supercoiled domain model and its connection to transcriptional bursting.
(**A**) Comparison of classical twin-supercoiled domain and a revisited model considering the apical localization of RNAP in −sc DNA. (**B**) Summary model connecting transcriptional induced torsional stress and transcriptional bursting in −sc DNA. (See Discussion for further description of the model).

Figure S1: Structural analysis of –sc plasmid under low salt and high salt conditions, related to Figure 1
(**A-B**) Representative cryo-ET map and model of the –sc plasmid (ΔLk centered on -15) under low salt (LS) and high salt (HS) conditions, respectively. Two exemplary particles (unbranched and branch) are displayed on the top panel, with their collection presented on the bottom, featuring writhe numbers. (**C-D**) Statistical analysis of plasmid radius of gyration (Rg) and writhe number using violin plots, respectively; N=64. (**E**) Schematic of inter-DNA spacing quantification (black bars) along the plectoneme axis (red dots). The inter-DNA spacing of the plectoneme was plotted from the plasmid apex toward the intermediate region; N=158 (**F**) Schematic illustrating the apical and intermediate regions of a plectoneme, accompanied by DNA curvature quantification. (**G**) Statistical analysis of DNA curvature near the apical and intermediate regions of the plectoneme under LS and HS conditions; N=158 (**H**) Distribution of the number of apices of the plasmid in LS and HS conditions; N=64. Statistics are calculated using a Mann-Whitney test, where *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; and ns, not significant.

Figure S2: Sub-tomogram averaging analysis of apically bound RNAP and dCas9, related to Figure 2

(**A-B**) Two orthogonal views of the sub-tomogram averaged RNAP map docked with PDB structure 6ALH. Averaging all RNAPs selected from the tomogram (N=1781) yielded a 14.1 Å resolution map (top), while using only plasmid-bound RNAPs after inspection (N=232) resulted in a 17.7 Å resolution map (bottom). (**C**) Superimposition of consensus TIC upstream DNA (black helix) and the TIC σ-factor (red) from PDBs 4YLN, 6CA0, 6JBQ, and 6N60 onto the apically stalled TEC on –sc DNA reveals a ~170° upstream DNA bend. (**D-E**) Sub-tomogram averaged dCas9 map docked with PDB structure 6O0X. Averaging all dCas9 particles selected from the tomogram (N=875) yielded a 17.6 Å resolution map (top), while the plasmid-bound subset (N=116) achieved a resolution of 18.9 Å (bottom). Resolutions were estimated by measuring the Fourier shell correlation (FSC) between two independently determined half-maps at 0.143. (**F**) Statistical analysis of global plasmid Rg and writhe number (N=90), along with local DNA curvature and inter-DNA spacing at apical sites (N=207), for plasmid only, plasmid with stalled RNAP, and plasmid with dCas9, respectively. Statistics are calculated using a Mann-Whitney test, where *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; and ns, not significant.

Figure S3: Quantification of plasmid geometry upon RNAP/dCas9 binding, related to Figure 3

(**A**) Generation of –sc DNA minicircles with a target site subjected to varying levels of constraint, defined by spring distances of 8, 6.4, and 4.8 nm, to modulate DNA relaxation. (**B**) Quantification of relaxed plasmid writhe, apical DNA curvature, and their correlation following MD simulation (sampled from frame 100 onward after relaxation; three replicates). (**C**) Predicted DNA plectoneme formation loci along the pUC19-T7A1U sequence. A control was generated using a polyA–Widom601–polyA sequence. (**D**) Overlay of apically bound RNAPs in TECs with RNA protrusion directions indicated by pink arrows. (**E**) Statistical analysis of global plasmid Rg and writhe number (N=79), along with local DNA curvature and inter-DNA spacing at apical sites (N=192) for sTECs and TECs. (**F**) Plectoneme branches were segmented (I), and their superhelical axes (II) and contour lengths (III) were determined. Writhe density was calculated as writhe number divided by segment contour length. Bottom right: writhe density distribution; N=187. (**G**) Plot of plectoneme branch length versus distance from the branch to the apical RNAP; N=38. Statistics are calculated using a Mann-Whitney test, where *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; and ns, not significant.

Figure S4. Cryo-EM analysis of the transcription elongation complex (TEC) on a –sc plasmid template, related to Figure 3

(**A**) Workflow of single-particle analysis and 3D reconstruction of TECs. Three final maps were obtained: the first (TEC1–SC, global resolution of 2.95 Å from 46,743 particles) and the second (TEC2–SC, global resolution of 2.94 Å from 54,869 particles) show particles with downstream DNA protrusion. The third class was not further analyzed due to the absence of DNA density, indicating likely unbound RNAPs. (**B**) Representative reference-free 2D class averages of TEC particles, displaying various TECs orientations. (**C**) Representative cryo-EM micrograph of TEC particles assembled on negatively supercoiled pUC19-T7A1U plasmid in vitreous ice. (**D**) Fourier shell correlation (FSC) curves between independently reconstructed even–odd maps. Resolution was estimated at the 0.143 FSC cutoff. (**E**) Cryo-EM density maps of TEC1–SC (left) and TEC2–SC (right), color-coded by local resolution, ranging from 3 Å (red) to yellow to 7 Å (blue). (**F**) Examination of RNAP active sites shows that both TEC1–SC and TEC2–SC are in a post-translocated state, with the trigger helix unfolded and the NTP absent.

Figure S5: Apical binding of dCas9 hinders DNA rotation, related to Figure 4
(**A**) DNA supercoiling relaxation assay in the absence (top) and presence (bottom) of a dCas9. Right panel: quantification of ΔLk changes over time following TopI addition. (**B**) Control experiment of A, with TopI replaced by hTopIB for DNA supercoiling relaxation. (**C**) Coarse-grained MD simulations of 525-bp –sc (ΔLk = –5) minicircle plasmid relaxation dynamics, initiated with different nick locations. Simulations were conducted without apical confinement (Model I) and with apical restriction (Models II–IV). Topological parameters (ΔTw, Wr, and ΔLk) are shown in rows 1–3 for each model. Each simulation was performed in triplicate, with averages indicated by black dashed lines. (**D**) Statistical analysis of plectoneme branch writhe density distribution; N=226. (**E**) Superimposition of apical DNA segments illustrating the dynamics of stalled RNAP and dCas9 when co-present on the plasmid.

Figure S6: Quantification of plasmid morphology with co-presence of RNAP and dCas9, and of RNAP and TopI, related to Figures 4 and 5

(**A**) Statistical analysis of global plasmid Rg and writhe number (N=107), along with local DNA curvature and inter-DNA spacing at apical sites (N=270) for sTEC-Cas and TEC-Cas particles. (**B**) Statistical comparison of the plectoneme writhe density distribution; N=198. (**C**) Mapping all bound RNAPs on pUC19-T7A1U templates using dCas9 as fiducial. For visualization, different plasmids and their associated RNAPs are displayed at distinct radii in the polar plot. RNAP on the transcriptional region, oriented downstream, is marked in pink; RNAP on the non-transcriptional region or pointing upstream, is colored in tan and blue, respectively; N=89 (**D–E**) Electrophoresis assay of RNAP transcription on –sc pUC19-T7A1U templates in the presence of dCas9 and in the presence of TopI over time, respectively. (**F**) Same statistical measurements as in (A), performed on TEC-TopI particles (90 particles and 212 apices total) . All statistics are calculated using a Mann-Whitney test, where *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001; and ns, not significant.

Figure S7: Top I and RNAP coupling slows supercoiling relaxation, and particle collections of dual-promoter TECs, related to Figures 5 and 6.

(**A**) 2D electrophoresis gel of –sc pUC19-T7A1U plasmid in the presence of RNAP and TopI, with quantification of the dominant ΔLk value. (**B**) 1D gel-based DNA supercoiling relaxation assay of plasmid in the presence of RNAP and a large excess of TopI. (**C**) DNA supercoiling relaxation assay of plasmid in the presence of RNAP core (lacking σ and unable to form stable transcription bubbles). (**D**) DNA supercoiling relaxation assay of plasmid in the presence of dEcoRI. 1D gel-based linking number quantification is shown in the right panel for B-D. (**E**) Particle collections of opposing-promoter TECs and tandem-promoter TECs, ordered by the number of bound RNAPs. Each dual-promoter TEC is also shown with its corresponding circular plasmid layout, annotated in red to indicate plectonemes extending into the positive supercoiling regime.