Manuscript information

Title

Crystal structure of *E. coli* Nissle 1917 flagellin reveals novel features that modulate

bacterial motility but not TLR5 recognition

Authors

Johanna Jakob[1,2,3,$], Michael B. Braun[4], Katharina Hipp[5], Iris Koch[5], Gaopeng Li[1], Pascal

Felgner[6], Maria Jose Giralt Zuniga[7,8], Hannah Raasch[7], Corinna Ghering-Khav[9,10], Andras

Szolek[1], Abdelhakim Boudrioua[11], Thomas Hagemann[2], Samuel Wagner[2,3,11,12], Thilo

Stehle[3,4], Liudmila Andreeva[9,10], Marc Erhardt[7,8], Michael Hensel[6], Julia-Stefanie Frick[2,11,§] ,

Alexander N. R. Weber[1,3,9,13,#]

Author affiliations

[1]Institute of Immunology, Department of Innate Immunity, University of Tübingen, Auf der

Morgenstelle 15, 72076 Tübingen, Germany

[2]Institute of Medical Microbiology and Hygiene, University Hospital Tübingen, Elfriede-

Aulhorn Straße 6, 72076 Tübingen, Germany

[3]Cluster of Excellence (EXC 2124). "Controlling Microbes to Fight Infections", University of

Tübingen, Tübingen, Germany

[4]Interfaculty Institute of Biochemistry, University of Tübingen, Auf der Morgenstelle 34, 72076

Tübingen, Germany

[5]Max Planck Institute for Biology Tübingen, Electron Microscopy Facility, Max-Planck-Ring 5,

72076 Tübingen, Germany

[6]Division of Microbiology, University of Osnabrück, Barbarastr. 11, 49076 Osnabrueck,

Germany

[7]Department of Molecular Microbiology, Institute of Biology, Humboldt-University Berlin,

Philippstr. 13, 10115 Berlin, Germany

[8]Max Planck Unit for the Science of Pathogens, Chariteplatz 1, 10117 Berlin, Germany

[9]iFIT – Cluster of excellence (EXC 2180). "Image-Guided and Functionally Instructed Tumor

Therapies", University of Tübingen, Germany

[10] Department of Medical Oncology and Pneumology (Internal Medicine VIII), University

Hospital Tübingen, Röntgenweg 11, 72076 Tübingen, Germany

[11]Interfaculty Institute of Microbiology and Infection Medicine (IMIT), Section of Cellular and

Molecular Microbiology, University of Tübingen, Elfriede-Aulhorn Straße 6, 72076 Tübingen,

Germany

[12]German Center for Infection Research (DZIF), Partner Site Tübingen, Tübingen, Germany

[13]German Cancer Consortium (DKTK), DKTK Partner Site Tübingen, 72076, Tübingen,

Germany.

[#] corresponding author: Prof. PhD Alexander N.R. Weber. Institute of Immunology,

Department of Innate Immunity, University of Tübingen, Auf der Morgenstelle 15, 72076

Tübingen, Germany; Tel: +49 7071 29 87623; email: alexander.weber@uni-tuebingen.de

[$] current address: Department of Dermatology, University Hospital Tübingen,

Liebermeisterstr. 20, 72076 Tübingen, Germany

[§] current address: MVZ Labor Ludwigsburg SE & Co. eGbR, Wernerstr. 33, 71636

Ludwigsburg

Short title

E. coli Nissle 1917 flagellin structure and motility

Address for correspondence

Prof. PhD Alexander N.R. Weber. Institute of Immunology, Department of Innate Immunity,

University of Tübingen, Auf der Morgenstelle 15, 72076 Tübingen, Germany; Tel: +49 7071

29 87623; email: alexander.weber@uni-tuebingen.de.

Abstract

The probiotic *E. coli* Nissle 1917 (EcN) strain is known to promote intestinal homeostasis via

flagellin, the protomer of its motility apparatus, the flagellum. The flagellin of EcN shows

atypical features, namely a hypervariable region (HVR), whose structure and significance

have remained elusive. We therefore determined the crystal structure of the *E. coli* Nissle

1917 flagellin FliC at a resolution of 1.2 Å which revealed an unusual domain architecture:

the canonical D1 domain was found connected by an extended linker to an extensive HVR

whose D2, D3 and D4 domains form an outer domain (OD) which surrounds the filament

core comprised of conserved domains D0-D1. Using both recombinant proteins and gene-

edited EcN strains expressing mutant flagellins, the functional requirement for these unique

features was subsequently studied for effects on immune recognition on intestinal epithelial

and immune cells, as well as on flagellar protein expression, assembly and bacterial motility.

While human and mouse TLR5 immune recognition of flagellar proteins or intact bacteria

was only moderately affected by removal of linker or D4, especially linker removal reduced

protein stability and bacterial motility in both soft agar and liquid media swimming assays.

Interestingly, depending on the environment, D4 or HVR removal had different effects on

motility and surface structure. Finally, a site-directed mutagenesis approach highlighted that

loss of TLR5 recognition strictly entails loss of motility but not vice versa. Our data indicate

that specific HVRs/OD might be relevant for motility of *E. coli* Nissle 1917 in specialized

environments, but not for immune recognition. Moreover, we find mutational tolerance is

greater for immune recognition than for motility, providing new insights into bacterial

adaptation to the host environment.

Keywords

Flagellin, flagellum, *E. coli* Nissle, Inflammatory Bowel Disease, TLR5, innate immunity

<u>Introduction</u>

Flagella-mediated motility is a common and essential fitness trait in many microbes and

enables bacteria to access nutrients or certain niches, and form biofilms while avoiding

toxicity [1]. The complex bacterial flagellar apparatus hinges on the protein flagellin as its

most abundant building block or protomer [2]. In *Escherichia coli*, an estimated 20,000-

30,000 subunits of *fliC*-encoded flagellin make up each of the several 10 µm-long and 0.02

µm-wide flagella [3]. The inner core of this assembly comprises two sheaths formed by the

D0 and D1 domains of flagellin. Central and outer regions of the flagellum are formed by

consecutive additional flagellin domains, e.g., D2, D3 and D4, originating from a single

polypeptide that folds back on itself in an antiparallel fashion, i.e., N- and C-termini join in the

D0 domain as two antiparallel alpha helices. Whereas D0 and D1 are highly conserved

across bacterial species, the other domains vary in number and residue length and are thus

referred to as 'hypervariable region' (HVR). Especially regions beyond the D3 domain show

great sequence and length variability: For example, *Salmonella enterica ssp. enterica*

serovar Typhimurium (ST) FliC contains only domains D2 and D3 [4, 5], whereas *E. coli*

Nissle 1917 (EcN), a well-known probiotic strain of *E. coli,* was found to contain a long HVR

of 326 amino acids with unknown structure [6]. Even for different E. coli strains, flagellin

sequence varies greatly (Fig. S1), with sequence identity in the HVR dropping as low as 29%

(Fig. S2).

In a vertebrate host, flagellin also influences bacterial immunogenicity, serving as a microbe-

associated molecular pattern (MAMP) sensed by the innate immune system via a pattern

recognition receptor (PRR). Specifically, residues in the D1 domain are sensed by the PRR

Toll-like receptor 5 (TLR5) [7]. Based on the crystal structure of FliC from *Salmonella*

enterica ssp. enterica serovar Dublin in complex with zebrafish TLR5 [8] and synthetic

biology studies [9], the FliC α-helices in the D1 region bind to TLR5 laterally, in the primary

interface between the leucine-rich repeat (LRR)-NT and LRR10 regions, with LRR9

protruding out as a loop thereby forming a binding cavity for FliC. A second interface involves

binding of the C-terminal αND1b to LRR12/13 of a second TLR5 molecule stabilizing

homodimer formation of two TLR5:flagellin complexes in a 2:2 quaternary complex [8]. For

signaling, a third binding interface is required that involves the D0 domain [10]. However, this

seems to be unimportant for binding. Recent structural comparisons of fish and mammalian

TLR5 indicate that this complex is likely an inhibitory arrangement [11, 12], so that the

structure of an active ligand-receptor complex for the mammalian system has yet to be

resolved. Mutagenesis studies indicate that residues 89-96, especially R90, and I411 within

the *Salmonella* FliC D1 domain interact with A268 in human TLR5 [10, 13]. Binding of TLR5,

which is expressed mainly on the basolateral sides of the intestinal epithelium, leads to

mitogen-activated protein kinase and MyD88-dependent activation of NF-κB, which

culminates in the secretion of pro-inflammatory cytokines (e.g., IL-6 and IL-8) and

antimicrobial peptides (e.g., human beta defensin 2, hBD2) [14-16]. Mucosal immune cells

also bear TLR5 and may respond to flagellin in case barrier integrity is compromised [17].

This is the case in inflammatory bowel disease such as Crohn's disease [18, 19]. Recent

work from our laboratories showed that flagellin from probiotic bacteria like EcN may have a

regulatory effect on intestinal inflammation, specifically in a DSS-mediated experimental

colitis mouse model [6]. We observed that EcN, but not other commensal *E. coli* strains,

ameliorates DSS colitis in a flagellin- and IL-22-dependent manner. Paradoxically, these

effects required TLR5 in conventional dendritic cells and yet involved the HVR, i.e. a region

outside the classical TLR5 recognition motif [6]. Deletion of flagellin or parts of the HVR

caused a loss of the probiotic effects. This begged the question whether structural features of

the EcN flagellin might modulate TLR5 recognition and other aspects of EcN fitness, e.g.,

motility, a key property of bacteria navigating the intestinal space, including the mucus layer

separating intestinal content from the (TLR5-expressing) intestinal barrier. The atomic

structure of EcN FliC has not been determined, but recent structural studies and its length

suggest it might adopt a fold identified in certain pathogenic and soil bacteria recently termed

'outer domain sheath' (ODS), including D2, D3 and an additional D4 domain [20].

Thus, although flagellin structure seems to have a profound effect on immune modulation

and motility, the structure-activity relationships are not well understood, especially when

considering potential "trade-offs" between immune recognition and motility. Sequence

variation that would enable immune evasion may be counteracted by the need to retain

motility; thus, changes at TLR5 binding interfaces are balanced against strong intradomain

(e.g. between alpha-helices in D0 and D1) as well as homodimer contacts to guarantee

flagellin folding and flagellar assembly [21]. One study recently investigated the effects of

mutations in ST FliC on recognition by human and mouse TLR5 [22]. Interestingly, in an *in*

vitro cell activation assay, recognition by human but not mouse TLR5 could be completely

evaded by a variant of ST FliC carrying a single point mutation (R90A). In comparison,

strains expressing ST FliC R90A retained 80% bacterial motility. In general, human TLR5

recognition was more sensitive to mutation compared to motility, whereas mouse TLR5

recognition was more robust and motility was lost more readily. In mice, but not in humans,

Salmonella would therefore struggle to avoid TLR5 recognition while retaining motility.

Another recent study focusing on human TLR5 found that a high proportion of flagellins in the

human microbiota, e.g., *Roseburia hominis* FlaB, are 'silent' because they bind to TLR5 but

fail to activate downstream signaling [23]. Engraftment of ST FliC D0 restored TLR5

activation in purified *Roseburia* FlaB *in vitro*, highlighting the contribution of this domain to

TLR5 activation [10, 23]. However, the stability and ability of these hybrid flagellins to form

flagella and the resulting effects on motility were not assessed. Thus, it is unknown whether

the 'silent' D0 domain of *Roseburia hominis* FlaB allows for motility in other flagellated

bacteria, such as *E. coli* or ST.

To gain a deeper understanding of the structural basis for the probiotic effect of the EcN

flagellin and the relationships between flagellin structure, TLR5 recognition and motility, we

first solved the crystal structure of EcN FliC and discovered that the HVR sequence adopts

another well resolved domain, D4. Moreover, and unexpectedly, D1 and D2 were separated

by an extended antiparallel β-sheet linker. We examined the effects of these unique

structural features on immune recognition by human and mouse TLR5 as well as motility.

This analysis showed that both unique features are redundant for TLR5 recognition and

secretion of flagellin from bacteria. However, both linker and D4 domain emerged as

essential for optimal bacterial motility in certain environments. EcN D4 appeared to represent

a novel OD. Although deletion of the linker and the D4 compromised motility, a complete

removal of the HVR was, surprisingly, tolerated, but depending on context. Our data indicate

that under *in vitro* conditions, ODSs are not essential for motility *per se*, but our study

highlights the unique D1-D2 linker in the flagellar EcN core as a critical context-dependent

determinant of bacterial motility. Moreover, using mutagenesis studies we found that loss of

TLR5 recognition strictly entails loss of motility but not vice versa, providing new insights into

host-bacterial   adaptation.

Results

The crystal structure of the atypical EcN flagellin contains an extensive ODS with a

rigid linker

To gain insight into the structure of EcN flagellin, especially its HVR, we expressed different

constructs of the entire EcN FliC in *E. coli* BL3 cells as His-tagged fusion proteins (see

Methods, Fig. S3 and Fig. S4) for subsequent crystallographic analysis. Whereas full length

and a D0 deletion construct (ΔD0, Δ1-47, Δ555-595) did not yield crystals, a construct with

additional deletion of D1 (ΔD0+D1, Δ1-175, Δ499-595) could be crystallized and diffracted.

As molecular replacements (ST FliC PDB 1IO1), as well as native and seleno-methionine

phasing did not yield results, heavy atom (Se-Urea) soaking was performed, allowing for

solving the native structure at 1.65 Å resolution. As the D1 is critical for TLR5 interaction,

another attempt to gain D1 information was made using additional shortened constructs, FliC

D1+2 (Δ1-61, Δ208-451, Δ542-595) and FliC D12(Δ1-61, Δ208-451<S, Δ555-595), which

resulted in crystals that diffracted to a resolution of 2.03 Å and 1.75 Å. Since the D2 domain

was present in both sets of constructs, it was superimposed with minimal differences (RMSD

0.4 Å over 275 aa), so that a combined model of the D1 to D4 domain was built (Fig. 1A).

The structure showed several typical features: firstly, a helical D1 domain with anti-parallel

coiled-coil structure that aligned well with the ST FliC (PDB 1IO1, Fig. S4E, RMSD 0.57 Å);

secondly, a globular D2 with a three stranded antiparallel β-sheet and one α-helix forming an

αβββ-sandwich; and thirdly, a globular D3 domain consisting of two αββ-sandwiches where

the four β-strands form a β-sheet with the two α-helices laying on the same side.

Surprisingly, another domain, D4, with a similar αββ-sandwich topology but 90° rotated β-

sheet orientation (Fig. 1B), was discovered. The tip of D4 showed an unusual fold, where

three strands form an antiparallel triangle, with β-sheet-like interactions, termed here β-

triangle. Thus, the HVR sequence previously described [6] contributed to a canonical D3 and

to a unique D4 domain. Compared to ST FliC, the positioning of D1-D3 deviated strongly.

This was primarily due to a linker separating D1 from D2, resulting in minimal contacts

between the two domains (Fig. 1C). Instead, in ST FliC, the transition between D1 and D2 is

seamless and generates numerous interactions between the D1 and D2 domain. The length

of the linker is about 30 Å, with the first 21 Å, from D1 to D2, stabilized by ten hydrogen

bonds and one ionic interaction between Q500 and K505. In the second, 9 Å-long part of the

linker, the N-terminal strand forms two hydrogen bonds with the C-terminal strand, which

itself forms a β-hairpin. This C-terminal β-hairpin also interacts with the D2 core fold, which

additionally stabilizes the linker region. Three structures of the FliC D12 construct were

solved from two crystals (one asymmetric unit contained two protomers). This allowed for a

structure alignment that helped to visualize the differences in the relative alternative positions

of D2 (Fig. 1D). The angle between the center of mass (COM) of D1, the Cα of P507 at the

tip of D1 and the COM of D2 varies between 130° and 136°. The angle between the Cα of

P507, the COM of D2 and the carbonyl C of T206, a point close to the transition to D3, varies

between 92° and 107°. Thus, the unusual linker would provide a certain degree of flexibility

between D1 and subsequent domains. We were intrigued to explore how these unique

features of the EcN FliC D1-D4 might affect flagellar structure and modelled on the published

cryo-EM structure of *Bacillus subtilis* flagellum (PDB 5WJT) using the D1 for alignment. The

two resulting models differ in their shape as a result of the different HVR domain orientations.

The model with the HVRs closer to the D1 resulted in a closed flagellum of 211 Å diameter,

where the HVRs form an outer ring-like structure. In the other model with 252 Å diameter, D3

and D4 are moved outwards, opening the gap between inner constant domain superstructure

and the outer HVR superstructure (Fig. 1E). In these predicted filaments most interactions

would occur between D0 and D1 domains. Between the two plausible models of the flagellar

assembly, which include the HVR, we noted a difference in the connectivity (Fig. 1F). In the

more 'closed' model, the flagellum would be narrower (211 Å) and HVR domains are closer

to the constant domains, with more putative interactions between different HVR domains.

The more 'open' model shows less contact between the domains but yields a wider flagellum

(252 Å). Collectively, our structural studies show EcN FliC adopts a flagellar fold with unique

features, most importantly the D1-D2 linker and the outer D4 domain, which are compatible

with the formation of a functional flagellum.

Linker and D4 domain are dispensable for TLR5 recognition

A *Salmonella* Dublin FliC:zebrafish TLR5 complex [8] has long been used as a blueprint for

flagellin recognition by human TLR5, although it has been proposed it may represent an

inhibitory complex (discussed in [12]). Nevertheless, to gain an insight into structural aspects

of EcN FliC interaction with TLR5, the EcN FliC and ST FliC structures were superimposed

(Fig. S5). This showed that the EcN and ST D1 domains interacted well with TLR5. However,

the EcN D2-D4 domains hardly interacted with TLR5. Of note, for the ST FliC, an agonist of

both human, mouse and zebrafish TLR5 [11], its D2-D3 were predicted to clash with the

second TLR5 molecule, providing further evidence that the described *S.* Dublin:TLR5

complex is unlikely to represent an active complex (Fig. S5).

We next sought to directly explore the relevance of the unique EcN structural features for

TLR5 recognition, using HEK293T cells stably expressing human or murine TLR5. These

were either challenged with purified flagellins, including the D0-D4 domains (full FliC), ΔD0,

ΔD4, Δlinker or ΔHVR, or exposed to EcN bacteria in which the chromosomal *fliC* had been

edited accordingly (see Methods). Purified ST FliC and/or aflagellated EcN (*ΔfliC*) were used

as controls, respectively. In line with earlier results, the D0 seemed strictly required for both

human and mouse TLR5 recognition. Interestingly, whereas the D4 was largely dispensable

for recognition of purified flagellins at equimolar concentration, deletion of the linker slightly

reduced IL-8 output (Fig. 2A, 2B). When bacteria expressing the corresponding FliC variants

were used to stimulate hTLR5 and mTLR5 HEK29eT reporter cells, no differences were

apparent for hTLR5 (Fig. 2C), but mTLR5 cells showed a slight reduction for ΔD4- or Δlinker-

FliC-expressing bacteria (Fig. 2D). Thus the unique structural features of EcN FliC only

moderately affect direct TLR5 recognition. As an alternative to HEK293T, we performed

similar experiments in more physiologically relevant intestinal epithelial systems, namely,

C2BBe1, an intestinal epithelial cell (IEC) line, and intestinal human organoid cultures, as

well as immune cells, namely, the human macrophage-like THP-1 cell line and primary

murine bone-marrow-derived dendritic cells (BMDCs). Intestinal C2BBe1 cells responded to

purified full-length FliC and ΔD4 FliC with robust IL-8 release, but Δlinker, ΔHVR and ΔD0

FliC did not trigger release significantly above background (Fig. S6A). Human beta defensin

1 (hBD1) mRNA was induced comparably little between flagellin preparations (Fig. S6B), but

hBD2 was strongly induced by all flagellins, with significantly more hBD2 mRNA induced by

the Δlinker flagellin version, and significantly lower by ΔD0 flagellin (Fig. S6C). In similar

experiments using whole bacteria, we noted comparable IL-8 release from C2BBe1 cells in

response to all but *ΔfliC* (aflagellated) and *fliC ΔTLR5rs* bacteria (which lack the TLR5

recognition epitope), suggesting a TLR5-dependent response. Of note, mouse commensal *E.*

coli strains, namely *E. coli* mpk, and *E. coli* MG1655 also triggered significantly lower IL-8

release (Fig. 3A), in line with earlier data [6]. HBD1 mRNA was not induced significantly (Fig.

S7A), but the strong hBD2 mRNA induction (>1000-fold) followed IL-8 results in that *fliC*

mutant strains Δ*TLR5rs* and Δ*fliC*, as well as *E. coli* mpk and MG1655 hardly induced any

hBD2 mRNA (Fig. 3B). Interestingly, in this assay ΔD4 FliC-expressing bacteria also did not

trigger hBD2 mRNA transcription (Fig. 3D). A preliminary test (n=1) in human enteroids

stimulated with purified flagellins showed comparable IL-8 release, except ΔHVR and ΔD0

flagellins, which induced significantly less IL-8 than WT EcN flagellin (Fig. S7B). Switching to

immune cells, which, unlike intestinal epithelial cells, are also competent for the LPS

receptor, TLR4, we focused on entire bacteria only, as even minor traces of LPS in the

purified protein preps would be expected to confound the data in a poorly tractable manner.

On the other hand, the analysis of whole bacteria enables the effects of flagellin to be

assessed in the context of other MAMPs naturally present in the same bacteria, and hence

provides a physiologically relevant context. In co-culture with THP-1 cells, different MOIs of

the EcN variants expectedly released very high levels of IL-8 and TNF, intermediate levels of

IL-6 and IL-10 and low levels of IFNβ and IL-1β, but there were no significant differences

between WT EcN and deletion mutants, mostly likely due to LPS acting as the overriding

MAMP (Fig. S8). Exposure of primary BMDCs (Fig. S9) to flagellin-expressing bacteria

resulted in downregulation of TLR5 (Fig. S10A), no effect on CD40 (Fig. S10B) and

upregulation of CD86 and MHC-II (Fig. S10C), although there were again no significant

differences between strains. The same applied to cytokine release, which was as strongly

induced by but did not vary between different flagellins (Fig. S10E-H). These data are

exemplified for IL-8 in the heat map of Fig. 3C. Collectively, we can conclude that immune

cells, due to the dominance of LPS, cannot distinguish different bacteria based on different

forms of flagellin. However, in cells with strict TLR5 dominance (transgenic TLR5-HEK cells

and epithelial cells), deletion of the linker had a subtle effect on recognition at the level of

purified proteins (albeit not as great as ΔD0). In good agreement with the structural analysis

(*cf.* Fig. 1E), neither the D1-D2 linker nor the D4 domain thus seemed to have a strong very

effect on immune recognition by immune and epithelial cells.

Linker and D4 domain affect flagellar expression, assembly and motility

Apart from immune recognition, the previously reported probiotic effect of EcN might also be

due to different swimming capabilities, e.g., different encroachment of the intestine. To

explore the effect of the deletions outlined above on motility, we first investigated whether

strains expressing the mutant flagellins chromosomally were able to secrete flagellin and

form flagella by assessing flagellin release in the culture media and negative-stain

transmission electron microscopy (TEM) micrographs of different EcN mutants. A preliminary

analysis of culture supernatants by SDS-PAGE and silver staining showed that all deletion

and mutation constructs were detectable and secreted to levels comparable with wild type

(WT) EcN flagellin, except ΔHVR and ΔTLR5rs flagellin (Fig. S11A). To assess the stability

of the proteins themselves, we conducted thermal stability assays with recombinant purified

proteins, which showed that deletion of D4, and even more so the linker, reduced stability,

whereas deletion of the entire HVR did not (Fig. S11B). To check if this had an effect on

flagellation, transmission electron microscopy (TEM) of all EcN mutant strains was

performed. All strains, including the *fliC ΔHVR*, showed flagella, apart from the *ΔfliC*

(expected) and the *fliC ΔTLR5rs* variant (Fig. S12), indicating this deletion might not only

abrogate the reported TLR5 recognition but also proper flagellar assembly. Next, the ability

for motility was assessed in soft agar assays [24](Fig. S13). Interestingly, quantification of

swimming radii (Fig. 4A; 4B) showed that *fliC ΔD4* and *fliC Δlinker* deletions were associated

with a significantly compromised swimming distance compared to EcN WT but retained

residual motility relative to the non-motile aflagellated *fliC* Δ*TLR5rs* and Δ*fliC* strains.

Surprisingly, a strain that lacked the entire HVR (*fliC* Δ*HVR*) was as motile as WT EcN,

suggesting that D4 in its full EcN FliC context contributes to motility and its truncation partly

destabilizes the flagellum, whereas full deletion of the HVR, i.e., deletion of T180-E500, can

yield a functional flagellum, despite lacking the outer domains. Interestingly, single-cell

motility analysis (Fig. 4C) confirmed an essential requirement for the linker in liquid

environment. A deletion of the D4 domain did not affect average swimming speed, whereas

deletion of the entire HVR significantly reduced swimming velocity in this assay. Thus, it

appeared that different environments (viscous soft agar vs. liquid) may require certain

flagellar features for optimal motility. One possibility is that different surface structures of the

flagellum create different levels of traction. Indeed, analysis of individual flagella by TEM

showed interesting differences in surface granularity, namely that, compared to WT the *fliC*

Δ*HVR* mutant showed a both thinner and smoother flagellar filament (Fig. 4D, 4E), whereas

the *fliC* Δ*linker* filament was only slightly thinner but retained a more granular surface

appearance. Moreover, when analyzing the wave form of these flagella, *fliC* Δ*HVR* and *fliC*

Δ*linker* flagella showed smaller distances between consecutive pitches (i.e. a smaller wave-

pitch distance; Fig. 4F). We speculate that especially in liquid environments, a smoother

surface makes motility less efficient, compared to movement in a more viscous soft agar.

Moreover, the low motility of the *fliC* Δ*linker* mutant in both environments (*cf.* Fig. 4A/4B vs

4C/D) – despite a more WT surface granularity but lower wave pitch distance, like in the *fliC*

Δ*HVR* mutant – may be explained by its significantly lower filament stability (*cf.* Fig. S11),

which would be expected to compromise flagellar performance independent of the

environment. Collectively, our analysis indicates that the HVR as a whole, the atypical linker

and D4 domain do not hinder TLR5 recognition, but they affect bacterial motility differently

depending on the environment by modulating protein stability, surface roughness and wave-

form of the filament.

Thus far, our data indicate that TLR5 recognition of EcN flagellin was less structurally

sensitive than bacterial motility, so that deletions in EcN flagellin that significantly affect or

'trade in' motility still did not evade full TLR5 recognition. Next, we sought to probe whether

more subtle modifications, such as point mutations, might generate EcN variants, in which

motility was retained but TLR5 recognition evaded, as shown for ST FliC [22]. In light of the

observed differences in sensitivity between human and murine TLR5, we exposed HEK293T

expressing either of these receptors to different EcN mutants, which were generated based

on comparisons of immunostimulatory (EcN and ST FliC) and TLR5-evasive flagellin

(*Campylobacter jejunii*) mutants at the proposed TLR5 interface [21]. Most single mutants

still caused significant secretion of IL-8, e.g. R93K, E94T and R530Q, 24 h after infection

(Fig. S14). R91T seemed to be the exception and emerged as crucial for recognition by

human and even murine TLR5, as it strongly reduced IL-8 secretion. We also tested

combinations of the previous mutations, but these only highlighted the dominance of R91T

for TLR5 recognition. We were curious about the effect of these mutations on motility (Fig.

S15 and Fig. 5A). Evidently, 'tampering with' R91 completely crippled EcN motility (Fig. 5A)

in that we were unable to identify any single or multi-mutants with abrogated TLR5

recognition but intact motility amongst the analyzed R91-mutated variants. Summarizing and

correlating relative (to WT) NF-κB activity of TLR5 HEK293T with motility data (Table S2 and

Fig. 5B/5C) suggested that greatest achievable reduction in TLR5 activity, to (only) about

70%, coincided with a motility reduction to 80% (R530Q) or even ~30% (R91T E94T). Any

further reduction of TLR5 activity below 60% resulted in a concomitant drop in motility to 40%

or less (red quadrant). Conversely, several mutants with less than 50% motility retained a

relatively high TLR5 reactivity (orange quadrant). This shows that motility seems far more

sensitive to mutations than TLR5 recognition is. Alternatively phrased, EcN undergoing FliC

mutations would probably be more likely to lose motility before avoiding TLR5 recognition.

Discussion

EcN represents one of the most extensively studied probiotic *E. coli* strains. Although

previous work implicated its flagellum, and especially its HVR, and TLR5 to be a critical

mediator of probiotic effects in experimental colitis [6], these properties have not yet been

linked to certain molecular or structural features. Here, we therefore determined the EcN

HVR structure, highlighted unique features compared to the prototypical ST FliC, but

probably not uncommon in *E. coli*. Most striking were the linker between D1 and D2 and the

additional D4 domain.

Regarding the latter, our data about the existence of an additional OD is in good agreement

with a recent cryo-EM study on flagellar filaments from enterohemorrhagic *Escherichia coli*

O157:H7, enteropathogenic *E. coli* O127:H6, *Achromobacter*, and *Sinorhizobium meliloti*

[20]. This study first described outer domain sheaths as a structural feature enabling cross-

linking of flagellin protomers via D4 dimerization, thus allowing for prolonged *E. coli* tumbling.

Moreover, it was speculated, albeit not proven, that a screw-like outer surface might provide

additional traction in high viscosity environments such as the intestine. Our structural data

revealed that probiotic EcN shares these structural features with pathogenic *E. coli* and that

the observed D4 dimerization also applies to EcN flagellar structure. In fact, HVRs long

enough to contain a D4 occur in many bacteria and the D4 sequence of EcN collects BLAST

hits, e.g., in pathogens like *Shigella* spp (data not shown).

Interestingly, although this feature was not reported or functionally assessed before, the

extended D1-D2 linker in EcN FliC is also present in atomic cryo-EM models of EHEC

O157:H7 and EPEC O127:H6*,* two pathogenic *E. coli* strains. These structures show a linker

of different amino acid sequences but similar length and angle to EcN (Fig. S16).

Interestingly, in the soil bacterium *Achromobacter,* the linker is also of a similar length and

angle; the flagellin of *Sinorhizobium meliloti,* another soil dweller, also shows this linker but

the angle is different. Our study suggests that in these species the linker may contribute to

optimal adaptation of motility to different environments. Further structure-function studies

may show by which mechanism rigid linkers in general contribute to motility across species.

Since both the linker and D4 domains individually emerged as affecting optimal motility, it

was unexpected that deletion of the entire HVR did not compromise motility in soft agar

assays to the same extent. This may be explained by the fact that functional flagella can be

formed by flagellins that entirely lack D2, D3 or D4 domains, i.e., HVRs (e.g. *Bacillus*

subtilis[25]), or harbor shorter HVRs: For example, *Pseudomonas aeruginosa* only contains a

D3 and it adopts a highly elongated structure that also engages in interdomain contacts [26].

Thus, the significance of any or small (D3 only) vs. extended (D3 and D4) HVRs remains

somewhat enigmatic and our data begs the question why EcN retains the biosynthetic

burden of an additional ~320 amino acids for an entire HVR that appears redundant for

motility. We speculate that in the intestine, a greater adaptability to different environments –

ranging from liquid environments to viscous intestinal mucus[27] – the HVR with a wider and

much more textured outer surface provides the EcN flagellum with better adaptability. Future

in vivo studies will be required to investigate this in detail.

In line with the TLR5 epitope mapping primarily to the D1 domain [13, 21], presence of linker,

D4 or the HVR did not affect TLR5 recognition, at least *in vitro*. This indicates that the TLR5-

mediated probiotic effect [6] is probably not dictated by unique HVR features of EcN flagellin

directly, i.e. by modulating interaction with TLR5. Rather, we hypothesize that the effect may

be indirect, e.g., EcN flagellins and resulting flagella may facilitate bacterial encroachment

onto the epithelium by enhancing motility through the intestinal mucus. This may trigger more

TLR5-signaling, which is required to mediate the probiotic effect. In line with this hypothesis,

flagellin-dependent TLR5 signaling was shown to mediate the symbiotic properties of another

gut commensal, *Roseburia hominis* [28] and thus, this phenomenon may be more common

than previously thought. More extensive *in vivo* analyses are thus warranted.

That EcN flagellin can activate TLR5 but structurally would not permit formation of a (TLR5-

EcN FliC)$_2$ complex modelled on the (zebrafish TLR5-Salmonella FliC)$_2$ [8] appears puzzling.

However, recent data from zebrafish and grass carp indicate that the structure of *Salmonella*-

TLR5 homodimers most likely represents an inhibitory complex [12], as TLR5a/TLR5b

heterodimers (not homodimers as in the crystal structure) are required to sense flagellin in

fish [11]. Thus, the structure of an activating complex with human TLR5 may generally differ

from the only known structure to date. Depending on the features of the engaging flagellin

[8], it might accommodate not only small HVRs but also longer HVRs connected to the D1 by

a rigid linker. Further structural studies of co-complexes may be required to unequivocally

confirm this.

Collectively, our study sheds light on structural features of the flagellin molecule of a primary

human probiotic *E. coli* strain, enabling insights into the trade-offs faced by such bacteria in

terms of immune recognition versus. motility in different environments. Clearly, environment-

dependent motility appears more structurally constrained than immune recognition. Unless

bacteria are prepared to accept complete loss of motility, TLR5 recognition accommodates

enough structural flexibility to avoid more subtle attempts if immune evasion by bacteria that

seek to retain motility. On the other hand, certain strains, e.g. *Campylobacter jejunii* [21],

seem to have reached TLR5 ignorance whilst retaining motility. We hypothesize that in

complex microbiomes trade-offs between immune recognition versus. motility are probably

finely tuned and highly species-specific, cautioning against oversimplification. Our analysis

highlights that a concomitant determination of structure-, motility- and immune-related

parameters may be necessary for additional individual flagellins and their host species to

deduce more general principles.

Materials and Methods

General materials

All used chemicals and reagents were purchased from Sigma unless otherwise stated. Kits

are listed in Supplementary table S1.

Gibson assembly, including generation of flagellin mutants

For the digestion of plasmid backbone, 2 µg plasmid were digested for 1 h at 37 °C using two

of the following enzymes, ApaI, BSu15I (ClaI) and/or HindIII fast Digest (ThermoFisher),

before enzyme activity was deactivated for 5 min at 95 °C. Digested plasmid fragments were

separated on 0.8 % agarose gels, fragments of the desired size cut and DNA was isolated

using GeneJET Gel Extraction Kit according to manufacturer's manual. Fragments were

dissolved in a volume of binding buffer equal to the weight (100 µl per 100 µg) at 60 °C for 10

min while shaking. DNA was bound to the column by centrifugation and then washed using

the supplied Wash Buffer. After a dry centrifugation to remove excess liquid, the DNA was

eluted using 30 µL ddH$_2$O. The generated backbone fragment was then used for Gibson

assembly. Gibson assembly was performed as published [29]. In brief, 25 µM primers

(Supplementary table S2) with an overlapping region of 30 nucleotides to the target insertion

site or another DNA fragment and 100 ng template DNA were used to amplify inserts using

the SuperFi II Polymerase according to manufacturer's instructions. The PCR run consisted

of an initial denaturation at 98 °C for 30 s, followed by 35 cycles of 10 s at 98 °C for

denaturation, 30 s at 60 °C for primer annealing and 30 s/kb at 72 °C for elongation before

the final elongation for 10 min at 72 °C. The PCR products were applied to a 1% agarose gel

and isolated from the gel as described for the digestion of the plasmid backbone. For the

actual Gibson Assembly reaction, all fragments (2 – 3 µL of each) and the vector backbone

(3 – 4 µL) were combined with 10 µL of Gibson Master Mix and incubated for 1 h at 60 °C.

The generated target vector was transformed into a suitable bacterial host cell line (either *E.*

coli DH5α or 116 λpir[+]) by heat shock. For this purpose, 6 – 10 µL of Gibson mix were

incubated with 50 µL chemically competent bacteria for 30 min on ice, followed by heat

shock at 42 °C for 40 s, 10 min on ice and incubation with 1 mL SOC medium at 37 °C for 1

h. Bacteria were plated on LB agar containing the suitable antibiotic. Colony formation was

seen after an overnight incubation at 37 °C.

Plasmid preparation

For protein production and purification, EcN FliC constructs in the pET-19b (Novagen)

vector were used. FliC E, FliC ETEV, FliC ∆D0D1, FliC ∆D4, FliC ∆D4s and

FliC ∆D3D4 insert sequences .**are listed in** Supplementary information**Supplementary**

figure legends (included in the actual supplementary figures)

Amino acid sequences for constructs used. The constructs FliC_∆D0 and FliC_∆D0D3D4

were synthesized by Biocat (Heidelberg, Germany) and inserted in pET-19b vectors using

restriction cloning with NcoI and BamHI. To reconstitute expression of FliCin fliC-deficient

bacteria, a pGEM-T backbone was used. For modifying the chromosomal FliC region,

pSB890 backbones were used. Plasmids used are standard subcloning, chemically

competent E. coli DH5α cells were used (Invitrogen). Plasmids were isolated using

commercial miniprep kits (Wizard Plus SV miniprep kit, Promega) and verified by automated

DNA sequencing (Microsynth Seqlab).

RNA isolation and quantitative PCR

RNA was isolated using the RNeasy Mini QIAcube protocol for animal tissue and cells with

DNAse digestion according to manufacturer's instructions. RNA was either directly reverse-

transcribed into cDNA or intermediately stored at –80 °C. For reverse transcription,the High-

Capacity RNA-to-cDNA-Kit (Thermo Fisher Scientific) was used according to manufacturer's

instructions, RNA concentrations were adjusted to 0.1 µg/µL. 10 µL of RT buffer and 1 µL of

RT enzyme were added per sample and incubated for 1 h at 37 °C before the enzyme was

inactivated for 5 min at 95 °C. 5 µL SYBR Green MasterMix (ROX, Roche), 0.3 µL of each

primer (10 µM, Supplementary Table S2) and 1 µL undiluted (for hBD1 and hBD2 qRT-PCR)

or 1:10 diluted (for GAPDH as housekeeping gene) cDNA were mixed with 3.4 µL ddH$_2$O per

reaction and run at a QuantStudio 7 Flex system. Thereby, an initial incubation at 50 °C for 2

min was followed by a denaturation step at 95 °C for 10 min. In the PCR stage, 40 cycles of

a denaturation at 95 °C for 15 s alternated with an elongation at 60 °C for 1 min. To generate

a melting curve, the samples were first heated to 95 °C for 15 s, then cooled to 60 °C,

stepwise heated with 0.05 °C/s increments to 95 °C and kept at 95 °C for 15 s. Data was

analyzed using the $\Delta\Delta C_T$ method.

Bacterial culture

Bacteria were cultured in LB medium with selective antibiotics if not stated otherwise.

Bacterial overnight cultures were set up from glycerol stocks, and day cultures by adding 50

µL overnight culture to 5 mL LB and grown while shaking at 120 rpm for 2 or 5 h for

stimulation or motility assay, respectively. For quantification, bacteria were centrifuged at

3200 × g for 5 min, supernatants discarded, and bacteria resuspended in 1 mL PBS.

Quantification was done by measuring the optical density (OD) at 600 nm assuming an OD_{600}

of 1 contains 5 x 10^8 colony forming units (CFUs) of *E. coli*. The concentration of bacteria

was calculated according to the desired final multiplicity of infection (MOI).

Soft agar motility assay

Soft agar plates were prepared from 5 g peptone, 2.5 g NaCl, 1.5 g agarose in 500 mL dH_2O

with 20 mL per Petri dish. Bacteria were cultured for 5 h in a day culture to reach a late

exponential growth phase. 3 µl of day culture were poked into the center of a soft agar plate

and incubated for 24 h at 37 °C. Pictures of the plates with a ruler were taken for

quantification. The swimming radii were measured at three positions using ImageJ and

averaged. To compare two strains on the same plate a pipet tip was dipped into the day

culture and poked into the agar at approximately $1/4^{th}$ of the plate diameter. The same

procedure was repeated with the second strain on the other half of the plate. Plates were

incubated 24 h before pictures were taken. Diameters were not quantified due to strains

hindering or influencing each other growth.

Bacterial single cell swimming assay in liquid media

The *E. coli Nissle* strains WT, *fliC ΔD4*, *fliC ΔLinker*, *fliC ΔHVR* and *ΔfliC::FCF*, all containing

a translational insertion in *malEK::PJ23119-mCherry* as a fluorescent marker were grown

overnight in LB at 37 °C and 180 rpm agitation. The next day, strains were diluted to an

OD_{600} of 0.15 and grown for 2 h at 37 °C and 180 rpm. OD_{600} was measured and samples

were diluted 1:1000 in PBS supplemented with 0.2% of glucose (w/v) and carefully loaded in

a self-made flow cell. Cell trajectories were recorded using a Ti-2 Nikon inverted microscope

equipped with a 20× objective, by imaging the phase contrast channel at 20 fps for 10 s.

Multiple videos were recorded and analyzed as follows: cell segmentation was performed in

Fiji [30] using the MicrobeJ plugin, version 5.13p [31]. Subsequent analyses were done using

custom Python scripts. For each detected particle, the centroid coordinates and area were

extracted using the skimage.measure module from scikit-image [32]. Particles were filtered

by area (5-150 pixels) to remove false detections. Tracking was performed on the resulting

binary masks using the Python package trackpy, version 0.6.4, with a search range of 5

pixels and a memory of 1 frame. For each tracked particle, centroid positions were used to

calculate instantaneous velocities from frame-to-frame displacements divided by the time

interval (50 ms). The mean velocity for each track was calculated as the average of its

instantaneous velocities. Only tracks with durations of at least 3 s were included in the

analysis. Based on the non-motile *ΔfliC* control, tracks with mean velocities below 5 μm/s

were considered non-motile (Brownian motion) and excluded from velocity statistics. For

each strain, data were pooled from multiple regions of the flow cell.

Negative stain transmission electron microscopy (TEM) of bacteria

For negative stain TEM of flagellated bacteria, bacterial day culture was inoculated with 500

μL overnight culture in 10 mL LB media. Bacteria were grown while shaking at 120 rpm, 37

°C for 4 h, adjusted to 1×10^9 CFU/mL and transferred to the Electron Microscopy Facility at

the Max Planck Institute for Biology Tübingen. The bacteria were shaken for another 1 h

before 1 – 1.5 mL of culture were spun down at 5000 × g for 3 min and the pellet was

resuspended in 100 μL of culture. 4 μL of resuspended bacteria were applied to glow-

discharged carbon-coated EM grids and incubated for 2 min. The grid was washed with three

drops of ddH$_2$O and three drops of 1% uranyl acetate. The 4th drop of 1% uranyl acetate was

incubated for 5 min before the liquid was removed and the grid was dried. Grids were

analyzed in a transmission electron microscope (Tecnai Spirit, Thermo Fisher Scientific,

Eindhoven, The Netherlands) equipped with a CMOS camera (TemCam-XF416, TVIPS,

Gilching, Germany).

Negative stain transmission electron microscopy (TEM) of isolated flagella

For negative-stain electron microscopy of isolated flagella, overnight cultures were used to

inoculate 100 ml LB medium to an OD$_{600}$ of 0.01. Bacteria were grown at 37 °C with shaking

at 80 rpm overnight. Cells were harvested by centrifugation at 2,000 × g for 20 min at 4 °C,

and the supernatant was discarded. The pellets were resuspended in 10 ml of HEPES (20

mM, 150 mM NaCl, pH 7.4) buffer per sample. To shear off the flagella, the bacterial

suspensions were transferred to new flasks and stirred at 500 rpm for 1 h at 4 °C. Bacterial

cells were removed by centrifugation at 8,000 × g for 30 min at 4 °C, repeated twice. The

supernatants were collected and concentrated using ultrafiltration units (Amicon 10 kDa

MWCO). Centrifugation was performed at 3,214 × g at 4 °C. The concentrated flagella

suspension was stored at 4 °C. For negative staining, 4 µl of the isolated flagella suspension

was applied to glow-discharged carbon-coated grids for 1 min and gently blotted away with

filter paper. This procedure was repeated three times with ddH$_2$O. Subsequently, 4 µl 2%

(w/v) phosphor-tungstic acid was applied for 45 s, replaced with 4 µl ddH$_2$O, and liquid was

directly removed by blotting with filter paper. The grids were allowed to air dry for 5 min.

Samples were examined using a JEOL JEM-2100 Plus transmission electron microscope

operated at 200 keV. Images were acquired at various magnifications. Flagella diameters

were measured using (Fiji is just) ImageJ (v1.54). Prior to measurements, background

subtraction was performed using the rolling ball method (2 px radius), and diameters were

determined from plot profiles.

Silver staining of SDS-PAGE gels

For visualization of low protein concentrations, the Pierce Silver Stain Kit (Thermo Fisher)

was used according to the manufacturer's instructions.

Protein production and purification for structural studies

Pre-cultures were inoculated with *E. coli* BL21 containing the appropriate plasmid and grown

overnight. 1-6 L LB media was inoculated 1:100 with overnight culture and grown at 37°C

while shaking until an OD_{600} of 0.6. Protein production was then induced with 0.2-0.5 mM

IPTG, and cells were grown at 25 ℃ overnight or 37 ℃ for 4 h. Cells were harvested by

centrifugation for 15 min at 9200 × g (Sorval RC 6+, Thermo Fisher Scientific). The pellets

were frozen in liquid nitrogen and stored at -80 °C. Cell pellets were resuspended in 5 mL

lysis buffer (50 mM Tris pH 8.0, 300 mM NaCl) per gram pellet wet-weight. The suspension

was supplemented with 1x "cOmplete" protease inhibitor (Roche, Basel, Switzerland) and

125-500 U Benzonase Nuclease (Merck KGaA, Darmstadt, Germany). Cells were lysed by

sonication with an amplitude of 40%, for 3 min, with on-off pulse cycles of 0.5 s (Digital

Sonifier 250, Branson Ultrasonics). The lysate was centrifuged for 45 min at 34'500 × g

(Sorval RC 6+, Thermo Fisher Scientific,), filtered (0.22 μm cutoff) and loaded on a 5 mL

HisTrap FF crude column (GE Healthcare) equilibrated with His-A buffer (50 mM Tris pH 8.0

300 mM NaCl) using an ÄKTAprime plus FPLC system (GE Healthcare). After sample

loading the column was washed with His-A and 2% His-B Buffer (50 mM Tris pH 8.0, 300

mM NaCl, 500 mM imidazole). Proteins were eluted using 10%, 20%, 30%, 60% and 100%

His-B Buffer steps. Fractions were pooled based on SDS-PAGE analysis. Pooled protein

fractions were dialyzed against SEC-buffer, or His_A buffer, using a Spectra/Por dialysis

membrane at 4° C overnight (Spectrum Laboratories Inc). If proteolytic cleavage of the His-

tag was attempted, 1-2 mL TEV protease (0.5 mg/mL, produced in our laboratory) was

added to the pooled fractions, before dialysis. The His-tagged TEV protease and undigested

protein were removed by a second IMAC. The flow through was pooled and yield and purity

assessed by SDS-PAGE. For size exclusion chromatography, pooled samples were filtered

(0.22 μm) and loaded on an equilibrated SEC-column. For preparative SEC a HiLoad 16/60

Superdex 200 or Superdex 75 column (Pharmacia) with a Pharmacia LKB GP250 plus

system were used. For analytical SEC a 3.2/30 Superdex 200 Increased column (GE

Healthcare) or a 3.2/30 Superdex 75 column (Pharmacia) was used with an Ettan LC system

(GE Healthcare). For crystallization a SEC-buffer   containing 20 mM HEPES pH 7.5 and150

mM NaCl was used. For stability experiments DPBS-buffer (0.90 mM $CaCl_2$, 0.49 mM $MgCl_2$,

2.67 mM KCl, 1.47 mM KH_2PO_4, 137.93 mM NaCl, 8.06 mM Na_2HPO_4) was used for SEC.

Crystallization

To determine an appropriate starting concentration for initial crystallization experiments, a

precipitation test was done. 0.5 µL of filtered (0.22 µm) protein solution was mixed on a glass

slide, with different precipitate (30% (w/v) PEG4000 and 3 M ammonium sulfate) solutions.

The protein concentration was stepwise increased until the protein precipitated within 1 min

with at least one of the two precipitants. For initial screens, commercially available kits were

used according to the manufacturers' instructions: JCSG, Morpheus (Molecular Dimensions),

Wizard I-IV (Emeraly BioSystems), Crystal Screens I and II and PEG ION (Hampton

Research). Crystallization screens were done using 96-well siting drop Intelli-plate (Art

Robbins Instruments, Sunnyvale, USA) with either a Gryphon (Art Robbins Instruments,

Sunnyvale, USA), or Freedom EVO (Tecan, Männedorf, Switzerland) crystallization robot.

Drops were set up in a one-to-one ratio of protein solution and mother liquor, with a total drop

size of 400 nL with the Gryphon, and 600 nL with the Freedom EVO. Crystals were optimized

using 4 × 6 (Hampton Research, Aliso Viejo, USA) hanging drop plates.

Structure determination

To optimize the cryo-protectant or to test diffraction capacity of a crystal, the in-house X-ray

system was used. This system is composed of a rotating copper anode X-ray source

(MicroMax-007HF (Rigaku, Sevenoaks, UK) producing CuKα radiation at λ = 1.5418 Å and a

mar345 image plate detector (marresearch, Norderstedt, Germany). The HATODAS II [33]

server was used to identify promising heavy atoms for initial soaking experiments. Therefore,

crystals were soaked for 5 and 10 min in crystallization solutions containing different heavy

atoms with a concentration of 10 mM or saturated if the solubility was lower. For Sm^{3+} and

UO_2^{2+} compounds various crystals were soaked with concentrations ranging from 2.5 mM to

10 mM and soaking time between 1 min and 24 h. Additionally, crystals were soaked with

3 mM $(NH_4)_2WS_4$ or $Lu(Ac)_3$ for 2 h, 1 mM Ta_6Br_{14} for 1 week, 2.5 or 5 mM $EuCl_3$ for 24 h or

1 mM $SeC(NH_2)_2$ for 10 min. Data sets were collected at the macromolecular crystallography

beamline X06DA-PXIII of the Swiss Light Source (Paul Scherrer Institute, Villigen,

Switzerland). Data sets for structure determination via molecular replacement were

collected with $\lambda = 1$ Å (12.3984 keV). For anomalous data the wavelength was adjusted to

specific values, according to the absorption edge of the element used. The XDS program

package was used to process (index, integrate, scale) the experimental diffraction image

data sets, XSCALE to scale and merge datasets, XDSCONV to convert different

crystallographic file formats [34]. From the PHENIX [35] program package XTRIAGE [36]

was used to analyse data sets and PHENIX.REFINE [37] was used for simulated annealing

and refinement. From the CCP4 [38] program package, POINTLESS [39] was used to

identify possible space groups, MATTHEWS_COEF [40] was used to evaluate the possible

unit cell content, CHAINSAW [41] to modify models for molecular replacement, PHASER [42]

and MOLREP [43] for molecular replacement, REFMAC5 [44] for refinement and COOT [45]

for model building and evaluation. PYMOL [46] was used for structure examination and figure

generation. Experimental phasing was done using SHELX C/D/E [47] and AUTOSHARP [48]

with SOLOMON [49] for density modification. MOLPROBITY [50] was used for model

evaluation. The following structure files were deposited to rcsb.org: [to follow]

Cultivation and stimulation of mammalian cells

HEK293T cells were grown in DMEM supplemented with 10% FCS and 1%

Penicillin/Streptomycin. mTLR5 HEK293T cells required the selective antibiotics Normocin

(50 mg/L) and Blasticidin (10 mg/L), while hTLR5 HEK293T cells required Blasticidin (10

mg/L), Hygromycin B (100 mg/L) and Tetracycline (12.5 mg/L) that were added freshly to the

medium. CaCo-2 (subclone C2BBe1) cells were cultured in DMEM supplemented with 10%

FCS, 1% non-essential amino acids (Gibco) and 1% Penicillin/Streptomycin. THP-1 cells

were cultured in RPMI 1640 (very low endotoxins), supplemented with 10% FCS, 1% L-

glutamine (gibco) and 1% Penicillin/Streptomycin. Murine bone marrow derived dendritic

cells (BMDCs) were cultured and differentiated in RPMI 1640 (very low endotoxins),

supplemented with 10% FCS, 4% GM-CSF conditioned medium (produced in our

laboratory), 1% non-essential amino acids (gibco), 1% Penicillin/Streptomycin and 0.5% β-

mercaptoethanol. Cells were incubated at 37°C 5% CO_2 and split every 3 - 4 days. **Error!**

Reference source not found.2×10^5 cells per well in a 24-well plate or 3×10^4 cells per well

of a 96 well plate were seeded in the respective culture medium. For stimulation with

recombinant FliC, cells were incubated ~ 40 h at 37 °C 5% CO_2 before medium was replaced

by medium containing recombinantly isolated FliC proteins for 24 h. In case of bacterial

infection, medium was replaced 24 h after seeding with antibiotics-free medium and cells

were incubated overnight at 37 °C 5% CO_2 before medium was exchanged to antibiotics-free

medium containing bacteria for a total of 24 h or 6 h in case of C2BBe1 cells. 1 h post

infection (p.I.) gentamicin was added to inhibit overgrowth of bacteria. For both stimulation

and infection, supernatants were taken, centrifuged at 3200 × g for 10 min, or max speed for

6 min and transferred to a new tube or plate. Cell-free supernatants were used for ELISA

analysis and C2BBe1 cells were lysed in 350 µL of RLT/1% β-mercaptoethanol and frozen at

–80 °C for RNA isolation. To differentiate THP-1 cells to a macrophage-like phenotype, 2 ×

10^5 cells per well of a 24-well plate were seeded in medium containing 100 nM PMA. Three

days later, PMA was removed and cells were left to rest for 72 h at 37 °C 5% CO_2 before

stimulation/infection (as described above).

Cultivation and stimulation of enteroids

To revive enteroids, cells were washed twice with pre-warmed DMEM before being in ice-

cold Matrigel. 4 drops of 10 µL Matrigel-enteroid mix were placed into one well of a 24 well

plate. A 10 min incubation at 37 °C followed to polymerize the Matrigel. 400 µL of enteroid

passage medium (growth medium [Advanced Dulbecco's modified Eagle medium/F12, 1 mM

HEPES, 1x Glutamax, 1x B27, 1 mM N-Acetylcysteine, 10 nM Gastrin, 50 ng/mL EGF, 10

mM Nicotinamide, 500 nM A83-01, 10 µM SB202190, 50% L-WRN-conditioned media

(contains Wnt3a, R-spondin 3, and Noggin)] + 10 µM Y27632, 250 nM CHIR99021) were

added. 2 – 3 days afterwards the medium was replaced by enteroid growth medium.

To split enteroids, Matrigel was dissociated by 1 mL of TrypLE express (gibco) and careful

pipetting ensured breaking. The cell-Matrigel mix was diluted 1:10 and incubated at 37°C

while pipetting until single cell dissociation was achieved. To inhibit trypsin, 100 µL of FCS

were added. Enteroids were pelleted at 300 × g for 5 min and resuspended in Matrigel and

dropped into a 24 well plate and cultivation was performed as described above. The medium

was replaced every 2 - 3 days by fresh enteroid growth medium and cells were split every 1

– 2 weeks, when they grew large or dense enough. For stimulation, the enteroids were

treated as if they were split. When enteroids in fresh Matrigel were distributed in 96 well

plates, one 10 µL drop was placed in one well and 200 µL of enteroid passaging medium

was added. 2 - 3 days later it was replaced by 200 µL of enteroid growth medium. Enteroids

were in culture for at least 4 days prior to stimulation. For stimulation, the medium was

replaced by recombinant FliC containing medium and left on cells for 18 h. Supernatants

were taken and used for IL-8 ELISA.

ELISA

Generally, ELISAs were performed according to manufacturer's protocols using commercially

available kits (Supplementary Table S1). If necessary, buffers were prepared accordingly.

Dilutions of reagents used were done as recommended by the manufacturer in suitable

buffers. Generally, sandwich ELISAs were performed. Thus, the plates were coated with a

suitable capture antibody overnight at 4 °C, before the coating solution was washed off in

four consecutive washing steps and replaced by the double standard volume (e.g. 200 µL or

80 µL) of the suitable blocking solution. All following incubations were at room temperature

while swaying. The blocking buffer was removed, the plates washed four times and the

standard protein, as well as the cell-free samples in a suitable dilution were applied and

incubated. Another four times wash followed, and the detection antibody fused to biotin was

incubated on the wells. After four more washes, the standard volume of a streptavidin-tagged

alkaline phosphatase (AP) or horseradish peroxidase (HRP) was added to the wells. An

incubation followed before the conjugate of enzyme and streptavidin was removed by a more

vigorous washing step according to the manufacturer's instructions. The substrate, either

para-nitrophenyl phosphate (pNPP) for AP or tetramethylbenzidine (TMB) for HRP, was

prepared and added to the wells. The incubation was performed in the dark at room

temperature without swaying for 20-30 min until the desired color change was reached. The

reaction was either stopped or not using 2N sulfuric acid. The absorption was read at a

wavelength of 405 nm for AP and 450 nm for HRP using a spectrophotometer.

Flow cytometry

For flow cytometric analysis, cells were carefully scraped off the wells and transferred to a V-

bottom 96 well plate and unspecific binding was blocked by adding 100 μL of either cell

culture supernatants of 2.4G2 cells secreting an antibody directed against the murine Fcγ

receptor for murine cells or a commercially available human Fc-Block (BD; diluted 1:20 in

FACS Buffer) for human cells for at least 15 min at room temperature. The cell pellet was

washed twice using PBS to remove FCS before staining for their viability using a fixable

viability dye (ThermoFisher) diluted 1:10,000 in PBS for another 15 min in the dark on ice or

at 4°C. After two more washing steps in PBS 1% FCS the staining solution containing the

desired antibodies in a 1:100 dilution (α-mCD11b BV421 (HL3, BD), α-mMHC-II PerCp-cy5.5

(M5/114.15.2, BD), α-mCD86 PE-cy7 (GL1, BD), α-mCD40 APC (3/23, BD), α-mTLR5 PE

(ACT5, BD) was added to the cells and incubated for 30 min in the dark at 4 °C or on ice.

The cells were washed another one time before resuspending them in 100 μL of FACS

Buffer. Measurements were done on a BD LSR Fortessa II. Analysis was done using FlowJo

v10.

Animals

C57BL/6J and C57BL/6N mice (Charles River Laboratories, Sulzfeld, Germany) were

ordered at an age of 6-8 weeks at least one week prior to start of the experiments. Mice were

housed under specific-pathogen-free (SPF) conditions in individually ventilated cages (IVCs).

All experiments were carried out in compliance with specific local institutional guidelines on

animal experiments under regular hygiene monitoring in accordance with the German animal

protection law and according to regulations by Gesellschaft für Versuchstierkunde/Society for

Laboratory Animal Science, GV-SOLAS, as well as according to the European Health Law of

the Federation of Laboratory Animal Science Associations, FELASA. All experiments were

registered and approved by the local federal authority, the Regierungspräsidium Tübingen,

under approval H01/20M.

Isolation, differentiation and stimulation of primary bone marrow derived dendritic

cells

Bone-marrow derived dendritic cells (BMDCs) were isolated and differentiated according to a

published protocol [51].Therefore, mice were sacrificed at an age of about 8 weeks, femurs

and tibias were isolated, freed of residual tissue and sterilized using ethanol while still intact.

Bones were cut open at both ends under sterile conditions, and a 0.45 mm syringe filled with

sterile PBS was used to flush out the bone marrow cells. Cells were spun down at 300 × g for

5 min, supernatant discarded and washed using fresh PBS. 2×10^6 cells were seeded per 10

cm Petri dish (not cell culture treated) in 10 mL of GM-CSF containing BMDC culture medium

as listed above**Error! Reference source not found.**. The cells were incubated at 37°C 5%

CO_2 for two days before another 10 mL of fresh BMDC culture medium was added to feed

the cells. Three days after the feeding, 10 mL of the culture supernatant was taken off the

cells, centrifuged at 300 × g for 5 min, supernatant was discarded, and cells resuspended in

10 mL fresh BMDC culture medium. Cells were subsequently transferred back to the Petri

dish and incubated two more days. To seed the cells for stimulation, cells were pooled and 2

$\times 10^6$ cells per well were seeded in a 6-well plate 1 h prior to stimulation in BMDC culture

medium. Either recombinantly isolated proteins at the indicated concentration or bacteria

grown for 2 h in a day culture were added at an MOI of 2. BMDC culture medium contains

penicillin and streptomycin, thus no gentamicin kill was additionally performed. BMDCs were

incubated for 16 h with FliCs or bacteria before supernatants were taken, centrifuged at max

speed for 5 min and transferred to a new tube. Cells were scraped off at 16 h after incubation

for flow cytometric analysis.

Statistical analysis

Collected data was visualized and statistically analyzed using GraphPad Prism version 8 and

10. First, Shapiro-Wilk test was deployed to test for normality in each dataset (e.g,. each

concentration of each construct or each MOI of each bacterium separately). If only few

datasets (less than 20% of datasets) were not normally distributed according to Shapiro-Wilk

test, normality was nevertheless assumed because of small sample sizes (often n=3), and

when normal distribution in other data sets of same stimulus often only varying in

concentration/MOI was confirmed. ANOVA, either ordinary one-way or 2-way ANOVA or

Kruskal-Wallis test for not-normally distributed data with Dunnett correction for multiple

testing, was chosen depending on the nature of the data to be analyzed as indicated in the

figure legends. Considering p values, the following scheme was applied: * $p \leq 0.05$, ** $p \leq$

0.01; *** $p \leq 0.001$; **** $p \leq 0.0001$. In some cases, two statistical tests were performed on

the same data set. One test was utilized to test for statistical differences of the sample

compared to unstimulated control (mock). Here, the same significance levels as described

above were used, but the symbol used was '#' instead of '*'. The '*' symbol was used to

indicate statistical differences between a given sample and either EcN WT, in case of

bacterial infection, or full-length EcN FliC, in case of stimulation with recombinantly isolated

proteins. Again, the same significance levels as described above were used with an '*'

symbol. The test used, the group to which the sample was tested against (sample versus

mock or sample versus EcN WT or sample versus full-length) as well as the symbol used (#

or *) to depict significance level are indicated in the figure legends.

<u>References</u>

1.   Erhardt, M., K. Namba, and K.T. Hughes, *Bacterial nanomachines: the flagellum and*
789   *type III injectisome.* Cold Spring Harb Perspect Biol, 2010. **2**(11): p. a000299.
790   2.   Einenkel, R., et al., *The structure of the complete extracellular bacterial flagellum*
791   *reveals the mechanism of flagellin incorporation.* Nat Microbiol, 2025. **10**(7): p. 1741-
792   1757.
793   3.   Macnab, R.M., *How bacteria assemble flagella.* Annual Review of Microbiology, 2003.
794   **57**: p. 77-100.
795   4.   Samatey, F.A., et al., *Structure of the bacterial flagellar protofilament and implications*
796   *for a switch for supercoiling.* Nature, 2001. **410**(6826): p. 331-7.
797   5.   Yonekura, K., S. Maki-Yonekura, and K. Namba, *Complete atomic model of the*
798   *bacterial flagellar filament by electron cryomicroscopy.* 2003.
799   6.   Steimle, A., et al., *Flagellin hypervariable region determines symbiotic properties of*
800   *commensal Escherichia coli strains.* PLoS Biol, 2019. **17**(6): p. e3000334.
801   7.   Hayashi, F., et al., *The innate immune response to bacterial flagellin is mediated by*
802   *Toll-like receptor 5.* Nature, 2001. **410**(6832): p. 1099-1103.
803   8.   Yoon, S.I., et al., *Structural basis of TLR5-flagellin recognition and signaling.* Science,
804   2012. **335**(6070): p. 859-64.
805   9.   Ivicak-Kocjan, K., et al., *Determination of the physiological 2:2 TLR5:flagellin*
806   *activation stoichiometry revealed by the activity of a fusion receptor.* Biochem
807   Biophys Res Commun, 2013. **435**(1): p. 40-5.
808   10.   Forstneric, V., et al., *The role of the C-terminal D0 domain of flagellin in activation of*
809   *Toll like receptor 5.* PLoS Pathog, 2017. **13**(8): p. e1006574.
810   11.   Liao, Z., et al., *Cyprinid-specific duplicated membrane TLR5 senses dsRNA as*
811   *functional homodimeric receptors.* EMBO Rep, 2022. **23**(8): p. e54281.
812   12.   Weber, A.N.R., *Fish TLR5 develops a taste for viral RNA.* EMBO Rep, 2022. **23**(8): p.
813   e55443.
814   13.   Andersen-Nissen, E., et al., *Evasion of Toll-like receptor 5 by flagellated bacteria.*
815   Proc Natl Acad Sci U S A, 2005. **102**(26): p. 9247-52.
816   14.   Rhee, S.H., et al., *Pathophysiological role of Toll-like receptor 5 engagement by*
817   *bacterial flagellin in colonic inflammation.* Proc Natl Acad Sci U S A, 2005. **102**(38): p.
818   13610-5.
819   15.   Schlee, M., et al., *Induction of human beta-defensin 2 by the probiotic Escherichia*
820   *coli Nissle 1917 is mediated through flagellin.* Infect Immun, 2007. **75**(5): p. 2399-407.
821   16.   Vijayan, A., et al., *Compartmentalized Antimicrobial Defenses in Response to*
822   *Flagellin.* Trends Microbiol, 2018. **26**(5): p. 423-435.
823   17.   Martini, E., et al., *Mend Your Fences: The Epithelial Barrier and its Relationship With*
824   *Mucosal Immunity in Inflammatory Bowel Disease.* Cell Mol Gastroenterol Hepatol,
825   2017. **4**(1): p. 33-46.
826   18.   Lodes, M.J., et al., *Bacterial flagellin is a dominant antigen in Crohn disease.* J Clin
827   Invest, 2004. **113**(9): p. 1296-306.
828   19.   Morgan, N.N., et al., *Crohn's Disease Patients Uniquely Contain Inflammatory*
829   *Responses to Flagellin in a CD4 Effector Memory Subset.* Inflamm Bowel Dis, 2022.
830   **28**(12): p. 1893-1903.
831   20.   Kreutzberger, M.A.B., et al., *Flagellin outer domain dimerization modulates motility in*
832   *pathogenic and soil bacteria from viscous environments.* Nat Commun, 2022. **13**(1):
833   p. 1422.
834   21.   Andersen-Nissen, E., et al., *A conserved surface on Toll-like receptor 5 recognizes*
835   *bacterial flagellin.* J Exp Med, 2007. **204**(2): p. 393-403.
836   22.   Forstneric, V., et al., *Distinctive Recognition of Flagellin by Human and Mouse Toll-*
837   *Like Receptor 5.* PLoS One, 2016. **11**(7): p. e0158894.
838   23.   Clasen, S.J., et al., *Silent recognition of flagellins from human gut commensal*
839   *bacteria by Toll-like receptor 5.* Sci Immunol, 2023. **8**(79): p. eabq7001.

24. Kearns, D.B., *A field guide to bacterial swarming motility.* Nat Rev Microbiol, 2010.
8(9): p. 634-44.
25. Wang, F., et al., *A structural model of flagellar filament switching across multiple*
bacterial species. Nat Commun, 2017. **8**(1): p. 960.
26. Nedeljkovic, M., et al., *An unbroken network of interactions connecting flagellin*
domains is required for motility in viscous environments. PLoS Pathog, 2023. **19**(5):
p. e1010979.
27. Troge, A., et al., *More than a marine propeller--the flagellum of the probiotic*
Escherichia coli strain Nissle 1917 is the major adhesin mediating binding to human
mucus. Int J Med Microbiol, 2012. **302**(7-8): p. 304-14.
28. Patterson, A.M., et al., *Human Gut Symbiont Roseburia hominis Promotes and*
Regulates Innate Immunity. Front Immunol, 2017. **8**: p. 1166.
29. Gibson, D.G., *Synthesis of DNA fragments in yeast by one-step assembly of*
overlapping oligonucleotides. Nucleic Acids Res, 2009. **37**(20): p. 6984-90.
30. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nat
Methods, 2012. **9**(7): p. 676-82.
31. Ducret, A., E.M. Quardokus, and Y.V. Brun, *MicrobeJ, a tool for high throughput*
bacterial cell detection and quantitative analysis. Nat Microbiol, 2016. **1**(7): p. 16077.
32. van der Walt, S., et al., *scikit-image: image processing in Python.* PeerJ, 2014. **2**: p.
e453.
33. Sugahara, M., et al., *Heavy-atom Database System: a tool for the preparation of*
heavy-atom derivatives of protein crystals based on amino-acid sequence and
crystallization conditions. Acta Crystallogr D Biol Crystallogr, 2005. **61**(Pt 9): p. 1302-
5.
34. Kabsch, W., *Xds.* Acta Crystallogr D Biol Crystallogr, 2010. **66**(Pt 2): p. 125-32.
35. Adams, P.D., et al., *PHENIX: a comprehensive Python-based system for*
macromolecular structure solution. Acta Crystallogr D Biol Crystallogr, 2010. **66**(Pt 2):
p. 213-21.
36. Zwart, P.H., R.W. Grosse-Kunstleve, and P.D. Adams, *Xtriage and Fest: automatic*
assessment of X-ray data and substructure structure factor estimation. CCP4
Newsletter Winter, 2005. **Contribution 7**.
37. Afonine, P.V., et al., *Towards automated crystallographic structure refinement with*
phenix.refine. Acta Crystallogr D Biol Crystallogr, 2012. **68**(Pt 4): p. 352-67.
38. Winn, M.D., et al., *Overview of the CCP4 suite and current developments.* Acta
Crystallogr D Biol Crystallogr, 2011. **67**(Pt 4): p. 235-42.
39. Evans, P.R., *An introduction to data reduction: space-group determination, scaling*
and intensity statistics. Acta Crystallogr D Biol Crystallogr, 2011. **67**(Pt 4): p. 282-92.
40. Matthews, B.W., *Solvent content of protein crystals.* J Mol Biol, 1968. **33**(2): p. 491-7.
41. Stein, N., *CHAINSAW: a program for mutating pdb files used as templates in*
molecular replacement. Journal of Applied Crystallography, 2008. **41**: p. 641-643.
42. McCoy, A.J., et al., *Phaser crystallographic software.* J Appl Crystallogr, 2007. **40**(Pt
4): p. 658-674.
43. Vagin, A. and A. Teplyakov, *Molecular replacement with MOLREP.* Acta Crystallogr D
Biol Crystallogr, 2010. **66**(Pt 1): p. 22-5.
44. Murshudov, G.N., A.A. Vagin, and E.J. Dodson, *Refinement of macromolecular*
structures by the maximum-likelihood method. Acta Crystallogr D Biol Crystallogr,
1997. **53**(Pt 3): p. 240-55.
45. Emsley, P., et al., *Features and development of Coot.* Acta Crystallogr D Biol
Crystallogr, 2010. **66**(Pt 4): p. 486-501.
46. Schrodinger, LLC, *The PyMOL Molecular Graphics System, Version 1.8.* 2015.
47. Sheldrick, G.M., *Experimental phasing with SHELXC/D/E: combining chain tracing*
with density modification. Acta Crystallogr D Biol Crystallogr, 2010. **66**(Pt 4): p. 479-
85.
48. Vonrhein, C., et al., *Automated structure solution with autoSHARP.* Methods Mol Biol,
2007. **364**: p. 215-30.

49.   Abrahams, J.P. and A.G. Leslie, *Methods used in the structure determination of*
bovine mitochondrial F1 ATPase. Acta Crystallogr D Biol Crystallogr, 1996. **52**(Pt 1):
p. 30-42.
50.   Chen, V.B., et al., *MolProbity: all-atom structure validation for macromolecular*
crystallography. Acta Crystallogr D Biol Crystallogr, 2010. **66**(Pt 1): p. 12-21.
51.   Lutz, M.B., et al., *An advanced culture method for generating large quantities of*
highly pure dendritic cells from mouse bone marrow. J Immunol Methods, 1999.
223(1): p. 77-92.

Acknowledgements

We thank Libera Lo Presti for excellent editorial support.

Funding

We thank Libera Lo Presti for excellent editorial support. ANRW was supported by the

University of Tübingen Medical Faculty, Deutsches Konsortium für Translationale

Krebsforschung (German Cancer Consortium, DKTK), the Deutsche

Forschungsgemeinschaft (German Research Foundation, DFG) Research Grant We-

4195/18-1 "ImMiGeNe - Interplay of immune parameters, microbiota and host genetics

during childhood stem cell transplantation", Grant We-4195/25-1 within the Priority Program

2225 "Exit Strategies", the Clusters of Excellence "iFIT – Image-Guided and Functionally

Instructed Tumor Therapies" (EXC-2180) and "CMFI – Controlling Microbes to Fight Infection

(EXC-2124). Gefördert durch die Deutsche Forschungsgemeinschaft (DFG) im Rahmen der

Exzellenzstrategie des Bundes und der Länder – EXC-2180 und EXC-2124. We also

gratefully acknowledge the Volkswagenstiftung support to AS and ANRW (project Innately

Human) and the support of the Open Access publishing funds of University of Tübingen and

the Medical Faculty Tübingen. ME acknowledges funding from the European Research

Council (ERC) under the European Union's Horizon 2020 research and innovation program

(grant agreement no. 864971), the Volkswagenstiftung for its support of the "Life?" program

(grant agreement no. 96732) and from the Max Planck Society as Max Planck Fellow. MH

was supported by the DFG by project HE1964/23-2 within the framework of priority program

SPP2225 'Exit Strategies', and P8 within SFB 1557 'Functional plasticity encoded by

membrane networks'. PF likes to thank the Hans Mühlenhoff-Stiftung for fellowship support

of his PhD. LA gratefully acknowledges funding by DFG (German Research Foundation)

Emmy Noether Programme (Project N. 520465503) and the Damon Runyon Cancer

Research Foundation and CRIS Cancer Foundation (DFS-57-23).

Author contributions (according to Credit guidelines)

Author initials	JJ	MB	KH	IK	GL	PF	MJ GZ	HR	CGK	AS	TH	SW	TS	LA	ME	MH	JSF
Conceptualization	X														X	X	
Data curation																	
Formal analysis	X	X	X			X	X	X		X			X	X			
Funding acquisition													X	X	X	X	
Investigation	X	X	X	X	X	X	X		X		X						
Methodology	X	X	X									X					
Project administration	X												X	X		X	
Resources	X										X	X				X	
Software																	
Supervision												X	X	X		x	
Validation	X	X	X													X	

Jakob, Braun et al., 2026 (confidential)

Visualization	X									X						X
Writing - original draft	X															X
Writing – review and editing	X	X	X	X	X	X	X	X	X	X		X	X	X	X	X

<u>Figure legends</u>

Figure 1: Structure of the EcN flagellin FliC shows a distinct D4 domain and D1-D2
linker.

(A) Cartoon and surface representations of the EcN FliC model comprising the D1 (orange),
D2 (green), D3 (blue) and D4 (yellow). **(B)** Side-view of the D4 β-triangle. **(C)** Interaction of
the two strands of the D1-D2 linker. The hydrogen bonds and ionic interaction are depicted
with black dotted lines. The distance between the D1 and D2 is depicted with a red dotted
line (21 Å) and a yellow dotted line (9 Å). The region close to D2 is stabilized by a β-hairpin.
Three different flagellum models. **(D)** Structure alignment of three EcN D12 structures, side
view with angles between center of mass (COM) of D1, Cα of P507, the COM of D2, and
carbonyl-C of T206, connected with a red line. The structures are shown in cartoon and
surface representations, with D1 (brown) and D2 (green). **(E)** Comparison of ST and EcN
FliC in flagellar context. Cartoon or surface representation axially viewed from the tip towards
the stator of the flagellums, with D0 (purple), D1 (orange), D2 (green), D3 (blue) and D4
(yellow). Top: ST. Middle: closed model of the EcN flagellum. Bottom: open model of the EcN
flagellum.

Figure 2: The D1-D2 linker affects TLR5 recognition of purified EcN flagellin, but not
flagellated bacteria

IL-8 cytokine ELISA of **(A, C)** human (h)TLR5 or **(B, D)** murine (m)TLR5-expressing
HEK293T cells stimulated with **(A, B)** recombinantly isolated FliC proteins or with **(C, D)**
gentamycin-killed bacteria for 24 h. Pooled data, n ≥ 3 experiments; statistics: two
independent tests were done: 2-way ANOVA with Dunnett correction sample versus mock
indicated by #; 2-way ANOVA with Dunnett correction sample versus full-length FliC
indicated by *

Figure 3: Subtle discrimination of purified EcN flagellins but not flagellated bacteria by
intestinal epithelial and immune cells

(A) IL-8 cytokine ELISA and **(B)** human beta defensin 2 (hBD2) expression of C2BBe1 cells

6 h after stimulation with gentamicin-killed bacteria. Pooled data, n ≥ 3; statistics: two

independent statistical tests were performed, 2-way ANOVA with Dunnett correction sample

versus mock indicated by #; 2-way ANOVA with Dunnett correction sample versus EcN WT

indicated by *. **(C)** Heat maps summarizing IL-8 cytokine ELISA data of THP-1, mTLR5-,

hTLR5-HEK293T and/or C2BBe1 following stimulation with recombinantly isolated FliC

proteins (left) and gentamicin-killed bacteria (right) after 24 h or 6 h (C2BBe1). Pooled data,

fractions of pie charts indicate technical replicates (THP-1 data) or mean value of n

independent experiments.

Figure 4: Linker and D4 domain affect motility.

(A) Soft agar swimming distances of FliC mutants depicted as mean of measured diameters

d, n ≥ 3; statistics: ordinary one-way ANOVA with Dunnett correction sample versus EcN WT

indicated by *. **(B)** Representative images for selected bacteria. **(C)** Mean single-cell

swimming velocities in liquid media of wild-type and mutant strains obtained from

exponentially growing cultures. Each dot represents the mean velocity of a trajectory of an

individual bacterium (≥3 s), with dots colored by replicate. At least 1521 individual trajectories

were analyzed per strain and replicate. **(D)** Representative TEM images of negative-stained

isolated flagella from wildtype (WT), hyperflagellated (hyperflag.), *fliC ΔD4*, *fliC Δlinker*, and

fliC ΔHVR strains after background subtraction. All samples were prepared as described in

Materials and Methods. **(E)** Average filament diameters were quantified using ImageJ and

are indicated for each strain. The diameter of 100 flagella per variant was determined by

calculating the mean value from three measurements performed along each filament. Scale

bar: 10 nm. **(F)** The average wave-pitch distances were quantified from 100 flagella per

strain using ImageJ. Scale bars as indicated. **(G)** Representative TEM images of isolated

flagella showing distinct wave-patterns between the different variants. Redlines indicate the

measured wave-pitch distances.

Figure 5: Point mutations in TLR5 interface impair motility but not TLR5 recognition

(A) Soft agar swimming distances of FliC mutants depicted as mean of measured diameters,

n ≥ 3. Statistics: ordinary one-way ANOVA with Dunnett correction sample versus KO:KI

indicated by *. **(B)** Graphical summary of normalized m- and hTLR5 activity plotted against

normalized motility for each mutant. Dashed line on the left indicates lower limit for motility

derived   from   bacterial   volume   on   plate   despite   lacking   motility.

Supplemental information

Supplementary figure legends (included in the actual supplementary figures)

Amino acid sequences for constructs used

All constructs for protein purification contained EcN FliC sequences fused N-terminally to an

8xHis tag (dark grey) connected by a linker (light grey) containing a TEV cleavage site

(green). Domain assignment is shown exemplarily for the full-length EcN FliC. Purple marks

the N and C-terminal D0 domain, pink the D1 domain, orange the linker region, dark red the

D2 domain, light red the D3 domain and blue the D4 domain. In the ΔD4 construct, a short

linker is inserted to connect the N- and C-terminal parts of the D3 (dark green).

Full-length EcN FliC:

1   MGHHHHHHHH SSGHITTENL YFQSMAQVIN TNSLSLITQN NINKNQSALS

SSIERLSSGL

61   RINSAKDDAA GQAIANRFTS NIKGLTQAAR NANDGISVAQ TTEGALSEIN

NNLQRIRELT

121   VQASTGTNSD SDLDSIQDEI KSRLDEIDRV SGQTQFNGVN VLAKDGSMKI

QVGANDGQTI

181   TIDLKKIDSD TLGLNGFNVN GSGTIANKAA TISDLTAAKM DAATNTITTT

NNALTASKAL

241   DQLKDGDTVT IKADAAQTAT VYTYNASAGN FSFSNVSNNT SAKAGDVAAS

LLPPAGQTAS

301   GVYKAASGEV NFDVDANGKI TIGGQEAYLT SDGNLTTNDA GGATAATLDG

LFKKAGDGQS

361   IGFNKTASVT MGGTTYNFKT GADAGAATAN AGVSFTDTAS KETVLNKVAT

AKQGTAVAAN

421   GDTSATITYK   SGVQTYQAVF   AAGDGTASAK   YADNTDVSNA   TATYTDADGE

MTTIGSYTTK

481   YSIDANNGKV   TVDSGTGTGK   YAPKVGAEVY   VSANGTLTTD   ATSEGTVTKD

PLKALDEAIS

541   SIDKFRSSLG   AIQNRLDSAV   TNLNNTTTNL   SEAQSRIQDA   DYATEVSNMS

KAQIIQQAGN

601 SVLAKANQVP QQVLSLLQG

ΔD4 EcN FliC:

1   MGHHHHHHHH   SSGHITTENL   YFQSMAQVIN   TNSLSLITQN   NINKNQSALS

SSIERLSSGL

61   RINSAKDDAA   GQAIANRFTS   NIKGLTQAAR   NANDGISVAQ   TTEGALSEIN

NNLQRIRELT

121   VQASTGTNSD   SDLDSIQDEI   KSRLDEIDRV   SGQTQFNGVN   VLAKDGSMKI

QVGANDGQTI

181   TIDLKKIDSD   TLGLNGFNVN   GSGTIANKAA   TISDLTAAKM   DAATNTITTT

NNALTASKAL

241   DQLKDGDTVT   IKADAAQTAT   VYTYNASAGN   FSFSNVSNGS   GSATDTASKE

TVLNKVATAK

301   QGTAVAANGD   TSATITYKSG   VQTYQAVFAA   GDGTASAKYA   DNTDVSNATA

TYTDADGEMT

361   TIGSYTTKYS   IDANNGKVTV   DSGTGTGKYA   PKVGAEVYVS   ANGTLTTDAT

SEGTVTKDPL

421   KALDEAISSI   DKFRSSLGAI   QNRLDSAVTN   LNNTTTNLSE   AQSRIQDADY

ATEVSNMSKA

481 QIIQQAGNSV LAKANQVPQQ VLSLLQG

ΔLinker EcN FliC:

1 MGHHHHHHHH SSGHITTENL YFQSMAQVIN TNSLSLITQN NINKNQSALS

SSIERLSSGL

61 RINSAKDDAA GQAIANRFTS NIKGLTQAAR NANDGISVAQ TTEGALSEIN

NNLQRIRELT

121 VQASTGTNSD SDLDSIQDEI KSRLDEIDRV SGQTQFNGVN VLAKDGSMKI

QVGANDGQTI

181 TIDLKKIDSD TLGLNGFNVN GNKAATISDL TAAKMDAATN TITTTNNALT

ASKALDQLKD

241 GDTVTIKADA AQTATVYTYN ASAGNFSFSN VSNNTSAKAG DVAASLLPPA

GQTASGVYKA

301 ASGEVNFDVD ANGKITIGGQ EAYLTSDGNL TTNDAGGATA ATLDGLFKKA

GDGQSIGFNK

361 TASVTMGGTT YNFKTGADAG AATANAGVSF TDTASKETVL NKVATAKQGT

AVAANGDTSA

421 TITYKSGVQT YQAVFAAGDG TASAKYADNT DVSNATATYT DADGEMTTIG

SYTTKYSIDA

481 NNGKVTVDSG TGTGKYAPKV GAEVYVSANG TLTTDATTKD PLKALDEAIS

SIDKFRSSLG

541 AIQNRLDSAV TNLNNTTTNL SEAQSRIQDA DYATEVSNMS KAQIIQQAGN

SVLAKANQVP

601 QQVLSLLQG

ΔHVR EcN FliC:

1 MGHHHHHHHH SSGHITTENL YFQSMAQVIN TNSLSLITQN NINKNQSALS

SSIERLSSGL

61   RINSAKDDAA   GQAIANRFTS   NIKGLTQAAR   NANDGISVAQ   TTEGALSEIN

NNLQRIRELT

121   VQASTGTNSD   SDLDSIQDEI   KSRLDEIDRV   SGQTQFNGVN   VLAKDGSMKI

QVGANDGQTI

181   TIDLKKIDSD   TLGLNGFNVN   GSGGTVTKDP   LKALDEAISS   IDKFRSSLGA

IQNRLDSAVT

241   NLNNTTTNLS   EAQSRIQDAD   YATEVSNMSK   AQIIQQAGNS   VLAKANQVPQ

QVLSLLQG

ΔD0 EcN FliC:

1   MGHHHHHHHH   SSGHITTENL   YFQSQAIANR   FTSNIKGLTQ   AARNANDGIS

VAQTTEGALS

61   EINNNLQRIR   ELTVQASTGT   NSDSDLDSIQ   DEIKSRLDEI   DRVSGQTQFN

GVNVLAKDGS

121   MKIQVGANDG   QTITIDLKKI   DSDTLGLNGF   NVNGSGTIAN   KAATISDLTA

AKMDAATNTI

181   TTTNNALTAS   KALDQLKDGD   TVTIKADAAQ   TATVYTYNAS   AGNFSFSNVS

NNTSAKAGDV

241   AASLLPPAGQ   TASGVYKAAS   GEVNFDVDAN   GKITIGGQEA   YLTSDGNLTT

NDAGGATAAT

301   LDGLFKKAGD   GQSIGFNKTA   SVTMGGTTYN   FKTGADAGAA   TANAGVSFTD

TASKETVLNK

361   VATAKQGTAV   AANGDTSATI   TYKSGVQTYQ   AVFAAGDGTA   SAKYADNTDV

SNATATYTDA

421   DGEMTTIGSY   TTKYSIDANN   GKVTVDSGTG   TGKYAPKVGA   EVYVSANGTL

TTDATSEGTV

481 TKDPLKALDE AISSIDKFRS SLGAIQNRLD SAVTNLNNTT TNLSEAQSRI QD

Construct	Mutation	M_W [kDa]	Extinction coefficient [M^{-1} cm^{-1}]	Source
Full-length EcN FliC	-	63.788	22350	This work
∆D4 EcN FliC	∆N255 - F371	52.850	17880	This work
∆Linker EcN FliC	∆S178 – A182 S499 – V503	62.885	22350	This work
∆HVR EcN FliC	∆T180 – E500	31.873	2980	This work, kind gift by T. Hagemann, University of Tübingen
∆D0 EcN FliC	∆M1 - G71 A580 -G619	54.663	20860	Kind gift by T. Hagemann, University of Tübingen

1088 **Supplementary tables**

1089 **Supplementary Table S1: Used kits**

KIT	Manufacturer	Catalog #
ELISA MAX™ Deluxe Set Human IL-8	BioLegend	431504
GeneJET Gel Extraction KIT	Thermo Fisher Scientific	K0692
GenElute bacterial genomic DNA isolation Kit	Merck	NA2100
Gibson assembly MasterMix	NEB	E2611S
High-Capacity RNA-to-cDNA-Kit	Thermo Fisher Scientific	4387406
Human IFNβ ELISA	R&D	DY814-05
Human IL-10 ELISA	BD	555157
Human IL-1β ELISA	R&D	DY201-05
Human IL-6 ELISA	R&D	DY206
Human TNF ELISA	BD	555212
Invitrogen Platinum SuperFi II Polymerase + 5 X Buffer	Thermo Fisher Scientific	12361050
Murine IL-10 ELISA	R&D	DY417-05
Murine IL-1β ELISA	R&D	DY401-05
Murine IL-6 ELISA	BD	555240
Murine TNF ELISA	BD	555268
Pierce™ Silver Stain Kit	Thermo Fisher Scientific	24612
PureYield Plasmid Mini Kit	Promega	A1222
PureYield™ Plasmid Midiprep System	Promega	A2496

RNAse-free DNAse Set (250)	QIAGEN	79256
RNeasy Mini QIAcube Kit	QIAGEN	74116

1090

1091   **Supplementary table S2: Primers**

Name	Use	Sequence 5' to 3'	Target
AWm733_f_R91T R93K E94T EcN FliC	mutagenesis	ttacagaccatcaaaaccctgacggttcaggc ttctaccgggactaactc	R91T R93K E94K EcN FliC
AWm734_r_R91T R93K E94T EcN FliC	mutagenesis	agaagcctgaaccgtcagggtttttgatggtctg taagttgttgttgatttcggacagc	R91T R93K E94K EcN FliC
AWm735_f_R530 Q EcN FliC	mutagenesis	ctgggggctatccaaaaccagttggattcc	R530Q mutation in EcN FliC
AWm736_r_R530 Q EcN FliC	mutagenesis	ggaatccaactggttttggatagcccccag	R530Q mutation in EcN FliC
AWm737_mut-Kit_R530Q_EcN FliC_f	mutagenesis	ccctgggggctatccaaaaccagttggattcc gccg	R530Q mutation in EcN FliC
AWm738_mut-Kit_R530Q_EcN FliC_r	mutagenesis	cggcggaatccaactggttttggatagccccc aggg	R530Q mutation in EcN FliC
AWm741_f EcN FliC TLR5rs wt	mutagenesis	cttacagcgtatccgtgagctgacggttcaggc ttctaccgggac	WT TLR5 recognition site in EcN FliC
AWm742_r EcN FliC TLR5rs wt	mutagenesis	gtcccggtagaagcctgaaccgtcagctcac ggatacgctgtaag	WT TLR5 recognition site in EcN FliC
AWm745_f_EcN	mutagenesis	cttacagaccatccgtgagctgacggttcagg	R91T EcN FliC

FliC R91T		cttctaccgggac	
AWm746_r_EcN FliC R91T	mutagenesis	gtcccggtagaagcctgaaccgtcagctcac ggatggtctgtaag	R91T EcN FliC
AWm751_f_EcN FliC R91T R93K	mutagenesis	cttacagaccatcaaagagctgacggttcagg cttctaccgggac	R91T R93K EcN FliC
AWm752_r_EcN FliC R91T R93K	mutagenesis	gtcccggtagaagcctgaaccgtcagctctttg atggtctgtaag	R91T R93K EcN FliC
AWm753_f_EcN FliC R91T E94T	mutagenesis	cttacagaccatccgtaccctgacggttcaggc ttctaccgggac	R91T E94T EcN FliC
AWm754_r_EcN FliC R91T E94T	mutagenesis	gtcccggtagaagcctgaaccgtcagggtac ggatggtctgtaag	R91T E94T EcN FliC
AWm755_f_EcN FliC R93K E94T	mutagenesis	cttacagcgtatcaaaaccctgacggttcagg cttctaccgggac	R93K E94T EcN FliC
AWm756_r_EcN FliC R93K E94T	mutagenesis	gtcccggtagaagcctgaaccgtcagggttttg atacgctgtaag	R93K E94T EcN FliC
AWq_026_GAPD H_F1	qPCR	agccacatcgctcagacac	GAPDH mRNA
AWq_027_GAPD H_R1	qPCR	gcccaatacgaccaaatcc	GAPDH mRNA
hBD2 for	qPCR	gactcagctcctggtgaagctc	human beta defensin 2 mRNA
hBD2 rev	qPCR	ctataccaccaaaaacacctggaag	human beta defensin 2 mRNA

hBD1 for	qPCR	gagatggcctcaggtggtaac	human beta defensin 1 mRNA
hBD1 rev	qPCR	ttggtaaagatcgggcaggc	human beta defensin 1 mRNA
AWs211_pET19b_backbone_f	sequencing	gagtattcaacatttccgtgtcg	pET19b backbone
AWs214_Cterm_Linker_sequencing_f	sequencing	agcgcaaaatatgccgataa	C term Linker EcN FliC
AWs215_Nterm_Linker_sequencing_f	sequencing	tacagcgtatccgtgagctg	N term Linker EcN FliC
AWs217_genomicFliC_f	sequencing	gataacagggttgacggcga	genomic FliC
AWs218_genomicFliC_r	sequencing	tgtaggcctgataagcgcag	genomic FliC
AWs219_pSB890_seq_f	sequencing	cgtcaccaaatgatgttattcc	pSB890
AWs220_pSB890_seq_r	sequencing	gttgagaagcggtgtaagtg	pSB890
AWs221_malEK_f_colony-seq	sequencing	tccatcctccttgcccctac	malEK cluster
AWs222_malEK_	sequencing	acgcggcttctgtgaactaa	malEK cluster

r_colony-seq				
AWs223_R531Q EcN FliC_f	sequencing	cgaaagtcggggctgaagta	R531Q mutation in EcN FliC	
AWx435_EcN-dLinker_Nterm_f	Gibson assembly	ggaattgtgagcggataacaattcccctctag	pET19b FliC	EcN
AWx436_EcN-dLinker_Nterm_r	Gibson assembly	cgccgctttattaccattcacgttaaaaccattc agccccag	pET19b FliC	EcN
AWx437_EcN-dLinker_HVR_for	Gibson assembly	tttaacgtgaatggtaataaagcggcgaccatt agcgacctg	pET19b FliC	EcN
AWx438_EcN-dLinker_HVR_rev	Gibson assembly	agtggatcttttgtagttgcatctgttgttaaagta ccattagcactaac	pET19b FliC	EcN
AWx439_EcN-dLinker_Cterm_f	Gibson assembly	caacagatgcaactacaaaagatccactgaa agctctgg	pET19b FliC	EcN
AWx440_EcN-dLinker_Cterm_r	Gibson assembly	catgtttgacagcttatcatcgataagctttaatg	pET19b FliC	EcN
AWx441_EcN-dLinker_for	colony PCR	cgtgaatggtaataaagcggcgac	EcN ΔLinker	FliC
AWx442_EcN-dLinker_rev	colony PCR	cagtggatcttttgtagttgcatctgttg	EcN ΔLinker	FliC
AWx443_dLinker _short_f	colony PCR	gtgaatggtaataaagcg	EcN ΔLinker	FliC
AWx444_dLinker _short_r	colony PCR	gatcttttgtagttgcatc	EcN ΔLinker	FliC
AWx465_chr-	Gibson	gttattccgcgaaatataatgaccctcttgtattg	chromosomal	

EcN_dLinker_f1	assembly	gcggtctggaaagtc	EcN FliC
AWx466_chr-EcN_dLinker_Nterm_r	Gibson assembly	tgtcaggtcgctaatggtcgccgctttattaccat tcacgttaaaaccattcagc	chromosomal EcN FliC
AWx467_chr-EcN_dLinker_HVR_f	Gibson assembly	ctggggctgaatggttttaacgtgaatggtaata aagcggcgaccattagcgac	chromosomal EcN FliC
AWx468_chr-EcN_dLinker_r1	Gibson assembly	ccaccgcggtggggcttttattgagcttgactgt gccttcgctagttgc	chromosomal EcN FliC
AWx469_chr-EcN_dLinker_f2	Gibson assembly	gttattccgcgaaatataatgaccctcttgacca ttagcgacctgacagc	chromosomal EcN FliC
AWx470_chr-EcN_dLinker_HVR_r	Gibson assembly	ttcatccagagctttcagtggatcttttgtagttgc atctgttgttaaagtaccattagcac	chromosomal EcN FliC
AWx471_chr-EcN_dLinker_Cterm_f	Gibson assembly	gctaatggtactttaacaacagatgcaactaca aaagatccactgaaagctctgg	chromosomal EcN FliC
AWx472_chr-EcN_dLinker_r2	Gibson assembly	ccaccgcggtggggcttttattgagcttgcgttc cgctacctcagtttc	chromosomal EcN FliC
AWx473_chr_Nterm_dLinker_colony_r	colony PCR	agtaccattagcactaac	EcN FliC ΔLinker
AWx474_chr_Cterm_dLinker_colony_f	colony PCR	gctgcaactaatactataac	EcN FliC ΔLinker

AWx509_gib_dHVR_EcN_r	Gibson assembly	agctttcagtggatcttttgttactgtgccaccgg aaccattcacgtt	EcN FliC ΔHVR
AWx510_gib_dHVR_EcN_f	Gibson assembly	ctgaatggttttaacgtgaatggttccggtggca cagtaacaaaagatccactg	EcN FliC ΔHVR
AWx515_gib_uni_890_f2	Gibson assembly	caagagggtcattatatttcgcg	pSB890
AWx516_gib_uni_890_r2	Gibson assembly	caagctcaataaaaagccccac	pSB890
AWx519_gib_malEK_890_f	Gibson assembly	gttattccgcgaaatataatgaccctcttgttattg accgcttccagacc	*E. coli* malEK cluster
AWx520_gib_malEK_890_r	Gibson assembly	ccaccgcggtggggctttttattgagcttgagcc agacttgctgacgatt	*E. coli* malEK cluster
AWx521_p15A-J23119_f	Gibson assembly	cctagggatatattccgcttcc	pdCas9_J23119
AWx522_p15A-J23119_r	Gibson assembly	agatcctttctcctctttagatcac	pdCas9_J23119
AWx523_RFP_hakims-plasmid_f	Gibson assembly	tactagtgatctaaagaggagaaaggatctat ggcttcctccgaagacgt	RFP
AWx524_RFP_hakims-plasmid_r	Gibson assembly	gtgagcgaggaagcggaatatatccctaggtt aagcaccggtggagtgacg	RFP
AWx525_mCherry_hakims-plasmid_f	Gibson assembly	tagtgatctaaagaggagaaaggatctatggg atccgtctccaagggcgag	mCherry
AWx526_mCherry_hakims-	Gibson assembly	gtgagcgaggaagcggaatatatccctaggtt acttgtacagctcgtccatgc	mCherry

plasmid_r

AWx527_malEK insert_f	Gibson assembly	cgcacataaaatcgccatga	malEK cluster in pSB890-malEK
AWx528_malEK insert_r	Gibson assembly	ctgtcaaagatctttagaatgcattatatccatct ccacattaccgcca	malEK cluster in pSB890-malEK
AWx529_J23119 _f	Gibson assembly	gatataatgcattctaaagatctttgacagc	fluorophore + J23119 promoter
AWx530_RFP_r	Gibson assembly	cttgcaaaaatcatggcgattttatgtgcgttaa gcaccggtggagtgacg	RFP
AWx531_mCherr y_r	Gibson assembly	cttgcaaaaatcatggcgattttatgtgcgttact tgtacagctcgtccatgc	mCherry
AWx534a_f_colo ny R91T R93K E94K EcN FliC	colony PCR	agaacgtggactccaacgtc	R91T R93K E94K EcN FliC
AWx535a_r_colo ny R91T R93K E94K EcN FliC	colony PCR	ctgaaccgtcagggttttgatggt	R91T R93K E94K EcN FliC
AWx536a_f_colo ny altern. silent TLR5rs	colony PCR	accatcaaaaccctgacggttcag	R91T R93K E94K EcN FliC
AWx537a_r_colo ny altern. silent TLR5rs	colony PCR	gttcacttcaccgcttgctg	R91T R93K E94K EcN FliC
AWx538a_ApaI	Gibson	ctatagggcgaattgggcccga	pSB890 MCS

recon for	assembly		
AWx539a_ClaI recon rev	Gibson assembly	cttgccgttgttagcatcgattg	R91T R93K E94K EcN FliC
AWx540_EcN FliC ND3 rev	Gibson assembly	gcagtttgagctgcatc	R531Q mutation in EcN FliC
AWx541_LacOp revcomp for	Gibson assembly	ctgtgtgaaattgttatccg	R531Q mutation in EcN FliC
AWx542_'LacOp for' rev	sequencing	cggataacaatttcacacag	R531Q mutation in EcN FliC
AWx543_synth. J23119+RBS_1_f	Gibson assembly	tgacagctagctcagtcctaggtataatactagt gatctaaagaggagaaagg	J23119 + RBS
AWx544_synth. J23119+RBS_2_f	Gibson assembly	gatataatgcattctaaagatctttgacagctag ctcagtcctagg	J23119
AWx545_elongat ed AWx539_r	Gibson assembly	ggcggtaatgtggagatggatataatgcattct aaagatctttgacagctagc	flourophore + J23119 promoter
AWx546_f gib EcN FliC TLR5rs wt	Gibson assembly	ttacagcgtatccgtgagctgacggttcaggctt ctaccgggactaactc	WT TLR5rs in EcN FliC
AWx547_r gib EcN FliC TLR5rs wt	Gibson assembly	agaagcctgaaccgtcagctcacggatacgc tgtaagttgttgttgatttcggacagc	WT TLR5rs in EcN FliC
AWx548_r_R91T _EcN FliC	Gibson assembly	agaagcctgaaccgtcagctcacggatggtct gtaagttgttgttgatttcggacagc	R91T EcN FliC
AWx549_f_R91T	Gibson	ttacagaccatccgtgagctgacggttcaggct	R91T EcN FliC

_EcN FliC	assembly	tctaccgggactaactc	
AWx550_r_R93K _EcN FliC	Gibson assembly	agaagcctgaaccgtcagctctttgatacgctg taagttgttgttgatttcggacagc	R93K EcN FliC
AWx551_f_R93K _EcN FliC	Gibson assembly	ttacagcgtatcaaagagctgacggttcaggct tctaccgggactaactc	R93K EcN FliC
AWx552_r_E94T _EcN FliC	Gibson assembly	agaagcctgaaccgtcagggtacggatacgc tgtaagttgttgttgatttcggacagc	E94T EcN FliC
AWx553_f_E94T _EcN FliC	Gibson assembly	ttacagcgtatccgtaccctgacggttcaggctt ctaccgggactaactc	E94T EcN FliC
AWx554_r_R91T-R93K_EcN FliC	Gibson assembly	agaagcctgaaccgtcagctctttgatggtctgt aagttgttgttgatttcggacagc	R91T R93K EcN FliC
AWx555_f_R91T-R93K_EcN FliC	Gibson assembly	ttacagaccatcaaagagctgacggttcaggc ttctaccgggactaactc	R91T R93K EcN FliC
AWx556_r_R91T-E94T_EcN FliC	Gibson assembly	agaagcctgaaccgtcagggtacggatggtct gtaagttgttgttgatttcggacagc	R91T E94T EcN FliC
AWx557_f_R91T-E94T_EcN FliC	Gibson assembly	ttacagaccatccgtaccctgacggttcaggct tctaccgggactaactc	R91T E94T EcN FliC
AWx558_r_R93K -E94T_EcN FliC	Gibson assembly	agaagcctgaaccgtcagggttttgatacgctg taagttgttgttgatttcggacagc	R93K E94T EcN FliC
AWx559_f_R93K-E94T_EcN FliC	Gibson assembly	ttacagcgtatcaaaaccctgacggttcaggct tctaccgggactaactc	R93K E94T EcN FliC

1092 **Supplementary table S3: Plasmids**

Name	Insert	Backbone	Resistance
pJJ001	full EcN FliC + 8xHis-tag TEV cleavable	pET19b	Ampicillin
pJJ002	none	pET19b	Ampicillin
pJJ003	EcN FliC ΔLinker	pET19b	Ampicillin
pJJ004	upstream chromosomal flanking region + N term of EcN FliC lacking N term Linker	pSB890	Tetracyclin
pJJ005	C term of EcN FliC lacking N term Linker + downstream chromosomal flanking region	pSB890	Tetracyclin
pJJ006	EcN FliC lacking HVR + chromosomal flanking regions	pSB890	Tetracyclin
pJJ008	EcN FliC R91T R93K E94T	pGEM-T easy	Ampicillin
pJJ007	intergenomic region + mal E and K cluster beginning	pSB890	Tetracyclin
pJJ009	EcN FliC R91T R93K E94T R531Q	pGEM-T easy	Ampicillin
pJJ010	malEK cluster containing J23119 mCherry in intergenomic region	pSB890	Tetracyclin
pJJ011	malEK cluster containing J23119 RFP in intergenomic region	pSB890	Tetracyclin
pJJ012	EcN FliC WT	pGEM-T easy	Ampicillin
pJJ013	EcN FliC R91T	pGEM-T easy	Ampicillin
pJJ014	EcN FliC R93K	pGEM-T easy	Ampicillin
pJJ015	EcN FliC E94T	pGEM-T easy	Ampicillin

pJJ016	EcN FliC R91T R93K	pGEM-T easy	Ampicillin
pJJ017	EcN FliC R91T E94T	pGEM-T easy	Ampicillin
pJJ018	EcN FliC R93K E94T	pGEM-T easy	Ampicillin
pJJ019	EcN FliC R531Q	pGEM-T easy	Ampicillin
pJJ020	EcN FliC R91T R531Q	pGEM-T easy	Ampicillin
pJJ021	EcN FliC R93K R531Q	pGEM-T easy	Ampicillin
pJJ022	EcN FliC E94T R531Q	pGEM-T easy	Ampicillin
pJJ023	EcN FliC R91T R93K R531Q	pGEM-T easy	Ampicillin
pJJ024	EcN FliC R91T E94T R531Q	pGEM-T easy	Ampicillin
pJJ025	EcN FliC R93K E94T R531Q	pGEM-T easy	Ampicillin

1093 **Supplementary table S4: Purification buffers**

1094 Buffers used for the purification of FliC. Two different size exclusion buffers were used, for

1095 crystallization experiments the SEC-buffer, and for biological experiments DPBS-buffer. The

1096 pH of the Buffers where adjusted at 4°C.

Buffer	Components
Lysis Buffer / His_A Buffer	50 mM Tris pH 8.0
	300 mM NaCl
His_B Buffer	50 mM Tris pH 8.0
	300 mM NaCl
	500 mM imidazole
SEC-Buffer	20 mM HEPES pH 7.5
	150 mM NaCl
DPBS-Buffer	0.90 mM $CaCl_2$
	0.49 mM $MgCl_2$
	2.67 mM KCl
	1.47 mM KH_2PO_4
	137.93 mM NaCl
	8.06 mM Na_2HPO_4

1097

Figure 1

A D2 D3 D4 D1

B β-triangle

C D1 D2

D 105° 136° 107° 134° 92° 130°

E
D0   D0 D1   D0 D1 D2   D0 D1 D2 D3   D0 D1 D2 D3

231 Å

ST

EcN

close
D0   D0 D1

211 Å

open
D0   D0 D1

252 Å

D0 D1 D2 D3 D4   D0 D1 D2 D3 D4

262 Å   211 Å

D0   D0 D1   D0   open   D0   closed

D1
D2
D3
D4

F

Figure 2

Figure 3

Figure 4

Figure 5