The nanoscale mobility of calcium channels is driven by readily releasable synaptic vesicles to support precise neurotransmission in live *C. elegans*

Yunke Zhao[1*], Dai Zhai[1] and Fabien Pinaud[1,2*]

[1]Department of Biological Sciences, University of Southern California, 1050 Child Way, Los Angeles, CA 90089, USA.

[2]Department of Physics and Astronomy, University of Southern California, 1050 Child Way, Los Angeles, CA 90089, USA.

*Correspondence should be addressed to Y.Z. (yunkezha@usc.edu) and F.P. (pinaud@usc.edu)

Abstract

Precise neurotransmission requires voltage-gated calcium channels (VGCCs) to be tightly coupled with synaptic vesicles (SVs) at active zones (AZs). As VGCCs are mobile in the fluid presynaptic membrane, it remained unclear how they spatiotemporally align with SVs for reliable synaptic release. Using super-resolution imaging and single-molecule tracking in *C. elegans*, we report that two diffusive behaviors of N-type VGCCs at AZs are governed by primed SVs via distinct mechanisms. VGCCs diffuse slowly in multiple nanodomains (∼100 nm) that spatially correlate with UNC-10/RIM, and the channel mobility and nanodomain numbers are coupled to SNAREs assembly and SV priming levels via RAB-3/UNC-10/UNC-13L tripartite complexes. Outside nanodomains, a faster mobility of VGCCs and their AZ distribution scale with UNC-13S-mediated priming under an antagonistic regulation by UNC-10 and TOM-1/Tomosyn. These findings reveal unexpected mechanisms whereby pools of readily releasable SVs actively control the nanoscale mobility and nanodomain organization of VGCCs to support precise neurotransmission.

Keywords

Nanoscale diffusion, Super-resolution imaging, Voltage-gated calcium channels, Active zone, Synaptic vesicle release, UNC-10, UNC-13, SNARE

Introduction

At the presynaptic active zone (AZ), exocytosis of synaptic vesicles (SVs) is spatiotemporally orchestrated by an evolutionarily conserved release machinery[1–3] that includes multi-subunits voltage-gated calcium channels (VGCCs), which provide crucial calcium influx for SV release[4,5]. In both mammals and *C. elegans* nematodes, VGCCs are recruited in close proximity to SVs mainly by the AZ modulator UNC-10/RIM, which interacts with the channel's α1 pore-forming subunit UNC-2/Ca$_v$2 and binds vesicular Rab-3 to dock SVs[6–9]. UNC-10/RIM also directly regulates UNC-13/Munc-13, which primes SVs to become readily releasable[10–12]. During priming, UNC-13/Munc-13 tethers SVs to the plasma membrane and coordinates the assembly of SNARE complexes by conformationally opening and aligning UNC-64/Syntaxin-1 (t-SNARE) with both SNAP-25 (t-SNARE) and vesicular SNB-1/VAMP (v-SNARE), which drive SV fusion upon calcium entry[13–16]. By interacting with UNC-13/Munc-13, UNC-10/RIM couples VGCCs to readily releasable SVs, awaiting calcium influx[17]. The local density of VGCCs and their coupling distances to SVs are therefore key determinants of the SV release probability and its efficacy[18,19]. Indeed, synapses having slow release kinetics are characterized by a relatively broad, microdomain coupling between VGCCs and SVs (>100 nm), while fast-releasing synapses adopt a much narrower, nanodomain coupling (<100 nm)[20,21].

While precise spatial coupling between VGCCs and SVs is critical for efficient neurotransmissions, maintaining such nanoscale proximities is complicated by the lateral diffusion of VGCCs in the fluid presynaptic membrane, as observed in diverse synaptic systems[22–24]. Indeed, the mobility of VGCCs at AZs has emerged as a new determinant of presynaptic release, with VGCC mobilities scaling with intracellular calcium levels and neural activities, and alteration of channel diffusion shaping short-term synaptic plasticity and homeostasis[23–25]. Because VGCCs are not static, one would expect their coupling distances to SVs and release probabilities to shift continuously. How the AZ organization maintains precise spatiotemporal coupling between mobile VGCCs and SVs for reliable synaptic releases remains unclear.

Here, we describe how readily releasable SVs actively drive the nanoscale mobility and spatial organization of N-type VGCCs within intact neuromuscular synapses of live *C. elegans*. Using single particle tracking and super-resolution imaging, we characterized two distinct diffusive behaviors of VGCCs at AZ. We show that VGCCs diffuse slowly within multiple intra-AZ nanodomains (∼100 nm) that are organized in the typical dense-projection-like landscape of *C.*

elegans and that spatially correlate with UNC-10. Outside these nanodomains, the channels adopt

a faster diffusion mode while being confined within the larger, diffraction-limited AZ puncta (~300

nm). We further show that the two mobilities of VGCCs are governed by SV priming and SNARE

complex assembly under the coordination of UNC-10, but by different UNC-13 isoforms.

Specifically, the slow mobility and nanodomain number of VGCCs are predominantly regulated

by UNC-13L-mediated priming through tripartite SV coupling complexes involving RAB-3,

UNC-10 and UNC-13L, while the fast mobility and AZ distribution of VGCCs outside

nanodomains scale with the pool of UNC-13S-primed SVs under antagonistic regulations by

UNC-10 and the t-SNAREs inhibitor TOM-1/Tomosyn. We propose that the heterogeneous

mobilities and spatial organizations of N-type VGCCs provide distinct channel-vesicle coupling

strategies for different pools of readily releasable SVs, offering a tunable mechanism to set SV

release probabilities.

Result

N-type VGCCs exhibit two distinct confined diffusive behaviors in live *C. elegans*

At neuromuscular junction (NMJ) synapses in *C. elegans*, the single N- and P/Q-type VGCCs

(UNC-2/Cav2) mediate both evoked and tonic synaptic releases[26]. The functions and surface

trafficking of UNC-2/Cav2 require the extracellular α2δ subunit UNC-36, whose own AZ

localization also depends on UNC-2/Cav2[26–28]. To study the dynamic distributions and nanoscale

mobilities of VGCC complexes, we used complementation activated light microscopy (CALM)

with single molecule sensitivity in intact living *C. elegans*[29,30]. We endogenously tagged UNC-

36/α2δ with a N-terminal dark split sfCherry1-10[31] (sCherry-UNC-36) using CRISPR/Cas9[32]. To

activate the fluorescent signals of the extracellular-facing sCherry-UNC-36, we introduced a

synthetic complementary peptide (sfCherry11) via pseudocoelomic microinjections[29,30], which

allows for specific detections of surface VGCC without interference from intracellular background

signals (Fig. 1a). We previously reported that N-terminal tagging of split-fluorescent proteins to

UNC-36/α2δ does not alter the conducting activities of VGCCs[29] and animals expressing sCherry-

UNC-36 had functional channels and exhibited normal locomotion compared to N2 Bristol wild-

type (WT) animals[33] (Fig. S1a). After fluorescence activation by CALM, surface sCherry-UNC-

36 showed puncta localization along the nerve cords as reported[29] and colocalized with

endogenous UNC-2/Cav2[27] (Fig. 1b; Fig S1b). To further confirm the AZ localization of sCherry-

UNC-36, we endogenously tagged the AZ marker UNC-10/RIM with 5xMyc and UNC-2/Cav2 with HA at the N-terminus. Those transgenic animals showed normal locomotion and synaptic release compared to WT (Fig. S1c and S1d). Consistent with previous findings[9,27,28], N-type VGCCs puncta represented by either surface sCherry-UNC-36 or HA-UNC-2 colocalized with Myc-UNC-10 puncta at AZs, and overlapped with NMJ boutons along the nerve cords (Fig. S1e and S1f).

In *C. elegans*, UNC-36/α2δ is also required for the function of L-type Egl-19/Cav1 channels in muscles, but is dispensable for Egl-19/Cav1-mediated tonic releases at NMJs[26,34]. To test whether synaptic sCherry-UNC-36 also represents L-type VGCCs, we probed the extent to which UNC-36/α2δ colocalizes with Egl-19/Cav-1 at nerve cords versus muscles. In contrast to UNC-2/Cav2, endogenous Egl-19/Cav1[35] showed a broad distribution along nerve cords, with little co-localization with either GFP-UNC-2 or surface sCherry-UNC-36 (Fig. S1f-S1h). This is consistent with previous findings that UNC-2/Cav2 and Egl-19/Cav1 localize to distinct synaptic regions[36,37] and aligns with the reported function of UNC-36/α2δ, which is required for UNC-2/Cav2 but not Egl-19/Cav1 mediated synaptic releases at NMJs[26]. The dispersed distribution of neuronal Egl-19/Cav1 was also in stark contrast to the puncta distribution of muscular Egl-19/Cav1, which extensively colocalized with surface sCherry-UNC-36 at the sarcolemma of muscles[29,30,34] (Fig. S1i and S1j). Together, these results indicate that UNC-36/α2δ is predominantly associated with UNC-2/Cav2 at AZs rather than with Egl-19/Cav1.

To track individual VGCCs by CALM, the sub-diffraction localizations of single sCherry-UNC-36 signals were extracted with a precision of ~20 nm and linked frame to frame to construct the diffusion trajectory[38] of each channel (Fig. 1b; Fig. S2a). At AZs, individual VGCCs exhibited heterogeneous diffusion dynamics with single-step sizes ranging from a few tens to hundreds of nanometers per frame (Fig. 1b; Fig. S2b; Movie S1-S2). To quantitatively define these different mobilities, we analyzed pooled trajectories (n=42,391 from 58 animals) based on the probability distribution of their squared displacements[29,30,39]. We found that VGCCs have two distinct confined diffusive behaviors at AZs in live *C. elegans* (Fig. 1c). The major fraction of VGCC (81%) exhibited a slow mobility, with a diffusion coefficient D_{slow} of 0.0075 ± 0.0010 μm^2/s and were confined within circular nanodomains 104 ± 23 nm in diameter. The remaining 19% of VGCCs displayed a five-fold faster diffusion ($D_{fast} = 0.038 \pm 0.008$ μm^2/s) and were confined within larger domains 289 ± 29 nm in diameter (Fig. 1c). Within individual VGCC trajectories,

alternations between slow and fast diffusions were observed (Fig. 1d; Fig. S2c), indicating that

D_{slow} and D_{fast} represent two diffusive behaviors of single channels rather than two independent

populations of VGCCs. Later observations also show that removal of AZ proteins that directly

interact with UNC-2 alters both diffusions, supporting that D_{fast} reflects a genuine faster mobility

of VGCC complexes rather than a transient dissociation of UNC-36 from UNC-2. While these

alternating diffusive behaviors of VGCCs in *C. elegans* are unique, their respective diffusion

coefficients are similar to those of Cav2 in different organisms. For instance, D_{slow} matches the

mean diffusion coefficient reported for Cac/Cav2 in *Drosophila* NMJs (0.0075 $\mu m^2/s$)[25], while

D_{fast} is similar to the mobility reported for Cav2.1 in rodent hippocampal synapses (0.04 $\mu m^2/s$)[23].

The small confinement size of VGCCs falls between those measured for rodent Cav2.1 (~80 nm)[23]

and Drosophila Cac/Cav2 (~117 nm)[25], while their larger confinement matches the expected size

of AZ in *C. elegans*[40]. This suggests that the nanoscale mobilities of VGCCs reflect intrinsic

structural features of AZs[41], which we studied next.

N-type VGCCs diffuse in multi-nanodomains that spatially correlate with UNC-10/RIM at

AZs

The hallmark of *C. elegans* AZ is an electron dense projection (DP) with branched ultrastructure

and ~ 100 nm bay-like sites for SV docking and release[42,43]. We therefore determined whether the

confinement of VGCCs in nanodomains reflects those AZ sub-structures by resolving the

nanoscale distribution of the channels and that of UNC-10 using stimulated emission depletion

(STED) microscopy. Interestingly, we found that VGCCs and UNC-10 form multiple co-localizing

nanodomains within individual AZ puncta, in agreement with our diffusion analyses (Fig. 1e and

1f; Fig. S3a-d). Notably, the spatial organization of these nanodomains closely resembled the bay-

like sites between DP branches[42], consistent with the known functions of UNC-10/RIM in docking

SVs and coupling VGCCs[1,3,44]. Similar to the variability in the number of docking bays across

different DPs[42], we also observed heterogeneities in nanodomain numbers per AZ. Most AZs

(~50%) contain 4-5 VGCC or UNC-10 nanodomains, 28-37% contain more than 6 nanodomains,

while 15-22% have 1-3 nanodomains (Fig. S3e). Quantitative analyses showed a significant

correlation between the numbers of VGCC and UNC-10 nanodomains, both exhibiting a median

of five nanodomains per AZ along the dorsal nerve cords (interquartile range [IQR]: 4/6) (Fig.1g).

While not strictly identical, the nanodomains of VGCCs and UNC-10 showed similar spatial

distribution, with a median nearest neighbor distance (NND) of 105 nm (IQR: 87/122nm) and 106

nm (IQR: 90/121nm), respectively (Fig. 1h). The size of VGCC nanodomains (median Area

Equivalent Diameter AED: 103 nm; IQR: 88/121nm) also aligned with the size of UNC-10

nanodomains (106 nm; IQR: 93/122nm) and importantly, closely matched the confinement size of

slow-diffusing VGCCs extracted by diffusion analyses (104 nm ± 23 nm) (Fig. 1c and 1h). This

indicates that the slow mobility (81%) of VGCCs reflects their diffusion within individual

nanodomains that spatially correlate with UNC-10.

Within each AZ, multiple VGCCs or UNC-10 nanodomains form an elliptic punctum similar to

the overall DP geometry established by electron microscopy[42,43] (Fig. 1f). The size of VGCC

puncta, defined by their long and short axes (median: 334 x 240 nm), strongly correlates with that

of UNC-10 (median: 339 x 248 nm) (Fig.1i and 1j; Fig. S3f). To adequately compare these

parameters with the circular confinement size of fast-diffusing VGCCs established by diffusion

analyses, we also calculated their area equivalent diameter (AED). The median AED of VGCCs

was 291 nm (IQR: 260/322nm) and that of UNC-10 was 301 nm (IQR: 265/340nm) (Fig. 1j). Both

sizes are consistent with the confined domain (289 ± 29 nm) of fast-diffusing VGCCs, suggesting

that the fast mobility (19%) reflects their exploration across the multi-nanodomain AZ area.

UNC-10 enhances the diffusion of VGCCs and promotes their multi-nanodomain

organization

To investigate whether VGCC nanodomains are organized by UNC-10, we examined their

nanoscale distribution in an *unc-10(md1117)*[8] null mutant. *Unc-10(md1117)* animals showed

severely impaired neurotransmission and fewer VGCC puncta along the nerve cords as previously

described[9,45] (Fig. 2a; Fig. S4a-4c). By STED, we found that, in the absence of UNC-10, VGCC

nanodomains persist (Fig. 2b), but their median number decreases significantly from 5 to 2 per

punctum (Fig. 2c). The percentage of VGCC puncta containing more than 3 nanodomains dropped

from 85% in WT animals to 11% in *unc-10(md1117)* mutants (Fig. 2d). Meanwhile, the NND

between VGCC nanodomains also significantly increased, resulting in a more dispersed

organization compared to WT (Fig. 2e). These results show that, while VGCC nanodomains form

independently of UNC-10, UNC-10 promotes their accumulation at AZs and maintains them in a

tight spatial organization.

We then studied the mobility of individual VGCCs in *unc-10(md1117)* mutants using CALM. The
two diffusive behaviors of VGCCs were still observed in these mutants and their relative fractions
remained unchanged compared to WT (Fig. 2f), suggesting that UNC-10 does not impact the
ability of VGGCs to diffuse inside or outside nanodomains. However, VGCCs were significantly
slower in *unc-10(md1117)* mutants, with both D_{slow} within nanodomains and D_{fast} across AZ being
reduced by half compared to WT (Fig. 2g). This is consistent with a similar decrease in Cav2
diffusion observed after removal of RIM at *Drosophila* NMJs[25]. The channels were also
significantly more confined within nanodomains (*unc-10*: 78 ± 23 nm *vs.* WT: 104 ± 23 nm), but
their diffusion range across AZ was maintained at WT levels (*unc-10*: 270 ± 28 nm *vs.* WT: $289 \pm$
29 nm) (Fig. 2h). Together, this indicates that UNC-10 enhances the mobilities of VGCCs both
inside and outside nanodomains, in addition to its role as a spatial organizer of VGCC
nanodomains at AZs.
203
The mobility and spatial organization of nanodomain-associated VGCCs are regulated by
UNC-10/RIM mediated coupling to synaptic vesicles
To understand how UNC-10, as a cytomatrix protein, enhances the mobility of VGCCs, we
investigated whether the channel diffusion reflects their physical coupling to SVs through UNC-
10. As UNC-10 docks SVs by binding to vesicular RAB-3[12,46], we examined if the mobility of
VGCCs would be impacted in *rab-3(js49)* mutants[46], which have SV docking defect similar to that
of *unc-10(md1117)* animals[47,48]. *Rab-3(js49)* mutants exhibited reduced neurotransmission[48],
showing impaired locomotion and increased resistance to aldicarb (Fig. 3a and 3b). However, the
loss of RAB-3 had no significant influence on the diffusion coefficients of VGCCs, suggesting
that docking defects alone do not account for the decreased channel mobilities observed in *unc-*
10(md1117) mutants (Fig. 3c and 3d; Fig. S5a). UNC-10 also binds the priming factor UNC-
13L/Munc-13, which physically bridges SVs with the presynaptic membrane[49,50]. We thus
examined the channel mobilities in *unc-10(nu487)* animals carrying K77E/K79E mutations, which
disrupt interactions between the ZnF domain of UNC-10 and UNC-13L[11,12,17]. *Unc-10(nu487)*
mutants were reported to have reduced release probability[17] and they showed mild uncoordinated
phenotypes with near-normal aldicarb-resistance (Fig. 3a and 3b). Again, the diffusion coefficients
of VGCCs in these mutants remained similar to WT (Fig. 3c and 3d; Fig. S5b), indicating that
dissociating UNC-10 from UNC-13L is not sufficient to reduce the channel mobilities.

Interestingly, when both interactions of UNC-10 with RAB-3 and UNC-13L were simultaneously disrupted in *rab-3;unc-10(nu487)* double mutants, VGCCs within nanodomains exhibited significantly reduced diffusion coefficients (D_{slow}) and increased confinement (double mutant: 74 ± 23 nm *vs.* WT: 104 ± 23 nm), phenocopying changes observed in *unc-10(md1117)* animals (Fig. 3c; Fig. S5c). On the contrary, D_{fast} for VGCCs outside nanodomains remained at WT levels (Fig. 3d). This indicates that the two diffusive behaviors of VGCCs are regulated by UNC-10 via distinct mechanisms, with the channel mobility in nanodomains being driven by UNC13L/RAB-3-dependent SV coupling but being independent of SV coupling or of UNC13L/RAB-3 in the rest of the AZ. We note that the larger diffusion range of VGCCs exploring the AZ was differentially impacted in the two single mutants, as it significantly decreased in the absence of RAB-3 (*rab-3*: 234 ± 30 nm *vs.* WT: 289 ± 29 nm), but increased in *unc-10(nu487)* mutants (*unc-10*: 322 ± 32 nm vs. WT: 289 ± 29 nm). These opposing effects offset each other in *rab-3;unc-10(nu487)* double mutants, where the diffusion range remained similar to WT (Fig. 3d). This suggests that the AZ area explored by VGCCs outside nanodomains is promoted by UNC-10/RAB-3-mediated SV docking while it is restricted by UNC-10/UNC-13L interactions. In *rab-3;unc-10(nu487)* double mutants, the neurotransmission and locomotion defects were significantly more severe than in both single mutants, suggesting that the interactions of UNC-10 with RAB-3 and UNC-13L are functionally complementary (Fig. 3a and 3b). Together, these findings show that the diffusion of nanodomain-associated VGCCs is driven by a tripartite UNC-10/RAB-3/UNC-13L SV coupling complex, which is essential for efficient synaptic release.

To test whether this SV coupling complex also participates in organizing VGCC nanodomains, we examined the distribution of VGCCs in *rab-3;unc-10(nu487)* double mutants by STED. We found that the distribution of VGCC nanodomains became more compact and elongated than in WT animals, with their NND and puncta shape (long/short axes ratio) being significantly altered (Fig. 3e-3g). The number of nanodomain per puncta was also significantly reduced, yet it was not impacted to the extent observed in the absence of UNC-10 (Fig. 3h and 3i). This indicates that SV coupling mediated by UNC-10 partially promotes the accumulations of VGCC nanodomains and shapes their spatial organization at AZs.

The mobility and AZ organization of VGCCs is governed by synaptic vesicle priming and SNARE complex assembly

As UNC-10/UNC-13L interactions couple VGCCs to readily releasable SVs[17], we next

investigated whether the channel mobility is regulated by SV priming. We examined the diffusion

of VGCCs in *unc-13(s69L-S-)* null mutants, where the absence of both UNC-13 isoforms abolishes

SV priming and synaptic release[49,51] (Fig. 4a; Fig. S6a). In *unc-13(s69L-S-)* animals, nanodomain-

associated VGCCs exhibited significantly reduced D_{slow} and increased confinement (*unc-13^{L-S-}*: 81

$\pm$ 21 nm *vs.* WT: 104 $\pm$ 23 nm), phenocopying changes observed in *rab-3;unc-10(nu487)* double

mutants (Fig. 4b; Fig. S6b). Because disrupting the interaction of UNC-13L with UNC-10 in *unc-*

10(nu487) animals did not influence the channel mobility, these results suggest that VGCC

diffusion in nanodomains is impacted by SV priming via both UNC-13L and UNC-13S rather than

by simple interaction between UNC-13L and UNC-10. To further test this, we studied the diffusion

of VGCCs in *unc-13(e51L-S+)* mutants, where priming levels are slightly restored by endogenous

expression of only the UNC-13S isoform, which lacks the UNC-10-binding domain, but contains

the core regions required for priming[51]. In *unc-13(e51L-S+)* animals, neurotransmission was

significantly improved compared to *unc-13(s69L-S-)* mutants (Fig. 4a; Fig. S6a), but it remained

defective compared to WT, consistent with UNC-13L being critical for efficient SV release[17,51].

When expressing the UNC-13S isoform, both D_{slow} and the confinement size of nanodomain-

associated VGCCs were partially restored, with D_{slow} remaining slower than in WT, albeit short of

statistical significance (p = 0.219, Fig. 4b; Fig. S6c and S6d). This indicates that moderately

improved SV priming levels by UNC13S expression is sufficient to promote the mobility of slow

VGCCs, but that UNC13L-mediated priming is required to restore WT-level diffusion in AZ

nanodomains. VGCCs outside nanodomains were impacted in similar ways. D_{fast} decreased in *unc-*

13(s69L-S-) mutants, albeit short of statistical significance (p = 0.178), and it was only partially

restored to WT-levels in *unc-13(e51L-S+)* animals (Fig. 4c). Compared to WT, the channel

diffusion range became significantly smaller when both UNC-13 isoforms were ablated (*unc-13^{L-}*

$^{S-}$: 240 $\pm$ 23 nm *vs.* WT: 289 $\pm$ 29 nm) but it was restored by the sole expression of UNC-13S (*unc-*

13^{L-S+}: 310 $\pm$ 28 nm *vs.* WT: 289 $\pm$ 29 nm; Fig. 4c). Together, these results indicate that UNC-13-

mediated SV priming modulates the mobility of VGCCs both within and outside nanodomains at

the AZ.

To investigate if the structural organization of VGCCs at AZs was also modulated by UNC-13-

mediated SV priming, we imaged VGCCs and UNC-10 by STED in the two *unc-13* mutants. For

both VGCCs and UNC-10, the number of nanodomains per AZ and the size of nanodomains were

significantly lower in *unc-13(s69L-S-)* mutants compared to WT (Fig. 4d-4f; Fig. S6f), with the

distribution of UNC-10 nanodomains being more compact, but that of VGCC nanodomains being

unchanged (NND, Fig. S6e). The mean AZ puncta size of VGCCs and UNC-10 were both

significantly reduced (AED, Fig. 4g), indicating that AZs were globally smaller when UNC-13

mediated SV priming was abolished, which led to a decrease in the diffusion range of fast VGCCs.

With the expression of UNC-13S in *unc-13(e51L-S+)* mutants, VGCC nanodomain numbers, their

size and the AZ puncta sizes were only partially restored compared to WT animals (Fig. 4d-4g;

Fig. S6f). This indicates that UNC-13S-mediated SV priming influences the spatial distribution of

VGCCs, but that UNC-13L-mediated priming is required to maintain a WT structural organization

of the channels at AZs, consistent with our diffusion measurements. For UNC-10, the sole

expression of UNC-13S fully restored the number of nanodomains per AZ and their NND to WT

levels (Fig. 4e; Fig. S6e). Yet, in the absence of UNC-13L, the nanodomain size of UNC-10 and

its AZ puncta size remained well below WT levels and non-significantly different from *unc-*

13(s69L-S-) mutants (Fig. 4f and 4g). These results indicate that the number and distribution of

UNC-10 nanodomains are also modulated by UNC-13S-mediated SV priming, while UNC13-L

mediated priming contributes to the size of individual UNC-10 nanodomains and that of the AZ.

Surprisingly, the apparent rescue of VGCC nanodomain confinement and AZ diffusion range to

WT levels in *unc-13(e51L-S+)* animals (Fig. 4c) did not fully correlate with the nanodomain and

AZ puncta sizes for UNC-10 in those mutants (Fig. 4g), suggesting that, in the absence of UNC-

13L, UNC-13S-mediated SV priming induces VGCCs to diffuse beyond spatial boundaries

defined by UNC-10.

We next investigated whether the mobility of VGCCs is modulated by SNARE complex assembly

during SV priming and whether SV coupling, at this molecular step, impacts the channels diffusion.

We utilized the transgenic expression of the botulinum neurotoxin serotype B light chain protease

(ScBoNTB), which blocks full SNARE complex assembly by cleaving the v-SNARE vesicular

SNB-1[52]. As previously reported, expressing ScBoNTB in GABAergic neurons silenced inhibitory

neurotransmissions[52] and 92% of transgenic animals showed "shrinker" phenotypes (Fig.4h). For

proper comparison, we first reanalyzed the mobility of VGCCs specifically at GABAergic

synapses in WT animals by selecting trajectories only located at the tip of GABAergic boutons

(Fig. S6g). GABAergic VGCCs also exhibited two diffusive behaviors, with respective fractions

and diffusion coefficients that were non-significantly different from those observed across all

buttons, although D_{fast} appeared slightly faster (Fig. 4i and 4j; Fig. S6h). Only the confinement

size of nanodomain-associated GABAergic VGCCs was smaller ($WT^{GABAergic}$: 90 ± 25 nm *vs.*

WT^{All}: 104 ± 23 nm) (Fig. 4i). We then tracked GABAergic VGCCs in ScBoNTB-expressing

animals. We found that both D_{slow} and D_{fast} significantly decreased, an indication that the mobility

of VGCCs inside and outside AZ nanodomains is indeed modulated by a molecular priming of

SVs that requires full SNARE assembly (Fig. 4i and 4j; Fig. S6i). The diffusion range of VGCCs

outside nanodomains was also significantly reduced ($ScBoNTB^{GABAergic}$: 207 ± 29 nm *vs.*

$WT^{GABAergic}$: 278 ± 42 nm; Fig. 4j). But the confinement size of nanodomain-associated VGCCs

was unchanged compared to GABAergic WT (Fig. 4i), suggesting that it is independent of SNARE

complex assembly. Altogether, these results demonstrate that, downstream of UNC-10 mediated

channel-vesicle coupling and as part of UNC-13 mediated priming, the complete assembly of

SNARE complexes that clamp SVs is a critical step that modulates VGCC diffusions at AZs.

The two mobilities of VGCCs are differentially modulated by UNC-13 isoforms and synaptic

vesicle priming levels

We then examined if the mobility of VGCCs would positively scale with SV priming levels by

tuning the balance between UNC-13 isoforms and the t-SNAREs inhibitor TOM-1/Tomosyn. In

C. elegans, removal of TOM-1 markedly upregulates the pool of primed SVs throughout synaptic

terminals[53,54], and also partially rescues synaptic release defects in *unc-13* mutants[51]. We therefore

compared the diffusion of VGCCs in *tom-1;unc-13*$^{L-S-}$ versus *tom-1;unc-13*$^{L-S+}$ double mutants,

where SV priming levels are progressively restored, and in *tom-1(nu468)* single mutants where

priming levels are largely augmented[51,54]. As previously reported, both double mutants exhibited

significantly improved locomotion and neurotransmissions compared to the corresponding *unc-13*

single mutants, with SV release rescue being stronger in the presence of UNC-13S (Fig. 5a; Fig.

S7a). We found that D_{slow} for nanodomain-associated VGCCs was partially restored in *tom-1;unc-*

13^{L-S-} animals, while their confinement size fully recovered to WT-level (*tom-1;unc-13*$^{L-S-}$: $111 \pm$

22 nm *vs.* WT: 104 ± 23 nm, Fig.5b), confirming that slow-diffusing VGCCs are modulated by

SV priming. However, despite enhanced SV priming levels, the mobility of VGCCs inside

nanodomains was not further improved by the expression of UNC-13S in *tom-1;unc-13*$^{L-S+}$ double

mutants, as it remained similar to *tom-1;unc-13*$^{L-S-}$ animals and to *unc-13(e51L-S+)* single mutants

(Fig. 5b; Fig. S7b-S7f). These results suggest that UNC-13S and TOM-1 have limited effect on

the mobility of nanodomain-associated VGCCs, which are primarily regulated by UNC-13L.

Indeed, the mobility of VGCCs in nanodomains did not further increase with augmented SV

priming levels in *tom-1(nu468)* single mutants (Fig. 5a and 5b; Fig. S7e).

On the contrary, for VGCCs outside nanodomains, both D_{fast} and their diffusion range significantly

increased in *tom-1(nu468)* animals (*tom-1*: 331 ± 33 nm *vs.* WT: 289 ± 29 nm; Fig. 5c). This

enhanced mobility was suppressed when UNC-13 was ablated in *tom-1;unc-13^{L-S-}* double mutants,

where D_{fast} was similar to that in *unc-13(s69L-S-)* single mutants (Fig. 5c). However, the increase

in D_{fast} was largely preserved by UNC-13S expression in *tom-1;unc-13^{L-S+}* double mutants, who

displayed significantly faster VGCC diffusion than in *tom-1;unc-13^{L-S-}* animals (Fig. 5c; Fig. S7d-

S7f). This indicates that the faster mobility of VGCCs outside nanodomains in *tom-1(nu468)*

mutants is primarily UNC-13S-dependent, conforming with a previously reported upregulation of

UNC-13S abundance in these animals[54]. Together, these findings show that the mobility of VGCCs

is predominantly governed by UNC-13L-mediated SV priming inside nanodomains, and that it

scales with a pool of TOM-1 regulated and UNC-13S-primed SVs across the rest of the AZ.

Interestingly, the diffusion range of VGCCs outside nanodomains was dramatically expanded in

tom-1;unc-13^{L-S-} double mutants (*tom-1;unc-13^{L-S-}*: 632 ± 112 nm *vs.* WT: 289 ± 29 nm; Fig. 5c),

in stark contrast with the reduction observed in *unc-13(s69L-S-)* single mutants. As the number of

membrane-contacting SVs located at the AZ periphery (400-600 nm away from the DPs)

substantially increases in *tom-1;unc-13^{L-S-}* animals[53], our observation suggests that VGCCs can

diffuse far across the synaptic area where UNC-13-independent SV priming is upregulated.

Concurrently, in the same *tom-1;unc-13^{L-S-}* animals, the size of UNC-10 puncta was significantly

reduced (Fig. 5e), indicating that, in the absence of UNC-13, the diffusion range of fast VGCCs is

set by the location of primed SVs rather than by the distribution of UNC-10 at AZ puncta. When

SV priming was controlled by UNC-13S in *tom-1;unc-13 $^{L-S+}$* double mutants, the diffusion range

of fast VGCCs was restored to WT levels (*tom-1;unc-13^{L-S+}*: 307 ± 37 nm *vs.* WT: 289 ± 29 nm),

together with the median puncta size of UNC-10 (Fig. 5c-5e). And when UNC-13S-mediated SV

priming was upregulated in *tom-1(nu468)* animals, this same diffusion range significantly

increased compared to WT (*tom-1*: 331 ± 33 nm *vs.* WT: 289 ± 29 nm; Fig. 5c), again exceeding

the median UNC-10 puncta size (Fig. 5c-5e). Together, these observations indicate that outside

nanodomains, the mobility of VGCCs is effectively governed by UNC-13S-mediated SV priming,

and that the channel diffusion range can expand when UNC-13-independent and spatially

imprecise SV priming is upregulated.

We also examined if increasing priming levels by TOM-1 removal modulates the AZ organization

of VGCC and UNC-10 nanodomains. We found that the defective nanodomain numbers and

puncta size of VGCCs we had observed when ablating the two UNC-13 isoforms were partially

restored by simultaneous removal of TOM-1 in *tom-1;unc-13^{L-S-}* double mutants, although they

both remained significantly lower than WT (Fig. 5d-5f). The median number and NND of UNC-

10 nanodomains were back to WT-level, yet UNC-10 puncta sizes did not recover (Fig. 5d-5f; Fig.

S7g-S7i). These results are similar to those found in *unc-13(e51L-S+)* mutants, where SV priming

levels are improved and comparable to those of *tom-1;unc-13^{L-S-}* animals[55]. Further improvements

of SV priming in *tom-1;unc-13 $^{L-S+}$* double mutants fully restored UNC-10 puncta sizes to WT

levels, but the organization of VGCC nanodomains remained partially defective (Fig. 5d-f),

confirming that UNC-13L is important for organizing VGCCs at the AZ. In *tom-1(nu468)* animals,

where both UNC-13 isoforms are present and UNC-13S-mediated priming is predominantly

upregulated, the spatial organizations of both VGCC and UNC-10 nanodomains were

indistinguishable from WT (Fig. 5d-5f). This suggests that VGCC and UNC-10 organizations scale

with SV priming levels up to a limit set by UNC-13L-activity. Collectively, these findings

demonstrate that UNC-13 mediated SV priming modulates the organization of VGCC and UNC-

10 nanodomains in a parallel and progressive manner. Notably the increase in UNC-10

nanodomain numbers is followed by an expansion of AZ puncta sizes, and the organization of

VGCC nanodomains exhibits greater sensitivity to priming levels than that of UNC-10.

Primed synaptic vesicles can modulate the mobilities and multi-nanodomain organization of

VGCCs downstream of UNC-10

As the diffusive behaviors of VGCCs are promoted by UNC-10 and SV priming, we tested whether

they both modulate VGCC mobility in the same molecular pathway by examining the channels

diffusion in *tom-1;unc-10* double mutants. Interestingly, the defective D_{slow} and confinement sizes

of nanodomain-associated VGCCs we observed in *unc-10(md1117)* single mutants were fully

restored to WT-level in *tom-1;unc-10* double mutants (Fig. 6a; Fig. S8a). Likewise, the diffusion

coefficient of VGCCs outside nanodomains was also significantly increased, with D_{fast} being faster

than in WT, albeit short of significance (p = 0.193, Fig. 6b), and being slower than the augmented

D_{fast} observed in *tom-1(nu468) single* mutants. Thus, upregulating SV priming levels can partially

bypass the role of UNC-10 in modulating the mobility of VGCCs. The diffusion range of fast

VGCCs in *tom-1;unc-10* double mutants showed significant increase compared to WT (*unc-*

10;tom-1: 336 ± 27 nm *vs.* WT: 289 ± 29 nm; Fig. 6b; Fig. S8a), phenocopying changes observed

in *tom-1(nu468)* animals and confirming that the diffusion range is set by primed SVs

independently of UNC-10. Together, it demonstrates that UNC-10 functions to couple mobile

VGCCs to primed SVs, which can directly modulate the channel mobilities downstream of UNC-

10. TOM-1 removal also largely rescued the impaired neurotransmissions in *unc-10(md1117)*

background (Fig. 6c) as previously reported[55]. Since UNC-10 is not required for molecular

priming[17,56], the rescue of synaptic releases through TOM-1 ablation in double mutants likely

stems from the upregulated SV priming levels and the restored VGCC mobility, which might

compensate for reduced SV docking[43].

Finally, we examined whether increasing priming levels can rescue the defective organizations of

VGCC nanodomains in the absence of UNC-10. In *tom-1;unc-10* animals, the number of VGCC

nanodomains per puncta was slightly but significantly increased compared to *unc-10(md1117)*

single mutants, yet it remained largely reduced compared to WT (Fig. 6d and 6e). In contrast, the

NND between nanodomains was non-significantly different from that in *unc-10(md1117)* animals,

showing a more dispersed organization compared to WT (Fig. 6f). These results demonstrate that

the multi-nanodomain organization of VGCCs at AZ requires UNC-10 and indicate that SV

priming levels regulate the spatial distribution of VGCC nanodomains through UNC-10.

Discussion

Using single channel tracking and super-resolution imaging in *C. elegans*, we dissected how

mobile VGCCs are spatiotemporally aligned with synaptic vesicles for precise neurotransmission.

We discovered that individual N-type VGCCs adopt distinct mobilities determined by their spatial

distribution and the molecular machinery coupling them to readily releasable SVs. VGCCs

tethered to primed SVs via a tripartite UNC-10/RAB-3/UNC-13L complex diffuse slowly in

multiple sub-AZ nanodomains that align with UNC-10/RIM scaffolds, whereas VGCCs coupled

to UNC-13S-primed SVs move significantly faster across the larger AZ puncta. The spatially

correlated multi-nanodomain arrangement of VGCCs and UNC-10 closely resembles Cav2 and

RIM1/2 nanoclusters observed at rodent hippocampal synapses[50,57], consistent with nematode

synapses exhibiting a multi-release-site organization similar to mammals[58]. Heterogeneity in

nanodomain number and size across synapses further points to a modular intra-AZ organization

that may enable local tuning of synaptic strength. Together, this nanoscale organization and the

distinct diffusion regimes of VGCCs likely give rise to diverse channel-vesicle coupling modes

that shape SV release probability.

In this context, UNC-10 emerges as a key regulator of VGCC mobility and nanodomain

organization. We found that UNC-10 promotes VGCC diffusion by coupling channels to primed

SVs, and that priming levels modulate both channel dynamics and multi-nanodomain organization.

As an SV docking factor and a direct modulator of VGCCs, UNC-10/RIM drives channel

accumulation at AZs[7,9,45], stabilizing them and securing their coupling distances to vesicles[8,18,59].

Consistent with these functions, UNC-10 facilitates the accumulation and tight organization of

multiple VGCC nanodomains at AZs. However, VGCCs can form nanodomains independently of

UNC-10 and, counterintuitively, UNC-10 enhances channel mobility. Thus, rather than simply

stabilizing VGCCs, UNC-10 modulates channel dynamics and shapes their nanotopography. The

influence of UNC-10 on VGCC diffusion appears conserved across species, as $Ca_V2.1$ isoforms

harboring or lacking RIM1/2 binding motifs exhibit different mobilities at rodent hippocampal

synapses[23], and Cac/Ca_V2 displays similarly slower diffusion in the absence of RIM at *Drosophila*

NMJs[25].

Importantly, at *C. elegans* NMJs, VGCCs within sub-AZ nanodomains and across the broader AZ

exhibit two different mobility regimes that both arise from coupling to primed SVs but are

governed by distinct molecular mechanisms. To dissect these mechanisms, we altered vesicle

priming through UNC-13 isoforms, the t-SNARE inhibitor TOM-1, and SNARE complex

blockade, and found that channel dynamics depend on UNC-13-mediated vesicle priming

following full SNARE assembly and SV-plasma membrane bridging. Within nanodomains,

VGCC dynamics are primarily driven by SV priming mediated through UNC-10/RAB-3/UNC-

13L complexes, with additional contributions from the UNC-13S isoform. Outside nanodomains,

however, this tripartite complex does not regulate VGCC mobility but instead influences the

channel distribution across AZ. In this region, VGCC dynamics scale with pools of UNC-13S-

primed or UNC-13-independent SVs, which are physiologically inhibited by TOM-1. Accordingly,

blocking SNARE assembly with ScBoNTB significantly reduces both VGCC diffusion and

restricts their AZ distribution, whereas removal of TOM-1 partially restores defective channel

dynamics in the absence of UNC-13 or even UNC-10. Thus, UNC-10-mediated SV coupling

drives VGCC diffusion following SNARE assembly, while primed SVs can further modulate

VGCC behavior beyond UNC-10-dependent coupling. These findings establish a causal link

between SV priming levels and channel dynamics and may mechanistically explain why conditions

that impair priming and SNARE assembly, such as intracellular Ca^{2+} chelation or silencing of

synaptic activity, are associated with reduced VGCC mobility but increased Syntaxin-1A/t-

SNARE diffusion[23,24,60].

Accordingly, the AZ distribution of fast VGCCs is primarily dictated by the location of primed

SVs rather than the multi-nanodomain organization of UNC-10. This is evident when simultaneous

removal of UNC-13 and TOM-1 leads to an expansion of VGCC diffusion range toward distal AZ

regions, where UNC-13-independent SV priming is upregulated[53]. The modest expansion

observed after disrupting UNC-10/UNC-13L interactions likely reflects a broadening in primed

SV distribution, with UNC-13L becoming less anchored at the AZ[55]. Conversely, removing RAB-

3 significantly reduces the diffusion range of VGCCs, as impaired SV docking limits their spatial

distribution across AZ. These observations indicate that VGCC topography is dynamically

instructed by the spatial distribution of primed SVs rather than solely imposed by AZ scaffolds.

Consistent with this relationship, SV priming levels progressively regulate the number and size of

VGCC and UNC-10 nanodomains, with the organization of VGCCs being more sensitive to

priming levels. These results complement previous findings that Ca_V2 clustering and vesicle

priming represent two distinct sub-assemblies tethered together by RIM[61]. Our data further

indicates that this tethering is itself regulated in a priming-level-dependent manner, with primed

vesicles reciprocally regulating UNC-10/RIM and directly shaping VGCC diffusion.

Collectively, our findings support a model in which primed SVs actively drive VGCC diffusion

through physical coupling, coordinating the movement of membrane-embedded channels with

that of adjacent primed SVs[62–66]. Such coordination likely secures the coupling distance between

the VGCC calcium donors and the SV-tethered synaptotagmin calcium sensors[1,67,68] thereby

ensuring precise neurotransmission within the highly dynamic environment of the AZ.

Furthermore, coupling distinct VGCC mobility to different pools of primed SVs suggests a level

of synaptic modularity that supports diverse release properties.

Placing this model in a broader context, studies at *Drosophila* and mammalian synapses[69–73],

together with functional analyses in *C. elegans* indicate that distinct channel-vesicle coupling

distances underlie different release properties, with fast release requiring UNC-13L and slow release depending on both UNC-13L and UNC-13S[55,74]. Coupling distance has conventionally been defined as the distance between docked SVs and the DP at the center of the AZ, where DP markers appear as homogeneous puncta in diffraction-limited images[27,45,56]. Our studies reveal a heterogeneous, multi-nanodomain topography of UNC-10 and VGCCs that closely resembles the architecture of DP[42]. This spatial organization, together with differences in VGCC mobility and their coupling to distinct UNC-13 isoforms, supports a multidimensional framework for generating diverse SV release properties. Within this framework, UNC-13L-primed SVs preferentially dock in nanodomains and tether slow-diffusing VGCCs, whose restricted mobility promotes concentrated calcium influx[24] for fast release. In contrast, UNC-13S-primed SVs dock outside nanodomains or at distal-AZ regions, where they couple to fast-diffusing VGCCs, whose higher mobility yields more diffuse calcium influx for slow release. This framework further implies that differences in channel mobility contribute to defining microdomain- and nanodomain-coupling modes[20].

Whether and how primed SVs tether fast-diffusing VGCCs outside nanodomains remains unclear. Because fast-diffusing VGCCs are not coupled to the tripartite UNC-10/RAB-3/UNC-13L complex, the assembled SNARE complex itself may represent a candidate. While mammalian Cav2 channels contain a "synprint" motif that directly binds SNARE complexes[4,75], UNC-2/Cav2 in *C. elegans* lacks such a sequence[76], and identifying the mechanisms that mediate coupling to fast-diffusing VGCCs will require further investigation. Despite this evolutionary divergence, the principle of vesicle-channel coupling may be more fundamental than the specific protein-protein interactions that mediate it[77]. Another open question is what determines the transition between slow and fast diffusion modes for VGCCs. It will be important to test whether coupled SV-VGCC units move in and out of VGCC nanodomains together, potentially supporting transitions between tightly primed (TS) and loosely primed (LS) states[78]. Addressing these questions will provide further insight into how synapses exploit VGCC dynamics to achieve both reliability and plasticity.

In conclusion, this study identifies a mechanism by which presynaptic VGCC diffusion is actively regulated by the release machinery and the priming status of SVs. The coupling of distinct channel mobilities to different pools of primed vesicles reveals a previously unrecognized layer of AZ

530 organization that fine-tunes neurotransmitter release. Ultimately, this work establishes a
531 mechanistic framework to investigate the role of VGCC dynamics in synaptic function and disease.
532

Acknowledgement
We thank Dr. Dion Dickman and Dr. Kaikai He for providing important resource support and
advice for STED imaging and analysis. We thank Dr. Hong Zhan for advice on CALM imaging
and genome editing of *C. elegans*, and Dr. Carolyn Phillips for resource support on genetics. We
also thank Dr. Jean-Louis Bessereau for kindly sharing the unc-25p::snb-1::gfp strain, Dr. Peri
Kurshan for her kind gifts of GFP-UNC-2 related strains. We additionally thank Dr. Joshua Kaplan
for sharing tom-1(nu468) related strains, Dr. Alexander Gottschalk for kindly providing the unc-
47p::eGFP::ScBoNTB strain and Dr. Zhitao Hu for sharing the unc-10 (nu487 K77/79E) strain.
This work was funded by NIH grant NS114911 from the National Institute of Neurological
Disorders and Stroke (NINDS). YZ was partially supported by the NIGMS training grant
T32GM118289-04 Chemistry Biology Interface Training Grant at the University of Southern
California. Some strains were obtained from the Caenorhabditis Genetics Center (CGC) that is
funded by the NIH Office of Research Infrastructure Programs (P40 OD010440).

Methods
C. elegans **culture and maintenance**
All *Caenorhabditis elegans* nematode strains were raised on nematode growth media (NGM)
plates at 20°C and fed on OP50 *Escherichia coli* as previously described[33]. All *C. elegans* used for
experiments were adult hermaphrodites. Worms were handled using a Nikon dissecting
microscope (SM2745). A full list of strains is available in Supplementary Table 1.
554

Molecular biology
For genotyping, worms were lysed in worm lysis buffer as previously described[30]. Worm lysates
were amplified using either 2xTaq PCR Premix (Bioland) or Phusion High-fidelity DNA
Polymerase (NEB). For CRISPR-Cas9 genome editing, direct microinjection of *in vitro*–
assembled CRISPR-Cas9 and single chimeric guide RNA (sgRNA) ribonucleoprotein complexes
into the gonads of hermaphrodite worms were performed as previously described[30,79]. Cas9

endonuclease was expressed *in vitro* using DE3 GOLD (Agilent) cells transformed with the

nm2973 plasmid[80] and was prepared to 10 μg/μl. Single-stranded DNA oligonucleotides (ssODNs)

were ordered from Integrated DNA Technology (IDT). DNA templates for sgRNA combining

tracrRNA and crRNA were amplified from two complementary ssODNs. sgRNA were transcribed

in vitro using a MEGA shortscript T7 kit (Thermo Fisher Scientific) and prepared at 2 μg/μl. Linear

homology-dependent repair templates (HDRT) containing 35-nucleotide homology arms were

amplified using Phusion HF PCR reactions and were prepared at 1 μg/μl. The *dpy-10* co-CRISPR

approach[81], which generates a roller phenotype was used for screening progeny with high edit

frequency. A total 15 μl mix of ribonucleoprotein complexes containing Cas9 endonuclease,

sgRNA and HDRT for the targeted genes and *dpy-10* were incubated at 37 °C for 10 min and spun

at 13,000 rpm for 2 min before gonad microinjections. To generate the sCherry::unc-36 transgenic

strain, linear HDRT were amplified using plasmids pcDNA3.1-sfCherry2(1-10)[82] (Addgene) as

PCR templates. For generating 5xMYC::unc-10 and HA::unc-2 strains or mutating the marker

gene *dpy-10*, ssODNs (IDT) were used as the HDRT. A silent mutation for the fifth Serine of UNC-

10 was introduced in the HDRT to prevent Cas9 recognition. All generated worm lines were

outcrossed at least three times to remove potential off-target mutations. A full list of primers,

sgRNAs and HDRT is available in Supplementary Table 2.

Microinjection

Microinjection was performed as previously described[30]. For genome editing, standard gonad

microinjection was done on young adult worms. For CALM imaging, synthetic complementary

peptide sfCherry2(11)-cysteine (NH$_2$-C-acp-linker GSGGGSTSYTIVEQYERAEARHSTGG-

COOH, LifeTein, LLC.75% purity) was first dissolved in DMSO to as a 30mM stock, which was

then diluted to 750μM with PBS buffer right before microinjections. For immunostaining of

extracellular sCherry-UNC-36, rabbit anti-mCherry (Abcam Cat# ab167453, RRID: AB_2571870)

was freshly diluted at 1:400 with injection buffer[83] (20 mM K$_2$HPO$_4$, 3 mM potassium citrate, 2%

PEG3000, pH 7.4). Either peptide or antibody solutions were spun at 15,000 rpm for 2 min before

microinjection into the pseudocoelomic cavity at the head region of staged 1-day adult worms.

Animals were recovered in recover buffer (5 mM HEPES pH 7.2, 3 mM CaCl$_2$, 3 mM MgCl$_2$, 66

mM NaCl, 2.4 mM KCl, 4% w/v glucose) drop on a NGM plate at room temperature for 16-20 h

before imaging or freeze cracking. Microinjection was done with an inverted IX50 Olympus

microscope, equipped with an Eppendorf FemtoJet micro-injector connected to FemtotipII

microinjection capillaries held on a Leitz manual micromanipulator.

Behavioral assays

Aldicarb acute paralysis assays and body bending thrashing assays were performed on staged 1-

day adult worms as previously described[84,85]. Numbers of body bending per minute were scored

in the M9 buffer added to 35 mm sterile NGM agar plates. Worms were first accommodated in

M9 buffer for 2 min and the number of sinusoidal curves made within a 2 min period were counted.

At least 15 worms were scored for each strain. For aldicarb assays, 6-well NGM plates containing

1 mM aldicarb were made and seeded with OP50 bacteria the day before the assays. For each strain,

30 animals were placed on plates and monitored every 30 min for 4 h. Worms were considered

paralyzed if their whole body including nose and tail regions show no movement after a gentle tap

on the head using a worm pick. For comparison to mutant animals, wildtype control worms were

cultured and scored simultaneously on the same batch of NGM or aldicarb plates. At least three

trials, with independent worm cultures, were performed for each assay. Data collections were

performed blinded to the animal strains. Data were analyzed using either GraphPad Prism software

version 8.0.2(263) or OriginPro software version 10.1.0.178.

Image acquisition for CALM

CALM imaging was performed as previously described[30]. 16-20h post peptide microinjection,

worms were anesthetized for 6-10 min in M9 buffer containing 30 mg/mL 2,3-butanedione

monoxime (BDM, Sigma) in a 35 mm plate. Anesthetized worms were then mounted in 8 μL BDM

solution on freshly prepared 2% agarose pads with high-precision Deckglaser microscope

coverslips (Marienfeld, 0117650). Worms were positioned with one of the nerve cords facing

upright by gentling sliding the coverslip without pressing down. The worm position was secured

by sealing the four quarters of the coverslip to the slide with minimal nail polish and stabilized for

10min before imaging. Anesthetized worms were imaged within 40 min, after which locomotion

recovered. Simultaneous dual-color imaging of Punc-25::SNB-1::GFP and sCherry-UNC-36 was

performed on an inverted Nikon Eclipse Ti-E microscope equipped with HiLO optics, a 100x 1.49

NA objective (Nikon), 488 and 561 nm fiber-coupled excitation lasers (Agilent), a multiband pass

ZET405/488/561/647 excitation filter (Chroma), a quad-band ZT405/488/561/647 dichroic mirror

(Chroma), two EMCCD cameras (Andor iXon), a dual-camera imaging adapter (Andor), an
emission-splitting FF560FDi01 dichroic mirror (Semrock), and emission filters at 525/50 nm
(Semrock) for GFP and 600/50 nm (Chroma) for sCherry imaging. Movies were acquired at a
frame rate of 80 ms/frame, with EMCCD gains of 300, and 1 x1 binning, leading to image pixel
sizes of 107 nm on each camera. Dual-channel alignment was done by imaging 40 nm diameter
TransFluoSphere beads (488/685, Invitrogen) at 1:10000 dilution. CALM imaging of individual
sCherry-UNC-36 was performed by continuous HiLO illumination of nerve cords marked by
either GFP-UNC-2 or Punc-25::SNB-1::GFP for 1-2 min per field of view from the tail to the head
regions of each animal. GFP channel was excited with minimal laser intensity to reduce
background. Single molecule detections of sCherry-UNC-36 showed typical blinking events,
single-step photobleaching, and sudden fluorescence appearance, as previously described[29,30].
Each strain condition was imaged with at least three independent cultures of worms. The numbers
of animals imaged for each strain are indicated in figure legends.
636

Tracking and diffusion analyses
Single molecule tracking was performed as previously described[29,30] using the Matlab software
SLIMfast, which is based on multiple-target tracing algorithms[86]. For sub-diffraction limit
localizations and trajectory assembly of sCherry-UNC-36, the following parameters were used for
all conditions: 107 nm/pixel, 80 ms/frame, 10^{-6} localization error, fixed PSF radius of 1.02 pixel
determined by theoretical calculation for a 1.49 NA and 615 nm maximum emission wavelength,
0.5-3 pixels search limit and maximum gap of 3 steps for linking positions. Only trajectories with
a minimal number of 3 steps were kept for diffusion analyses. Diffusion analyses of neuronal
VGCCs were performed on sCherry-UNC-36 trajectories located along the nerve cords or, in the
case of only GABAergic VGCCs, that overlapped with synaptic boutons marked by Punc-
25::SNB-1::GFP. For each strain condition, trajectories from multiple worms were pooled together
and analyzed using the PDSD method[30] to differentiate diffusive behaviors, and determine the
numbers, fractions and mean square displacements of each diffusion regime. Mean square
displacement curves for the first 10 steps were plotted for each behavior and fitted with a 2D
circularly confined diffusion model[29] to compute the diffusion coefficients and radius of
confinement. OriginPro software version 10.1.0.178 was used for data analysis.

Immunofluorescence staining for confocal and STED microscopy

Worms were harvested at stage 2-day adult or 18h post antibody microinjection for immunostaining of extracellular sCherry-UNC-36. Worms were mounted on slides with ice cold Milli-Q water and frozen on dry ice for 1 hour before freeze-cracking as previously described[87]. Fixation was done overnight at 4 °C in microcentrifuge tubes with phosphate-buffered saline (PBS) containing 1% paraformaldehyde (PFA). For HA-UNC-2 immunostaining, fixation was reduced to 20 min at room temperature. Fixation was followed by three 20 min washes with PBS at 4 °C and blocking in blocking solution (PBS, 5% BSA, 0.5% Triton X-100, 1mM EDTA pH 8, 0.05% Sodium azide) for 1 h at room temperature or overnight at 4 °C. Incubations with primary and secondary antibodies were each done overnight at 4 °C in antibody solution (PBS, 2.5% BSA, 0.5% Triton X-100, 1mM EDTA pH 8, 0.05% Sodium azide). Three 20 min washes with PBST (0.5% Tween) were performed at 4 °C between each incubation and one 5 min final wash with PBS was done before mounting. Samples were mounted using high-precision Deckglaser coverslips (Marienfeld, 0117650) and ProLong Glass Antifade Mountant (Invitrogen). Mounted samples were cured for 24 h at room temperature before imaging. The primary antibodies used were rabbit anti-mCherry (Abcam Cat# ab167453, RRID: AB_2571870; 1:400), mouse anti-MYC (Cell Signaling Technology Cat# 2276, RRID: AB_331783; 1:1000), rat anti-HA (Roche Cat# 11867423001, RRID: AB_390918; 1: 100) and chicken anti-GFP (Aves Labs Cat# GFP-1010, RRID: AB_2307313; 1:200). The secondary antibodies used were goat anti-rabbit STAR RED (Abberior Cat# STRED-1002-500UG, RRID: AB_2833015; 1:400), goat anti-mouse STAR RED (Abberior Cat# STRED-1001-500UG, RRID: AB_3068620; 1:600), goat anti-mouse Alexa Fluor 594 (Jackson Immuno Research Labs Cat# 115-585-146, RRID: AB_2338881; 1:500), goat anti-rat Alexa Fluor 594 (Thermo Fisher Scientific Cat# A-11007, RRID:AB_10561522; 1:200) and goat anti-chicken IgY Alexa Fluor 594 (Thermo Fisher Scientific Cat# A-11042, RRID:AB_2534099; 1:500).

Image acquisition and analyses for confocal and STED microscopy

Both confocal and STED imaging were performed on a Nikon Eclipse FN1 upright microscope equipped with an Abberior STEDYCON system, four excitation lasers (640, 561, 488, and 405 nm), a pulsed STED laser of 775 nm, and three avalanche photodiode detectors operating in single photon counting mode. 2D STED images were acquired using a 100× Nikon Plan APO 1.45 NA

oil immersion objective, fixed pixel size of 15 nm, a dwell time set at 10 µs, 5x line accumulation

in photon counting mode, and time gating set at 1 ns with a 6 ns width for all channels. Dual-color

STED images were acquired using 775 nm depletion beam with the same depletion power for both

STAR RED and Alexa Fluor 594 fluorophores. Confocal scans with 1x line accumulation for all

channels were done before STED scans, using 5x line accumulation for the same channels. To

overlay STED and confocal images of Punc-25::SNB-1::GFP, 5x line accumulation was also used.

Acquisition settings for each channel were identical in all conditions. Imaging field of views with

sizes of 10 µm or 20 µm x 3µm were used. 5-15 different animals were imaged for each condition.

Only dorsal nerve cords were imaged as cell bodies often interfered with imaging at ventral nerve

cords.

For figures display, raw STED images were deconvolved with the Huygens Essential version 24.10

software (Scientific Volume Imaging, The Netherlands, http://svi.nl) using the standard mode and

theoretical point spread functions of the STED microscope. For quantitative analysis of UNC-36

and UNC-10, raw images were processed with the Savitzky Golay denoise filter[88] of the Space

Noise Reduce plugin in ImageJ/Fiji, setting the parameter order at 7 and the width at 5.

Segmentations and measurements of puncta and nanodomains were done with the Particle Sizer

plugin[89] in ImageJ/Fiji, using the single particle mode for puncta and the watershed mode for

irregular structures mode for nanodomain. AZs with planar orientation where both UNC-36 and

UNC-10 signals could be clearly resolved were manually selected for measurements.

Segmentations parameters were optimized using both WT animals and mutants with severe

phenotypes and were kept identical for each protein and all conditions. The number of

nanodomains and their nearest neighborhood distance (NND) were quantified by treating each

nanodomain as a theoretical point spread function to localize its center position. The images were

first processed with the Rolling Ball Background Subtraction algorithm[90] built in ImageJ/Fiji,

using with a 10-pixel rolling ball radius, and center localizations and NND measurements were

done in the Picasso software[91]. The same settings were used for all conditions. All measurements

were done on 2D STED images taken from staged 2-day adult worms. Data analysis and statistics

were performed using OriginPro software version 10.1.0.178.

References

1. Emperador-Melero, J. & Kaeser, P. S. Assembly of the presynaptic active zone. *Curr. Opin. Neurobiol.* **63**, 95–103 (2020).

2. Katz, B. Quantal Mechanism of Neural Transmitter Release. *Science* **173**, 123–126 (1971).

3. Südhof, T. C. Neurotransmitter Release: The Last Millisecond in the Life of a Synaptic Vesicle. *Neuron* **80**, 675–690 (2013).

4. Dolphin, A. C. & Lee, A. Presynaptic calcium channels: specialized control of synaptic neurotransmitter release. *Nat. Rev. Neurosci.* **21**, 213–229 (2020).

5. Nanou, E. & Catterall, W. A. Calcium channels, synaptic plasticity, and neuropsychiatric disease. *Neuron* **98**, 466–481 (2018).

6. Jánosi, B. *et al.* RIM and RIM-Binding Protein Localize Synaptic CaV2 Channels to Differentially Regulate Transmission in Neuronal Circuits. *J. Neurosci.* **44**, e0535222024 (2024).

7. Kaeser, P. S. *et al.* RIM Proteins Tether Ca2+ Channels to Presynaptic Active Zones via a Direct PDZ-Domain Interaction. *Cell* **144**, 282–295 (2011).

8. Koushika, S. P. *et al.* A post-docking role for active zone protein Rim. *Nat. Neurosci.* **4**, 997–1005 (2001).

9. Kushibiki, Y., Suzuki, T., Jin, Y. & Taru, H. RIMB-1/RIM-Binding Protein and UNC-10/RIM Redundantly Regulate Presynaptic Localization of the Voltage-Gated Calcium Channel in *Caenorhabditis elegans*. *J. Neurosci.* **39**, 8617–8631 (2019).

10. Betz, A. *et al.* Functional Interaction of the Active Zone Proteins Munc13-1 and RIM1 in Synaptic Vesicle Priming. *Neuron* **30**, 183–196 (2001).

11. Deng, L., Kaeser, P. S., Xu, W. & Südhof, T. C. RIM Proteins Activate Vesicle Priming by Reversing Autoinhibitory Homodimerization of Munc13. *Neuron* **69**, 317–331 (2011).

12. Dulubova, I. *et al.* A Munc13/RIM/Rab3 tripartite complex: from priming to plasticity? *EMBO J.* **24**, 2839–2850 (2005).

13. Rizo, J. Molecular Mechanisms Underlying Neurotransmitter Release. *Annu. Rev. Biophys.* **51**, 377–408 (2022).

14. Jahn, R. & Scheller, R. H. SNAREs — engines for membrane fusion. *Nat. Rev. Mol. Cell Biol.* **7**, 631–643 (2006).

15. Richmond, J. E., Weimer, R. M. & Jorgensen, E. M. An open form of syntaxin bypasses the requirement for UNC-13 in vesicle priming. *Nature* **412**, 338–341 (2001).

16. Südhof, T. C. & Rothman, J. E. Membrane Fusion: Grappling with SNARE and SM Proteins. *Science* **323**, 474–477 (2009).

17. Liu, H. *et al.* Heterodimerization of UNC-13/RIM regulates synaptic vesicle release probability but not priming in C. elegans. *eLife* **8**, e40585 (2019).

18. Dittman, J. S. & Ryan, T. A. The control of release probability at nerve terminals. *Nat. Rev. Neurosci.* **20**, 177–186 (2019).

19. Nakamura, Y. *et al.* Nanoscale Distribution of Presynaptic Ca2+ Channels and Its Impact on
Vesicular Release during Development. *Neuron* **85**, 145–158 (2015).

20. Eggermann, E., Bucurenciu, I., Goswami, S. P. & Jonas, P. Nanodomain coupling between
Ca2+ channels and sensors of exocytosis at fast mammalian synapses. *Nat. Rev. Neurosci.* **13**,
7–21 (2012).

21. Guerrier, C. & Holcman, D. The First 100 nm Inside the Pre-synaptic Terminal Where Calcium
Diffusion Triggers Vesicular Release. *Front. Synaptic Neurosci.* **10**, 23 (2018).

22. Mercer, A. J., Chen, M. & Thoreson, W. B. Lateral mobility of presynaptic L-type calcium
channels at photoreceptor ribbon synapses. *J. Neurosci.* **31**, 4397–4406 (2011).

23. Heck, J. *et al.* Transient Confinement of CaV2.1 Ca2+-Channel Splice Variants Shapes
Synaptic Short-Term Plasticity. *Neuron* **103**, 66-79.e12 (2019).

24. Schneider, R. *et al.* Mobility of calcium channels in the presynaptic membrane. *Neuron* **86**,
672–679 (2015).

25. Ghelani, T. *et al.* Interactive nanocluster compaction of the ELKS scaffold and Cacophony
Ca^{2+} channels drives sustained active zone potentiation. *Sci. Adv.* **9**, eade7804 (2023).

26. Tong, X.-J. *et al.* Retrograde Synaptic Inhibition Is Mediated by α-Neurexin Binding to the
α2δ Subunits of N-Type Calcium Channels. *Neuron* **95**, 326-340.e5 (2017).

27. Oh, K. H., Xiong, A., Choe, J., Richmond, J. E. & Kim, H. Active Zone Trafficking of
CaV2/UNC-2 Channels Is Independent of β/CCB-1 and α2δ/UNC-36 Subunits. *J. Neurosci.*
43, 5142–5157 (2023).

28. Saheki, Y. & Bargmann, C. I. Presynaptic CaV2 calcium channel traffic requires CALF-1 and
the α2δ subunit UNC-36. *Nat. Neurosci.* **12**, 1257–1265 (2009).

29. Zhan, H. *et al.* In vivo single-molecule imaging identifies altered dynamics of calcium
channels in dystrophin-mutant C. elegans. *Nat. Commun.* **5**, 4974 (2014).

30. Zhao, Y. & Pinaud, F. In Vivo Single-Molecule Tracking of Voltage-Gated Calcium Channels
with Split-Fluorescent Proteins in CRISPR-Engineered C. elegans. in *Single Molecule*
Microscopy in Neurobiology (eds Yamamoto, N. & Okada, Y.) vol. 154 11–37 (Springer US,
New York, NY, 2020).

31. Feng, S. *et al.* Improved split fluorescent proteins for endogenous protein labeling. *Nat.*
Commun. **8**, 370 (2017).

32. Tzur, Y. B. *et al.* Heritable custom genomic modifications in Caenorhabditis elegans via a
CRISPR-Cas9 system. *Genetics* **195**, 1181–1185 (2013).

33. Brenner, S. THE GENETICS OF *CAENORHABDITIS ELEGANS*. *Genetics* **77**, 71–94 (1974).

34. Laine, V., Frokjaer-Jensen, C., Couchoux, H. & Jospin, M. The alpha1 subunit EGL-19, the
alpha2/delta subunit UNC-36, and the beta subunit CCB-1 underlie voltage-dependent calcium
currents in Caenorhabditis elegans striated muscle. *J. Biol. Chem.* **286**, 36180–36187 (2011).

35. McDonald, K. *et al.* Using CRISPR knock-in of fluorescent tags to examine isoform-specific
expression   of   EGL-19   in   C.   elegans.

https://doi.org/10.17912/MICROPUB.BIOLOGY.000858   (2023)
doi:10.17912/MICROPUB.BIOLOGY.000858.

36. Mueller, B. D. *et al.* CaV1 and CaV2 calcium channels mediate the release of distinct pools of
synaptic vesicles. *eLife* **12**, e81407 (2023).

37. Zhao, J., Gao, L., Nurrish, S. & Kaplan, J. M. Post-synaptic GABAA receptors potentiate
transmission by recruiting CaV2 channels to their inputs. *Cell Rep.* **42**, 113161 (2023).

38. Serge, A., Bertaux, N., Rigneault, H. & Marguet, D. Dynamic multiple-target tracing to probe
spatiotemporal cartography of cell membranes. *Nat. Methods* **5**, 687–694 (2008).

39. Schutz, G. J., Schindler, H. & Schmidt, T. Single-molecule microscopy on model membranes
reveals anomalous diffusion. *Biophys. J.* **73**, 1073–1080 (1997).

40. Watanabe, S. *et al.* Ultrafast endocytosis at Caenorhabditis elegans neuromuscular junctions.
eLife **2**, e00723 (2013).

41. Ackermann, F., Waites, C. L. & Garner, C. C. Presynaptic active zones in invertebrates and
vertebrates. *EMBO Rep.* **16**, 923–938 (2015).

42. Kittelmann, M. *et al.* Liprin-α/SYD-2 determines the size of dense projections in presynaptic
active zones in *C. elegans*. *J. Cell Biol.* **203**, 849–863 (2013).

43. Stigloher, C., Zhan, H., Zhen, M., Richmond, J. & Bessereau, J.-L. The Presynaptic Dense
Projection of the *Caenorhabiditis elegans* Cholinergic Neuromuscular Junction Localizes
Synaptic Vesicles at the Active Zone through SYD-2/Liprin and UNC-10/RIM-Dependent
Interactions. *J. Neurosci.* **31**, 4388–4396 (2011).

44. Calahorro, F. & Izquierdo, P. G. The presynaptic machinery at the synapse of C. elegans. *Invert.*
Neurosci. **18**, 4 (2018).

45. Oh, K. H., Krout, M. D., Richmond, J. E. & Kim, H. UNC-2 CaV2 Channel Localization at
Presynaptic Active Zones Depends on UNC-10/RIM and SYD-2/Liprin-α in *Caenorhabditis*
elegans. *J. Neurosci.* **41**, 4782–4794 (2021).

46. Nonet, M. L. *et al. Caenorhabditis elegans rab-3* Mutant Synapses Exhibit Impaired Function
and Are Partially Depleted of Vesicles. *J. Neurosci.* **17**, 8061–8073 (1997).

47. Weimer, R. M. *et al.* UNC-13 and UNC-10/Rim Localize Synaptic Vesicles to Specific
Membrane Domains. *J. Neurosci.* **26**, 8040–8047 (2006).

48. Gracheva, E. O., Hadwiger, G., Nonet, M. L. & Richmond, J. E. Direct interactions between
C. elegans RAB-3 and Rim provide a mechanism to target vesicles to the presynaptic density.
Neurosci. Lett. **444**, 137–142 (2008).

49. Dittman, J. S. Unc13: a multifunctional synaptic marvel. *Curr. Opin. Neurobiol.* **57**, 17–25
(2019).

50. Sakamoto, H. Synaptic weight set by Munc13-1 supramolecular assemblies. *Nat Neurosci* **21**,
(2018).

51. Hu, Z., Tong, X.-J. & Kaplan, J. M. UNC-13L, UNC-13S, and Tomosyn form a protein code
for fast and slow neurotransmitter release in Caenorhabditis elegans. *eLife* **2**, e00967 (2013).

52. Liu, Q. *et al.* A Photoactivatable Botulinum Neurotoxin for Inducible Control of
Neurotransmission. *Neuron* **101**, 863-875.e6 (2019).

53. Gracheva, E. O. *et al.* Tomosyn Inhibits Synaptic Vesicle Priming in Caenorhabditis elegans.
PLoS Biol. **4**, e261 (2006).

54. McEwen, J. M., Madison, J. M., Dybbs, M. & Kaplan, J. M. Antagonistic Regulation of
Synaptic Vesicle Priming by Tomosyn and UNC-13. *Neuron* **51**, 303–315 (2006).

55. Hu, Z., Tong, X.-J. & Kaplan, J. M. UNC-13L, UNC-13S, and Tomosyn form a protein code
for fast and slow neurotransmitter release in Caenorhabditis elegans. *eLife* **2**, e00967 (2013).

56. Krout, M. *et al. C. elegans* Clarinet/CLA-1 recruits RIMB-1/RIM-binding protein and UNC-
13 to orchestrate presynaptic neurotransmitter release. *Proc. Natl. Acad. Sci.* **120**,
e2220856120 (2023).

57. Sakamoto, H. *et al.* Synapse type-specific molecular nanoconfigurations of the presynaptic
active zone in the hippocampus identified by systematic nanoscopy. Preprint at
https://doi.org/10.1101/2022.03.11.483942 (2022).

58. Maschi, D. & Klyachko, V. A. Spatiotemporal Regulation of Synaptic Vesicle Fusion Sites in
Central Synapses. *Neuron* **94**, 65-73.e3 (2017).

59. Südhof, T. C. The presynaptic active zone. *Neuron* **75**, 11–25 (2012).

60. Bademosi, A. T. *et al.* In vivo single-molecule imaging of syntaxin1A reveals
polyphosphoinositide- and activity-dependent trapping in presynaptic nanoclusters. *Nat.*
Commun. **7**, 13660 (2016).

61. Emperador-Melero, J. *et al.* Distinct active zone protein machineries mediate Ca2+ channel
clustering and vesicle priming at hippocampal synapses. *Nat. Neurosci.* **27**, 1680–1694 (2024).

62. Gaffield, M. A., Rizzoli, S. O. & Betz, W. J. Mobility of Synaptic Vesicles in Different Pools
in Resting and Stimulated Frog Motor Nerve Terminals. *Neuron* **51**, 317–325 (2006).

63. Park, H., Li, Y. & Tsien, R. W. Influence of Synaptic Vesicle Position on Release Probability
and Exocytotic Fusion Mode. *Science* **335**, 1362–1366 (2012).

64. Forte, L. A., Gramlich, M. W. & Klyachko, V. A. Activity-Dependence of Synaptic Vesicle
Dynamics. *J. Neurosci.* **37**, 10597–10610 (2017).

65. Shtrahman, M., Yeung, C., Nauen, D. W., Bi, G. & Wu, X. Probing Vesicle Dynamics in Single
Hippocampal Synapses. *Biophys. J.* **89**, 3615–3627 (2005).

66. Miki, T., Midorikawa, M. & Sakaba, T. Direct imaging of rapid tethering of synaptic vesicles
accompanying exocytosis at a fast central synapse. *Proc. Natl. Acad. Sci.* **117**, 14493–14502
(2020).

67. Südhof, T. C. The presynaptic active zone. *Neuron* **75**, 11–25 (2012).

68. Kobbersmed, J. R., Berns, M. M., Ditlevsen, S., Sørensen, J. B. & Walter, A. M. Allosteric
stabilization of calcium and phosphoinositide dual binding engages several synaptotagmins in
fast exocytosis. *eLife* **11**, e74810 (2022).

69. Bohme, M. A. Active zone scaffolds differentially accumulate Unc13 isoforms to tune Ca2+
channel-vesicle coupling. *Nat Neurosci* **19**, (2016).

70. Nakamura, Y. *et al.* Nanoscale Distribution of Presynaptic Ca2+ Channels and Its Impact on
Vesicular Release during Development. *Neuron* **85**, 145–158 (2015).

71. Rebola, N. *et al.* Distinct Nanoscale Calcium Channel and Synaptic Vesicle Topographies
Contribute to the Diversity of Synaptic Function. *Neuron* **104**, 693-710.e9 (2019).

72. Vyleta, N. P. & Jonas, P. Loose Coupling Between Ca^{2+} Channels and Release Sensors at a
Plastic Hippocampal Synapse. *Science* **343**, 665–670 (2014).

73. Reddy-Alla, S. Stable positioning of Unc13 restricts synaptic vesicle fusion to defined release
sites to promote synchronous neurotransmission. *Neuron* **95**, (2017).

74. Zhou, K., Stawicki, T. M., Goncharov, A. & Jin, Y. Position of UNC-13 in the active zone
regulates synaptic vesicle release probability and release kinetics. *eLife* **2**, e01180 (2013).

75. Zamponi, G. W. Regulation of Presynaptic Calcium Channels by Synaptic Proteins. *J.*
Pharmacol. Sci. **92**, 79–83 (2003).

76. Mathews, E. A. *et al.* Critical Residues of the *Caenorhabditis elegans unc-2* Voltage-Gated
Calcium Channel That Affect Behavioral and Physiological Properties. *J. Neurosci.* **23**, 6537–
6545 (2003).

77. Tan, C., Wang, S. S. H., De Nola, G. & Kaeser, P. S. Rebuilding essential active zone functions
within a synapse. *Neuron* **110**, 1498-1515.e8 (2022).

78. Neher, E. & Brose, N. Dynamically Primed Synaptic Vesicle States: Key to Understand
Synaptic Short-Term Plasticity. *Neuron* **100**, 1283–1291 (2018).

79. Paix, A., Folkmann, A. & Seydoux, G. Precision genome editing using CRISPR-Cas9 and
linear repair templates in C. elegans. *Methods* **121–122**, 86–93 (2017).

80. Hua Fu, B. X., Hansen, L. L., Artiles, K. L., Nonet, M. L. & Fire, A. Z. Landscape of
target:guide homology effects on Cas9-mediated cleavage. *Nucleic Acids Res.* **42**, 13778–
13787 (2014).

81. Arribere, J. A. *et al.* Efficient Marker-Free Recovery of Custom Genetic Modifications with
CRISPR/Cas9 in *Caenorhabditis elegans*. *Genetics* **198**, 837–846 (2014).

82. Feng, S. *et al.* Improved split fluorescent proteins for endogenous protein labeling. *Nat.*
Commun. **8**, 370 (2017).

83. Gottschalk, A. & Schafer, W. R. Visualization of integral and peripheral cell surface proteins
in live Caenorhabditis elegans. *J. Neurosci. Methods* **154**, 68–79 (2006).

84. Mahoney, T. R., Luo, S. & Nonet, M. L. Analysis of synaptic transmission in Caenorhabditis
elegans using an aldicarb-sensitivity assay. *Nat. Protoc.* **1**, 1772–1777 (2006).

85. Nawa, M. & Matsuoka, M. The Method of the Body Bending Assay Using Caenorhabditis
elegans. *BIO-Protoc.* **2**, (2012).

86. Sergé, A., Bertaux, N., Rigneault, H. & Marguet, D. Dynamic multiple-target tracing to probe
spatiotemporal cartography of cell membranes. *Nat. Methods* **5**, 687–694 (2008).

87. Duerr, J. Immunohistochemistry. *WormBook* https://doi.org/10.1895/wormbook.1.105.1 (2006)
doi:10.1895/wormbook.1.105.1.

88. Savitzky, Abraham. & Golay, M. J. E. Smoothing and Differentiation of Data by Simplified
Least Squares Procedures. *Anal. Chem.* **36**, 1627–1639 (1964).

89. Wagner, T. & Eglinger, J. thorstenwagner/ij-particlesizer: v1.0.9 Snapshot release. Zenodo
https://doi.org/10.5281/ZENODO.820296 (2017).

90. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods*
9, 676–682 (2012).

91. Schnitzbauer, J., Strauss, M. T., Schlichthaerle, T., Schueder, F. & Jungmann, R. Super-
resolution microscopy with DNA-PAINT. *Nat. Protoc.* **12**, 1198–1228 (2017).

Figure 1. **N-type VGCCs exhibit two distinct diffusive behaviors within multiple nanodomains that correlate with UNC-10 at AZ.**

(a) Schematic of CALM by pseudocoelomic microinjection of synthetic sfcherry (11) peptides to activate extracellular split sfCherry (1-10) tagged to UNC-36 and detect surface-localized VGCC complexes at *C. elegans* NMJ. **(b)** Sum projection of a dual-color SM CALM movie by HILO light sheet imaging of sCherry-UNC-36 (red) along a dorsal nerve cord (DNC) marked with GABAergic Punc-25::SNB-1::GFP (blue, top). Corresponding VGCC trajectory maps color-coded by time (Tmin = blue; Tmax = red, middle) and zoom of selected insets (bottom). Scale bars, 1 µm (middle) and 100 nm (bottom). **(c)** Ensemble mean square displacement (MSD ± SEM) of slow (black squares) and fast (blue dots) fractions for VGCCs (n = 42,391; 58 animals). MSD curves

are fitted with a confined 2D diffusion model and the fit 95% confidence bands are shown together
with the diffusion coefficient (D ± SEM) and the domain diameter (∅ ± localization precision) for
each fraction. **(d)** Representative trajectories of heterogeneous VGCC diffusive behaviors color-
coded by instantaneous diffusion coefficients ($D_{instantaneous}$). Trajectory start: yellow dot; trajectory
stop: red dot; scale bar: 100 nm. **(e)** Confocal and dual-color STED images of immunostained
surface sCherry-UNC-36 (red) and 5xMYC-UNC-10 (cyan) along dorsal nerve cords (DNC)
marked by Punc-25::SNB-1::GFP (blue). Scale bars, 1 μm. **(f)** Zoomed insets of STED images in
(e). Scale bars, 100 nm. **(g)** Correlation between the number of UNC-36 (red) and UNC-10 (cyan)
nanodomains per AZ (n = 516; 10 animals). Linear regression indicates proportional scaling
(y=1.03x, R^2: 0.932). Notched boxes: median ± interquartile range (IQR); whiskers: 1.5xIQR.
Dashed line marks the median number of UNC-36 nanodomains. **(h)** Nearest-neighbor distance
(NND) and area-equivalent circular diameter (AED) of nanodomains for UNC-36 (red; n =1416
NND; 993 AED, 10 animals) and UNC-10 (cyan; n = 1454 NND; 786 AED; 10 animals). Notched
boxes: median ± IQR; whiskers: 1.5xIQR. Dashed line marks the confinement size of the slow
regime for VGCCs in (C). **(i, j)** Puncta sizes of UNC-36 (red) and UNC-10 (cyan) (n = 516; 10
animals). Notched boxes: median ± IQR; whiskers: 1.5xIQR. Dashed line marks the confinement
size of the fast regime for VGCCs in (c).
960

961
962

Figure 2. UNC-10 enhances VGCCs diffusion and promotes the multi-nanodomain organization of VGCC.

(a) Percentage of paralyzed animals over time (mean ± SEM) in aldicarb-sensitivity assays, comparing WT and *unc-10(md1117)* genotypes (n = 90). Two-way ANOVA and Tukey's tests, ****p<0.0001. **(b)** Representative STED images of immunostained surface sCherry-UNC-36 (red) overlayed with confocal images of Punc-25::SNB-1::GFP (blue) at the dorsal nerve cords of WT or *unc-10(md1117)* animals. Scale bars, 1 µm and 100 nm for insets. **(c-e)** Number (c), distribution (d) and nearest-neighbor distance NND (e) of UNC-36 nanodomains per diffraction-limited puncta in WT (gray; n = 1076 puncta; 1416 NND; 10 animals) and *unc-10 (md1117)* mutants (red; n = 634 puncta; 855 NND; 7 animals). Notched boxes: median ± interquartile range (IQR); whiskers: 1.5xIQR. Comparisons to WT by Kruskal-Wallis and Dunn's test, ****p < 0.0001. **(f-h)** Mobility of VGCCs in WT (gray; n= 42,391 trajectories; 58 animals) and *unc-10 (md1117)* mutant (red; n= 18,645 trajectories; 34 animals) with fractions (f, mean ± SEM), diffusion coefficients (g, mean ± SEM), and nanodomain confinement size or diffusion range (h, mean diameter ± localization precision) for the channels' slow (top row, black) and fast (bottom row, blue) diffusive behaviors. Comparisons to WT by two-tailed T-test. ****p < 0.0001; **p<0.01; *p<0.05; ns, not significant.

980
981
Figure 3. The mobility and spatial organization of nanodomain-associated VGCCs is
regulated by UNC-10/RIM mediated SV coupling
(a) Numbers of body bending per min (mean ± SD) in trashing assays for animals (n = 36) of
indicated genotypes. Comparisons by Kruskal-Wallis and Dunn's test, ****p < 0.0001. **(b)**
Percentage of paralyzed animals over time (mean ± SEM) in aldicarb-sensitivity assays, for the
indicated genotypes (n = 90). Comparisons to WT at time points by two-way ANOVA and Tukey's
tests, ****p<0.0001; ns, not significant. **(c-d)** Mobility of VGCCs in WT (gray; n = 42,391
trajectories; 58 animals), *rab-3(js49)* (red; n = 33,175 trajectories; 107 animals), *unc-10 (K77/79E)*
(blue; n = 29,278 trajectories; 26 animals) and *rab-3; unc-10(K77E/K79E)* mutants (brown; n =
24,455 trajectories; 19 animals), including diffusion coefficients (mean ± SEM), nanodomain
confinement size or diffusion range (mean diameter ± localization precision) and fractions (mean
± SEM) for the channels' slow (c) and fast (d) diffusive behaviors. Comparisons by two-tailed T-
test. ****p<0.0001; ***p<0.001; **p<0.01; *p<0.05; ns, not significant. **(e)** Representative STED

images of immunostained surface sCherry-UNC-36 (red) overlayed with confocal images of Punc-
25::SNB-1::GFP (blue) at the dorsal nerve cords of WT or *rab-3; unc-10(K77E/K79E)* animals.
Scale bars, 1 μm and 100 nm for insets. **(f-i)** Quantifications of UNC-36 nanodomain nearest-
neighbor distance NND (f), puncta long/short axe ratio (g), nanodomain numbers per puncta (h)
and their distribution (i) in STED images from WT (gray; n = 1416 NND; 1076 puncta; 10 animals)
and *rab-3; unc-10(K77E/K79E)* mutants (brown; n = 509 NND; 244 puncta; 8 animals). Notched
boxes: median ± interquartile range (IQR); whiskers: 1.5xIQR. Comparisons to WT by Kruskal-
Wallis and Dunn's test, ****$p < 0.0001$; **$p < 0.01$; *$p < 0.05$.
1003

DNC / sCherry-UNC-36 / 5xMYC-UNC-10 / Punc-25::SNB-1-::GFP

Figure 4. The mobility and AZ organization of VGCCs is governed by SV priming and SNARE complex assembly.

(a) Percentage of paralyzed animals over time (mean ± SEM) in aldicarb-sensitivity assays, for the indicated genotypes (n = 90). Comparisons to WT at time points or across genotypes at t = 240 min by two-way ANOVA and Tukey's tests, ****p<0.0001. **(b-c)** Mobility of VGCCs in WT (gray; n = 42,391 trajectories; 58 animals), *unc-13L-S- (s69)* (red; n = 31,355 trajectories; 35 animals) and *unc-13L-S+ (e51)* mutants (blue; n = 19,899 trajectories; 34 animals), including diffusion coefficients (mean ± SEM) and nanodomain confinement size or diffusion range (mean diameter ± localization precision) for the channels' slow (b) and fast (c) diffusive behaviors. Comparisons by two-tailed T-test. ****p<0.0001; ***p<0.001; **p<0.01; *p<0.05; ns, not significant. **(d)** Representative dual-color STED images of immunostained surface sCherry-UNC-36 (red) and

5xMYC-UNC-10 (cyan) overlayed with confocal images of Punc-25::SNB-1::GFP (blue) at the
dorsal nerve cords of WT, *unc-13L-S- (s69)* and *unc-13L-S+ (e51)* animals. Scale bars, 1 μm. **(e-**
g) Quantifications of UNC-36 and UNC-10 nanodomain numbers per AZ (e), nanodomain sizes
(f), and puncta size at AZ (g) in WT (gray; n = 516 AZ puncta; 993 UNC-36 and 786 UNC-10
nanodomains; 10 animals), *unc-13L-S- (s69)* (red; n = 320 AZ puncta; 607 UNC-36 and 641 UNC-
10 nanodomains; 6 animals) and *unc-13L-S+ (e51)* mutants (blue; n = 205 AZ puncta; 655 UNC-
36 and 473 UNC-10 nanodomains; 4 animals). Notched boxes: median ± interquartile range (IQR);
whiskers: 1.5xIQR. Comparisons by Kruskal-Wallis and Dunn's test, ****$p < 0.0001$; **$p<0.01$;
*$p<0.05$; ns, not significant. **(h)** Percentage of animals displaying normal (grey) or "shrinker"
(green) locomotory phenotypes (mean ± SEM) in WT and transgenic animals expressing
ScBoNTB in GABAergic synapses (n = 50), over 3 independent batches of animals. **(i-j)** Mobility
of VGCCs in WT animals at all boutons (dark gray; n = 42,391 trajectories; 58 animals), in WT
animals at GABAergic boutons only (light gray; n = 19,034 trajectories; 42 animals) and in
ScBoNTB-expressing animals at GABAergic boutons only (green; n = 11,549 trajectories; 36
animals), including fractions (mean ± SEM), diffusion coefficients (mean ± SEM), and
nanodomain confinement size or diffusion range (mean diameter ± localization precision) for the
channels' slow (i) and fast (j) diffusive behaviors. Comparisons by two-tailed T-test. ****$p<0.0001$;
***$p<0.001$; **$p<0.01$; *$p<0.05$; ns, not significant.
1035

1036
1037
Figure 5. Distinct mobilities and nanodomain organization of VGCCs are differentially modulated by SV priming levels.
(a) Percentage of paralyzed animals over time (mean ± SEM) in aldicarb-sensitivity assays, for the indicated genotypes (n = 90). Comparisons to WT at time points t = 120min and across genotypes at t = 240 min by two-way ANOVA and Tukey's tests, ****p<0.0001; ***p<0.001; **p<0.01. **(b-c)** Mobility of VGCCs in WT (gray; n = 42,391 trajectories; 58 animals), *unc-13L-S- (s69)* (red; n = 31,355 trajectories; 35 animals), *unc-13L-S- (s69);tom-1(nu468)* (brown; n = 25,309 trajectories; 15 animals), *unc-13L-S+ (e51);tom-1(nu468)* mutants (cyan; n = 18,963 trajectories; 15 animals), and *tom-1(nu468)* mutants (green; n = 34,062 trajectories; 25 animals), including diffusion coefficients (mean ± SEM) and nanodomain confinement size or diffusion range (mean diameter ± localization precision) for the channels' slow (b) and fast (c) diffusive behaviors. Comparisons by two-tailed T-test. ***p<0.001; **p<0.01; *p<0.05; ns, not significant. **(d)** Representative dual-color STED images of immunostained surface sCherry-UNC-36 (red) and 5xMYC-UNC-10 (cyan) overlayed with confocal images of Punc-25::SNB-1::GFP (blue) at the dorsal nerve cords of WT, *unc-13L-S-(s69)*, *unc-13L-S-(s69);tom-1(nu468)*, *unc-13L-S+ (e51);tom-1(nu468)* and *tom-1(nu468)* animals. Scale bars, 1 μm. **(e-f)** Quantifications of UNC-36 and UNC-10 puncta size at

AZ (e) and nanodomain numbers per AZ (f) in WT (gray; n = 516 AZ puncta; 10 animals), *unc-13L-S-(s69)* (red; n = 320 AZ puncta; 6 animals), *unc-13L-S- (s69);tom-1(nu468)* (brown; n = 349 AZ puncta; 10 animals)*, unc-13L-S+ (e51);tom-1(nu468)* mutants (cyan; n = 276 AZ puncta; 8 animals) and *tom-1(nu468)* mutants (green; n = 529 AZ puncta; 10 animals). Notched boxes: median ± interquartile range (IQR); whiskers: 1.5xIQR. Comparisons by Kruskal-Wallis and Dunn's test, ****p < 0.0001; *p<0.05; ns, not significant.

1061
Figure 6. Primed SVs modulate the mobilities and multi-nanodomain organization of
VGCCs downstream of UNC-10.
(a-b) Mobility of VGCCs in WT (gray; n= 42,391 trajectories; 58 animals); *unc-10 (md1117)* (red;
n= 18,645 trajectories; 34 animals) and *unc-10(md1117);tom-1(nu468)* (brown; n = 20,882
trajectories; 26 animals) mutants, including diffusion coefficients (mean ± SEM), nanodomain
confinement size or diffusion range (mean diameter ± localization precision) and fractions (mean
± SEM) for the channels' slow (a) and fast (b) diffusive behaviors. Comparisons by two-tailed T-
test. ****p < 0.0001; ***p<0.001; **p<0.01; *p<0.05; ns, not significant. **(c)** Percentage of
paralyzed animals over time (mean ± SEM) in aldicarb-sensitivity assays, for the indicated
genotypes (n = 90). Comparisons to WT at time points or across genotypes at t = 240 min by two-
way ANOVA and Tukey's tests, ****p<0.0001; ns, not significant. **(d)** Representative STED
images of immunostained surface sCherry-UNC-36 (red) overlayed with confocal images of Punc-
25::SNB-1::GFP (blue) at the dorsal nerve cords of WT, *unc-10 (md1117)* and *unc-*
10(md1117);tom-1(nu468) animals. Scale bars, 1 μm and 100 nm for insets. **(e-f)** Quantifications
of UNC-36 nanodomain numbers per puncta (e; left), distribution (e; right) and nearest-neighbor
distance NND (f) in WT (gray; n = 1076 puncta; 1416 NND; 10 animals), *unc-10 (md1117)* (red;
n = 634 puncta; 855 NND; 7 animals) and *unc-10(md1117);tom-1(nu468)* (brown; n = 1087 puncta;
748 NND; 8 animals) mutants. Notched boxes: median ± interquartile range (IQR); whiskers:

1.5xIQR. Comparisons by Kruskal-Wallis and Dunn's test, ****$p < 0.0001$; **$p<0.01$; ns, not
significant.