Proliferation/Differentiation in intestinal organoids as a balance of ligand-modulated EGFR traffic

Caracci, MO[1*], Seidler, S, Muñoz-Nava, LM[1], Soetje, B[1], Michel, K[1], Bastiaens, PIH[1#].

[1]Department of Systemic Cell Biology, Max Planck Institute for Molecular Physiology, Dortmund, Germany
*Corresponding author
#Deceased

Summary

Expression and signaling competency at the plasma membrane of EGFR drives crypt proliferation vs villus differentiation by medium ligand-composition, aiding mouse intestinal organoids self-renewal and regeneration.

Abstract

Epidermal Growth factor (EGF) signaling is associated with (oncogenic) proliferation. Conversely, EGF-family ligands are able to trigger a differentiation program in cultured cells, an effect attributed to ligand affinity and EGFR phosphorylation. How EGF/EGFR driven proliferation-differentiation dynamics underlie tissue self-renewal has not been addressed. We show that culturing mouse small intestinal organoids (mSIOs) without EGF enhanced EGFR expression and base phosphorylation while maintaining a balanced development of proliferative crypts and differentiated villi. Addition of EGF or EREG triggers receptor endocytosis, reducing cell-surface and expression levels. While EGF promoted crypt proliferation, EREG promoted both proliferation and villus differentiation compared to untreated controls. Removal or re-introduction of EGF or EREG proved sufficient to induce development comparable to constant presence of ligands over 96h. Sub-saturating concentrations of EGF led to increased villus differentiation, resembling EREG treatments, suggesting that control over EGFR

34   endocytic cycle ultimately regulates the balance of proliferation and differentiation in

35   mSIOs

Introduction

EGF/EGFR signaling is essential for proliferation and self-renewal of the intestinal stem cells. Aberrant activity of EGF downstream signaling has been consistently associated with oncogenic transformations. EGF is a fundamental growth factor for many in vitro organoid culture systems (van der Vaart et al., 2021; Mahe et al., 2013; Wang et al., 2025; Hu et al., 2018). In mouse small intestinal organoids (mSIOs), Paneth cells secrete paracrine EGF within the stem cell niche in crypts (Sato et al., 2011a). Nevertheless, additional growth factors, including EGF, are added to the culture medium, mimicking the complexity of stromal compartment surrounding the intestinal epithelia (Sato et al., 2009).

EGFR is mainly expressed in the intestinal crypts (Yang et al., 2017) and inhibition of tyrosine kinase activity in mature organoids led to cell cycle arrest and quiescence of LGR5+ stem cells (Basak et al., 2017). EGFR inhibition also enhanced differentiation towards entero-endocrine cells, suggesting that perturbations to EGF/EGFR activation during distinct developmental stages could aid differentiation in mSIOs. Other EGF-family ligands such as NRG1, EREG and HB-EGF can replace EGF in mSIOs culture promoting organoid development (Lemmetyinen et al., 2023; Jardé et al., 2020; Chen et al., 2012), notably epithelial KO of NRG1 does not affect differentiation of secretory linage cells (Jardé et al., 2020). While NRG1 does not bind EGFR, EREG binds both EGFR and ERBB4 and has been used in human intestinal organoids to enhance cellular differentiation and crypt development (Childs et al., 2023).

Under basal medium conditions, mSIOs show both crypt proliferation and villus differentiation in two distinct, spatially separated compartments. In contrast, human intestinal organoids feature less cellular and structural diversity. Several remarkable efforts have been taken to enhance human organoid complexity by modulating cell signaling, including EGF inhibitors and EGF-family ligands. Nevertheless, the spatio-temporal responses of dynamic intracellular signaling networks to extracellular ligands have been largely overlooked. (Yang et al., 2025; Fujii et al., 2018; Sato et al., 2011b; He et al., 2022).

It is well stablished that EGFR signaling plays an integral role in partially opposing
phenotypic outcomes, including migration, apoptosis, proliferation and differentiation
and is generally acknowledged that these responses are a result of different affinities
among EGF-family ligands and their capacity to induce receptor phosphorylation
(Abud et al., 2021; Jendrisek et al., 2026). For instance, high affinity ligands such as
EGF, HB-EGF and TGFa, all produced in the intestinal epithelia, are often associated
to cell proliferation. In other hand, low affinity ligands such as EREG and AREG, which
are expressed by stromal cells in the intestine, are associated to cell differentiation
and migration (Jendrisek et al., 2026; Lemmetyinen et al., 2023). Interestingly, EGF
triggered responses are dependent on ligand concentration, where high, receptor
saturating concentrations promote cellular proliferation while low, sub-saturating
concentrations promote migratory behavior in cultured cells (Brüggemann et al., 2021;
Joshi et al., 2023). The fact the EGF-EGFR signaling is able to trigger multiple
phenotypical responses underscores the importance of this ligand-receptor pair in
development. However, these properties have mainly been demonstrated in
immortalized cell lines and this decision-taking has not yet been explored in complex
developing tissues under homeorhetic conditions.
86
Therefore, we studied mSIOs in early developmental stages during self-renewal and
regeneration from seeded single crypts and their subsequent maturation. For this, we
cultured mSIOs with either EGF or EREG to examine the differential effect of ligand
availability on sustaining organoid crypt and villus development. We find that
maintenance of plasma membrane (PM) EGFR localization drives villus development
regardless of ligand affinity.
93
94
95 Results/Discussion
96
97 <u>EGF promotes crypt proliferation and EREG promotes villus differentiation</u>
98
In order to understand how the balance within crypt proliferation and villus
differentiation is triggered by distinct EGF-family growth factors we stained organoids
with CD44, a bonafide crypt marker (Sumigray et al., 2018), and enterocyte marker

AldoB to identify the differentiated villus region (Serra et al., 2019). Images were
acquired by CLSM and maximum intensity projections from multiple confocal stacks
were masked and quantified for area, as surrogates for crypt and villus growth
triggered by our treatments (Figure 1A and B). Intestinal organoid crypts were grown
for 96h in control NR medium (Noggin, R-SPO1) or NR medium supplemented with
either EGF (50 ng/mL), EREG (50 ng/mL). EGF and EREG as high and low affinity
ligands of EGFR respectively, are known to induce receptor phosphorylation and
endocytosis (Freed et al., 2017).
110
EGF and EREG addition significantly enhanced Organoid growth compared to NR
medium, however, Crypt Area was differentially enhanced by EGF, while Crypt and
Villus area was enhanced by EREG (Figure 1C) The relative ratio of Crypt-to-Villus
(CD44/ALDO) showed that EGF strongly favored the Crypt proliferation over Villus
differentiation, while NR medium and EREG maintained balanced development of
Crypt and Villus, despite the overall larger size of EREG treated organoids relative to
NR treatment (Figure 1D).
118
Crypt number was also greatly enhanced by EGF, while EREG only had small effect
in increasing the number of crypts relative to NR organoids (Figure S1A). Crypt
establishment is a direct result of apical constriction by Paneth cells(Langlands et al.,
2016). Paneth cells are found in the crypt, interspaced between Intestinal Stem cells
(ISCs) and their abundance is often correlated to ISCs numbers (Tallapragada et al.,
2021). Immuno-fluorescence staining showed that neither EGF or EREG enhanced
the number of Paneth cell, regardless of the effect observed in crypt area (Figure S1B).
126
Results suggests that both EGF and EREG are capable of inducing organoid
development through distinctive regulation of the proliferation/differentiation balance
with EGF promoting CD44+ crypts enlargement over differentiation into ALDO+ villus
region. On the other hand, EREG promotes growth of both crypt and villus regions,
thus suggesting that EGF-family ligands promote mSIOs development through
distinctive downstream mechanisms.
133
134 <u>EGF and EREG mediated endocytosis reduces EGFR PM localization and expression.</u>
135

EGF and EREG are known to bind, induce endocytosis and phosphorylation of EGFR.

EREG is also a ligand of ERBB4, however expression of this receptor is not detected

in small intestinal epithelia or stroma (Lemmetyinen et al., 2023). Saturating EGF

concentrations are commonly used in Organoid culture (Shankaran et al., 2021)at

which most of the PM receptor is internalized and undergo lysosomal degradation as

demonstrated in cultured immortalized cells (Brüggemann et al., 2021). In contrast,

EREG binding to EGFR promotes recycling back to the plasma membrane (Roepstorff

et al., 2009). In order to quantify end-point surface levels of EGFR, organoids were

incubated for 30 min with 50 ng/mL of EGF-647 before fixation and staining (Figure

1E). We observe that most of the internalized EGF-647 is located in the crypt regions

with few vesicles being observable in the villus regions, this is consistent with the

literature stating that EGFR is highly expressed in cycling cells within the intestinal

crypt (Yang et al., 2017). EGF-647 endocytosis was observed in organoids grown in

NR or EREG medium, organoids grown with unlabeled EGF for 96h showed very low

endocytosis suggesting reduced levels of PM EGFR (Figure 1F). Absence of ligands

allowed for accumulation of surface EGFR throughout the 96h developmental window

and EREG mediumted recycling back to the plasma membrane maintained high

surface levels.

EGF-647 endocytosis showed co-localization with late endosome marker Rab7

consistent with the use of saturating amounts of ligand (Figure 1F; Figure S2A and B).

DLS (Digital Lightsheet) imaging of EGF-647 endocytosis shows an initial association

to the crypts basal membrane followed by the entrance of vesicles travelling adjacent

to the inner membranes towards the apical side (Figure S2C and Video1). While basal

to apical trafficking is clearly observed we cannot discard the possibility of apical

endocytosis of EGF-647 from the lumen, as cell extrusions within the crypts are

observable in the time frame of our experiments which could allow EGF-647 to seep

into the luminal compartment. Transcytosis of EGF from basal to apical has been

proposed in MDCK epithelial layers (Brändli et al., 1991), however little is known of

the mechanisms involved

As a PM receptor, EGFR phosphorylation is highly dependent on its interaction with

ligands in the extracellular medium, nevertheless gene amplification enhancing

expression levels or point mutations affecting inhibitory mechanisms can lead to EGFR

phosphorylation in absence of ligand (Stanoev et al., 2018; Arteaga and Engelman, 2014). Immunoblotting of both total and phosphorylated EGFR Y1068 showed that EGF and EREG grown organoids had a significant reduction in EGFR protein expression compared to NR grown organoids (Figure 1G). Immuno-fluorescence staining pEGFR Y1068 and CD44 after treatments, showing an increase in phosphorylation staining only in crypts growth without EGF or EREG while organoid grown with either ligand showed reduced staining (Figure 1H).

EGFR endocytic cycle is fundamental to regulate receptor phosphorylation, allowing for degradation in late endosomes or the interaction with protein tyrosine phosphatases (PTPs) within endomembrane compartments (Baumdick et al., 2015; Joshi et al., 2023; Stanoev et al., 2018). In turn, NR grown organoids show an accumulation of pEGFR likely due to endogenously expressed EGF by Paneth cells and amplification by autocatalytic activation of PM bound receptors. It is important to consider that medium changes every 48h imply 2 "pulses" of fresh EGF or EREG in a 96h developmental window, in between pulses, biosynthetic and endocytic mechanisms allow for the replenishment of surface EGFR (Scharaw et al., 2016).Surprisingly while EREG treatment did allow for significantly higher EGFR PM levels, total levels were greatly diminished, despite this ligand promoting receptor recycling over late endosome trafficking (Figure 1G and H).

<u>Temporal modulation of EGF-family ligands during development alters villus differentiation</u>

In vitro cell culture often promotes proliferating conditions which ultimately allow for biomass expansion required for biochemical assays, and organoid culture is no exception. In the conception of ENR medium, addition of EGF and R-SPO1 were sufficient to induce organoid growth from single crypts and the addition of Noggin proved instrumental for organoid passaging over time (Sato et al., 2009). While lower concentrations of EGF (e.g 10 ng/mL) were also tested in this seminal work with similar results, little consideration was given to organoid developmental paths under evolving availability of EGF in regards to ligand concentration or duration of stimuli.

To test the temporal requirements of EGF/EGFR signaling in organoid development,
organoids were grown for 96h with either EGF or EREG. However a complete medium
exchange with NR medium (No EGF or EREG) was done after 1h or after 48h of
incubation (Figure 2A). Developing from single crypts, organoids activate a
regeneration program that initially drives proliferation into a fetal-like state where
differentiation and stem cells markers are mostly absent (Sprangers et al., 2021),
budding within the organoid usually appears at the 48h mark where mature crypts will
develop (Sato et al., 2009) (Figure S3A). After 48h in culture, most organoids feature
a round morphology that is entirely CD44 positive, consistent with an undifferentiated
proliferating organoid (Serra et al., 2019). About 25% of them showed AldoB staining,
suggesting initial establishment of a villus region. The percentage of organoids
showing AldoB staining increases with EREG treated crypts, as shown for 96h
organoids grown with EREG. Overall, total area of organoids was not changed in any
of the treatments regardless of presence or absence of AldoB staining (Figure S3B-
E). EGF-647 endocytosis was also clearly observed in NR and EREG grown organoids
but not in EGF grown organoids as shown for 96h organoids. Endocytosis was
possible in organoids displaying either crypt or round morphology, often confined to
proliferating cells as noted by Ki67 staining (Figure S3F).
221
Regardless of EGF or EREG removal time-frame, organoids showed comparable total
area after 96h in culture. Moreover, 1h and 48h of EGF containing medium was
sufficient to induce crypt growth comparable to 96h with constant EGF. EREG induced
villus differentiation showed a similar trend as EGF with crypt development.
Surprisingly, 1h EGF treatment not only induced crypt growth, but also enhanced villus
differentiation to levels similar to the ones observed in EREG treatment (Figure2B and
C). Removal of EGF or EREG did not affect crypt number, however, removal of EGF
increased Paneth cell number compared to 96h control (FigureS1C and D).
Alternatively, organoids were treated for 96h and 1h with 50ng/mL of NRG1 (Figure
S1I), An EGF-family ligand know to bind ERBB3 and ERBB4 but not EGFR. NRG1
was able to induce organoid growth as previously shown (Jardé et al., 2020;
Lemmetyinen et al., 2023). Notably, 1h treatment with NRG1 and subsequent
withdrawal failed to induce organoid growth to 96h levels, with a significant growth-
increase relative to NR-grown organoids (Figure S1J and K). Therefore EGFR is likely
to induce robust signaling responses under EGF and EREG treatments.

96h and 48h incubation with EGF led to a near complete depletion of surface EGFR,

as very low EGF-647 endocytosis was detected., In contrast, 1h incubation with EGF

exhibited high levels of endocytosis suggesting high levels of surface EGFR, similar

to EREG treatments which increased over EREG-starvation times (Figure 2D). EGFR

phosphorylation was enhanced by EGF or EREG removal compared to the 96h

treatments further highlighting that phosphorylation of EGFR requires PM localization

(Figure S1E).

In a complementary experiment, single crypts were seeded in NR medium and then

ligands were re-introduced after 48h or 1h of growth factor starvation. (Figure 2E).

EGF and EREG-containing medium increased organoids total area and crypt area. In

contrast to early EREG treatment, addition of EREG after growth factor starvation

enhanced crypt growth similar to EGF and at the same time increased villus growth as

previously established (Figure 2F-G). Likewise, there was a marked increase in the

number of organoids featuring multiple crypts, although only EGF treatment enhanced

the number of Paneth cells (Figure S1F and G).

As previously shown, EGF and EREG reintroduction into the culture medium greatly

reduced endocytosis of labelled EGF-647 with EREG treated organoids retaining

some endocytic capacity consistent with EGFR recycling towards the plasma

membrane (Figure 2H). EGF or EREG removal from the culture medium allowed for

increased phosphorylation of EGFR, likewise, re- introducing EGF or EREG at 48h

after seeding significantly reduced phosphorylation, while brief 1h starvation did not

change phosphorylation levels compared to the NR condition (Figure S1H).

Altogether, these results highlight the remarkable robustness of EGFR signaling in

response to high affinity and low affinity ligands, effect that could not be replicated by

NRG1 treatment, which does not bind EGFR. EGF or EREG treatment at any point

tested was able to reduce EGFR surface levels compared to NR and lead to similar

total area growth, with 1h treatments upon seeding being sufficient to induce crypt and

villus development respectively.

<u>Sub-saturating concentrations of EGF enhance both crypt proliferation and villus</u>
<u>differentiation</u>
272
Given that organoids can develop in EGF-free medium, albeit at smaller size of crypt
and villus; and that EREG, a low affinity ligand of EGFR, induces organoid
development while maintaining high levels of surface EGFR, we tested if lower, non-
saturating concentrations of EGF could sustain organoid growth. Single crypts were
cultured for 96h with EGF concentrations decreasing by one order of magnitude of
starting at 50 ng/mL to 0.05 ng/mL (Figure 3A). NR treatment (0 ng/mL of EGF) showed
consistent development compared to 50 ng/mL as shown in previous experiments,
however, concentrations in between showed remarkable differences in development.
281
282 All EGF treatments enhanced organoid total area compared to NR, however crypt
283 growth was significantly enhanced starting at 0.5 ng/mL. Notably, villus differentiation
284 was increased starting at 0.05 ng/mL before being reduced at 50 ng/mL (Figure 3B).
285 Likewise, CD44/ALDO ratio was enhanced towards crypt development starting at 0.5
286 ng/mL and increased at 50ng/mL as villus area is reduced (Figure 3C).
287
288 EGF-647 endocytosis also showed a negative correlation between surface EGFR and
289 EGF concentration. Notably, concentrations that enhanced villus development showed
290 surface EGFR levels similar to EREG treatments (Figure 3D).
291
292 <u>Multiple pulses of EGF deplete EGFR surface expression reducing villus differentiation</u>
293
Removing saturating amounts of EGF from the culture medium, EREG treatment and
sub-saturating EGF concentration all are able to maintain a high level of surface EGFR
expression compared to long term EGF treatment. A combination of biosynthesis and
endocytic cycling is constantly supplying EGFR to the cell surface, EGF has also been
shown to increase EGFR mRNA levels (Scharaw et al., 2016), thus constant turnover
is likely critical for the phenotypical outcomes we observe.
300
To further test this idea, organoids were grown for 96h, however medium was
exchanged only 1, 2 or 4 times during this time window in order to constantly deplete
surface EGFR in developing organoids (Figure 4A). Total Area of the organoids

remained unchanged in EGF and EREG regardless of the number of medium
changes, showing that fresh medium addition does not change overall organoid
growth (Figure 4B). Contrastingly, Crypt area was increased while Villus area was
reduced by multiple EGF medium changes, an effect that was not observed with EREG
treatment that maintained Crypt Area in all treatments. 4x EREG treatments
significantly reduced Villus Area compared to 96h and 48h, however this change did
not significantly affect Crypt/Villus balance (Figure 4C and D). As expected, multiple
EGF treatments greatly reduced EGF-647 endocytosis while EREG and NR
treatments maintained similar levels of endocytosis in all treatments (Figure 4E).

<u>Concluding Remarks</u>

Human plasma levels of EGF range between 0.04 and 0.1 ng/mL (Alshammari et al.,
2026), whereas, concentrations of up to 300 ng/mL have been identified in rodent milk
and is essential for early gastrointestinal development in newborns. After weaning and
with age, EGF levels in the intestine are greatly reduced (Schaudies et al., 1989;
Fisher and Lakshmanan, 1990). EGF expression is also upregulated in Paneth cells
upon injury in the intestinal epithelia (Calafiore et al., 2023). Absolute concentrations
of EGF in tissues could also be obscured by tissue geometry, such as the confined
space within an intestinal crypt, allowing for a higher concentration of ligand within a
lower volume (Tschumperlin et al., 2004; Schmick and Bastiaens, 2014) In all, the
intestinal tissue is exposed to changing of ligands throughout development, with higher
levels often associated with epithelial regeneration processes interspersed within
periods of low ligand availability with balanced proliferation and differentiation. In turn,
periods of low ligand availability enhance long term stability of EGFR expression and
enrichment at the plasma membrane, making intestinal crypts susceptible to changes
in the concentration of ligands in the environment.

Saturating amounts of EGF might be instrumental for passaging and expanding
intestinal organoids, especially in when seeding single stem cells (Boretto et al., 2024).
However, seeding single crypts allows for a pre-stablished cellular niche to trigger
initial organoid regeneration, even in the absence of medium EGF. Over 48h, the
organoids establish a fetal-like morphology which was not influenced by EGF or EREG
as organoids developed equally in total area regardless of presence or absence of

ligand. Similar developmental profiles could be induced by re-introducing EGF or EREG after 48h and by removing ligands after 1h of incubation upon crypt seeding, highlighting the robustness of EGFR mediated signaling.

To couple phenotype to EGFR activity, we incubated organoids with labelled EGF-647 and examined endocytosis as measurement of the amount of EGFR present in the PM after treatments. It is well established in the literature that receptor-bound EGF is released under acidic pH of endosomal compartments. Whether this applies to EREG and EREG-triggered recycling produces ligand-less surface receptors is unknown. Labelled EGF has been extensively used to quantify surface levels and/or total receptor levels in cell lines (Stanoev et al., 2018; Joshi et al., 2023; Roepstorff et al., 2009).This process usually include incubations and washes with ice-cold buffers. These treatments are not possible in organoids where cold washes could compromise the integrity of the Matrigel matrix sustaining organoid morphology and the intracellular cytoskeletal network.

We observe that EGF treatment might promote receptor degradation, thus reducing EGFR endocytosis of EGF-647, it is also possible that smaller vesicles associated to clathrin mediated endocytosis of EGFR monomers could not be detectable by CLSM over the background signal. In other hand, higher levels of surface receptor in NR and EREG treatment enhance non-clathrin mediated endocytosis of EGFR clusters within larger vesicles and higher number of fluorescent molecules which facilitate detection.

On the other hand EREG, a low affinity EGFR ligand that has been shown to promote sustained EGFR phosphorylation and differentiation in cultured cells (Freed et al., 2017). After 96h of growth we were unable to observe increased phosphorylation of EGFR in EREG treated organoids, pEGFR levels quantified by immunofluorescence were often comparable to EGF treated organoids, also total levels of EGFR were reduced by EREG 96h treatment. EREG-induced EGFR endocytosis leads to rapid recycling back to the plasma membrane and reduced ubiquitination compared to EGF treatment; however, a fraction of EREG-bound receptors shows colocalization with late endosomal markers (Roepstorff et al., 2009; Deguchi et al., 2024) which could account for degradation.

Methods

Mouse intestinal organoids culture

AdDF+++ medium containing Advanced Dulbecco's modified Eagle's medium/F12 supplemented with 100 u/mL penicillin/streptomycin, 10 mM HEPES, 2mM Glutamax. ENR medium consists of AdDF+++ supplemented with 1x B27 (Life Technologies) and 1.25 mM N-acetylcysteine, 50ng/mL mEGF (Peprotech, 315-09), 500 ng/mL R-Spondin 1 (Peprotech, 315-32) and 100 ng/mL Noggin (Peprotech, 250-38). Alternatively, medium without EGF was used as "NR medium". Cryopreserved mouse small intestinal organoids were purchased form STEMCELL Technologies (Cat #70931) and grown in Matrigel 30 μL droplets in 24 well plates in ENR medium for 96-120h before splitting by mechanical disruption with yellow pipette tips. Alternatively for treatments, NR medium was supplemented with 50 ng/mL of EREG (Biotechne, 1195-EP-025,) or 50 ng/mL NRG1 (Biotechne-377-HB-050/CF).

Immuno-staining

Single organoid crypts were seeded in 20 μL Matrigel droplets in x-well culture chambers 8 wells on coverslips (Sarstedt, 94.6190.802). After 96h of growth organoids were fixed for 40 min in 4% PFA. Then washed 3 times with IF-buffer (0.2% Triton X100, 0.05% Tween20) for 5 min and permeabilized with PBS 0.5% Triton X100 for 1h. Organoids were then blocked with IF-Buffer supplemented with 3% BSA. Alternatively, for staining of phosphorylated EGFR (pEGFR), BSA was replaced by 50% Goat Serum (Invitrogen, 50197Z) and 50% IF buffer for both blocking and antibody staining. Primary antibodies CD44 (1:500; BD Pharmigen™, 560568), Lysozyme (1:1000, DAKO, A0099) and Aldolase (1:500, ab153828, Abcam), pEGFR (1:250,EP774Y,Abcam), Ki-67 (1:500, 55060, BD Biosciences) , Rab7 (1:500, D95F2, Cell Signaling Technology) were incubated overnight with agitation at 4°C. Organoids were washed 3 times with IF Buffer for 5 min and then incubated for 1h with Alexa Fluor secondary antibodies (1:500,Thermo-Fisher) and DAPI (1:1000). Alexa Fluor 488 (A11008, A11001), Alexa Fluor 546 (A11081, A11030) and Alexa Fluor 647

405 (A31573, A21247) secondary antibodies were used. After washing with IF buffer,
406 samples were kept in Clearing Solution (50% v/v Glycerol, 2.5M Fructose) up to
407 imaging.
408

<u>Confocal Laser Scanning Microscopy (CLSM) imaging</u>

Organoids were imaged using Leica Confocal TCS SP8 with a 20x/0.75 NA air
objective. 2 μm spaced z-stacks were acquired in the fluorescence channels suitable
for each sample. Area quantifications were performed using binary masks thresholded
from Maximum Intensity Projections of either DAPI, CD44 or AldoB representing total,
crypt and villus area respectively. Alternatively, organoid crypts were imaged with
63x1.4 NA oil-immersion objective for pEGFR and CD44 imaging.

<u>EGF-647 endocytosis</u>

The His-CBD-Intein-(Cys)-hEGF-(Cys) plasmid was kindly provided by Prof. Luc
Brunsveld, University of Technology, Eindhoven. EGF-647 was purified in house as
previously described (Sonntag et al., 2014; Joshi et al., 2023). For endocytosis in
organoids, medium was aspirated from x-well culture chambers, 8 wells on coverslips
(94.6190.802, Sarstedt) and rinsed with warm AdDF+++ once. Subsequently,
organoids were incubated for 30 min with 50ng/mL of EGF-647 before fixation with 4%
PFA. Organoids were imaged in Leica Confocal SP8 with 63x oil immersion objective,
crypts were identified by Ki-67 staining. Cellular area was masked with Ki67 and Rab7
staining and the number vesicles were quantified using the find maxima command in
Fiji.

<u>Immunoblotting</u>

Organoid samples were pelleted, resuspended in modified RIPA buffer (50 mM Tris
HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA, 1 mM EGTA, 0.5% Triton X-100, and 0.5%
sodium deoxycholate) supplemented with Complete Mini EDTA-free protease inhibitor
(Roche Applied Science) and phosphatase inhibitor cocktail 2 and 3 (1:100, P5726
and P0044, Sigma–Aldrich), frozen in liquid nitrogen and stored in -70°C freezer.

Samples were then dissociated using 10-gauge needle syringes and centrifuged at 12,000 rpm for 15 min at 4°C. Protein concentration was determined by BCA (Thermo Scientific, 23235) and 50 μg of protein was loaded in 8% SDS/PAGE gels and then transferred into PVDF membranes. Membranes were then blocked in intercept buffer and incubated with primary antibody overnight. EGF Receptor (C75B9) (1:1000, Cell Signaling Technology, 2646) for total EGFR, EGFR Y1068 (AP774Y) (1:500; Abcam, 40815) for phospho EGFR (pEGFR), and α-Tubulin (1:10000; Merck, T6074,) were used. Secondary antibodies IRDye 680 donkey anti-Rabbit IgG (1:10000), IRDye 800 donkey anti-Mouse IgG (1:10000) and IRDye 800 donkey anti-Rabbit IgG (1:10000) (LI-COR Biosciences) were used.

Digital Lightsheet (DLS) Microscopy

Organoids were grown in U-Shaped Capillaries 20 x 1.5 mm (Leica, 158007061) inside Glass-Bottom 35 mm MatTek Dishes (MatTek Life Sciences, P35G-1.5-10-C) for 96h in NR medium. Before imaging, organoids were treated for 1h with 1:500 Cell Mask™ Green Actin Tracking stain (Invitrogen, A57243) and Hoechst 33342 (Invitrogen, C10340 G). Afterwards organoids were washed twice with warm AdDF+++ medium and returned to fresh NR medium. Organoids were imaged in Leica TCS SP8 Digital Lightsheet (DLS) DMi850 using HC PL FLUOTAR 5x/0.15 excitation objective and FLUOTAR L25x/0.95 W DLS detection objective. 50ng/mL of EGF-647 was added after calibration and ROI selection.

Statistical Analyses

Analysis was performed with at least 3 biological replicates (N=3) unless otherwise stated. Scatter plots show mean±95%CI and bar graphs show mean±SD. For single group comparisons Krustal-Wallis with Dunn's multiple comparisons test was used. For multiple group comparisons Two-way ANOVA with Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown. All analyses and graphs were implemented using Prism GraphPad

Acknowledgments

This project was supported by Max Planck Society for the Advancement of Science. We thank Dr. Malte Schmick for invaluable comments in the production of this manuscript. Special thanks to Dr. Sven Müller for technical support with confocal and DLS microscopy. Figure 1E was created with BioRender.com.

MOC: conceptualization, data curation, formal analysis, experimental methodology, visualization, writing and editing, SS: experimental methodology, editing, LMMN: conceptualization, writing and editing, BS: conceptualization, data curation, visualization, editing, KM: experimental methodology, editing. PHIB: Conceptualization, funding acquisition, supervision, project administration.

Prof. Dr. Philippe I.H. Bastiaens, passed away on May 15, 2025. All authors agree that his inclusion as a co-author is appropriate to honor his intellectual contribution to this work.

The authors state that this research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Figure 1. EGFR surface depletion by EGF promotes Crypt proliferation while EREG promotes both proliferation and differentiation. (A) Maximum intensity projections (top) were binarized (bottom) to quantify the area of DAPI (nuclei, blue), CD44 (Green) and AldolaseB (Magenta) used to quantify "Total", "Crypt" and "Villus" area respectively (B-H) Organoids were grown, with NR medium or supplemented with either EGF (50ng/mL) or EREG (50ng/mL) for 96h with a complete medium change at 48h. (B) Organoids were immuno-stained with CD44 (Green) AldoB (Magenta) and DAPI (Blue). Scale Bar 100 μm. (C) Quantification of the areas as shown in (A). Scatter plots with mean±95%CI. N=3; n=57-67 organoids/condition; significance was calculated by Kruskal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown (D) Relative ratio of CD44 to ALDO areas from (C). Scatter plot with mean±95%CI; significance was calculated by Kruskal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown. Values above 1 suggests organoid development tends towards crypt proliferation while values below 1 tend to villus differentiation (E) Alexa Fluor-647 labelled EGF (50 ng/mL) is added to organoids after 96h of growth, after 30 min of incubation organoids were fixed and stained. Alexa Fluor-647 positive endosomes indicate the level of surface EGFR and consecutive endocytosis upon ligand binding. (F) (Left) Representative micrographs of organoids after immunofluorescence staining against Ki67 (Yellow) and Rab7 (Magenta). EGF-647 internalized through EGFR endocytosis is labelled in (Green). Scale bar 20μm. (Right) EGF-647 vesicles per area as scatter plots mean±95%CI; N=3, n=28-37 crypts/condition; significance was calculated by Kruskal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown. (G) (Left) Immunoblot of pEGFR, EGFR and a-TUB with approximate MW in kDa. (Right) Bar Graphs of mean±SD of pEGFR or EGFR relative to a-TUB. N=4. (H) (Left) Ratio of pEGFR to CD44 Scatter plots of mean±95%CI. N=3, n=32-60 crypts/condition; significance was calculated by Kruskal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown. (Right) Representative micrographs of organoids after immunofluorescence staining against pEGFR (top) and CD44 (bottom). Scale bar 20μm.

Figure 2. Removing EGF or EREG after crypt seeding does not modify developmental trayectories. (A-D) Organoids were grown, with NR medium supplemented with either EGF (50ng/mL) or EREG (50ng/mL) for 48h or 1h, then complete medium was

replaced with NR medium until fixation at 96h. (B) Scatter plots showing mean±95%CI, N=3, n=57-79 organoids/condition of masked Total, CD44+ and ALDO+ areas. Significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown. (C) Relative ratio of CD44 to ALDO areas from (B). Scatter plot mean±95%CI of relative ratio within CD44/ALDO areas. Values above 1 suggests organoid development tends towards crypt proliferation while values below 1 tend to villus differentiation. Significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test. p-values <0.05 are shown. (D) EGF-647 vesicles per area as scatter plots showing mean±95%CI; N=3, n=28-41 crypts/condition; significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown. (E-H) Organoids were grown for 1h and 48h with NR, then complete medium was replaced with NR medium supplemented with either EGF (50ng/mL) or EREG (50ng/mL). (F) Scatter plots showing mean±95%CI, N=3, n=57-73 organoids/condition of masked Total, CD44+ and ALDO+ areas. Significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown, ns non-significant. (G) Relative ratio of CD44 to ALDO areas from (B). Scatter plot mean±95%CI of relative ratio within CD44/ALDO areas. Values above 1 suggests organoid development tends towards crypt proliferation while values below 1 tend to villus differentiation. Significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown. (H) EGF-647 vesicles per area as scatter plots showing mean±95%CI; N=3, n=20-38 crypts/condition; significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown.

Figure 3. Subsaturating concentrations of EGF trigger both crypt and villus growth. (A-D) Organoids were grown, with NR medium, supplemented with EGF at different concentrations: 50, 5, 0.5 and 0.05 ng/mL or EREG 50 ng/mL for 96h before fixation and immno-staining. (B) Scatter plots showing mean±95%CI, N=3, n=44-71 organoids per condition, significance was calculated by Kruskal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown (C) Relative ratio of CD44 to ALDO areas from (B). Scatter plot mean±95%CI of relative ratio within CD44/ALDO areas. Values above 1 suggests organoid development tends towards crypt proliferation while values

below 1 tend to villus differentiation. significance was calculated by Kruskal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown (D) EGF-647 vesicles per area as scatter plots showing mean±95%CI; N=3, n=39-48 crypts/condition; significance was calculated by Kruskal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown

Figure 4. Multiple EGF or EREG medium changes reduces villus differentiation. (A-E) Organoids were grown, with NR medium or supplemented with either EGF (50ng/mL) or EREG (50ng/mL) for 96h with a complete medium change every 24h (4x), 48h (2x) or 96h (1x). (B-C) Scatter plots showing mean±95%CI, N=3, n=47-67 organoids/condition of masked areas corresponding to (B) total Area and (C) CD44+ and ALDO+ areas; significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown. (D) Relative ratio of CD44 to ALDO areas from (C). Scatter plot mean±95%CI of relative ratio within CD44/ALDO areas. Values above 1 suggests organoid development tends towards crypt proliferation while values below 1 tend to villus differentiation. significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown (E) EGF-647 vesicles per area as scatter plots showing mean±95%CI; N=3, n=21-40 crypts/condition; significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown.

Figure S1. Effect of EGF/EREG temporal modulation in Crypt number, Paneth cell number and pEGFR staining. (A-B) Organoids were grown with NR medium or supplemented with either EGF (50ng/mL) or EREG (50ng/mL) for 96h with a complete medium change at 48h (A) Percentage of Organoids displaying either 1-2 or 3 or more crypts. N=3 (B) (Left) Organoids were immuno-stained with CD44 (Green) and Lysozyme (Lyz, Magenta). Scatter plots showing mean±95%CI of Paneth cells number per organoids was quantified in each condition. Quantification of n=43-56 organoids/condition, N=3, significance was calculated by Krustal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown. (C-E) Organoids were grown, with NR medium supplemented with either EGF (50ng/mL) or EREG (50ng/mL) for 48h or 1h, then complete medium was replaced with NR medium until fixation at 96h.

(C) Percentage of Organoids displaying either 1-2 or 3 or more crypts. N=3 (D) Scatter plots showing mean±95%CI of Paneth cells number per organoids was quantified in each condition. Quantification of n=42-61 organoids per condition, N=3, significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown (E) Ratio of pEGFR to CD44 Scatter plots of mean±95%CI. N=3, n=21-49 crypts/condition; significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown. (F-H) Organoids were grown for 1h and 48h with NR, then complete medium was replaced with NR medium supplemented with either EGF (50ng/mL) or EREG (50ng/mL). (F) Percentage of Organoids displaying either 1-2 or 3 or more crypts. N=3 (G) Scatter plots showing mean±95%CI of Paneth cells number per organoids was quantified in each condition. Quantification of n=50-69 organoids/condition, N=3, significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown (H) Ratio of pEGFR to CD44 Scatter plots of mean±95%CI. N=2-3, n=39-54 crypts/condition; significance was calculated by Two-way ANOVA and Sidak's multiple comparisons test multiple comparisons test. p-values <0.05 are shown. (I) Organoids were grown, with NR medium or NR medium supplemented with NRG1 (50ng/mL) for 96h or 1h. After 1h complete medium was replaced with NR medium until fixation at 96h. (J) Organoids were immuno-stained with CD44 (Green) AldoB (Magenta) and DAPI (Blue). Scale Bar 100 μm. (J-K) Scatter plots showing mean±95%CI of masked areas corresponding to total, CD44+, ALDO+ and ratio Quantification of n=58-65 organoids/condition, N=2-3, significance was calculated by Krustal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown.

Figure S2. Organoids grown for 48h often display no villus region and retain EGF-647 endocytosis in proliferating cells. (A) Brightfield time lapse imaging of developing organoid from seeded crypts in EGF medium showing standard developmental trajectories between seeding (0h), 24h, 48h, and 96h (B) Organoids were grown, with NR medium or supplemented with either EGF (50ng/mL) or EREG (50ng/mL) for 48h before fixing and staining. (C) Percentage of Organoids showing ALDO+ villus region after 48h of development, N=3. (D) Scatter plots of masked areas corresponding to total area, for all organoids (Left) and organoids with ALDO+ region showing

mean±95%CI, N=3, n=45-70 organoids/condition, (Right). Quantification of, significance was calculated by Kruskal-Wallis with Dunn's multiple comparisons test. p-values <0.05 are shown, ns non-significant. (E) Organoids were immuno-stained with CD44 (Green) AldoB (Magenta) and DAPI (Blue). Organoids ALDO- and ALDO+ are shown for each treatment. Scale Bar 100 μm. (F) Alexa Fluor-647 labelled EGF (50ng/mL) is added to organoids after 48h of growth, after 30min of incubation organoids were fixed and stained with Ki67 (Yellow), Rab7 (Magenta) and DAPI (Blue). EGF-647 is show in (Green). Representative images of Round or Crypt morphologies are represented for each treatment.

Figure S3. EGF-647 endocytosis in organoid crypts co-localizes with late endosomal marker Rab7. (A-B) Organoids were grown, with NR medium or supplemented with either EGF (50ng/mL) or EREG (50ng/mL) for 96h with a complete medium change at 48h. (A) Organoids were immuno-stained for Ki67 (Yellow) and Rab7 (Magenta), Yellow boxes denote zoom-in areas shown in the row below. (B) 50ng/mL of Alexa Fluor-647 labelled EGF (Green) is added to organoids after 96h of growth, after 30min of incubation organoids were fixed and stained. Yellow boxes denote zoom-in areas shown in the row bellow with EGF-647 (Green) and Rab7 (Magenta). (C) Digital Lightsheet Microscopy was used to see EGF-647 endocytosis in an organoid crypt grown for 96h in NR medium. Images showing Cell Mask, Hoechst and EGF-647 every 10 minutes. Red triangles show cells protruding into the lumen. For more details see *Methods* and (Video1)

Video1. EGF-647 endocytosis in NR grown organoid crypt using Digital Light Sheet imaging. Panel showing time-lapse acquisition of EGF-647 endocytosis for 52 minutes after addition of ligand. Timestamp in minutes (min) in the upper-left corner. EGF-647 (Left panel), Hoechst (Middle panel) and CellMask (Right panel).

References

Abud, H.E., W.H. Chan, and T. Jardé. 2021. Source and impact of the EGF family of ligands on intestinal stem cells. *Front. Cell Dev. Biol.* 9:685665.

Alshammari, Y., H. Ali, M.B. Assas, R. Alshammari, M. Assas, L. Koti, R. Nizam, and F. Al-Mulla. 2026. Reduced plasma epidermal growth factor levels reflect poor glycaemic status in type 2 diabetes. *Endocr. Connect.* 15.

Arteaga, C.L., and J.A. Engelman. 2014. ERBB receptors: from oncogene discovery to basic science to mechanism-based cancer therapeutics. *Cancer Cell*. 25:282–303.

Basak, O., J. Beumer, K. Wiebrands, H. Seno, A. van Oudenaarden, and H. Clevers. 2017. Induced quiescence of Lgr5+ stem cells in intestinal organoids enables differentiation of hormone-producing enteroendocrine cells. *Cell Stem Cell*. 20:177–190.

Baumdick, M., Y. Brüggemann, M. Schmick, G. Xouri, O. Sabet, L. Davis, J.W. Chin, and P.I.H. Bastiaens. 2015. EGF-dependent re-routing of vesicular recycling switches spontaneous phosphorylation suppression to EGFR signaling. *Elife*. 4:e12223.

Boretto, M., M.H. Geurts, S. Gandhi, Z. Ma, N. Staliarova, M. Celotti, S. Lim, G.-W. He, R. Millen, and E. Driehuis. 2024. Epidermal growth factor receptor (EGFR) is a target of the tumor-suppressor E3 ligase FBXW7. *Proceedings of the National Academy of Sciences*. 121:e2309902121.

Brändli, A.W., E.D. Adamson, and K. Simons. 1991. Transcytosis of epidermal growth factor. The epidermal growth factor receptor mediates uptake but not transcytosis. *Journal of Biological Chemistry*. 266:8560–8566.

Brüggemann, Y., L.S. Karajannis, A. Stanoev, W. Stallaert, and P.I.H. Bastiaens. 2021. Growth factor–dependent ErbB vesicular dynamics couple receptor signaling to spatially and functionally distinct Erk pools. *Sci. Signal.* 14:eabd9943.

Calafiore, M., Y.-Y. Fu, P. Vinci, V. Arnhold, W.Y. Chang, S.A. Jansen, A. Egorova, S. Takashima, J. Kuttiyara, and T. Ito. 2023. A tissue-intrinsic IL-33/EGF circuit promotes epithelial regeneration after intestinal injury. *Nat. Commun.* 14:5411.

Chen, C.-L., X. Yu, I.O.A. James, H. Zhang, J. Yang, A. Radulescu, Y. Zhou, and G.E. Besner. 2012. Heparin-binding EGF-like growth factor protects intestinal stem cells from injury in a rat model of necrotizing enterocolitis. *Laboratory investigation*. 92:331–344.

Childs, C.J., E.M. Holloway, C.W. Sweet, Y.-H. Tsai, A. Wu, A. Vallie, M.K. Eiken, M.M. Capeling, R.K. Zwick, and B. Palikuqi. 2023. EPIREGULIN creates a developmental niche for spatially organized human intestinal enteroids. *JCI Insight*. 8.

Deguchi, E., S. Lin, D. Hirayama, K. Matsuda, A. Tanave, K. Sumiyama, S. Tsukiji, T. Otani, M. Furuse, and A. Sorkin. 2024. Low-affinity ligands of the epidermal growth factor receptor are long-range signal transmitters in collective cell migration of epithelial cells. *Cell Rep.* 43.

Fisher, D.A., and J. Lakshmanan. 1990. Metabolism and effects of epidermal growth factor and related growth factors in mammals. *Endocr. Rev.* 11:418–442.

Freed, D.M., N.J. Bessman, A. Kiyatkin, E. Salazar-Cavazos, P.O. Byrne, J.O. Moore, C.C. Valley, K.M. Ferguson, D.J. Leahy, and D.S. Lidke. 2017. EGFR ligands differentially stabilize receptor dimers to specify signaling kinetics. *Cell*. 171:683–695.

Fujii, M., M. Matano, K. Toshimitsu, A. Takano, Y. Mikami, S. Nishikori, S. Sugimoto, and T. Sato. 2018. Human intestinal organoids maintain self-renewal capacity and cellular diversity in niche-inspired culture condition. *Cell Stem Cell*. 23:787–793.

He, G.-W., L. Lin, J. DeMartino, X. Zheng, N. Staliarova, T. Dayton, H. Begthel, W.J. van de Wetering, E. Bodewes, and J. van Zon. 2022. Optimized human intestinal organoid model reveals interleukin-22-dependency of paneth cell formation. *Cell Stem Cell*. 29:1333–1345.

Hu, H., H. Gehart, B. Artegiani, C. LÖpez-Iglesias, F. Dekkers, O. Basak, J. van Es, S.M.C. de Sousa Lopes, H. Begthel, and J. Korving. 2018. Long-term expansion of functional mouse and human hepatocytes as 3D organoids. *Cell*. 175:1591–1606.

Jardé, T., W.H. Chan, F.J. Rossello, T.K. Kahlon, M. Theocharous, T.K. Arackal, T. Flores, M. Giraud, E. Richards, and E. Chan. 2020. Mesenchymal niche-derived neuregulin-1 drives intestinal stem cell proliferation and regeneration of damaged epithelium. *Cell Stem Cell*. 27:646–662.

Jendrisek, G., D. Mesa, A. Conte, M.G. Malabarba, and S. Sigismund. 2026. Beyond the membrane: rethinking EGFR signaling in physiology and cancer. *Cellular and Molecular Life Sciences*.

Joshi, M.S., A. Stanoev, J. Huebinger, B. Soetje, V. Zorina, L. Roßmannek, K. Michel, S.A.H. Müller, and P.I.H. Bastiaens. 2023. The EGFR phosphatase RPTPγ is a redox-regulated suppressor of promigratory signaling. *EMBO J.* 42:e111806.

Langlands, A.J., A.A. Almet, P.L. Appleton, I.P. Newton, J.M. Osborne, and I.S. Näthke. 2016. Paneth cell-rich regions separated by a cluster of Lgr5+ cells initiate crypt fission in the intestinal stem cell niche. *PLoS Biol.* 14:e1002491.

Lemmetyinen, T.T., E.W. Viitala, L. Wartiovaara, T. Kaprio, J. Hagström, C. Haglund, P. Katajisto, T.C. Wang, E. Domènech-Moreno, and S. Ollila. 2023. Fibroblast-derived EGF ligand neuregulin 1 induces fetal-like reprogramming of the intestinal epithelium without supporting tumorigenic growth. *Dis. Model. Mech.* 16:dmm049692.

Mahe, M.M., E. Aihara, M.A. Schumacher, Y. Zavros, M.H. Montrose, M.A. Helmrath, T. Sato, and N.F. Shroyer. 2013. Establishment of gastrointestinal epithelial organoids. *Curr. Protoc. Mouse Biol.* 3:217–240.

Roepstorff, K., M.V. Grandal, L. Henriksen, S.L.J. Knudsen, M. Lerdrup, L. Grøvdal, B.M. Willumsen, and B. Van Deurs. 2009. Differential effects of EGFR ligands on endocytic sorting of the receptor. *Traffic*. 10:1115–1127.

Sato, T., J.H. Van Es, H.J. Snippert, D.E. Stange, R.G. Vries, M. Van Den Born, N. Barker, N.F. Shroyer, M. Van De Wetering, and H. Clevers. 2011a. Paneth cells constitute the niche for Lgr5 stem cells in intestinal crypts. *Nature*. 469:415–418.

Sato, T., D.E. Stange, M. Ferrante, R.G.J. Vries, J.H. Van Es, S. Van Den Brink, W.J. Van Houdt, A. Pronk, J. Van Gorp, and P.D. Siersema. 2011b. Long-term expansion of epithelial organoids from human colon, adenoma, adenocarcinoma, and Barrett's epithelium. *Gastroenterology*. 141:1762–1772.

Sato, T., R.G. Vries, H.J. Snippert, M. Van De Wetering, N. Barker, D.E. Stange, J.H. Van Es, A. Abo, P. Kujala, and P.J. Peters. 2009. Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche. *Nature*. 459:262–265.

Scharaw, S., M. Iskar, A. Ori, G. Boncompain, V. Laketa, I. Poser, E. Lundberg, F. Perez, M. Beck, and P. Bork. 2016. The endosomal transcriptional regulator RNF11 integrates degradation and transport of EGFR. *Journal of Cell Biology*. 215:543–558.

Schaudies, R.P., J. Grimes, D. Davis, R.K. Rao, and O. Koldovsky. 1989. EGF content in the gastrointestinal tract of rats: effect of age and fasting/feeding. *American Journal of Physiology-Gastrointestinal and Liver Physiology*. 256:G856–G861.

Schmick, M., and P.I.H. Bastiaens. 2014. The interdependence of membrane shape and cellular signal processing. *Cell*. 156:1132–1138.

Serra, D., U. Mayr, A. Boni, I. Lukonin, M. Rempfler, L. Challet Meylan, M.B. Stadler, P. Strnad, P. Papasaikas, and D. Vischi. 2019. Self-organization and symmetry breaking in intestinal organoid development. *Nature*. 569:66–72.

Shankaran, A., K. Prasad, S. Chaudhari, A. Brand, and K. Satyamoorthy. 2021. Advances in development and application of human organoids. *3 Biotech*. 11:257.

Sonntag, M.H., J. Ibach, L. Nieto, P.J. Verveer, and L. Brunsveld. 2014. Site-specific protection and dual labeling of human epidermal growth factor (hEGF) for targeting, imaging, and cargo delivery. *Chemistry–A European Journal*. 20:6019–6026.

Sprangers, J., I.C. Zaalberg, and M.M. Maurice. 2021. Organoid-based modeling of intestinal development, regeneration, and repair. *Cell Death Differ.* 28:95–107.

Stanoev, A., A. Mhamane, K.C. Schuermann, H.E. Grecco, W. Stallaert, M. Baumdick, Y. Brüggemann, M.S. Joshi, P. Roda-Navarro, and S. Fengler. 2018. Interdependence between EGFR and phosphatases spatially established by vesicular dynamics generates a growth factor sensing and responding network. *Cell Syst.* 7:295–309.

Sumigray, K.D., M. Terwilliger, and T. Lechler. 2018. Morphogenesis and compartmentalization of the intestinal crypt. *Dev. Cell*. 45:183–197.

Tallapragada, N.P., H.M. Cambra, T. Wald, S.K. Jalbert, D.M. Abraham, O.D. Klein, and A.M. Klein. 2021. Inflation-collapse dynamics drive patterning and morphogenesis in intestinal organoids. *Cell Stem Cell*. 28:1516–1532.

Tschumperlin, D.J., G. Dai, I. V Maly, T. Kikuchi, L.H. Laiho, A.K. McVittie, K.J. Haley, C.M. Lilly, P.T.C. So, and D.A. Lauffenburger. 2004. Mechanotransduction through growth-factor shedding into the extracellular space. *Nature*. 429:83–86.

van der Vaart, J., L. Bosmans, S.F. Sijbesma, K. Knoops, W.J. van de Wetering, H.G. Otten, H. Begthel, I.H.M. Borel Rinkes, J. Korving, and E.G.W.M. Lentjes. 2021. Adult mouse and human organoids derived from thyroid follicular cells and modeling of Graves' hyperthyroidism. *Proceedings of the National Academy of Sciences*. 118:e2117017118.

Wang, Y., Y. Guo, P. Wang, J. Liu, X. Zhang, Q. Liu, L. Wei, C. Xu, and J. Qin. 2025. An engineered human placental organoid microphysiological system in a vascular niche to model viral infection. *Commun. Biol.* 8:669.

Yang, L., X. Wang, X. Zhou, H. Chen, S. Song, L. Deng, Y. Yao, and X. Yin. 2025. A tunable human intestinal organoid system achieves controlled balance between self-renewal and differentiation. *Nat. Commun.* 16:315.

Yang, Y.-P., H. Ma, A. Starchenko, W.J. Huh, W. Li, F.E. Hickman, Q. Zhang, J.L. Franklin, D.P. Mortlock, and S. Fuhrmann. 2017. A chimeric Egfr protein reporter mouse reveals Egfr localization and trafficking in vivo. *Cell Rep.* 19:1257–1267.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure S1.

A

NR　EGF　EREG

Ki67

Rab7

B +EGF-647 (30min)

NR　EGF　EREG

Ki67

Rab7

EGF-647

C

| | 0 min | 10 min | 20 min | 30 min | 40 min | 50 min |

EGF-647

Hoechst

CellMask

Figure S2.

Figure S3.