Evolutionary persistence of a highly prevalent multicopy mitochondrial-derived nuclear insertion (Mega-NUMT) in Neotropical *Drosophila* flies

Authors:

Merce Montoliu-Nerin[1,4], Anton Strunov[2,5], Eleanor Heyworth[1], Daniela I. Schneider[2], Julia Thoma[2,6], Aurélie Hua-Van[3], Cécile Courret[3,7], Lisa Klasson[1], Wolfgang J. Miller[2]

Corresponding authors:

Lisa Klasson

Wolfgang J. Miller

Affiliations:

1. Molecular Evolution, Department of Cell and Molecular Biology, Science for Life Laboratory, Uppsala University, Uppsala, Sweden
2. Lab Genome Dynamics, Department Cell & Developmental Biology, Center for Anatomy and Cell Biology, Medical University of Vienna, Vienna, Austria
3. Laboratoire Évolution, Génomes, Comportement et Écologie, CNRS, IRD, Université Paris-Saclay, Gif-sur-Yvette, France
4. Tree of Life Programme, Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, Cambridgeshire, CB10 1SA, United Kingdom
5. Centre for Microbiology and Environmental Systems Science, University of Vienna, Djerassiplatz 1, 1030, Vienna, Austria
6. Department of Biosciences, Faculty of Medicine/Dentistry, Danube Private University, Förthofstraße 4, 3500 Krems, Austria
7. Institute for Integrative Biology of the Cell (I2BC), Université Paris-Saclay, CEA, CNRS, 91198, Gif-sur-Yvette, France.

Merce Montoliu-Nerin: ORCID: 0000-0002-5200-0411
Anton Strunov: ORCID: 0000-0002-3574-6481
Eleanor Heyworth: ORCID: 0000-0001-8184-9556
Daniela I. Schneider: ORCID: 0000-0002-7740-8827
Julia Thoma: ORCID: 0000-0002-4222-7380
Aurélie Hua-Van: ORCID: 0000-0001-7859-2788
Cécile Courret: ORCID: 0000-0001-5849-8014
Lisa Klasson ORCID: 0000-0002-5874-7153
Wolfgang Miller: ORCID: 0000-0002-5044-6276

Abstract

Background: Although strict maternal transmission of mitochondria is a general feature of animals and humans for ensuring homogeneity in mitochondrial DNA (mtDNA) across generations, exceptions were reported in the recent past. For example, some extremely rare but spectacular cases of heteroplasmy and paternal transmission in humans have questioned the universal evolutionary principle. Hence, as an alternative, the Mega-NUMT concept was coined to explain this discovery and was thereafter partly proven to exist. This concept expands on the quite common transfer of mtDNA fragments to the nucleus (NUMTs) by considering the existence of multicopy mitochondrial nuclear insertions. Mega-NUMT reports are currently restricted to a few cases in animals, including humans. However, even in humans, their detailed genomic organization, natural prevalence, and potential biological functions remain unclear.

Methodology/Principal Findings: Here, we discovered that up to 60 full-sized mitochondrial genomes are integrated into the nuclear genome of the neotropical fruit fly *Drosophila paulistorum* using long-read sequencing and confirmed their presence by *in situ* hybridization. The copies are organized in one cluster on chromosome 3, which we, due to its similarity with the Mega-NUMT concept, designated the "*Dpau* Mega-NUMT". Contrary to the rarity in humans, this Mega-NUMT is found at high prevalence (40%) in both long-term laboratory lines and natural *D. paulistorum* populations of different semispecies. Additionally, the mitochondrial copies in the Mega-NUMT cluster are phylogenetically separated from the current mitotypes of *D. paulistorum*. Together, these observations suggest long-term maintenance of the Mega-NUMT in nature. Hence, we propose that the *Dpau* Mega-NUMT may have been transferred to the nuclear genome before *D. paulistorum* semispecies radiation and maintained at relatively high prevalence in nature by balancing selection due to yet undetermined functions.

Conclusions/Significance: To our knowledge, this is the first verified existence and detailed dissection of a Mega-NUMT outside cats and humans. We show that Mega-NUMTs can be persistent in nature, even at high prevalence, potentially due to balancing selection. Our findings strengthen the importance of high-quality long-read sequencing technologies for deciphering complex repeat-rich genomic regions to deepen our understanding of the dynamics of genome evolution within genomic "dark matter".

Introduction

Mitochondrial DNA (mtDNA) has been widely used as a genetic marker for molecular systematics of animals for more than 30 years (Harrison, 1989). The high copy number of mtDNA per cell, as well as its rapid mutation rate, makes it particularly informative for investigating phylogenetic relationships and evolutionary history (Brown et al., 1979). Mitochondrial markers have been extensively utilized to resolve taxonomic uncertainties and to identify cryptic species (Hebert et al., 2003; Elías-Gutiérrez et al., 2008). Additionally, the use of mitochondrial data allows the reconstruction of phylogeographic patterns, providing insights into historical population dynamics and speciation events (Avise et al., 1983; Baião et al., 2023).

Studies from the pre-genomic era using Southern blots and PCR have revealed the existence of nuclear mitochondrial DNA segments (NUMTs), described as fragments of mtDNA that have become integrated into the nuclear genome (Lopez et al., 1994; Hazkani-Covo & Graur, 2007). Although the mechanisms of integration are not yet fully described, a common explanation outlines the insertion of mitochondrial sequences during the repair of double-strand chromosomal breaks (Blanchard & Schmidt, 1995, 1996; Bensasson et al., 2001). NUMTs have primarily been found in genomic regions with high levels of repetitive elements (Behura, 2007; Tsuji et al., 2012; Dayama et al., 2014; Schiavo et al., 2017; Wang et al., 2020) and distinctive GC content (Tsuji et al., 2012; Dayama et al., 2014; Schiavo et al., 2017). After their integration, NUMTs often degrade quickly, as the different genetic codes in the nuclear and mitochondrial genomes prevent proper translation of the genes (Bensasson et al., 2001). Nonetheless, some functions associated with NUMTs have been found, including the formation of new exons (Noutsos et al., 2007) and regulatory elements (Vendrami et al., 2022).

The presence of NUMTs can lead to misinterpretations when studying mitochondrial genomes (Calvignac et al., 2011), as it can result in erroneous inferences of relationships between lineages (Kress et al., 2015; DeSalle & Goldstein, 2019). Furthermore, the presence of NUMTs may lead to wrongful inference of heteroplasmy (Parr et al., 2006; Balciuniene & Balciunas, 2019), which is the coexistence of more than one mtDNA sequence variant within an individual. The misinterpretations arise from the difficulty in distinguishing NUMTs from real heteroplasmic variants when using techniques such as PCR. However, the high copy number of mtDNA compared to nuclear loci containing NUMTs may commonly suffice to mask the NUMT signal. Even so, one must be cautious (Parfait et al., 1998), as several studies have reviewed investigations in which high copy numbers of NUMTs in the genome had been interpreted as heteroplasmy (Hirano et al.,

1997; Salas et al., 2005; Yao et al., 2006; Lutz-Bonengel et al., 2021). To describe such multicopy NUMTs, the term Mega-NUMT was coined as a theoretical concept in response to a claim of paternal transmission of mitochondria in seemingly heteroplasmic humans (Balciuniene & Balciunas, 2019; Lutz-Bonengel et al., 2021). However, large multicopy NUMTs, covering approximately half of the mt-genome and with an estimated 38-76 copies, had also previously been found in the domestic cat and its close relatives (Lopez et al., 1994) and later also identified in several cat sequences using long-read sequencing, albeit in somewhat fewer copies (Patterson et al., 2023).

In this study, we focus on the Neotropical fly *Drosophila paulistorum* spp., commonly used as a study system for the evolutionary process of sympatric speciation (Burla et al., 1949; Dobzhansky & Spassky, 1959; Dobzhansky et al., 1964). The *D. paulistorum* superspecies belongs to the D. willistoni group and is composed of six semispecies, named Amazonian (AM), Andean Brazilian (AB), Centro-American (CA), Interior (IN), Orinocan (OR), and Transitional (TR), presenting different levels of pre- and post-mating incompatibilities (Dobzhansky & Spassky, 1959). Phylogenetic incongruences between the nuclear and mitochondrial DNA of *D. paulistorum* spp. have been described (Gleason et al., 1998; Robe et al., 2010; Baião et al., 2023), suggesting past introgression events. Additionally, a recent phylogenomic analysis of *D. paulistorum* spp. revealed that their mitochondrial genomes are polyphyletic and split into two major clades, α and β (Baião et al., 2023).

Here, we identify a *D. paulistorum* line from the OR semispecies, O11, with an apparently fixed presence of the two different mitotypes, α and β. However, our analysis revealed that in addition to the presence of a β mitochondrion, O11 has a NUMT consisting of around 60 copies of nearly complete mt-genomes with high similarity to the α mitotype on chromosome 3. We designate this nuclear insertion as the *Dpau* Mega-NUMT. Furthermore, we found that this Mega-NUMT is present in at least one line from three of the six described *D. paulistorum* semispecies and has a relatively high prevalence of around 40% in fly populations from French Guiana. Taken together, these data suggest that the *Dpau* Mega-NUMT was present before or just after the split of the semispecies and may be maintained in nature by balancing selection.

Results

The two mitotypes, α and β, are fixed in individual flies of the *D. paulistorum* line O11

We sequenced PCR products of the mitochondrial *Cytochrome Oxidase I* (*COI*) gene from several *D. paulistorum* lines for mt-barcoding. We found multiple robust double peaks in the resulting chromatograms derived from single flies of the OR line O11, suggesting the presence of both α and β mitotypes within the same individual. The presence of a *Hin*dIII restriction site for the α-mitotype but not ß (**Fig. 1A**) was used to perform diagnostic RFLP/*Hin*dIII Southern blots on DNA from single O11 flies (Junakovic, 2004) probed with the *COI* cloned plasmid (**Fig. 1B** and **Table S1, Fig. S1A**). We observed that the O11 line again showed a clear heteroplasmic pattern, with both α and β mitotypes represented in individual flies at approximately similar abundances, suggesting almost equal copy numbers (**Fig. 1B**). In contrast, single flies of line C2 belonging to the CA semispecies and line A28 from the AM semispecies were homoplasmic and carried either the α or the β mitotype, respectively (**Fig. 1B**). Furthermore, we performed *COI*-specific RFLP/*Hin*dIII-PCR on multiple single females and males (**Table S1, Fig. S1**) which showed a consistency on the presence of both α and β mitotypes in all tested O11 flies (**Fig. 1C**), which suggested fixation in this long-term lab line collected in the 1960s.

Figure 1. Presence of two mitotypes in the *D. paulistorum* Orinocan line O11. (A) Chromatogram derived from *COI* direct Sanger sequencing of one O11 female covering the α-diagnostic *Hin*dIII restriction

site (underlined). The asterisk highlights the double peak with Guanine (G) for the α- and Adenine (A) for

the β-mitotype. (**B**) Single-fly *CO1* Southern blot digested with *Hin*dIII and probed with cloned *CO1*. While

the Amazonian females of the A28 line (n=3) present only the β-mitotype (4.8kb) and the females of the

Centro-American line C2 (n=2) only the α-mitotype (2.4kb), all randomly picked Orinocan O11 females

(n=3) present both the α- and β-mitotypes (4.8 and 2.4kb). (**C**) α-diagnostic *CO1* RFLP/*Hin*dIII-PCR screen

(**Fig. S1**) on O11 single flies (n=20 for each sex). Undigested (U) and digested (D) CO1-PCR fragments per

fly are shown exemplary for two individual specimens (magnification at the bottom right in C). All tested

single flies (40/40) exhibited clear double bands after *Hin*dIII digestion.

Subsequently, we measured the copy numbers of both α and β mitochondrial (mt) genomes, relative

to the nuclear genome, using discriminant qPCR on the *CO3* gene (see Materials and Methods).

The quantification of both the α and β mt genomes was in multiple tissues of both females and

males and resulted in similar average values of 27.2±13.9 α and 31.6±22.7 β mt genome copies per

nuclear genome (**Fig. 2**). The β mt-genome copy numbers were strongly affected by tissue (**Table**

S2; F-value=27.694, p<0.001, F-test, **Fig. 2A**, **Table S2**) and a more subtly, yet significantly, by

age and sex combined (**Table S2**; F-value=6.66, p<0.05, F-test), and by sex alone (**Table S2**; F-

value=4.714, p<0.05, F-test), as expected based on mitochondrial function. In contrast, the number

of copies of the α mt-genome was not significantly affected by any of the tested parameters (**Fig.**

2B, **Table S2**), suggesting that the α mitotype is not a typical mitochondrion.

Figure 2. **qPCR quantification of α and β mitotypes in *D. paulistorum* O11 flies**. 120 flies were analyzed

to assess the number of α and β mitotype copies relative to the nuclear genome copies using discriminant

qPCR. Boxplots show the distribution of values, mean, and standard deviation for the (**A**) β mitotype and

the (**B**) α mitotype.

Discovery of a Mega-NUMT with long reads

Given the seemingly fixed high abundance of two mitotypes in one individuum and unresponsive behaviour of the α mitotype in different tissues of the *D. paulistorum* O11 line, we sequenced the O11 genome using Oxford Nanopore technology (ONT). The assembly of the nuclear genome of O11 was 231 Mb (111 contigs, N50 = 18.6 Mb) (**Table S3**), with a BUSCO completeness of 99% (diptera_odb, **Table S4**). Using the mitochondrial genome from the O11 line (β mitotype, Baião et al., 2023), we searched the nuclear assembly and found 12 mt-genome copies clustered at the end of one contig (ctg000940). From here on, we refer to this cluster as the *Dpau* Mega-NUMT, given its similarity to the theoretical Mega-NUMT concept. Further analysis of the 12 NUMT copies revealed a conserved synteny of the mitochondrial genes across most copies (**Fig. 3**). Four of the NUMT copies have lost genes compared to the mt-genome, and in all NUMT copies, many of the genes have suffered frameshift mutations (**Fig. 3**, **Table S5**). When comparing the gene length to the fraction of intact gene copies per protein-coding gene, we found a significant inverse correlation, suggesting that the pseudogenization process is random (**Fig. S2**). However, given the complexity of the Mega-NUMT, it is possible that the sequence of each gene copy in the assembly is not 100% accurate. Nevertheless, at least one copy of each of the 13 mitochondrial protein-coding genes is intact in the Mega-NUMT, albeit with the mitochondrial genetic code (**Table S5**).

A phylogeny based on the complete set of genes and pseudogenes in the *Dpau* Mega-NUMT, together with the published mitochondrial genomes of *D. paulistorum* spp. and other willistoni group species (Baião et al., 2023), shows that all individual NUMT copies in the Mega-NUMT are found in one clade, which is sister to the α mitotype of the *D. paulistorum* C2 and MS lines (**Fig. 3**) belonging to the CA and AB semispecies, respectively (**Table 1, Table S6**). Hence, we refer to these NUMT copies as α-NUMTs. Importantly, though, the separation of the α mitotype and α-NUMTs into different clades, and the monophyly of within these clades, suggests that none of the α-NUMT copies in the *Dpau* Mega-NUMT are recent insertions from an extant α mt-genome. This is further supported by the fact that the O11 line carries the β mitotype, as does the majority of *D. paulistorum* semispecies and lines.

Figure 3. Phylogenetic position and synteny of NUMT copies of *D. paulistorum* O11 and closely related mitochondrial genomes. Maximum likelihood phylogeny based on the complete set of genes and pseudogenes of the NUMT copies found in the Mega-NUMT of *D. paulistorum* O11, together with the mitochondrial genes in *D. paulistorum* and closely related species. Black dots indicate nodes with 100 bootstrap supports. The α and β mitotype clades are annotated on the tree. Mitochondrial genomes of *D. paulistorum* O11 (green) and *D. paulistorum* MS (blue) were used as references for the β and α mitotypes. The grey ribbons between sequences represent homologous regions of the same direction, while red ribbons represent inversions. The 13 mitochondrial protein-coding genes are shown as colored arrows, and rRNA and tRNA genes as blocks. Intact genes are filled, and pseudogenes only have a colored outline.

Further, we analyzed the genomic region where the *Dpau* Mega-NUMT is located. The Mega-NUMT is directly flanked by an AT-rich 36 bp satellite sequence (**Fig. 4** and **Fig. S3**) and the gene

FOXO encoding the Forkhead box O transcription factor (Jünger et al., 2003), located in Muller

element E on chromosome 3, as deduced by comparison to *D. melanogaster*. The same 36 bp

satellite DNA is also found inside the Mega-NUMT, downstream of α-NUMT 3 (**Fig. 4** and **Fig.**

S3). The positions of *FOXO* and the Mega-NUMT next to each other were also confirmed by DNA-

hybridizations on polytene chromosomes of larval salivary glands of O11, and the bright

polytenized signal of the Mega-NUMT (**Fig. S4**) implies its euchromatic state since it is

polytenized and in close genomic vicinity to the essential host gene *FOXO*.

Figure 4. Graphic representation of the end of contig000940 in the assembly of *D. paulistorum* O11.

The region is divided into three rows to facilitate visualization. The position of the nuclear gene *FOXO* and

each α-NUMT is represented with a white box, which also contains colored lines that represent the position

of each protein-coding and RNA gene. The positions of the satellite and repeat elements are shown below

in the order: The AT-rich 36-bp satellite, *Gypsy*, *TcMariner*, *Helitron*, *Copia*-like transposons, and simple

repeats.

All α-NUMT copies except α-NUMT 3 have the same orientation in the *Dpau* Mega-NUMT (**Fig.**

4). The sequence of α-NUMT 3 also contains a rearrangement not seen in any of the other copies

of the contig (**Fig. 3** and **4**). Additionally, the 12 α-NUMT copies in the O11 nuclear genome

assembly are surrounded by repeat elements, and in some cases, insertions of transposable elements

are present within an α-NUMT sequence (**Fig. 4**). Specifically, we found remnants of the LTR

retrotransposon *Gypsy* within the α-NUMT copies 6, 9, and 11 and of the LTR element *Copia* in α-NUMT copies 5 and 9 (**Fig. 4**). In addition, we found short regions with sequence similarity to the DNA transposon Tc Mariner in six of the α-NUMTs and to the rolling-circle DNA transposable element, *Helitron*, in nine of the α-NUMTs, as well as between α-NUMT copies when a non-NUMT sequence was found between them (**Fig. 4**).

Number of α-NUMT copies in the genome

Since the discriminant qPCR results indicated that more than 12 α-NUMT copies could be present, we investigated the coverage of ONT reads in the Mega-NUMT region of O11 to assess the true number of α-NUMT copies. When mapping the reads to the assembly, we also included the β mt-genome from Baião et al., (2023) and the genome of the bacterial symbiont *Wolbachia* of O11 from Papachristos et al., (2025) to prevent mapping of reads from the real mitochondrion or *Wolbachia* to the Mega-NUMT. Additionally, to be as stringent as possible in our assessment, only one mapping per read was allowed. The read coverage over the *Dpau* Mega-NUMT shows a clear discordance with the average coverage across the genome (**Fig. 5**). The *Dpau* Mega-NUMT has an average coverage of 402±159, while the genome assembly as a whole has an average coverage of 80.8±77.4 (**Fig. 5**). Using these numbers, we thus infer that the genome contains around 60 whole mt-genome α-NUMTs (**Fig. 5**). We note that the coverage over the different α-NUMT copies varies substantially. Especially high coverage was found on α-NUMT 8, with 17 times higher coverage than the genome average, followed by α-NUMTs 7, 9, and 10, with ten, seven-, and six-times higher coverage than the genome average, respectively. These results indicate that the unassembled α-NUMT copies resemble α-NUMT copies 7-10 the most. Specifically, we note that both α-NUMT 7 and 8 contain the full mt-genome and have no intervening sequence, suggesting that many of the unresolved copies are also full mt-genomes with no intervening sequences (**Figs. 4** and **5**).

We note that the 60 α-NUMT copies estimated from long read mapping are more than the ca. 30 we estimated by our qPCR results. As α-NUMT 3 lacks *CO3* completely, only 11 of the 12 copies in our assembly have a binding site for the primer CO3_SNP_1 used in qPCR for determining the copy numbers (**Fig. 2**, **Table S1**). Hence, although we do not know exactly what the remaining copies look like, this indicates that the discrepancy in total copy numbers of the α-NUMT between the coverage and qPCR results might be explained by a lack of primer binding in some of the α-NUMT copies.

Figure 5. Coverage quantification of α-NUMTs in the nuclear genome of *D. paulistorum* O11. The x-axis represents the end region of contig000940 of the genome assembly of *D. paulistorum* O11, which includes the nuclear gene *FOXO*, and the 12 assembled α-NUMTs. Coverage was obtained from mapping Nanopore reads to the genome assembly. The average and standard deviation of the coverage over each α-NUMT and the *FOXO* gene are plotted in the same color as used for the box depicting each feature. The total number of α-NUMT copies in the Mega-NUMT is estimated from the average over all the assembled α-NUMT copies in relation to the average coverage of the whole genome assembly.

Mega-NUMT visualization on *D. paulistorum* chromosome.

To validate the genomic findings of the Mega-NUMT localization on the 3rd chromosome of *D. paulistorum* O11, we performed DNA fluorescent *in situ* hybridization (FISH) on mitotic chromosomes of larval brains using mitochondria-specific probes and a probe for the Mega-NUMT-linked satellite (see **Fig. 4** and the Materials and Methods section). In O11, we observed two satellite signals (red) on the 3rd chromosome—one in the (peri)centromeric region and a second in the subtelomeric region—as well as a Mega-NUMT signal (green) adjacent to the subtelomeric satellite signal (**Fig. 6A**), in agreement with the assembly data (**Fig. 4**). In contrast, the *D. paulistorum* line FG295 (**Table S6**), which also belongs to the OR semispecies (**Table 1**), shows only a single satellite signal in the (peri)centromeric region of the third chromosome, but no Mega-NUMT signal (**Figure 6B**).

Figure 6. DNA-FISH on mitotic chromosomes from larval brains of the *D. paulistorum* O11 and FG295 lines. Probes targeting the Mega-NUMT (green) and the Mega-NUMT–linked satellite (red) were used. The scale bar represents 5 μm. The inset highlights the third chromosome, showing two satellite signals and one Mega-NUMT signal in O11 (**A**), but only a single satellite signal in FG295 (**B**). The schematic on the right summarizes the localization of the satellite (red) and the Mega-NUMT (green) in each strain.

Additionally, to follow Mega-NUMT dynamics in interphase and dividing cells, we investigated the localization of the *Dpau* Mega-NUMT in early embryonic mitosis by mt-DNA FISH (**Fig. S5**). Contrary to larval brains that were generated by squash fixation and thereby washing off the cytoplasm (**Fig. 6**), in our whole mount confocal mt-DNA-FISH assays on O11 embryos, we observed staining of mitochondria in the cytoplasm and the chromosome-associated Mega-NUMT signal as two dots, with some cases of a fused, larger dot (arrowheads and arrow, respectively, **Fig. S5A-D, F**). Each dot consists of two parts with a small bridge in between,

suggesting somatic pairing (inset of **Fig. S5B**). Based on our results on metaphase chromosomes from 3rd instar larvae (**Fig. 6**), we conclude that each dot represents a Mega-NUMT signal located on one of the duplicated arms of chromosome 3, with two smaller parts representing sister chromatids. At the anaphase stage, we could clearly observe the separation of sister chromatid signals, creating a 4-dot pattern (**Fig. S5E**). Altogether, systematic DNA-FISH at different stages of early embryo mitosis reconfirmed the presence of only one Mega-NUMT signal on chromosome 3 of the O11 line.

Prevalence of the Mega-NUMT across *D. paulistorum* semispecies and populations

To test if the *Dpau* Mega-NUMT is restricted to only the O11 line, we analyzed α-NUMT prevalence across different *D. paulistorum* lines. We used RFLP-PCR from adult flies to assess the prevalence and abundance of α-NUMTs (**Table 1**). In addition to O11, another OR line, FG111, presented α-NUMT, but it appeared not to be fixed in the population, as it was only found in 10 out of 40 flies that originated from an isofemale (**Table 1**). In yet another OR line, FG103, the Mega-NUMT was lost between the first test (2014-15) and the last one (2018) (**Table 1**). Two AM lines, FG16 and FG572, also presented a non-fixed Mega-NUMT (**Table 1**), while two lines from the AB semispecies, MS and Yellow, both carried the Mega-NUMT in all tested flies (**Table 1**). To sum up, we found the Mega-NUMT in several OR lines, as well as in the AM and AB semispecies, but not in the few representatives of the other three semispecies of *D. paulistorum* (CA, IN and TR). Possibly, this is caused by a sampling bias, since none of these three semispecies were found in our collections from French Guiana between 2014 and 2018 (**Table 1** and **Table S7**). In addition to O11, we only found fixation of the *Dpau* Mega-NUMT in the old long-term stocks of MS and Yellow that were maintained at small population sizes in the laboratory since the 1960s or earlier. Our finding that the Mega-NUMT is not fixed in isofemale lines from our recent collections in French Guiana suggests that it is frequently hemizygous in wild populations.

Table 1. α-NUMT abundance and prevalence in *D. paulistorum* lines based on *CO1* RFLP-PCR from adult flies. Abundance is given in percentage of α-NUMTs compared to the mitochondrial copies per fly (**Fig S1**) and is presented separately for females (left) and males (right). Prevalence of the Mega-NUMT among tested individuals is presented both as a percentage and in absolute numbers (in parentheses). Abbreviations: Spp: Semispecies, AB: Andean-Brazilian, AM: Amazonian, CA: Centro-American, IN: Interior, OR: Orinocan, TR: Transitional, Coll: Collection year, ND: Not determined. Further details about the fly lines are provided in **Table S6**.

Line	Coll.	Spp.	Mitotype	α-NUMT detected	α-NUMT abundance (%)[#]	α-NUMT prevalence
O11	1957	OR	ß	Yes	30.7±3.7/36.8±3.6	100% (39/39)
FG103	2014	OR	ß	Yes	ND[§]	0% (0/40)[§]
FG111	2014	OR	ß	Yes	13.4±1.1/18.3±2.9	25% (10/40)
FG295	2014	OR	ß	-	0	0% (0/40)
TP37	2009	OR	ß	-	0	0% (0/20)
POA1	2003	OR	ß	-	0	0% (0/20)
RP	1995	OR	ß	-	0	0% (0/20)
A28	1952	AM	ß	-	0	0% (0/40)
FG16	2018	AM	ß	Yes	57.2±8.6/66.5±7.2	55% (22/40)
FG18	2018	AM	ß	-	0	0% (0/40)
FG572	2015	AM	ß	Yes	45.3±8.1/63.7±7.0	90% (36/40)
MS[*]	1962	AB	α	Yes	21.8±3.1/36.0±4.1	100% (40/40)
Yellow[*]	1960	AB	α	Yes	20.4±4.7/27.4±4.3	100% (40/40)
C2	1954	CA	α	-	0	0% (0/20)
White	1960	CA	α	-	0	0% (0/20)
L1	1958	IN	ß	-	0	0% (0/20)
SM	1956	TR	ß	-	0	0% (0/20)

[#]Abundance of the α-NUMT is given in percentage of females and males determined by *CO1* RFLP-PCR.

[§]The line FG103 presented α-NUMT in 2014/15 (determined by Sanger and Illumina sequencing) but has lost it afterwards (verified by *CO1* RFLP-PCR and Sanger sequencing in 2018/19).

[*]To distinguish αNUMTs from the α-mitotype of AB semispecies, the α-diagnostic *Eco*RV site of *CO1* was used.

To evaluate the prevalence of the *Dpau* Mega-NUMT in nature, we screened a total of 80 *D. paulistorum* spp. isofemale lines at generation F1 post-collection that were sampled between 2014 and 2024 from five different locations in French Guiana (**Fig. 7**; **Table S7**).

Fig. 7. Natural prevalence of Mega-NUMTs in *D. paulistorum* spp. populations in French Guiana. All 80 fly samples were collected between 2014 and 2024 at five different locations (red dots, and **Table S7**). The individual pie charts are in reference to locality and collection date and show the actual number of Mega-NUMT-positive (green) and negative (gray) specimens. Species designation and Mega-NUMT-typing per established *D. paulistorum* spp isoline (n=80) was performed on multi-fly DNA at generation F1 post-collection by *CO1* PCR and direct sequencing. The presence of robust double peaks in their chromatograms at α-NUMT-diagnostic positions was used in combination with cloning and sequencing.

Sampled flies were established as isofemale lines and barcoded at F1 post-collection as members of the *D. paulistorum* species complex by *CO1* PCR followed by direct Sanger sequencing and/or cloning of *CO1* (**Table S7**). Similar to O11 (**Fig 1A**), the presence of robust double peaks in the chromatograms from direct sequencing was used for Mega-NUMT typing, suggesting that 34 of the 80 isofemale lines harbor α-NUMTs (**Table S7**). The high prevalence of α-NUMT of 43% in our recent collections from *D. paulistorum* populations, together with its presence in at least three of the six semispecies (**Table 1**), strongly suggests their ancestral evolutionary state in this neotropical species complex and its stable maintenance both in nature and over long evolutionary time.

Discussion

In animal species, mtDNA is normally transmitted strictly uniparentally from mother to progeny (Ladoukakis & Zouros, 2017), with rare exceptions (Breton et al., 2007). However, reports of mitochondrial heteroplasmy and biparental transmission in humans challenged this view (Schwartz & Vissing, 2002; Luo et al., 2018) and prompted skepticism by the scientific community (McWilliams & Suomalainen, 2019; Salas et al., 2020). Consequently, the theoretical Mega-NUMT concept, proposing nuclear integration of multiple copies of whole mitochondrial genomes, was coined as an alternative explanation for such unorthodox findings (Balciuniene & Balciunas, 2019). Although this concept has been verified in humans (Wei et al., 2020; Lutz-Bonengel et al., 2021; Pagnamenta et al., 2021; Wei et al., 2022), similar phenomena have so far only been described in a limited number of animal systems where only partial regions of the mt-genome were found at multicopy (Lopez et al., 1994; Patterson et al., 2023; Biró et al., 2024), but not in Drosophila (Rogers & Griffiths-Jones, 2012), despite a relatively high prevalence of NUMTs in insects (Hebert et al., 2023). On the other hand, nuclear insertions of whole bacterial genomes larger than 1 Mb, like the α-proteobacterium *Wolbachia*, have been repeatedly detected in invertebrate genomes, including insects and nematodes (Kondo et al., 2002; Hotopp et al., 2007; Nikoh et al., 2008; Klasson et al., 2014; and recently reviewed in Keeling, 2024).

The Mega-NUMT of *D. paulistorum* spp.

In this study, we found that multiple whole mitochondrial genomes are integrated in the nuclear genome of the O11 line from the *D. paulistorum* species complex, designated α-NUMTs because of their close phylogenetic relationship with the cytoplasmic α-mitotype of the Centro-American and Andean-Brazilian semispecies of *D. paulistorum* (Baião et al., 2023). These α-NUMTs localize in a nuclear cluster in the Müller element E on chromosome 3 with up to 60 nearly complete 15kb mitochondrial genomes, potentially with a total size of 900 kb. Given the agreement between this finding and the Mega-NUMT concept, we named this cluster the "*Dpau* Mega-NUMT". Similar to earlier NUMT reports (Behura, 2007; Tsuji et al., 2012; Dayama et al., 2014; Schiavo et al., 2017; Wang et al., 2020; Biró et al., 2024), the *Dpau* Mega-NUMT of the O11 line is located in a genomic region containing other repetitive elements, *i.e.,* a chromosome 3-specific satellite and several different TEs (**Fig. 6A, B**), which suggests it might have served as a "safe haven" for such sequences (Werren, 2011). Such repetitive sequences may have enlarged the *Dpau* Mega-NUMT beyond 1 Mb in size, which is larger than the 641 kb NUMT recently discovered in the genome of

Arabidopsis thaliana, consisting of a partially duplicated large mitochondrial genome (Fields et
al., 2022) and currently one of the largest described fully sequenced NUMTs. Additionally,
contrary to currently described human Mega-NUMTs with prevalence below 0,5% (Wei et al.,
2020; Bai et al., 2021; Lutz-Bonengel et al., 2021), the *Dpau* Mega-NUMT is found at high
prevalence, around 40% in long-term laboratory lines (3/8; **Table 1**) and 42.5% (34/80) in *D.*
paulistorum spp. flies collected between 2014 and 2024 at different locations in French Guiana
(**Fig. 7**; **Table S7**). Furthermore, the *Dpau* Mega-NUMT is present in members of at least three of
the six semispecies of *D. paulistorum*, either in homo- or hemizygosity (**Table 1**). Finally, the α-
NUMTs form a separate monophyletic clade to the exclusion of the current α mitotype (**Fig. 3**),
indicating that the insertion did not involve any of the extant mitotypes but possibly a more
ancestral one. Therefore, we proposed that the *Dpau* Mega-NUMT might have been inserted in the
nuclear genome before or just after the semispecies radiation, estimated to have occurred between
1.2-2.1 MYs ago (Zanini et al., 2018; Kim et al., 2022).

Integration and chromosomal position
The overall well-conserved head-to-tail structure of the *D. paulistorum* α-NUMT cluster suggests
that the *Dpau* Mega-NUMT might have integrated as a concatemer of multiple mt-genomes. As
earlier proposed by Balciuniene & Balciunas, 2019, the mechanism of generating Mega-NUMTs
of complete mt-genomes may be explained by replication via a rolling circle intermediate. Such
mt-rolling circles have, for example, been reported in yeast, nematodes, and humans (Pohjoismäki
et al., 2009; Ling et al., 2016; Ling & Yoshida, 2020), but not in Drosophila until now. The
switching between conventional circular to rolling circle replication might be quite feasible because
key components of the mitochondrial transcription and replication apparatus are derived from the
T-odd lineage of bacteriophage rather than from an α-Proteobacterium, as the endosymbiont
hypothesis would predict (Filée et al., 2002; Shutt & Gray, 2006). In addition, rolling circle
replication (RCR) has been described in Drosophila for *histone* and the *Stellate* and *Suppressor of*
Stellate (Su(Ste)) genes (Cohen et al., 2005; Cohen & Segal, 2009). Moreover, *Helitron* transposons
like *DINE-1* elements that can reach extremely high copy numbers in *D. melanogaster* (Thomas et
al., 2014) also propagate via RCR (reviewed in Barro-Trastoy & Köhler, 2024).

420   The nuclear integration site of the mt-concatemer in the chromosome 3 of *D. paulistorum*
spp. is interesting, since it is a fusion between Muller elements E and F, orthologous to
chromosome arm 3R and chromosome 4 of the melanogaster group, respectively (Muller, 1940;
Spassky & Dobzhansky, 1950). The location of the *Dpau* Mega-NUMT in the O11 line next to the

satellite DNA (**Fig 6A**), also present in the acrocenter of the fusion autosome E/F (**Fig. 6B**),

potentially suggests a chromosomal inversion event splitting the satellite in two, but not affecting

the upstream gene *FOXO*. Such paracentric inversions are quite commonly found in the *D.*

paulistorum species complex (Kastritsis, 1967) and can even play adaptive roles (Krimbas &

Powell, 1992), like for the successful recent colonization of urban environments (Valiati & Valente,

1997). In addition, inversions are enriched on chromosome 3 and often maintained in

heterozygosity, potentially under balancing selection (Gromko & Richmond, 1978). These earlier

findings suggest that chromosome 3 of *D. paulistorum* spp. may be an evolutionary hotspot for

chromosomal rearrangements like inversions and translocations, but also mitochondrial-derived

insertions like the *Dpau* Mega-NUMT as reported here.

Why is the *Dpau* Mega-NUMT maintained at high prevalence and over evolutionary time?

Contrary to currently described human Mega-NUMTs, which are extremely rare in prevalence, the

Dpau Mega-NUMT appears to be evolutionarily persistent since it is present in lines from at least

three different *D. paulistorum* semispecies, *i.e.*, Amazonian, Andean Brazilian and Orinocan.

Additionally, it is present at a relatively high prevalence of potentially more than 40% in natural

populations (**Figure 7**; **Table S7**). Even so, the fact that the Mega-NUMT is not fixed in *D.*

paulistorum except for old laboratory stocks (**Fig. 1**; **Table 1**), and that natural populations are

mainly hemizygous, suggests balancing selection rather than an essential function.

Although almost half of the mitochondrial genes in Mega-NUMT present in the 12 resolved

α-NUMT copies in our assembly (**Table S5**) still appear to encode intact ORFs (albeit with the mt

genetic code), we believe that they are unlikely to be under selection for having the same function

as when they were present in the mt-genome. Mainly because the mt genetic code will cause

problems during translation, but also since the pseudogenizing mutations appear to have occurred

randomly throughout the genes (**Fig. S2**), suggesting neutrality. However, we can't exclude that

some of the DNA in the Mega-NUMT might be transcribed and serve yet undetermined functions.

Potentially, the TEs within and between α-NUMT units, especially remnants of LTR

retrotransposons like *Copia* and *Gypsy* (**Fig. 4**), may contribute to expression, as LTRs are well

known to harbor cell- and tissue-specific *cis*-regulatory elements that affect expression of

neighboring host genes (McDonald et al., 1997). Alternatively, TEs can also create *de novo*

regulatory regions acting *in trans*, so-called piRNA clusters (recently reviewed in Pritam and

Signor, 2025, which, if expressed in the antisense orientation of the transposon, can serve as

efficient epigenetic silencers of homologous TEs via the piRNA pathway (Brennecke et al., 2008).

Indeed, *de novo* creation of such a repressive piRNA cluster has recently been demonstrated in *D.*

melanogaster (Rafanel et al., 2025).

Another possibility is that the Mega-NUMT provides a structural function. Other studies

have seen that NUMTs are located close to centromeric or telomeric regions (Puertas & González-

Sánchez, 2020), suggesting that they might be involved in their function. However, the *Dpau*

Mega-NUMT of O11 is not found in either the centromeric or telomeric chromosomal region of

chromosome 3, according to our FISH. Even so, a yet undetermined structural role for the *Dpau*

Mega-NUMT is still possible.

Further experiments will be necessary to elucidate any potential function and fitness

benefits of the Mega-NUMT in the *D. paulistorum* species complex.

Conclusions

To our knowledge, this is the first verified existence and detailed dissection of a Mega-NUMT

outside cats and humans. Moreover, we show that Mega-NUMTs can persist at high prevalence in

nature and over relatively long time periods, suggesting balancing selection. Our findings

strengthen the importance of high-quality long-read sequencing technologies for deciphering

complex repeat-rich genomic regions and mobile DNAs to deepen our understanding of genomic

"dark matter". Finally, it is expected that with the rapidly increasing number of high-quality

genomes, the *Dpau* Mega-NUMTs will not remain a unicorn. Instead, similar Mega-NUMTs will

very likely be found in further eukaryotic genomes.

Data availability

478 All data will be available upon publication.

Acknowledgements

We thank Tom Martin for performing the DNA extraction for Nanopore sequencing of *D.*

paulistorum O11. ONT sequencing of O11 was performed by the Uppsala Genome Center (UGC)

in Uppsala, Sweden. The facility is part of the National Genomics Infrastructure (NGI) Sweden

and the Science for Life Laboratory. Some of the data handling was enabled by resources provided

by the Swedish National Infrastructure for Computing (SNIC) at UPPMAX. UGC and UPPMAX

are supported by the Swedish Research Council. This work was supported by the Swedish Research

Council VR grant 2014–4353 to LK and by the Austrian Science Fund FWF grants P28255B22,

FW613A0501 and FW613A0502 to WJM. We also thank the Nouragues research field station

(managed by CNRS), which benefits from "Investissement d'Avenir" grants managed by Agence

Nationale de la Recherhe (AnaEE France ANR-11-INBS-0001; Labex CEBA ANR-10-LABX-25-

01) to WJM.

Contributions

494   LK and WJM conceived and designed the study. MMN performed assemblies, annotations and

495   phylogenetic analyses. EH performed qPCRs. JT and DIS performed line screening and RFLP

496   analyses. AS and CC performed FISH. AH-V and WJM collected flies. LK performed

497   bioinformatic analyses. MMN, WJM and LK wrote the paper with contributions from CC and AS.

498   All authors read and approved the manuscript.

Material and Methods

Fly strains

502   All fly lines used in the present study are listed in **Table S6**, including their collection date and

503   semispecies designation. All flies were kept at 25±1°C on Formula 4-24 Drosophila instant food

504   (Carolina, USA) with a 12 hrs. light-dark cycle.

Drosophila collections from French Guiana 2014-2024

Field samples of neotropical Drosophila specimens were collected in French Guiana at five

different geographic locations between 2014 and 2024 (**Table S7**) by trapping live flies with

banana/yeast baits in plastic PET bottles over a 48 hrs. time period. Taxonomic grouping - at least

at the species-group level - was performed by eye under the stereo loupe by using FlyNap. Trapped

inseminated single females were isolated into individual small single fly culture vials supplemented

with Carolina Instant Fly-Food, shipped back to the Medical University of Vienna to establish

consecutive isofemale lines in WM´s lab. Upon egg deposition and the emergence of F1 larvae,

DNA was isolated from the G0 mother for single fly DNA extraction with the Gentra Puregene

Tissue kit (Qiagen, Germany) and mitochondrial *CO1*-barcoding. PCR was performed with the

primer pair CO1-F and CO1-univR (**Table S1**), followed by direct Sanger sequencing and cloning

(for further details, see Madi-Ravazzi et al., 2021).

This study was conducted under ABS Permits No. ABSCH-IRCC-FR-245930-1 and ABSCH-IRCC-FR-257164-1 issued by the French Ministry of Ecological Transition on 25 Jun 2018 and 16 Sep 2021 respectively, in compliance with the Nagoya Protocol and French Biodiversity Law (2016).

Single fly Southern blots

Single fly Southern blots were performed following the protocol by Junakovic, 2004. In brief, DNA was extracted from single individuals, digested with the restriction enzyme *Hin*dIII and processed for electrophoresis on a vertical 0.8% agarose gel. After vacuum-blotting onto a positively charged nylon membrane, fragments were hybridized with a P[32]-labelled mitochondrial *CO1* probe derived from a plasmid. Detection was performed by exposing the membranes to X-ray films at -80°C.

Restriction Fragment Length Polymorphism-PCR (RFLP-PCR)

Total DNA was extracted from single or multiple flies using the Gentra Puregene Tissue kit (Qiagen, Germany). Flies were homogenized using a TissueLyser LT (Qiagen, Germany) at 50 Hz for 50 sec and DNA was consequently isolated following the manufacturer's protocol. DNA concentration was determined with a NanoDrop OneC Spectrophotometer (Thermo Scientific, USA). RFLP-PCR for the mitochondrial Cytochrome c oxidase genes *CO1* and *CO2* (**Figure S1**) was set up in 20 µl reactions containing 1x Promega reaction buffer, 2.5 mM $MgCl_2$, 0.5 µM of each primer (**Table S1**), 200 µM of each dNTP, and 0.025 U of GoTaq G2 DNA Polymerase. *CO1*/*CO2* reactions were performed using the following thermal profile: initial denaturation for 2 min at 95°C followed by 30 cycles consisting of 45 sec denaturation at 95°C, 45 sec annealing at the appropriate primer conditions T_m°C (**Table S1**) and 30 sec extension at 72°C. Final extension was carried out at 72°C for 10 min. PCR products were consequently used for RFLP (digest plus undigested control) using the restriction enzymes *Hin*dIII and *Eco*RV-HF for *CO1*, and *Msp*I for *CO2* (**Figure S1**). Reactions were performed in 20 µl volume containing 10 or 20 U of enzyme, 1x reaction buffer and 10 µl PCR product. After overnight incubation at 37°C, enzymes were inactivated by either heat (80°C for 20 min) or by adding EDTA-containing gel loading dye. 10 µl of undigested and digested samples were then run on a 1.5% agarose gel and post-stained with 10 mg/ml ethidium bromide. Imaging was performed using a Molecular Imager ChemiDocTM XRS Imaging System and the intensities of obtained PCR bands were quantified using Image Lab 5.2.1 software (Bio-Rad).

Quantifying Mega-NUMT abundance within a fly

Undigested samples representing the sum of both mitotypes were set as a reference band with

relative quantity 1. The relative levels of mitotypes were then quantified by summing up the two

digested fragments plus the undigested fragment. Due to technical differences in band intensity,

however, the sum of all fragments did not result in a relative quantity of 1. Hence, the actual sum

value was set to 100% to determine the percentage of mitotype levels. Smaller fragments with a

length up to approximately 100 bp were not detectable after ethidium bromide staining and we

hence used the correction factor Cf = undigested [bp] / digested [bp] to estimate their relative

quantities.

Quantitative PCR (qPCR)

DNA was extracted by homogenizing whole flies in 200µl Chelex solution at 5% concentration in

distilled water. A total of 10µl of proteinase K (10mg/ml) was added per sample and the vials were

incubated for 6 hours at 56°C. After digestion, vials were centrifuged at 16,200 x g for 3 minutes,

and the supernatant was transferred to a new vial. DNA was stored at 4°C and freeze-thaw cycles

were avoided when possible. Samples were run in duplicate using SYBR Green reagent on a

StepOnePlus Real Time PCR machine (Applied Biosystems, USA). Technical repeats with >1 Ct

difference were rerun or discarded. Each reaction contained 10µl Maxima SYBR Green/ROX

qPCR Master Mix (2x), 1µl of forward primer, 1µl of reverse primer, 6µl nuclease-free water and

2µl of the DNA sample. Discriminant mitochondrial primers cycling conditions were 95°C for 20

seconds, followed by 40 cycles of: 95°C for 3 seconds and 55°C for 30 seconds (**Table S1**).

Discriminant primers used linked base technology to increase specificity (Exiqon, Denmark). Melt

curves and the fly control gene *rps17* were run on each plate to control for successful single-peak

amplification and to control for DNA quality. All primers had >90% efficiency. For absolute

quantification, each 96-well qPCR plate was analyzed using StepOne Software v2.2.2 (Applied

Biosystems, USA) and Ct values were obtained by comparing each primer sample to a single

standard curve of known concentration and using identical threshold and baseline levels for each

primer target across plates. Standard curves were created by amplifying positive control samples

using PCR, calculating DNA concentrations using QBIT, and then serially diluting the sample 1:10

with distilled water to create a 5-sample curve comprising known concentrations decreasing from

10pmol/ ml. Samples with Ct values over 30 were classed as negative, confirmed by our negative

controls. As a result, the detection limit was 50 copies per sample for *rps17*, 115 for *CO3*

(coxSNP_1) and 735 for *CO3* (coxSNP_2). DNA concentration was calculated for each sample by

comparing the sample Ct values back to the equation of the standard curve. DNA copy numbers

were then calculated by using the DNA product length to determine DNA concentration. To control for fly size and extraction efficiency, copy numbers for each sample were normalized using the fly housekeeping gene as a control, giving a final value in terms of mitochondrial genome copies to fly nuclear genome copy ratio. Since *rps17* is located on Muller element D, which in *D. paulistorum* is fused to Muller element A and part of the X chromosome, the copy numbers differ between males and females. Hence, male *rps17* numbers were doubled. We used ANOVA in R version 3.3.2 (R Core Team, 2016) when testing for significant changes in copy numbers of the α-NUMT and the real β mitochondrial genome between ages and sexes of different tissues. The data was log-transformed to increase normality.

DNA extraction, genome sequencing and assembly

DNA extractions and Nanopore sequencing

Total DNA for Oxford Nanopore (ONT) sequencing of O11 was extracted from many adult flies of both sexes using the MagAttract HMW DNA Kit (Qiagen, Germany) with only minor modifications of the manufacturer's instructions. Several similar independent DNA extractions were then pooled, cleaned and concentrated with the Ampure XP beads clean-up protocol (Beckman Coulter, USA). The Circulomics SRE XS kit (Pacific Biosciences, USA) was then used to eliminate short DNA fragments before sequencing. A DNA library was prepared using the Ligation Sequencing kit (Oxford Nanopore Technologies, UK) and sequenced on an R9.4 PromethION flowcell (FLO-PRO002) at the Uppsala Genome Center, Uppsala, Sweden. Basecalling was performed with Guppy 4.3.4 and the HAC model. A total of 1.4M reads were generated, with 19.54 Gb passed bases (**Table S3**).

D. paulistorum O11 genome assembly

Reads were subsampled with Filtlong v.0.2.1 (https://github.com/rrwick/Filtlong, last visited 18th March 2022) to lower the overall coverage of the reads to facilitate the assembly process. After testing different settings (**Table S3**), the O11 Nanopore reads were filtered to a coverage of 50X, giving priority to the quality of the reads (--target_bases 12500000000 –min_length 1000 – mean_q_weight 10). The whole genome assembly was built from the dataset of subsampled reads using NextDenovo v.2.5.0 (Hu et al., 2024; https://github.com/Nextomics/NextDenovo, last visited 18th March 2022) with a read cutoff of 1 Kb, and setting the genome size to 250 Mb. The genome assembly was then polished with both the same Nanopore reads and Illumina reads from Baião et al., 2023 (available via NCBI Bioproject PRJNA643793). First, Nanopore reads were mapped to

the genome assembly with Minimap2 (Li, 2018) using the map-ont mode and bam files were sorted and indexed with samtools (Danecek et al., 2021). Then, the pipeline PEPPER-Margin-DeepVariant r0.4 (Shafin et al., 2021) was used to call variants. The variants were filtered with bcftools v1.15.1-21 (Danecek et al., 2021) using the following thresholds: QUAL>=30 && FMT/DP>=10 && FMT/GQ>=30 && FMT/VAF>=0.8, and remaining SNPs were corrected in the genome with bcftools v1.15.1-21 (Danecek et al., 2021). Illumina reads were aligned to the genome assembly using BWA mem v0.7.17-r1198-dirty (Li, 2013) and sorted with samtools v1.14 (Danecek et al., 2021), and two runs of Pilon v1.24 (Walker et al., 2014) were used for correction. BUSCO v5.2.2 with diptera_odb10 and augustus species fly was run to check genome assembly completeness, and Quast v5.0.2 (Mikheenko et al., 2018) was run with default settings to quantitatively assess the genome assembly stats. Scripts can be found in https://github.com/mmontonerin/Dpaulistorum_Mega-NUMT.

Detection and annotation of α-NUMT copies in the genome assembly

The O11 β mt-genome (Baião et al., 2023) was used as a query when running BLASTn v2.12.0+ (Camacho et al., 2009) against the genome assembly. Hits over 500 bp long and with 90% identity to the mitochondrial sequence were inspected, and the 12 mt-genome copies in the Mega-NUMT region were identified on contig ctg000940. The position of each gene in the α-NUMT copy in the Mega-NUMT was obtained by using gene sequences from the *D. willistoni* mt-genome (GD-H4-1 line) as queries in a BLASTn search. Each inferred gene was subsequently checked manually in Artemis v.16 (Rutherford et al., 2000) to identify pseudogenes and adjust the position if needed. Scripts can be found in https://github.com/mmontonerin/Dpaulistorum_Mega-NUMT.

Phylogenomic analysis of the α-NUMT copies

The 13 mitochondrial protein-coding genes from all α-NUMT copies and the genes from the real mt-genomes from several species of the willistoni group (see **Fig. 3**), previously published in Baião et al., 2023 (available via NCBI Bioproject PRJNA643793), were aligned using MAFFT v.7.407 (Katoh & Standley, 2013) and subsequently trimmed with trimAl v1.4.1 (Capella-Gutiérrez et al., 2009) using the setting -gt 0.1. The python script gene_stitcher.py (https://github.com/ballesterus/Utensils/blob/master/geneStitcher.py, last visited 12 March 2026) was used to concatenate the trimmed alignments. The concatenated alignment was used to infer a maximum likelihood phylogeny in IQTree2 v2.2.0 (Minh et al., 2020) with one partition per gene

and the parameter settings -m MFP -b 100 -T AUTO. Scripts can be found in
https://github.com/mmontonerin/Dpaulistorum_Mega-NUMT.

Analyzing insertion sequences in α-NUMTs
Finally, we used a snakemake pipeline for repeat discovery and annotation in the genome assembly,
which includes prediction of repeat content and transposable elements with RepeatModeler v1.0.8
(https://github.com/Dfam-consortium/RepeatModeler, last visited 12 March 2026). The pipeline
also creates a database based on UniProt/Swissprot (The UniProt Consortium, 2025) protein dataset
without transposable elements that initial predictions can be blasted against and removed if
matching. By this, protein-coding genes in multiple copies are avoided in our final repeat library.
The final repeat library was then used to mask the genome assembly with RepeatMasker v4.0.7
(https://github.com/Dfam-consortium/RepeatModeler, last visited 12 March 2026). Scripts can be
found in https://github.com/mmontonerin/Dpaulistorum_Mega-NUMT.

Collection of staged fly embryos for DNA extraction and DNA FISH
D. paulistorum embryos at early (0-30 min AEL), mid (3-4 hrs. AEL) and late (18-20 hrs. AEL)
stages of development were collected on homemade medium (agar, molasses, cornmeal and yeast
medium) in plastic Petri dishes (3 x 3 cm) using collection chambers at 25°C. Prior to collecting
embryos, 1–2-week-old flies were kept in these chambers for 2-3 days to adapt to the new
environment. Embryos were then collected by gently washing them off the plate onto a mesh with
water. A subset of embryos was further processed for DNA FISH (see below). The rest was
manually picked with a dissection needle and transferred immediately to an ice-cold lysis buffer
(Qiagen, Germany) and processed for DNA extraction or stored at -20°C for no more than one
week. In total, we collected 20 embryos for each sample with at least five biological replicates.

Fluorescent in situ hybridization (FISH) for Mega-NUMT visualization
We performed DNA FISH on various fly tissues using both satellite and Mega-NUMT probes. The
Mega-NUMT probe was generated by direct labeling via PCR with modified dNTPs (Alexa Fluor
488 dUTP; Jena Bioscience, Germany). We used a mixture of five mitochondrial loci (*CO1*, *CO2*,
ND1, *ND2* and *ND4*) of *D. paulistorum*, cloned beforehand in pTZ57R/5 vector (Thermo
Scientific, USA). For targeting the satellite sequence adjacent to the Mega-NUMT, we used a
primary oligo probe coupled with a sec6 adaptor (satO11-
CACACGCTCTCCGTCTTGGCCGTGGTCGATCAttttttttttTTTATAAAAATTAATACTGAG
GACAAACTGAGGACT) and the secondary probe Sec6 coupled with Cy3 (Sec6-

/5Cy3/aTGATCGACCACGGCCAAGACGGAGAGCGTGTGaa). The probe for the nuclear single copy gene *FOXO* gene was generated in the same way as the Mega-NUMT probe, but by only using three probes covering the *D. paulistorum FOXO* gene and labelled with Alexa Fluor 594 dUTP (Jena Bioscience, Germany).

To visualize Mega-NUMT in mitotically active neuroblasts of third instar larvae, we dissected their brains in PBS and incubated for 15 min in 0.5% sodium citrate. Then we fixed the brains for 15 min in 4% formaldehyde and 45% acetic acid before squashing them between the slide and coverslip. Squashed samples were immediately immersed in liquid nitrogen and transferred afterwards in 100% ethanol for 5 min. Then we air dried slides for at least 1 hour before proceeding to the hybridization. For the hybridization, we used 20 pmol of satellite probes coupled with 80 pmol of the sec6 probe and approximately 150 ng of Mega-NUMT probe in 50 μL of hybridization buffer (50% formamide, 10% dextran sulfate, 2xSSC). We heated slides for 5 min at 95°C to denature and incubated them overnight at 37°C in a humid chamber. We then washed the slides 3 times for 5 min with 4xSSCT and 3 times for 5 min with 0.1xSSC before mounting in SlowFade with DAPI and sealing with nail polish.

To visualize Mega-NUMT in mitotically active early embryos (0-3 h), staged embryos were collected as stated above, dechorionated in 50% bleach and fixed for 20 min in 3.7% paraformaldehyde in PBX (PBS and 0.15% Triton X-100) mixed with heptane (1:1 v/v). Then the embryos were postfixed in ice-cold methanol and further processed for FISH following a protocol by Gemkow et al., 1996 with slight modifications. Additionally, salivary glands and polytene chromosomes were used for Mega-NUMT visualization. The glands were dissected in PBS and fixed for 20 min in 3.7% paraformaldehyde in PBX (PBS and 0.15% Triton X-100). After washing three times in PBX, the tissues were kept in 70% ethanol until being used for DNA FISH. Polytene chromosomes were prepared according to (Pimpinelli et al., 2010) and processed for DNA FISH as described above.

Images were taken with an Olympus FW3000 laser scanning confocal microscope (Olympus Corporation, Japan) using a 60x objective and further processed with Photoshop CS6 (Adobe CS, USA), adjusting levels for each channel.

Bibliography

Avise, J. C., Shapira, J. F., Daniel, S. W., Aquadro, C. F., & Lansman, R. A. (1983).

Mitochondrial DNA differentiation during the speciation process in Peromyscus.

Molecular Biology and Evolution, *1*(1), 38–56.

https://doi.org/10.1093/oxfordjournals.molbev.a040301

Bai, R., Cui, H., Devaney, J. M., Allis, K. M., Balog, A. M., Liu, X., Schnur, R. E.,

Shapiro, F. L., Brautbar, A., Estrada-Veras, J. I., Hochstetler, L., McConkie-Rosell, A.,

McDonald, M. T., Solomon, B. D., Hofherr, S., Richard, G., & Suchy, S. F. (2021).

Interference of nuclear mitochondrial DNA segments in mitochondrial DNA testing

resembles biparental transmission of mitochondrial DNA in humans. *Genetics in*

Medicine, *23*(8), 1514–1521. https://doi.org/10.1038/s41436-021-01166-1

Baião, G. C., Schneider, D. I., Miller, W. J., & Klasson, L. (2023). Multiple

introgressions shape mitochondrial evolutionary history in Drosophila paulistorum and

the Drosophila willistoni group. *Molecular Phylogenetics and Evolution*, *180*, 107683.

https://doi.org/10.1016/j.ympev.2022.107683

Balciuniene, J., & Balciunas, D. (2019). A Nuclear mtDNA Concatemer (Mega-NUMT)

Could Mimic Paternal Inheritance of Mitochondrial Genome. *Frontiers in Genetics*, *10*.

https://www.frontiersin.org/articles/10.3389/fgene.2019.00518

Barro-Trastoy, D., & Köhler, C. (2024). Helitrons: Genomic parasites that generate

developmental novelties. *Trends in Genetics*, *40*(5), 437–448.

https://doi.org/10.1016/j.tig.2024.02.002

Behura, S. K. (2007). Analysis of Nuclear Copies of Mitochondrial Sequences in

Honeybee (Apis mellifera) Genome. *Molecular Biology and Evolution*, *24*(7), 1492–

1505. https://doi.org/10.1093/molbev/msm068

Bensasson, D., Zhang, D.-X., Hartl, D. L., & Hewitt, G. M. (2001). Mitochondrial

pseudogenes: Evolution's misplaced witnesses. *Trends in Ecology & Evolution*, *16*(6),

314–321. https://doi.org/10.1016/S0169-5347(01)02151-6

Biró, B., Gál, Z., Fekete, Z., Klecska, E., & Hoffmann, O. I. (2024). Mitochondrial

genome plasticity of mammalian species. *BMC Genomics*, *25*(1), 278.

https://doi.org/10.1186/s12864-024-10201-9

Blanchard, J. L., & Schmidt, G. W. (1995). Pervasive migration of organellar DNA to the

nucleus in plants. *Journal of Molecular Evolution*, *41*(4), 397–406.

https://doi.org/10.1007/BF00160310

Blanchard, J. L., & Schmidt, G. W. (1996). Mitochondrial DNA migration events in yeast

and humans: Integration by a common end-joining mechanism and alternative

perspectives on nucleotide substitution patterns. *Molecular Biology and Evolution*, *13*(3),

537–548. https://doi.org/10.1093/oxfordjournals.molbev.a025614

Brennecke, J., Malone, C. D., Aravin, A. A., Sachidanandam, R., Stark, A., & Hannon, G.

J. (2008). An Epigenetic Role for Maternally Inherited piRNAs in Transposon Silencing.

Science, *322*(5906), 1387–1392. https://doi.org/10.1126/science.1165171

Breton, S., Beaupré, H. D., Stewart, D. T., Hoeh, W. R., & Blier, P. U. (2007). The

unusual system of doubly uniparental inheritance of mtDNA: Isn't one enough? *Trends in*

Genetics, *23*(9), 465–474. https://doi.org/10.1016/j.tig.2007.05.011

Brown, W. M., George, M., & Wilson, A. C. (1979). Rapid evolution of animal

mitochondrial DNA. *Proceedings of the National Academy of Sciences*, *76*(4), 1967–

1971. https://doi.org/10.1073/pnas.76.4.1967

Burla, H., da Cunha, A. B., Cordeiro, A. R., Dobzhansky, Th., Malogolowkin, C., &

Pavan, C. (1949). The Willistoni Group of Sibling Species of Drosophila. *Evolution*, *3*(4),

300–314. https://doi.org/10.2307/2405716

Calvignac, S., Konecny, L., Malard, F., & Douady, C. J. (2011). Preventing the pollution

of mitochondrial datasets with nuclear mitochondrial paralogs (numts). *Mitochondrion*,

11(2), 246–254. https://doi.org/10.1016/j.mito.2010.10.004

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., &

Madden, T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*,

10(1), 421. https://doi.org/10.1186/1471-2105-10-421

Capella-Gutiérrez, S., Silla-Martínez, J. M., & Gabaldón, T. (2009). trimAl: A tool for

automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*,

25(15), 1972–1973. https://doi.org/10.1093/bioinformatics/btp348

Cohen, S., Agmon, N., Yacobi, K., Mislovati, M., & Segal, D. (2005). Evidence for

rolling circle replication of tandem genes in Drosophila. *Nucleic Acids Research*, *33*(14),

4519–4526. https://doi.org/10.1093/nar/gki764

Cohen, S., & Segal, D. (2009). Extrachromosomal Circular DNA in Eukaryotes: Possible

Involvement in the Plasticity of Tandem Repeats. *Cytogenetic and Genome Research*,

124(3–4), 327–338. https://doi.org/10.1159/000218136

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O.,

Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years

of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008.

https://doi.org/10.1093/gigascience/giab008

Dayama, G., Emery, S. B., Kidd, J. M., & Mills, R. E. (2014). The genomic landscape of

polymorphic human nuclear mitochondrial insertions. *Nucleic Acids Research*, *42*(20),

12640–12649. https://doi.org/10.1093/nar/gku1038

DeSalle, R., & Goldstein, P. (2019). Review and Interpretation of Trends in DNA

Barcoding. *Frontiers in Ecology and Evolution*, *7*.

https://www.frontiersin.org/articles/10.3389/fevo.2019.00302

Dobzhansky, T., & Spassky, B. (1959). Drosophila paulistorum, a cluster of species in

statu nascendi*†. *Proceedings of the National Academy of Sciences*, *45*(3), 419–428.

https://doi.org/10.1073/pnas.45.3.419

Dobzhansky, Th., Ehrman, L., Pavlovsky, O., & Spassky, B. (1964). The superspecies

drosophila paulistorum*. *Proceedings of the National Academy of Sciences*, *51*(1), 3–9.

https://doi.org/10.1073/pnas.51.1.3

Elías-Gutiérrez, M., Jerónimo, F. M., Ivanova, N. V., Valdez-Moreno, M., & Hebert, P.

D. N. (2008). DNA barcodes for Cladocera and Copepoda from Mexico and

Guatemala, highlights and new discoveries. *Zootaxa*, *1839*(1), Article 1.

https://doi.org/10.11646/zootaxa.1839.1.1

Fields, P. D., Waneka, G., Naish, M., Schatz, M. C., Henderson, I. R., & Sloan, D. B.

(2022). Complete Sequence of a 641-kb Insertion of Mitochondrial DNA in the

Arabidopsis thaliana Nuclear Genome. *Genome Biology and Evolution*, *14*(5), evac059.

https://doi.org/10.1093/gbe/evac059

Filée, J., Forterre, P., Sen-Lin, T., & Laurent, J. (2002). Evolution of DNA Polymerase

Families: Evidences for Multiple Gene Exchange Between Cellular and Viral Proteins.

Journal of Molecular Evolution, *54*(6), 763–773. https://doi.org/10.1007/s00239-001-

0078-x

Gemkow, M. J., Buchenau, P., & Arndt-Jovin, D. J. (1996). FISH in whole-mount

Drosophila embryos. RNA: Activation of a transcriptional locus, DNA: gene architecture

and expression. *Bioimaging*, *4*(2), 107–120. https://doi.org/10.1002/1361-

6374(199606)4:2%3C107::AID-BIO8%3E3.0.CO;2-Q

Gleason, J. M., Griffith, E. C., & Powell, J. R. (1998). A MOLECULAR PHYLOGENY

OF THE DROSOPHILA WILLISTONI GROUP: CONFLICTS BETWEEN SPECIES

CONCEPTS? *Evolution*, *52*(4), 1093–1103. https://doi.org/10.1111/j.1558-

5646.1998.tb01836.x

Gromko, M. H., & Richmond, R. C. (1978). MODES OF SELECTION MAINTAINING

AN INVERSION POLYMORPHISM IN DROSOPHILA PAULISTORUM. *Genetics*,

88(2), 357–366. https://doi.org/10.1093/genetics/88.2.357

Harrison, R. G. (1989). Animal mitochondrial DNA as a genetic marker in population and

evolutionary biology. *Trends in Ecology & Evolution*, *4*(1), 6–11.

https://doi.org/10.1016/0169-5347(89)90006-2

Hazkani-Covo, E., & Graur, D. (2007). A Comparative Analysis of numt Evolution in

Human and Chimpanzee. *Molecular Biology and Evolution*, *24*(1), 13–18.

https://doi.org/10.1093/molbev/msl149

Hebert, P. D. N., Bock, D. G., & Prosser, S. W. J. (2023). Interrogating 1000 insect

genomes for NUMTs: A risk assessment for estimates of species richness. *PLOS ONE*,

18(6), e0286620. https://doi.org/10.1371/journal.pone.0286620

Hebert, P. D. N., Cywinska, A., Ball, S. L., & deWaard, J. R. (2003). Biological

identifications through DNA barcodes. *Proceedings of the Royal Society of London.*

Series B: Biological Sciences, *270*(1512), 313–321.

https://doi.org/10.1098/rspb.2002.2218

Hirano, M., Shtilbans, A., Mayeux, R., Davidson, M. M., DiMauro, S., Knowles, J. A., &

Schon, E. A. (1997). Apparent mtDNA heteroplasmy in Alzheimer's disease patients and

in normals due to PCR amplification of nucleus-embedded mtDNA pseudogenes.

Proceedings of the National Academy of Sciences, *94*(26), 14894–14899.

https://doi.org/10.1073/pnas.94.26.14894

Hotopp, J. C. D., Clark, M. E., Oliveira, D. C. S. G., Foster, J. M., Fischer, P., Torres, M.

C. M., Giebel, J. D., Kumar, N., Ishmael, N., Wang, S., Ingram, J., Nene, R. V., Shepard,

J., Tomkins, J., Richards, S., Spiro, D. J., Ghedin, E., Slatko, B. E., Tettelin, H., &

Werren, J. H. (2007). Widespread Lateral Gene Transfer from Intracellular Bacteria to

Multicellular Eukaryotes. *Science*, *317*(5845), 1753–1756.

https://doi.org/10.1126/science.1142490

Hu, J., Wang, Z., Sun, Z., Hu, B., Ayoola, A. O., Liang, F., Li, J., Sandoval, J. R.,

Cooper, D. N., Ye, K., Ruan, J., Xiao, C.-L., Wang, D., Wu, D.-D., & Wang, S. (2024).

NextDenovo: An efficient error correction and accurate assembly tool for noisy long

reads. *Genome Biology*, *25*(1), 107. https://doi.org/10.1186/s13059-024-03252-4

Junakovic, N. (2004). Southern Blot Analysis of Individual Drosophila Flies. In W. J.

Miller & P. Capy (Eds), *Mobile Genetic Elements: Protocols and Genomic Applications*

(pp. 41–57). Humana Press. https://doi.org/10.1385/1-59259-755-6:041

Jünger, M. A., Rintelen, F., Stocker, H., Wasserman, J. D., Végh, M., Radimerski, T.,

Greenberg, M. E., & Hafen, E. (2003). The Drosophila Forkhead transcription factor

FOXO mediates the reduction in cell number associated with reduced insulin signaling.

Journal of Biology, *2*(3), 20. https://doi.org/10.1186/1475-4924-2-20

Kastritsis, C. D. (1967). A comparative study of the chromosomal polymorphs in the

incipient species of the Drosophila paulistorum complex. *Chromosoma*, *23*(2), 180–202.

https://doi.org/10.1007/BF00331112

Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software

Version 7: Improvements in Performance and Usability. *Molecular Biology and*

Evolution, *30*(4), 772–780. https://doi.org/10.1093/molbev/mst010

Keeling, P. J. (2024). Horizontal gene transfer in eukaryotes: Aligning theory with data.

Nature Reviews Genetics, *25*(6), 416–430. https://doi.org/10.1038/s41576-023-00688-5

Kim, B. Y., Wang, J. R., Miller, D. E., Barmina, O., Delaney, E., Thompson, A.,

Comeault, A. A., Peede, D., D'Agostino, E. R., Pelaez, J., Aguilar, J. M., Haji, D.,

Matsunaga, T., Armstrong, E., Zych, M., Ogawa, Y., Stamenković-Radak, M., Jelić, M.,

Veselinović, M. S., … Petrov, D. A. (2022). Correction: Highly contiguous assemblies of

101 drosophilid genomes. *eLife*, *11*, e78579. https://doi.org/10.7554/eLife.78579

Klasson, L., Kumar, N., Bromley, R., Sieber, K., Flowers, M., Ott, S. H., Tallon, L. J.,

Andersson, S. G. E., & Dunning Hotopp, J. C. (2014). Extensive duplication of the

Wolbachia DNA in chromosome four of Drosophila ananassae. *BMC Genomics*, *15*(1),

1097. https://doi.org/10.1186/1471-2164-15-1097

Kondo, N., Nikoh, N., Ijichi, N., Shimada, M., & Fukatsu, T. (2002). Genome fragment

of Wolbachia endosymbiont transferred to X chromosome of host insect. *Proceedings of*

the National Academy of Sciences, *99*(22), 14280–14285.

https://doi.org/10.1073/pnas.222228199

Kress, W. J., García-Robledo, C., Uriarte, M., & Erickson, D. L. (2015). DNA barcodes

for ecology, evolution, and conservation. *Trends in Ecology & Evolution*, *30*(1), 25–35.

https://doi.org/10.1016/j.tree.2014.10.008

Krimbas, C. B., & Powell, J. R. (1992). *Drosophila Inversion Polymorphism*. CRC Press.

Ladoukakis, E. D., & Zouros, E. (2017). Evolution and inheritance of animal

mitochondrial DNA: Rules and exceptions. *Journal of Biological Research-Thessaloniki*,

24(1), 2. https://doi.org/10.1186/s40709-017-0060-4

Li, H. (2013). *Aligning sequence reads, clone sequences and assembly contigs with BWA-*

MEM (arXiv:1303.3997). arXiv. https://doi.org/10.48550/arXiv.1303.3997

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*,

34(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Ling, F., Niu, R., Hatakeyama, H., Goto, Y., Shibata, T., & Yoshida, M. (2016). Reactive

oxygen species stimulate mitochondrial allele segregation toward homoplasmy in human

cells. *Molecular Biology of the Cell*, *27*(10), 1684–1693.

https://doi.org/10.1091/mbc.E15-10-0690

Ling, F., & Yoshida, M. (2020). Rolling-Circle Replication in Mitochondrial DNA

Inheritance: Scientific Evidence and Significance from Yeast to Human Cells. *Genes*,

11(5). https://doi.org/10.3390/genes11050514

Lopez, J. V., Yuhki, N., Masuda, R., Modi, W., & O'Brien, S. J. (1994). Numt, a recent

transfer and tandem amplification of mitochondrial DNA to the nuclear genome of the

domestic cat. *Journal of Molecular Evolution*, *39*(2), 174–190.

https://doi.org/10.1007/BF00163806

Luo, S., Valencia, C. A., Zhang, J., Lee, N.-C., Slone, J., Gui, B., Wang, X., Li, Z., Dell,

S., Brown, J., Chen, S. M., Chien, Y.-H., Hwu, W.-L., Fan, P.-C., Wong, L.-J., Atwal, P.

S., & Huang, T. (2018). Biparental Inheritance of Mitochondrial DNA in Humans.

Proceedings of the National Academy of Sciences, *115*(51), 13039–13044.

https://doi.org/10.1073/pnas.1810946115

Lutz-Bonengel, S., Niederstätter, H., Naue, J., Koziel, R., Yang, F., Sänger, T., Huber, G.,

Berger, C., Pflugradt, R., Strobl, C., Xavier, C., Volleth, M., Weiß, S. C., Irwin, J. A.,

Romsos, E. L., Vallone, P. M., Ratzinger, G., Schmuth, M., Jansen-Dürr, P., … Parson,

W. (2021). Evidence for multi-copy Mega-NUMTs in the human genome. *Nucleic Acids

Research*, *49*(3), 1517–1531. https://doi.org/10.1093/nar/gkaa1271

McDonald, J. F., Matyunina, L. V., Wilson, S., King Jordan, I., Bowen, N. J., & Miller,

W. J. (1997). Ltr retrotransposons and the evolution of eukaryotic enhancers. *Genetica*,

100(1), 3–13. https://doi.org/10.1023/A:1018392117410

McWilliams, T. G., & Suomalainen, A. (2019). Mitochondrial DNA can be inherited

from fathers, not just mothers. *Nature*, *565*(7739), 296–297.

https://doi.org/10.1038/d41586-019-00093-1

Mikheenko, A., Prjibelski, A., Saveliev, V., Antipov, D., & Gurevich, A. (2018).

Versatile genome assembly evaluation with QUAST-LG. *Bioinformatics*, *34*(13), i142–

i150. https://doi.org/10.1093/bioinformatics/bty266

Minh, B. Q., Schmidt, H. A., Chernomor, O., Schrempf, D., Woodhams, M. D., von

Haeseler, A., & Lanfear, R. (2020). IQ-TREE 2: New Models and Efficient Methods for

Phylogenetic Inference in the Genomic Era. *Molecular Biology and Evolution*, *37*(5),

1530–1534. https://doi.org/10.1093/molbev/msaa015

Muller, H. J. (1940). An analysis of the process of structural change in chromosomes

ofDrosophila. *Journal of Genetics*, *40*(1), 1–66. https://doi.org/10.1007/BF02982481

Nikoh, N., Tanaka, K., Shibata, F., Kondo, N., Hizume, M., Shimada, M., & Fukatsu, T.

(2008). Wolbachia genome integrated in an insect chromosome: Evolution and fate of

laterally transferred endosymbiont genes. *Genome Research*, *18*(2), 272–280.

https://doi.org/10.1101/gr.7144908

Noutsos, C., Kleine, T., Armbruster, U., DalCorso, G., & Leister, D. (2007). Nuclear insertions of organellar DNA can create novel patches of functional exon sequences. *Trends in Genetics*, *23*(12), 597–601. https://doi.org/10.1016/j.tig.2007.08.016

Pagnamenta, A. T., Wei, W., Rahman, S., & Chinnery, P. F. (2021). Biparental inheritance of mitochondrial DNA revisited. *Nature Reviews Genetics*, *22*(8), 477–478. https://doi.org/10.1038/s41576-021-00380-6

Papachristos, K., Miller, W. J., & Klasson, L. (2025). A co-speciation dilemma and a lifestyle transition with genomic consequences in Wolbachia of Neotropical Drosophila. *BMC Genomics*, *27*(1), 41. https://doi.org/10.1186/s12864-025-12340-z

Parfait, B., Rustin, P., Munnich, A., & Rötig, A. (1998). Coamplification of Nuclear Pseudogenes and Assessment of Heteroplasmy of Mitochondrial DNA Mutations. *Biochemical and Biophysical Research Communications*, *247*(1), 57–59. https://doi.org/10.1006/bbrc.1998.8666

Parr, R. L., Maki, J., Reguly, B., Dakubo, G. D., Aguirre, A., Wittock, R., Robinson, K., Jakupciak, J. P., & Thayer, R. E. (2006). The pseudo-mitochondrial genome influences mistakes in heteroplasmy interpretation. *BMC Genomics*, *7*(1), 185. https://doi.org/10.1186/1471-2164-7-185

Patterson, E. C., Lall, G. M., Neumann, R., Ottolini, B., Sacchini, F., Foster, A. P., Jobling, M. A., & Wetton, J. H. (2023). Defining cat mitogenome variation and accounting for numts via multiplex amplification and Nanopore sequencing. *Forensic Science International: Genetics*, *67*, 102944. https://doi.org/10.1016/j.fsigen.2023.102944

Pimpinelli, S., Bonaccorsi, S., Fanti, L., & Gatti, M. (2010). Fluorescent In Situ Hybridization (FISH) of Mitotic Chromosomes from Drosophila Larval Brain. *Cold Spring Harbor Protocols*, *2010*(3), pdb.prot5391. https://doi.org/10.1101/pdb.prot5391

Pohjoismäki, J. L. O., Goffart, S., Tyynismaa, H., Willcox, S., Ide, T., Kang, D.,

Suomalainen, A., Karhunen, P. J., Griffith, J. D., Holt, I. J., & Jacobs, H. T. (2009).

Human Heart Mitochondrial DNA Is Organized in Complex Catenated Networks

Containing Abundant Four-way Junctions and Replication Forks*. *Journal of Biological*

Chemistry, *284*(32), 21446–21457. https://doi.org/10.1074/jbc.M109.016600

Pritam, S., & Signor, S. (2025). Evolution of piRNA-guided defense against transposable

elements. *Trends in Genetics*, *41*(5), 390–401. https://doi.org/10.1016/j.tig.2024.11.011

Puertas, M. J., & González-Sánchez, M. (2020). Insertions of mitochondrial DNA into the

nucleus—Effects and role in cell evolution. *Genome*, *63*(8), 365–374.

https://doi.org/10.1139/gen-2019-0151

R Core Team. (2016). *R: A language and environment for statistical computing*

[Computer software]. R Foundation for Statistical Computing, Vienna, Austria.

https://www.R-project.org/

Rafanel, B., Protsenko, L., Handler, D., Brennecke, J., & Senti, K.-A. (2025). *Antisense*

transposon insertions into host genes trigger piRNA mediated immunity (p.

2025.07.28.667215). bioRxiv. https://doi.org/10.1101/2025.07.28.667215

Robe, L. J., Cordeiro, J., Loreto, E. L. S., & Valente, V. L. S. (2010). Taxonomic

boundaries, phylogenetic relationships and biogeography of the Drosophila willistoni

subgroup (Diptera: Drosophilidae). *Genetica*, *138*(6), 601–617.

https://doi.org/10.1007/s10709-009-9432-5

Rogers, H. H., & Griffiths-Jones, S. (2012). Mitochondrial Pseudogenes in the Nuclear

Genomes of Drosophila. *PLOS ONE*, *7*(3), e32593.

https://doi.org/10.1371/journal.pone.0032593

Rutherford, K., Parkhill, J., Crook, J., Horsnell, T., Rice, P., Rajandream, M.-A., &

Barrell, B. (2000). Artemis: Sequence visualization and annotation. *Bioinformatics*,

16(10), 944–945. https://doi.org/10.1093/bioinformatics/16.10.944

Salas, A., Schönherr, S., Bandelt, H.-J., Gómez-Carballa, A., & Weissensteiner, H.

(2020). Extraordinary claims require extraordinary evidence in asserted mtDNA

biparental inheritance. *Forensic Science International: Genetics*, *47*, 102274.

https://doi.org/10.1016/j.fsigen.2020.102274

Salas, A., Yao, Y.-G., Macaulay, V., Vega, A., Carracedo, Á., & Bandelt, H.-J. (2005). A

Critical Reassessment of the Role of Mitochondria in Tumorigenesis. *PLOS Medicine*,

2(11), e296. https://doi.org/10.1371/journal.pmed.0020296

Schiavo, G., Hoffmann, O. I., Ribani, A., Utzeri, V. J., Ghionda, M. C., Bertolini, F.,

Geraci, C., Bovo, S., & Fontanesi, L. (2017). A genomic landscape of mitochondrial

DNA insertions in the pig nuclear genome provides evolutionary signatures of

interspecies admixture. *DNA Research*, *24*(5), 487–498.

https://doi.org/10.1093/dnares/dsx019

Schwartz, M., & Vissing, J. (2002). Paternal Inheritance of Mitochondrial DNA. *New

England Journal of Medicine*, *347*(8), 576–580. https://doi.org/10.1056/NEJMoa020350

Shafin, K., Pesout, T., Chang, P.-C., Nattestad, M., Kolesnikov, A., Goel, S., Baid, G.,

Kolmogorov, M., Eizenga, J. M., Miga, K. H., Carnevali, P., Jain, M., Carroll, A., &

Paten, B. (2021). Haplotype-aware variant calling with PEPPER-Margin-DeepVariant

enables high accuracy in nanopore long-reads. *Nature Methods*, *18*(11), 1322–1332.

https://doi.org/10.1038/s41592-021-01299-w

Shutt, T. E., & Gray, M. W. (2006). Bacteriophage origins of mitochondrial replication

and transcription proteins. *Trends in Genetics*, *22*(2), 90–95.

https://doi.org/10.1016/j.tig.2005.11.007

Spassky, B., & Dobzhansky, T. (1950). Comparative genetics of Drosophila willistoni.

Heredity, *4*(2), 201–215. https://doi.org/10.1038/hdy.1950.15

The UniProt Consortium. (2025). UniProt: The Universal Protein Knowledgebase in

2025. *Nucleic Acids Research*, *53*(D1), D609–D617.

https://doi.org/10.1093/nar/gkae1010

Thomas, J., Vadnagara, K., & Pritham, E. J. (2014). DINE-1, the highest copy number

repeats in Drosophila melanogaster are non-autonomous endonuclease-encoding rolling-

circle transposable elements (Helentrons). *Mobile DNA*, *5*(1), 18.

https://doi.org/10.1186/1759-8753-5-18

Tsuji, J., Frith, M. C., Tomii, K., & Horton, P. (2012). Mammalian NUMT insertion is

non-random. *Nucleic Acids Research*, *40*(18), 9073–9088.

https://doi.org/10.1093/nar/gks424

Valiati, V. H., & Valente, V. L. S. (1997). Chromosomal polymorphism in urban

populations of Drosophila paulistorum. *Brazilian Journal of Genetics*, *20*, 567–581.

https://doi.org/https://doi.org/10.1590/S0100-84551997000400004

Vendrami, D. L. J., Gossmann, T. I., Chakarov, N., Paijmans, A. J., Litzke, V., Eyre-

Walker, A., Forcada, J., & Hoffman, J. I. (2022). Signatures of Selection on Mitonuclear

Integrated Genes Uncover Hidden Mitogenomic Variation in Fur Seals. *Genome Biology*

and Evolution, *14*(7), evac104. https://doi.org/10.1093/gbe/evac104

Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C.

A., Zeng, Q., Wortman, J., Young, S. K., & Earl, A. M. (2014). Pilon: An Integrated Tool

for Comprehensive Microbial Variant Detection and Genome Assembly Improvement. *PLOS ONE*, *9*(11), e112963. https://doi.org/10.1371/journal.pone.0112963

Wang, J.-X., Liu, J., Miao, Y.-H., Huang, D.-W., & Xiao, J.-H. (2020). Tracking the Distribution and Burst of Nuclear Mitochondrial DNA Sequences (NUMTs) in Fig Wasp Genomes. *Insects*, *11*(10), Article 10. https://doi.org/10.3390/insects11100680

Wei, W., Pagnamenta, A. T., Gleadall, N., Sanchis-Juan, A., Stephens, J., Broxholme, J., Tuna, S., Odhams, C. A., Fratter, C., Turro, E., Caulfield, M. J., Taylor, J. C., Rahman, S., & Chinnery, P. F. (2020). Nuclear-mitochondrial DNA segments resemble paternally inherited mitochondrial DNA in humans. *Nature Communications*, *11*(1), 1740. https://doi.org/10.1038/s41467-020-15336-3

Wei, W., Schon, K. R., Elgar, G., Orioli, A., Tanguy, M., Giess, A., Tischkowitz, M., Caulfield, M. J., & Chinnery, P. F. (2022). Nuclear-embedded mitochondrial DNA sequences in 66,083 human genomes. *Nature*, *611*(7934), 105–114. https://doi.org/10.1038/s41586-022-05288-7

Werren, J. H. (2011). Selfish genetic elements, genetic conflict, and evolutionary innovation. *Proceedings of the National Academy of Sciences*, *108*(supplement_2), 10863–10870. https://doi.org/10.1073/pnas.1102343108

Yao, Y.-G., Salas, A., Bravi, C. M., & Bandelt, H.-J. (2006). A reappraisal of complete mtDNA variation in East Asian families with hearing impairment. *Human Genetics*, *119*(5), 505–515. https://doi.org/10.1007/s00439-006-0154-9

Zanini, R., Müller, M. J., Vieira, G. C., Valiati, V. H., Deprá, M., & Valente, V. L. da S. (2018). Combining morphology and molecular data to improve Drosophila paulistorum (Diptera, Drosophilidae) taxonomic status. *Fly*, *12*(2), 81–94. https://doi.org/10.1080/19336934.2018.1429859