Exploratory 16S rRNA Metagenomic Analysis of Soil Microbial Communities in Agroecosystems of

North-Central Argentina

Ana Leticia Guzmán[a], Antonella Marozzi[b], Cecilia Peralta[c], Eleodoro E. Del Valle[d], Leonardo Castoldi[a]

and Leopoldo Palma[c]*

[a]Centro de Investigaciones y Transferencia de Villa María (CIT-VM), Consejo Nacional de Investigaciones

Científicas y Técnicas (CONICET), Universidad Nacional de Villa María, 5900 Villa María, Argentina.

[b]Instituto de Investigaciones en Biodiversidad y Medioambiente (INIBIOMA) – CONICET –

UNCo, Argentina.

[c]Laboratorio de Control Biotecnológico de Plagas, Instituto BIOTECMED, Departamento de Genética,

Universitat de València, Burjassot, València, 46100, Spain.

[d]ICiagro Litoral, CONICET, Facultad de Ciencias Agrarias, Universidad Nacional del Litoral, Kreder 2805,

Esperanza S3080, Argentina

*Corresponding author: Leopoldo Palma (leopoldo.palma@uv.es)

Instituto Académico Pedagógico de Ciencias Básicas y Aplicadas, Universidad Nacional de Villa María,

Avenida Arturo Jauretche 1555, Villa maría (5900), Córdoba, Argentina. Tel.: +54-(0353)-4539100

extension 3069; Fax: +54-(0353)-4539117

Abstract:

Agriculture has modified the soil structure due to the influence of external factors and processes

that affect microbial biodiversity. Metagenomics is a fundamental tool for the study of soil microbial

diversity because it provides information about the ecosystem diversity, including both the microorganisms

that cannot be isolated in culture media and those that are no longer viable in the analyzed sample. In this

work, six soil samples obtained from agroecosystems of central and northern Argentina were subjected to

a preliminary 16S metagenomic analysis. Copiotrophic bacteria (Proteobacteria and Actinobacteria) were

dominant and one of the samples had a dominance of an oligotrophic Phylum (Acidobacteria). Our findings

support previous evidence from traditionally managed agroecosystems and provide new insights into the

diversity of soil microbiomes in Argentine regions outside the Pampas. Finally, we analyzed the most

common genera with relevant species to agronomy, both beneficial and pathogenic, and their abundance

and diversity in the sequenced samples.

Keywords: 16S rRNA metagenomic sequencing, bacteriome, ecosystem degradation, biodiversity, Chaco

region

Introduction

The effects of anthropogenic activities are evident in landscapes and biological diversity, as well as at the intensification of erosion and climate change. These factors remark the need to modify production methods and their links with nature in order to promote a conscious and sustainable exploitation that prevents forcing the limits of soil capacity for providing its environmental services (Aizen et al., 2009; Guimarães, 1998). At the global scale, Argentina is the eighth largest country and has a low density of inhabitants, with population being concentrated in a few and quite focused spaces. Since the 20^{th} century, the country's economy has been based mainly on exports of rural goods (Aizen et al., 2009), which are developed in more than 80% of its area and are related mainly to agricultural, livestock, and forestry practices (Bran et al., 2017).

The development of intensive agricultural practices can be analyzed at different scales, with the ecosystem-complexes scale being the minimum representation of the predominant landscape of an ecological region in our homeland country Argentina (Aizen et al., 2009; Bran et al., 2017). This is an iterating approach, since modifications in them have been detrimental to environmental quality at all the other different scales. During the 1990-2006 period, a shift in the productive structure occurred towards soybean as the main Argentine crop, with a yield increase of approximately 45% (Aizen et al., 2009). The modifications that occurred in these territories are not only at the landscape scale but also at the level of the environmental services they provide, like the loss of soil quality, water absorption, and the alteration of biodiversity (Aizen et al., 2009).

Land use patterns have several effects on soil biodiversity, food webs, and ecosystem services (Bran et al., 2017). Agricultural activity has replaced the biological processes that regulate soil structure through external factors, such as the application of fertilizers or synthetic pesticides. In this sense, changes driven by different management practices alter soil properties, causing a direct effect on microbial communities, also known as soil microbiome. For example, modifications produced by fertilizers change the soil chemical properties that may have an influence over the presence and prevalence of some bacteria groups. preventing the growth of others (Dai et al., 2018). Furthermore, chemical fertilizers have different

effects than those of organic fertilizers. Likewise, the overuse of nitrogen (N) causes the loss of organic

carbon, with an impact on microbial biodiversity, that also increases nitrification and leaching of nitrate to

nitrogen and reaching groundwater sources (Nowinski et al., 2008; Seneviratne, 2009; Ramirez et al., 2010).

Among the traditional practices, differences were also observed between fertilization with N alone and with

N complemented with phosphorus (P) and potassium (K) (Dai et al., 2018). The modification of microbial

communities due to the application of different agricultural practices has an impact on the agroecosystems,

since microorganisms participate in soil formation through nutrient inputs (e.g. phosphorus and nitrogen),

degradation of recalcitrant organic matter, and mineral weathering (Dubey et al., 2019). In this regard, a

better understanding of the factors determining the structure of soil microbiomes may contribute to a better

understanding of soil dynamics and change the approach of the most widely used traditional agricultural

practices.

Microbial diversity should be considered an indicator of soil health and an important factor to

ensure agriculture sustainability (Kraut-Cohen et al., 2020). Metagenomic analyses allow us to identify the

microorganisms that cannot be isolated and grown in culture media; consequently, the information obtained

using method has become a fundamental tool to correlate factors that have an influence on soil microbiota

in an evolving environment (Dubey et al., 2019). These analyses have been conducted in agricultural

systems to assess variations in soil biodiversity relative to management type (Rascovan et al., 2013). In

Argentina, some metagenomic studies were done in the Pampas agroecosystems (Fierer et al., 2013;

Rascovan et al., 2013); other productive regions of the country, however, have not been subjected to these

analyses.

In this work, we conducted an exploratory metagenomic analysis of the bacterial communities

present in agricultural soil samples belonging to north-central region of Argentina, including the Chaco and

Pampas ecosystem complexes.

Material and methods

Area of analysis

Soil samples were collected from different agroecosystems cultivated with soybean (*Glycine max*
L.) and corn (*Zea mays* L.) in Argentina (Morello et al., 2012). The agroecosystems were located in the
three most populated ecoregions of the country under an extensive agricultural production that causes
significant modifications of the landscape structures.

The Dry Chaco ecoregion is part of the Great South American Chaco and is characterized by a
large flat terrain with a slight slope towards the east of Cordoba? province. The climate conditions are
widely varied, with the highest summer temperatures exceeding 47 Celsius degrees (°C), and the lowest
winter temperatures dropping to below –6 °C. Cumulative volumetric rainfall ranges from 400 to 800 mm
per year and predominant soil types include Mollisols, Entisols, and Alfisols. The landscapes of this
ecoregion have also suffered different transformations after cotton cells were consolidated, technological
changes of agriculturalization processes have been incorporated for soybean production and cattle raising
in the areas where soybean cannot be cultivated. The soils are poorly enriched, with a slight tendency to be
saline. In the Central Subhumid Chaco Complex, the physiognomy is dominated by halophytic savannahs
and Chaco forest, which were fragmented by land use conversion to extensive agriculture (Morello et al.,
2012) (Figure 1).

The Espinal ecoregion is represented by a flat to gently undulating terrain, including a lesser extent
low mountain ranges and loessial to sandy soils, Mollisols, Vertisols, and Alfisols that stand out in the
Northern Complexes, with a slightly more humid and warmer climate. Cumulative volumetric rainfall
ranges between 500 and 1000 mm per year, decreasing from north to south. This ecoregion is characterized
by the presence of low xerophytic forests, including alternating grasslands and savannas. The loss of the
natural physiognomy in this ecoregion is one of the most important of the country, and has occurred due to
processes of anthropization associated with crop production and urbanization. This region includes the
Pampa Pedemontana Complex, which is present almost entirely in Córdoba province (Morello et al., 2012).
The Pampas ecoregion is located the east-central region of Argentina and is an extensive plain with
soft undulations in some areas and a slope from West to East towards the Atlantic Ocean. The predominant

climate is subtropical, with a humidity gradient that decreases from East to West and Southwest; cumulative

volumetric rainfall ranges from 1200 mm to 700 mm annually. Average annual temperature ranges from

20 to 14 °C, decreasing latitudinally. The predominant soils are Mollisols, mostly Argiudols, Haplustols,

Hapludols, and Natracuols (Morello et al., 2012). The dominant vegetation type is the grass steppe and the

species composition varies according to the local climatic and soil characteristics. This ecoregion has the

longest history of agricultural and livestock use in the country and is divided into two sub-regions; our

study was conducted in the Undulating Pampas complex, which is located in the north and is predominantly

a grassland. However, most of the area has been converted into farmlands, and the pastures were replaced

with crops, except in some relictual spaces, where agricultural activity cannot be developed (Morello et al.,

2012).

Sample collection and analysis

Soil samples were taken with a tubular soil sampler, from the first 20 cm and as a composite of 10

random sub-samples of ~100 g each. We collected six samples from agroecosystems with traditional

agricultural management, mainly of mollisols soils. Four samples were obtained from the Central Subhumid

Chaco Complex (Los Juríes, Las Breñas Napenay and Concepción del Bermejo) and the other two were

taken from the Pampa Pedemontana (National University of Villa María, UNVM), and Pampa Ondulada

Complexes (Arrecifes). The samples were taken in February 2018 from soybean cultivated soils, whereas

UNVM was under corn management.

DNA purification, 16S rRNA sequencing and analysis

Soil samples were homogenized in a high-speed mixer (High-speed Universal Disintegrator, Pro-

Lab Diagnosis) and DNA extraction was performed using the microbiome DNA purification kit PureLink™

(Invitrogen). To verify the correct DNA extraction, total DNA electrophoresis was performed on a 1%

agarose gel, using SYBR Safe DNA staining (Thermo Fisher Scientific) and quantified by a PICODROP

spectrophotometer (PICO 100 l Spectrophotometer). An Illumina library was constructed from the purified

DNA and sequenced using the high-performance Illumina sequencing technology of the MySeq platform

(Macrogen). The conserved regions V3 and V4 of the ribosomal DNA encoding the 16S rRNA gene were

amplified by PCR using forward and reverse primers. The obtained raw reads were then trimmed (low

quality trimmed regions were removed), the paired reads were then merged and a last filtering step was

performed to remove reads shorter than 50 bp. After reads conditioning, the taxonomic assignment of the

amplicons was performed using suitable tools included in Geneious R11 (www.geneious.com). Obtained

raw reads were merged at fastq format and stored at NCBI's Sequence Read Archive (SRA) (Leinonen et

al., 2011) under Bioproject Accession No. PRJNA722523.

Diversity indexes (number of genera, Shannon index, and equitability) were calculated to compare

the samples using R's package "vegan" (Oksanen et al., 2019). Moreover, the frequency of the dominant

phylum was analyzed to compare the results of this exploratory analysis with those obtained in other

productive systems of Argentina.

Genus abundance data, expressed as % read count, were obtained with Geneious R11 software and

the 16 S biodiversity tool. These results were used to compare genus diversity and abundance in each sample

of those genera grouped according to their agronomic relevance, namely as plant-beneficial bacteria,

following a categorization reported by Basharat Ali (2019). The plant-beneficial group of bacteria includes

members of different genera such as *Burkholderia*, *Enterobacter*, *Stenotrophomonas*, *Pseudomonas*,

Herbaspirillum, *Ralstonia*, *Bradyrhizobium*, *Ochrobactrum*, *Bacillus*, *Rhizobium* and *Staphylococcus* (Ali,

2019). For the plant-pathogenic group of bacteria, we used the top 10 list of bacterial plant pathogens

described by Mansfield et al., (2012), which includes species belonging to the genera *Pseudomonas*,

Ralstonia, *Agrobacterium*, *Xanthomonas*, *Erwinia*, *Xylella*, *Dickeya* and *Pectobacterium* (Mansfield et al.,

2012).

Results

The samples with the highest bacterial diversity were those taken from C. del Bermejo and UNVM,

without differences among diversity indexes. The Shannon index for C. del Bermejo and UNVM was 5.41

and 5.50, respectively. The values observed for this index in the other sites were Las Breñas 5.18; Napenay

5.05; Los Juríes 5.24 and Arrecifes 5.37. The equitability index values were: C. del Bermejo 0.73; Las

Breñas 0.70; Napenay 0.69; Los Juríes 0.71; Arrecifes 0.73; UNVM 0.74.

The most predominant phyla in all samples were Actinobacteria, Acidobacteria, and Proteobacteria,

showing an incidence greater than 15% (Fig. 2). Proteobacteria was the phylum with the highest incidence

in samples of C. del Bermejo, Los Juríes, Arrecifes, and UNVM, with more than 40% assigned by

comparison with each amplicon. Actinobacteria was the dominant Phylum in the sample obtained in

Napenay and Acidobacteria in the sample belonging to Las Breñas. Verrucomicrobia was highly

represented in the sample from Arrecife with respect to all the other samples, but it was not dominant in

any of them.

In addition, the most abundant genera in the sequenced samples were, as per their order of

abundance (% of sequence count), *Bradyrhizobium*, *Bacillus*, *Rhizobium* and *Pseudomonas* (Figure 3). In

samples from C. del Bermejo, Napenay, Los Juríes and Arrecife, the genus *Bradyrhizobium* was the most

abundant, followed by *Bacillus, Rhizobium* and *Pseudomonas*. Samples from Las Breñas and UNVM

showed a major abundance of the genus *Bacillus* followed by *Bradyrhizobium* and *Rhizobium.* The UNVM

sample showed also the most abundant proportion of *Pseudomonas* bacteria (10%). Minor genera in terms

of abundance were detected in the six samples and included *Burkholderia Stenotrophomonas*,

Herbaspirillum, *Xanthomonas* and *Ralstonia*. None of the following genera were detected in any of the six

samples: *Enterobacter*, *Agrobacterium*, *Erwinia*, *Xylella*, *Dickeya* and *Pectobacterium* (Figure 3).

Discussion

In this work, six samples obtained from the most productive agricultural areas of Argentina were

analyzed using 16S metagenomics. The samples corresponding to C. del Bermejo and UNVM showed the

highest biodiversity indices (Shannon index and Pielou index). Variations in these samples would be due

to the influence of management on bacterial biodiversity, a topic that has been widely discussed in the

literature (Dai et al., 2018; Huang et al., 2019). The relationships between N:P and soil available phosphorus

have been indicated as an influential factor in these variables. The soil nutrient content should be evaluated

in future studies to better understand the observed variations. N and NPK fertilization systems have

different effects on the dominance of some phyla over others (Dai et al., 2018; Huang et al., 2019). Some

research works propose the importance of evaluating changes in the proportions of copiotrophic bacteria,

such as Proteobacteria, which increase their frequency due to fertilization, and the proportion of
oligotrophic bacteria like Acidobacteria, which decreases under these conditions (Fierer et al., 2013). Since
NPK fertilization is frequent in agricultural management, the high frequency of Proteobacteria in the
analyzed samples might be expected. In experiments with different fertilization systems, a predominance
of this phylum has been observed (Huang et al. 2019; Dai et al. 2018; Zou et al. 2015), with a frequency
similar to the observed in this work.

The species of the phylum Actinobacteria are also characterized by being copiotrophic organisms,
and a high frequency is usually identified in the same environments as those where Proteobacteria is present
(Fierer et al., 2013; Huang et al., 2019). The increase of Proteobacteria and Actinobacteria related to high
nitrogen fertilization was observed, regardless of soil types or vegetation present (Fierer et al., 2013).
Moreover, Actinobacteria include spore-forming and gram-positive bacteria that can survive in
environments where soils are under extreme acidification processes; therefore, the relatively higher
presence of this phylum in the samples of C. del Bermejo and Napenay might be an indicator of the
occurrence of such processes (Dai et al., 2018).

In productive systems, there is a low frequency of oligotrophic phyla, like Acidobacteria and
Verrucomicrobia (Fierer et al., 2013; Fierer et al., 2012). In this work, Acidobacteria exhibited a relatively
high frequency in the samples, whereas Verrucomicrobia showed a low frequency. The frequency of this
phylum was relatively high in Las Breñas sample. Acidobacteria subdivisions 4 and 6 were the most
abundant. These... were proposed to be correlated positively with soil pH (Kielak et al., 2016); therefore,
the relatively higher frequency of this phylum and in particular subdivisions suggests a higher pH than in
the other samples.

The phylum Verrucomicrobia had a low frequency in most of the samples, being more abundant in
Arrecife and UNVM. Verrucomicrobia used to be widely present in US grasslands (up to 50%) before they
were managed with chemical fertilizers (Fierer et al., 2013). In this regard, the low frequency of
Verrucomicrobia in the analyzed samples is consistent with the hypothesis that under agricultural
management, the frequency of this phylum is low. Studies carried out in Argentine Pampas agroecosystems

proposed that one of the major regulators of microbial biodiversity is water availability as well as some soil

properties such as silt, organic matter, nitrogen content, pH, and salinity (Dubey et al., 2019). It was also

proposed that managed systems are more heterogenic than non-cultivated environments due to changes

caused by crop rotation and fertilization. It was also proposed that the dominant bacteria in the Pampas soils

might have the capacity to adapt to these changes (Carbonetto et al., 2014). Studies in Uruguay found a

higher frequency of phylum Verrucomicrobia in unmanaged soil samples when compared to maize-grown

soils (Federici Rodriguez et al., 2017). The authors found frequencies greater than 25% for Acidobacteria

and Proteobacteria and less than 20% for Verrucomicrobia in cultivated soils, whereas in uncultivated soils

the frequencies of these phyla were more than 30% Acidobacteria, between 20 and 25% Proteobacteria and

more than 25% Verrucomicrobia (Federici Rodriguez et al., 2017). Our results are similar to those obtained

by these authors.

The abundance and distribution of bacteria with agronomic importance were also analyzed. The

most common and abundant genera found were *Bradyrhizobium*, *Rhizobium*, *Bacillus* and *Pseudomonas.*

Bradyrhizobium and *Rhizobium* (rhizobia) are gram negative nitrogen-fixing soil bacteria that infect

legume plants (e.g. soybean, peanut, etc.) and induce nodulation, favoring the conversion of molecular

nitrogen (N_2) from the air into ammonia, nitrates, and other nitrogenous compounds necessary for the plant

metabolic functions (Verma et al., 2020). In this symbiotic association, rhizobia receives carbon-rich

organic compounds that are used for the production of energy. This particular group of bacteria are specially

needed for legume plants in soils with poor sources of N_2.

The third most abundant genus found was the *Bacillus*, which includes two plant-beneficial species,

B. subtilis and *B. thuringiensis*, as the most relevant species for agriculture. *B. subtilis* has been defined as

a GRAS (Generally Recognized as Safe) microorganism and used as a model bacterium for different studies

for several years. This has also been the bacterium of choice in the industry because of its friendly genetic

manipulation and other biotechnological properties that make it suitable for large scale fermentations and

the biotechnological production of proteins and enzymes (Zhang et al., 2020). *B. subtilis* has also been used

as plant growth promoting rhizobacteria (PGPR) and as a biocontrol agent of phytopathogenic organisms

such us bacteria (Bais et al., 2004), fungi (Chen et al., 2013) and nematodes (Backer et al., 2018). *B.*

thuringiensis is the gram-positive bacterium most widely studied and used for the biological control of

invertebrate pests in agriculture. This bacterium harbors insecticidal genes encoding δ-endotoxins (Cry

and Cyt proteins) and other vegetative insecticidal proteins (VIP) exhibiting toxic activity against a wide

range of agricultural pests as well as human disease vectors (mosquitoes) and nematodes (Palma et al.,

2014). The genes encoding Cry and Vip3 proteins have been successfully used for the construction of

transgenic plants with resistance to insect pests in agriculture; thus, these plants are used in integrated pest

management programs and for reducing the use of chemical pesticides (Heckel, 2020).

Finally, *Pseudomonas* includes both plant-beneficial and plant-pathogenic species, among the most

frequent and abundant genera found in the sequenced samples. Species belonging to this genus showed

PGPR potential (Qessaoui et al., 2019), whereas other such as *Pseudomonas entomophila* exhibited

interesting insecticidal potential for the control of pests in agriculture (Dieppois et al., 2015). However,

Pseudomonas also harbors plant-pathogenic species, such as different pathovars of *Pseudomonas syringae*

(Mansfield et al., 2012). This species was voted as number one into the top 10 bacterial plant pathogens in

terms of its importance (Mansfield et al., 2012).

Minor genera in terms of abundance but with agricultural importance were *Burkholderia*,

Stenotrophomonas, *Herbaspirillum*, *Ralstonia* and *Xanthomonas*. The genus *Burkholderia* includesboth

plant-beneficial and plant-pathogenic species. *Burkholderia phenazinium*, *Burkholderia megapolitana*, and

Burkholderia bryophila were described for their interesting antifungal activities, whereas *Burkholderia*

caryophylli and *Burkholderia gladioli* are well known plant pathogens (Eberl and Vandamme, 2016).The

Stenotrophomonas genus includes some species exhibit interesting biotechnological properties, such as

Stenotrophomonas maltophilia, due to their potential plant growth-promoting effects for the biological

control of some phytopathogenic fungi (Wolf et al., 2002). Some species of the *Herbaspirillum* genus are

N_2-fixing endophytic bacteria (Schmid et al., 2006). *Ralstonia* and *Xhantomonas* include several plant-

pathogenic species included into the top 10 list of bacterial plant pathogens (Mansfield et al., 2012), with

Ralstonia solanacearum, *Xanthomonas oryzae* pathovar *oryzae*, *Xanthomonas campestris* pathovars and

Xanthomonas axonopodis pv. *manihotis* ranking second, fourth, fifth and sixth, respectively (Mansfield et
al., 2012).

The presence of *Enterobacter* was not detected in this study, although *Enterobacter* species may
be found in different environmental samples (Davin-Regli et al., 2019). *Agrobacterium tumefaciens*, the
most representative plant-pathogenic bacterium species of the genus, and the third in the top 10 list of
bacterial plant pathogens (Mansfield et al., 2012), was also not detected into the analyzed samples. Species
belonging to Erwinia, *Xylella*, *Pectobacterium* and *Dickeya* were also included in the top ten list of plant-
pathogenic bacteria, ranking 7th, 8th , 9th and 10th, respectively, with *Erwinia amylovora*, *Xylella fastidiosa*,
Dickeya dadantii, *Dickeya* solani and *Pectobacterium carotovorum* and *P. atrosepticum* being the most
representative plant-pathogenic species of these genera (Mansfield et al., 2012); these species were also
absent or not detected in the analyzed samples

Conclusions

The results obtained in this work are the first ones focusing on the bacteriome from Argentine soils
that are subjected to agricultural management in areas outside the Pampas region. Although data provided
here are preliminary and should be considered exploratory, they provide the basis for a high-throughput
metagenomic study involving more than 200 samples obtained from different agricultural regions from
Argentina. In addition, the present results suggest that the analyzed sites had similar behavior to other
systems under agricultural management, providing basic information for future analyses about the evolution
of the microbiota in these agricultural systems.

Acknowledgements

This work was supported by grants PIC-UNVM 2017-2019 from Universidad Nacional de Villa

María, PIP-2017-2019 GI from CONICET and PICT-2017-0087 from Ministerio de Ciencia, Tecnología e

Innovación (MinCyT) Argentina.

Bibliography

Aizen, M. A., et al., 2009. Expansión de la soja y diversidad de la agricultura argentina. Ecología Austral. 19, 45-54.

Ali, B., 2019. Functional and Genetic Diversity of Bacteria Associated with the Surfaces of Agronomic Plants. Plants (Basel). 8.

Backer, R., et al., 2018. Plant Growth-Promoting Rhizobacteria: Context, Mechanisms of Action, and Roadmap to Commercialization of Biostimulants for Sustainable Agriculture. Front Plant Sci. 9, 1473.

Bais, H. P., et al., 2004. Biocontrol of *Bacillus subtilis* against infection of *Arabidopsis* roots by *Pseudomonas syringae* is facilitated by biofilm formation and surfactin production. Plant Physiol. 134, 307-19.

Bran, D., et al., Los indicadores de calidad de suelo como un componente de la sustentabilidad de los agroecosistemas. In: M. G. Wilson, (Ed.). Ediciones INTA, Entre Ríos, 2017, pp. 15-17.

Carbonetto, B., et al., 2014. Structure, composition and metagenomic profile of soil microbiomes associated to agricultural land use and tillage systems in Argentine Pampas. PLoS ONE. 9.

Chen, Y., et al., 2013. Biocontrol of tomato wilt disease by *Bacillus subtilis* isolates from natural environments depends on conserved genes mediating biofilm formation. Environ Microbiol. 15, 848-864.

Dai, Z., et al., 2018. Long-term nitrogen fertilization decreases bacterial diversity and favors the growth of Actinobacteria and Proteobacteria in agro-ecosystems across the globe. Global Change Biology. 24, 3452-3461.

Davin-Regli, A., et al., 2019. Enterobacter spp.: Update on Taxonomy, Clinical Aspects, and Emerging Antimicrobial Resistance. Clin Microbiol Rev. 32.

Dieppois, G., et al., *Pseudomonas entomophila*: A Versatile Bacterium with Entomopathogenic

Properties. In: J.-L. Ramos, et al., Eds.), Pseudomonas: Volume 7: New Aspects of

Pseudomonas Biology. Springer Netherlands, Dordrecht, 2015, pp. 25-49.

Dubey, A., et al., 2019. Soil microbiome: a key player for conservation of soil health under

changing climate. Biodiversity and Conservation. 28**,** 2405-2429.

Eberl, L., Vandamme, P., 2016. Members of the genus *Burkholderia*: good and bad guys.

F1000Res. 5.

Federici Rodriguez, M. T., et al., 2017. Soil and Rhizosphere Bacterial Diversity in Maize Agro-

Ecosystem. Sustainable Agriculture Research. 6**,** 35-35.

Fierer, N., et al., 2013. Reconstructing the microbial diversity and function of pre-agricultural

tallgrass prairie soils in the United States. Science. 342**,** 621-624.

Fierer, N., et al., 2012. Comparative metagenomic, phylogenetic and physiological analyses of soil

microbial communities across nitrogen gradients. ISME Journal. 6**,** 1007-1017.

Guimarães, R. P., 1998. Aterrizando un cometa: indicadores territoriales de sustentabilidad.

Heckel, D. G., 2020. How do toxins from *Bacillus thuringiensis* kill insects? An evolutionary

perspective. Arch Insect Biochem Physiol 104**,** 1-12.

Huang, Q., et al., 2019. The 19-years inorganic fertilization increased bacterial diversity and

altered bacterial community composition and potential functions in a paddy soil. Applied

Soil Ecology. 144**,** 60-67.

Kielak, A. M., et al., 2016. The ecology of Acidobacteria: Moving beyond genes and genomes.

Frontiers in Microbiology. 7**,** 1-16.

Kraut-Cohen, J., et al., 2020. Effects of tillage practices on soil microbiome and agricultural

parameters. Science of the Total Environment. 705**,** 135791-135791.

Leinonen, R., et al., 2011. The sequence read archive. Nucleic Acids Res. 39**,** D19-21.

Mansfield, J., et al., 2012. Top 10 plant pathogenic bacteria in molecular plant pathology. Mol

Plant Pathol. 13**,** 614-29.

Morello, J., et al., 2012. Ecorregiones y complejos ecosistémicos Argentinos. Orientación Gráfica

Editora.

Oksanen, J., et al., vegan: Community Ecology Package. 2019.

Palma, L., et al., 2014. *Bacillus thuringiensis* toxins: an overview of their biocidal activity. Toxins

(Basel). 6**,** 3296-325.

Qessaoui, R., et al., 2019. Applications of New Rhizobacteria *Pseudomonas* Isolates in

Agroecology via Fundamental Processes Complementing Plant Growth. Sci Rep. 9**,** 12832.

Rascovan, N., et al., 2013. The PAMPA datasets: A metagenomic survey of microbial

communities in Argentinean pampean soils. Microbiome. 1**,** 1-6.

Schmid, M., et al., The Genus *Herbaspirillum*. In: M. Dworkin, et al., Eds.), The Prokaryotes:

Volume 5: Proteobacteria: Alpha and Beta Subclasses. Springer New York, New York,

NY, 2006, pp. 141-150.

Verma, R., et al., Chapter 4 - Rhizobium. In: N. Amaresan, et al., Eds.), Beneficial Microbes in

Agro-Ecology. Academic Press, 2020, pp. 37-54.

Wolf, A., et al., 2002. *Stenotrophomonas rhizophila* sp. nov., a novel plant-associated bacterium

with antifungal properties. Int J Syst Evol Microbiol. 52**,** 1937-1944.

Zhang, X., et al., 2020. Applications of *Bacillus subtilis* Spores in Biotechnology and Advanced

Materials. Appl Environ Microbiol. 86.

 Figure captions

 Figure 1

368   **Figure 2**

372   **Figure 3**

373   Los Juries   Arrecife   VIIla María (UNVM)

Figure legends

Figure 1. Schematic map showing the sampled geographical regions.

Figure 2. Taxonomic profiles of north-center regions of Argentina. The most abundant taxa are shown in the bars chart and shows the dynamics of variation in the microbial communities (Phyla) in the analyzed samples.

Figure 3. Comparison among the most relevant plant-beneficial and plant-pathogenic groups of bacteria. A pie graph was constructed for each sample by calculating the % read count for a given genus within the group of bacteria defined in this study as agronomically relevant. Read counts and mean confidence values were calculated using 16S Biodiversity tool included in Geneious R11 (www.geneious.com). Only those genera exhibiting mean confidence values >50 are shown.