Genomic and functional characterization of *Bacillus* sp. strain Bva_UNVM-123 reveals
resistance and environmental adaptation traits.

Cecilia Peralta[1], Diego Herman Sauka[2], Verónica Felipe[3,4], Sergio Redondo-Moreno[1],
Eleodoro E. Del Valle[5] and Leopoldo Palma[1*]

[1]Laboratorio de Control Biotecnológico de Plagas, Instituto BIOTECMED, Departamento de
Genética, Universitat de València, 46100 València, Spain.

[2]Instituto Nacional de Tecnología Agropecuaria (INTA), Instituto de Microbiología y Zoología
Agrícola, 1712 Castelar, Argentina.

[3]Universidad Nacional de Villa María. Instituto Académico Pedagógico de Ciencias Básicas y
Aplicadas. Villa María, Argentina.

[4]CONICET-Universidad Nacional de Villa María. Instituto Multidisciplinario de Investigación
y Transferencia Agroalimentaria y Biotecnológica (IMITAB). Villa María, Argentina

[5]ICiagro Litoral, CONICET, Facultad de Ciencias Agrarias, Universidad Nacional del Litoral,
Kreder 2805, Esperanza S3080, Argentina.

*Correspondence: leopoldo.palma@uv.es

Abstract
The *Bacillus* genus comprises physiologically versatile, endospore-forming bacteria widely
distributed in natural environments. In this study, we report the isolation and genomic
characterization of strain Bva_UNVM-123, recovered from agricultural soil in Pergamino,
Argentina. Whole-genome sequencing using Illumina technology yielded a 5.1 Mbp draft
genome assembled in 67 contigs with a GC content of 36%. Comparative genomic analyses
using the TYGS server and digital DNA–DNA hybridization (dDDH) values supported its
classification as a potentially novel species within the *Bacillus* sensu lato (s.l.) group. Genome
annotation revealed 4,866 protein-coding genes, including multiple determinants conferring
resistance to antibiotics (e.g., fosfomycin, tetracycline, beta-lactams) and toxic heavy metals
(e.g., arsenic, cadmium, mercury), supporting its potential application in bioremediation.

Additionally, PathogenFinder predicted a low probability of human pathogenicity (0.207),
reinforcing its safety for environmental use. Functional classification based on Swiss-Prot
further supported a metabolically versatile profile and revealed the presence of resistance-
related categories associated with environmental adaptation. This study adds to the growing
knowledge of environmental *Bacillus* species and their biotechnological potential.

Keywords: *Bacillus* spp.; genomic sequence; antibiotic resistance; heavy metal resistance; soil
microbiome; NCBI; genome assembly; Firmicutes; endospore-forming bacteria;
bioremediation.

Introduction

Species belonging to the *Bacillus* genus are known for their remarkable natural properties, which range from the synthesis of various toxins (e.g., cereulide) to the production of proteins and enzymes with numerous biotechnological applications, including fermentation, biodegradation, and the production of secondary metabolites. Many of these organisms can synthesize a wide range of bioactive compounds, including antibiotics (e.g., bacitracina), enzymes (amylases, proteases, lipases, etc.), and biopolymers, which can be used in industrial applications [Herrmannet al., 2024].

The genes responsible for these biosynthetic pathways are often located on extrachromosomal DNA elements, such as plasmids or bacteriophages, which can harbor additional virulence factors including toxins and antibiotic resistance genes [Bennett, 2008].

These plasmids are typically transferable, and through horizontal gene transfer (HGT), they can be transmitted to related bacterial species [Saukaet al., 2024]. Horizontal transfer of genetic material between bacteria plays a crucial role in the spread of antibiotic resistance, as it allows for rapid dissemination of resistance traits across different bacterial populations [Kaplan, 2014, Partridgeet al., 2018]. This transfer can occur between pathogenic and non-pathogenic bacteria, and in the case of non-pathogenic species, it can result in the presence of antibiotic resistance genes even in bacteria that do not cause disease [Martínez, 2008]. Notably, recent studies emphasize the role of non-pathogenic soil bacteria in harboring mobile resistance elements that may be horizontally transferred to clinical pathogens under environmental stress conditions. Such non-pathogenic bacteria, isolated from various ecosystems like soil, water, or animal manure, are of great interest because they can serve as reservoirs for resistance genes that may eventually be transferred to pathogenic organisms under selective pressure [Larsson and Flach, 2022].

Antibiotic resistance genes are essential mechanisms used by pathogenic bacteria to evade the effects of antibiotics, a challenge that has become a major global health concern. However, they are not only found in pathogenic bacteria but are also commonly detected in non-pathogenic environmental microorganisms, including species used in biotechnology and environmental applications [Amos, 2013]. This characteristic is particularly interesting from an environmental management perspective because non-pathogenic, antibiotic-resistant bacteria could potentially be utilized in bioremediation strategies. For example, certain *Bacillus* strains with intrinsic resistance to antibiotics can be employed in bioremediation efforts to clean

up environments contaminated by antibiotics, particularly those originating from hospitals,

dairy farms, and livestock operations, where residues of antibiotics often persist in soil and

water [Gulet al., 2022]. This provides a valuable opportunity for using these bacteria in the

development of new biotechnological tools aimed at mitigating the environmental impact of

antibiotic contamination, which is a growing concern in public health and environmental

science [Aprejaet al., 2022].

In addition to antibiotic resistance, several *Bacillus* species are known to tolerate or resist

toxic heavy metals such as arsenic, cadmium, lead, copper, and mercury. This resistance is

often mediated by efflux systems, metal-binding proteins, and enzymatic detoxification

pathways encoded in their genomes [Prabhakaranet al., 2016, Margaryanet al., 2021]. These

properties make them promising candidates for bioremediation strategies aimed at

decontaminating polluted environments. The capacity to thrive in metal-contaminated soils,

together with their ability to form resilient endospores, enhances their potential in long-term or

large-scale remediation projects [Singh, 2014].

The capacity of *Bacillus* species to adapt to diverse environments, combined with their

resistance to antibiotics and heavy metals, makes them ideal candidates for further research in

the field of bioremediation. By leveraging their natural resistance mechanisms, it may be

possible to engineer more efficient systems for cleaning up contaminated environments,

particularly those impacted by human activity.

In this work, we aimed to sequence and analyze the genome of strain Bva_UNVM-123, a

newly isolated *Bacillus* strain from soil, to explore its taxonomic identity and biotechnological

potential. The assembled draft genome sequence of Bva_UNVM-123 provides insights into its

genomic makeup, revealing potential resistance genes against antibiotics and heavy metals,

which may contribute to its potential as a bioremediation agent. This study contributes to a

deeper understanding of the genetic factors underlying resistance in *Bacillus* species, further

supporting their application in environmental remediation. Additionally, genome-based

analyses suggest that strain Bva_UNVM-123 may represent a novel species within the *Bacillus*

sensu lato (s.l.) group, for which the name *Bacillus pergaminensis* has been proposed.

Materials and Methods

Isolation and Morphological Characterization

The strain Bva_UNVM-123 was isolated from a soil sample collected from soybean field
in Pergamino, Buenos Aires Province (Argentina), as part of a screening program aimed at
identifying entomopathogenic bacteria (e.g., *Bacillus thuringiensis*) as previously described
[Iriarteet al., 1998].

Those strains showing a *B. thuringiensis*-like (matte white and with uneven borders)
phenotype were reisolated in axenic cultures and observed by phase contrast microscopy
(Nikon Eclipse Ti, Japan). Gram staining was performed using a commercial kit (Biopack,
Buenos Aires, Argentina), following the manufacturer's instructions. In addition, the strain was
also examined using Scanning Electron Microscopy (SEM) at the Comprehensive Center for
Electron Microscopy (CONICET-Universidad Nacional de Tucumán, Argentina). The strain is
maintained as glycerol suspension (50% v/v) at −80 °C.

The type strain Bva_UNVM-123[t] was deposited in the International Culture Collection of
the Institute of Microbiology and Agricultural Zoology at INTA Castelar (IMYZA-INTA)
(access number pending).

Biochemical Characterization and Antibiotic Resistance Analyses

NaCl tolerance was evaluated by culturing the strain on TSA medium supplemented with
NaCl concentrations ranging from 2.5% to 7.5% (w/v) and incubating for 72 hours. Anaerobic
growth was assessed using the candle jar technique. Motility was tested on semisolid agar
medium (0.75% w/v). Hemolytic activity was tested on blood agar plates supplemented with
5% sheep blood (Britania Laboratories, Argentina) incubated at 37 °C for 48 hours.

The sensitivity of strain Bva_UNVM-123 to various antibiotics, chloramphenicol (30 µg),
novobiocin (5 µg), ampicillin (10 µg), tetracycline (30 µg), and penicillin (10 U) (Britania
Laboratories, Argentina), was assessed on Mueller–Hinton agar using the disc diffusion
method. Oxidase and catalase activities were determined using standard microbiological
procedures. Additional physiological and biochemical characterizations were conducted using
the API 50CHB, API ZYM, and API 20A systems (bioMérieux), following the protocols
provided by the manufacturer.

Fatty acid profiling of the strain was performed using the MIDI Sherlock Microbial
Identification System (version 6.2B) following the methodology described by Ferrari et al.,
(2018). The sample was analyzed using the RTSBA6 method, and fatty acids were identified
based on their equivalent chain length (ECL) and retention time [Ferrariet al., 2018].

DNA isolation, Genome assembly and Annotation and Species Delimitation

For DNA extraction, strain Bva_UNVM-123 was grown in LB broth for 16 hours. Genomic DNA, including chromosomal and plasmid DNA, was extracted using the Wizard Genomic DNA Purification Kit (Promega, Madison, WI, USA), following the manufacturer's protocol for gram-positive bacteria. The extracted DNA was analyzed by electrophoresis in 1% agarose gels, stained with SYBR Safe (ThermoFisher Scientific, Waltham, MA, USA), and quantified using a PICODROP PICO 100 μL spectrophotometer.

The extracted DNA was subjected to PCR to confirm its belonging to the *Bacillus* genus [Wuet al., 2006]. The PCR assay was performed using the methodology described by Wu et al. (2006), which involves a group-specific primer set designed for *Bacillus* species identification.

Then, the purified DNA was used to construct a pooled Illumina sequencing library, which was sequenced at the Genomics Unit of the National Institute of Agricultural Technology (INTA, Argentina) using Illumina high-throughput sequencing technology. Raw sequencing reads were processed to remove low-quality regions and assembled into contigs using Velvet within Geneious version R11 software (www.geneious.com) with default parameters for *de novo* assembly.

Assembly statistics were evaluated using QUAST v4.5 [Gurevichet al., 2013], and genome completeness and contamination were assessed using CheckM [Parkset al., 2015]. Gene predictions and functional annotations were done via PGAP [16] and validated through the RAST server [Azizet al., 2008].

Genome annotation was performed using the NCBI Prokaryotic Genome Annotation Pipeline (2023 release) and the RAST server [Azizet al., 2008].

Species delimitation was carried out using the Type (Strain) Genome Server [Meier-Kolthoff and Goker, 2019] and PubMLST [Jolleyet al., 2018]. Secondary metabolites searches were performed using antiSmash server [Blinet al., 2019].

Functional classification based on Swiss-Prot homology

RAST server predicted proteins were additionally queried against a local UniProtKB/Swiss-Prot database [Boutetet al., 2007] using BLASTp [Altschulet al., 1990] with an e-value cutoff of 1e-10, retaining only the top-scoring hit for each query. Significant hits were classified into broad functional groups according to keyword-based annotation mining, and category frequencies were plotted using the ggplot2 package in R [Wickham, 2016].

Results

Morphological Characterization

The strain formed opaque colonies with a matte texture and creamy white color after
overnight growth on TSA medium at 37 °C during 72 h. Microscopic examination revealed
rod-shaped vegetative cells with terminal deforming spores (Figure 1A and B). The terminal
positioning of the endospore is consistent with known morphological traits in environmental
Bacillus species [Voset al., 2011].

Figure 1. Microscopic analysis of strain Bva_UNVM-123 grown on TSA medium (72 h/37
°C). A) Morphological characterization using phase-contrast microscopy and B) Scanning
electron microscopy. A scale bar (200 nm) was included in SEM images to reflect structural
dimensions.

Biochemical Characterization

Cells are Gram-stain-positive, facultatively anaerobic, non-motile, and form terminal
endospores. The strain was able to grow on nutrient agar (NA), casein–collagen–yeast extract
agar (CCY), tryptic soy agar (TSA), and brain heart infusion agar. Optimal growth was
observed on TSA at 37 °C and the strain was non-hemolytic. The strain tested positive for both
catalase and oxidase activities and when cultured in the presence of chloramphenicol, did not
exhibit measurable inhibition zones.

Additional physiological and enzymatic tests were carried out using the API 50CHB, API
ZYM, and API 20A systems. However, the results obtained were inconclusive due to limited
growth under assay conditions, including incubation at 30 °C and 37 °C.

The fatty acid profile of strain Bva_UNVM-123 consisted primarily of branched-chain
saturated fatty acids, a chemotaxonomic trait typical of gram-positive bacteria, particularly
members of the genus *Bacillus* or related taxa. The major components included iso- and
anteiso-fatty acids such as iso-C15:0, anteiso-C15:0, iso-C16:0, and iso-C17:0. A total of
61.21% of the detected fatty acids were identified, with iso-C15:0 representing the most
abundant component (32.46%). It is worth noting that obtaining the fatty acid profile was
technically challenging due to the limited growth of the strain under standard laboratory

conditions (e.g., LB medium), which may have affected biomass yield and detection sensitivity.
The complete fatty acid profile is presented in Table 1.

Table 1. Key Major fatty acid components of strain Bva_UNVM-123. Only those representing
≥3% of the total identified fatty acids are shown. Summed Feature 4 includes fatty acids that
could not be resolved individually by the system.

Fatty Acid	Composition (%)	Other Characteristics
iso-C15:0	32.46	Most abundant component
anteiso-C15:0	7.28	-
iso-C16:0	5.21	-
C16:0	4.39	Also known as palmitic acid
iso-C17:0	6.16	-
anteiso-C17:0	3.69	-
Summed Feature 4	5.20	Includes C17:1 iso I / anteiso B isomers

Genome Characterization and Species Identification

The Illumina sequencing produced 14,532,725 high-quality paired end reads, resulting in
67 assembled contigs with a total genome size of 5,119,019 bp in size and 36% G+C content
(Table 2).

Table 2. Key features statistics and quality analysis of the draft genome assembly of
Bva_UNVM-123 strain obtained with QUAST, CheckM and the NCBI Prokaryotic Genome
Annotation Pipeline (PGAP).

Feature statistics	Value
Assembled Genome Size (bp)	5,119,099
Contig Number	67
Largest contig (bp)	583,230
N50 (bp)	341,242
N's	5074
GC content (%)	36
Suspected Contamination (%)*	3.85
Estimated Completeness (%)*	95.85
Estimated Coverage (×)	400
PGAP Predicted Genes	5139
PGAP Predicted Protein Coding Genes	4866

*CheckM determinations.

Table 3 highlights multiple genes potentially related to antibiotic resistance, identified
using the RAST annotation server. The presence of genes conferring resistance to fosfomycin
(fosB), tetracyclines, chloramphenicol, fosmidomycin, kanamycin, oxetanocin A, and various
beta-lactams (e.g., beta-lactamase class A-like and penicillin-binding proteins) suggests that
strain Bva_UNVM-123 possesses a broad resistance profile.

Table 3. Putative annotated genes related to antibiotic resistance found at the genome sequence
of Bva_UNVM-123 strain using RAST server.

Product	Type	Copies
Efflux ABC transporter (ATP-binding/permease)	Multidrug resistance	45
Fosmidomycin resistance protein	Fosfomycin resistance	2
Similar to tetracycline resistance protein	Tetracyclin resistance	1
Similar to tetracycline resistance, MFS efflux pump	Tetracyclin resistance	1
Chloramphenicol/florfenicol resistance efflux pump	Chloramphenicol resistance	1
Zwittermicin A resistance protein	Zwittermicin A resistance	1
Kanamycin kinase	Kanamycin resistance	1
Oxetanocin A resistance protein	Oxetanocin A resistance	1
Beta-lactamase (Class A/C-like)	Beta-lactam resistance	12
Vancomycin response proteins (VanR/VanS)	Glycopeptide resistance	5
Penicillin-binding protein 1A/1B	Penicillin resistance	1
Penicillin acylase	Penicillin resistance	1
Penicillin-binding protein 4B	Penicillin resistance	1

Table 4 lists genes potentially associated with resistance to toxic heavy metals, including
arsenic, cadmium, zinc, cobalt, lead, mercury, and tellurite. These genes encode various efflux
pumps, ATPases, and regulatory proteins, such as TehB, ArsD, ACR3, and CzcD, which are
known to facilitate resistance and detoxification [Bruinset al., 2000, Nies, 2003].

Table 4. Putative heavy-metal resistance genes found at the genome sequence of Bva_UNVM-
123 strain using RAST server.

Product	Metal	Copies
Tellurium resistance proteins (TerD, TerB)	Tellurium	5
Arsenical resistance proteins (Acr3, ArsR, ATPase)	Arsenic	9
Cadmium/Zinc/Cobalt transporters (P-type ATPase, CzcD)	Cd/Zn/Co	7
Copper chaperones and transporters (CopZ, CopA)	Copper	4
Zinc ABC transporter systems (ZnuA/B/C, Zur)	Zinc	5
Manganese efflux pump (MntP)	Manganese	1

Lead, Mercury, and other heavy-metal transporters	Pb/Hg/Cd/Zn	6
Multicopper oxidases and polyphenol oxidases	Copper	1
Aluminum resistance protein	Aluminum	1

The Swiss-Prot-based functional classification revealed a proteome dominated by housekeeping and metabolic functions, together with a relevant proportion of transport-associated proteins and a smaller but clearly detectable set of resistance-related proteins. In particular, the identification of both antibiotic- and heavy metal-associated categories agrees with the resistance determinants detected by RAST and reinforces the proposed ecological adaptation and bioremediation potential of strain Bva_UNVM-123.

Metabolism represented the largest functional category (30.4%), followed by other/poorly resolved functions (30.3%), unknown proteins (12.2%), information processing (10.5%), and transport-related proteins (8.6%). Resistance-related categories, including heavy metal resistance (1.2%) and antibiotic resistance (0.4%), were also detected (Figure 2).

Figure 2. Functional classification of predicted proteins of strain Bva_UNVM-123 based on BLASTp searches against the Swiss-Prot database. Proteins were grouped into broad functional categories based on annotation keywords. Percentages are calculated relative to the total number of proteins with significant Swiss-Prot matches.

It should be noted that Swiss-Prot represents a curated but limited database, and therefore the proportion of annotated proteins reflects high-confidence functional assignments rather than the total functional potential of the genome.

238   Concerning species identification, preliminary PCR analysis showed that isolate 123

239 belongs to the genus *Bacillus* (Figure 3).

Figure 3. PCR analysis of strain Bva_UNVM-123 for the Bacillus genus showing a positive

PCR amplicon (positive control: *B. thuringiensis* HD1, 123: Bva_UNVM-123 strain).

The genome-based phylogenetic tree revealed that strain Bva_UNVM-123 clusters within

a lineage comprising members of the *Bacillus* sensu lato (s.l.) group, showing closest affiliation

to *Lysinibacillus* and related genera (Figure 4). This placement is consistent with the complex

and recently revised taxonomy of the *Bacillus* group and supports its classification as a distinct

genomic lineage.

Figure 4. Genome-based phylogenetic tree inferred using the GBDP method via the TYGS

server, showing the position of strain Bva_UNVM-123 among closely related taxa within the

Bacillus sensu lato (s.l.) group. The strain clusters in proximity to members of the genera

Lysinibacillus, *Cytobacillus*, and *Neobacillus*. Branch lengths are scaled in terms of GBDP

distance, and bootstrap values (>60%) based on 100 pseudoreplicates are indicated at the nodes.

In silico sequence typing using the PubMLST database identified alleles for multiple loci

(n = 53); however, the allelic profile did not match any previously defined sequence type,

suggesting that strain Bva_UNVM-123 represents a novel sequence type. This result is
consistent with the phylogenomic analyses and further supports the distinct genomic position
of the strain.

Based on PathogenFinder 1.1, the strain was predicted as non-pathogenic, with a human
pathogenicity probability of 0.207, indicating its safety for environmental applications
[Cosentinoet al., 2013].

Discussion

After comprehensive genomic and phenotypic analyses, including phylogenomic
reconstruction and digital DNA–DNA hybridization (dDDH) values below the 70% species
threshold, strain Bva_UNVM-123 likely represents a novel species within the *Bacillus* sensu
lato (s.l.) group. These findings suggest that it constitutes a distinct genomic lineage within this
taxonomic framework, for which the name *Bacillus pergaminensis* has been proposed.

The genome of strain Bva_UNVM-123 reveals multiple adaptive traits that support its role
in bioremediation. In line with this, preliminary phenotypic assays showed that strain
Bva_UNVM-123 exhibited no measurable inhibition zones against chloramphenicol and
novobiocin, suggesting intrinsic tolerance that may correspond to the resistance genes found in
the genome. Nonetheless, further MIC assays are necessary to validate this resistance. The
relatively high proportion of uncharacterized proteins highlights the presence of potentially
novel functions, which may represent unexplored metabolic or adaptive capabilities.

In addition, the presence of antibiotic resistance genes such as fosB, tetA, bla-like beta-
lactamases, and multidrug efflux systems suggests long-term exposure to antibiotic residues in
soil environments. This aligns with findings that environmental *Bacillus* species often act as
reservoirs for antibiotic resistance genes due to selective pressure in natural and agricultural
soils [Martínez, 2008, Larsson and Flach, 2022]. This resistance could be determinant and
particularly relevant for bioremediation applications, where bacteria must survive in antibiotic-
contaminated environments. These results are consistent with the resistance determinants
identified through RAST annotation (Tables 3 and 4), further supporting the functional
potential of the strain for survival in contaminated environments.

The use of API-based systems for metabolic profiling did not yield conclusive results. This
outcome is frequently observed with environmental isolates, which often exhibit poor growth
or atypical metabolic patterns when tested in commercial platforms originally designed for
clinical or fast-growing laboratory strains. These limitations highlight the need to optimize test

conditions or employ alternative phenotypic strategies when characterizing environmentally adapted bacteria. For instance, even in industrial contexts such as dairy processing environments, phenotypic microbiological techniques have shown limited reliability for environmental strains like *Pseudomonas* spp. and *Bacillus* spp., often resulting in inconclusive species-level identifications [Wiedmannet al., 2000]. Consequently, in this study, species identification relied primarily on genome-based approaches, which provided higher taxonomic resolution and reliability.

Additionally, the fatty acid profile of strain Bva_UNVM-123 did not match any entry in the RTSBA6 reference library, which is consistent with the genomic evidence suggesting novelty. However, the total response signal obtained in both replicates was below the recommended threshold for reliable identification. This limitation was most likely due to the poor growth of the strain under the conditions used for biomass collection, rather than to methodological issues. Further optimization of culture conditions is therefore needed to obtain sufficient biomass for accurate chemotaxonomic profiling.

Moreover, the strain harbors a diverse array of heavy metal resistance genes, including *arsD*, *acr3*, *czcD*, and *tehB*, which confer tolerance to arsenic, cadmium, mercury, and tellurite. These genes are commonly found in metal-contaminated environments and are considered key biomarkers for bacterial adaptation and survival in such niches [Bruinset al., 2000, Nies, 2003, Margaryanet al., 2021]. The abundance and diversity of these genes suggest that strain Bva_UNVM-123 is well adapted to metal-contaminated environments. This further supports its potential use in bioremediation, particularly in soils impacted by industrial or agricultural pollutants. The presence of both antibiotic and heavy metal resistance-related functional categories, even at relatively low proportions, is particularly relevant, as these traits are often co-selected in environmental bacteria exposed to anthropogenic pressures. This supports the hypothesis that strain Bva_UNVM-123 may be adapted to environments impacted by agricultural and chemical inputs.

The non-pathogenic nature of the strain, as predicted by PathogenFinder, along with the absence of virulence determinants and hemolytic activity, strongly suggests its suitability for environmental application. This safety profile is crucial for any candidate microorganism intended for bioremediation purposes, as environmental deployment requires strict biosafety assessments [Cosentinoet al., 2013].

319   Importantly, the genome of the strain Bva_UNVM-123 also provides a valuable resource
320   for future biotechnological exploration. Its genetic repertoire could be exploited not only for
321   bioremediation of antibiotic- and metal-polluted sites but also for the potential production of
322   industrial enzymes or bioactive secondary metabolites, as often observed in *Bacillus* spp.
323   [Blinet al., 2019, Herrmannet al., 2024].
324
325

Author contributions

Conceptualization, L.P., methodology, D.H.S, C.P and L.P.; formal analysis, D.H.S, C.P, V.F. and L.P.; investigation, D.H.S; C.P, V.F., E.E.D.V and L.P.; resources, L.P.; data curation, L.P.; writing—original draft preparation, L.P.; writing, review and editing, L.P. and D.H.S; visualization, L.P.; supervision, L.P.; project administration, L.P.; funding acquisition, L.P. All authors have read and agreed to the published version of the manuscript.

Conflicts of interest

The authors declare there are no conflicts of interest.

Funding information

This work was supported by MCIN/AEI and the European Union through the Ramón y Cajal programme (RYC2023-043507-I). L.P. was supported by the Ramón y Cajal research contract (RYC2023-043507-I).

Data availability

The unprocessed genome sequencing data have been deposited in the NCBI database under the BioSample accession SAMN18815149, within the BioProject PRJNA723377. The corresponding genome assembly is accessible through the NCBI Whole Genome Shotgun (WGS) database under the accession number JAGTPU000000000.

Acknowledgements

Leopoldo Palma gratefully acknowledges the Spanish Ministry of Science, Innovation, and Universities, the Spanish State Research Agency, and the European Union for funding his Ramón y Cajal contract (grant ref. RYC2023-043507-I).

Appendix A. Supplementary data

Supplementary data associated with this article, including the TYGS taxonomic report, RAST annotation tables, Swiss-Prot BLAST results, and scripts used for functional classification, are available at Zenodo (DOI: 10.5281/zenodo.19456917).

References

1. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990 1990/10/05/;215(3):403–10.

2. Amos GC. Environmental reservoirs of antibiotic resistance. University of Warwick; 2013.

3. Apreja M, Sharma A, Balda S, Kataria K, Capalash N, Sharma P. Antibiotic residues in environment: antimicrobial resistance development, ecological risks, and bioremediation. Environmental Science and Pollution Research. 2022:1–17.

4. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, et al. The RAST Server: rapid annotations using subsystems technology. BMC Genom. 2008 Feb 8;9(75):1–15.

5. Bennett PM. Plasmid encoded antibiotic resistance: acquisition and transfer of antibiotic resistance genes in bacteria. British journal of pharmacology. 2008;153(S1):S347–S57.

6. Blin K, Pascal Andreu V, de Los Santos ELC, Del Carratore F, Lee SY, Medema MH, et al. The antiSMASH database version 2: a comprehensive resource on secondary metabolite biosynthetic gene clusters. Nucleic Acids Res. 2019 Jan 8;47(D1):D625–D30.

7. Boutet E, Lieberherr D, Tognolli M, Schneider M, Bairoch A. UniProtKB/Swiss-Prot: the manually annotated section of the UniProt KnowledgeBase. Plant bioinformatics: methods and protocols. Springer; 2007. p. 89–112.

8. Bruins MR, Kapil S, Oehme FW. Microbial resistance to metals in the environment. Ecotoxicology and environmental safety. 2000;45(3):198–207.

9. Cosentino S, Voldby Larsen M, Møller Aarestrup F, Lund O. PathogenFinder-distinguishing friend from foe using bacterial whole genome sequence data. PloS one. 2013;8(10):e77302.

10. Ferrari AE, Ravnskov S, Wall LG. Crop rotation in no-till soils modifies the soil fatty acids signature. Soil use and management. 2018;34(3):427–36.

11. Gul S, Hussain S, Khan H, Khan KI, Khan S, Ullah S, et al. Chapter 4 - Advances in bioremediation of antibiotic pollution in the environment. In: Kumar S, Hashmi MZ, editors. Biological Approaches to Controlling Pollutants. Woodhead Publishing; 2022. p. 49–78.

383   12.   Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for

384   genome assemblies. Bioinformatics. 2013 Apr 15;29(8):1072–5.

385   13.   Herrmann LW, Letti LAJ, de Oliveira Penha R, Soccol VT, Rodrigues C, Soccol CR.

386   *Bacillus* genus industrial applications and innovation: First steps towards a circular

387   bioeconomy. Biotechnology Advances. 2024;70:108300.

388   14.   Iriarte J, Bel Y, Ferrandis MD, Andrew R, Murillo J, Ferré J, et al. Environmental

389   distribution and diversity of *Bacillus thuringiensis* in Spain. Syst Appl Microbiol.

390   1998 Mar;21(1):97–106.

391   15.   Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics:

392   BIGSdb software, the PubMLST.org website and their applications. Wellcome Open

393   Res. 2018;3(124):2–20.

394   16.   Kaplan T. The role of horizontal gene transfer in antibiotic resistance. Eukaryon.

395   2014;10:80–81.

396   17.   Larsson DJ, Flach C-F. Antibiotic resistance in the environment. Nature Reviews

397   Microbiology. 2022;20(5):257–69.

398   18.   Margaryan A, Panosyan H, Birkeland N-K. Heavy metal resistance in prokaryotes:

399   Mechanism and application. Microbial communities and their interactions in the

400   extreme environment. 2021:273–313.

401   19.   Martínez JL. Antibiotics and antibiotic resistance genes in natural environments.

402   Science. 2008;321(5887):365–67.

403   20.   Meier-Kolthoff JP, Goker M. TYGS is an automated high-throughput platform for

404   state-of-the-art genome-based taxonomy. Nat Commun. 2019 May 16;10(1):1–10.

405   21.   Nies DH. Efflux-mediated heavy metal resistance in prokaryotes. FEMS

406   microbiology reviews. 2003;27(2-3):313–39.

407   22.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing

408   the quality of microbial genomes recovered from isolates, single cells, and

409   metagenomes. Genome Res. 2015 Jul;25(7):1043–55.

410   23.   Partridge SR, Kwong SM, Firth N, Jensen SO. Mobile genetic elements associated

411   with antimicrobial resistance. Clinical microbiology reviews.

412   2018;31(4):10.1128/cmr. 00088–17.

413   24.   Prabhakaran P, Ashraf MA, Aqma WS. Microbial stress response to heavy metals in

414   the environment. Rsc Advances. 2016;6(111):109862–77.

25. Sauka DH, Del Valle EE, Palma L. *Bacillus toyonensis* biovar Thuringiensis: an
overlooked entomopathogen? bioRxiv. 2024:2024.07. 21.604280.

26. Singh R. Microorganism as a tool of bioremediation technology for cleaning
environment: a review. Proceedings of the International Academy of Ecology and
Environmental Sciences. 2014;4(1):1.

27. Vos P, Garrity G, Jones D, Krieg NR, Ludwig W, Rainey FA, et al. Bergey's manual of
systematic bacteriology: Volume 3: The Firmicutes. New York: Springer New York,
2009.; 2011.

28. Wickham H. Data analysis. ggplot2: elegant graphics for data analysis. Springer;
2016. p. 189–201.

29. Wiedmann M, Weilmeier D, Dineen Sean S, Ralyea R, Boor Kathryn J. Molecular
and Phenotypic Characterization of *Pseudomonas* spp. Isolated from Milk. Applied
and Environmental Microbiology. 2000 2000/05/01;66(5):2085–95.

30. Wu X-Y, Walker MJ, Hornitzky M, Chin J. Development of a group-specific PCR
combined with ARDRA for the identification of *Bacillus* species of environmental
significance. Journal of Microbiological Methods. 2006 2006/01/01;64(1):107–19.

431