Draft Genome Sequence of *Bacillus pergaminensis* sp. nov. strain Bva_UNVM-123: A
Promising Candidate for Bioremediation.

Cecilia Peralta[1], Diego Herman Sauka[2], Verónica Felipe[3,4], Eleodoro E. Del Valle[5] and
Leopoldo Palma[1*]

[1]Laboratorio de Control Biotecnológico de Plagas, Instituto BIOTECMED, Departamento de
Genética, Universitat de València, 46100 València, Spain.
[2]Instituto Nacional de Tecnología Agropecuaria (INTA), Instituto de Microbiología y Zoología
Agrícola, 1712 Castelar, Argentina.
[3]Universidad Nacional de Villa María. Instituto Académico Pedagógico de Ciencias Básicas y
Aplicadas. Villa María, Argentina.
[4]CONICET-Universidad Nacional de Villa María. Instituto Multidisciplinario de Investigación
y Transferencia Agroalimentaria y Biotecnológica (IMITAB). Villa María, Argentina
[5]ICiagro Litoral, CONICET, Facultad de Ciencias Agrarias, Universidad Nacional del Litoral,
Kreder 2805, Esperanza S3080, Argentina.

*Correspondence: leopoldo.palma@uv.es

Abstract

The *Bacillus* genus comprises physiologically versatile, endospore-forming bacteria
widely distributed in natural environments. In this study, we report the isolation and genomic
characterization of strain Bva_UNVM-123, recovered from agricultural soil in Pergamino,
Argentina. Whole-genome sequencing using Illumina technology yielded a 5.1 Mbp draft
genome assembled in 67 contigs with a GC content of 36%. Comparative genomic analyses
using the TYGS server and digital DNA–DNA hybridization (dDDH) values supported its
classification as a novel species within the genus *Bacillus*. We propose the name *Bacillus
pergaminensis* sp. nov. for this taxon, with strain Bva_UNVM-123 as the type strain. Genome
annotation revealed 4,866 protein-coding genes, including multiple determinants conferring
resistance to antibiotics (e.g., fosfomycin, tetracycline, beta-lactams) and toxic heavy metals
(e.g., arsenic, cadmium, mercury), supporting its potential application in bioremediation.
Additionally, PathogenFinder predicted a low probability of human pathogenicity (0.207),
reinforcing its safety for environmental use. This study adds to the growing knowledge of
environmental *Bacillus* species and their biotechnological potential.

Keywords: *Bacillus* spp.; genomic sequence; antibiotic resistance; heavy metal resistance; soil microbiome; NCBI; genome assembly; Firmicutes; endospore-forming bacteria; bioremediation.

Introduction

Species belonging to the *Bacillus* genus are known for their remarkable natural properties, which range from the synthesis of various toxins (e.g., cereulide) to the production of proteins and enzymes with numerous biotechnological applications, including fermentation, biodegradation, and the production of secondary metabolites. Many of these organisms can synthesize a wide range of bioactive compounds, including antibiotics (e.g., bacitracina), enzymes (amylases, proteases, lipases, etc.), and biopolymers, which can be used in industrial applications [1].

The genes responsible for these biosynthetic pathways are often located on extrachromosomal DNA elements, such as plasmids or bacteriophages, which can harbor additional virulence factors including toxins and antibiotic resistance genes [2].

These plasmids are typically transferable, and through horizontal gene transfer (HGT), they can be transmitted to related bacterial species [3]. Horizontal transfer of genetic material between bacteria plays a crucial role in the spread of antibiotic resistance, as it allows for rapid dissemination of resistance traits across different bacterial populations [4, 5]. This transfer can occur between pathogenic and non-pathogenic bacteria, and in the case of non-pathogenic species, it can result in the presence of antibiotic resistance genes even in bacteria that do not cause disease [6]. Notably, recent studies emphasize the role of non-pathogenic soil bacteria in harboring mobile resistance elements that may be horizontally transferred to clinical pathogens under environmental stress conditions. Such non-pathogenic bacteria, isolated from various ecosystems like soil, water, or animal manure, are of great interest because they can serve as reservoirs for resistance genes that may eventually be transferred to pathogenic organisms under selective pressure [7].

Antibiotic resistance genes are essential mechanisms used by pathogenic bacteria to evade the effects of antibiotics, a challenge that has become a major global health concern. However, they are not only found in pathogenic bacteria but are also commonly detected in non-pathogenic environmental microorganisms, including species used in biotechnology and environmental applications [8]. This characteristic is particularly interesting from an environmental management perspective because non-pathogenic, antibiotic-resistant bacteria could potentially be utilized in bioremediation strategies. For example, certain *Bacillus* strains with intrinsic resistance to antibiotics can be employed in bioremediation efforts to clean up environments contaminated by antibiotics, particularly those originating from hospitals, dairy farms, and livestock operations, where residues of antibiotics often persist in soil and water [9].

This provides a valuable opportunity for using these bacteria in the development of new

biotechnological tools aimed at mitigating the environmental impact of antibiotic

contamination, which is a growing concern in public health and environmental science [10].

In addition to antibiotic resistance, several Bacillus species are known to tolerate or resist

toxic heavy metals such as arsenic, cadmium, lead, copper, and mercury. This resistance is

often mediated by efflux systems, metal-binding proteins, and enzymatic detoxification

pathways encoded in their genomes [11, 12]. These properties make them promising candidates

for bioremediation strategies aimed at decontaminating polluted environments. The capacity to

thrive in metal-contaminated soils, together with their ability to form resilient endospores,

enhances their potential in long-term or large-scale remediation projects [13].

The capacity of *Bacillus* species to adapt to diverse environments, combined with their

resistance to antibiotics and heavy metals, makes them ideal candidates for further research in

the field of bioremediation. By leveraging their natural resistance mechanisms, it may be

possible to engineer more efficient systems for cleaning up contaminated environments,

particularly those impacted by human activity.

In this work, we aimed to sequence and analyze the genome of strain Bva_UNVM-123, a

newly isolated *Bacillus* strain from soil, to explore its taxonomic identity and biotechnological

potential. The assembled draft genome sequence of Bva_UNVM-123 provides insights into its

genomic makeup, revealing potential resistance genes against antibiotics and heavy metals,

which may contribute to its potential as a bioremediation agent. This study will contribute to a

deeper understanding of the genetic factors underlying the resistance of Bacillus species,

furthering their application in environmental remediation efforts.

Materials and Methods

Isolation and Morphological Characterization

The strain Bva_UNVM-123 was isolated from a soil sample collected from soybean field

in Pergamino, Buenos Aires Province (Argentina), as part of a screening program aimed at

identifying entomopathogenic bacteria (e.g., *Bacillus thuringiensis*) as previously described

[14].

Those strains showing a *B. thuringiensis*-like (matte white and with uneven borders)

phenotype were reisolated in axenic cultures and observed by phase contrast microscopy

(Nikon Eclipse Ti, Japan). Gram staining was performed using a commercial kit (Biopack,

Buenos Aires, Argentina), following the manufacturer's instructions. In addition, the strain was

also examined using Scanning Electron Microscopy (SEM) at the Comprehensive Center for

Electron Microscopy (CONICET-Universidad Nacional de Tucumán, Argentina). The strain is
maintained as glycerol suspension (50% v/v) at −80 °C. The strain Bva_UNVM-123 was also
deposited in the International Culture Collection of the Institute of Microbiology and
Agricultural Zoology at INTA Castelar (IMYZA-INTA) (access number pending).

Biochemical Characterization and Antibiotic Resistance Analyses

NaCl tolerance was evaluated by culturing the strain on TSA medium supplemented with
NaCl concentrations ranging from 2.5% to 7.5% (w/v) and incubating for 72 hours. Anaerobic
growth was assessed using the candle jar technique. Motility was tested on semisolid agar
medium (0.75% w/v). Hemolytic activity was tested on blood agar plates supplemented with
5% sheep blood (Britania Laboratories, Argentina) incubated at 37 °C for 48 hours.

The sensitivity of strain Bva_UNVM-123 to various antibiotics, chloramphenicol (30 µg),
novobiocin (5 µg), ampicillin (10 µg), tetracycline (30 µg), and penicillin (10 U) (Britania
Laboratories, Argentina), was assessed on Mueller–Hinton agar using the disc diffusion
method. Oxidase and catalase activities were determined using standard microbiological
procedures. Additional physiological and biochemical characterizations were conducted using
the API 50CHB, API ZYM, and API 20A systems (bioMérieux), following the protocols
provided by the manufacturer.

Fatty acid profiling of the strain was performed using the MIDI Sherlock Microbial
Identification System (version 6.2B) following the methodology described by Ferrari et al.,
(2018). The sample was analyzed using the RTSBA6 method, and fatty acids were identified
based on their equivalent chain length (ECL) and retention time [15].

DNA isolation, Genome assembly and Annotation and Species Delimitation

For DNA extraction, strain Bva_UNVM-123 was grown in LB broth for 16 hours. Genomic
DNA, including chromosomal and plasmid DNA, was extracted using the Wizard Genomic
DNA Purification Kit (Promega, Madison, WI, USA), following the manufacturer's protocol
for gram-positive bacteria. The extracted DNA was analyzed by electrophoresis in 1% agarose
gels, stained with SYBR Safe (ThermoFisher Scientific, Waltham, MA, USA), and quantified
using a PICODROP PICO 100 µL spectrophotometer.

The extracted DNA was subjected to PCR to confirm its belonging to the *Bacillus* genus
[16]. The PCR assay was performed using the methodology described by Wu et al. (2006),
which involves a group-specific primer set designed for Bacillus species identification.

Then, the purified DNA was used to construct a pooled Illumina sequencing library, which
was sequenced at the Genomics Unit of the National Institute of Agricultural Technology

(INTA, Argentina) using Illumina high-throughput sequencing technology. Raw sequencing
reads were processed to remove low-quality regions and assembled into contigs using Velvet
within Geneious version R11 software (www.geneious.com, accessed on 10 November 2023)
with default parameters for de novo assembly. Assembly statistics were evaluated using
QUAST v4.5 [17], and genome completeness and contamination were assessed using CheckM
[18]. Gene predictions and functional annotations were done via PGAP [16] and validated
through the RAST server [19].

Genome annotation was performed using the NCBI Prokaryotic Genome Annotation
Pipeline (2023 release) and the RAST server [19]. Species delimitation was carried out using
the Type (Strain) Genome Server [20] and PubMLST [21]. Secondary metabolites searches
were performed using antiSmash server [22].

Results

Morphological Characterization

The strain formed opaque colonies with a matte texture and creamy white color after
overnight growth on TSA medium at 37 °C during 72 h. Microscopic examination revealed
rod-shaped vegetative cells with terminal deforming spores (Figure 1A and B). The terminal
positioning of the endospore is consistent with known morphological traits in environmental
Bacillus species [23].

Figure 1. Microscopic analysis of strain Bva_UNVM-123 grown on TSA medium (72 h/37
°C). A) Morphological characterization using phase-contrast microscopy and B) Scanning
electron microscopy. A scale bar (200 nm) was included in SEM images to reflect structural
dimensions.

Biochemical Characterization

Cells are Gram-stain-positive, facultatively anaerobic, non-motile, and form terminal
endospores. The strain was able to grow on nutrient agar (NA), casein–collagen–yeast extract
agar (CCY), tryptic soy agar (TSA), and brain heart infusion agar. Optimal growth was
observed on TSA at 37 °C and the strain was non-hemolytic. The strain tested positive for both

catalase and oxidase activities and when cultured in the presence of chloramphenicol, did not exhibit measurable inhibition zones.

Additional physiological and enzymatic tests were carried out using the API 50CHB, API ZYM, and API 20A systems. However, the results obtained were inconclusive due to limited growth under assay conditions, including incubation at 30 °C and 37 °C.

The fatty acid profile of strain Bva_UNVM-123 consisted primarily of branched-chain saturated fatty acids, a chemotaxonomic trait typical of gram-positive bacteria, particularly members of the genus Bacillus or related taxa. The major components included iso- and anteiso-fatty acids such as iso-C15:0, anteiso-C15:0, iso-C16:0, and iso-C17:0. A total of 61.21% of the detected fatty acids were identified, with iso-C15:0 representing the most abundant component (32.46%). The complete fatty acid profile is presented in Table 1.

Table 1. Key Major fatty acid components of strain Bva_UNVM-123. Only those representing ≥3% of the total identified fatty acids are shown. Summed Feature 4 includes fatty acids that could not be resolved individually by the system.

Fatty Acid	Composition (%)	Other Characteristics
iso-C15:0	32.46	Most abundant component
anteiso-C15:0	7.28	-
iso-C16:0	5.21	-
C16:0	4.39	Also known as palmitic acid
iso-C17:0	6.16	-
anteiso-C17:0	3.69	-
Summed Feature 4	5.20	Includes C17:1 iso I / anteiso B isomers

Genome Characterization and Species Identification

The Illumina sequencing produced 14,532,725 high-quality paired end reads, resulting in 67 assembled contigs with a total genome size of 5,119,019 bp in size and 36% G+C content (Table 2).

Table 2. Key features statistics and quality analysis of the draft genome assembly of Bva_UNVM-123 strain obtained with QUAST [17], CheckM [18] and the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) [24].

Feature statistics	Value
Assembled Genome Size (bp)	5,119,099
Contig Number	67
Largest contig (bp)	583,230
N50 (bp)	341,242
N's	5074
GC content (%)	36
Suspected Contamination (%)*	3.85
Estimated Completeness (%)*	95.85
Estimated Coverage (×)	400
PGAP Predicted Genes	5139
PGAP Predicted Protein Coding Genes	4866

*CheckM determinations.

Table 3 highlights multiple genes potentially related to antibiotic resistance, identified
using the RAST annotation server. The presence of genes conferring resistance to fosfomycin
(fosB), tetracyclines, chloramphenicol, fosmidomycin, kanamycin, oxetanocin A, and various
beta-lactams (e.g., beta-lactamase class A-like and penicillin-binding proteins) suggests that
strain Bva_UNVM-123 possesses a broad resistance profile.

Table 3. Putative annotated genes related to antibiotic resistance found at the genome sequence
of Bva_UNVM-123 strain using RAST server [19].

Product	Type	Copies
Efflux ABC transporter (ATP-binding/permease)	Multidrug resistance	45
Fosmidomycin resistance protein	Fosfomycin resistance	2
Similar to tetracycline resistance protein	Tetracyclin resistance	1
Similar to tetracycline resistance, MFS efflux pump	Tetracyclin resistance	1
Chloramphenicol/florfenicol resistance efflux pump	Chloramphenicol resistance	1
Zwittermicin A resistance protein	Zwittermicin A resistance	1
Kanamycin kinase	Kanamycin resistance	1

Oxetanocin A resistance protein	Oxetanocin A resistance	1
Beta-lactamase (Class A/C-like)	Beta-lactam resistance	12
Vancomycin response proteins (VanR/VanS)	Glycopeptide resistance	5
Penicillin-binding protein 1A/1B	Penicillin resistance	1
Penicillin acylase	Penicillin resistance	1
Penicillin-binding protein 4B	Penicillin resistance	1

Table 4 lists genes potentially associated with resistance to toxic heavy metals, including
arsenic, cadmium, zinc, cobalt, lead, mercury, and tellurite. These genes encode various efflux
pumps, ATPases, and regulatory proteins, such as TehB, ArsD, ACR3, and CzcD, which are
known to facilitate resistance and detoxification [25, 26].

Table 4. Putative heavy-metal resistance genes found at the genome sequence of Bva_UNVM-
123 strain using RAST server.

Product	Metal	Copies
Tellurium resistance proteins (TerD, TerB)	Tellurium	5
Arsenical resistance proteins (Acr3, ArsR, ATPase)	Arsenic	9
Cadmium/Zinc/Cobalt transporters (P-type ATPase, CzcD)	Cd/Zn/Co	7
Copper chaperones and transporters (CopZ, CopA)	Copper	4
Zinc ABC transporter systems (ZnuA/B/C, Zur)	Zinc	5
Manganese efflux pump (MntP)	Manganese	1
Lead, Mercury, and other heavy-metal transporters	Pb/Hg/Cd/Zn	6
Multicopper oxidases and polyphenol oxidases	Copper	1
Aluminum resistance protein	Aluminum	1

Concerning species identification, preliminary PCR analysis showed that isolate 123
belongs to the genus *Bacillus* (Figure 2).

Figure 2. PCR analysis of strain Bva_UNVM-123 for the Bacillus genus showing a positive
PCR amplicon (positive control: *B. thuringiensis* HD1, 123: Bva_UNVM-123 strain).

The low average branch support observed in the GBDP phylogenetic tree generated by the
TYGS server likely reflects the high genomic divergence of strain Bva_UNVM-123 and the
lack of closely related reference genomes. Despite this, digital DNA–DNA hybridization
(dDDH) values support its classification as a potentially novel species (Figure 3).

Figure 3. Genome-based phylogenetic tree of Bacillus sp. strain Bva_UNVM-123 and related
taxa, constructed using the GBDP method via the TYGS server. Branch lengths are scaled in
terms of GBDP distance. Bootstrap values indicate the average support for each branch, based
on 100 pseudoreplicates.

Based on PathogenFinder 1.1, the strain was predicted as non-pathogenic, with a human
pathogenicity probability of 0.207, indicating its safety for environmental applications [27].

Discussion

After a comprehensive genomic and phenotypic analyses, including phylogenomic
reconstruction and digital DNA-DNA hybridization (dDDH) values below the 70% species
threshold, we propose that strain Bva_UNVM-123 represents a novel species within the genus
Bacillus, for which the name *Bacillus pergaminensis* sp. nov. is proposed, with strain
Bva_UNVM-123 as the type strain.

The genome of *B. pergaminensis* sp. nov. Bva_UNVM-123 reveals multiple adaptive traits
that support its role in bioremediation. In line with this, preliminary phenotypic assays showed
that strain Bva_UNVM-123 exhibited no measurable inhibition zones against chloramphenicol
and novobiocin, suggesting intrinsic tolerance that may correspond to the resistance genes
found in the genome. Nonetheless, further MIC assays are necessary to validate this resistance.

In addition, the presence of antibiotic resistance genes such as fosB, tetA, bla-like beta-
lactamases, and multidrug efflux systems suggests long-term exposure to antibiotic residues in
soil environments. This aligns with findings that environmental *Bacillus* species often act as
reservoirs for antibiotic resistance genes due to selective pressure in natural and agricultural
soils [6, 7]. This resistance could be determinant and particularly relevant for bioremediation
applications, where bacteria must survive in antibiotic-contaminated environments.

The use of API-based systems for metabolic profiling did not yield conclusive results. This
outcome is frequently observed with environmental isolates, which often exhibit poor growth
or atypical metabolic patterns when tested in commercial platforms originally designed for
clinical or fast-growing laboratory strains. These limitations highlight the need to optimize test
conditions or employ alternative phenotypic strategies when characterizing environmentally
adapted bacteria. For instance, even in industrial contexts such as dairy processing
environments, phenotypic microbiological techniques have shown limited reliability for
environmental strains like *Pseudomonas* spp. and *Bacillus* spp., often resulting in inconclusive
species-level identifications [28]. Consequently, in this study, species identification relied
primarily on genome-based approaches, which provided higher taxonomic resolution and
reliability.

Additionally, the fatty acid profile of strain Bva_UNVM-123 did not match any entry in
the RTSBA6 reference library, which is consistent with the genomic evidence suggesting
novelty. However, the total response signal obtained in both replicates was below the
recommended threshold for reliable identification. This limitation was most likely due to the
poor growth of the strain under the conditions used for biomass collection, rather than to
methodological issues. Further optimization of culture conditions is therefore needed to obtain
sufficient biomass for accurate chemotaxonomic profiling.

Moreover, the strain harbors a diverse array of heavy metal resistance genes, including
arsD, acr3, czcD, and tehB, which confer tolerance to arsenic, cadmium, mercury, and tellurite.
These genes are commonly found in metal-contaminated environments and are considered key
biomarkers for bacterial adaptation and survival in such niches [11, 25, 26]. The abundance
and diversity of these genes suggest that strain Bva_UNVM-123 is well adapted to metal-
contaminated environments. This further supports its potential use in bioremediation,
particularly in soils impacted by industrial or agricultural pollutants.

The non-pathogenic nature of the strain, as predicted by PathogenFinder, along with the
absence of virulence determinants and hemolytic activity, strongly suggests its suitability for

environmental application. This safety profile is crucial for any candidate microorganism
intended for bioremediation purposes, as environmental deployment requires strict biosafety
assessments [27].

Importantly, the genome of *B. pergaminensis* sp. nov. also provides a valuable resource for
future biotechnological exploration. Its genetic repertoire could be exploited not only for
bioremediation of antibiotic- and metal-polluted sites but also for the potential production of
industrial enzymes or bioactive secondary metabolites, as often observed in *Bacillus* spp. [1,
22].

Author contributions

274 Conceptualization, L.P., methodology, D.H.S, C.P and L.P.; formal analysis, D.H.S, C.P, V.F.
275 and L.P.; investigation, D.H.S; C.P, V.F., E.E.D.V and L.P.; resources, L.P.; data curation, L.P.;
276 writing—original draft preparation, L.P.; writing, review and editing, L.P. and D.H.S;
277 visualization, L.P.; supervision, L.P.; project administration, L.P.; funding acquisition, L.P. All
278 authors have read and agreed to the published version of the manuscript.

Conflicts of interest

280 The authors declare there are no conflicts of interest.

Funding information

282 This research was funded by Ministerio de Ciencia, Tecnología e Innovación (FONCYT), Plan
283 Argentina Innovadora 2020, grant number PICT 2017-0087, by the Consejo Nacional de
284 Investigaciones Científicas y Técnicas (CONICET), grant number PIP 2017-2019 GRUPO
285 INVES and the Instituto de Investigación of the Universidad Nacional de Villa María, grant
286 number PIC 2017-2019.

Data availability

288 The unprocessed genome sequencing data have been deposited in the NCBI database under the
289 BioSample accession SAMN18815149, within the BioProject PRJNA723377. The
290 corresponding genome assembly is accessible through the NCBI Whole Genome Shotgun
291 (WGS) database under the accession number JAGTPU000000000.

Acknowledgements

293 Leopoldo Palma gratefully acknowledges the Spanish Ministry of Science, Innovation, and
294 Universities, the Spanish State Research Agency, and the European Union for funding his
295 Ramón y Cajal contract (grant ref. RYC2023-043507-I).

References

1. **Herrmann LW, Letti LAJ, de Oliveira Penha R, Soccol VT, Rodrigues C et al.** *Bacillus* genus industrial applications and innovation: First steps towards a circular bioeconomy. *Biotechnology Advances* 2024;70:108300.

2. **Bennett PM**. Plasmid encoded antibiotic resistance: acquisition and transfer of antibiotic resistance genes in bacteria. *British journal of pharmacology* 2008;153(S1):S347–S357.

3. **Sauka DH, Del Valle EE, Palma L**. *Bacillus toyonensis* biovar Thuringiensis: an overlooked entomopathogen? *bioRxiv* 2024:2024.2007. 2021.604280.

4. **Partridge SR, Kwong SM, Firth N, Jensen SO**. Mobile genetic elements associated with antimicrobial resistance. *Clinical microbiology reviews* 2018;31(4):10.1128/cmr. 00088–00017.

5. **Kaplan T**. The role of horizontal gene transfer in antibiotic resistance. *Eukaryon* 2014;10:80–81.

6. **Martínez JL**. Antibiotics and antibiotic resistance genes in natural environments. *Science* 2008;321(5887):365–367.

7. **Larsson DJ, Flach C-F**. Antibiotic resistance in the environment. *Nature Reviews Microbiology* 2022;20(5):257–269.

8. **Amos GC**. *Environmental reservoirs of antibiotic resistance*. University of Warwick; 2013.

9. **Gul S, Hussain S, Khan H, Khan KI, Khan S et al.** Chapter 4 - Advances in bioremediation of antibiotic pollution in the environment. In: Kumar S, Hashmi MZ (editors). *Biological Approaches to Controlling Pollutants*: Woodhead Publishing; 2022. pp. 49–78.

10. **Apreja M, Sharma A, Balda S, Kataria K, Capalash N et al.** Antibiotic residues in environment: antimicrobial resistance development, ecological risks, and bioremediation. *Environmental Science and Pollution Research* 2022:1–17.

11. **Margaryan A, Panosyan H, Birkeland N-K**. Heavy metal resistance in prokaryotes: Mechanism and application. *Microbial communities and their interactions in the extreme environment* 2021:273–313.

12. **Prabhakaran P, Ashraf MA, Aqma WS**. Microbial stress response to heavy metals in the environment. *Rsc Advances* 2016;6(111):109862–109877.

13.   **Singh R**. Microorganism as a tool of bioremediation technology for cleaning

environment: a review. *Proceedings of the International Academy of Ecology and*

Environmental Sciences 2014;4(1):1.

14.   **Iriarte J, Bel Y, Ferrandis MD, Andrew R, Murillo J et al.** Environmental

distribution and diversity of *Bacillus thuringiensis* in Spain. *Syst Appl Microbiol*

1998;21(1):97–106.

15.   **Ferrari AE, Ravnskov S, Wall LG**. Crop rotation in no-till soils modifies the soil

fatty acids signature. *Soil use and management* 2018;34(3):427–436.

16.   **Wu X-Y, Walker MJ, Hornitzky M, Chin J**. Development of a group-specific PCR

combined with ARDRA for the identification of *Bacillus* species of environmental

significance. *Journal of Microbiological Methods* 2006;64(1):107–119.

17.   **Gurevich A, Saveliev V, Vyahhi N, Tesler G**. QUAST: quality assessment tool for

genome assemblies. *Bioinformatics* 2013;29(8):1072–1075.

18.   **Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW**. CheckM:

assessing the quality of microbial genomes recovered from isolates, single cells, and

metagenomes. *Genome Res* 2015;25(7):1043–1055.

19.   **Aziz RK, Bartels D, Best AA, DeJongh M, Disz T et al.** The RAST Server: rapid

annotations using subsystems technology. *BMC Genom* 2008;9(75):1–15.

20.   **Meier-Kolthoff JP, Goker M**. TYGS is an automated high-throughput platform for

state-of-the-art genome-based taxonomy. *Nat Commun* 2019;10(1):1–10.

21.   **Jolley KA, Bray JE, Maiden MCJ**. Open-access bacterial population genomics:

BIGSdb software, the PubMLST.org website and their applications. *Wellcome Open Res*

2018;3(124):2–20.

22.   **Blin K, Pascal Andreu V, de Los Santos ELC, Del Carratore F, Lee SY et al.** The

antiSMASH database version 2: a comprehensive resource on secondary metabolite

biosynthetic gene clusters. *Nucleic Acids Res* 2019;47(D1):D625–D630.

23.   **Vos P, Garrity G, Jones D, Krieg NR, Ludwig W et al.** *Bergey's manual of*

systematic bacteriology: Volume 3: The Firmicutes: New York: Springer New York, 2009.;

2011.

24.   **Tatusova T, DiCuccio M, Badretdin A, Chetvernin V, Nawrocki EP et al.** NCBI

prokaryotic genome annotation pipeline. *Nucleic acids research* 2016;44(14):6614–6624.

25.   **Nies DH**. Efflux-mediated heavy metal resistance in prokaryotes. *FEMS microbiology*

reviews 2003;27(2-3):313–339.

26.   **Bruins MR, Kapil S, Oehme FW**. Microbial resistance to metals in the environment.

Ecotoxicology and environmental safety 2000;45(3):198–207.

27.   **Cosentino S, Voldby Larsen M, Møller Aarestrup F, Lund O**. PathogenFinder-

distinguishing friend from foe using bacterial whole genome sequence data. *PloS one*

2013;8(10):e77302.

28.   **Wiedmann M, Weilmeier D, Dineen Sean S, Ralyea R, Boor Kathryn J**.

Molecular and Phenotypic Characterization of *Pseudomonas* spp. Isolated from Milk. *Applied*

and Environmental Microbiology 2000;66(5):2085–2095.

369