**Proteome analyses reveal Endoplasmic Reticulum stress-induced changes in**

**protein abundance associated with Ube2j2 deficiency in human cell culture**

Dahlberg, C. L.[1,2,*], Zinkgraf, M.[2,*], Laugesen, S.H.[1], Søltoft, C.L.[1], Ginebra, Q.[1], Bennett, E.P.[3],

Hartmann-Petersen, R.[1], and Ellgaard, L.[1,*]

Affiliations:

[1] Department of Biology, Linderstrøm-Lang Centre for Protein Science, University of Copenhagen,

Copenhagen 2200, Denmark

[2] Department of Biology, Western Washington University, Bellingham, Washington, USA, 98225

[3] Genome Editing Department, Nucleic Acid Research, Novo Nordisk A/S, Måløv 2760, Denmakr

*Corresponding authors: Dahlberg, C. (dahlbec@wwu.edu), Zinkgraf, M. (zinkgrm@wwu.edu),

Ellgaard, L. (lellgaard@bio.ku.dk)

Running title: Proteomic analysis of cells lacking Ube2j2

Keywords: Ube2j2, unfolded protein response, ER-associated degradation, ubiquitin proteasome

system, protein degradation, E2 ubiquitin-conjugating enzyme, proteomics

## Abstract

The unfolded protein response (UPR) helps reinstate cellular proteostasis upon an accumulation of misfolded proteins in the endoplasmic reticulum (ER), in part through ER-associated degradation (ERAD). Ube2j2 is an ER-localized E2 ubiquitin-conjugating enzyme that participates in ERAD. We used mass spectrometry analysis of cultured U2OS cells to investigate how the loss of Ube2j2 affects the cellular proteome in response to tunicamycin-induced ER stress. We constructed a network of twelve statistically distinct modules of protein abundance profiles across conditions. We describe the Gene Ontology annotations for each module along with the "hub gene" proteins whose abundance levels most closely adhere to each module's protein abundance profile. Our analysis identifies known Ube2j2-associated pathways (e.g., the UPR and ERAD) and cellular functions that were previously unassociated with Ube2j2 (e.g., RNA metabolism, ER-Golgi transport, and cell-cycle progression). These data are available via ProteomeXchange with identifier PXD076153 and provide avenues for further investigation into the cellular functions of Ube2j2 under basal and ER-stressed conditions.

## Introduction

Protein ubiquitylation plays a central role in cell physiology by regulating a vast number of protein activities. The ubiquitin-proteasome system is well-established as a central axis for regulating targeted protein abundance, and it is also crucial in maintaining protein homeostasis (proteostasis) in conjunction with molecular chaperones. The general structure of interactions in the "ubiquitin cascade" utilizes increasing specificity to ensure that substrates are ubiquitylated properly. In mammalian cells, two E1 ubiquitin-activating enzymes are responsible for the ATP-dependent charging of ubiquitin, which is then transferred to one of approximately 40 E2 ubiquitin-conjugating enzymes (E2s) (van Wijk and Timmers 2010; Parashar et al. 2025). Ubiquitin-charged E2s interact directly with unique combinations of the hundreds of E3 ubiquitin-protein ligases (E3s) to transfer ubiquitin onto substrate molecules (Ye and Rape 2009; Stewart et al. 2016; Gundogdu and Walden 2019). Ubiquitylation can also be reversed by deubiquitylating enzymes (DUBs) (Fischer 2003; Sowa et al. 2009). Depending on the number and pattern of ubiquitin modifications, substrate proteins may be re-localized, activated, repressed, or degraded. Thus, specific cohorts of E2 conjugases and E3 ligases can have wide-ranging effects on cellular pathways.

E2 enzymes provide an initial element of subcellular and temporal regulation to the ubiquitylation cascade through spatial restriction (van Wijk et al. 2009; Ye and Rape 2009; van Wijk and Timmers 2010; Witus et al. 2021). E2s with specific subcellular localizations like the endoplasmic reticulum (ER), cytosol, nucleus, or mitochondria, are positioned to interact with E3 ligases to facilitate ubiquitylation of more narrowly defined classes or groups of proteins (Stewart et al. 2016; Christianson and Carvalho 2022). Nonetheless, some E3-containing complexes, such as those that function in ER-associated degradation (ERAD), target diverse groups of substrates (Carvalho et al. 2006; Ravid et al. 2006; Sato et al. 2009; Sharninghausen et al. 2024) despite being spatially constrained.

The ERAD machinery is transcriptionally controlled downstream of the Unfolded Protein Response (UPR), which coordinates transcriptional and translational responses to proteotoxic stress caused by e.g., protein misfolding and aggregation, hypoxia, and oxidative stress (reviewed in Hetz et al., 2015). The overall aim of the UPR is to reinstate cellular homeostasis and, should this fail, initiate apoptosis. ERAD complexes identify and remove misfolded proteins from the ER lumen and membrane and target them for degradation by the proteasome (reviewed in Christianson et al., 2023; Christianson

and Carvalho, 2022). This process is dependent on an expanding suite of recognized E3 ligases that
partner with ER-localized E2 conjugases.

The ER-localized E2 enzymes Ubc6 and Ubc7 were originally identified in yeast as critical members
of the ERAD machinery, and the mammalian orthologs, Ube2j1/Ube2j2 and Ube2g1/Ube2g2 were
subsequently identified (Lenk et al. 2002; Wang et al. 2009; Abdul Rehman et al. 2024; Swarnkar et
al. 2024). Ubc6 and Ubc7 facilitate E3-dependent ubiquitylation by interacting with the E3s Doa10
and Hrd1, respectively (Kreft and Hochstrasser 2011; Zattas and Hochstrasser 2015; Lips et al. 2020;
Mehrtash and Hochstrasser 2022; Christianson et al. 2023). In addition, they participate in sequential
ubiquitylation, whereby Ubc6 first catalyzes the addition of ubiquitin to an [atypical] hydroxylated
amino acid (Ser or Thr) and Ubc7 then extends a ubiquitin chain via [typical] lysine modification
(Weber et al. 2016). Ube2j2, in particular, displays a preference for hydroxylated amino acids (Wang
et al. 2009; Abdul Rehman et al. 2024; Swarnkar et al. 2024), whereas Ube2j1 can utilize
hydroxylated residues *in vivo* (Burr et al. 2013). Also similar to yeast, Ube2j2 and Ube2g2 interact
with distinct ER-localized E3 ligases, including MARCH6 and HRD1, the mammalian orthologs to
Doa10 and Hrd1, respectively, along with RNF139 and RNF145 (reviewed in Christianson and
Carvalho 2022; Vrentzou et al. 2025). The conserved partnership between ERAD E2s and E3s is an
indicator of their central role in cellular function.

Human Ube2j1 and Ube2j2 are similar (43% amino acid sequence similarity, 27% identity), and their
similar (though not identical) biochemical activity suggests overlapping functions in the cell.
However, investigations into the activities and targets of Ube2j2 in mammalian cells highlight its
unique targets and physiological roles (Younger et al. 2006; van den Boomen et al. 2014; van de
Weijer et al. 2014; Glaeser et al. 2018; Liu et al. 2019; Cremer et al. 2021; Wolf et al. 2021; Yu et al.
2025). Recently, Ube2j2 has also been shown to contribute to progression and drug resistance in
cancer models (Kumari and Kumar 2023; Nakao et al. 2023; Lin et al. 2024). Thus, Ube2j2 plays a
distinct role in mammalian protein homeostasis and cell physiology.

To uncover cellular pathways associated with Ube2j2, we first used mass spectrometry to quantify
the proteomes of Ube2j2 CRISPR-derived knockout cells under control and tunicamycin-induced ER
stress conditions. We then analyzed the proteomic data using a computational network-based
procedure that combined the *weighted gene co-expression analysis* (WGCNA) (Langfelder and

Horvath, 2008) and differentially expressed proteins (DEP) (Robinson et al. 2010) approaches to
model changes in protein abundance and hereby provide a systems-level view of molecular
differences between treatments (Serin et al. 2016; Vella et al. 2017). Thus, we were able to (1) identify
sets of proteins (modules) with similar abundance profiles across experimental conditions, (2) define
module eigengenes (representative profiles that summarize the behavior of any given module,
calculated as the first principal component of the protein abundance data) and relate these to
experimental treatments, (3) annotate the modules using Gene Ontology (GO) enrichment analysis to
extract biological meaning (Ashburner et al. 2000), and (4) find key drivers in the different modules
(so-called hub genes) by identifying highly connected proteins that had strong significance in the
DEP analysis. To avoid confusion with the term "hub protein" (a central protein in a protein-protein
interaction network), we here employ the term "hub gene", which originates from the WGCNA
approach, even though the present analysis is applied to protein abundance. Such hub genes provide
potential targets for future study into the biological processes associated with Ub2j2 expression and/or
ER stress.

## Results

We first used CRISPR-Cas to create a cell line of U2OS cells, C6β (denoted KO or C6 in raw data sets), lacking a region of 432 base pairs spanning the 5′ UTR and first coding exon of the *UBE2J2* gene and not displaying any visible Ube2j2 expression as probed by Immunoblotting (**Figure S1**). Then, to gain insight into changes in protein abundance as a result of ER-stress duration and the presence or absence of Ube2j2, we grew (in quadruplicate) wild-type and Ube2j2 KO cells in either dimethyl sulfoxide (DMSO; vehicle control) or 10 μg/mL tunicamycin – a cell permeable agent that prevents N-glycosylation in the ER and thereby leads to protein misfolding and activates the UPR – for 6 or 18 hours. Upon flash-freezing, the cells were then processed for and analyzed by quantitative mass spectrometry.

The resulting 32 data sets contained 5903 unique protein sequences, and the quality of samples was gauged based on the coefficient of variation of replicates, which ranged from 11.24 to 18.77% within a sample group. Samples with a coefficient of variation of 10-20% are generally considered reproducible for downstream analyses (Cho et al. 2021). The WGCNA approach defined 12 statistically separate protein abundance profiles each comprising 200-1029 proteins (**Figure 1a, Supplementary File 1**). These 12 modules were annotated to describe the expression and functional properties of the proteins within each module and their association to experimental factors. As visualized by the eigengene values for each module (**Figure 1b**), variation in protein levels across genotypes was the primary factor in defining modules, followed by tunicamycin treatment, and to a lesser extent time (**Figure 1c, Supplementary File 2**). Eigengenes represent the absolute value of changes in abundance, so the magnitude of the change, rather than the direction of change, in protein abundance is reflected in the eigengene pattern. The identified modules also had significant enrichment for specific GO categories in biological processes, cellular components and to a lesser extent molecular functions (**Supplementary File 3**), allowing additional insight into the functional properties of modules.

Four pairwise contrasts were selected to understand the effects that tunicamycin treatment has on the Ube2j2 KO cells: KO_Tu_18hr-vs-WT_Tu_18hr; KO_Crtl_18hr-vs-WT_Ctrl_18hr; KO_ Crtl _6hr-vs-KO_Ctrl_18hr; WT_ Crtl _6hr-vs-WT_ Crtl _18hr. These four comparisons were chosen for their potential to provide insight into differences in protein abundance based on changes of a single variable at a time (genotype, drug treatment, or time in culture). We chose to focus on treatment conditions at

18 hours, since this late timepoint was expected to provide maximal changes in protein abundance.
Further dissection of the modules showed that 9 out of the 12 modules were associated with at least
one of these selected pairwise contrasts of interest (**Figure 1d, Supplementary File 2**). Below, we
focus on four modules or groups of modules whose patterns of protein abundance levels support our
current understanding of the functions of Ube2j2 in protein homeostasis at the ER and offer new
insights into cellular protein networks associated with the loss of Ube2j2 and/or ER stress.

**Pink module: Response to tunicamycin is not affected by loss of Ube2j2**
The pink module includes proteins whose abundance is affected strongly by tunicamycin-induced
stress but not by cellular genotype. The eigengene for the pink module shows stable abundance over
time and between the wild-type and knockout cells (**Figure 1b**). However, treatment with
tunicamycin leads to a strong up or down regulation of these proteins, relative to the control,
regardless of genetic background. Proteins included in the pink module represent well-characterized
members of ER-stress pathways such as HERP, HSPA5 (BiP), and VIMP (SelS) (**Figure 2a-c**), all
of which also showed clear upregulation in response to tunicamycin via immunoblotting (**Figure 2d**).
In this manner, the pink module provides an important validation of our mass spectrometry data.
**Figure 2e** shows the distribution of proteins in the pink module annotated by biological process
ontology. These are enriched for protein processing, folding, and homeostasis as well as protein
synthesis and metabolism. Processes related to cytokinesis and chromosome segregation are also
identified in this module.

**Magenta module: Loss of Ube2j2, regardless of treatment with tunicamycin, drives changes in**
**protein abundance**
The magenta module includes 1029 proteins whose abundance is dependent on the presence or
absence of Ube2j2 (p-value < 0.001), with little-to-no additional effect of tunicamycin or time
(**Figure 1b**). Functional analysis of the magenta module identified significant associations with terms
from the cellular components and biological processes, and to a lesser extent molecular function
ontologies (**Supplementary File 2).** Based on this analysis, the magenta module was enriched with
three main categories (**Figures 3a and 3b**). First, actin-binding proteins, including some that are
involved with vesicular transport and exocytosis; second, regulation of mitotic cell cycle involving
spindle localization and segregation of sister chromatids; and third, processes that are important for
protein abundance regulation and homeostasis. To further narrow down the list of proteins in the

magenta module, we conducted a hub gene analysis and identified 135 proteins that were highly
connected in the magenta module and significantly associated with protein abundance differences
between Ube2j2 KO versus wildtype cell lines (**Figure 3c**). GO enrichment analysis of magenta hub
genes by cellular component (**Figure 3d**) found similar results to the module level analysis and
suggest that hub genes play a critical role in defining the functions associated with actin and the
regulation of the proteasome.  Interestingly, Ube2g2, the closely related E2 enzyme that works in
tandem with Ube2j2 in yeast (Weber et al. 2016; Mehrtash and Hochstrasser 2022) is also included
in this module. This suggests that at least some of the proteins in the magenta module may directly
compensate for the loss of Ube2j2's enzymatic activity. Thus, at least two major cellular components
– actin-based motility and cellular structure, and protein degradation – are influenced by the presence
or absence of Ube2j2, regardless of tunicamycin-induced ER stress.
194    
**Blue and black modules: Loss of Ube2j2 phenocopies treatment with tunicamycin**
The blue and black modules include proteins whose abundance in response to tunicamycin treatment
is phenocopied by the loss of Ube2j2 (**Figure 1b**). Proteins that are members of the black and blue
modules are affiliated with a broad set of biological processes including protein-complex interactions,
translation initiation, and protein and RNA-localization (black module, **Figure 4a**) and vesicular and
ER-to-Golgi transport, electron transport, and nucleotide metabolism (blue module, **Figure 4b**).
201    
While there are no identified hub genes for the black module, the blue module includes 20 hub genes
(**Figure 4c)**. Among these, a cluster of COPI coatomer proteins – COPB1, COPB2, COPG1, and
COPG2 – are striking due to their roles in Golgi-to-ER transport. In addition, protein homeostasis
components including the proteasome adapter and scaffold protein ECM29, UFM1- and SUMO-
activating proteins (UBA5 and SAE2, respectively), and the E3 ubiquitin ligase, listerin, are blue
module hub genes, along with the translation initiation factor EIF2A, which plays a central role in
regulating the PERK branch of the UPR (Harding et al. 2000; Hetz et al. 2015b). Thus, many blue
module hub genes have roles in proteostasis. Only four of the 20 hub genes for the blue module have
significant differences between 18 hours of tunicamycin treatment in wildtype cells and 6 hours of
basal conditions in mutant cells. Overall, we find that the blue and black modules describe proteins
whose abundance in the absence of Ube2j2 is similar to that of ER stress conditions, and for which
additional ER stress does not amplify the effect of the loss of Ube2j2.
214    

**Purple, midnightblue, and brown modules: Effect of time in culture is abrogated in the absence of Ube2j2**

The three related modules purple, midnightblue, and brown describe a pattern wherein time-dependent changes in protein abundance under control conditions are attenuated when Ube2j2 is missing (**Figure 1b**). This could represent a change in cell-cycle progression or disparate reactions to time in culture.

In all three modules, proteins involved with ribosomal metabolism (including biogenesis and rRNA processing) are highly represented (**Figure 5**), suggesting a previously unappreciated role for Ube2j2 in ribosome biosynthesis and regulation. The GO annotations of proteins in the purple, midnightblue, and brown modules suggest that Ube2j2 may also impact protein translation during cellular growth in culture, or potentially during advancement of the cell cycle (**Figure 5**).

Only the brown module has associated hub genes and about one third of these 23 hub genes are directly associated with the ribosomal metabolism described by the annotation of the entire module. This suggests that those proteins will be strong candidates for future research into the role of Ube2j2 in regulating cellular health during culturing and growth.

## Discussion

Here, we engineered a line of U2OS cells lacking Ube2j2 and profiled their proteome alongside wild-type cells under four different conditions. Our analysis resulted in 12 modules that describe patterns of changes in protein abundance across nearly 6000 protein sequences. We identified four modules/groups of modules that are characterized by interactions between cellular genotype, treatment condition, or time in culture. GO annotation of the proteins that most strongly drive the eigengene pattern for these modules (hub genes) show that defined classes of biological functions are associated with the magenta, black and blue modules, and that three closely related modules (brown, midnightblue, and purple) show overlapping representation of the same or similar biological function.

There are, to our knowledge, two other proteomic datasets available for cells lacking wildtype Ube2j2 levels. One investigation showed that differentially downregulated proteins in Ube2j2 knockout testes cells were enriched in gamete-specific proteins (Yu et al. 2025). The other investigation, a large-scale transcriptomic and proteomic study, found that the combined knockdown of Ube2j2 and the E1 ligase UBA1 reduced K48-linked ubiquitylation and resulted in the concomitant downregulation of Ube2D1 and Ube2E3 mRNA levels and Ube2D1 and Ube2E1 protein levels in HEK293T cells (Hunt et al. 2023). The present data set is unique because it probes the response of cells to the stress that Ube2j2 would normally help to alleviate.

Our dataset and analysis add to previously published data sets by combining the complete knockout background with a network-based perspective on changes to the proteome in response to two variables, time and drug-induced stress. Our analyses provide a suite of protein-abundance profiles that reveal both expected and unpredicted changes across the proteome in the absence of Ube2j2. Our results highlight UPR and ubiquitin-dependent protein homeostasis networks (highly represented in the pink module and magenta modules (Figures 2 and 3)). The network analysis also suggests Golgi-to-ER trafficking pathways (magenta, blue, and black modules, Figures 3 and 4) and RNA metabolism pathways (brown, purple, and midnightblue modules, Figure 5) that are dependent on Ube2j2. In addition, we were interested in any changes in abundance of the Ube2j2 paralog, Ube2j1, and the two Ubc7 homologs, Ube2g1 and Ube2g2 in this dataset. Ube2j1 and Ube2g1 are found in the brown module, which makes them members of a cohort of proteins whose reaction to time in culture is modulated by Ube2j2 (Figure 1), while Ube2g2 is a hub gene for the magenta module (Figure 3). These results suggest different types of compensation by ER-associated E2 enzymes in

264    the absence of Ube2j2. Together, the WGCNA analysis, eigengene profiles, and GO annotations that

265    accompany our data sets suggest inroads for future research by providing insights into changes in

266    protein abundance that are based entirely on the presence or absence of Ube2j2 or through complex

267    interactions between Ube2j2 and tunicamycin-induced stress and/or time in culture.

## Materials and methods

**Generation of CRISPR-Cas9–Mediated UBE2J2 knockout cell line**

Human osteosarcoma (U2OS) cells were subjected to two rounds of CRISPR-mediated gene editing. Cells were transiently co-transfected with plasmids encoding Cas9 and single guide (sg) RNAs targeting UBE2J2 (sgRNA#3: 5′-CAGTAAGAGGGCTCCGACCA-3′ and sgRNA#1: 5′-CCGGGTCTTTCTTAATGCGA-3′) using polyethylenimine transfection reagent. sgRNA sequences were designed and kindly provided by Dr. Dick van den Boomen in the lab Prof. Paul J. Lehner, University of Cambridge, UK. After 24 hours, cells were selected with 10 µg/µl Puromycin and 100 µg/µl Hygromycin B to enrich for transfected cells. To enhance selection efficiency and avoid false positives, cells were split three days post-transfection. Selection medium was replaced once non-transfected controls were eliminated. Due to variable sgRNA efficiency and to obtain a homogenous population of cells, single-cell clones were isolated by limiting dilution (0.3 cells/well in 96-well plates), expanded, and harvested.

Genomic DNA was extracted from potential knockout clones using QuickExtract™ DNA Extraction Solution 1.0 (Epicentre), following the manufacturer's instructions. Briefly, ~9,000 cells were pelleted at 500 × g for 5 min, washed in phosphate buffered saline (PBS), and resuspended in 50 µL QuickExtract™. Extracts were incubated at 70°C for 30 minutes with intermittent vortexing, followed by 95°C for 5 min.

The extracted genomic DNA was used as template for PCR amplification prior to TOPO cloning and Sanger sequencing. PCR reactions were performed using the Phusion DNA polymerase (Thermo Fisher Scientific) with a primer pair designed to amplify a region of the gene covering the sgRNA target site (FWD: 5′-GGCCAGGGCATAAGAGTTTATTTAC-3′, REV: 5′-TACTGCTCTGTCGAAACACTGC-3′), yielding an expected amplicon size of 1472 bp. PCR products were verified by agarose gel electrophoresis and subsequently ligated into the pCR™-Blunt II-TOPO™ vector using the Zero Blunt™ TOPO™ PCR Cloning Kit (Thermo Fisher Scientific), following the manufacturer's protocol. The ligation products were transformed into competent *E. coli* DH5α cells, and ten colonies per reaction were picked for overnight culture. Plasmid DNA was isolated using mini-prep procedures and subjected to Sanger sequencing (Eurofins) using the M13

299  reverse -29 primer (5′-CAGGAAACAGCTATGACC-3′). Sequencing analysis revealed a 432 bp

300  deletion in the targeted *UBE2J2* locus (**Supplementary Figure 1**).

**Generation of a monoclonal anti-VIMP antibody**

303  Mouse monoclonal antibodies were raised to human VIMP using a previously described protocol

304  (Hendil 2005). For immunization, we used the His$_6$-lipoFF-tagged cytosolic region (residues 49–189)

305  of human VIMP with the Sec residue at position 188 replaced with a Cys residue. The protein was

306  produced recombinantly as previously described (Christensen et al. 2012).

**Cell culture, tunicamycin treatment, and harvest**

Wild-type or knock-out U2OS cells were grown maintained in Dulbecco's Modified Eagles Medium

supplemented with L-glutamine and heat-treated fetal bovine serum. Cells were passaged regularly,

and experiments were performed with cells that had been passaged fewer than 20 times. For drug

treatment experiments prior to immunoblotting, cells were treated by adding 10 µg/mL tunicamycin

(final concentration) directly to growth media followed by incubation for the indicated times. As a

control, 0.01% DMSO (vehicle control, final concentration) was added to cells, which were then

incubated for the longest incubation time.

**Protein concentration determination, SDS-PAGE, and immunoblotting**

Cells were harvested by placing plates on ice and washing with PBS, followed by lysis on ice in 40

uL radioimmunoprecipitation assay (RIPA) buffer supplemented with phenylmethylsulfonyl fluoride

(0.01%) and cOmpleteMini protease inhibitor mixture (1 tablet per 10 mL of buffer, according to

manufacturer's directions). Cells were scraped off from the plates and sonicated for five seconds on

ice. Cell lysates were centrifuged for 7.5 minutes at 12,500 rpm at 4°C and supernatant was removed

to fresh tubes for analysis. Protein content of cleared lysates was determined by quantitative

Bicinchoninic Acid (BCA, Sigma Aldrich) anlysis in 96-well plate format using bovine serum

albumin (Sigma Aldrich) as standard. All dilutions for the standard curve and sample measurements

were done in triplicates. The BCA was measured at 562 nm using a plate reader (Tecan) and

equivalent portions of lysate were run on acrylamide gels and transferred to polyvinylidene fluoride

for immunoblotting. Immunostaining was performed with the following primary antibodies, which

were diluted in Tris buffered saline plus 0.05% Tween containing 5% non-fat milk or bovine serum

albumin: rabbit anti-BiP/GRP78 (Sigma, GL-19; RRID:AB_477030) (1:1000), rabbit anti-HERP

(kind gift from L. Hendershot) (1:1000), rabbit anti-Ube2j2 (kind gift from L. Hendershot) (1:1000),

mouse anti-VIMP (described above) (1:1000). Primary antibodies were detected with secondary

antibodies conjugated to Horseradish Peroxidase (Pierce) diluted 1:100,000 in Tris buffered saline

plus 0.05% Tween containing 5% non-fat milk. Blots were visualized using Pierce Femto ECL

according to manufacturer's directions.

**Cell treatment, growth and harvest for mass spectrometry**

Wild-type or knockout cells were grown to approximately 50% confluency in 10 cm plates. Cells

were grown as described above and then treated by replacing original growth media with media

supplemented with 0.01% DMSO (vehicle control) or 10 µg/mL tunicamycin in DMSO followed by

incubation for 6 or 18 hours. Media was removed by aspiration and plates were moved onto ice. Cells

were scraped off plates into PBS supplemented with protease inhibitor (cOmpleteMini protease

inhibitor mixture was dissolved in PBS according to manufacturer's direction). Cells were spun

gently (5 min at 2000 rpm, at 4°C) and washed twice with ice cold PBS supplemented with protease

inhibitors. Cell pellets were snap frozen in liquid nitrogen and stored at -80°C.

**Mass spectrometry procedure**

In preparation for mass spectrometry, cell pellets were lysed in lysis buffer (1% sodium deoxycholate,

100 mM Tris-HCl, pH 8.5) and incubated for 10 min at 99°C, followed by sonication in a Bioruptor

Pico (20 cycles, 30 s on/30 s off, ultra-low frequency). Lysates were cleared by centrifugation,

reduced with 5 mM TCEP for 15 min at 55°C, and alkylated with 20 mM chloroacetamide for 30 min

at room temperature. Proteins were digested for 16 h at 37°C using trypsin and LysC at a 1:100

(enzyme:protein) ratio. Peptides were acidified with 1% trifluoroacetic acid and desalted using SDB-

RPS StageTips.

Peptides were separated on a 50 cm × 75 µm ID home-packed analytical column containing ReproSil-

Pur C18-AQ resin (1.9 µm; Dr. Maisch) using an EASY-nLC 1200 ultra-high-pressure system.

Samples were loaded in buffer A (0.1% formic acid) and separated using a non-linear gradient from

5% to 60% buffer B (0.1% formic acid, 80% acetonitrile) over 50 min at a flow rate of 300 nL/min.

Peptides were introduced via a CaptiveSpray source with a 10 µm emitter into a timsTOF Pro2 mass

spectrometer (Bruker) operated in diaPASEF mode.

MS data were acquired over a 100–1700 m/z range using the standard long-gradient diaPASEF
method, comprising 16 diaPASEF scans with two 25 Da windows per ramp, covering a total mass
range of 400–1201 Da and an ion mobility range of 1.43–0.6 1/$K_0$. Collision energy decreased linearly
from 59 eV at 1/$K_0$ = 1.6 to 20 eV at 1/$K_0$ = 0.6 Vs cm$^{-2}$. Both accumulation and PASEF ramp times
were set to 100 ms, resulting in a total cycle time of 1.17 s.

Quantification of raw data was performed using Spectronaut v15.5 (Bruderer et al. 2015) in directDIA
mode using the *Homo sapiens* UniprotKB FASTA (UP000005640; TaxID 9606).
Carbamidomethylation of cysteine was set as a fixed modification; oxidation of methionine and
protein N-terminal acetylation were set as variable modifications. Up to two missed cleavages were
allowed, and a minimum peptide length of seven amino acids was required. The false discovery rate
(FDR) at the PSM, peptide, and protein group level was set to 0.01. The raw data processing was
implemented in the automated analysis pipeline of the Clinical Knowledge Graph (Santos et al. 2022)
and exported as label-free quantitative (LFQ) intensities. The LFQ intensities were normalized by
log2 transformation and proteins with less than at least 2 valid values in at least one group were
filtered out. Missing values were imputed using the MinProb approach (width=0.2 and shift=1.8,
(Lazar et al. 2016)) and with Mixed imputation.

**Bioinformatics analysis**
The quantifications from 5903 proteins were used to identify differentially expressed proteins and
generate weighted gene co-expression network analysis.

The differential expression analysis was done using edgeR (v3.40.2; (Chen et al. 2025) and the protein
intensities were normalized with the trimmed mean of M-values method. Generalized linear models
were used to determine if differences in protein intensities were associated with the main effects from
the experimental design: genotypes (wildtype and Ube2j2 KO) and cell treatments (vehicle control
and tunicamycin) at 6 or 18 hours. Pairwise contrasts for DEPs were specified using the emmeans
package (v1.8.7; Lenth and Piaskowski 2017) in R and focused on the effect of tunicamycin treatment
has on Ube2j2 mutants through time (KO_Tu_18hr-vs-WT_Tu_18hr; KO_Ctrl_18hr-vs-
WT_Ctrl_18hr ; KO_Ctrl_6hr-vs-KO_Ctrl_18hr; WT_Ctrl_6hr-vs-WT_Ctrl_18hr). To determine if
a protein was significantly differential expressed with main experiential factors or pairwise contrasts
a Benjamini-Hochberg adjusted p-value was less than 0.05. DEPs with significant pairwise contrasts
were also used as a criterion for hub gene analysis.

Co-expression analysis and module identification were conducted using functions from WGCNA

(v1.72-1; (Langfelder et al. 2008). The soft threshold was determined as the value producing a >90%

model fit to the scale-free topology and low mean connectivity. Co-expression relationships were

calculated using unsigned Pearson's correlation coefficients raised to the soft threshold and grouped

using hierarchical clustering of dissimilarity among the unsigned topological overlap matrix. Modules

were identified using dynamicTreeCut (v1.62-1; (Langfelder et al. 2008)) with parameters of

minimum module size of 100, deep split value of 2, and cut height of 0.99. Module eigengenes were

calculated as the first principal component of the expression matrix and modules with correlated

(>0.80) eigengene values were collapsed because these modules contained proteins that strongly

covary in abundance across samples.

The biological meaning of the co-expression modules was investigated using three approaches. First,

we tested if module eigengenes were associated with the main effects of the experimental conditions

using generalized linear models. Specifically, we tested if module eigengenes were significantly

associated with differences between genotypes (wildtype and Ube2j2 KO) or cell treatments (vehicle

control and tunicamycin) at 6 or 18 hours. Second, functional annotation of the modules as conducted

using gene ontology (GO) enrichment analysis. GO enrichment for each ontology (biological rocess,

molecular function, or cellular component) was performed using the clusterProfiler package (v4.17.0;

(Yu et al. 2012) and bioconductor Human annotation database (org.Hs.eg.db; v3.18). Visualization

of GO enrichment results were done using enrichplot (v1.29.1) and simplifyEnrichment (v1.6.1; (Gu

and Hübschmann, 2023). GO terms were determined to be significant if the Benjamini-Hochberg

adjusted p-value was less than 0.05. Third, hub genes within modules were identified by examining

the interaction between module membership and the –log10 p-values for at least one of the specific

pairwise contrasts. Module membership was calculated as the absolute of the Pearson's correlation

between the module eigengene and protein LFQ intensities. The threshold for module membership

was having a greater than 0.85 correlation, and the DEP threshold was a –log10 adjusted p-value of

greater than 2.

**Data availability**

Cell lines are available upon request. Please refer to the Reagent Table for futher details about cell

lines and antibodies. Supplemental files available at FigShare. Supplementary File 1 contains the

module assignments and accession numbers for proteins in the dataset. Supplementary File 2 contains

the raw and adjusted pvalues used to generate Figure 1C & 1D in this manuscript. Supplementary

File 3 contains the GO annotations for proteins in each module. Code generated for the analyses

described can be found at: https://github.com/mzinkgraf/Ube2j2 . The mass spectrometry proteomics

data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository

(Perez-Riverol et al. 2025) with the dataset identifier PXD076153.

**Acknowledgements**

We acknowledge Dick J. H. van den Boomen (Cambridge Institute for Medical Research, University

of Cambridge, UK) for the design of sgRNA sequences and Linda M. Hendershot (St. Jude Children's

Research Hospital, Memphis, USA) for kindly providing us with rabbit anti-HERP and rabbit anti-

Ube2j2 antibodies.

**Funding**

Mass spectrometry-based proteomics analyses were performed by the Proteomics Research

Infrastructure (PRI) at the University of Copenhagen (UCPH), supported by the Novo Nordisk

Foundation (NNF) (grant agreement number NNF19SA0059305). CLD was supported by funding

from the American Scandinavian Foundation and the Fulbright U.S. Scholar Program. Support for

this research was also provided by the Lizzi og Mogens Staal Fond (grant number 2024-1314). LE

was supported by the Department of Biology, University of Copenhagen, and a sabbatical grant from

the Lundbeck Foundation (grant number R500-2024-1904).

**Conflict of interest**

The authors declare no conflicts of interest.

**Author contributions**

Conceptualization, C.L.D. and L.E.; Data curation, C.L.D. and M.Z.; Formal Analysis, M.Z.; Funding

acquisition, C.L.D. and L.E.; Investigation, C.L.D., S.H.L., C.L.S., Q.G., E.B., R.H.P. ; Project

administration, C.L.D. and L.E.; Supervision, L.E.; Visualization, C.L.D., M.Z., and S.H.L.;

writing—original draft preparation, C.L.D., M.Z., and L.E.; writing—review and editing, all authors.

All authors have read and agreed to the published version of the manuscript.

**Abbreviations**

DEP, differentially expressed protein; DMSO, dimethyl sulfoxide; DUB, deubiquitylating enzyme;

ER, endoplasmic reticulum; ERAD, ER-associated degradation; GO, gene ontology; KO, knockout;

LB, lysogeny broth; LFQ, label-free quantitative; MS, mass spectrometry; PBS, phosphate buffered

saline; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; UPR, unfolded

protein response.; WGCNA, weighted gene co-expression analysis; WT, wildtype.

**Figure 1. Overview of proteomic data gathered from WT and Ube2j2 knockout (KO) cells in four conditions.**

(**a**) Dendrogram showing the relatedness of eigengenes between modules (named by color). (**b**) Line graphs of eigengene expression from the wildtype (WT) and Ube2j2 mutant (KO) genotypes across treatment (control and tunicamycin) and time (6 and 18 hours). (**c**) Heatmap of –log10 p-value of module eigengene significance for main effects from an ANOVA model. (**d**) Heatmap of –log10 p-

value of the comparisons of interest for each of the modules. The smaller the p-value (the warmer the

color), the more significant the difference between conditions.

ALT TEXT: Graphical figure with panels labeled a through d. The figure includes a

dendrogram that shows the relationship between different protein abundance modules

(named by colors). The eigen gene that describes the changes in protein abundance in

wildtype and Ube2j2 cells lines is shown next to each module. Two heat maps show the

statistical relevance of different conditions and pairwise comparisons between conditions

for each module with darker colors representing lower p-values.

**Figure 2. Quantitative relative expression data and immunoblot validation for individual proteins in the pink module.** Graphs show changes in protein abundance for (**a**) BiP, (**b**) HERP, and

(**c**) VIMP in wildtype and Ube2j2 deletion cells in each of the four conditions (DMSO control at 6

and 18 hours, or tunicamycin treatment at 6 and 18 hours). (**d**) Representative immunoblot of U2OS

cells following treatment with tunicamycin for the indicated times. Cell lysates were normalized by

a BCA assay prior to SDS-PAGE loading, and blots were probed for the three UPR-associated

proteins and Ube2j2. (**e**) GO enrichment analysis of proteins present in the pink module using the

biological processes ontology. The graph shows the top 20 most significant terms (adjusted p-value

of <0.05) and the count of module proteins observed in each term.

ALT TEXT: Three box-and-whisker plots labeled a through c show how the abundance of

three individual proteins change in wildtype and Ube2j2 knockout cells under different

conditions. An immunoblot labeled d shows corresponding biochemical data for the

same three proteins.

**Figure 3. Functional annotation and hub gene analysis for the magenta module.** GO enrichment analysis of all protein in the magenta module using the (**a**) cellular component and (**b**) biological processes ontology. The graphs show the most significant terms (adjusted p-value of <0.05) and the count of module proteins observed in each term. (**c**) Hub genes (upper right quadrant) in the magenta module were identified as genes having both a module membership greater then 0.85 and significant (adjusted p-value <0.05) differential abundance with at least one of the pairwise comparisons that relate to changes in protein abundance between KO vs WT cells over time. Proteins that are in the magenta module but are not hub genes are displayed in muted colors in the remaining three quadrants

of the graph. (**d**) GO enrichment analysis of all proteins present in the magenta module using the cellular component ontology. The graph shows the top 20 most significant terms (adjusted p-value of <0.05) and the count of module proteins observed in each term.

ALT TEXT: Three graphs (a, b, d) show the distribution of GO annotations across proteins and hub genes in the magenta module. Circles representing the number of proteins with any given GO annotation are colored according to their statistical significance, with red and blue representing higher and low statistical significance, respectively. Graph c is divided into quadrants. The upper right quadrant includes subset of hub genes in the magenta module accumulate differently depending on their exposure to tunicamycin in Ube2j2 cells.

520 **Figure 4. Analysis of the black and blue modules. (a)** GO enrichment of proteins present in the

521 black module using the biological process **(b)** GO enrichment of proteins present in the blue module

522 using the biological process. Graphs show the top significant (adjusted p-value < 0.05) terms and the

count of module proteins observed in each term. (**c**) Identification of hub genes (upper right quadrant) in the blue module showing proteins with high module membership (>0.85) and significant association with at least one pairwise comparison that relate to changes in protein abundance between KO vs WT cells over time.

ALT TEXT: Two graphs (a, b) show the distribution of GO annotations across proteins and hub genes in the black and blue modules, respectively. Circles representing the number of proteins with any given GO annotation are colored according to their statistical significance, with red and blue representing higher and low statistical significance, respectively. Graph c is divided into quadrants. The upper right quadrant includes subset of hub genes in the blue module accumulate differently depending on their exposure to tunicamycin in Ube2j2 cells.

**Figure 5.** GO enrichment analysis of proteins present in the (**a**) purple, (**b**) midnightblue, and (**c**)

brown modules using the biological process ontology. Terms are significant with an adjusted p-value

of <0.05.

ALT TEXT: Three graphs labeled a, b, and c show the distribution of GO annotations

across proteins and hub genes in the purple, midnightblue, and brown modules. Circles

representing the number of proteins with any given GO annotation are colored according

to their statistical significance, with red and blue representing higher and low statistical

significance, respectively.

545
546

 **Supplementary Figure 1. Generation and verification of Ube2j2 KO cell line.** (**a**) Nucleotide

 sequence of the mutated *UBE2J2* clone, illustrating a 432 bp deletion that spans both non-coding

 (grey) and coding (black) regions. Base numbering is relative to the start of exon 2, the first coding

 exon. (**b**) Schematic representation of the Ube2j2 protein domain architecture. The deleted region is

 indicated by black dashed lines in the protein and by an orange dashed line in the upstream nucleic

 acid sequence. Numbers refer to amino acid residues; TM: transmembrane domain; ER:

 endoplasmic reticulum lumen localization. (**c**) Immunoblot analysis of protein lysates from wild-

 type (WT) and *UBE2J2* knockout (KO) U2OS cells. β-actin serves as a loading control.

 ALT TEXT: A figure with three panels labeled a, b and c. a and c show the DNA

 sequence and schematic representations of the Ube2j2 knock-out in the cells used in this

 study. c shows the immunoblot that confirmed the successful removal of the Ube2j2

 protein in the cells used in this study.

## References

Abdul Rehman SA et al. 2024. Discovery and characterization of noncanonical E2-conjugating enzymes. Science Advances. 10(13):eadh0123 [accessed 2025 Oct 5]. https://www.science.org/doi/10.1126/sciadv.adh0123. https://doi.org/10.1126/sciadv.adh0123

Ashburner M et al. 2000. Gene Ontology: tool for the unification of biology. Nat Genet. 25(1):25–29 [accessed 2025 Oct 31]. https://www.nature.com/articles/ng0500_25. https://doi.org/10.1038/75556

van den Boomen DJH et al. 2014. TMEM129 is a Derlin-1 associated ERAD E3 ligase essential for virus-induced degradation of MHC-I. Proceedings of the National Academy of Sciences. 111(31):11425–11430 [accessed 2025 Oct 5]. https://www.pnas.org/doi/full/10.1073/pnas.1409099111. https://doi.org/10.1073/pnas.1409099111

Bruderer R et al. 2015. Extending the Limits of Quantitative Proteome Profiling with Data-Independent Acquisition and Application to Acetaminophen-Treated Three-Dimensional Liver Microtissues *[S]. Molecular & Cellular Proteomics. 14(5):1400–1410 [accessed 2025 Oct 5]. https://www.mcponline.org/article/S1535-9476(20)32885-1/abstract. https://doi.org/10.1074/mcp.M114.044305

Burr ML et al. 2013. MHC class I molecules are preferentially ubiquitinated on endoplasmic reticulum luminal residues during HRD1 ubiquitin E3 ligase-mediated dislocation. Proceedings of the National Academy of Sciences. 110(35):14290–14295 [accessed 2025 Oct 5]. https://www.pnas.org/doi/full/10.1073/pnas.1303380110. https://doi.org/10.1073/pnas.1303380110

Carvalho P, Goder V, Rapoport TA. 2006. Distinct Ubiquitin-Ligase Complexes Define Convergent Pathways for the Degradation of ER Proteins. Cell. 126(2):361–373 [accessed 2025 Oct 5]. https://www.cell.com/cell/abstract/S0092-8674(06)00857-9. https://doi.org/10.1016/j.cell.2006.05.043

Chen Y et al. 2025. edgeR v4: powerful differential analysis of sequencing data with expanded functionality and improved support for small counts and larger datasets. Nucleic Acids Research. 53(2):gkaf018 [accessed 2025 Oct 5]. https://academic.oup.com/nar/article/doi/10.1093/nar/gkaf018/7973897. https://doi.org/10.1093/nar/gkaf018

Cho K-C et al. 2021. Evaluation of the sensitivity and reproducibility of targeted proteomic analysis using data independent acquisition for serum and cerebrospinal fluid proteins. J Proteome Res. 20(9):4284–4291 [accessed 2025 Oct 31]. https://pmc.ncbi.nlm.nih.gov/articles/PMC8631582/. https://doi.org/10.1021/acs.jproteome.1c00238

Christensen LC et al. 2012. The human selenoprotein VCP-interacting membrane protein (VIMP) is non-globular and harbors a reductase function in an intrinsically disordered region. J Biol Chem. 287(31):26388–26399. https://doi.org/10.1074/jbc.M112.346775

Christianson JC, Carvalho P. 2022. Order through destruction: how ER-associated protein degradation contributes to organelle homeostasis. The EMBO Journal. 41(6):e109845 [accessed 2024 Feb 19]. https://www.embopress.org/doi/full/10.15252/embj.2021109845. https://doi.org/10.15252/embj.2021109845

Christianson JC, Jarosch E, Sommer T. 2023. Mechanisms of substrate processing during ER-
associated protein degradation. Nat Rev Mol Cell Biol. 24(11):777–796.
https://doi.org/10.1038/s41580-023-00633-8

Cremer T et al. 2021. The ER-embedded UBE2J1/RNF26 ubiquitylation complex exerts
spatiotemporal control over the endolysosomal pathway. Cell Reports. 34(3):108659 [accessed
2025 Oct 5]. https://linkinghub.elsevier.com/retrieve/pii/S221112472031648X.
https://doi.org/10.1016/j.celrep.2020.108659

Fischer JA. 2003. Deubiquitinating enzymes: their roles in development, differentiation, and
disease. Int Rev Cytol. 229:43–72 [accessed 2012 Mar 15].
http://www.ncbi.nlm.nih.gov/pubmed/14669954

Glaeser K et al. 2018. ERAD-dependent control of the Wnt secretory factor Evi. The EMBO
Journal. 37(4):e97311 [accessed 2025 Oct 5].
https://www.embopress.org/doi/10.15252/embj.201797311.
https://doi.org/10.15252/embj.201797311

Gu Z, Hübschmann D. 2023. *SimplifyEnrichment* : A Bioconductor Package for Clustering and
Visualizing Functional Enrichment Results. Genomics, Proteomics & Bioinformatics. 21(1):190–
202 [accessed 2025 Oct 31]. https://academic.oup.com/gpb/article/21/1/190/7274174.
https://doi.org/10.1016/j.gpb.2022.04.008

Gundogdu M, Walden H. 2019. Structural basis of generic versus specific E2–RING E3 interactions
in protein ubiquitination. Protein Science. 28(10):1758–1770 [accessed 2025 Oct 5].
https://onlinelibrary.wiley.com/doi/10.1002/pro.3690. https://doi.org/10.1002/pro.3690

Harding HP et al. 2000. Regulated translation initiation controls stress-induced gene expression in
mammalian cells. Mol Cell. 6(5):1099–1108. https://doi.org/10.1016/s1097-2765(00)00108-8

Hendil KB. 2005. Development and use of antiproteasome monoclonal antibodies. Methods
Enzymol. 398:439–453. https://doi.org/10.1016/S0076-6879(05)98036-1

Hetz C, Chevet E, Oakes SA. 2015a. Proteostasis control by the unfolded protein response. Nat Cell
Biol. 17(7):829–838 [accessed 2025 Oct 2]. https://www.nature.com/articles/ncb3184.
https://doi.org/10.1038/ncb3184

Hetz C, Chevet E, Oakes SA. 2015b. Proteostasis control by the unfolded protein response. Nat Cell
Biol. 17(7):829–838. https://doi.org/10.1038/ncb3184

Hunt LC et al. 2023. An adaptive stress response that confers cellular resilience to decreased
ubiquitination. Nat Commun. 14(1):7348 [accessed 2025 Oct 5].
https://www.nature.com/articles/s41467-023-43262-7. https://doi.org/10.1038/s41467-023-43262-7

Kreft SG, Hochstrasser M. 2011. An unusual transmembrane helix in the endoplasmic reticulum
ubiquitin ligase Doa10 modulates degradation of its cognate E2 enzyme. J Biol Chem.
286(23):20163–20174. https://doi.org/10.1074/jbc.M110.196360

Kumari S, Kumar P. 2023. Identification and characterization of putative biomarkers and
therapeutic axis in Glioblastoma multiforme microenvironment. Front Cell Dev Biol. 11 [accessed

2025 Oct 5]. https://www.frontiersin.org/journals/cell-and-developmental-biology/articles/10.3389/fcell.2023.1236271/full. https://doi.org/10.3389/fcell.2023.1236271

Langfelder P, Zhang B, Horvath S. 2008. Defining clusters from a hierarchical cluster tree: the Dynamic Tree Cut package for R. Bioinformatics. 24(5):719–720 [accessed 2025 Oct 5]. https://academic.oup.com/bioinformatics/article/24/5/719/200751. https://doi.org/10.1093/bioinformatics/btm563

Lazar C et al. 2016. Accounting for the Multiple Natures of Missing Values in Label-Free Quantitative Proteomics Data Sets to Compare Imputation Strategies. J Proteome Res. 15(4):1116–1125 [accessed 2025 Oct 5]. https://pubs.acs.org/doi/10.1021/acs.jproteome.5b00981. https://doi.org/10.1021/acs.jproteome.5b00981

Lenk U et al. 2002. A role for mammalian Ubc6 homologues in ER-associated protein degradation. Journal of Cell Science. 115(14):3007–3014 [accessed 2025 Oct 5]. https://doi.org/10.1242/jcs.115.14.3007. https://doi.org/10.1242/jcs.115.14.3007

Lenth RV, Piaskowski J. 2017. emmeans: Estimated Marginal Means, aka Least-Squares Means. 2.0.1 [accessed 2026 Feb 15]. https://CRAN.R-project.org/package=emmeans. https://doi.org/10.32614/CRAN.package.emmeans

Lin S et al. 2024. The UBE2J2/UBE2K-MARCH5 ubiquitination machinery regulates apoptosis in response to venetoclax in acute myeloid leukemia. Leukemia. 38(3):652–656 [accessed 2025 Oct 5]. https://www.nature.com/articles/s41375-024-02178-x. https://doi.org/10.1038/s41375-024-02178-x

Lips C et al. 2020. Who with whom: functional coordination of E2 enzymes by RING E3 ligases during poly-ubiquitylation. The EMBO Journal. 39(22):e104863 [accessed 2021 Apr 26]. http://www.embopress.org/doi/full/10.15252/embj.2020104863. https://doi.org/10.15252/embj.2020104863

Liu X et al. 2019. Human cytomegalovirus evades antibody-mediated immunity through endoplasmic reticulum-associated degradation of the FcRn receptor. Nat Commun. 10(1):3020 [accessed 2025 Oct 5]. https://www.nature.com/articles/s41467-019-10865-y. https://doi.org/10.1038/s41467-019-10865-y

Mehrtash AB, Hochstrasser M. 2022. Elements of the ERAD ubiquitin ligase Doa10 regulating sequential poly-ubiquitylation of its targets. iScience. 25(11) [accessed 2025 Oct 5]. https://www.cell.com/iscience/abstract/S2589-0042(22)01623-6. https://doi.org/10.1016/j.isci.2022.105351

Nakao F et al. 2023. Targeting a mitochondrial E3 ubiquitin ligase complex to overcome AML cell-intrinsic Venetoclax resistance. Leukemia. 37(5):1028–1038 [accessed 2025 Oct 5]. https://www.nature.com/articles/s41375-023-01879-z. https://doi.org/10.1038/s41375-023-01879-z

Parashar S, Kaushik A, Ambasta RK, Kumar P. 2025. E2 conjugating enzymes: A silent but crucial player in ubiquitin biology. Ageing Research Reviews. 108:102740 [accessed 2025 Oct 2]. https://www.sciencedirect.com/science/article/pii/S1568163725000868. https://doi.org/10.1016/j.arr.2025.102740

Perez-Riverol Y et al. 2025. The PRIDE database at 20 years: 2025 update. Nucleic Acids Res.
53(D1):D543–D553. https://doi.org/10.1093/nar/gkae1011

Ravid T, Kreft SG, Hochstrasser M. 2006. Membrane and soluble substrates of the Doa10 ubiquitin
ligase are degraded by distinct pathways. EMBO J. 25(3):533–543.
https://doi.org/10.1038/sj.emboj.7600946

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR : a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 26(1):139–140 [accessed 2026
Jan 19]. https://academic.oup.com/bioinformatics/article/26/1/139/182458.
https://doi.org/10.1093/bioinformatics/btp616

Santos A et al. 2022. A knowledge graph to interpret clinical proteomics data. Nat Biotechnol.
40(5):692–702 [accessed 2025 Oct 5]. https://www.nature.com/articles/s41587-021-01145-6.
https://doi.org/10.1038/s41587-021-01145-6

Sato BK, Schulz D, Do PH, Hampton RY. 2009. Misfolded membrane proteins are specifically
recognized by the transmembrane domain of the Hrd1p ubiquitin ligase. Mol Cell. 34(2):212–222.
https://doi.org/10.1016/j.molcel.2009.03.010

Serin EAR, Nijveen H, Hilhorst HWM, Ligterink W. 2016. Learning from Co-expression
Networks: Possibilities and Challenges. Front Plant Sci. 7 [accessed 2025 Dec 22].
http://journal.frontiersin.org/Article/10.3389/fpls.2016.00444/abstract.
https://doi.org/10.3389/fpls.2016.00444

Sharninghausen R, Hwang J, Dennison DD, Baldridge RD. 2024. Identification of ERAD-
dependent degrons for the endoplasmic reticulum lumen Glick BS, Ron D, editors. eLife.
12:RP89606 [accessed 2025 Oct 5]. https://doi.org/10.7554/eLife.89606.
https://doi.org/10.7554/eLife.89606

Sowa ME, Bennett EJ, Gygi SP, Harper JW. 2009. Defining the human deubiquitinating enzyme
interaction landscape. Cell. 138(2):389–403 [accessed 2012 Mar 15].
http://www.ncbi.nlm.nih.gov/pubmed/19615732. https://doi.org/10.1016/j.cell.2009.04.042

Stewart MD, Ritterhoff T, Klevit RE, Brzovic PS. 2016. E2 enzymes: more than just middle men.
Cell Res. 26(4):423–440. https://doi.org/10.1038/cr.2016.35

Swarnkar A et al. 2024. Determinants of chemoselectivity in ubiquitination by the J2 family of
ubiquitin-conjugating enzymes. The EMBO Journal. 43(24):6705–6739 [accessed 2025 Oct 5].
https://www.embopress.org/doi/full/10.1038/s44318-024-00301-3. https://doi.org/10.1038/s44318-
024-00301-3

Vella D et al. 2017. From protein-protein interactions to protein co-expression networks: a new
perspective to evaluate large-scale proteomic data. J Bioinform Sys Biology. 2017(1):6 [accessed
2025 Dec 22]. https://bsb-eurasipjournals.springeropen.com/articles/10.1186/s13637-017-0059-z.
https://doi.org/10.1186/s13637-017-0059-z

Vrentzou A et al. 2025. UBE2J2 sensitizes the ERAD ubiquitination cascade to changes in
membrane lipid saturation. Nat Commun. 16(1):8973. https://doi.org/10.1038/s41467-025-64777-1

Wang X et al. 2009. Ube2j2 ubiquitinates hydroxylated amino acids on ER-associated degradation
substrates. J Cell Biol. 187(5):655–668 [accessed 2025 Oct 5].
https://doi.org/10.1083/jcb.200908036. https://doi.org/10.1083/jcb.200908036

Weber A et al. 2016. Sequential Poly-ubiquitylation by Specialized Conjugating Enzymes Expands
the Versatility of a Quality Control Ubiquitin Ligase. Mol Cell. 63(5):827–839.
https://doi.org/10.1016/j.molcel.2016.07.020

van de Weijer ML et al. 2014. A high-coverage shRNA screen identifies TMEM129 as an E3 ligase
involved in ER-associated protein degradation. Nat Commun. 5:3832.
https://doi.org/10.1038/ncomms4832

van Wijk SJL et al. 2009. A comprehensive framework of E2–RING E3 interactions of the human
ubiquitin–proteasome system. Molecular Systems Biology. 5(1):317 [accessed 2025 Oct 5].
https://doi.org/10.1038/msb.2009.76. https://doi.org/10.1038/msb.2009.76

van Wijk SJL, Timmers HTM. 2010. The family of ubiquitin-conjugating enzymes (E2s): deciding
between life and death of proteins. The FASEB Journal. 24(4):981–993 [accessed 2024 Nov 26].
https://onlinelibrary.wiley.com/doi/abs/10.1096/fj.09-136259. https://doi.org/10.1096/fj.09-136259

Witus SR et al. 2021. BRCA1/BARD1 site-specific ubiquitylation of nucleosomal H2A is directed
by BARD1. Nat Struct Mol Biol. 28(3):268–277 [accessed 2025 Oct 5].
https://www.nature.com/articles/s41594-020-00556-4. https://doi.org/10.1038/s41594-020-00556-4

Wolf LM, Lambert AM, Haenlin J, Boutros M. 2021. EVI/WLS function is regulated by
ubiquitylation and is linked to ER-associated degradation by ERLIN2. J Cell Sci.
134(16):jcs257790 [accessed 2025 Oct 5]. https://doi.org/10.1242/jcs.257790.
https://doi.org/10.1242/jcs.257790

Ye Y, Rape M. 2009. Building ubiquitin chains: E2 enzymes at work. Nat Rev Mol Cell Biol.
10(11):755–764 [accessed 2025 Oct 5]. https://www.nature.com/articles/nrm2780.
https://doi.org/10.1038/nrm2780

Younger JM et al. 2006. Sequential Quality-Control Checkpoints Triage Misfolded Cystic Fibrosis
Transmembrane Conductance Regulator. Cell. 126(3):571–582 [accessed 2025 Oct 31].
https://linkinghub.elsevier.com/retrieve/pii/S0092867406009081.
https://doi.org/10.1016/j.cell.2006.06.041

Yu G, Wang L-G, Han Y, He Q-Y. 2012. clusterProfiler: an R Package for Comparing Biological
Themes Among Gene Clusters. OMICS: A Journal of Integrative Biology. 16(5):284–287 [accessed
2025 Oct 31]. https://journals.sagepub.com/doi/10.1089/omi.2011.0118.
https://doi.org/10.1089/omi.2011.0118

Yu X et al. 2025. UBE2J2 is essential for the progression of meiosis prophase I during
spermatogenesis in mice. iScience. 28(8) [accessed 2025 Oct 5].
https://www.cell.com/iscience/abstract/S2589-0042(25)01139-3.
https://doi.org/10.1016/j.isci.2025.112878

757 Zattas D, Hochstrasser M. 2015. Ubiquitin-dependent protein degradation at the yeast endoplasmic
758 reticulum and nuclear envelope. Crit Rev Biochem Mol Biol. 50(1):1–17.
759 https://doi.org/10.3109/10409238.2014.959889