A faster incubation explains Usutu leading West Nile in temperate Europe

Robert S. Paton [†*a], Maisie Vollans [*a], Lorna Glenn [b], Martyn Fyles [a], Alexander G.C. Vaux [c], Jolyon M. Medlock [c], Julie Day [a] & Thomas Ward [a]

a. Infectious Disease Modelling Team, Analysis and Intelligence Assessment, Chief Data Officer Group, UKHSA, London, UK
b. Institute of Infection, Veterinary and Ecological Sciences, Faculty of Health and Life Sciences, University of Liverpool, Liverpool, UK
c. Medical Entomology and Zoonoses Ecology, Centre for Climate and Health Security, UKHSA, Salisbury, UK

† Corresponding author address: rob.paton2@ukhsa.gov.uk.
* Co-first authors.

Abstract

Usutu virus (USUV) is a mosquito-borne flavivirus that has recently expanded northwards in Europe and

become endemic in the UK [1-3]. USUV emergence often precedes the closely related West Nile virus

(WNV), potentially reflecting differences in epidemiological parameters [4, 5]. One key parameter is the

extrinsic incubation period (EIP), the time required for a mosquito to become infectious following an

infected blood meal. Here we present the first ever estimate of the temperature-dependent EIP for USUV

in the vector *Culex pipiens molestus*. We were able to quantify the shortening of the EIP with temperature

by re-analysing published laboratory data with bespoke Bayesian model that accounted for key features

of the experimental design. Under typical UK summer temperatures, the median EIP (EIP_{50}) of USUV is

shorter than that of WNV, and the potential transmission season of USUV is both longer and

geographically more extensive. Under RCP8.5 climate projections, WNV transmission suitability is

expected to match or exceed current USUV levels between 2055 and 2065, highlighting the future threat

to the UK from emerging mosquito-borne pathogens. Our findings support USUV as a precursor for WNV

in northern Europe and provide a robust characterisation of a key epidemiological parameter of USUV,

enabling accurate modelling of its transmission dynamics.

Key words: Usutu; Vector-borne disease; EIP; WNV; Bayesian statistics; *Culex pipiens molestus; Survival model; Culex pipiens pipiens*; RCP8.5; Climate change.

Introduction

Usutu virus (USUV) is an arbovirus that primarily infects birds, causing high mortality and mass die-offs

in species such as the Eurasian blackbird, *Turdus merula* [6]. The virus is maintained in an enzootic

transmission cycle involving avian hosts and ornithophilic mosquitoes with birds acting as amplifying

hosts and reservoirs of the virus [7]. Spillover events can occur to humans and other mammals, which

are dead-end hosts and therefore do not contribute to onward transmission [8]. The dominant vector of

USUV in Europe are mosquitoes from the *Culex pipiens* species complex [4], with *Culex pipiens molestus*

biotype considered a potential bridge vector that facilitates the spillover of USUV from birds to humans

in Europe, due to its more diverse biting habits including both birds and mammals [9].

Clinical symptoms of USUV infection are not well characterised in humans, due to the relatively low

number of reported cases in comparison to other arboviruses, such as West Nile virus (WNV) [10, 11]. In

the cases that have been reported, USUV infections are generally asymptomatic or result in mild

symptoms. However, the virus can in rare cases cause severe neurological symptoms (including

meningitis and meningoencephalitis) which are more likely to occur in immunocompromised individuals

[8]. Consequently, USUV is a notable public health concern.

In recent years, USUV has undergone substantial range expansion in Europe. USUV is thought to have

been introduced to Europe from Africa multiple times since the 1950s via the movement of infectious

migratory birds [12]. The first documented case of USUV in birds in Europe dates from 1996 [13], where

retrospective analyses of dead Eurasian Blackbirds (*Turdus merula*) were tested in Italy. Since its

introduction, USUV has become widespread across Europe. This includes Northern Europe, where the

United Kingdom first detected USUV in both birds and mosquitoes in 2020 [1, 2]. There is evidence to

suggest that USUV overwinters in the UK, with autochthonous transmission among birds and vertical

transmission to mosquito eggs in the *Culex pipiens* complex [3]. To date, there have been no published

reports of autochthonous human infections in the UK.

WNV was retrospectively detected in *Aedes vexans* sampled from Gamston, Nottinghamshire, UK from samples collected in 2023 [14], but with no subsequent reports of autochthonous transmission to either birds, or humans. This reflects broader trends observed across Central and Northern Europe, where USUV is often either transiently detected or becomes established endemically in avian populations several years before the first detection of WNV in birds or humans. This has been reported in several countries such as Austria [15, 16], Germany [17, 18] and the Netherlands [19, 20]. Similarly, USUV has been shown to overwinter and become endemic in multiple European countries, including Belgium and the UK, before autochthonous transmission of WNV was reported [3, 21].

The repeated observation that USUV circulation precedes WNV [4, 5] emergence may indicate differences in their transmission dynamics. However, there has been little quantitative work to test this hypothesis. A major barrier to comparative analysis between USUV and WNV is the poor characterisation of key epidemiological parameters for USUV. A key parameter that is yet to be characterised for USUV is the extrinsic incubation period (EIP), which is the time taken for a mosquito to become infectious after ingesting an infected blood meal.

During the EIP, the virus must overcome multiple tissue barriers within the vector, evade immune responses and disseminate to the mosquito salivary glands [22-24]. Empirically, the EIP is generally defined as the time from ingestion of an infectious blood meal to viral dissemination to either the mosquito's legs or salivary glands. The detection of viral material in the mosquito body is typically used as evidence of infection only, not dissemination. Detection of virus in the legs is often preferred for determining the EIP, as it provides a more reliable proxy for the feasibility of transmission via mosquito saliva, given the technical challenges associated with saliva extraction [25].

The EIP is highly temperature sensitive: higher temperatures lead to faster rates of viral replication within the mosquito, shortening the time required for dissemination to the salivary glands [26-28]. As mosquitoes are poikilotherms, their internal body temperature is largely determined by ambient environmental conditions, thus external temperature strongly influences how rapidly mosquitoes

become infectious. The EIP is not deterministic and there is substantial variation among individual mosquitoes, making it important to statistically characterise individual heterogeneity. The specific relationship between EIP and temperature has been characterised for multiple arboviruses and differs among them [26-28].

Here, we reanalyse experimental data from three previously published studies [9, 29, 30] in which adult *Cx. pipiens molestus* mosquitoes were experimentally infected with live USUV and maintained under different, constant temperature conditions. We develop a bespoke Bayesian model to estimate the temperature dependent extrinsic incubation period for USUV. This framework mainly relies on body positivity data, applying a modifier to account for the delay between body infection and dissemination. We use this model, alongside a previously published model of the temperature-dependent West Nile virus EIP [28], to compare the feasibility of transmission of each virus in the UK. This allows us to examine whether differences in the EIP may help explain the relative success of USUV in the UK. For both USUV and WNV, we evaluate a minimum condition for transmission: the probability the EIP could complete in the average mosquito lifespan, giving opportunity for onward infection. We combine these analyses with climate projections to explore how the relative success of WNV and USUV may change under a high-emissions scenario, RCP 8.5 [31]. To our knowledge, this work provides the first characterisation of the temperature-dependent EIP of USUV and the first quantitative comparison of this key epidemiological parameter with WNV.

Results

We find that the extrinsic incubation period of Usutu virus becomes shorter at higher temperatures. The

median EIP (EIP_{50}), defined as the time after which 50% of mosquitoes are expected to be infectious, fell

from 68.06 (95% CrI: 38.62 to 141.04) at 17 °C to 12.35 days (95% CrI: 8.10 to 17.09) at 25°C, as seen in

Figure 3. We compared this estimated median function for the EIP_{50} of USUV to that of WNV [28], finding

that USUV was 37.75% (95% CrI: 3.38 to 55.68) faster between the 17°C to 25°C temperature range. There

was only 2% overlap in the posterior draws, indicating a high level of confidence that USUV has a faster

EIP than WNV.

To estimate the temperature-dependent USUV EIP, we modelled empirical infection data from a total of

565 *Cx. pipiens molestus* mosquitoes across 6 constant temperature treatments ranging from 17 to 25°C,

from three previously published laboratory studies [9, 29, 30], as summarised in **Figure 1**. In these

studies, adult mosquitoes were fed blood meals infected with USUV and maintained at these constant

temperatures representative of UK climatic conditions. Subsets of mosquitoes were sampled at pre-

determined intervals post blood meal and tested for viral RNA using RT-qPCR.

Two of the studies provided temperature-stratified time series of whole-body positivity [9, 29]. The third

study [30] reported both body and leg / wing test positivity at a single temperature. Dissemination to the

leg or wing is commonly used as an experimental proxy for completion of the extrinsic incubation period

[25]. We used data from the third study [30] to estimate the delay between initial body infection and

systemic dissemination. We then apply this modifier to the whole-body positivity data to estimate the

temperature-dependent EIP [9, 29, 30].

As RT-qPCR amplifies viral RNA regardless of origin, it cannot differentiate between RNA contained in the

infectious blood meal, and virus that has passed tissue barriers and is replicating in mosquito tissue.

Consequently, body positivity is expected to be uniformly high immediately after blood feeding, even

though infection in the midgut epithelium has not yet occurred. To account for this, we modelled body

positivity as the combined output of two processes: (i) transient positivity due to blood meal RNA, and (ii)

infection positivity from the virus replicating in mosquito tissue.

Model selection supported a model structure where the EIP_{50} varied log-linearly with temperature and the

probability of infection was study-specific but did not vary with temperature. The parameters estimated

by the model for this log linear function are given in **Table 2**, alongside the other parameters estimated

by the model. The looic results are shown in **Table 1**, where the lowest value indicates the preferred

model structure. It is worth reflecting that the model including both a log-linear median EIP *and* a logit-

linear probability of infection scores similarly to the chosen model. It is intuitive that the infection

probability ρ would vary across temperatures. However, given the number of temperature treatments and

mosquitoes per treatment in the experiments, it is unsurprising that this extra parameter does not explain

enough variation to merit inclusion in the model.

In **Figure 2** we show the model fit and the inferred processes: the estimated model components (**A-C**),

and how the structure fit to the qPCR positivity of the data (**D-E**). We estimate an exponential decline in

the rate of positive tests from the bloodmeal, falling to 50% after around 2.5 days and becoming negligible

towards two weeks post infection (**Figure 2A**). This detection rate includes both live virus that has not yet

infected the mosquito and RNA that may not successfully infect the mosquito. In **Figure 2B**, we show

how the cumulative probability of infection (described by the estimated Weibull probability density

function) changes across the minimum and maximum experimental temperatures, with significantly

steeper increases following infection observed across the relatively modest six-degree temperature

interval used in the study. Combining these two functions (and the study specific probability of infection,

ρ_S), the model can describe the trends in body PCR positivity well, shown in **Figure E**. An initially high

positivity following infection declines due to bloodmeal RNA being metabolised. Further along the

infection timeline (particularly in hotter treatments), this decline is mitigated by an increase in the

genuine positivity from infections. The model predicts a high probability of infection and low variance

across study specific probabilities of infection (**Table 2**, all over 95%). This suggests that the experimental

results from different authors were compatible. However, the model also explores lower infection rates,

down to approximately 80%. This means that model posteriors reflect more incomplete infection rates in

the credible intervals. **Figure D** shows the model fit to the leg / wing positivity data used to estimate the

correction for disseminated infections.

We used our characterisation of the USUV EIP to compare the likelihood of USUV transmission and WNV

transmission in the UK. Our findings suggest that in the temperate climate experienced in the UK, USUV

transmission is more feasible than WNV transmission. For both viruses, we estimated a simple, data-

driven metric: the probability that the EIP could complete within the average vector lifespan. We

estimated the temperature-dependent survival in the mosquito biotype used in the empirical EIP studies

(*Cx. pipiens molestus* for USUV and *Cx. pipiens pipiens* for WNV), using published experimental data [32,

33]. We combined these with the USUV EIP characterised in this study and our previously published

estimates of the WNV EIP [34]. It was regrettable that we were unable to compare the viruses in the same

biotype of *Culex pipiens*, but we determined it was better to reflect the specific thermal biology of the

biotype used in the corresponding laboratory studies rather than ignoring these differences altogether.

In **Figure 4** we compare the approximate USUV EIP in this paper with the values for WNV estimated by

Vollans *et al*. (2024) [28]. We compare the pathogens across UK summer months (June to September),

averaging across 2019 to 2024 temperatures. Then, we calculated the mean number of days in each

month where it was more than 50% likely that the EIP would complete within the average vector lifespan.

USUV exhibits a shorter monthly average EIP across the UK than WNV; the USUV EIP fell as low as

approximately 45.97 days in parts of central and southeastern England in August, compared to a

minimum of approximately 62.38 days for WNV. There were a greater number of days across the average

summer season where the EIP of USUV could complete within the lifespan of *Cx. pipiens molestus* than

for WNV in *Cx. pipiens pipiens*. For example, in August there were an estimated 22.33 days where there

was a greater than 50% chance of the USUV EIP completing in the lifespan of *Cx. pipiens molestus*,

compared to only 13.50 days for WNV in *Cx. pipiens pipiens*. Results for the survival models used to

produce these plots are given in **Figure S1** and **Table S2**. In brief, the average lifespan of *Cx. pipiens*

molestus fell from 75.80 days (95% CrI: 70.55 to 80.99) at 15°C to 18.39 days (95% CrI: 14.11 to 22.78) at

30°C. This compared to 93.36 days (95% CrI: 77.38 to 114.33) and 15.89 days (13.46 to 18.84) across the

same temperature range for *Cx. pipiens pipiens.*

In **Figure 5** we summed the number of days in the year where there was a greater than 50% chance of the

EIP completing within the vector lifespan. Averaging across 2019-2024 there were 59 days where the

USUV EIP could complete within the average lifespan of *Cx. pipiens molestus* compared to 24 days for

WNV in *Cx. pipiens pipiens*. Approximately 37.34% of the UK landmass experienced at least 14 days

where there was a 50% chance of the USUV EIP completing in the vector lifespan, compared to only

7.70% for WNV. For USUV, 23.29% of the country experienced 28 days' worth of risk and 6.99% 42 days

of risk, primarily concentrated in the south and east of England. Nowhere in the UK experienced either of

these levels of risk for WNV.

In **Figure 6 and 7** we use RCP8.5 climate projections for the UK to try to quantify how the transmission

dynamics of USUV and WNV could change as temperatures increase under this specific emissions

pathway. Values presented are averages of twelve probabilistic model runs. The number of days where

there is a 50% chance or more of the EIP completing in the mosquito lifespan increases from a national

average of 28.96 days in 2025 to 59.22 days in 2075 for USUV, compared to 13.99 days and 35.20 for WNV

for the same years. The projected scenario suggests that by 2075, 72.58% of the UK will experience at

least 28 days of USUV transmission risk, expanding into Scotland, Wales and Northern Ireland. The same

statistic is 50.35% for WNV, an expansion which is a doubling of the 24.17% share of the country in 2025.

Figure 7 differences the number of days transmission risk for USUV in 2025 from the number of days for

WNV in each projected year; with Usutu currently endemic, this highlights when a similar level of

endemic transmission risk could be reached. These results suggest that between 2055 and 2065 WNV

will experience a similar number of days at the 50% risk threshold as USUV in 2025.

Discussion

Usutu virus (USUV) is an emerging mosquito-borne virus of growing public health concern, with a

geographic range that has expanded northwards across Europe in recent years. Despite this, key

epidemiological parameters governing USUV transmission remain poorly characterised. Consequently,

modelling of USUV has relied on surrogate parameters from other members of the Japanese encephalitis

virus complex, commonly West Nile virus (WNV) [35, 36]. This introduces substantial uncertainty, limiting

our ability to accurately capture USUV-specific transmission dynamics. Across Northern Europe, USUV

transmission (often alluded to by die offs in avian hosts) precedes the detection of WNV, making it a

priority that we understand why this is the case. Here, we address this knowledge gap by providing the

first characterisation of the temperature-dependent extrinsic incubation period (EIP) of USUV in the

bridge vector *Culex pipiens molestus*. Using this model, we compared the feasibility of USUV and WNV

transmission in the UK, potentially elucidating why USUV is currently endemic while WNV has not yet

known to have established.

Across the temperature range examined (17 to 25°C), we find that the median USUV EIP decreases log-

linearly with temperature, more than halving from 68.06 days (95% CrI: 38.62 to 141.04) at 17°C to 12.35

days (95% CrI: 8.10 to 17.09) at 25°C. The strong thermal sensitivity of the USUV EIP is expected:

mosquitoes are poikilotherms, and their internal temperature varies with ambient temperature.

Consequently, higher environmental temperatures can accelerate viral replication and dissemination,

leading to shorter EIPs. Characterising temperature-dependent parameters for vector-borne pathogens

is essential for assessing transmission risk: the EIP is a key metric underpinning measures such as

vectorial capacity and the basic reproduction number (R_0) [37].

We used our characterisation of the USUV EIP to compare the likelihood of USUV transmission and WNV

transmission in the UK. Our findings suggest that in the temperate climate experienced in the UK, USUV

transmission is more feasible than WNV transmission. For both viruses, we estimated a simple, data-

driven metric: the probability that the EIP could complete within the average vector lifespan. We did this

by characterising temperature-dependent survival in the mosquito biotype used in the empirical EIP

studies (*Cx. pipiens molestus* for USUV and *Cx. pipiens pipiens* for WNV) and combining these with the

USUV EIP characterised in this study and previously published estimates of the WNV EIP [28]. It was

regrettable that we were unable to compare the viruses in the same *Culex pipiens* biotype, but we

determined it was better to reflect the specific thermal biology of the biotype used in the corresponding

laboratory studies rather than ignoring these differences altogether.

We concluded with high confidence that the median EIP of USUV was 37.75% (95% CrI: 3.38 to 55.68)

shorter than that of WNV, at least across the 17 to 25°C temperature conditions explored here. This

supports the observation that USUV tends to establish earlier than WNV in temperate regions of Europe

[5, 38]. Although both viruses circulate primarily in enzootic cycles between birds and *Culex* mosquitoes,

USUV has been more successful in cooler climates such as the Netherlands [38], Denmark [39],

Germany [40] and the UK [1, 3]. Until now, there has been no clear quantitative explanation for these

differences. Our results suggest that the success of USUV in cooler regions may, in part, be explained by

a faster EIP in temperature climates, allowing the virus to infect and disseminate within the lifespan of

the vector. Indeed, it is suggested that the outcome of WNV strain competition in the northern United

States was determined by an emergent strain replicating faster at colder temperatures [28, 41]. This

suggests that faster replication is a valuable attribute in more temperate climates.

We then examined the risk of transmission of WNV and USUV under RCP 8.5 climate projections. Our

results suggest a progressive geographic expansion in the region of thermal suitability for both pathogens

from 2025 until 2075. We show that by between 2055 and 2065 WNV could meet or exceed the number

of days where the EIP is likely to complete in the mosquito lifespan compared to USUV in 2025. We must

emphasize that meeting or exceeding the risk we present for USUV in 2025 does not guarantee or

preclude the establishment of WNV in the UK; it is simply an indicative benchmark that if another

mosquito borne pathogen (USUV) can become endemic under these conditions, then it is credible that

another that meets these conditions also could.

Our USUV EIP model is the first to provide a robust method for estimating the EIP from qPCR body-

positivity data. Our model primarily relies on temperature-dependent whole body positivity data

measured by RT-qPCR [5, 21], supplemented by a single-temperature study reporting body and leg / wing

positivity [30]. While plaque assays are the gold standard for assessing mosquito infectivity, qPCR offers

practical advantages: it is faster, cheaper, more scalable, and requires lower biosafety containment [42,

43], and has been used to study EIP in other arboviruses such as Zika virus [44]. While saliva or leg / wing

samples would better reflect viral dissemination, saliva extraction is technically challenging [25], and

temperature-dependent leg / wing positivity data is not available for USUV. This framework demonstrates

how body positivity data can be leveraged by modelling blood-meal RNA decay to approximate true

infection and applying a modifier to account for the delay between body infection and dissemination.

Our analysis relied on empirical data from three laboratory studies with experimental designs that

required specific statistical treatment. This meant we have had to make a series of compromises and

assumptions, as with any analysis. First, this model parameterisation relied on data collected across a

relatively narrow temperature range, highlighting the necessity for further empirical work to validate

applicability across broader temperature regimes, including future UK climates and other countries.

Second, the USUV EIP data were collected in a "bridge vector", *Cx. pipiens molestus.* This biotype is

interesting as it is important for human spillover risk, with diverse biting habits including both birds and

mammals [45]. However, as vector competence can vary by biotype and species [9], it would be useful

to collect high quality empirical data on USUV EIP in *Cx. pipiens pipiens*, the primary biotype maintaining

USUV in the avian reservoir. Third, our model relies heavily on body positivity data, and the modifier used

to approximate dissemination is based on a single temperature treatment. Temperature-dependent data

on EIP proxies such as leg dissemination, in our view, remain a more appealing way of measuring EIP than

either saliva or body positivity [28] and this remains an important avenue for further empirical work.

Fourth, the empirical data used to characterise the WNV EIP and USUV EIP models vary. The WNV EIP

model is based on data from plaque assays of mosquito legs [28, 41], while the USUV EIP model uses

predominantly body positivity data from RT-qPCR, supplemented by leg positivity data. Additionally, the

WNV EIP is characterised in colonies of *Cx. pipiens pipiens*, but the USUV EIP is estimated in colonies of

Cx. pipiens molestus. Finally, the lab studies use different temperature treatments, rearing regimes,

mosquito strains and USUV lineages. While there are terms in the model to try and account for inter-study

variability (the study-specific probability of infection), this may not fully capture the extent of the

underlying experimental variation. These differences have motivated us to be clear about the limitations

of our approach and the methodological differences that may complicate comparisons across

pathogens and vectors.

We aimed to explore the impact of climate change on the transmission dynamics of USUV and WNV,

however we are limited by the available climate scenarios available. Our modelling metrics required a

daily timestep and high spatial resolution, and the Met Office only provides daily climate change

projections at 12km by 12km for a single emissions scenario, Representative Concentration Pathway

(RCP) 8.5. RCP 8.5 represents a trajectory in which emissions continue largely unmitigated. This ties our

projections to a pessimistic future for USUV and WNV transmission risk based on this high-emissions

scenario. In addition, our projections assume no evolutionary or ecological adaptation of either the

mosquito vectors or the viruses. The extrinsic incubation period (EIP) and mosquito survival models are

parameterised using data from contemporary populations and are applied directly to projected

temperature changes. As a result, potential future changes in vector or viral thermal responses are not

captured.

All coarse-scale analyses of arbovirus infection and ecology must approximate local dynamics. To

estimate the EIPs and adult mosquito lifespans across the UK, and through time, we use the ambient air

temperature. This implicitly assumes that mosquitoes experience these predicted air temperatures

directly, and that this temperature determines viral dissemination and adult mosquito survival. This

assumption may be violated by mosquitoes, which may moderate their experienced temperature across

heterogeneous microhabitats. For instance, the *Cx. pipiens molestus* biotype commonly inhabits

sheltered, belowground human-made structures such as subways, basements and sewage

infrastructure year-round. Consequently*, Cx. pipiens molestus* may experience warmer and more

thermally stable conditions than those captured by ambient air temperature, especially during cooler

periods. Our temperature-based estimates may therefore underestimate transmission potential in some

settings, by underestimating the experienced temperature of the pathogen in a vector routinely seeking

shelter in locations with warmer temperatures. Furthermore, our analysis does not account for spatial

variation in mosquito or avian host abundance. Because the primary transmission cycles of both USUV

and WNV occur between mosquitoes and birds, realised transmission risk depends not only on

temperature suitability but also on local host and vector densities. This is particularly the case for the

establishment of WNV and reintroductions of USUV, where infected migratory birds from the continent

can trigger cycles of infection. Our results should therefore be interpreted as characterising the climatic

feasibility of transmission, rather than predicting the precise transmission intensity or vector habitat

suitability.

Conclusion

In this paper we approximate the temperature-dependent USUV EIP for the first time. This research

contributes essential knowledge to a poorly characterised pathogen of increasing relevance to public

health, particularly in Northern Europe. This USUV EIP can be used to estimate transmission risk and

public health implications of the virus, with measures such as the basic reproduction number reliant on

accurate pathogen-specific measures. Our evidence supports USUV as a precursor for WNV in Northern

Europe and suggests that WNV endemicity will become increasingly likely with climate change, matching

the dynamics of USUV by 2065 at the latest.

References

1.   Folly, A.J., et al., *Detection of Usutu virus infection in wild birds in the United Kingdom, 2020*. Eurosurveillance, 2020. **25**(41): p. 2001732.
2.   Lawson, B., et al., *Combining host and vector data informs emergence and potential impact of an Usutu virus outbreak in UK wild birds*. Scientific Reports, 2022. **12**(1): p. 10298.
3.   Folly, A.J., et al., *Evidence for overwintering and autochthonous transmission of Usutu virus to wild birds following its redetection in the United Kingdom*. Transboundary and Emerging Diseases, 2022. **69**(6): p. 3684–3692.
4.   Nikolay, B., *A review of West Nile and Usutu virus co-circulation in Europe: how much do transmission cycles overlap?* Transactions of The Royal Society of Tropical Medicine and Hygiene, 2015. **109**(10): p. 609–618.
5.   Laverdeur, J., et al., *West Nile and Usutu viruses: current spreading and future threats in a warming northern Europe*. Frontiers in Virology, 2025. **Volume 5 - 2025**.
6.   Ashraf, U., et al., *Usutu Virus: An Emerging Flavivirus in Europe*. Viruses, 2015. **7**(1): p. 219–238.
7.   Roiz, D., et al., *Evidence that Passerine Birds Act as Amplifying Hosts for Usutu Virus Circulation*. EcoHealth, 2019. **16**(4): p. 734–742.
8.   Cadar, D. and Y. Simonin, *Human Usutu Virus Infections in Europe: A New Risk on Horizon?* Viruses, 2023. **15**(1): p. 77.
9.   Pilgrim, J., et al., *UK mosquitoes are competent to transmit Usutu virus at native temperatures*. One Health, 2024. **19**: p. 100916.
10.   Agliani, G., et al., *Pathological features of West Nile and Usutu virus natural infections in wild and domestic animals and in humans: A comparative review*. One Health, 2023. **16**: p. 100525.
11.   Lustig, Y., et al., *Surveillance and Diagnosis of West Nile Virus in the Face of Flavivirus Cross-Reactivity*. Frontiers in Microbiology, 2018. **Volume 9 - 2018**.
12.   Engel, D., et al., *Reconstruction of the Evolutionary History and Dispersal of Usutu Virus, a Neglected Emerging Arbovirus in Europe and Africa*. mBio, 2016. **7**(1): p. e01938–15.
13.   Weissenböck, H., et al., *Usutu Virus, Italy, 1996*. Emerging Infectious Diseases, 2013. **19**(2): p. 274–277.
14.   Bruce, R.C., et al., *Detection of West Nile virus via retrospective mosquito arbovirus surveillance, United Kingdom, 2025*. Eurosurveillance, 2025. **30**(28): p. 2500401.
15.   Wodak, E., et al., *Detection and molecular analysis of West Nile virus infections in birds of prey in the eastern part of Austria in 2008 and 2009*. Veterinary Microbiology, 2011. **149**(3): p. 358–366.
16.   Weissenböck, H., et al., *Usutu virus activity in Austria, 2001–2002*. Microbes and Infection, 2003. **5**(12): p. 1132–1136.
17.   Bergmann, F., et al., *Reconstruction of the molecular evolution of Usutu virus in Germany: Insights into virus emersion and circulation*. PLoS Negl Trop Dis, 2023. **17**(10): p. e0011203.
18.   Ziegler, U., et al., *West Nile virus epizootic in Germany, 2018*. Antiviral Research, 2019. **162**: p. 39–43.
19.   Rijks, J., et al., *Widespread Usutu virus outbreak in birds in the Netherlands, 2016*. Eurosurveillance, 2016. **21**(45): p. 30391.
20.   Vlaskamp, D.R., et al., *First autochthonous human West Nile virus infections in the Netherlands, July to August 2020*. Eurosurveillance, 2020. **25**(46): p. 2001904.
21.   Benzarti, E., et al., *Usutu Virus Epizootic in Belgium in 2017 and 2018: Evidence of Virus Endemization and Ongoing Introduction Events*. Vector-Borne and Zoonotic Diseases, 2019. **20**(1): p. 43–50.
22.   Carpenter, A. and R.J. Clem, *Factors Affecting Arbovirus Midgut Escape in Mosquitoes*. Pathogens, 2023. **12**(2): p. 220.
23.   Erlandson, M.A., U. Toprak, and D.D. Hegedus, *Role of the peritrophic matrix in insect-pathogen interactions*. Journal of Insect Physiology, 2019. **117**: p. 103894.

24. Lim, E.X.Y., et al., *Mosquitoes as Suitable Vectors for Alphaviruses*. Viruses, 2018. **10**(2): p. 84.

25. Gloria-Soria, A., D.E. Brackney, and P.M. Armstrong, *Saliva collection via capillary method may underestimate arboviral transmission by mosquitoes*. Parasites & Vectors, 2022. **15**(1): p. 103.

26. Chan, M. and M.A. Johansson, *The Incubation Periods of Dengue Viruses*. PLOS ONE, 2012. **7**(11): p. e50972.

27. Johansson, M.A., et al., *Incubation periods of Yellow fever virus*. Am J Trop Med Hyg, 2010. **83**(1): p. 183–8.

28. Vollans, M., et al., *Modelling the temperature dependent extrinsic incubation period of West Nile Virus using Bayesian time delay models*. J Infect, 2024. **89**(6): p. 106296.

29. Seechurn, N., et al., *Impact of temperature on vector competence of Culex pipiens molestus: implications for Usutu virus transmission in temperate regions*. Parasites & Vectors, 2025. **18**(1): p. 310.

30. Holicki, C.M., et al., *German Culex pipiens biotype molestus and Culex torrentium are vector-competent for Usutu virus*. Parasit Vectors, 2020. **13**(1): p. 625.

31. Met Office Hadley, C., *UKCP18 Global Climate Model Projections for the entire globe*. 2018.

32. Spanoudis, C.G., et al., *Effect of Temperature on Biological Parameters of the West Nile Virus Vector Culex pipiens form 'molestus' (Diptera: Culicidae) in Greece: Constant vs Fluctuating Temperatures*. Journal of Medical Entomology, 2019. **56**(3): p. 641–650.

33. Andreadis, S.S., O.C. Dimotsiou, and M. Savopoulou-Soultani, *Variation in adult longevity of Culex pipiens f. pipiens, vector of the West Nile Virus*. Parasitology Research, 2014. **113**(11): p. 4315–4319.

34. Vollans, M., et al., *Modelling the temperature dependent extrinsic incubation period of West Nile Virus using Bayesian time delay models*. Journal of Infection, 2024: p. 106296.

35. Rubel, F., et al., *Explaining Usutu virus dynamics in Austria: model development and calibration*. Prev Vet Med, 2008. **85**(3-4): p. 166–86.

36. Cheng, Y., et al., *Evaluating the risk for Usutu virus circulation in Europe: comparison of environmental niche models and epidemiological models*. Int J Health Geogr, 2018. **17**(1): p. 35.

37. Smith DL, B.K., Hay SI, Barker CM, Scott TW, McKenzie FE, *Ross, Macdonald, and a Theory for the Dynamics and Control of Mosquito-Transmitted Pathogens*. PLOS Pathogens, 2012. **8**(4): p. e1002588.

38. Münger, E., et al., *One Health approach uncovers emergence and dynamics of Usutu and West Nile viruses in the Netherlands*. Nature Communications, 2025. **16**(1): p. 7883.

39. Gelskov, L.V., et al., *First detection of Usutu virus in wild birds in Denmark, 2024*. Scientific Reports, 2026. **16**(1): p. 5156.

40. Schopf, F., et al., *Circulation of West Nile virus and Usutu virus in birds in Germany, 2021 and 2022*. Infectious Diseases, 2025. **57**(3): p. 256–277.

41. Kilpatrick, A.M., et al., *Temperature, Viral Genetics, and the Transmission of West Nile Virus by Culex pipiens Mosquitoes*. PLOS Pathogens, 2008. **4**(6): p. e1000092.

42. Di Giuseppe, M., et al., *Can qPCR replace the standard plaque assay in the ASTM F1671 to assess personal protective equipment barrier performance?* Int J Cloth Sci Technol, 2024. **36**(3): p. 385–392.

43. Bae, H.-G., et al., *Detection of yellow fever virus: a comparison of quantitative real-time PCR and plaque assay*. Journal of Virological Methods, 2003. **110**(2): p. 185–191.

44. Winokur, O.C., et al., *Impact of temperature on the extrinsic incubation period of Zika virus in Aedes aegypti*. PLoS Negl Trop Dis, 2020. **14**(3): p. e0008047.

45. Linda, K., et al., *Bloodmeal, Host Selection, and Genetic Admixture Analyses of <i>Culex pipiens</i> Complex (Diptera: Culicidae) Mosquitoes in Chicago, IL*. Journal of Medical Entomology, 2019. **57**(1): p. 78–87.

46. Tsujimoto, H. and Z.N. Adelman, *An 11-point time course midgut transcriptome across 72 h after bloodfeeding provides detailed temporal resolution of transcript expression in the arbovirus vector, Aedes aegypti*. Genome Res, 2023. **33**(9): p. 1638–1648.

47. Lord, J.S. and M.B. Bonsall, *Mechanistic modelling of within-mosquito viral dynamics: Insights into infection and dissemination patterns*. PLoS Comput Biol, 2023. **19**(10): p. e1011520.

48. Vehtari, A., A. Gelman, and J. Gabry, *Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC*. Statistics and Computing, 2017. **27**(5): p. 1413–1432.

49. Stan Development Team, *Stan Modeling Language Users Guide and Reference Manual, 2.32*. 2023.

50. R Core Team, *R: A Language and Environment for Statistical Computing*. 2023, R Foundation for Statistical Computing.

51. *ECMWF Reanalysis v5 (ERA5)*. 2025.

52. Gorelick, N., et al., *Google Earth Engine: Planetary-scale geospatial analysis for everyone*. Remote Sensing of Environment, 2017. **202**: p. 18–27.

Tables (1-2)

Table 1: Model selection results for Usutu EIP model using looic.

	Model structure	looic
1	Approximate median EIP constant across temperatures.	181.54
2	**Approximate median EIP varies log-linearly with temperature.**	**148.56**
3	Approximate median EIP constant across temperatures, the probability of becoming infected varies logit-linearly with temperature.	157.64
4	Approximate median EIP varies log-linearly with temperature and the probability of becoming infected varies logit-linearly with temperature.	150.23

Table 2: Parameters estimates for the Usutu EIP model.

Parameter	Value (posterior median with 95% credible intervals)
Log-linear temperature dependent median EIP	
β_0	3.19 (95% CrI: 2.99 to 3.43)
β_1	-0.55 (95% CrI: -0.8 to -0.36)
$\overline{T}$	21.03
$sd(T)$	2.55
Intercept correction to reflect disseminated infections	
β_2	0.16 (95% CrI: -0.17 to 0.5)
Weibull shape	
α	1.44 (95% CrI: 0.97 to 2.11)
Bloodmeal positivity	
k	0.24 (95% CrI: 0.19 to 0.31)
Infection rates	
$\rho_{S=\text{Pilgrim}}$	95.67% (95% CrI: 77.44 to 99.69)
$\rho_{S=\text{Seechurn}}$	96.00% (95% CrI: 81.47 to 99.72)
$\rho_{S=\text{Holicki}}$	96.14% (95% CrI: 83.91 to 99.76)
$\overline{\rho}$	3.12 (95% CrI: 1.80 to 4.54)
σ_ρ	0.51 (95% CrI: 0.02 to 2.69)
Beta-binomial overdispersion	
θ	28.09 (95% CrI: 13.32 to 66.53)

Figures (1-6)

Figure 1: Data overview.

An overview of the experimental data discussed in this study. Points indicate proportions calculated for each sample and thick / thin lines denote the binomially distributed 80 / 95% credible intervals for each proportion. **Panel A** shows body data from Pilgrim *et al.* (2024) and **panel B** Seechurn *et al.* (2025) (both sets of results refer to *Culex pipiens molestus*). Columns show different temperature treatments (note that temperature conditions are slightly different for each study). In **panel C** we show results for Holicki *et al.* (2020) for body positive PCR tests while **panel D** shows leg / wing positivity.

Figure 2: Model fit to data.

Panel A shows the model estimated probability of a PCR positive test resulting from viral RNA in the bloodmeal. Points and intervals show test data from "day 0" post infectious bloodmeal (which are all exclusively bloodmeal positive) and the 20°C treatment from Seechurn *et al*. (2025), where we it appeared tests reflected almost exclusively bloodmeal positivity. *Panel B* shows the Weibull cumulative density function for the approximate EIP for USUV solved at the temperature extremes from the laboratory data. In *panel C* we show the cumulative probability corrected for disseminated infection. *Panel D* shows the model fit to the 25°C leg / wing positivity data from Holicki *et al*. (2020). Finally, *panel E* shows the model fit to the whole-body positivity data in each experimental treatment, where black points show the raw proportions and black lines are the credible intervals on these proportions. The model fit in *panel C* reflects the combination of the bloodmeal positivity function from *panel A*, the temperature dependent cumulative probability function in *panel B* and the study specific probability of infection ρ_s.

Figure 3: Temperature sensitive extrinsic incubation period of USUV.

Modelled median approximate extrinsic incubation period (EIP_{50}). The posterior median is shown as a line while 80% and 95% credible intervals are shown as dark and light shaded ribbons, respectively. Parameters for this function are given in **Table 2**.

Figure 4: Seasonal variation in transmission risk.

Panel A shows the monthly EIP_{50} of USUV (top row) and WNV (bottom row) in the UK. This is an inter-year average from 2019 to 2024. For both pathogens, these values were calculated using the log-linear function describing the EIP_{50} for each virus on daily temperature data. We calculated the monthly geometric mean across years from these daily values to give an average. Hotter colours indicate faster average EIP_{50} values and cooler colours slower EIP_{50} values. Grey regions show values above 150 days. *Panel B* shows the number of days where there was a greater than 50% chance that the estimated EIP could complete in the average mosquito lifespan for the temperatures in that month. Again, these values are the 2019 to 2024 average. Note that each pathogen is paired with the specific survival model for the species used in the corresponding experiments (*Cx. pipiens molestus* for USUV and *Cx. pipiens pipiens* for WNV). Hotter colours indicate more days above the 50% probability threshold.

Figure 5: Annual variation in transmission risk.

The colour temperature shows the number of days in each year (2019 to 2024) where there is a greater than 50% chance of virus EIP completing in the average mosquito lifespan of *Cx. pipiens molestus* (USUV, top row) and *Cx. pipiens pipiens* (WNV, bottom row) in the UK. The final column shows the 2019 to 2024 average.

Figure 6: Expanding and increasing transmission risk under climate change.

The maps show the number of days in each year where there is a greater than 50% chance of the EIP of the virus completing in the average mosquito lifespan of *Cx. pipiens molestus* (USUV, top row) and *Cx. pipiens pipiens* (WNV, bottom row) in the UK. Each column shows a decade from 2025 up until 2075 for RCP8.5. Values are the average of twelve probabilistic runs for each given year. There is stochastic variation in individual years that causes particular years to deviate from the general trend show across decadal increments. A white contour line delineates the region where there is a least a month (28 days) of the year where the 50% probability condition is expected to be met. Note the small area of white in Northern Ireland is Lough Neagh.

Figure 7: Evolving timeline of WNV transmission risk between 2025 and 2075, compared to USUV risk in 2025.

The maps show the difference between the transmission risk of WNV compared to USUV in 2025, where blue indicates lower risk, white equivalent risk and red a higher risk. We calculated the number of days in 2025 where the EIP of USUV had a 50% or greater chance of completing in the mosquito lifespan for each grid cell. This was then subtracted from the equivalent value for WNV in each year to give the number of days less or more than the current USUV transmission risk. Note the small area of white in Northern Ireland is Lough Neagh.

Methods

Laboratory protocols

This analysis used data from three laboratory studies: Pilgrim *et al.* (2024) [9], Seechurn *et al.* (2025) [29]

and Holicki *et al.* (2020) [30].

Pilgrim *et al.* (2024) [9] and Seechurn *et al.* (2025) [29] experimentally infected cohorts of *Culex pipiens*

molestus with the UK-endemic strain of USUV (London, 2020) via infectious blood meals. Pilgrim *et al.*

(2024) [9] infected a total of 187 mosquitoes, while Seechurn *et al.* (2025) [29] infected 336 mosquitoes.

Following infection, mosquitoes were divided into groups, and held at multiple different, constant

temperatures representative of the UK climate (17°C to 23°C). At predefined days post-infection (dpi),

subsets of mosquitoes were sampled, and bodies and saliva tested for viral RNA using quantitative

reverse transcriptase PCR (RT-qPCR).

A limitation of both studies is that the saliva positivity shows no clear trend through time, or across

temperatures. This observation is consistent with a previous study which demonstrated that forced

salivation assays have technical challenges and poor reproducibility, often leading to an underestimation

of transmission compared with using viral dissemination to mosquito legs as a proxy [25]. For these

reasons, our previously published model describing the EIP of WNV used leg positivity as a proxy for

disseminated infection as opposed to saliva [28]. In this study we therefore do not model the saliva data

from Pilgrim *et al.* (2024) [9] or Seechurn *et al.* (2025) [29], instead supplementing the body positivity data

with another source.

Holicki *et al.* (2020) [30] followed a similar experimental procedure to Pilgrim *et al.* (2024) [9] and

Seechurn *et al.* (2025) [29] but for only one temperature treatment (25°C). A total of 42 *Cx. pipiens*

molestus from German and Serbian colonies were infected with USUV Africa 2 strain and tested for viral

RNA using RT-qPCR. A crucial difference however is that this study analysed viral RNA separately in

mosquito bodies and the legs / wings. This study therefore contributed an additional 25°C condition to

model the process of whole-body infection, alongside data we could use to parametrise disseminated

infection in the legs and wings, which we explain later in the methods.

Data treatment and interpretation

PCR assays amplify viral RNA, irrespective of whether it is live, infectious virus or inert viral material. This

contrasts with other vector competence methods, such as plaque assays, which assess whether

infectious virus can be cultured from a sample. For example, in Kilpatrick *et al.* (2008) [41] *Culex pipiens*

pipiens mosquitoes were infected with WNV and plaque assays were used to test whole body, leg and

saliva at multiple dpi. In our published reanalysis [28] of these plaque assay data, we modelled the

temperature-dependent EIP of WNV by treating positive tests as interval censored observations (with

dissemination occurring sometime between ingestion of an infectious bloodmeal and sampling) and

negative tests as right censored (dissemination having not yet occurred) observations. Our model

assumed that positivity of disseminated infection is low soon after infection, when tests often precede

dissemination, and increases over time as the virus disseminates throughout the mosquito. We found

that positivity increases more quickly at higher temperatures as viral replication is faster. We make similar

assumptions in this work.

Estimating the temperature-dependent EIP for USUV presents additional challenges, as available data

are from RT-qPCR outputs rather than plaque assay. First, the body positivity data show a complex

pattern of positivity following infection, as seen in **Figure S1**. Body positivity is initially 100%

(demonstrated by Seechurn *et al.* (2025) [29], who sampled mosquitoes on day zero post infection)

followed by a decline soon after, before subsequently increasing. This trend could be explained by an

artefact of the experimental design: initial RT-qPCR positivity reflects viral RNA from the bloodmeal that

has not yet infected the mosquito. As the bloodmeal is metabolised, it is expected this source of viral

RNA decreases, resulting in the observed decline in positivity. Subsequent increases in positivity reflect

the virus infecting the midgut and replicating: this infection process is what we aim to uncover. This

motivates an interesting modelling problem, whereby positivity is driven by both bloodmeal derived RNA

and true infection, changing in representation through time.

Second, the body positivity data from Seechurn *et al*. (2025) [29] shows an unexpected trend with

temperature. Mosquitoes held at 19 and 22°C show higher positivity across all dpi sampling points than

those held at 20°C. As viruses typically replicate faster as temperatures increase, the observed dip in

positivity at 20°C is unlikely to be genuine signal and we therefore assume that the 20°C experimental

treatment by in large failed. However, RNA would still be returned as positive in early tests due to residual

viral RNA from the blood meal, even in the absence of a viable infection. We therefore treat the data from

the 20°C infection treatment in Seechurn *et al*. (2025) [29] as reflecting exclusively bloodmeal derived

positivity, with no infections possible. This helps us to parameterise the decay in positivity associated

with viral RNA only.

To address these challenges, we decided to model the process of mosquito whole body infection using

six temperature treatments: 17, 21, 23°C from Pilgrim et al. (2024) [9] in combination with the 18 and

22°C temperature from Seechurn *et al*. (2025) [29] and 25°C treatment from Holicki *et al*. (2020) [30].

These data were available from the original publications and were extracted for each dpi and temperature

condition. Then we used the leg and wing data available from Holicki *et al*. (2020) [30] to model the

process of the mosquito developing a disseminated infection, by estimating a modifier to the body

infection process to reflect the delay in the infection becoming systemic.

Model overview

Each observed mosquito body is tested on a particular day (d_i where i denotes the unique individual

mosquito) and receives either a positive (y_i^+) or negative (y_i^-) RT-qPCR test result. The test result

determines whether the mosquito became infected between the start of the study and the day it was

observed (positive when observed; interval censored) or may become infected in the future (negative

when observed; right censored). However, positive results also reflect residual viral RNA in the blood

meal, while negative results may include mosquitoes that never develop infection, even if given an

infinite time horizon.

Bloodmeal positive tests

370   To account for residual viral RNA from the blood meal, we define an exponential decline in the probability

371   of a positive test resulting from the blood meal, starting the day after ingestion:

$$\pi(d_i) = e^{-k\, d_i}$$

372   The strictly positive parameter k describes the rate of decline from 100% bloodmeal positive rate on day

373   zero (because infections will not begin immediately) with $\pi(d_i)$ the probability of a bloodmeal positive

374   result on day d_i. We chose this functional form to reflect changes in RNA concentration through time in

375   a parsimonious way [46, 47]

Incomplete infection rates

Not all mosquitoes will necessarily develop an infection: variation in mosquito immunity, microbiome

and the viral load in the blood meal may mean that, even with unlimited time, some mosquitoes may

remain uninfected. This is relevant for right censored observations, where it is unknown whether an

infection will ever develop. We therefore estimate a study specific parameter, ρ_s, which gives the

probability that a mosquito would ever become infected (irrespective of time). This also allows the model

to reflect differences in the two experimental protocols in delivering an infectious dose to sampled

mosquitoes.

Model structure

In our model we use a Weibull distribution (parametrised by a shape parameter α and scale parameter s)

to describe the variation in the approximate EIP across individual mosquitoes. We chose this specific

distribution as our recent work supported this as the best descriptor of the West Nile virus extrinsic

incubation period in *Culex pipiens* [28].

We assume that the median time to infection $\mu T'$ varies log-linearly with temperature. We standardise

experimental temperature by the mean $(\overline{T})$ and standard deviation $(sd(T))$ to improve convergence

properties in the model:

$$T' = \frac{T - \overline{T}}{sd(T)},$$

390 and the log-median is modelled as:

$$\log[\mu(T')] = \beta_0 + \beta_1 T'.$$

391 For a Weibull distribution with shape α and scale $\phi(T')$, the median satisfies $\mu T' = \phi(T')(\log(2))^{1/\alpha}$.

392 Therefore, the scale parameter is:

$$\phi(T') = \frac{\mu T'}{(\log(2))^{\frac{1}{\alpha}}} = \frac{\exp(\beta_0 + \beta_1 T')}{(\log(2))^{\frac{1}{\alpha}}}.$$

393 We model the process of positives due to residual viral RNA from the bloodmeal using data from the 20°C

394 temperature treatment in Seechurn *et al.* (2025) [29], where infection rates were very low:

$$P(y^+ \mid d_i) = \pi(d_i).$$

Mosquito bodies testing positive in other temperature treatments are a mixture of 'true positives'

(mosquitoes that have developed infection) and 'bloodmeal positives' (residual RNA detected from the

blood meal). First, we define $F[d_i \mid \alpha, \phi(T')]$ as the cumulative probability of the Weibull distribution,

modelling the interval censored observation of a true positive (interval censoring occurs as an infection

could have occurred at any point between ingestion of the infectious blood meal, and the time of the

positive test result). Then we define bloodmeal positive tests as A (where the probability is $P(A|d_i) =$

e^{-kd_i}) and true positives as B (where the probability is given by $P(B \mid d_i) = \rho_{s_i} F[d_i \mid \alpha, \phi(t)]$).

As A and B are not mutually exclusive $(A \cup B)$ we model positive tests as:

$$A \cup B = A + B - A \cap B.$$

403   We assume independence between residual blood-meal RNA detectability (event A) and the occurrence

404   of true infection by day d (event B), so that $P(A \cap B) = P(A|d)\,P(B|d)$.

405   Under the assumption that A and B are independent, $P(A \cap B) = P(A)P(B)$. This allows us to write the

406   probability of an observed positive test on day d_i in study s_i in temperature treatment T'_i as:

$$P(y_i^+ \mid d_i, s_i, T'_i) = \pi(d_i) + \rho_{s_i} F[\, d_i \mid \alpha, \phi(T'_i)\,] - \pi(d_i)\rho_{s_i} F[\, d_i \mid \alpha, \phi(T')\,].$$

407   Here, ρ_{s_i} was modelled as a random effect, with each level corresponding to one of the three studies:

$$logit(\rho_{s_i}) = \bar{\rho} + \rho'_{s_i},$$

$$\rho'_{s_i} \sim \mathcal{N}(0, \sigma_\rho).$$

Wing and leg positivity

408   We assume that being body positive is a prerequisite for testing positive in the wings / legs, and that

409   disseminated infections cannot be a result of residual viral RNA from the bloodmeal. Wing and leg

410   positive tests (x_i^+) are therefore the outcome of the overall infection rate ρ_{s_i} and a cumulative probability

411   of infection. As we have no temperature dependent wing / leg positivity data, we rely on a modified body

412   infection process to describe this cumulative probability:

$$log[\mu_d] = \beta_0 + \beta_2 + \beta_1 T',$$

$$\phi_d(T') = \frac{e^{\mu_d(T')}}{log(2)^{\frac{1}{\alpha}}}.$$

$$P(x_i^+ \mid d_i, s_i, T'_i) = \rho_{s_i} F[\, d_i \mid \alpha, \phi_d(T'_i)\,].$$

413   Here the median μ_d is a modified version of μ with an additional parameter β_2 which allows the model to

414   correct the median response for body infections to account for the further delay associated with

415   disseminated infection. This adjustment was applied to the median response, so relies on the same scale

416   parameter α as the body infection process.

Likelihood

417 To capture additional variation associated with test outcomes, we modelled the likelihood ($\mathcal{L}$) as a Beta-

418 binomial process. Here, the mean probability p and dispersion parameter θ give the two positive shape

419 parameters (s_1 and s_2) of a Beta distribution allowing additional variation in outcomes across the number

420 of positive tests (n) out of the total tests (N):

$$p = \begin{cases} P(y^+ \mid d_i) \\ P(y_i^+ \mid d_i, s_i, T_i') \\ P(x_i^+ \mid d_i, s_i, T_i') \end{cases}$$

$$s_1 = p\,\theta,$$

$$s_2 = (1 - p)\,\theta,$$

$$\mathcal{L} = \text{BetaBinomial}(n \mid N, s_1, s_2).$$

Model selection

421 We compare the model structure described above to alternative formulations that reflect different

422 hypotheses. First, we developed a model where the approximate EIP_{50} did not change with temperature,

423 such that:

$$\log[\mu] = \beta_0,$$

$$\log[\mu_d] = \beta_0 + \beta_2.$$

424 Second, we assume that the approximate EIP_{50} was fixed across temperatures, but the probability of ever

425 becoming infected is a function of temperature:

$$\text{logit}[\rho(T')] = \bar{\rho} + \rho'_{s_i} + \delta T'.$$

Here, a logit-linear function temperature dependent process ($\delta T'$) modifies the study specific probability

of infection. This structure allows the model to explain differences in temperature treatments using the

probability of infection only.

A final (and most complex model) allows the approximate EIP_{50} and ρ to vary with temperature. Models

were compared using the leave-one-out cross validation information criterion, "looic" [48]. Models with

looic scores within two standard errors were considered equivalent.

All models were written in the Bayesian programming language "stan" [49] and fit using Hamiltonian

Monte Carlo methods interfaced through the statistical programming language "R" [50]. Priors used in

the model are given in the **Supplementary Table 1**.

Comparison to WNV

To determine how the transmission season for USUV compared to WNV in the UK, we compared the

approximate EIP estimated in this paper to our previously published temperature dependent function for

the median EIP of WNV [28]. We extracted hourly temperature values from ERA5-Land [51] using Google

Earth Engine [52] for the years 2019 to 2024. We created a 5km-by-5km daily raster by resampling the

mean of native-resolution pixels overlapping each output grid and averaging hourly values. For each

pathogen, we solved the log-linear function describing the median EIP on daily temperatures for each

year. We then calculated the geometric mean of these EIPs for each month in each pixel across the five

years.

The values of EIPs in themselves are difficult to interpret in isolation, particularly when comparing across

different vectors and pathogens. We therefore extracted data on *Cx. pipiens pipiens* and *Cx. pipiens*

molestus longevity at different temperatures so we could benchmark the duration of EIPs against the

lifespans of the vectors.

Survival models

Our study analyses the EIP of USUV in *Cx. pipiens molestus* while the model for the EIP of WNV used *Cx.*

pipiens pipiens[28]. To accurately represent the dissemination process for both viruses we use a bespoke

vector survival model for each vector / virus pair. This is challenging, as a component of the differences

in the probability metric we calculate will be due to the vector, rather than the virus. We felt, however,

that it was preferable to accurately represent the vector pathogen pair, as opposed to using the same

survival model that would unavoidably be incorrect for one EIP model.

We used two experimental studies to estimate the temperature dependent survival of *Cx. pipiens*

molestus and *Cx. pipiens pipiens* [32, 33]. In the first, Andreadis *et al.* (2019) [33] reared cohorts of *Cx.*

pipiens pipiens under different temperature conditions (15, 20, 25, 27.5 and 30°C) with 20 males and 20

females in each temperature condition. We digitised data for each temperature treatment from Figure 1c

for females only, allowing us to extract pseudo row-level data. A second study by Spanoudis *et al.* (2018)

[32] examined more temperature conditions for *Cx. pipiens molestus* (15, 17.5, 20, 22.5, 25, 27.5, 30, and

32.5) but we could only extract mean and standard error from Figure 4.

We model observed lifespans, L, with a Weibull distribution where the inverse scale parameter $(1/_\omega)$ is a

log-linear function of standardised temperature and φ is the shape parameter:

$$log(\omega) = \gamma_0 + \gamma_1 \left(\frac{T - \overline{T}}{sd(T)} \right),$$

$$L \sim Weibull\left(\varphi, 1/_\omega\right).$$

The intercept and gradient of the function are given by γ_0 and γ_1 respectively. We assume that the same

shape parameter can be used to describe *Cx. pipiens molestus* survival times as was used to describe

Cx. pipiens pipiens survival times but allow the temperature dependent function to vary across the two

biotypes. As we only had mean and standard errors for each *Cx. pipiens molestus* temperature

treatment, we implemented these data as a soft constraint prior on the predicted mean $(E(\overline{L}))$ of the

Weibull distribution:

$$log(\omega_{mol}) = \gamma_0 + \gamma_2 + (\gamma_1 + \gamma_3) \left(\frac{T - \overline{T}}{sd(T)} \right),$$

$$E(\overline{L}) = \frac{1}{\omega_{mol}} \Gamma \left(1 + \frac{1}{\varphi} \right)$$

$$E(\overline{L}) \sim \mathcal{N}(\overline{L}_{mol}, SE_{mol}).$$

468   Here, γ_2 and γ_3 are the intercept and gradient adjustment for the temperature dependent response for

469   *Cx. pipiens molestus.* The inputs $\bar{L}_{mol}$ and SE_{mol} are the mean and standard error from each

470   experimental treatment. Priors for this model are given in **Table S4**.

Transmission risk

We use these survival models to determine the predicted average lifespan of each mosquito biotype and

estimate the probability of the EIP completing within the vector's lifespan across space and time. To

estimate this, we calculated the average mosquito lifespan at the given temperature and used the

corresponding temperature dependent pathogen cumulative probability function to estimate the

probability of the EIP completing in that lifespan. We term the number of days where there is more than

a 50% chance of competing within the average vector lifespan as "days of transmission risk".

Forecasting the impact of climate change

We used the Met Office projections of RCP8.5 [31] for each year in decadal increments between 2025

and 2075 to assess how the transmission risk increases under climate change. Daily temperatures are

only produced for RCP8.5, restricting our selection to this scenario. Projections are produced for each

day across 12 model runs at a 12km x 12km resolution. We calculated the number of days across each

year and model run where there was a 50% chance or more of the EIP completing in the average mosquito

lifespan. Daily values in each scenario were smoothed with a seven-day rolling average to prevent

stochastic hotspots from inflating risk. Values for each model run were averaged to give a yearly projected

estimate under RCP8.5 [31].

Author contributions

485   RP, MV and JD conceived the study. RP designed and wrote the model with advice, input and review

486   from MV, LG, MF and TW. LG, MV and RP extracted and interpreted the published data. MV and JD

487   handled project administration. RP, MV and LG wrote the manuscript. All authors reviewed and

488   contributed to the final manuscript.

Supplementary materials

Parameter	Prior
Log-linear temperature dependent median EIP	
β_0	$\mathcal{N}(\log(30), 0.33)$
β_1	$\mathcal{N}(-0.33, 0.25)$
Intercept correction to reflect disseminated infections	
β_2	$\mathcal{N}(\log(1.25), 0.25)$
Weibull shape	
α	$\text{Lognormal}(\log(1.5), 0.33)$
Bloodmeal positivity	
k	$\text{Lognormal}(\log(0.27), 0.25)$
Infection rates	
ρ_S	$\mathcal{N}(0, \sigma_\rho)$
$\bar{\rho}$	Model 1 & 2: $\mathcal{N}(2.994, 0.75)$ Model 3 & 4: $\mathcal{N}(0.5, 1)$
σ_ρ	$Exponential(1)$
δ	$\mathcal{N}(1, 0.5)$
Beta-binomial overdispersion	
θ	$\text{Lognormal}(\log(20), 0.5)$

Table S1: Priors used for the Usutu EIP model.

Parameter	Prior	Value (posterior median with 95% credible intervals)
γ_0	$\mathcal{N}(0, 10)$	-3.65 (95% CrI: -3.75 to -3.55)
γ_1	$\mathcal{N}(0, 5)$	0.64 (95% CrI: 0.54 to 0.75)
γ_2	$\mathcal{N}(0, 5)$	0.01 (95% CrI: -0.15 to 0.19)
γ_3	$\mathcal{N}(0, 5)$	-0.13 (95% CrI: -0.27 to 0.01)
ω	$Exponential(1)$	1.97 (95% CrI: 1.68 to 2.3)

Table S2: Priors used for the *Culex pipiens pipiens* and *Culex pipiens molestus* survival models.

Figure S1: We estimated the temperature dependent survival functions for *Cx. pipiens pipiens* and *Cx. pipiens molestus*. We digitised row-level data for the lifespan of *Cx. pipiens pipiens* from Andreadis *et al. (*2014) and used priors from plots in Spanoudis *et al.* (2018) to adjust the model for *Culex pipiens molestus*. In Panel A we show the temperature dependent mean lifespan of each vector. Points and intervals show the priors used for *Cx. pipiens molestus* (mean lifespan as points and standard error as bars) while these points and intervals reflect summaries of the pseudo row level data extracted for *Cx. pipiens pipiens*. In *Panel B* we show the model fit for the pseudo row level data for *Cx. pipiens pipiens*.