nELAVL phosphorylation by CDKL5 regulates RNA metabolism and condensates communication to promote experience-dependent maturation of the visual cortex

Shiyang Yuan[1,22], Yao Zhu[1,22], Zhongyu Zheng[1,22], Hei Matthew Yip[1,22], Maggie See Wing Chan[1], Zhongjie Zhang[1], Yue Chai[1], Kyle Robert Jenks[2], Katya Tsimring[2], Gregg Robert Heller[2], Jose Carlos Zepeda[3], Marco Celotto[2,4], Hung Kwok Hung[5], Yangyang Duan[5], Shun-Fat Lau[5], Crystal Wing Yan Ho[5], Hayley Wing Sum Tsang[5], Amy Kit Yu Fu[5], Hei Ming Lai[6], Andrew John Kwok[7,8,9], Ho Ko[7,8,9], Chunbin Gu[8,10,11], Billy Wai-Lung Ng[8, 9,10,11], Jinyue Liao[12], Sin Hang Fung[1], Qin Cao[1], Stephen Kwok-Wing Tsui[1], Taeyun Ku[13], Liming Tan[14,15], Andrea Colarusso[16], Keren Lasker[16], Yihui Wan[17], Jinwei Zhu[17,18], Jacque Pak Kan Ip[1, 9, 19,20,21]*

[1]School of Biomedical Sciences, The Chinese University of Hong Kong, Hong Kong, China
[2]Department of Brain and Cognitive Sciences, Picower Institute for Learning and Memory, Massachusetts Institute of Technology, Cambridge, MA, USA
[3]Department of Pharmacology, Vanderbilt University, Nashville, TN, USA
[4] Institute for Neural Information Processing, Center for Molecular Neurobiology (ZMNH), University Medical Center Hamburg-Eppendorf (UKE), Hamburg, Germany
[5]Division of Life Science, State Key Laboratory of Molecular Neuroscience and Molecular Neuroscience Center, The Hong Kong University of Science and Technology, Hong Kong, China
[6]Department of Chemical Pathology, The Chinese University of Hong Kong, Hong Kong, China
[7]Department of Medicine and Therapeutics, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China
[8]Li Ka Shing Institute of Health Sciences, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China
[9]Gerald Choa Neuroscience Institute, The Chinese University of Hong Kong, Hong Kong, China
[10]Guangdong-Hong Kong-Macao Joint Laboratory for New Drug Screening, School of Pharmacy, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China
[11]Peter Hung Pain Research Institute, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, China
[12]Department of Ophthalmology, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong, China.
[13]Graduate School of Medical Science and Engineering, Korea Advanced Institute of Science and Technology, Daejeon, Republic of Korea
[14]Shenzhen Key Laboratory of Neuropsychiatric Modulation, Shenzhen-Hong Kong Institute of Brain Science, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China
[15]Guangdong Provincial Key Laboratory of Brain Connectome and Behavior, the Brain Cognition and Brain Disease Institute, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China
[16]Department of Integrative Structural and Computational Biology, Scripps Research, La Jolla, California, USA
[17]Bio-X Institutes, Key Laboratory for the Genetics of Developmental and Neuropsychiatric Disorders, Shanghai Jiao Tong University, Shanghai, China
[18]Department of Nephrology, Shanghai Sixth People's Hospital Affiliated to Shanghai Jiao Tong University School of Medicine, Shanghai, China.

[19]Kunming Institute of Zoology - The Chinese University of Hong Kong (KIZ-CUHK) Joint Laboratory of Bioresources and Molecular Research of Common Diseases, The Chinese University of Hong Kong
[20]CUHK Gerald Choa Neuroscience Institute – Oujiang Laboratory Joint Laboratory for Neuroscience and Neurology, Wenzhou, Zhejiang Province, China.
[21] Lead contact
[22]These authors contributed equally to this work.
*Correspondence: jacqueip@cuhk.edu.hk

Summary

Mutations in Cyclin-Dependent Kinase-Like 5 (CDKL5) cause CDKL5 deficiency disorder (CDD), an X-linked neurodevelopmental condition. Through a phosphoproteomic screen, we identified the neuron-specific nELAVL family of RNA-binding proteins as direct activity-dependent substrates of CDKL5. In support of this regulatory axis, single-nuclei transcriptomics of *Cdkl5* knockout (KO) cortices revealed an enriched reduction in activity-dependent mRNAs. Mechanistically, we show that nELAVL proteins undergo phase separation to form biomolecular condensates, the size of which is gated by CDKL5 phosphorylation. Loss of CDKL5 leads to enlarged nELAVL condensates, which exhibit reduced binding affinity for the target mRNA *Fos*, resulting in its accelerated degradation. This disruption extends to inter-condensate communication: phosphodeficient nELAVL show diminished interaction with P-bodies, which themselves become enlarged in CDD mutant iNeurons. Functionally, the absence of nELAVL phosphorylation recapitulates the deficits in experience-dependent visual function observed in *Cdkl5* KO mice. Our findings establish a critical molecular mechanism by which CDKL5-mediated phosphorylation governs mRNA metabolism by tuning the properties of nELAVL condensates and their communication with other biomolecular condensates, ultimately promoting experience-dependent maturation of the visual cortex.

Introduction

CDD is a severe, X-linked neurodevelopmental disorder caused by loss of function mutations in the *CDKL5* gene[1]. Initially categorized as an early-onset-seizure variant of Rett syndrome, CDD is now recognized as a distinct condition with its own ICD-10 code (G40.42). CDD is characterized by a range of neurodevelopmental features including early-onset epilepsy, motor dysfunction, learning disabilities, autistic features and severe neurodevelopmental delay[2,3]. CDKL5 is a serine/threonine kinase highly expressed in the brain[4]. *Cdkl5* knockout (KO) mice have been shown to exhibit autistic-like phenotypes, impairments in motor control and fear memory[5], as well as reduced amplitude of visually evoked potentials and diminished visual acuity[6]. *Cdkl5* KO also exhibit impaired dendritic spine development and microtubule dynamics[7,8]. Phosphorylation targets of CDKL5 include microtubule-binding proteins EB2 and MAP1S, transcriptional regulators ELOA and SOX9, and the voltage-gated Ca^{2+} channel Cav2.3[7,9-12]. However, clinically relevant phenotypes of CDD rodent model such as motor deficit or cortical visual impairment were not fully recapitulated in the phospho-deficient knock-in mouse models of the abovementioned CDKL5 phosphorylation targets[7,13], suggesting that critical CDKL5-mediated functions or phosphorylation targets remain undiscovered.

Here, we present evidence that CDKL5 regulates the refinement of binocular visual circuits in mouse V1 through a previously unidentified phosphorylation target. We performed an unbiased phosphoproteomic screen and identified a family of RNA-binding proteins

(RBPs), the neuronal Embryonic Lethal, Abnormal Vision-Like proteins (nELAVLs), as
direct substrates of CDKL5. The nELAVL proteins, identified as being recruited into
ribonucleoprotein (RNP) granules[14-16], play crucial roles in regulating RNA stability,
localization, and activity-dependent translation[17,18].

Considering that post-translational modifications, such as phosphorylation, can
impact the phase transition of RNP granules, we postulated that CDKL5-mediated
phosphorylation would modulate the propensity of nELAVLs to form condensates.
Supporting this hypothesis, we found that nELAVL positive condensates are aberrantly
enlarged in *Cdkl5* KO neurons. Furthermore, we observed that CDKL5-mediated
phosphorylation also affects the size of other biomolecular condensates, including those
labelled with Processing (P)-body marker GW182 and stress granules marker TIA1,
suggesting that CDKL5 is a key gatekeeper for fine-tuning condensate properties. Consistent
with this notion, we found that mRNA levels of various activity-dependent genes were
reduced in the V1 of *Cdkl5* KO mice, linking these molecular changes to the impaired
binocular matching of visual responses. Mechanistically, phospho-deficient mutant of
ELAVL3 exhibited reduced binding affinity with its target mRNA *Fos* resulting in faster
degradation. Functionally, we demonstrate that the orientation selectivity and binocular
matching of visual responses is impaired in *Cdkl5* KO mouse V1 neurons with the ipsilateral
eye responses more severely affected. Furthermore, we showed that AAV-mediated
expression of nELAVL phospho-deficient in mouse bV1 recapitulated the deficits observed
in *Cdkl5* KO mice.

Taken together, these findings are consistent with a model of cortical visual
impairment in CDD in which enlarged nELAVL condensates hampers activity-dependent
mRNA regulation, thereby impairing orientation selectivity and binocular matching of V1
neurons.
126
127 **Results**
128
129 **CDKL5 phosphorylates nELAVLs in an activity-dependent manner**
130
Cortical visual impairment is a phenomenon common amongst neurodevelopmental
disorders[6,19-25]. In particular, amongst the 75% of individuals diagnosed with Cyclin-
Dependent Kinase-Like 5 (CDKL5) deficiency disorder (CDD)[4,6,21,26-30], an X-linked
neurodevelopmental disorder attributed by loss of function mutations in the *CDKL5* gene[1],
cortical visual impairment correlates with delayed developmental milestones and has
therefore been proposed to be a potential biomarker[21]. Given the clinical relevance of the
primary visual cortex (V1), we first sought to identify relevant direct substrates of CDKL5 by
applying an unbiased phosphoproteomic screen using V1 tissue from WT and *Cdkl5* KO
mice. We used Tandem Mass Tag-based phosphoproteomics analysis and identified 9048
phosphopeptides on 3146 phosphoproteins in V1 (Figure 1A; table S1). Among them, 119
phosphopeptides were upregulated and 105 downregulated in *Cdkl5* KO mice, corresponding
to 88 and 73 phosphoproteins, respectively (Figure 1B). The number and distribution of
phosphorylation sites on all identified proteins were analyzed (Figure S1A & B). Gene
Ontology (GO) analysis showed that the downregulated phosphopeptides in V1 were
predominantly related to synaptic function or structural constituent of the postsynaptic
density (Figure 1C). Seven of the downregulated phosphorylated proteins contained the
CDKL5 consensus phosphorylation sequences, Arg-Pro-X-Ser-Ala (RPXS*A) or Arg-Pro-X-
Ser (RPXS*)[31], suggesting they are potentially direct substrates of CDKL5 (Figure S1C).
nELAVLs were the most significantly down-regulated phosphorylated proteins among these,
and the phosphorylation sites on ELAVL2 (S119), 3 (S119) and 4 (S131) are evolutionary

conserved among human, mouse, and rat (Figure 1D). ELAVL2, ELAVL3 and ELAVL4
have 80% sequence homology, are predominantly expressed in the nervous system and are
therefore collectively referred as nELAVLs[32]. nELAVLs bind to AU-rich elements in a wide
range of target mRNAs to regulate RNA stability, localization, and translation[17,18,33].
To validate that CDKL5 can phosphorylate nELAVLs, we generated a
phosphospecific antibody targeting these identified sites. We also obtained a wild-type (WT)
EGFP-nELAVLs expression construct and generated a phospho-deficient mutant (nELAVLs
SA) with the putative serine phosphorylation site replaced by alanine. We confirmed the
specificity of nELAVLs antibodies via Western Blot (Figure S1D). Co-expression of CDKL5
and EGFP-ELAVL2, 3 or 4 in HEK 293T cells all showed nELAVL phosphorylation, while
phosphorylation was absent when co-expressing CDKL5 and the phospho-deficient mutants
(Figure 1E). We also found that nELAVL phosphorylation was consistently reduced in the
visual, motor and frontal cortex of *Cdkl5* KO mice (Figure 1F & G). These results show that
nELAVL proteins are previously unidentified and *bona fide* CDKL5 phosphorylation targets.
Visual experience plays an essential role in V1 development, including the refinement
and matching of input from the contra and ipsi eyes[34]. To test whether phosphorylation of
nELAVLs by CDKL5 is experience-dependent, mice were kept in the dark for 16 hours and
re-exposed to light for 0, 1 or 2 hours. nELAVLs phosphorylation was significantly higher
after 1-hour of light re-exposure but not after 2-hour light re-exposure, suggesting the level of
nELAVLs phosphorylation is experience-dependent within a tight time window (Figure 1H
& I). Is experience-dependent phosphorylation in V1 specific to nELAVLs, or is this a
general feature of CDKL5 substrates? EB2 and MAP1S are two previously reported CDKL5
substrates that are involved in microtubule regulation[10]. We measured the phosphorylation
level of MAP1S and EB2 in V1, and found that the phosphorylation level of MAP1S and
EB2 were indeed decreased in *Cdkl5* KO mice compared to WT (Figure S1E & G). However,
MAP1S and EB2 phosphorylation levels did not show similar changes as nELAVLs
phosphorylation after 1-hour and 2-hour light re-exposure, suggesting MAP1S and EB2
phosphorylation in V1 is not experience-dependent *in vivo* (Figure S1F & H).
179
180 **CDKL5 deletion alters expression of activity-dependent genes in V1.**
181
Given that nELAVLs are RNA binding proteins involved in the regulation of mRNA stability
and translation[33], an important question is how the aberrant phosphorylation of nELAVLs
observed in CDKL5 deficiency impacts the expression of relevant mRNAs within mouse V1.
Thus, we first used bulk RNA sequencing to profile the transcriptomic differences between
WT and *Cdkl5* KO mouse V1. We identified 218 downregulated genes and 272 upregulated
genes in *Cdkl5* KO mice when compared to WT mice (Figure 2A & S2A; table S2). Eight
selected differentially expressed genes (DEGs) identified by bulk RNA sequencing, including
Arc, Fos, Npas4, Pcdh19, Homer1, Gad1, Sst, and Pvalb, were validated using real-time
qPCR (Figure S2B). All the DEGs were submitted to STRING to construct a Protein-Protein
interaction (PPI) network. The MCC algorithm in CytoHubba plugin was used to identify hub
genes in the PPI network. The top ten genes with the highest MCC scores, including *Fos*,
Nr4a1, *Egr1*, *Fosb*, *Junb*, *Dusp1*, *Egr2*, *Egr3*, *Fosl2* and *Btg2*, were visualized (Figure 2B).
These ten genes are all immediate early genes, suggesting that immediate early genes may
play an important role in CDD pathophysiology. Consistently, the Gene Set Enrichment
Analysis suggests that many activity-dependent neuronal functions, which are known to be
driven by visual experience in V1, are impaired in *Cdkl5* KO mice, including regulation of
synapse organization, glutamatergic synapses and dendritic spine development (Figure S2C).
To validate whether CDKL5 deficiency could lead to dendritic spine deficits in V1,
sparsely labeled layer 2/3 and layer 5 excitatory neurons were analyzed by applying epitope-

preserving Magnified Analysis of Proteome (eMAP) (Figure S3A-F). In layer 2/3 neurons,
we found that the total dendritic spine density of basal and apical dendrites was lower in
Cdkl5 KO mice. In contrast, in layer 5 neurons, we only observed a significant difference in
the spine density of basal dendrites (Figure S3G-J). To further investigate the changes in
spine morphology, we compared the density of mushroom-shaped, thin, and stubby spines as
well as filopodia between WT and *Cdkl5* KO mice. In layer 2/3 neurons, we found that
mushroom-shaped spine density was reduced in *Cdkl5* KO mice on both apical and basal
dendrites. In layer 5 neurons, only the density of mushroom-shaped spines on basal dendrites
decreased in *Cdkl5* KO mice while on apical dendrites, the density of filopodia was higher in
Cdkl5 KO (Figure S3G-J). These findings indicate that *Cdkl5* KO mice have altered spine
density and morphology on excitatory neurons, with a more pronounced effect observed in
layer 2/3 neurons.
To investigate potential differential effects on specific neuronal layers like layers 2/3
and 5, we conducted single nucleus (sn) RNA sequencing to compare the transcriptomes of
WT and *Cdkl5* KO V1. Nuclei from 6 snRNA libraries of V1 were collected from 6 WT and
6 *Cdkl5* KO mice and the gene expression matrices were filtered to remove low-quality cells
and cells containing a high proportion of mitochondrial transcripts (>5%). Doublets were
identified using Scrublet[35] and removed in the downstream analysis (Figure S4A) and
Harmony[36] was applied for batch correction (Figure S4B). We obtained 62051 high-quality
cells from the 6 snRNA libraries and combined them for the downstream analysis.
Dimensionality reduction and Leiden clustering were applied to generate different cell classes,
subclasses and types (Figure 2C & S4E-G). Classes consisted of glutamatergic neurons
(37289 cells), GABAergic neurons (7318 cells) and non-neuronal cells (17444 cells). The
UMAP of overlapping WT and *Cdkl5* KO libraries showed no gross change in cell types
(Figure S4C & D).
Next, we performed differential gene expression analysis between the nuclei from
WT and *Cdkl5* KO V1. Interestingly, we identified 69 activity-dependent genes among the
458 total DEGs in *Cdkl5* KO mice (Figure 2D; table S3), and ~50% of these overlapped with
activity-dependent DEGs identified in bulk RNA sequencing (Figure S4O), and among these
29 activity-dependent DEGs, 22 are overlapped with nELAVL-binding mRNAs identified
previously[33] (Figure S4P), again suggesting that activity-dependent plasticity might be
disrupted by loss of CDKL5. We found that CDKL5, and its phosphorylation targets the
nELAVLs, are mainly expressed in glutamatergic neurons (Figure 2E & S4H) which make
up the majority of cells in our dataset (Figure S4D). Therefore, we further analyzed
glutamatergic subclasses (Figure S4I), and plotted the number of DEGs in the individual
glutamatergic cell types (Figure 2F; table S4). L2/3, L4 and L6 cell types have the most
DEGs in *Cdkl5* KO mice. We intersected DEGs in different subtypes with activity-dependent
genes and found that significant enrichment among DEGs in L2/3 sublayers (Figure 2G).
L2/3 glutamatergic neurons exhibit spatial segregation and display graded gene
expression across three sublayers (A, B, and C). L2/3_A is adjacent to the pia, L2/3_C
borders L4 and L2/3_B is positioned between L2/3_A and L2/3_C. This segregation
represents an activity-dependent maturation of the layer and can be disrupted by dark
rearing[37]. The laminar distribution of these sublayer neurons has been suggested to related to
different downstream projecting targets and functions[38,39]. Given the deficits in layer 2/3
neurons identified through dendritic spine analysis, we were intrigued to investigate whether
transcriptomic maturation of layer 2/3 neurons is influenced by the deletion of CDKL5.
Therefore, we separated L2/3 glutamatergic neurons from WT and *Cdkl5* KO V1 into three
types (A, B, and C) (Figure S4J & K) according to the sublayer-specific marker
Cdh13, *Trpc6*, and *Chrm2*[37]. We found that the cell proportions across three sublayers of
neurons was unchanged (Figure S4L), but significant transcriptomic alterations were

observed. Again, we highlighted several activity-dependent downregulated DEGs (Figure 2H)
including *Homer1, Syt4, Tuba1a, Nptx2, Per1, Arc, Prickle1,* as well as the L2/3_A sublayer
marker *Cdh13,* which are all documented to associate with neuronal plasticity[40-47]. Among
the highlighted DEGs, *Homer1, Syt4, Tuba1a* were found to be common across all sublayers
of L2/3. *Cdh13* was uniquely identified as a DEG in sublayer L2/3_A, as expected, while
Nptx2 and *Per1* were specific DEGs in L2/3_B. Arc was found to be significantly
downregulated in L2/3_B and L2/3_C whereas *Prickle1* was exclusively identified as a DEG
in L2/3_C. These findings suggest distinct molecular signatures associated with each
sublayer, indicating potential disruptions in the specialized functional roles within L2/3
sublayers following CDKL5 deletion, without affecting the overall cell proportions of layer
2/3 neurons.
In addition, we also analyzed inhibitory neurons and found their composition also did
not show any significant difference between groups (Figure S4M). However, when
computing DEGs in different types of inhibitory neurons, we found that most DEGs were
specific to Pvalb+ inhibitory neurons (Figure S4N).
To validate whether the activity-dependent DEGs identified in bulk and snRNA
sequencing are decreased in *Cdkl5* KO mice, we used fluorescence in-situ hybridization
(FISH). We found that mRNA expression levels of *Fos*, *Homer1* and *Arc* were indeed
significantly reduced in *Cdkl5* KO mice while *Camk2a*, an excitatory neuron marker, was
unaltered (Figure 2I & J). Immunostaining results also showed that the expression of c-fos
was decreased significantly in layers 2/3, 4 and 6 in *Cdkl5* KO mice (Figure S4Q & R). Of
note, the *Fos* mRNA punctum volume in *Cdkl5* KO mice appeared larger than those in WT
mice (Figure 2I & J).
Taken together, the structural abnormalities in dendritic spines (Figure S3) align with
the decreased expression of activity-dependent genes and excitatory synaptic mRNA as
revealed in our transcriptomic analysis (Figure 2A). Notably, 21 out of 29 dysregulated
activity-dependent genes (highlighted in Figure 2D), including *Fos* and *Homer1*, are direct
binding targets of nELAVLs identified through CLIPseq analysis[33]. These observations led
us to postulate that the CDKL5-mediated phosphorylation of nELAVLs modulates the
mRNA levels of activity-dependent genes, thereby influencing the refinement of the visual
circuit.
282
283 **Loss of CDKL5-mediated phosphorylation alters nELAVLs phase separation**
284
We next sought to investigate how phosphorylation of nELAVLs by CDKL5 affects mRNA
expression in mouse V1. While CDKL5 itself was found to undergo phase separation[48], we
therefore evaluate the phase separation propensity of CDKL5's kinase target nELAVLs. We
employed the LLPhyScore algorithm[49] to analyze a composite dataset comprising these target
sequences alongside a benchmark set of experimentally verified LLPS-positive drivers (e.g.,
FUS, TDP-43, TIA-1) and a broad background of negative control sequences (Figure S5).
The comparative ranking analysis indicates that nELAVL proteins possess a significant
intrinsic potential for phase separation, distinct from non-phase-separating background
proteins. While canonical "scaffold" proteins such as FUS and TDP-43 cluster at the apex of
the distribution with near-maximal scores ($\approx$1.0) and top-tier rankings (Rank < 200), the
nELAVLs occupy a prominent intermediate-to-high probability tier. nELAVLs exhibit scores
ranging from approximately 0.70 to 0.85, positioning them within the top quartile of the
distribution (approximate ranks 700–1300) (Figure S5). Although they do not reach the
saturation levels of primary drivers, their significant deviation from the tail of the distribution
suggests they encode sufficient multivalent interaction valency to participate in phase
separation, likely functioning as client proteins recruited into condensates or driving phase

transitions under specific physiological conditions rather than acting as constitutive scaffolds.
For example, post-translational modification like phosphorylation can finetune the phase
separation features of proteins[50]. To examine whether nELAVLs show different protein
distribution with or without CDKL5, we employed expanded V1 tissue of WT and *Cdkl5* KO
mice by eMAP. We immunolabelled nELAVLs using a pan-nELAVLs antibody and
ELAVL3 using a ELAVL3 specific antibody available (Figure S1D). We found that the
immunolabelled signals formed cytoplasmic puncta in V1 neurons (Figure 3A). Although the
number of puncta in each neuron showed no significant change, the volume of these puncta
were significantly increased in volume in *Cdkl5* KO V1 neurons (Figure S6A). By fraction
analysis, we also showed that the volume fraction distribution of these puncta in each neuron
has significantly right shifted in *Cdkl5* KO mice, providing evidence that each of the KO V1
neurons contains larger nELAVL punta as compared to WT V1 neurons (Figure 3B). A
similar phenotype was observed in the *CDKL5* knockdown primary neurons (Figure S6B &
C). These unexpected findings suggested that nELAVLs are involved in droplet-like puncta
in cells, and the punctum size is regulated by phosphorylation of nELAVLs by CDKL5.
To test the dependence of droplet-formation on nELAVL phosphorylation by CDKL5,
we employed phospho-mimetic and phospho-deficient mutations of nELAVLs (SE and SA
mutants with the consensus CDKL5 phosphorylation motif of -RPSSA- was substituted with
-RPSEA- and -RPSAA-, respectively), to evaluate their capacity to form the droplet-like
puncta. We overexpressed GFP-tagged nELAVL-SE and nELAVL-SA in both primary
neurons and HEK 293T cells (Figure 3C & S6D, G & I). We first verified that the exogenous
nELAVL proteins showed equal expression by immunoblotting (Figure S6E). Subsequently,
through stimulated emission depletion (STED) microscopy, we visualized the cellular
distribution of nELAVL proteins and showed that the volume of ELAVL3-SA and ELAVL4-
SA puncta were significantly larger than that of ELAVL3-SE and ELAVL4-SE punta,
respectively, in both primary neurons and HEK 293T cells (Figure 3C & S6F-J), supporting a
critical role of CDKL5-mediated phosphorylation in regulating nELAVL punctum size.
The liquid-like nature of the nELAVL puncta was corroborated through live-cell
imaging. As we observed similar droplet-like puncta among three nELAVLs, we selected
ELAVL3 as a key example to demonstrate their biophysical property. Smaller ELAVL3-SA
puncta were found to rapidly coalesce into larger ones over time, suggesting a liquid-like
nature (Figure 3D). To evaluate the fluid-like property of the nELAVL puncta, fluorescence
recovery after photo-bleaching (FRAP) were performed in HEK 293T expressing ELAVL3
SA and SE mutants (Figure 3E & F). The FRAP assays showed that both the SA and SE
mutants within these puncta were capable of exchanging with the surrounding cytoplasm. It
is worth noting that the recovery rate of SA was much slower than that of SE; within 40
seconds, the recovery rate of SA was approximately 35%, while that of SE was around 60%
(Figure 3E & F). These data indicated that nELAVL is a member of cytosolic condensates
and that its phosphorylation state modulates the properties of this condensate.
We next purified the recombinant full-length ELAVL3 protein and sparsely labeled
the purified ELAVL3 with the Cy3 fluorophore. Cy3-labeled ELAVL3 protein formed
spherical droplets in solution in a concentration-dependent manner (Figure 3G). These
droplets fused quickly with each other over time (Figure 3H). The number of ELAVL3
droplets decreased when the salt concentration was elevated (Figure 3I), indicating that
electrostatic interaction may contribute to the droplet's formation. The phase diagram
demonstrated that the droplets could form at the minimal concentration of 2.5μM, which is
close to physiological condition (Figure 3J). The *in vitro* FRAP experiment showed that the
ELAVL3 droplets exhibited recovery subsequent to photobleaching through protein exchange
between the condensed droplets and surrounding solution (Figure 3K), suggesting that the
ELAVL3 droplets possess properties characteristic of a phase separated state.

Taken together, these biophysical features of nELAVLs are reminiscent of biomolecular condensates in biological systems, indicating that nELAVLs can undergo phase separation *in vitro* and in cells. Moreover, their phase separation characteristics can be modulated by the CDKL5-dependent phosphorylation.

CDD patient iPSC-derived neurons form aberrant nELAVLs condensates and altered crosstalk with P-bodies and stress granules.

To investigate whether nELAVL condensates behavior is altered in the context of CDD, we aimed to characterize the nELAVL condensates in female patient-derived iPSC lines carrying the pathogenic variant 1648C>T (p.R550*) of CDKL5. As controls, we utilized a WT allele-expressing control line from the same patient, as well as another base editing-corrected isogenic control line (Figure 4A). We confirmed that CDKL5 expression was absent in the R550* line-derived neurons (iNeurons) (Figure 4B), rendering it a comparable model to KO. CDKL5 expression was restored in the isogenic control line (Figure 4C). As expected, the phosphorylation levels of nELAVLs were significantly reduced in R550* iNeurons and rescued in the isogenic line (Figure 4D & E). STED images of iNeurons revealed that the volume of ELAVL3 and nELAVL condensates were both significantly increased in the R550* iNeurons than in the control line, and this phenotype was rescued in the isogenic line (Figure 4F & G & S7A). These results demonstrated that nELAVL condensates is regulated by CDKL5-mediated phosphorylation in human CDD iNeurons, suggesting that our observations from CDKL5 deficient mouse neurons are translatable.

RNP granules have emerged as crucial regulators of many cellular processes, including transcription and translation[51]. Among the RNP granules, P-bodies and stress granules are well documented to regulate RNA processing and are dysregulated in human diseases, including several neurodevelopmental disorders[52]. In addition, aberrant RBPs co-condensation has been reported in neurodegenerative conditions[14]. We therefore asked whether dysregulated nELAVL condensates may trigger aberrations in other RNP condensates, such as P-bodies and stress granules to influence RNA processing. To test this, we co-immunolabelled nELAVLs with P-body marker GW182 and stress granules marker TIA1 in CDD iNeurons. We found that P-bodies and stress granules volumes were significantly increased in R550* group and such abnormal increase was rescued in isogenic control group (Figure 4H & I & S7B-D).

To further investigate the impact of nELAVL phosphorylation on P-bodies and stress granules dynamics, we co-expressed ELAVL3-SA with GW182 or TIA1 in HT22 cells (Figure S7E & G). Similar to the effects observed in iNeurons, the expression of ELAVL3-SA led to an increased size of GW182 and TIA1 puncta, while the expression of ELAVL3-SE led to decreased size (Figure S7E-H). We next hypothesized that the phosphorylation status of ELAVL3 modulates its interaction with GW182. To test this, we performed co-immunoprecipitation and found that the interaction between GW182 and ELAVL3-SE was significantly stronger than that with the phospho-dead ELAVL3-SA mutant (Figure 4J & K). These results suggest that phosphorylation of nELAVLs not only mediates the formation of their own condensates but also modulates their interaction with other biomolecular condensates, such as P-bodies.

CDKL5 modulate mRNA processing by regulating nELAVL condensates

P-bodies and stress granules are RNP granules known to regulate mRNA storage, decay, and translation[52], and dynamically sequester mRNAs as a regulatory site during cellular processes[53-55]. To study the impact of nELAVL phosphorylation on RNA processing,

we employed FISH assay and puro-PLA assay to examine the mRNA level and translation of
Fos, a known binding target of nELAVLs[56]. We found reduced *Fos* mRNA puncta in
ELAVL3-SA expressed primary neurons compared to ELAVL3-SE expressed neurons
(Figure 5A& B). From the puro-PLA assay, we dual-labelled puromycin and Fos protein after
the puromycin treatment and found reduced *Fos* translation in ELAVL3-SA expressed
primary neurons compared to ELAVL3-SE expressed neurons (Figure S7I & J).
Cell-free Electrophoretic Mobility Shift Assay (EMSA) reveals different binding
affinities between nELAVL-SE and SA to *Fos* RNA. The recombinant nELAVL proteins
formed protein-RNA complexes with *Fos*-Cy5 RNA (Figure 5C). The binding affinity of
nELAVL-SE to *Fos* was significantly higher than that of nELAVL-SA to *Fos*, indicating that
the phosphorylation status of nELAVLs modulates their binding affinity to target RNA
(Figure 5C).
To test whether the mRNA stability was altered, we used Actinomycin D to inhibit
the transcription at different time point (0, 5, 10, 30, 60 mins) before FISH assay on iNeurons.
We found that in R550* iNeurons, the half-life of *Fos* mRNA was significantly shorter than
in Control iNeurons (Control: 10.61 vs. R550*: 3.304), suggesting increased mRNA
degradation in CDD iNeurons (Figure 5 D & E).
Taken together, these findings provide evidence for a mechanistic link between
CDKL5-mediated nELAVL phosphorylation and the formation and function of RNP granules
which plays important role in mRNA processing.
421
422 **CDKL5 deletion impairs visually guided behavior and visually evoked functional**
423 **responses**
424
To determine whether the *Cdkl5* KO mice exhibits behavioral deficits related to cortical
visual impairment, we employed the visual cliff task[57,58], which is designed to detect
impairments in stereoscopic depth perception. In this task, animals were placed on a central
platform within an apparatus containing two distinct areas: a "safe side" and a "cliff side"
(the later consisting of a glass plate positioned above a sharply dropped, patterned floor)
(Figure 6A, left). WT mice with intact stereoscopic vision consistently avoided the "cliff"
side, descending approximately 80% from the "safe" side (Figure 6A, right). On the other
hand, we found that *Cdkl5* KO mice descended significantly more times from the 'cliff' side,
suggesting impaired stereoscopic depth perception (Figure 6A, right).
434
To investigate how CDKL5 deletion influences neuronal responses and computations in the
mouse binocular visual cortex (bV1), we conducted two-photon calcium imaging of layer 2/3
neurons expressing the calcium indicator GCaMP6f. The visual stimuli were drifting
sinusoidal gratings (8 directions, with 45° increments[59], 0.08, 0.04 and 0.02 cycle per degree
(cpd) at 2Hz temporal frequency) with a 3-second grey screen between grating stimuli,
followed by 4-second of natural scenes (Figure 6B). During imaging, mice viewed drifting
grating and natural scenes stimuli presented independently to each eye (contralateral [contra]
or ipsilateral [ipsi] only eye viewing) or to both eyes (binocular viewing) (Figure 6C & D &
J). Consistent with established findings, we observed bV1 neurons exhibited varying
responses and preferences for stimulus direction to binocular, ipsi eye, or contra eye viewing
(Figure 6C & D & J)[60,61]. WT and *Cdkl5* KO mice did not differ in their percentage of
binocularly responsive or monocularly responsive neurons (Table S5). Individual neurons in
V1 are often preferentially activated by specific orientations of grating. Therefore, we next
measured these preferences, quantified as an orientation selectivity index (OSI). *Cdkl5* KO
mice showed robust and significant decrease in OSI across all three imaging conditions
(binocular, contra eye only and ipsi eye viewing) at high spatial frequency (0.08cpd) as

compared to WT mice, suggesting that they are overall less selective (Figure 6E). A significant difference in OSI was observed across all three tested spatial frequencies, with 0.08cpd having the largest OSI decrease in *Cdkl5* KO mice compared to WT (Figure 6E, S8A & K). There is no difference in neuron response amplitude between WT and *Cdkl5* KO mice except ipsi eye viewing condition with 0.02cpd (Table S6).These results suggested that the cortical visual deficits in *Cdkl5* KO V1 neurons are more sensitive to higher spatial frequency (Figure S8K).

Neurons in bV1 that receive inputs from both eyes (contra+ipsi responsive) match specific visual responses from each eye to integrate their visual information, thereby establishing an accurate representation of binocular visual space. The binocular neurons are crucial to visual functions such as disparity and depth perception[62], and the matching of visual information is highly sensitive to disruption of visual experience or plasticity in development[63,64]. We therefore examined if binocular matching was altered upon CDKL5 deletion. We first examined the relationship between OSI and orientation matching (measured by pairwise signal correlations of their calcium responses between the contra eye and ipsi eye) in WT mice (Figure 6F, left). We observed that neurons with a high OSI (OSI>0.5) in both their contra eye and ipsi eye responses exhibited stronger contra-ipsi orientation matching (blue region in Figure 6F, left). On the other hand, neurons with a low OSI (OSI<0.5) in both their contra eye and ipsi eye responses exhibited weaker contra-ipsi orientation matching (red region in Figure 6F, Left), consistent with published findings[60,65]. In *Cdkl5* KO mice, we observed a significantly reduced fraction of neurons with high OSI and high contra-ipsi orientation matching (blue region in Figure 6F & G) and an increased fraction of neurons with low OSI and low contra-ipsi orientation matching (red region in Figure 6F & G) as compared to WT mice. This suggested that there was a specific impairment of ipsi eye alignment to contra and binocular orientation preferences in *Cdkl5* KO bV1 neurons. Additionally, we examined the difference in preferred orientation between contra and ipsi responses among binocular neurons in WT and *Cdkl5* KO mice. Of note, contra-ipsi (C/I) response correlation and difference in preferred orientation between contra and ipsi responses are two negatively correlated measurements to examine binocular matching (Figure S8B). We observed that *Cdkl5* KO mice showed significant decrease in C/I responses correlation and significant increase in difference in preferred orientation in grating stimuli at spatial frequency 0.08cpd (Figure 6H & I). Consistent with the OSI results, a milder binocular matching deficit was observed in *Cdkl5* KO bV1 neurons at spatial frequency 0.04cpd (Figure S8C-F), and no significant difference was observed at 0.02cpd (Figure S8G-J). In addition, bV1 neurons in *Cdkl5* KO mice exhibited decreased C/I correlation in response to natural scene stimuli, which represent a more complex visual stimulus class (Figure 6K). In sum, the deletion of CDKL5 results in decreased orientation selectivity and contra-ipsi response matching in the bV1 neurons. These deficits suggest there is impaired integration of ipsi eye input, which matures after contra eye inputs are already established and is experience dependent[66].

CDKL5 deletion disrupts circuit-wide visual information processing

To investigate how these single-neuron changes affected the binocular circuit, we computed the information encoding at the network level. We applied a supervised support vector machine (SVM) decoder to estimate the information encoding at a populational level, discriminating all four orientations or two natural scenes in our imaging paradigm. The decoding accuracy in the *Cdkl5* KO mice was consistently lower than in the WT mice at fixed population sizes (Figure S8M), suggesting a reduction in visual information in *Cdkl5* KO mice. To evaluate the binocular matching in the population level, we also trained a

decoder based on contra only responses and applied this decoder to the ipsi only responses from the same neuronal populations (Figure S8L). At a spatial frequency of 0.08 and 0.04cpd, contra trained decoding accuracy of ipsi responses was significantly lower in *Cdkl5* KO mice than in WT mice, suggesting that visual information contained in the contra responses was less well matched to ipsi responses in *Cdkl5* KO mice. Thus, the deletion of CDKL5 impairs the encoding of visual information at network level, with more pronounced deficits in ipsi eye responses and matching.

Fine balance of nELAVLs phosphorylation level is necessary for normal visual processing

Finally, to determine if cortical visual deficits in *Cdkl5* KO mouse bV1 could be recapitulated by altering nELAVL phosphorylation, we again performed *in vivo* two-photon calcium imaging in layer 2/3 neurons in the bV1 of WT mice injected with virus containing the nELAVL-WT (WT), the nELAVL-SE phospho-mimetic (SE) or nELAVL-SA phospho-deficient (SA), and also compared with the group without expressing exogenous nELAVL proteins (Ctrl) (Figure 6L & S9A). We found no change in the overall proportion of responsive neurons across the different viewing conditions (Table S5). However, OSI was decreased in each viewing condition for nELAVL-WT, SE and SA injected mice when compared to Ctrl, with OSI lower in SA than in SE injected mice for all viewing conditions in grating stimuli at spatial frequency 0.08, 0.04 and 0.02cpd (Figure 6M & S9B & E), while neuron response amplitude is no difference between SE and SA injected mice (Table S6). Intriguingly, WT injected mice showed higher OSI than SA mice but lower than the OSI of SE mice, suggesting nELAVL phosphorylation level modulates orientation selectivity. We next examined binocular matching in these mice. We observed that both WT and SA injected mice showed decrease in C/I response correlation and increase in difference in preferred orientation in grating stimuli at spatial frequency 0.08cpd, as compared to Ctrl (Figure 6N & O). On the other hand, no significant difference was observed between SE-injected mice and Ctrl group (Figure 6N & O). These suggested that overexpression of nELAVL-WT or their phosphodeficient mutant, but not the phosphomimetic mutant, led to impairment of orientation matching. Similar to *Cdkl5* KO mice, a mild effect was observed at spatial frequency 0.04cpd and no significant difference was observed at spatial frequency 0.02cpd (Figure S9C-D & F-G). For the more complex natural scene stimuli, C/I response correlation was decreased in nELAVL-WT, SE and SA injected mice when compared to Ctrl (Figure 6P), suggesting that binocular matching of complex stimuli is more sensitive to the expression level and phosphorylation status of nELAVLs. At the population level, decoding accuracy was decreased across viewing conditions in WT, SE and SA injected mice at spatial frequency 0.08cpd and natural scene when compared to Ctrl (Figure S9H & I). In sum, expression of nELAVL proteins with different phosphorylation status in V1 are sufficient to replicate the cortical visual deficits observed in *Cdkl5* KO mice.

Discussion

Sensory processing deficits are a prevailing feature in numerous syndromic developmental disorders. Within the spectrum of developmental disorders, individuals with CDD are notable for their high rate of cortical visual impairment. In this study, utilizing a CDD mouse model, we delineated visual processing deficits of orientation tuning, interocular and monocular-binocular matching, and information encoding in the bV1. Our findings indicate that fundamental features of cortical visual processing, both at the single-neuron and network levels, is compromised in CDD mice, potentially impacting higher-order processing stages

encompassing recognition, identification, and memory for visual patterns[67]. The binocular orientation mismatch observed in CDD mice likely diminishes their ability to differentiate similar stimuli and hinders the accurate representation of visual space.

At eye-opening in mice at P13-14, bV1 is predominantly composed of contra responsive cells which are already orientation selective[34], while ipsi and contra+ipsi responsive cells emerge after eye-opening. The contra+ipsi responsive population later becomes orientation matched by experience-dependent plasticity during the ocular dominance critical period[61]. Our work reveals that in adult *Cdkl5* KO mice, both orientation selectivity and contra-ipsi orientation matching compromised, with more severe deficits in ipsi driven responses. L2/3 excitatory neurons in bV1 have been shown to undergo structural synaptic remodelling during the critical period dependent on experience-driven changes in gene expression[68]. We found alterations in L2/3 synaptic density and morphology, as well as the transcriptomic profile of *Cdkl5* KO mouse V1, with gene expression changes most prominent in L2/3 (Figure 2F-H). While the overall cell composition of V1 did not change, mRNA for the canonical immediate early gene *Fos*, a marker for neuronal activation and a direct binding target of nELAVLs, was identified as a hub gene for downregulated activity-dependent DEGs in *Cdkl5* KO mice (Figure 2B). Together, our data suggests that L2/3-mediated circuit impairment in *Cdkl5* KO mice is caused by dysregulated activity-dependent gene expression through the loss of CDKL5-dependent phosphorylation of nELAVLs. The deficits in contra-ipsi matching towards the contra eye further suggest that CDKL5-dependent phosphorylation of nELAVLs is most crucial during the developmental ocular dominance critical period. Given that cortical visual impairment is a well-reported clinical feature in CDD patients and has been suggested to be employed as a functional biomarker of CDD[6,26], it will be of clinical interest to assess whether CDD patients also experience deficits in interocular integration.

Despite the identification of several physiological substrates, it remained unclear how CDKL5 dysfunction leads to cortical visual impairment at the molecular, cellular and neural circuit levels. Here we identified nELAVLs as direct substrates of CDKL5 and demonstrated that CDKL5-mediated phosphorylation can affect the propensity of nELAVLs to undergo phase separation. Our FRAP assay revealed a decreased exchange rate within ELAVL3 condensates in the absence of CDKL5-mediated phosphorylation, suggesting a more compact internal structure and less dynamic property of the condensates. CDKL5-dependent phosphorylation introduces negatively charged phosphate groups into the RPSS*A motif on nELAVLs, modulating charge-patterning critical for phase separation. This charge redistribution disrupts the balance of attractive (e.g., cation-π, π-π) and repulsive forces within nELAVLs, akin to phosphorylation-driven phase separation regulation in other RBPs including FUS and TDP-43 are known to undergo phase separation[69]. The negative charge introduced by phosphorylation may also disrupt hydrogen bonding with adjacent arginine/lysine residues[70], altering RNA-binding domain conformation. Additionally, as activity-dependent DEGs, *Fos* and *homer1* are direct nELAVLs binding targets[33], and our data demonstrated lower mRNA level of these DEGs, low RNA:RBP ratios may also modulate condensate size[71]. Given that impaired nELAVL phosphorylation was also observed in the frontal cortex and motor cortex of Cdkl5 KO mice, it will be of great interest to examine the roles of nELAVLs dysregulation in other clinically relevant phenotypes of CDKL5 deficiency, including motor and social deficits.

Our data highlight a CDKL5-mediated phosphorylation switch of nELAVLs that regulates the crosstalk between the nELAVL condensates and other biomolecular condensates. Stress granules and P-bodies are RNP granules that regulate mRNA processing[52-55]. The mRNAs sequestered within stress granules or P-bodies can be translationally repressed and these mRNAs may either re-enter the translation process after

granule disassembly[53,54,72,73] or undergo RNA decay[74-77]. In CDD models as well as in scenarios where nELAVLs are non-phosphorylated, we observed reduced levels of nELAVLs-binding mRNA concomitant with an augmentation in the volume of nELAVLs, stress granules, and P-bodies puncta. These results suggest that the loss of CDKL5 function in CDD patients leads to a downregulation of nELAVL phosphorylation, which in turn results in abnormal phase separation of nELAVLs. Such aberrant nELAVL phase separation then contributes to dysregulated stress granules and/or P-bodies, which sequester and prevent the activity-dependent translation of key mRNAs essential for synaptic plasticity, ultimately contributing to impaired refinement of neural circuits during the pathological, developmental progression of CDD (Figure S10). To elucidate how aberrant RNP condensates—including P-bodies and stress granules—coordinately dysregulate mRNA expression and translation in CDKL5 deficiency will be our future direction.

Altered dynamics of stress granules and P-bodies has been suggested to contribute to the pathology of several neurodegenerative conditions including amyotrophic lateral sclerosis, frontotemporal dementia, Alzheimer's disease and Parkinson's disease[78-80]. Genetic and functional studies also suggest that nELAVLs, stress granules, and P-bodies are strongly implicated in neurodevelopmental disorders. ELAVL3 and ELAVL4 null mice display deficits in learning, motor function, and neuronal maturation[81-83]. Differential gene expression analysis in ELAVL2 knockdown primary human neurons also reveals autism related alterations[84]. ELAVL3 has also been identified as an autism-associated gene in multiple large-scale genetic studies[85-87], and both ELAVL2 and ELAVL3 are listed as autism candidate genes by the Simons Foundation Autism Research Initiative. Similarly, pathogenic variants of genes encoding for stress granules and P-bodies core proteins such as G3BP1, UBAP2L and TNRC6B have been identified in patients with neurodevelopmental disorders[88,89]. Given the pivotal roles of nELAVLs in a wide range of neurological disorders, it is reasonable to speculate that the dysregulation of CDKL5-mediated phosphorylation of nELAVLs, and the subsequent formation of aberrant nELAVL condensates, may also contribute to the etiology of other diseases.

In summary, our findings shed light on the physiological significance of CDKL5-mediated nELAVLs phase separation, as well as their interplay with other biomolecular condensates, in the context of activity-dependent visual circuit refinement. These findings deepen our understanding of the link between CDKL5 loss of function and cortical visual impairment in CDD, and potentially other pathology. Crucially, our discovery also opens up possibilities for exploring how phosphorylation governs the crosstalk among various biomolecular condensates which have broad and profound implications for normal brain function as well as neurodevelopmental and neurodegenerative disorders.

Acknowledgments

We thank Gavin Siu, Josie Lai, Joaquim Vong and Cara Kwong for their technical assistance. We also thank Core Laboratories, School of Biomedical Sciences, the Chinese University of Hong Kong as well as Biosciences Central Research Facility, the Hong Kong University of Science and Technology for their technical support. We thank Hei Man Kim Chow for sharing HEK 293T and HT22 cells. We thank Hovy Wong and Kwok-On Lai for critical comments on the manuscript.

Author contributions

S.Y. performed surgeries, two-photon imaging, phosphoproteomic and eMAP experiments. Y.Z. performed RNA extraction, real-time PCR, FISH, bulk and snRNA-seq data analysis. Zhongyu Z. performed cell line, primary neuron and iPSC experiments, STED imaging and quantification. H.Y. processed and analyzed the two-photon imaging data. Zhongjie Z. carried out population analysis for two-photon imaging data. K.J., K.T., G.H., J.Z., Marco C. and L.T. contributed to two-photon imaging data analysis. Maggie C. and Y.C. performed Western blot, with inputs from H.H.. Y.D., S.L. and A.F. contributed to snRNA-seq experiments. C.H., H.T. and Zhongyu Z. performed FRAP experiments. H.L. and H.K. provided expertise on the FISH experiment. J.L., A.K., S.F., Q.C. and S.T. contributed to snRNA-seq analysis. T.K. provided expertise on eMAP experiments. A.C. and K.L. provided expertise on cellular condensates experiments. C.G. and B.N. computed the phase separation propensity of nELAVLS. Y.W. and J.Z. performed the in vitro phase separation assay. J.I. conceived and supervised all aspects of the experiments. S.Y., Y.Z., Zhongyu. Z., H.Y. and J.I. wrote the paper. All authors edited the manuscript.

Funding

This work was supported by Brain Science and Brain-like Intelligence Technology-National Science and Technology Major Project (2022ZD0214400; J.I., J.Z), Research Grants Council (RGC) of Hong Kong SAR (C4005-24Y, J.I.; 24117220, J.I.), a research grant from the University of Pennsylvania Orphan Disease Center on behalf of the Loulou Foundation (J.I.; K.L.), Guangdong Basic and Applied Basic Research Foundation (2022B1515130007; J.I.), the National Research Foundation of Korea (RS-2023-00264980; T.K.), European Union's Horizon research and innovation programme grant agreement No. 101206609, Marie Sklodowska-Curie Global Fellowship - AstroError (M.C.), Lo Kwee-Seong Biomedical Research Fund (J.I.), Faculty Innovation Awards (FIA2020/A/04) from the Faculty of Medicine, CUHK (J.I.) and Gerald Choa Neuroscience Institute Research Program Fund (J.I.).

Declaration of interests

The authors declare no competing interests.

Methods

Key Resources Table

Antibodies	Source	Catlog #
Rabbit anti-ELAVL2	Proteintech	14008-1-AP
Rabbit anti-nELAVL pS	Biomatik	custom-made
Rat anti-EB2	Abcam	ab45767

Rabbit anti-EB2 pS222	Covalab	00117741
Rabbit anti-MAP1S	Sigma	HPA050934
Rabbit anti-MAP1S pS812	Covalab	00118044
Mouse monoclonal anti-GAPDH	Invitrogen	AM4300
Mouse monoclonal anti-CDKL5	Santa Cruz	sc-376314
Rabbit monoclonal anti-c-fos	Synaptic Systems	226008
Chicken polyclonal anti-GFP	Invitrogen	A10262
Rabbit polyclonal anti-ELAVL3 (HuC)	Proteintech	55047-1-AP
Mouse monoclonal anti-ELAVL4 (HuD)	Santa Cruz	sc-48421/sc-28299
Mouse monoclonal anti-GW182	Santa Cruz	sc-56314
Mouse monoclonal anti-TIA1	Santa Cruz	sc-166247
Goat anti-chicken, Alexa 488	Invitrogen	A32931
Goat anti-rabbit, Alex 555	Invitrogen	A32732
Goat anti-mouse, Alex 647	Invitrogen	A32728
Goat anti-rabbit, Alex 647	Invitrogen	A32733
Abberior STAR ORANGE	Abberior	STORANGE
Abberior STAR RED	Abberior	STRED

Bacterial and virus strains	Source	Catlog #
rAAV-hSyn-GCaMp6f-WPRE-hGH polyA	Brain VTA	PT-0147
AAV9-hSyn-nElavl-P2A-DsRed-WPRE-BGH-polyA	Vigene	/

AAV9-hSyn-nElavl(SE)-P2A-DsRed-WPRE-BGH-polyA	Vigene	/
AAV9-hSyn-nElavl(SA)-P2A-DsRed-WPRE-BGH-polyA	Vigene	/
pCAG-FLEX-EGFP-WPRE	Addgene	51502
pENN.AAV.CamKII 0.4.Cre.SV40	Addgene	105558

Experimental models: Organisms/strains	Source	Catlog #
Mouse: *Cdkl5* KO	The Jackson Laboratory	Strain #:021967; RRID:IMSR_JAX:021967
Mouse: C57BL/6J	Laboratory Animal Services Centre, The Chinese University of Hong Kong	/

Software and algorithms	Source	Catlog #
MATLAB	Mathworks	https://www.mathworks.com/products/matlab.html
GraphPad Prism 9	GraphPad Software Inc	https://www.graphpad.com/scientific-software/prism/
Fiji	Open source	https://imagej.net/software/fiji/
Suite2p	Open source	https://suite2p.readthedocs.io/en/latest/
Imaris	Oxford Instruments	https://imaris.oxinst.com/
Python	Anaconda3	https://www.anaconda.com
R	R version 4.1.2	https://www.r-project.org/

685
686 **Resource availability**
687
688 *Leading contact*
689 Further information and requests for resources and reagents should be directed to and will be
690 fulfilled by the lead contact (jacqueip@cuhk.edu.hk).
691
692 *Materials availability*
693 This study did not generate new unique reagents.
694
695 *Data and code availability*

696 Any additional information to reanalyze the data reported in this paper is available upon
697 request. All raw and processed snRNA-seq datasets reported in this study will be available
698 before publication. Analysis code generated during this study are available upon reasonable
699 request.
700
701

Methods details

Mouse line

Cdkl5 KO mouse line[5] (B6.129(FVB)-*Cdkl5 tm1.1Joez /J,* strain no: 02196, Jackson
Laboratory) was imported from the Jackson Laboratory. A colony was maintained at the
Laboratory Animal Services Centre of the Chinese University of Hong Kong and housed in
the Animal Holding Core of the School of Biomedical Sciences, the Chinese University of
Hong Kong. *Cdkl5* heterozygous females (+/-) were used as breeders with C57BL/6 wildtype
males to produce WT and *Cdkl5* KO littermates. Our study mainly focused on investigating
hemizygous male mice as our model to avoid potential confounds induced by mosaic
expression of CDKL5 in heterozygous females resulting from random X chromosome
inactivation. C57BL/6 mice were used for nELAVLs, nELAVLs-SE, nELAVLs-SA virus
injection for two-photon calcium imaging. Mice were group-housed (no more than five mice
per cage) with a standard light/dark cycle of 12/12 hours with access to food and water *ad
libitum*. All experimental procedures were approved by the Animal Experimentation Ethics
Committee of The Chinese University of Hong Kong.

Genotyping

A PCR-based method was used to detect deletions in *Cdkl5* exon 6 for genotyping (Phire
Tissue Direct PCR Master Mix, Thermo Fisher Scientific). Dilution Buffer and DNA Release
from the genotyping kit were mixed for genomic DNA extraction for 5 minutes at 98 °C, 20
µl Dilution Buffer and 1 µl DNA Release for each sample. 1 µl of genomic DNA extracted
from the ear tissue was used for PCR. Primers for genotyping are forward: 5'-
CCACCCTCTCAGTAAGGCAGCAG-3' and reverse: 5'-
GTCCTTTTGCCACTCAATTCCATCC-3'[5]. PCR reactions contained 0.5 µl of each primer,
10 µl 2X Phire Tissue Direct Master Mix and 8 µl nuclease-free water. PCR was performed
using the following program: 1) 95°C for 3 min, 2) 98°C for 20 s, 3) 60°C for 15 s, 4) 72°C
for 1 min, 5) 72°C for 2 min, 6) 4°C for 10 minand finish. Repeat steps 2-4 for 27 times. The
WT allele showed a 653 base pair product, and the *Cdkl5* KO allele showed a 305 base pair
product.

Phosphoproteomics

V1 of 4 pairs WT and of *Cdkl5* KO littermates (8-week old) were processed for TMT
labeling by Shanghai Applied Protein Technology Co., Ltd. For protein extraction and
digestion, tissues were separately lysed in SDT buffer (4% SDS, 100 mM Tris–HCl, and 1
mM DTT [pH 7.6]) and quantified using the BCA Protein Assay Kit. Digested peptides of
each sample were sonicated by C18 Cartridges and concentrated by vacuum centrifugation,
followed by reconstituting in 40 µl of 0.1% (v/v) formic acid. 280 nm UV light spectral
density was used to estimate the peptide content. The peptide mixture of each sample (100 µg)
was labeled with TMT (Thermo Fisher Scientific) reagent. Phosphorylated peptides were
subjected to TiO2 bead-based enrichment (90% enrichment efficiency) followed by LC-
MS/MS analysis in Thermo Scientific Q Exactive HF mass spectrometer (Thermo Scientific)
that was coupled to Easy nLC (Proxeon Biosystems, now Thermo Fisher Scientific) for 120
min. The peptides were loaded onto a reverse phase trap column (Thermo Scientific Acclaim
PepMap100, 100 µm*2 cm, nanoViper C18) connected to the C18-reversed phase analytical

column (Thermo Scientific Easy Column, 10 cm long, 75 μm inner diameter, 3μm resin) in
buffer A (0.1% Formic acid) and separated with a linear gradient of buffer B (84%
acetonitrile and 0.1% Formic acid) at a flow rate of 300 nl/min controlled by IntelliFlow
technology. The mass spectrometer was operated in positive ion mode. MS data was acquired
using a data-dependent top10 method dynamically choosing the most abundant precursor ions
from the survey scan (300–1800 m/z) for HCD fragmentation. Automatic gain control (AGC)
target was set to 3e6, and maximum inject time to 10 ms. Dynamic exclusion duration was
40.0 s. Survey scans were acquired at a resolution of 70,000 at m/z 200 and resolution for
HCD spectra was set to 17,500 at m/z 200, and isolation width was 2 m/z. Normalized
collision energy was 30 eV and the underfill ratio, which specifies the minimum percentage
of the target value likely to be reached at maximum fill time, was defined as 0.1%. The
instrument was run with peptide recognition mode enabled. For TiO2 enrichment, samples
were reconstituted in pre-cooled IAP Buffer (1.4ml) and followed by adding TiO2 beads,
shaken for 40 minutes and centrifuged to remove the supernatants. Beads were put into the
tips and washed with the washing buffer 3 to 6 times. Phosphopeptides were eluted and
concentrated using an elution buffer, followed by dissolving in formic acid (20 μl 0.1%) for
MS analysis. For LC-MS/MS analysis, a reverse phase trap column connected to a C18-
reversed phase analytical column was used to load the peptides. A positive ion mode was
used for the mass spectrometer, and MS data were acquired using a data-dependent top10
method. Peptides and proteins were mapped by Proteome Discoverer 2.4 (Thermo
Electron). The parameters set as follow: Enzyme: Trypsin; Max Missed Cleavages: 2;
Variable modifications: Oxidation (M), Phospho (ST) / Phospho (Y); Peptide Mass Tolerance:
± 20 ppm; Fragment Mass Tolerance: 0.1Da; Database pattern: Decoy; Peptide FDR: ≤0.01;
Protein FDR: ≤0.01; Experimental Bias: Normalizes all peptide ratios by the median protein
ratio. The median protein ratio should be 1 after the normalization. Downregulated genes
meeting the criterion of $p < 0.05$ were selected for GO analysis with all genes we identified in
phosphoproteomics as background genes. Top 10 GO terms were mapped, and sequences
were annotated using gProfiler.
774
Western blotting
Samples were collected and homogenized in RIPA lysis buffer supplemented with protease
and phosphatase inhibitors, and were incubated for 30 min at 4°C. The resultant lysates were
cleared by centrifugation at 20,000g for 20 min at 4°C. Proteins were denatured by 1x sample
buffer supplemented with 0.1M DTT and incubated at 95°C for 5-7 min. Samples were then
resolved in 10% Tri-glycine SDS-PAGE gel. Proteins were then transferred onto a PVDF
membrane at 100V for 2 hr on ice. The membranes were blocked by 5% non-fat milk in TBS
and probed with primary antibody diluted in 2% non-fat milk in TBST at 4°C overnight:
nELAVL (1:4000, Proteintech), pS-nELAVL (1:1000, Biomatik, rabbit polyclonal
phosphospecific antibodies were raised against SYARPS(pS)ASIRDAN) , EB2 (1:2000,
Abcam), pS222-EB2 (1:2000, Covalab), MAP1S (1:1000, Sigma Aldrich), pS812-MAP1S
(1:1000, Covalab), GAPDH (1:7500, Life Technologies), CDKL5 (1:500, Santa Cruz).
Following incubation with 1:5000 HRP-conjugated secondary antibody (2% in non-fat milk
in TBST) for 1 hr at room temperature, membranes were washed with 1x TBST for five times,
and 10 min for each wash. Signal was detected using enhanced chemiluminescent substrate.
The expression of targeted proteins was quantified by ImageJ software.
791
Immunohistochemistry
Mice were perfused with ice-cold phosphate-buffered saline (PBS) and 4% paraformaldehyde
transcardially. Brains were collected and incubated in 4% paraformaldehyde at 4°C overnight.

Brain was sectioned by vibratome and then subjected to immunostaining. Brain sections were
blocked for 1 hr in PBS with 2% horse serum and 0.4% Triton-100 followed by an overnight
incubation at 4°C with rabbit anti-c-fos primary antibody (1: 2500, Synaptic Systems). Brain
sections were washed and incubated with a secondary antibody (1: 1000, Invitrogen) for 1 hr
at RT. After PBS rinsing, slices were mounted on glass slides. Confocal microscopy (TCS
SP8, Leica) was used for imaging.
801
RNA extraction and real-time PCR
Total RNA from mouse V1 was extracted using NucleoSpin RNA kit (Macherey–Nagel),
following the manufacturer's instructions. Reverse transcription of RNA into cDNA was
conducted using a PrimeScript RT Reagent Kit (RR037A, Takara Bio). Quantitative real-time
PCR was conducted with a Premix Ex Taq Kit (RR390A, Takara Bio) using ABI ViiA7 Real
Time PCR System (Applied Biosystems). The mRNA expression was normalized to that of
Gapdh.
809

mouse-Gapdh-Forward primer	TCATGACCACAGTCCATGCC
mouse-Gapdh-Reverse primer	CAGATCCACGACGGACACAT
mouse-Arc-Forward primer	GGAGGGAGGTCTTCTACCGT
mouse-Arc-Reverse primer	CTACAGAGACAGTGTGGCGG
mouse-Fos-Forward primer	CGAAGGGAACGGAATAAG
mouse-Fos-Reverse primer	CTCTGGGAAGCCAAGGTC
mouse-Homer1a-Forward primer	GAAGTCGCAGGAGAAGATG
mouse-Homer1a-Reverse primer	TGATTGCTGAATTGAATGTGTACC
mouse-Homer1b-Forward primer	ACACCCGATGTGACACAGAA
mouse-Homer1b-Reverse primer	CACTGCTTCACATTGGCAGT
mouse-Homer1c-Forward primer	GAGAAGTCGCAGGAGAAGATG
mouse-Homer1c-Reverse primer	TTGCTGAACTAGCATGAGAGAG
mouse-Npas4-Forward primer	CACTCGCAAGGGTGTCTTCT
mouse-Npas4-Reverse primer	AATCCAGGTAGTGCTGCCAC
mouse-Pcdh19-Forward primer	CAGCTTTCGCATTACGGCTC
mouse-Pcdh19-Reverse primer	TAGGTGGACTCGCCAAACAC
mouse-Gad1-Forward primer	TGCTTCCAGCTAAGAACGGG
mouse-Gad1-Reverse primer	CAGAACCTTGGTGGAGCGAT
mouse-Sst-Forward primer	CCCCAGACTCCGTCAGTTTC

mouse-Sst-Reverse primer	GGCTCCAGGGCATCATTCTC
mouse-Pvalb-Forward primer	GCAGACTCCTTCGACCACAA
mouse-Pvalb-Reverse primer	CCTTCTTCACCTCATCCGGG

Bulk RNA sequencing and data analysis

811
RNA was extracted from V1 tissues of 6 pairs WT and *Cdkl5* KO mice (8-week-old) using
NucleoSpin RNA column (Macherey–Nagel), and RNA integrity was assessed by Agilent
4200 TapeStation System. mRNA enrichment fragmentation and cDNA library construction
were performed by Novogene Co. The transcriptomic profile for each sample was generated
by Novogene Co utilizing the Illumina Novaseq 6000 platform, 150bp paired-end sequencing.
For RNA sequencing, quality control of the datasets was firstly performed using FastQC[90].
Reads alignment was performed by HISAT2[91] and counting was conducted by
FeatureCounts[92]. Differential gene expression analysis was performed using R package
DESeq2[93]. DEGs between WT and *Cdkl5* KO mice were selected using the following criteria:
foldchange $>$ 1.2 or $<$ −0.83 and *padj* $<$ 0.05. STRING was used to construct PPI
network for DEGs. The PPI network with minimally required interaction score of high
confidence (0.7) were visualized in Cytoscape (3.10.0). Maximal Clique Centrality (MCC)
algorithm in CytoHubba was used to calculate and identify hub genes. DEGs with
log2Foldchange were subjected to GSEA, which was performed by clusterProfiler[94] that
enabled GO term enrichment analysis.

Isolation of V1 nuclei for snRNA-seq

828
Nuclei from 6 pairs of frozen WT and *Cdkl5* KO mouse (8-week-old) V1 samples were
isolated following previously established protocol[95]. WT and *Cdkl5* KO mice were
euthanized as described above. Mouse V1 was dissected in ice-cold DPBS and immediately
snap-frozen in liquid nitrogen. Frozen V1 tissues from two animals were pooled and
transferred to a prechilled Dounce homogenizer containing ice-cold homogenization buffer.
This buffer consisted of 250 mM sucrose, 25 mM KCl, 5 mM MgCl2, 20 mM tricine-KOH
(pH 7.8), 1 mM dithiothreitol, 0.15 mM spermine, 0.5 mM spermidine, 5 μg/mL actinomycin,
0.16% Nonidet P-40, 0.04% bovine serum albumin, protease inhibitors, and 200 U/mL
RNAsin. The tissues were gently homogenized on ice using a loose pestle until no visible
solid fragments remained. The lysate was subsequently combined with OptiPrep (Sigma) at a
ratio of 1:1 (v/v) and centrifuged at 10,000 $\times$ *g* for 20 min at 4 °C. The supernatant was
removed and the nuclei were washed once with Dulbecco's modified Eagle medium/F12
supplemented with 5% fetal bovine serum. The nuclei were resuspended in Dulbecco's
modified Eagle medium/F12 supplemented with 10% fetal bovine serum and 200 U/mL
RNAsin. The resuspended nuclei were counted on a hemocytometer using Trypan blue
labeling and diluted to 400 nuclei per microliter. Nuclei were recounted on an automated cell
counter (RWD) before further processing.

snRNA-seq library preparation

847
snRNA-seq libraries of WT and *Cdkl5* KO mouse V1 samples were constructed by
Chromium Single Cell 3′ Library Kit v3 (1000078; 10× Genomics). 40 μL suspension of
diluted nuclei for each sample was loaded and processed according to manufacturer's
instructions. The libraries were subjected to 150bp paired-end sequencing on an Illumina
NovaSeq 6000 system (Novogene Co), obtaining a minimum of 200 GB raw data for each
library.

 snRNA-seq analysis
The demultiplexed FASTQ files were mapped to the GRCm38 genome reference using Cell
Ranger (version 7.1.0) with the default settings. The transcriptomic analysis was performed
using a Python package called SCANPY[96]. Raw gene expression matrices from 6 snRNA-seq
libraries (each library was derived from two animals) were combined for analysis. For quality
control, the following criteria was used to exclude low-quality cells or outliers: n_genes >
6500 or < 700, percent_mito > 5%, and n counts > 40,000. Genes detected in less than 8 cells
were removed[37]. Cells were normalized using *scanpy.pp.normalize_per_cell* with default
setting, followed by log-transformation.
864
After quality control, highly variable genes (HVGs) were identified using
scanpy.pp.highly_variable_genes with default setting. Harmony[36] was employed for batch
correction. The top 40 principal components (PCs) were used to compute a nearest-neighbor
graph on the cells using *scanpy.pp.pca* function. Leiden algorithm[97] was applied to do
clustering with default setting and the graph was finally embedded in 2D via the Uniform
Manifold Approximation and Projection (UMAP) algorithm[98]. Canonical marker genes
which were reported in previous studies[37] were used for annotation. Doublets were identified
using Scrublet[35]. Clusters that expressed markers of more than two classes or contained more
than 50% doublets were discarded.
874
To further identify types in each class, this combined gene expression matrix was divided
into 3 matrices: glutamatergic neurons, GABAergic neurons and non-neuronal cells. The raw
matrix of glutamatergic neurons was further processed for highly variable gene selection,
batch correction, dimensionality reduction and clustering as described above and visualized
by UMAP. Previously reported type-specific markers[37] were used for type annotation of
glutamatergic neurons.
881
882 Differential expression analysis was performed to assess the transcriptomic alterations
883 between WT and *Cdkl5* KO mouse V1, and to identify markers in each subclass or type.
884 Differential expression analysis in this study was achieved using
885 *scanpy.tl.rank_genes_groups* function in the SCANPY package. Differentially expressed
886 genes were defined as: adjusted P value < 0.05 and a log2 fold change > 0.25 or < −0.25.
887
888 Enrichment for activity-dependent genes in visual cortex[99] was calculated using the R
889 package GeneOverlap. The lists of DEGs for each cell type were tested, utilizing all genes
890 existing in snRNA seq from this study as the background. *P* values from Fisher exact
891 test were reported.
892
Fluorescent in situ hybridization (FISH)
WT and *Cdkl5* KO mice (8-week-old) were euthanized and perfused as described. The brains
were removed and fixed in 4% PFA for 24hr at 4 °C, followed by sucrose gradient until
dehydration. Mouse V1 tissues were dissected under microscope and embedded in optimal
cutting temperature compound (OCT, Tissue-Tek), and frozen at -80°C. The V1 tissues were
sectioned at 20 μm with a cryostat (Epredia Cryostar NX70 Cryostat). The sections were
fixed with 4% PFA for 10 min, washed 3 times with PBS and dehydrated using 70% ethanol.
The sections were permeabilized by 8% SDS in PBS for 15 min at room temperature. The
sections were then labelled according to the manufacturer's protocol[100] (Molecular
Instruments). DAPI was stained before imaging. Probes for *Fos* and *Arc* were commercially
designed by the manufacturer (Molecular Instruments). Probe for *Homer1* and *Camk2a* were
custom designed and the target sequences are listed below:

Homer1-1
GGCTATCAGCCCATTGGCCAAACTTTTGAGATGTTTTAGTAAATGTCATG
Homer1-2
GCCGAGCAGCTTCCTTAAATTCCTGAAACTTTTCTGCGAATTTTGAAAGA
Homer1-3 GCTGAAGATAGGTTGCTCCCC
Homer1-4 GGTGTGATGGTGCTATTTATTATTGCCTTTGAGCCATCTAAACTG
Homer1-5 AAGGGGTACTGGTCAGCTCC
Camk2a-1 CCACAGCGTGAGAAAGAGCAGCAT
Camk2a-2 TTCCCCAGGGCCTCTGGTTCAA
Camk2a-3 ATCTGCCATTTGCCGTCCCTGC
915
916 FISH assay on rat primary hippocampal neurons and iNeurons were performed with
917 ViewRNA Fos probe following manufacturer's protocol (Rat: Catalog number: VX-06,
918 Assay ID: VC1-11292-VCP; Human: Catalog number: VX-01, Assay ID: VA1-11504-VC;
919 Thermo Fisher Scientific).
920
921 Leica SP8 confocal microscope with a 40×, NA 0.85□objective (SBS core lab, CUHK) was
922 used for imaging. V1 cortical areas were imaged with optical sectioning of 0.5 μm and 15
923 optical sections. Channels were imaged sequentially to avoid any optical crosstalk. The
924 number and volume of puncta within cells were quantified using the Imaris reconstruction
925 software (v.9.0, Bitplane).
926
Epitope-preserving Magnified Analysis of Proteome (eMAP), imaging and analysis
We performed eMAP as previously described[101,102]. Mice under anesthesia were perfused
with PBS followed by 4% paraformaldehyde. Brain tissues were collected and post-fixed in 4%
paraformaldehyde overnight at 4□. Brain tissues were then embedded in MAP solution (30%
acrylamide, 10% sodium acrylate, 0.1% bisacrylamide, and 0.03% VA-044 in PBS) under
hypoxic conditions. The embedded samples were then immersed in hydration solution (0.02%
sodium azide in PBS) and sectioned at 60 μm using a Leica VT1000S vibratome. Lipid was
removed by incubation in clearing solution (6% SDS, 0.1 M phosphate buffer, 50 mM
sodium sulfite, 0.02% sodium azide in deionized water [pH 7.4]) for 4 hours at 37□. Samples
were then washed with PBST (0.1% Triton X-100 and 0.02% sodium azide in PBS) and
stained with primary anti-GFP (A10262, Invitrogen) antibody diluted in PBST overnight at
37□. After washing with PBST, samples were incubated with secondary antibodies diluted in
PBST overnight at 37□ and then washed again with PBST. The samples were expanded and
mounted in 0.01× PBS immediately before imaging. Approximately 3× total linear expansion
was achieved, and imaging was performed under a Leica SP8 confocal microscope using a
20X lens with a 0.75 numerical aperture or 63X water immersion lens with a 1.2 numerical
aperture. For dendritic spine morphology analysis, FIJI (version 2.14) was used to measure
head diameter (D_{head}), neck diameter (D_{neck}), and the length (L_{length}) of dendritic spines
according to dendritic spine classification criteria previously described[103].
946
Site-directed mutagenesis
Wildtype sequences of ELAVL 2 to 4 were directly cut out by corresponding restriction
enzymes and ligated to in-frame EGFP-C vectors. Then, by using them as templates, SE
mutants of Elavls 2 to 4 were built using traditional Primer Extension Mutagenesis with two-
stage nested PCR[104]. While SA mutants were built using one-stage PCR and Dpn1 digestion
method[105].
Addgene plasmids carrying wild-type sequences of ELAVL 2 to 4:
ELAVL 2: pENTRY4_HuB (65752);

ELAVL 3: pENTRY4_HuC (65753);
ELAVL 4: pENTRY4_HuD (65754).
Primers for SE mutants:
ELAVL2:
(first-stage PCR1)   gcccgaattcggatggaaacacaactgtctaat
atctctgatagaagcttcacttgggcgagcata
(first-stage PCR2)   tatgctcgcccaagtgaagcttctatcagagat
ggtacctcgagaggctttgtgcgttttgtttgt
(second-stage PCR)   gcccgaattcggatggaaacacaactgtctaat
ggtacctcgagaggctttgtgcgttttgtttgt
ELAVL 3:
(first-stage PCR1)   gcccgaattcggatggtcactcagatactgggg
ccggatggatgcttcactgggtctggc
(first-stage PCR2)   gccagacccagtgaagcatccatccgg
ggtacctcgagacgccttgtgctgtttgctggt
(second-stage PCR)   gcccgaattcggatggtcactcagatactgggg
ggtacctcgagacgccttgtgctgtttgctggt
ELAVL 4:
(first-stage PCR1)   accgagatctctatggttatgataattagcacc
ccctgattgaggcctcgctcggacgggc
(first-stage PCR2)   gcccgtccgagcgaggcctcaatcaggg
taccgtcgactggacttgtgggctttgttggt
(second-stage PCR)   gcccgtccgagcgaggcctcaatcaggg
taccgtcgactggacttgtgggctttgttggt
979
Primers for SA mutants:
ELAVL 2:   cctatgctcgcccaagtgcagcttctatcagagat
atctctgatagaagctgcacttgggcgagcatagg
ELAVL 3:   atgccagacccagtgcagcatccatccgg
ccggatggatgctgcactgggtctggcat
ELAVL 4:   tgcccgtccgagcgctgcctcaatcag
Ctgattgaggcagcgctcggacgggca
987

LLPhyScore algorithm
The algorithm computed a composite LLPS_score for each entry by integrating eight distinct
biophysical features derived from PDB statistical potentials, including solvent accessibility
and pi-pi interaction frequencies. We subsequently constructed a global rank-ordered
distribution curve by sorting all sequences by their predicted scores. This framework allowed
for a quantitative, comparative assessment of the ELAVL family's biophysical driving forces
relative to both established high-confidence scaffolds and the general baseline of the
proteome.
996

Cell line culture, transfection and drug treatment
The primary hippocampal neurons were prepared from embryonic day 18.5 rat embryos.
Hippocampal neurons (1×10^5) were plated on 10 µg/ml poly-D-lysine (Sigma Aldrich)
coated 18 mm coverslips in 12-well plates, fed with Neurobasal medium (21103-049, Gibco)
supplemented with 2% B27 (17504-044, Gibco) and 1% GlutaMAX (35050-061, Gibco). We
performed calcium phosphate transfection on div 11 and fixed the neuron for immunostaining
following previously described protocol on div 17[106]. The HEK 293T and HT22 cells were
grown in Dulbecco's Modified Eagles Medium (DMEM) supplemented with 10% of fetal

bovine serum and 1% of penicillin/streptomycin solution up to 40 passages. We used the
general PEI transfection protocol by adding 0.4 μg DNA/coverslip, transfect overnight and
perform immunostaining. For Actiomycin D treatment, we replenish the iNeurons with
neurobasal medium containing 2 μg/mL Actinomycin D at different time point, and fixed
them together for FISH assay.
1010
1011

iPSC culture
Human iPSC lines OR00005 (p.R550* mutant allele clone) and OR00006 (WT allele clone)
were purchased from Coriell Institute. The two lines were reprogrammed from a
heterozygous CDD female patient carrying a pathogenic variant 1648C>T (p.R550*) of
CDKL5 in one allele as well as a WT allele. To generate adenine base editor (ABE)-
corrected iPSCs, we electroporated 5 μg ABEmax plasmid with a guide RNA to correct the
mutation using Human Stem Cell Nucleofector 1 (Lonza, VPH-5012) according to the
manufacturer's protocols. ABE-corrected iPSC clones were selected with puromycin and
confirmed through sanger sequencing. iPSCs were cultured in Essential 8 (E8) Medium
(Thermo Fisher Scientific, A1517001) on 6-well cell culture dishes (SPL, 30006) coated with
Geltrex (Thermo Fisher Scientific, A1414402) diluted 1:100 in DMEM/F-12 (Thermo Fisher
Scientific, 11330032). E8 medium was daily replenished. When iPSC colonies demonstrated
mature morphology, the iPSCs were either clump passaged with EDTA (Thermo Fisher
Scientific, 25300054) for routine maintenance or differentiation into cortical neurons. For
clump passaging with EDTA, iPSCs were washed with Dulbecco's PBS (DPBS; Millipore
Sigma, D8537) and then incubated with EDTA for 5–7 min at 37 ; the EDTA solution was
then aspirated and replaced with E8; iPSC colonies were then gently detached and
resuspended gently and centrifuged at 300 g for 3 min. Then the supernatant was discarded,
and cell pellets were resuspended with E8 + Y-27632 (Sigma Aldrich, Y0503) and passaged
at 1:10–1:40 dilutions in E8 + Y-27632, with Y-27632 removed the next day.
1032

Two-dimensional neuron differentiation
For differentiation of iPSCs into cortical neurons, published protocols were followed with
minor modifications[107]. iPSCs were plated on 6-well Geltrex-coated plates and suspended
lentivirus with rtTA and Ngn2-GFP-puro vectors 2 hours later in E8 medium containing 10
μM Y-27632 and infected the iPSCs for 24 hours. After the colonies demonstrated mature
morphology (Day 0), the culture medium was replaced with N2 medium (DMEM/F-12, 1 ×
N2 and 1 × NEAA; Thermo Fisher Scientific) containing 2 μg/mL doxycycline (Sigma
Aldrich). On day 1, the culture medium was replaced with N2 medium containing 2 μg/mL
doxycycline and 1 μg/mL puromycin (Thermo Fisher Scientific, A1113803). On day 3, the
induced cells were dissociated with Accutase (Thermo Fisher Scientific, A1110501) and
plated them onto 6-well PDL/laminin coated plates at 1.0×10^6 cells per well or 24-well
plates with coverslips at 5×10^4 cells per coverslip in B27 medium (Neurobasal medium, 1 ×
B27, 1 × GlutaMAX and 0.2 μg/ml laminin, Thermo Fisher Scientific) containing 2 μg/mL
doxycycline, 1 μg/mL puromycin. For further experiments, from day 7 until day 28, half of
the medium was removed every 4 days and was replaced with B27 medium containing 0.5
μg/mL doxycycline, 10 μg/mL BDNF and 10 μg/mL GDNF.
1049

1050 *Immunocytochemistry*
1051 Cells grown on coverslips or chamber slides were fixed with 4% paraformaldehyde (Sigma,
1052 F8775) for 15 min at room temperature. Then the cells were permeabilized with 0.1% Triton
1053 X-100 in DPBS for 5 min at RT and blocked with 1% BSA in DPBS for 20 min. Primary
1054 antibodies were incubated with the cells overnight at 4℃. The following primary antibodies

were used: anti-Elavl2 (Proteintech, 14008-1), anti-Elavl3 (Proteintech, 55047-1), anti-Elavl4 (Santa Cruz, sc-48421 and sc-28299), anti-GFP (Invitrogen, A10262), anti-TIA1 (Santa Cruz, sc-166247), anti-GW182 (Santa Cruz, sc-56314). After washing three times in DPBS, the cells were incubated with secondary antibodies and DAPI for 1 hr at room temperature. The following secondary antibodies were used: Alexa Fluor 488 (Invitrogen, A32931), Alexa Fluor 555 (Invitrogen, A32732), Alexa Fluor 568 (Invitrogen, A21124), Alexa Fluor 594 (Invitrogen, A32742), Alexa Fluor 647 (Invitrogen, A32728, A32733), STAR Orange (Abberior, STORANGE-1001, STORANGE-1002) and STAR Red (Abberior, STRED-1001, STRED-1002). After washing for 3 times, the samples were mounted with anti-fade mounting medium.

Confocal images were acquired using Leica SP8 microscope with 63x objective, Z-series images were collected using z-serial scanning mode (20 optical sections at 0.3 µm steps). STED images were acquired using abberior STED microscopy system at 50 nm resolution, z-series images were collected using z-serial scanning mode (100 optical sections at 75nm per step).

Quantification of puncta number, volume and fraction distribution
The number and volume of puncta within cells were quantified using the Imaris reconstruction software (v.9.0, Bitplane). 3D reconstruction was performed on high-resolution confocal/STED images. The border of cells and nucleus was determined by surface module in the software, rendering with 0.1-µm smoothing, and disconnected processes were merged with the cell body to create a single surface. Then the spots module was used to measure the number and volume of the fluorescence puncta. The puncta within cells were limited by distance between puncta and cell surface/Nucleus surface. The fraction distribution analysis was done by fraction fining puncta within each cell by their volume, following by non-linear regression.

Protein expression and purification
Human *ELAVL3* (GenBank: NM_001420.4) gene was cloned into a modified pET32a vector with an N-terminal Trx-His$_6$-tag. Full-length ELAVL3 was expressed in *Escherichia coli* BL21 (DE3) host cells at 37°C for 4 hr induced by 0.2 mM isopropyl-β-D-thiogalactoside (IPTG). Cells were harvested and resuspended in buffer containing 50 mM Tris, pH 8.0, 1 M NaCl, 5% Glycerol, 2 mM β-ME and 10 mM imidazole, and then lysed by high pressure homogenizer at 4°C. After centrifugation at 18,000 rpm for 20 min, the supernatants were loaded onto the Ni^{2+}-NTA agarose affinity column and eluted with the buffer containing 50 mM Tris, pH 8.0, 1 M NaCl and 500 mM imidazole, and then purified by a size-exclusion chromatography in the buffer containing 50 mM Tris, pH 8.0, 1 M NaCl, 5% Glycerol, 1mM EDTA and 1 mM DTT.

Protein labeling with fluorophore
Purified ELAVL3 proteins (~5 mg/mL) were dissolved in the buffer containing 300 mM NaCl, 100 mM NaHCO$_3$, pH 8.3, 4 mM β-ME. Cy3 NHS ester (AAT Bioquest) was dissolved in DMSO and incubated with the corresponding protein (molar ratio ~1:1) at room temperate for 1 hr. The labeling reaction was quenched by the buffer of 200 mM Tris, pH 8.2, and the labeled protein was purified with a HiTrap desalting column with the buffer containing 50 mM Tris, pH 8.0, 1 M NaCl , 5% Glycerol, 1mM EDTA and 1 mM DTT. Fluorescence labeling efficiency was determined by Nanodrop 2000 (Thermo Fisher).

In vitro phase separation assay
The stock ELAVL3 protein was centrifuged at 16,873 g for 10 min at 4°C to remove any precipitations and diluted to designed concentrations and buffer conditions. Subsequently, all the samples were placed on ice before phase separation assay. For microscope-based assays, each sample was injected into a home-made chamber for fluorescence imaging (Leica TCS SP8).

Fluorescence recovery after photo-bleaching assay
For FRAP of ELAVL3 SA and SE overexpressed HEK 293T cells, a circular ROI was bleached by 488 nm laser beam at 37□. The photobleaching was acquired with 50 cycles of 488 nm laser at 10% power and imaging every 0.5 s for 5 min after bleaching. The fluorescence intensity of the bleached area was normalized to the intensity of the same area before bleaching. The recovery curves were corrected with the region without photobleaching in the same frames.

For Cy3-labeled protein droplets, ROI was bleached by a 561 nm laser beam at room temperature. A dwell time of 1.2 s and 100% of laser power was used during bleaching. Fluorescence recovery was monitored for 300 s at 5 s intervals. For each experiment, the fluorescence intensity of a neighboring droplet with similar size to the bleached one was also recorded for intensity correction. Background intensity was subtracted before data analysis. The pre-bleaching ROI intensity was normalized to 100%.

Co-Immunoprecipitation
Protein extraction was performed in HEK 293T cells using a NP40 buffer (50 mM Tris-HCl buffer (pH 7.4), 150 mM NaCl, 2 mM MgCl, 1 mM EDTA, 0.5% (v/v) NP40, 1 mM DTT and complete EDTA-free protease inhibitor cocktail). Pre-clearing of protein extract was performed by 30 minutes R/T incubation with Protein A magnetic beads (Life Technologies).

Pre-cleared extract was incubated with 2 µg of GW182 antibody (sc-56314; Santa Cruz Biotechnology) for O/N at 4°C, then performed immunoprecipitation with Protein A magnetic beads by 30 minutes R/T incubation. Samples were washed 5 times using NP40 buffer and protein complex were released by resuspending the beads in 1x sample buffer supplemented with 0.1M DTT and incubated at 95°C for 5-7 min.

Electrophoresis Mobility Shift Assay (EMSA)
nELAVL SE and SA mutants were cloned and overexpressed in E.coli. Proteins were purified by Gene Universal Inc., (Chuzhou, Anhui). Fos-Cy5 au-rich RNA oligos (sequence: AUAUUUAUAUUUUUAUUUUAUUUUUUU) was synthesized by Techdragen. The binding buffer for EMSA consisted of 60 mM KCl, 300 mM NaCl, 2 mM MgCl2, 20 mM HEPES (pH 8.0) 1 mM EDTA, 1 mM DTT, 10 % Glycerol, 0.5 mg/mL of tRNA, and 2 fmol of Fos-Cy5 RNA per reaction. Reactions were equilibrated for 1 h at room temperature before loading on a Tris-glycine gel (to minimize complex dissociation), running 45 mins at 100 V. All gel shifts were done at least three times. Bands were quantified using ImageJ. Fraction bounds were calculated by plotting the ratio of RNA-protein complex against total RNA. Lines were obtained by nonlinear regression.

Puromycin Proximity Ligation Assay (Puro-PLA)
The Puro-PLA were performed on rat primary hippocampal neurons expressed with ELAVL3-SE/SA probing Fos protein and puromycin with kit (Duolink®□ In Situ Red

Starter Kit Mouse/Rabbit, catalog number: DUO92101, Sigma Aldrich) following manufacturer's instructions.

Craniotomy and AAV injection
6-week-old mice were anesthetized with isoflurane mixed with air (3% for induction, 2% for surgery) and placed on a stereotaxic apparatus (RWD Instruments). A 38 °C heating pad was used for maintaining animal body temperature. Ophthalmic ointment (Bausch & Lomb corner gel) was applied to the eyes for eye lubrication. Ethanol and betadine swabs were used to sterilize the skin above the skull before making an incision. A ~4 mm craniotomy window was made over the left V1 (center coordinates were 3.2 mm ML and 1 mm AP relative to lambda) by a dental drill as described previously[108]. A glass pipette (RWD Instruments) was pulled to a fine tip by a puller (PC-100, Narishige), and filled with mineral oil in the whole pipette and AAV in the front tip. The filled glass pipette was lowered into the V1. At a depth of 0.1-0.2 mm below the dura, virus was injected slowly (120nl in 5 min) into V1 using a microinjector (R480 Nanoliter injection pump, RWD Instruments). After the injection, pipette was kept for 5 minutes before being withdrawn to prevent the backflow of virus. A two-layer glass window was made by sticking a 3 mm and 5 mm-diameter cover glass (#1 thickness, Warner Instruments) using UV sensitive optical adhesive. The glued glass window was placed under slight downward pressure on the exposed brain. The 5 mm was attached to the skull and the 3 mm was fitted into the craniotomy. A tissue-friendly adhesive (3M Vetbond) was applied to seal around the surrounding skull. A custom-made stainless steel head plate was attached to the skull with dental adhesive resin cement. Dental cement was used to seal the rest of the exposed skull. Two-photon imaging was conducted 2 weeks after craniotomy window implantation.

Viral vectors
For *in vivo* calcium imaging, rAAV-hSyn-GCaMp6f-WPRE-hGH-polyA (titer: ~5.7x10^{12}vg/ml, BrainVTA Co.,Ltd) was used. For sparse labeling of neurons for eMAP, a mixture of diluted pENN.AAV.CamKII 0.4.Cre.SV40 (titer: 1.8x10^{13}vg/ml, Addgene: 105558, 1:10000-20000 dilution in PBS) and cre-dependent EGFP virus (titer: 1x10^{13} vg/ml, Addgene: 51502) were used. nELAVL-WT AAV was generated with a mix (1:1:1) of ELAVL2-WT, ELAVL3-WT and ELAVL4-WT plasmid (titer: 2.45×10^{13}vg/ml, Vigene Biosciences, lnc.). nELAVLs-SE AAV was generated with a mix (1:1:1) of ELAVL2-S119E, ELAVL3-S119E and ELAVL4-S131E plasmid (titer: 6.60×10^{13}vg/ml, Vigene Biosciences, lnc.). nELAVLs-SA virus was generated with a mix (1:1:1) of ELAVL2-E119A, ELAVL3-E119A and ELAVL4-E131A plasmid (titer: 8.02×10^{13}vg/ml, Vigene Biosciences, lnc.).

Two-photon in vivo imaging with visual stimuli
The Psychophysics Toolbox in MATLAB was used to generate drifting grating stimuli and natural scene[109,110]. For each stimuli trial, there was a 3s blank period (gray at mean luminance) followed by a 1s drifting sinusoidal grating (mean luminance: ~50 lx) with three different spatial frequencies (0.08, 0.04 and 0.02 cycles per degree) at temporal frequency of 2 Hz followed a 4s blank period with 4s natural scene. Eight drifting directions were separated by 45 degrees, and ten trials were displayed. An LCD monitor (21.5-inch, 75 Hz refresh rate) was placed 18 cm centered in front of the mouse's eyes to display the stimuli. The monitor covered a field of view of ~106 degrees horizontal and ~58 degrees vertical. Mice were head-fixed, and the left V1 was imaged by a two-photon microscope (Scientifica, U.K.) with a resonant scanning module controlled by ScanImage built-in Matlab software. The Coherent Discovery laser was used at 920 nm for excitation through a Nikon water immersion objective (16X, N.A. = 0.8, working distance: 3.0 mm). The power was less than

100 mW for all imaging. The signal was detected by a photomultiplier with bandwidth of
480-560nm. Images were acquired at ~7.5 Hz (30 Hz averaged to 4) over 512x512 pixels of
each view with bidirectional scanning. BNC DAQ (BNC-2110, National Instruments) was
used to synchronize the two-photon microscope recording and visual stimuli display. Mice
were habituated with the imaging set-up for two to three days (30 min daily) before data
collection. Neurons of L2/3 at 100-250μm below dura were imaged. To record visually
evoked responses from different eyes, an opaque black card blocker was placed in front of
the eye according to imaging conditions. 3 to 6 fields of view (FOVs) were collected for each
mouse.
1212
Image processing and data analysis for two-photon microscopy
To extract the calcium signal of each neuron at different imaging conditions (i.e., spatial
frequencies 0.02, 0.04 and 0.08cpd, and ways of eye blocking), all images of the same FOV
were imported and analyzed by suite2p (version 0.10.1) at the same time. The parameters of
suite2p were set as follow: tau of 0.7, denoise of 1, diameter of 9, anatomical only of 1,
maximum iterations of 1, frames per second of 7.5 and 191 minimum neuropil pixels. To
confirm the stereotaxic coordinates reliably targeted bV1, we ensured at least 10% of putative
neurons (regions of interest (ROIs)) extracted from each imaging session elicited time-locked,
visually evoked responses from the ipsi eye, similar to other published reports[60,65].
1222
The debris-like ROIs were removed manually ("*iscell*" from 1 to 0). The intensities of the
remaining neurons were saved in .mat files for data analysis in MATLAB (version R2021a).
The raw intensities of neurons were adjusted with neuropil signal according to $F_{adj} = F_{raw} -$
$0.7 \times F_{neu}$[111]. Normalization of adjusted intensity was done by transforming into modified Z-
score $Z = \frac{F_{Adj} - mean\ of\ F_{OFF}}{Standard\ deviation\ of\ F_{OFF}}$[65], where F_{OFF} is the 1-second adjusted signal in blank
period just before each visual stimulus of that trial. The neuron with more than 20% of trials
(i.e., > 2 trials in drifting grating and > 6 trials in natural scene) with $Z > 2$ at any stimulus,
as well as the trial-average calculated for all time-point, $Z_{trial\_avg} > 2$ in that stimulus, was
counted as stimulus-responsive. Tuning curve was obtained by averaging the Z_{trial_avg} in
each grating direction. To analyze the orientation selectivity of grating-responsive neurons,
the response $R(\theta_k)$ of orientation k is calculated by averaging the Z_{trial_avg} of opposite
directions. The orientation selectivity index (OSI) was defined as the norm of $\vec{L} =$
$\frac{\sum_k R(\theta_k)\exp(2i\theta_k)}{\sum_k R(\theta_k)}$ and the preferred orientation was half of the corresponding angle of $\vec{L}$, where
θ_k was expressed in radian[112]. The similarity of their signal was calculated as the distance
correlation between their tuning curves for neuron responding to drifting grating stimuli in
any two viewing conditions[113]. The confidence interval of mean difference of OSI
distribution between WT and *Cdkl5* KO mice was estimated with bootstrapped distribution
under significantly large sampling (i.e., 9999 times).
1241
Considering that a neuron can contribute to population encoding regardless of their individual
responses to the stimuli, all extracted putative neurons from suite2p were thus included in the
population analysis. To quantify stimulus information encoded in activities of the neural
populations, we measured the performance of a classifier in the prediction of different
grating[114]. A support vector machine classifier was trained by 5-fold cross-validation[115].
Since the number of trials was limited in our study, the trained classifier would tend to overfit
the training set, which could lead to an underestimation of the performance of the classifier
for the validation set. Therefore, during training, the early stop was applied. Then the
classifier's performance was evaluated by accuracy on the validation set. To evaluate the

binocular matching in the neural population, decoder was trained with contralateral data set
and tested on both contra- and ipsilateral signal.
1253
Visual cliff task
The visual cliff task assessed mice's preference for descending a platform in an open field.
The field consisted of an open field (40 x 40 x 40 cm) with transparent plexiglass floor
positioned on the edge of a laboratory bench: half of the box rested on the tabletop (the "safe"
side), while the other half extended over the floor 80 cm below (the "cliff" side)[57]. Both sides
featured patterns visible through the plexiglass floor, a black and white checkerboard cover
(3.5 x 3.5 cm squares) was placed on the bench top and extended vertically down to the floor,
creating the illusion of a cliff. Prior to each trial, mice were transported to the testing room
and habituated in their home cages for at least 30 minutes. At the start of each trial, a mouse
was placed on a platform (3 cm in height, 5 cm in diameter) at the center of the box, facing
the middle line between the two sides. The side from which the mouse descended was
recorded by a camera. Each mouse underwent 5 trials, and the open field was cleaned
between trials. The percentage of descents from the cliff side was calculated for each mouse.
1267
1268

References

Montini, E. *et al.* Identification and characterization of a novel serine-threonine kinase gene from the Xp22 region. *Genomics* **51**, 427-433, doi:10.1006/geno.1998.5391 (1998).

Kalscheuer, V. M. *et al.* Disruption of the serine/threonine kinase 9 gene causes severe X-linked infantile spasms and mental retardation. *Am J Hum Genet* **72**, 1401-1411, doi:10.1086/375538 (2003).

Olson, H. E. *et al.* Cyclin-Dependent Kinase-Like 5 Deficiency Disorder: Clinical Review. *Pediatr Neurol* **97**, 18-25, doi:10.1016/j.pediatrneurol.2019.02.015 (2019).

Leonard, H. *et al.* CDKL5 deficiency disorder: clinical features, diagnosis, and management. *Lancet Neurol* **21**, 563-576, doi:10.1016/S1474-4422(22)00035-7 (2022).

Wang, I. T. *et al.* Loss of CDKL5 disrupts kinome profile and event-related potentials leading to autistic-like phenotypes in mice. *Proc Natl Acad Sci U S A* **109**, 21516-21521, doi:10.1073/pnas.1216988110 (2012).

Mazziotti, R. *et al.* Searching for biomarkers of CDKL5 disorder: early-onset visual impairment in CDKL5 mutant mice. *Hum Mol Genet* **26**, 2290-2298, doi:10.1093/hmg/ddx119 (2017).

Sampedro-Castaneda, M. *et al.* Epilepsy-linked kinase CDKL5 phosphorylates voltage-gated calcium channel Cav2.3, altering inactivation kinetics and neuronal excitability. *Nat Commun* **14**, 7830, doi:10.1038/s41467-023-43475-w (2023).

Zhou, A., Han, S. & Zhou, Z. J. Molecular and genetic insights into an infantile epileptic encephalopathy - CDKL5 disorder. *Front Biol (Beijing)* **12**, 1-6, doi:10.1007/s11515-016-1438-7 (2017).

Munoz, I. M. *et al.* Phosphoproteomic screening identifies physiological substrates of the CDKL5 kinase. *EMBO J* **37**, doi:10.15252/embj.201899559 (2018).

Baltussen, L. L. *et al.* Chemical genetic identification of CDKL5 substrates reveals its role in neuronal microtubule dynamics. *EMBO J* **37**, doi:10.15252/embj.201899763 (2018).

Khanam, T. *et al.* CDKL5 kinase controls transcription-coupled responses to DNA damage. *EMBO J* **40**, e108271, doi:10.15252/embj.2021108271 (2021).

Kim, J. Y. *et al.* A kinome-wide screen identifies a CDKL5-SOX9 regulatory axis in epithelial cell death and kidney injury. *Nat Commun* **11**, 1924, doi:10.1038/s41467-020-15638-6 (2020).

Lopes, A. T., Janiv, O., Claxton, S. & Ultanir, S. K. CDKL5's role in microtubule-based transport and cognitive function. *bioRxiv*, 2024.2008.2028.610038, doi:10.1101/2024.08.28.610038 (2024).

De Santis, R. *et al.* Mutant FUS and ELAVL4 (HuD) Aberrant Crosstalk in Amyotrophic Lateral Sclerosis. *Cell Rep* **27**, 3818-3831 e3815, doi:10.1016/j.celrep.2019.05.085 (2019).

Markmiller, S. *et al.* Context-Dependent and Disease-Specific Diversity in Protein Interactions within Stress Granules. *Cell* **172**, 590-604 e513, doi:10.1016/j.cell.2017.12.032 (2018).

Bowles, K. R. *et al.* ELAVL4, splicing, and glutamatergic dysfunction precede neuron loss in MAPT mutation cerebral organoids. *Cell* **184**, 4547-4563 e4517, doi:10.1016/j.cell.2021.07.003 (2021).

Bolognani, F., Contente-Cuomo, T. & Perrone-Bizzozero, N. I. Novel recognition motifs and biological functions of the RNA-binding protein HuD revealed by

genome-wide identification of its targets. *Nucleic Acids Res* **38**, 117-130,
doi:10.1093/nar/gkp863 (2010).

18   Colombrita, C., Silani, V. & Ratti, A. ELAV proteins along evolution: back to the
nucleus? *Mol Cell Neurosci* **56**, 447-455, doi:10.1016/j.mcn.2013.02.003 (2013).

19   Ip, J. P. K., Mellios, N. & Sur, M. Rett syndrome: insights into genetic, molecular and
circuit mechanisms. *Nat Rev Neurosci* **19**, 368-382, doi:10.1038/s41583-018-0006-3
(2018).

20   Banerjee, A. *et al.* Jointly reduced inhibition and excitation underlies circuit-wide
changes in cortical processing in Rett syndrome. *Proc Natl Acad Sci U S A* **113**,
E7287-E7296, doi:10.1073/pnas.1615330113 (2016).

21   Demarest, S. T. *et al.* CDKL5 deficiency disorder: Relationship between genotype,
epilepsy, cortical visual impairment, and development. *Epilepsia* **60**, 1733-1742,
doi:10.1111/epi.16285 (2019).

22   Goel, A. *et al.* Impaired perceptual learning in a mouse model of Fragile X syndrome
is mediated by parvalbumin neuron dysfunction and is reversible. *Nat Neurosci* **21**,
1404-1411, doi:10.1038/s41593-018-0231-0 (2018).

23   Townsend, L. B., Jones, K. A., Dorsett, C. R., Philpot, B. D. & Smith, S. L. Deficits
in higher visual area representations in a mouse model of Angelman syndrome. *J*
Neurodev Disord **12**, 28, doi:10.1186/s11689-020-09329-y (2020).

24   Galli, J. *et al.* Neurovisual profile in children affected by Angelman syndrome. *Brain*
Dev **45**, 117-125, doi:10.1016/j.braindev.2022.10.003 (2023).

25   Boggio, E. M. *et al.* Visual impairment in FOXG1-mutated individuals and mice.
Neuroscience **324**, 496-508, doi:10.1016/j.neuroscience.2016.03.027 (2016).

26   Olson, H. E. *et al.* Cerebral visual impairment in CDKL5 deficiency disorder: vision
as an outcome measure. *Dev Med Child Neurol* **63**, 1308-1315,
doi:10.1111/dmcn.14908 (2021).

27   Brock, D. *et al.* Cerebral Visual Impairment in CDKL5 Deficiency Disorder
Correlates With Developmental Achievement. *J Child Neurol* **36**, 974-980,
doi:10.1177/08830738211019284 (2021).

28   Lupori, L. *et al.* Site-specific abnormalities in the visual system of a mouse model of
CDKL5 deficiency disorder. *Hum Mol Genet* **28**, 2851-2861,
doi:10.1093/hmg/ddz102 (2019).

29   Pizzo, R. *et al.* Lack of Cdkl5 Disrupts the Organization of Excitatory and Inhibitory
Synapses and Parvalbumin Interneurons in the Primary Visual Cortex. *Front Cell*
Neurosci **10**, 261, doi:10.3389/fncel.2016.00261 (2016).

30   Quintiliani, M. *et al.* Cortical Visual Impairment in CDKL5 Deficiency Disorder.
Front Neurol **12**, 805745, doi:10.3389/fneur.2021.805745 (2021).

31   Eyers, P. A. A new consensus for evaluating CDKL5/STK9-dependent signalling
mechanisms. *EMBO J* **37**, doi:10.15252/embj.2018100848 (2018).

32   Mulligan, M. R. & Bicknell, L. S. The molecular genetics of nELAVL in brain
development and disease. *Eur J Hum Genet* **31**, 1209-1217, doi:10.1038/s41431-023-
01456-z (2023).

33   Scheckel, C. *et al.* Regulatory consequences of neuronal ELAV-like protein binding
to coding and non-coding RNAs in human brain. *Elife* **5**, doi:10.7554/eLife.10421
(2016).

34   Crair, M. C., Gillespie, D. C. & Stryker, M. P. The role of visual experience in the
development of columns in cat visual cortex. *Science* **279**, 566-570,
doi:10.1126/science.279.5350.566 (1998).

35　Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of
Cell Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289,
doi:10.1016/j.cels.2018.11.005 (2019).
36　Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296, doi:10.1038/s41592-019-0619-0 (2019).
37　Cheng, S. *et al.* Vision-dependent specification of cell types and function in the
developing cortex. *Cell* **185**, 311-327 e324, doi:10.1016/j.cell.2021.12.022 (2022).
38　Kim, E. J. *et al.* Extraction of Distinct Neuronal Cell Types from within a Genetically
Continuous Population. *Neuron* **107**, 274-282 e276,
doi:10.1016/j.neuron.2020.04.018 (2020).
39　O'Toole, S. M., Oyibo, H. K. & Keller, G. B. Molecularly targetable cell types in
mouse visual cortex have distinguishable prediction error responses. *Neuron* **111**,
2918-2928 e2918, doi:10.1016/j.neuron.2023.08.015 (2023).
40　Jenks, K. R. *et al.* Arc restores juvenile plasticity in adult mouse visual cortex. *Proc*
Natl Acad Sci U S A **114**, 9182-9187, doi:10.1073/pnas.1700866114 (2017).
41　Brakeman, P. R. *et al.* Homer: a protein that selectively binds metabotropic glutamate
receptors. *Nature* **386**, 284-288, doi:10.1038/386284a0 (1997).
42　Barber, C. F., Jorquera, R. A., Melom, J. E. & Littleton, J. T. Postsynaptic regulation
of synaptic plasticity by synaptotagmin 4 requires both C2 domains. *J Cell Biol* **187**,
295-310, doi:10.1083/jcb.200903098 (2009).
43　Aiken, J., Buscaglia, G., Bates, E. A. & Moore, J. K. The alpha-Tubulin gene
TUBA1A in Brain Development: A Key Ingredient in the Neuronal Isotype Blend. *J*
Dev Biol **5**, doi:10.3390/jdb5030008 (2017).
44　Gu, Y. *et al.* Obligatory role for the immediate early gene NARP in critical period
plasticity. *Neuron* **79**, 335-346, doi:10.1016/j.neuron.2013.05.016 (2013).
45　Kwapis, J. L. *et al.* Epigenetic regulation of the circadian gene Per1 contributes to
age-related changes in hippocampal memory. *Nat Commun* **9**, 3323,
doi:10.1038/s41467-018-05868-0 (2018).
46　Paradis, S. *et al.* An RNAi-based approach identifies molecules required for
glutamatergic and GABAergic synapse development. *Neuron* **53**, 217-232,
doi:10.1016/j.neuron.2006.12.012 (2007).
47　Ban, Y. *et al.* Prickle promotes the formation and maintenance of glutamatergic
synapses by stabilizing the intercellular planar cell polarity complex. *Sci Adv* **7**,
eabh2974, doi:10.1126/sciadv.abh2974 (2021).
48　Li, M. *et al.* CDKL5 modulates structural plasticity of excitatory synapses via liquid-
liquid phase separation. *bioRxiv*, 2025.2003.2016.643581,
doi:10.1101/2025.03.16.643581 (2025).
49　Cai, H., Vernon, R. M. & Forman-Kay, J. D. An interpretable machine learning
algorithm to predict disordered protein phase separation based on biophysical
interactions. *bioRxiv*, 2022.2007.2006.499043, doi:10.1101/2022.07.06.499043
(2022).
50　Li, J. *et al.* Post-translational modifications in liquid-liquid phase separation: a
comprehensive review. *Mol Biomed* **3**, 13, doi:10.1186/s43556-022-00075-2 (2022).
51　Ripin, N. & Parker, R. Formation, function, and pathology of RNP granules. *Cell* **186**,
4737-4756, doi:10.1016/j.cell.2023.09.006 (2023).
52　Youn, J. Y. *et al.* Properties of Stress Granule and P-Body Proteomes. *Mol Cell* **76**,
286-294, doi:10.1016/j.molcel.2019.09.014 (2019).
53　Hubstenberger, A. *et al.* P-Body Purification Reveals the Condensation of Repressed
mRNA Regulons. *Mol Cell* **68**, 144-157 e145, doi:10.1016/j.molcel.2017.09.003
(2017).

| 1416 | 54 | Wilbertz, J. H. *et al.* Single-Molecule Imaging of mRNA Localization and Regulation
| 1417 | | during the Integrated Stress Response. *Mol Cell* **73**, 946-958 e947,
| 1418 | | doi:10.1016/j.molcel.2018.12.006 (2019).
| 1419 | 55 | Kodali, S. *et al.* RNA sequestration in P-bodies sustains myeloid leukaemia. *Nat Cell*
| 1420 | | *Biol* **26**, 1745-1758, doi:10.1038/s41556-024-01489-6 (2024).
| 1421 | 56 | Sato, K. & Kotani, T. Visualizing the translational activation of a particular mRNA in
| 1422 | | zebrafish embryos using in situ hybridization and proximity ligation assay. *STAR*
| 1423 | | *Protoc* **5**, 102951, doi:10.1016/j.xpro.2024.102951 (2024).
| 1424 | 57 | De Jesus-Cortes, H. *et al.* Using the visual cliff and pole descent assays to detect
| 1425 | | binocular disruption in mice. *bioRxiv*, doi:10.1101/2023.05.29.542767 (2024).
| 1426 | 58 | Boone, H. C. *et al.* Natural binocular depth discrimination behavior in mice explained
| 1427 | | by visual cortical activity. *Curr Biol* **31**, 2191-2198 e2193,
| 1428 | | doi:10.1016/j.cub.2021.02.031 (2021).
| 1429 | 59 | Chen, T. W. *et al.* Ultrasensitive fluorescent proteins for imaging neuronal activity.
| 1430 | | *Nature* **499**, 295-300, doi:10.1038/nature12354 (2013).
| 1431 | 60 | Tan, L., Tring, E., Ringach, D. L., Zipursky, S. L. & Trachtenberg, J. T. Vision
| 1432 | | Changes the Cellular Composition of Binocular Circuitry during the Critical Period.
| 1433 | | *Neuron* **108**, 735-747 e736, doi:10.1016/j.neuron.2020.09.022 (2020).
| 1434 | 61 | Chang, J. T., Whitney, D. & Fitzpatrick, D. Experience-Dependent Reorganization
| 1435 | | Drives Development of a Binocularly Unified Cortical Representation of Orientation.
| 1436 | | *Neuron* **107**, 338-350 e335, doi:10.1016/j.neuron.2020.04.022 (2020).
| 1437 | 62 | Tan, L. M., Ringach, D. L. & Trachtenberg, J. T. The Development of Receptive
| 1438 | | Field Tuning Properties in Mouse Binocular Primary Visual Cortex. *Journal of*
| 1439 | | *Neuroscience* **42**, doi:10.1523/Jneurosci.1702-21.2022 (2022).
| 1440 | 63 | Espinosa, J. S. & Stryker, M. P. Development and plasticity of the primary visual
| 1441 | | cortex. *Neuron* **75**, 230-249, doi:10.1016/j.neuron.2012.06.009 (2012).
| 1442 | 64 | Wang, B. S., Sarnaik, R. & Cang, J. Critical period plasticity matches binocular
| 1443 | | orientation preference in the visual cortex. *Neuron* **65**, 246-256,
| 1444 | | doi:10.1016/j.neuron.2010.01.002 (2010).
| 1445 | 65 | Jenks, K. R. & Shepherd, J. D. Experience-Dependent Development and Maintenance
| 1446 | | of Binocular Neurons in the Mouse Visual Cortex. *Cell Rep* **30**, 1982-1994 e1984,
| 1447 | | doi:10.1016/j.celrep.2020.01.031 (2020).
| 1448 | 66 | Smith, S. L. & Trachtenberg, J. T. Experience-dependent binocular competition in the
| 1449 | | visual cortex begins at eye opening. *Nat Neurosci* **10**, 370-375, doi:10.1038/nn1844
| 1450 | | (2007).
| 1451 | 67 | Ukita, J. Causal importance of low-level feature selectivity for generalization in
| 1452 | | image recognition. *Neural Netw* **125**, 185-193, doi:10.1016/j.neunet.2020.02.009
| 1453 | | (2020).
| 1454 | 68 | Sun, Y. J., Espinosa, J. S., Hoseini, M. S. & Stryker, M. P. Experience-dependent
| 1455 | | structural plasticity at pre- and postsynaptic sites of layer 2/3 cells in developing
| 1456 | | visual cortex. *Proc Natl Acad Sci U S A* **116**, 21812-21820,
| 1457 | | doi:10.1073/pnas.1914661116 (2019).
| 1458 | 69 | Song, J. Molecular Mechanisms of Phase Separation and Amyloidosis of ALS/FTD-
| 1459 | | linked FUS and TDP-43. *Aging Dis* **15**, 2084-2112, doi:10.14336/AD.2023.1118
| 1460 | | (2024).
| 1461 | 70 | Liu, N., Guo, Y., Ning, S. & Duan, M. Phosphorylation regulates the binding of
| 1462 | | intrinsically disordered proteins via a flexible conformation selection mechanism.
| 1463 | | *Commun Chem* **3**, 123, doi:10.1038/s42004-020-00370-5 (2020).
| 1464 | 71 | Maharana, S. *et al.* RNA buffers the phase separation behavior of prion-like RNA
| 1465 | | binding proteins. *Science* **360**, 918-921, doi:10.1126/science.aar7366 (2018).

1466	72	Di Stefano, B. *et al.* The RNA Helicase DDX6 Controls Cellular Plasticity by
1467		Modulating P-Body Homeostasis. *Cell Stem Cell* **25**, 622-638 e613,
1468		doi:10.1016/j.stem.2019.08.018 (2019).
1469	73	Courel, M. *et al.* GC content shapes mRNA storage and decay in human cells. *Elife* **8**,
1470		doi:10.7554/eLife.49708 (2019).
1471	74	Marcelo, A., Koppenol, R., de Almeida, L. P., Matos, C. A. & Nobrega, C. Stress
1472		granules, RNA-binding proteins and polyglutamine diseases: too much aggregation?
1473		*Cell Death Dis* **12**, 592, doi:10.1038/s41419-021-03873-8 (2021).
1474	75	Sheth, U. & Parker, R. Decapping and decay of messenger RNA occur in cytoplasmic
1475		processing bodies. *Science* **300**, 805-808, doi:10.1126/science.1082320 (2003).
1476	76	Dar, S. A. *et al.* Full-length direct RNA sequencing uncovers stress granule-
1477		dependent RNA decay upon cellular stress. *Elife* **13**, doi:10.7554/eLife.96284 (2024).
1478	77	Blake, L. A., Watkins, L., Liu, Y., Inoue, T. & Wu, B. A rapid inducible RNA decay
1479		system reveals fast mRNA decay in P-bodies. *Nat Commun* **15**, 2720,
1480		doi:10.1038/s41467-024-46943-z (2024).
1481	78	Hallacli, E. *et al.* The Parkinson's disease protein alpha-synuclein is a modulator of
1482		processing bodies and mRNA stability. *Cell* **185**, 2035-2056 e2033,
1483		doi:10.1016/j.cell.2022.05.008 (2022).
1484	79	Bentmann, E., Haass, C. & Dormann, D. Stress granules in neurodegeneration--
1485		lessons learnt from TAR DNA binding protein of 43 kDa and fused in sarcoma. *FEBS*
1486		*J* **280**, 4348-4370, doi:10.1111/febs.12287 (2013).
1487	80	Wolozin, B. & Ivanov, P. Stress granules and neurodegeneration. *Nat Rev Neurosci*
1488		**20**, 649-666, doi:10.1038/s41583-019-0222-5 (2019).
1489	81	Ince-Dunn, G. *et al.* Neuronal Elav-like (Hu) proteins regulate RNA splicing and
1490		abundance to control glutamate levels and neuronal excitability. *Neuron* **75**, 1067-
1491		1080, doi:10.1016/j.neuron.2012.07.009 (2012).
1492	82	Akamatsu, W. *et al.* The RNA-binding protein HuD regulates neuronal cell identity
1493		and maturation. *Proc Natl Acad Sci U S A* **102**, 4625-4630,
1494		doi:10.1073/pnas.0407523102 (2005).
1495	83	DeBoer, E. M. *et al.* Prenatal deletion of the RNA-binding protein HuD disrupts
1496		postnatal cortical circuit maturation and behavior. *J Neurosci* **34**, 3674-3686,
1497		doi:10.1523/JNEUROSCI.3703-13.2014 (2014).
1498	84	Berto, S., Usui, N., Konopka, G. & Fogel, B. L. ELAVL2-regulated transcriptional
1499		and splicing networks in human neurons link neurodevelopment and autism. *Hum*
1500		*Mol Genet* **25**, 2451-2464, doi:10.1093/hmg/ddw110 (2016).
1501	85	Zhou, X. *et al.* Integrating de novo and inherited variants in 42,607 autism cases
1502		identifies mutations in new moderate-risk genes. *Nat Genet* **54**, 1305-1319,
1503		doi:10.1038/s41588-022-01148-2 (2022).
1504	86	Satterstrom, F. K. *et al.* Large-Scale Exome Sequencing Study Implicates Both
1505		Developmental and Functional Changes in the Neurobiology of Autism. *Cell* **180**,
1506		568-584 e523, doi:10.1016/j.cell.2019.12.036 (2020).
1507	87	Rodin, R. E. *et al.* The landscape of somatic mutation in cerebral cortex of autistic
1508		and neurotypical individuals revealed by ultra-deep whole-genome sequencing. *Nat*
1509		*Neurosci* **24**, 176-185, doi:10.1038/s41593-020-00765-6 (2021).
1510	88	Jia, X. *et al.* De novo variants in genes regulating stress granule assembly associate
1511		with neurodevelopmental disorders. *Sci Adv* **8**, eabo7112,
1512		doi:10.1126/sciadv.abo7112 (2022).
1513	89	Eising, E. *et al.* A set of regulatory genes co-expressed in embryonic human brain is
1514		implicated in disrupted speech development. *Mol Psychiatry* **24**, 1065-1078,
1515		doi:10.1038/s41380-018-0020-x (2019).

1516 90 Andrews, S. (Babraham Bioinformatics, Babraham Institute, Cambridge, United
1517   Kingdom, 2010).
1518 91 Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome
1519   alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**,
1520   907-915, doi:10.1038/s41587-019-0201-4 (2019).
1521 92 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program
1522   for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930,
1523   doi:10.1093/bioinformatics/btt656 (2014).
1524 93 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and
1525   dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550,
1526   doi:10.1186/s13059-014-0550-8 (2014).
1527 94 Yu, G., Wang, L. G., Han, Y. & He, Q. Y. clusterProfiler: an R package for
1528   comparing biological themes among gene clusters. *OMICS* **16**, 284-287,
1529   doi:10.1089/omi.2011.0118 (2012).
1530 95 Lau, S. F., Cao, H., Fu, A. K. Y. & Ip, N. Y. Single-nucleus transcriptome analysis
1531   reveals dysregulation of angiogenic endothelial cells and neuroprotective glia in
1532   Alzheimer's disease. *Proc Natl Acad Sci U S A* **117**, 25800-25809,
1533   doi:10.1073/pnas.2008762117 (2020).
1534 96 Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene
1535   expression data analysis. *Genome Biol* **19**, 15, doi:10.1186/s13059-017-1382-0 (2018).
1536 97 Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing
1537   well-connected communities. *Sci Rep* **9**, 5233, doi:10.1038/s41598-019-41695-z
1538   (2019).
1539 98 Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP.
1540   *Nat Biotechnol*, doi:10.1038/nbt.4314 (2018).
1541 99 Hrvatin, S. *et al.* Single-cell analysis of experience-dependent transcriptomic states in
1542   the mouse visual cortex. *Nat Neurosci* **21**, 120-129, doi:10.1038/s41593-017-0029-5
1543   (2018).
1544 100 Choi, H. M., Beck, V. A. & Pierce, N. A. Next-generation in situ hybridization chain
1545   reaction: higher gain, lower cost, greater durability. *ACS Nano* **8**, 4284-4294,
1546   doi:10.1021/nn405717p (2014).
1547 101 Ku, T. *et al.* Multiplexed and scalable super-resolution imaging of three-dimensional
1548   protein localization in size-adjustable tissues. *Nat Biotechnol* **34**, 973-981,
1549   doi:10.1038/nbt.3641 (2016).
1550 102 Park, J. *et al.* Epitope-preserving magnified analysis of proteome (eMAP). *Sci Adv* **7**,
1551   eabf6589, doi:10.1126/sciadv.abf6589 (2021).
1552 103 Zhao, J. *et al.* Specific depletion of the motor protein KIF5B leads to deficits in
1553   dendritic transport, synaptic plasticity and memory. *Elife* **9**, doi:10.7554/eLife.53456
1554   (2020).
1555 104 Reikofski, J. & Tao, B. Y. Polymerase chain reaction (PCR) techniques for site-
1556   directed mutagenesis. *Biotechnol Adv* **10**, 535-547, doi:10.1016/0734-9750(92)91451-
1557   j (1992).
1558 105 Fisher, C. L. & Pei, G. K. Modification of a PCR-based site-directed mutagenesis
1559   method. *Biotechniques* **23**, 570-571, 574, doi:10.2144/97234bm01 (1997).
1560 106 Ip, J. P. *et al.* alpha2-chimaerin controls neuronal migration and functioning of the
1561   cerebral cortex through CRMP-2. *Nat Neurosci* **15**, 39-47, doi:10.1038/nn.2972
1562   (2011).
1563 107 Ye, T. *et al.* Efficient manipulation of gene dosage in human iPSCs using
1564   CRISPR/Cas9 nickases. *Commun Biol* **4**, 195, doi:10.1038/s42003-021-01722-0
1565   (2021).

108   El-Boustani, S. *et al.* Locally coordinated synaptic plasticity of visual cortex neurons
in vivo. *Science* **360**, 1349-1354, doi:10.1126/science.aao0862 (2018).
109   Brainard, D. H. The Psychophysics Toolbox. *Spat Vis* **10**, 433-436 (1997).
110   Pelli, D. G. The VideoToolbox software for visual psychophysics: Transforming
numbers into movies. *Spatial Vision* **10**, 437-442, doi:Doi 10.1163/156856897x00366
(1997).
111   Stringer, C., Pachitariu, M., Steinmetz, N., Carandini, M. & Harris, K. D. High-
dimensional geometry of population responses in visual cortex. *Nature* **571**, 361-365,
doi:10.1038/s41586-019-1346-5 (2019).
112   Mazurek, M., Kager, M. & Van Hooser, S. D. Robust quantification of orientation
selectivity and direction selectivity. *Front Neural Circuits* **8**, 92,
doi:10.3389/fncir.2014.00092 (2014).
113   Székely, G. J., Rizzo, M. L. & Bakirov, N. K. Measuring and testing dependence by
correlation of distances. *The Annals of Statistics* **35**, 2769-2794, 2726 (2007).
114   Kafashan, M. *et al.* Scaling of sensory information in large neural populations shows
signatures of information-limiting correlations. *Nat Commun* **12**, 473,
doi:10.1038/s41467-020-20722-y (2021).
115   Pedregosa, F. *et al.* Scikit-learn: Machine learning in Python. *the Journal of machine*
Learning research **12**, 2825-2830 (2011).

Fig 1

Figure 1. nELAVLs are phosphorylation substrates of CDKL5.

(A) Workflow of phosphoproteomic screening in WT and *Cdkl5* KO primary visual (V1) cortices. V1 tissues were harvested for isobaric peptide labelling, followed by peptide pooling and phosphopeptide enrichment. The samples were then quantitatively identified by liquid chromatography-tandem mass spectrometry (LC-MS/MS).

(B) Volcano plot showing the $\log_2$ (fold change) and $-\log_{10}$ (p-value) of each gene comparing WT and *Cdkl5* KO mice. Differentially expressed phosphopeptides with fold change > 1.2 and < 0.83 are highlighted in pink and blue, respectively. Dashed lines indicate $p < 0.05$ and a fold change > 1.2 or < 0.83. Each point represents one peptide, those with the CDKL5 consensus motif (RPXS*A/RPXS*) are labeled with gene name.

(C) Gene Ontology (GO) term enrichment of down-regulated differentially expressed genes between WT and *Cdkl5* KO mice.

(D) Conserved consensus serine phosphorylation sequence in ELAVL2 (119), ELAVL3 (119) and ELAV*L*4 (131) in human, mouse and rat.

(E) Immunoblotting of immunoprecipitates for phosphorylated serine of ELAVL2, ELAVL3 and ELAVL4 in CDKL5-expressing HEK 293T cell. The mutation of serine to alanine at the RPXS*A motif prevented the phosphorylation.

(F & G) Levels of pS-nELAVLs in visual, frontal and motor cortices were virtually undetectable in *Cdkl5* KO mice. Visual: n=4 mice, frontal and motor: n=3 mice. Unpaired t-test (two-tailed).

(H & I) Increased pS-nELAVLs in visual cortex in WT mice after dark environment (16hr) with 0hr, 1hr and 2hr light induction. n=5 mice. Unpaired t-test (two-tailed). Dot lines indicate the same batch of experiments. Data are presented as mean ± SEM, ** $p < 0.01$. *** $p < 0.001$. **** $p < 0.0001$.

Fig 2

Figure 2. CDKL5 deletion alters activity-dependent gene expression.

(A) Volcano plot showing the $\log_2$ (fold change) and $-\log_{10}$ (FDR-adjusted p-value) of each gene comparing WT and *Cdkl5* KO mouse V1. Differentially expressed genes (DEGs) with $\log_2$ (fold change) > 0.25 and < -0.25 are highlighted in pink and blue, respectively. Activity-dependent genes are highlighted in green. The dashed line indicates FDR-adjusted p-value < 0.05 and $\log_2$ (fold change) > 0.25 and < -0.25.

(B) Hub genes of DEGs between WT and *Cdkl5* KO mouse V1 were identified from PPI network by maximal clique centrality (MCC) algorithm. Edges represent the protein-protein associations. The shading from red to yellow depicts MCC scores from high to low.

(C) UMAP plot visualizing annotated subclasses of WT and *Cdkl5* KO mouse V1.

(D) Matrix plot showing activity-dependent DEGs identified between WT and *Cdkl5* KO mouse V1. 29 overlapped activity-dependent DEGs between bulk and snRNA sequencing are underlined.

(E) Dot plot showing *Cdkl5* and *nElavls* gene expression across subclasses of WT and *Cdkl5* KO mouse V1.

(F) Number of DEGs between WT and *Cdkl5* KO mouse V1 in individual cell types of glutamatergic neurons, genes up- and down-regulated in *Cdkl5* KO compared with WT shown in pink and blue bars, respectively. DEGs were defined as genes with $|\log_2$ fold change$| > 0.25$ (light colored bars) or > 0.58 (dark colored bars) and FDR-adjusted p-value < 0.05.

(G) Dot plot visualizing enrichment results of activity-dependent genes with DEGs. Enrichment was calculated separately for DEGs identified in individual cell types of glutamatergic neurons. Dot size is the number of intersected genes. Color is shaded by $-\log_{10}$ (P value) (one-sided Fisher's exact test). 'ns' indicates P value ≥ 0.05.

(H) Dot plot visualizing *Homer1, Syt4, Tuba1a, Cdh13, Nptx2, Per1, Arc and Prickle1* gene expression levels in sublayers of Layer 2/3 glutamatergic neurons in WT and *Cdkl5* KO mouse V1.

(I) FISH images of mRNA expression of *Fos, Homer1, Arc, Camk2a* in WT and *Cdkl5* KO mouse V1. Scale bars, 20μm.

(J) (Top) Bar plot quantifying *Fos, Homer1, Arc, Camk2a* mRNA expression levels. The quantification of mRNA numbers was normalized to the number of DAPI+ cells per image. (Bottom) Quantification of *Fos, Homer1, Arc, Camk2a* mRNA punctum volume. *Fos* and *Homer1*: n=4 mice, *Arc and Camk2a*: n=3 mice. Mann-Whitney test (two-tailed). Data are presented as mean ± SEM, * p < 0.05. *** p < 0.001. **** p < 0.0001.

Fig 3

Figure 3. Loss of phosphorylation by CDKL5 alters nELAVL phase separation.

(A & B) With epitope-preserving Magnified Analysis of Proteome (eMAP), ELAVL3 (magenta) and nELAVL (red) puncta in *Cdkl5* KO mice showed large-shifted punctum volume distribution among neurons. Scale bar, 10 μm and 3 μm for zoomed-in images (right) of ELAVL3 and nELAVL. WT: n=21 neurons from 3 mice; *Cdkl5* KO: n=27 neurons from 3 mice. Left: Mann-Whitney test (two-tailed); Right: Two-way ANOVA with Sidak's multiple comparisons test on main effect of WT and *Cdkl5* KO.

(C) Super-resolution STED microscopy of overexpressed WT and SE/SA mutant of ELAVL3 in rat primary hippocampal neurons. Compared to SE, SA mutant showed large-shifted punctum volume distribution among neurons. Scale bar, 5 μm. n=4 cells from 3 batches of experiments. Left: One-way ANOVA followed by Turkey's multiple comparisons test; Right: Two-way ANOVA with Sidak's multiple comparisons test on main effect of SE and SA.

(D) Time-lapse images showing that ELAVL3 droplets fused with each other over time in ELAVL3-SA-overexpressing HEK 293T cells. Scale bar, 3 μm.

(E) Confocal images showing fluorescence recovery between ELAVL3-SA and -SE puncta. Bleaching area radius ~ 2.5 μm. Images represent the ELAVL3 puncta before bleaching and right after bleaching (5s, 40s and 60s). Scale bar, 5μm

(F) Fluorescence recovery rate of bleached SA puncta was slower than those of bleached cytoplasm or bleached SE puncta. n=20 (unbleached cytoplasm), 12 (bleached cytoplasm), 24 (bleached SA puncta) and 17 (bleached SE puncta) cells from 3 batches of experiments. Two-way ANOVA with Sidak's multiple comparisons test.

(G) Purified full length ELAVL3 protein underwent phase separation at indicated concentrations. ELAVL3 was sparsely labelled by Cy3 fluorophore at 1%. Scale bar, 5μm.

(H) Time-lapse images showing that ELAVL3 droplets fused with each other over time. Scale bar, 5 μm.

(I) The number of ELAVL3 droplets decreased when NaCl concentration increased. Scale bar, 5 μm.

(J) Phase separation diagram showing phase separation properties within demonstrated NaCl concentration and ELAVL3 concentration. The highlighted red dots: phase separation; grey dots: no phase separation.

(K) Fluorescence recovery analysis showing that ELAVL3 in condensed droplets dynamically exchanged with those in the dilute phase. Data are presented as mean ± SEM, **** p < 0.0001.

Fig 4

Figure 4. CDD patient iPSC-derived neurons forms aberrant nELAVL condensates and altered crosstalk with P-bodies.

(A) An illustration of CDD patient iPSC-derived cortical neurons (iNeurons). iPSCs lines: R550* = CDKL5 deficient, control = wildtype (WT) and isogenic control = R550* that has undergone adenine base editing (ABE) to restore CDKL5 expression.

(B & C) CDKL5 expression levels were comparable between control and isogenic control iNeurons but absent in R550* iNeurons. n=3. One-way ANOVA followed by Tukey's multiple comparisons test.

(D & E) pS-nELAVLs levels were comparable between control and isogenic control iNeurons but reduced in R550* iNeurons. Control: n=6, R550*: n=7, isogenic control: n=8. One-way ANOVA followed by Tukey's multiple comparisons test.

(F & G) STED microscopy showed large-shifted punctum volume distribution of ELAVL3 (green) and nELAVLs (red) in R550* iNeurons, in comparison to control and isogenic control iNeurons. Scale bar, 5 μm. n=6 cells from 3 batches of experiments. Left: One-way ANOVA with Tukey's multiple comparisons test; Right: Two-way ANOVA with Sidak's multiple comparisons test on main effect of Control, R550* and Iso control; Multiple comparisons: top: $p < 0.0001$ for Control vs R550*, $p < 0.0001$ for R550* vs Iso control; bottom: $p < 0.0001$ for Control vs R550*, $p < 0.0001$ for R550* vs Iso control.

(H & I) STED microscopy showed large-shifted punctum volume distribution of GW182 (red) in R550* iNeurons, in comparison to control and isogenic control iNeurons. Scale bar, 5μm. n=6 cells from 3 batches of experiments. Left: One-way ANOVA with Tukey's multiple comparisons test; Right: Two-way ANOVA with Sidak's multiple comparisons test on main effect of Control, R550* and Iso control; Multiple comparisons: top: $p < 0.0001$ for Control vs R550*, $p < 0.0001$ for R550* vs Iso control; bottom: $p < 0.0001$ for Control vs R550*, $p < 0.0001$ for R550* vs Iso control.

(J) Co-immunoprecipitation of GW182 and ELAVL3-SE/SA from cytoplasmic extracts of HEK 293T cells. Immunoprecipitation was performed using an anti-GW182 antibody. ELAVL3 bands were revealed with an anti-ELAVL3 antibody.

(K) Quantification results of Co-immunoprecipitation, the GW182 capture more ELAVL3-SE than ELAVL3-SA. n=3 batches of experiments. Unpaired t-test (two-tailed). Data are presented as mean ± SEM, * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$.

Fig 5

Figure 5. CDKL5 modulate mRNA processing by regulating nELAVL condensates

(A & B) FISH images of mRNA expression of *Fos* in ELAVL3-SE/SA expressing rat primary neurons. Compared to SE, ELAVL3-SA expressing neurons showed less *Fos* RNA puncta. Scale bar, 5μm. n=10 (SE) and 8 (SA) from 4 batches of experiments. Mann-Whitney test (two-tailed).

(C) EMSA showing different concentration of nELAVL-SE and -SA protein binding with *Fos* RNA. At higher protein concentration, the *Fos* RNA bound fraction in SE group is significantly higher than SA group. n=3. Two-way ANOVA with Sidak's multiple comparisons test.

(D & E) FISH images of mRNA expression of *Fos* in Control and R550* iNeurons after 2 μg/mL Actinomycin D (Act D) treatment. Compared to control group, CDD iNeurons showed shorter *Fos* mRNA half-life. Scale bar, 5μm. n=12 iNeurons from 3 batches of experiments. Two-way ANOVA with Sidak's multiple comparisons test on main effect of Control and R550*. Data are presented as mean ± SEM, ** p < 0.01; **** p < 0.0001.

Fig 6

Figure 6. CDKL5 deletion leads to impaired visual function and nELAVL phosphorylation is necessary for visual processing.

(A) Graphical design of the visual cliff task with measurements. Animals were placed on a platform (3 cm height) in the middle of the open field (left). Percentage of descents from the cliff side per mouse (5 trials per mouse) between WT and Cdkl5 KO mice (right). WT: n=16 mice, *Cdkl5* KO: n=16 mice. Mann-Whitney U test (two-tailed).

(B) Head-fixed wildtype control (WT) mice and *Cdkl5* KO were presented, either to one eye or both eyes, with visual stimuli of different directional grating patterns. In the binocular visual cortex (bV1), calcium signal in layer 2/3 (L2/3) neurons were live-imaged with two-photon laser microscopy.

(C) Field of views (FOVs) showing grating and natural scene responsive neurons in WT (left) and *Cdkl5* KO (right) mice. Scale bar, 20 μm.

(D) Traces showing grating responsive neurons in WT and *Cdkl5* KO mice across 3 viewing conditions, which are binocular (Binoc), contralateral (Contra) and ipsilateral (Ipsi) viewing relatively to the brain hemisphere imaged. Yellow regions indicate grating presentation periods. Gray lines represent the Z-score from all trials and the colored lines show trial average.

(E) Boxplots showing the orientation selective index (OSI) of *Cdkl5* KO mice were reduced across Binoc, Contra and Ipsi conditions under grating (s.f. 0.08cpd). Binoc OSI: 870 neurons from 6 WT and 698 neurons from 6 *Cdkl5* KO mice. Contra OSI: 794 neurons from 6 WT and 693 neurons from 6 *Cdkl5* KO. Ipsi OSI: 629 neurons from 6 WT and 476 neurons from 6 *Cdkl5* KO. The centerlines represent median values, and the whiskers connect the nonoutlier minimum and maximum values to 0.25 and 0.75 quartiles respectively. Outliers are values greater than 1.5 interquartile range away from the quartiles. Mann-Whitney U test (two-tailed).

(F) Scatter plots comparing OSI and correlation of grating (s.f. 0.08cpd) responsive neurons in WT mice and *Cdkl5* KO mice. Color bar shows the correlation between tuning curves. 301 neurons from 6 WT and 211 neurons from 6 *Cdkl5* KO. Mann-Whitney U test (two-tailed).

(G) Boxplots showing fraction of neurons per FOV within the blue region (OSI > 0.5) and red region (OSI < 0.5) from E. 26 FOV from 6 WT mice and 21 FOV from 6 *Cdkl5* KO mice. Mann-Whitney U test (two-tailed).

(H) Distribution of Contra and Ipsi eye correlation between WT and *Cdkl5* KO under grating (s.f. 0.08cpd) stimuli. 301 neurons from 6 WT and 211 neurons from 6 *Cdkl5* KO. Mann-Whitney U test (two-tailed).

(I) Distribution of difference in preferred orientation between WT and *Cdkl5* KO under grating (s.f. 0.08cpd) stimuli. 301 neurons from 6 WT and 211 neurons from 6 *Cdkl5* KO. Mann-Whitney U test (two-tailed).

(J) Traces showing natural scene responsive neurons in WT and *Cdkl5* KO mice across 3 viewing conditions. Pink regions indicate natural scene presentation periods. Gray lines represent the Z-score from all trials and the colored lines show trial average.

1889

1890 (K) Distribution of contra and ipsi eye correlation between WT and *Cdkl5* KO under natural
1891 scene stimuli. 529 neurons from 6 WT and 274 neurons from 6 *Cdkl5* KO. Mann-Whitney U
1892 test (two-tailed).

1893

1894 (L) Mice were virally injected to express GCaMP6f (Ctrl), GCaMP6f and nELAVL (WT),
1895 GCaMP6f and nELAVL-SE (SE) or GCaMP6f and nELAVL-SA (SA). Head-fixed mice
1896 were presented, either to one eye or both eyes, with visual stimuli of different directional
1897 grating patterns and natural scene. In the binocular visual cortex (bV1), calcium signal in
1898 layer 2/3 (L2/3) neurons were live-imaged with two-photon laser microscopy.

1899

1900 (M) Boxplots showing OSI were reduced in WT, SE and SA mice across Binoc, Contra and
1901 Ipsi conditions under grating (s.f. 0.08cpd). Binoc OSI: 870 neurons from 6 Ctrl, 670 neurons
1902 from 5 WT injected, 532 neurons from 4 SE injected and 782 neurons from 5 SA injected
1903 mice. Contra OSI: 794 neurons from 6 Ctrl, 715 neurons from 5 WT injected, 500 neurons
1904 from 4 SE injected and 852 neurons from 5 SA injected mice. Ipsi OSI: 629 neurons from 6
1905 Ctrl, 574 neurons from 5 WT injected, 384 neurons from 4 SE injected and 613 neurons from
1906 5 SA injected mice. Kruskal-Wallis test followed by Tukey's multiple comparisons test.

1907

1908 (N) Boxplots showing Contra and Ipsi correlation under grating (s.f. 0.08cpd) stimuli in Ctrl,
1909 WT injected, SE injected and SA injected mice. 301 neurons from 6 Ctrl, 253 neurons from 5
1910 WT injected, 159 neurons from 4 SE injected and 275 neurons from 5 SA injected mice.
1911 Kruskal-Wallis test followed by Tukey's multiple comparisons test.

1912

1913 (O) Boxplots showing difference in preferred orientation under grating (s.f. 0.08cpd) stimuli
1914 in Ctrl, WT injected, SE injected and SA injected mice. 301 neurons from 6 Ctrl, 253 neurons
1915 from 5 WT injected, 159 neurons from 4 SE injected and 275 neurons from 5 SA injected
1916 mice. Kruskal-Wallis test followed by Tukey's multiple comparisons test.

1917

1918 (P) Boxplots showing Contra and Ipsi correlation under natural scene stimuli in Ctrl, WT
1919 injected, SE injected and SA injected mice. 529 neurons from 6 Ctrl, 185 neurons from 5 WT
1920 injected, 428 neurons from 4 SE injected and 443 neurons from 5 SA injected mice. Kruskal-
1921 Wallis test followed by Tukey's multiple comparisons test. Data are presented as mean $\pm$
1922 SEM, * $p < 0.05$. ** $p < 0.01$. *** $p < 0.001$. **** $p < 0.0001$.

1923