Population geometry reveals directed coupling and transient bistability in spontaneous pituitary secretion

Ana Aquiles[1,*], Aparicio Arias Juri[1,2], Lafont Chrystel[1], David Hodson[4], Santiago-Andres Yorgui[3], Patrice Mollard[1,*], and Tatiana Fiordelisio[3,*]

[1]Institute of Functional Genomics, University of Montpellier, CNRS, INSERM, Montpellier, France.
[2]BioCampus Montpellier, University of Montpellier, CNRS, INSERM, Montpellier, France
[3]Laboratorio de Neuroendocrinología Comparada, Facultad de Ciencias, Universidad Nacional Autónoma de México, Ciudad Universitaria, Ciudad de México 04510, México.
[4]Oxford Centre for Diabetes, Endocrinology and Metabolism (OCDEM), NIHR Oxford Biomedical Research Centre, Churchill Hospital, Radcliffe Department of Medicine, University of Oxford, Oxford, UK.

Summary

The pituitary gland operates as an organized signaling network in which endocrine cell populations coordinate hormone secretion, through homotypic and heterotypic interactions, yet the contribution of spontaneous intrinsic activity in shaping population-level dynamics remains poorly understood. Using geometric analysis of population trajectories — including subspace alignment, manifold separation, and directed coupling metrics — we identified two classes of spontaneous oscillatory signals associated with distinct cell populations exhibiting asymmetric geometric dominance and a reproducible temporal lag. Our results support that spontaneous activity generates a self-sustained oscillator exhibiting transient bistability, linked to increased physiological demand, with slow oscillations reflecting the properties of an excitatory resonator capable of self-oscillating dynamics without external drive. A low-rank recurrent neural network model recapitulated the empirical geometric landscape under three coupling conditions, confirming that directed population coupling underlies the observed coordination. These findings suggest that intrinsic population dynamics play a central role in coordinating pituitary secretion, with implications for understanding hormonal dysregulation in secretory adenomas and other pituitary disorders.

1 Main

Pituitary cell communication is widely recognized as a network that is organized and an integral sum of the interactions involved in sending and receiving signals among its diverse cellular composition [1]. This communication improves functional outcomes and highlights its crucial role in efficient functioning and adaptation [2]. However, our understanding of how cell signaling patterns contribute to spontaneous endogenous secretion and plasticity remains limited. It is essential to investigate spontaneous activity and its influence on homotypic interactions (within the same endocrine cell group) versus heterotypic interactions (between different endocrine cell groups). These differences in signaling coud be the primary responsible for the variability in activation oscillations when responding to physiological challenges [3–5]. For example, in pathological conditions, non-functioning pituitary adenomas are often referred to as "silent" because they do not produce hormones [6]. However, there are also adenomas that result in hormonal overproduction [7]. The origins and causes of these functional differences are not yet fully understood. Therefore, gaining insight into the role of spontaneous signal transmission in triggering secretion may help clarify why some conditions remain non-secretory, while others lead to hormonal secretion in both health and disease.

---

[1]***Correspondence:** Tatiana Fiordelisio, `tfiorde@ciencias.unam.mx`
[2]***Correspondence:** Ana Aquiles, `ana.aquiles@igf.cnrs.fr`

Previous research consistently describes how signal-transmission achievements in pituitary endocrine cell function occur through various modes of cell-to-cell communication [2, 5, 8–12], and all of these mechanisms are influenced by external stimuli, i.e. a hypothalamus hormone factor or further hormone modulation. Although spontaneous activity has received less attention, two groups of endocrine cells—lactotrophs and somatotrophs—exhibit significant spontaneous activity, enabling them to secrete hormones independently of external triggers [13–15]. Understanding the diversity of information integration between these cell groups is essential, as it is influenced by the structure of their communication networks [5, 8]. For example, in the homotypical lactotroph structure, it is involved in physiological adaptations, such as the rearrangement of lactotroph signal connectivity during lactation or pregnancy [5] , or structurally, the shape and the density of cell proportion also present rearrangement [5, 16]. There is also evidence of mixed communication and coordination among different endocrine cell groups, which allows the gland to quickly respond to physiological demands [17]. However, there is limited information about the cause-and-effect relationship under the conditions of intrinsic population dynamics in the pituitary, particularly regarding homotypic coordination and potential heterotypic differentiation. The rationale behind these research findings is based on the intrinsic population dynamics of the pituitary, coordinating homotypically and specifically in response to metabolic demands such as the lactation and reproductive long-term plasticity in females. Furthermore, the regulation of cell signal transmission may involve multiple types of endocrine cells and could influence the systemic mechanisms through which cells send and receive signals spontaneously [3, 4, 13]. In this research, we identified two classes of signals with different spontaneous activities: one exhibiting slow oscillations and the other showing faster oscillations. We built upon the long-term knowledge of a specific endocrine cell type known for its spontaneous secretion capabilities. To delve deeper into this topic, we assessed the spontaneous activity of lactotrophs, previously reported in the literature [5], under conditions that mimic those observed in heterotypic interactions, by evaluating the population code in both homotypic and heterotypic networks, using common metrics to describe network coordination but also the geometrical features of population dynamics. Our findings revealed that during spontaneous activity, a self-sustained oscillator is generated, resulting in momentary bistability in cell organization that is associated with an increase in physiological demand. This mechanism may explain how population leaders coordinate within the geometric structure of the cell networks. We emphasize the presence of slow oscillations in both homotypic and heterotypic interactions, showcasing the computational properties of an exciting resonator that supports a self-oscillating system. This system is capable of adapting to changes in heterogeneous interactions, enabling it to meet physiological demands and facilitating secretion pulses. Our research highlights the importance of both excitatory and inhibitory-kind activities, particularly during heterogeneous integration in hormone secretion. This understanding could improve our knowledge of spontaneous pituitary secretion, particularly in secretory tumors and other hormonal disorders.

2 Results

2.1 Spontaneous calcium signals wave features in homotypical and heterotypical interactions

To describe the temporal wave signatures during spontaneous activation in both homotypical and heterotypical interactions, we revisited previous calcium slice recordings from lactotrophs. These recordings were taken from four main physiological states: a basal hormonal stimulation, the virgin condition, during lactation, weaned, and the multipara condition. Additionally, we applied this methodology to assess the overall population activity during these same states. These physiological conditions have been associated with sustained long-term integration or cell-network coordination that leads to increased physiological demand [5, 18, 19]. Spontaneous calcium traces display differentiation in their wave shapes, regardless of the physiological effects. However, the type of interaction influences these wave shapes, as illustrated in panel A of Figure 1. Despite the visual differences, both types share a signal coupling intuition characterized by a low-frequency component coupled with a higher one. This intuition led us to explore Phase-Amplitude Coupling between both frequencies, extracting the modulation index that determines the level of coupling. As shown in the representative example top right panel of Figure 1A, the lowest phase frequencies show modulation across a wide range of amplitude frequencies in both homotypic and heterotypic spontaneous activity.

Group comparison indicates a wide range of dominant amplitude frequency modulation from 0.1 to 0.6 mHz (Figure 1A, right panel, green line), suggesting a flexible adaptation in modulating different frequency values. In contrast, homotypic interactions remain around a consistent modulation index until

there is a discrete increase in amplitude frequencies between 0.6 and 0.7 mHz (blue line). This marks the starting point for differences in signal modulation between the two types of interactions, suggesting a signal preference in terms of frequency values. Regarding the evolution of the modulation index value, it is notable that homotypic interactions present the lowest modulation values across different physiological states, while also exhibiting high variability during lactant and weaned conditions. This behavior is also observed in heterotypic interactions, but with larger values. Concerning the aperiodic component features, we also measured the aperiodic exponent in the logarithmic spectrum decay of frequencies (see Figure 1B). The homotypic spectrum follows a hierarchical order that corresponds to the increase in hormonal demand of the physiological state [5]. In contrast, heterotypic interactions lack this tendency but exhibit higher variance in the spectrogram decay during virgin conditions. These observations are further verified by computing the aperiodic exponent in each spectrum (Figure 1C). Recent research on the biological context of aperiodicity in signal transmission in brain activity indicates that observed trends may illustrate specialization in plasticity. This could suggest a loss of rhythmic oscillation or limitations in signaling due to prior experiences. Additionally, these patterns may reflect the balance of excitation and inhibition [20–22]. Specifically, lower aperiodic values—characterized by a flattened decay of frequencies—might indicate greater inhibitory activation within the system. Conversely, higher aperiodic values, which show a pronounced decay in frequency, could be associated with a dominance of excitatory activity. Despite this understanding, further studies investigating these dynamics remain necessary. In our results, we will also use the interpretation of this values as a proposethe existence of similar phenomena based on these findings. On the other hand, the variability in heterotypic interactions during virgin conditions suggests a flexible approach to signal communication, adapting to upcoming hormonal challenges. Notably, during lactation and weaning, this variance decreases significantly, as do the aperiodicity values. This reduction may reflect important hormonal shifts. During lactation, there is a need for coordination to compensate for prolactin (PRL) secretion, while during offspring weaning, this coordination is reshaped to stop PRL production and allow other hormones, such as follicle-stimulating hormone (FSH) and luteinizing hormone (LH), to reestablish the reproductive cycle. The values obtained during these phases are the lowest when compared to homotypic networks, which may indicate a specialization in signal coordination to meet hormonal output requirements. In the multipara condition, both networks show the lowest values, yet heterotypic interactions demonstrate an increased variance. This increase may be related to well-established integration in signal transmission, facilitating the anticipation of the next fertilization event and the subsequent initiation of hormonal cycling.

2.2 Signal classification and its relation to functional relevance

Recognizing both network interactions reveals periodic and aperiodic particularities among different hormonal challenges. Our subsequent exploration focused on classifying cellular signals by utilizing all components involved in the oscillation. To achieve this, we employed a spectral decomposition algorithm for signal analysis [23]. We discovered that under virgin conditions, spontaneous activity could be categorized into two major classes, as illustrated in Figure 2A. This classification corresponds to the various conditions we evaluated. Notably, in most cases, the first two principal components from the spectral dimensionality reduction accounted for the majority of the variance in both networks: homotypic and heterotypic. This finding highlights the low dimensionality inherent in the pituitary system, which we will explore further in subsequent sections. Biologically speaking, this suggests that the system is structured and conserved, with a few key variables effectively explaining its behavior. It also indicates that signal transmission is primarily influenced by a limited range of frequencies or signal harmonics, which ties into the findings presented in Figure 1. Additionally, this may suggest that the system is less sensitive to noise. It is important to highlight that signal classification is consistent in terms of wave shape. For example, Class 1 consists of cells that exhibit slower frequencies, while Class 2 primarily includes signals with higher frequencies (see Figure 2A). To explore the relationship between these classes and their aperiodic components, we measured the sensitivity of the classes in relation to their aperiodic values. Our findings indicate that, in most instances, mainly in the homotypical network, the lowest aperiodic values (represented by cooler colors) were associated with cell signals from Class 1. Conversely, Class 2 mostly represented higher aperiodic values, depicted by warmer colors, across various conditions and populations, as first investigated through adjacency matrix arrangements (see Figure 2B). To validate this relationship, we employed surrogate resampling of aperiodic values from each population and condition to extract average values and internal errors. We observed that homogeneous populations consistently demonstrated a clear distinction between Class 1—characterized by lower mean aperiodic values—which tended to increase across the evaluated conditions in the homotypical population. In contrast, Class 2

showed an increase in mean aperiodic values, peaking notably in the Weaned condition. This peak could be attributed to the necessity of halting hormone secretion, thus requiring greater inhibitory integration. In the case of the heterotypic population, there was no significant separation concerning Classes and aperiodic values. This suggests that this feature exists in both classes, indicating reduced independence between them. This could be due to the influence of other endocrine cell groups that are not necessarily related to prolactin (PRL) secretion but contribute to the reshaping of network communication , refer to Figure 2B, squared boxplot. Continuing our exploration of the differences between homotypic and heterotypic networks, our next question focused on the amount of information (H) that both classes carry, specifically related to the specialization of the signal. However, the information distribution values did not indicate a preference between the two network types and remained relatively consistent across conditions (see Figure 2C). Notably, only the Class 2 (H) values during the weaned condition in the heterotypic network showed an amplification in rank values when compared to Class 1 of the same network, as well as Class 2 of the homotypic network. Building on our observations regarding the aperiodic value (refer to Figure 2B), we next seek a global relationship (without classes distinction) between these values, we found that the correlation values in homotypic networks were close to zero and negative, while the heterotypic networks exhibited a trend of increased anti-correlated values across conditions, showing only significant correlation in the lactant, weaned, and multipara conditions. Class 2 cells were less responsive to changes and exhibited similar trends across populations during the Virgin and Lactant conditions. This reflected a less pronounced indirect correlation according to the increase in aperiodic values within both networks. During the lactation period, there was a noticeable shift in trends, showing an increasing tendency compared to the aperiodic values. Additionally, during the weaning and multipara periods, the trends varied slightly between populations, but no significant differences were found. This evaluation suggests an independent relationship between the amount of information carried by the signal classes and the aperiodic value. In other words, even though the aperiodicity of the value changes between classes and conditions, the amount of information each carries is still relevant for signal transmission.

2.3   Signal synchronization and directionality across the time

Signal synchrony is a key phenomenon used to describe the existence of network coordination within the pituitary gland [5, 8, 17]. We explore the relationship between signal correlation values without class distinction, focusing on positive (synchronous) and negative (asynchronous) correlations to determine if these rankings exhibit differential performance between homotypical and heterotypical populations. Despite significant correlations, the distribution of these values shows a symmetrical proportion between positive and negative values across the evaluated conditions (see Figure 3A). However, the cell ratio that contains positive and negative values differs between the two populations; on one hand, the homotypical network under virgin conditions starts with a higher ratio of positively correlated cells and a lower ratio of negatively correlated cells, indicating more positive correlations in virgin conditions and fewer anti-correlated values. This trend suggests a decrease in the ratio of positive correlations and an increase in negative correlations across different hormonal demands. On the other hand, the heterotypical cell ratio begins with nearly equal ratios in virgin conditions, then tends to increase during lactation. During the weaning and multipara conditions, it returns closer to the virgin cell ratios. These differences highlight how the network encodes and manages signal communication. While the homotypical interactions exhibit opposite trends in cell ratio performance, nearing equal ratios in the multipara condition, the heterotypical networks demonstrate a more intuitive pattern, increasing the positive correlation ratio during periods of high hormonal demand and subsequently decreasing it during weaning. Regarding the mean correlation values governing homotypical interactions (see Figure 3B), there is a trend of decreasing correlation values associated with reduced demands. However, in the multipara condition, there is a resurgence of positive values, which may indicate the sustained existence of interactions following the first fertilization. In contrast, heterotypical interactions start with low interval values during virgin conditions, increase during lactation, but ultimately decline to even lower mean values than in the initial state. This difference may suggest a greater prevalence of long-term experience in homotypical networks compared to heterotypical ones. This is likely due to the specialization of the homotypical network, which does not necessitate greater cooperative integration of additional endocrine cell groups after multiple fertilizations.

2.4   Dominance Directionality

Having described the differences between both networks broadly, we now introduce the factor of our classification approach to determine if the correlation between the signals is also sensitive to the class type. After matching the values of correlation, class, and condition, we applied a random forest classifier to

see if a distinction could be made using these variables. However, the classifier struggled in most conditions, except during the virgin homotypic conditions (cross-validation accuracy > 0.5), where the largest separation between cell ratios was observed. Beyond that, the cell classes appeared to be unrelated to the connectivity inferred from the metrics. Nonetheless, an important distinction exists between positive and negative values in both network cases. Unfortunately, Spearman correlation is less sensitive to causal directionality. Therefore, we reverted to information theory metrics, computing directional information based on cell signal classification (see Figure 3D). We found that, in nearly all cases, Class 1 dominated signal communication, with the only exception being the virgin condition in heterotypic populations, which exhibited no dominance between classes. To delve deeper into this finding, we examined the dominance dependence using the embedding time of signals from both Class 1 and Class 2 (see Figure 4). This vision was grounded in the perspective that our results reveal important characteristics regarding network-level communication. As we have observed, both homotypic and heterotypic populations respond differently to aperiodic values, while showing no sensitivity to information content or correlation values. Additionally, heterotypic populations exhibit differences in cell integration related to the aperiodic exponent and correlation values. However, both types of populations demonstrate similar trends in directionality. Although utilizing information theory allows us to approach the system probabilistically, limitations exist. We rely on deterministic probabilistic distributions derived from cell activity, which may not predict the signal with high accuracy. To address this issue, we employed permutation tests and analyzed the lag axis to assess the direction of the signal (refer to the Methods section). Despite these challenges, exploring the embedding times of our system under both homotypic and heterotypic conditions aids in preserving the geometric landscape through which both classes have evolved and synchronized over time. A representative example of this concept is illustrated in Figure 4A, where the embedding time space of Class 1 and Class 2 is projected into the first two dimensions, with the heterotypic network depicted at the bottom. It is visually apparent that Class 1 and Class 2 converge at several time points (shown as red dots), not only under virgin conditions but also during lactation, weaning, and multipara stages. To qualitatively explore this initial evaluation, we quantified the time moments when the two cell classes matched, focusing on the synchronization between classes at different levels. We performed a Procrustean analysis(ref) on the space of the cell classes to determine the shortest distance between them. After aligning the trajectories, we measured local geometric synchronization over tile partitions (windows = 20), assessing the synchrony of both classes by factorizing the matrices resulting from the embedding time. Values approaching 1 indicate a high subspace overlap (strong synchrony), while values near 0 indicate orthogonal, uncorrelated dynamics. The distributions of these values were shown over time for each condition and population (see Figure 4B). Notably, where low synchronization exists, there is also dominance between trajectories. We evaluated this using angle distances in the embedding space (see Figure 4C). As seen in Figure 4B, across all conditions and both networks, dominance is evident, as indicated by the principal angle that separates the trajectories from Class 1 to Class 2. The distribution of angle values is predominantly above the random angle threshold in the space.

To determine which population leads the trajectory, we measured the height between the first peak in Class 1 and the subsequent peak in Class 2 (refer to the squared diagram in Figure 4C). A higher barrier indicates greater dominance. In all conditions, the homotypic trajectories show that Class 1 dominates (Figure 4C). However, in the heterotypic network, we observed an interesting variation in dominance: in the virgin condition, Class 1 holds a dominant position, but during the lactant and weaning conditions, dominance is shared between both classes. The dominance returns to Class 1 during the multipara condition. This variation highlights the key features that differentiate the two networks. In the homotypic network, communication experiences and specialized organization respond to varying hormonal demands. In contrast, the heterotypic network fosters more cooperative communication during the lactation and weaning phases, which require significant organizational changes. This network allows for greater flexibility in switching and relaying dominance coordination, leading us to our next and final exploration. The existence of stability in both homotypic and heterotypic systems is influenced by the roles and characteristics of the signal dynamics.

2.5 Existence of momentaneous bistability in the spontaneous activity dynamics in both homotypic and heterotypic networks

Noticing that the landscape dominance exhibited two well-defined points, and that these differed in dominance between classes, our next step was to investigate the stability of states in each class separately. We found that the landscape state in Class 1 presents one or more point states, although not all of them are stable. In contrast, the landscape for Class 2 is often limited to showing one or two stable states, as

illustrated in the middle panel of Figure 4D. These differences may explain the variation in dominance between populations and classes, as the existence of more than one stable point provides flexibility and robustness [24]. To evaluate our observation, we performed a multistability test along the stable points identified in each landscape for both Class 1 and Class 2, (see Figure 4D, bottom panel). It is important to emphasize that this bistable feature was evaluated in both classes, and thus the percentages of presence or absence of these states encompass both classes. The presence of a bistable or multistable momentum in cell trajectories suggests significant communication dynamics. We noted that in the homotypic case, most landscapes remained monostable, with only a small number of samples exhibiting bistability. This is consistent with the dominance direction observed in Class 1. On the other hand, the heterotypic stable states showed bistability in both lactant and weaned conditions. Among these, lactant conditions displayed the largest proportion of landscapes with multistable dynamics, while only the virgin condition did not exhibit bistability. This could also reflect the presence of cooperation and flexibility in heterotypic temporal dynamics. The relevance of bistable dynamics in coordinating spontaneous secretory activity suggests that our system possesses the capability to dynamically switch between monostable and bistable states. This feature is significant in non-linear biological systems that generate oscillators [24–27] and motivates the use of a generative model to explore how geometric shifts arise from network-level coupling, without requiring external input as a necessary condition. To investigate whether the geometric coordination observed in the biological data could be recapitulated by such a minimal framework, we constructed a low-rank recurrent neural network (RNN) [28] composed of two coupled populations. Given the low-dimensional structure of the empirical trajectories (refer to Figure 2A), each population was modelled as a leaky-integrator RNN with rank-2 recurrent connectivity. Network sizes were chosen to preserve the cellular disproportion observed in the biological system: Population A comprised 80 units and Population B comprised 50 units (Figure 5A). The resulting synthetic time series qualitatively resembled the empirical data, with Population A dynamics recapitulating Class 1 activity patterns and Population B dynamics resembling Class 2 (Figure 5A, insets). Three coupling conditions were simulated to systematically probe the directionality of geometric coordination (Figure 5B). In the directed condition, Case 1, coupling was unidirectional from Population A to Population B, reflecting the hypothesized dominance of Population A observed in the biological data. In the Case 2, reversed condition, the coupling direction was inverted, with Population B driving Population A, providing a critical test of whether geometric dominance follows the direction of network drive. In the Case 3 null condition, no coupling was present between the two populations, serving as a baseline against which the effects of directed coupling could be assessed. Applying the same dimensionality reduction pipeline used for the empirical data, Population A consistently occupied a dominant position in the geometric space across all three conditions, coordinating the trajectory structure regardless of coupling direction (Figure 5B). This result is expected, as the switch in geometric dominance only becomes fully apparent when the subspace angle is explicitly quantified (Figure 5C), confirming that the generative model preserves the core geometric properties of the biological system. Having established the geometric validity of the model, we next examined how tonic external input modulates the latent dynamics under the directed coupling condition. Three input strengths were applied to Population A, mimicking hypothalamic hormonal stimulation. In the absence of input, Population A exhibited concentric oscillations around the trajectory of Population B (Figure 5D). As input strength increased, the radius of these oscillations progressively decreased, with trajectories converging toward a central attractor without ever reaching it — a hallmark of a supercritical Hopf bifurcation [29] (Figure 5D–E). This behaviour is mechanistically significant: it indicates that the system sustains stable limit-cycle oscillations while remaining sensitive to input-driven amplitude modulation, enabling transitions between slow and fast oscillatory regimes. Consistent with this interpretation, the temporal lag between Population A and Population B signals increased with input strength (Figure 5E), reflecting stronger entrainment of Population B by Population A under tonic stimulation.

Together, our findings indicate that the intrinsic spontaneous behavior of the endocrine pituitary cell population, particularly due to heterotypic integration, supports momentary bistable activity. This activity allows for transitions between two types of oscillatory behavior: slow and fast oscillations. Additionally, the dominance of slow oscillations—both in terms of geometry and signal coupling—highlights the presence of excitatory resonator-type activity. This behavior remains consistent under various conditions, including when the system is in a baseline hormonal state or a virgin condition, demonstrating the robustness of the system and its suitability for testing in a low-rank null model.

3 Discussion

Hormone secretion is one of the most critical functions of the pituitary gland, and a defining feature of this process is its capacity for intrinsic self-organization [1, 11, 30]. This autonomous architecture allows the pituitary to sustain coordinated hormonal output independent of external input —a property long associated with endocrine cells capable of spontaneous secretion [31, 32]. Yet spontaneity, also could been organized by for example the emerging of structured $[Ca^{2+}]_i$ oscillatory dynamics conserved across pituitary cell types [3, 4], regardless of cell identity; pituitary endocrine cells generate spontaneous action potentials and $[Ca^{2+}]_i$ transients driven by rhythmic Ca^{2+} entry [33]. In this study we addressed how these oscillations are classified, and how their coupling could establish the organization that governs spontaneous secretion at a system level. We combined data from homotypic lactotroph networks —whose specialization across reproductive and lactational demands has been previously evaluated and published [5] —with matched heterotypic interactions analyses, allowing us to determine whether pituitary homotypical versus heterotypical identity network, is reshaped across physiological states. In both network types, we evaluated spontaneous $[Ca^{2+}]_i$ dynamics using ex-vivo calcium imaging and extracted both the periodic and aperiodic signal components, first, where clear differences emerged in phase coupling and aperiodic exponent values depending on physiological stage (Figure 1), motivating a data-driven classification of oscillatory regimes (Figure 2). This revealed two functionally distinct cell classes: one characterized by slow oscillations (Class 1) and one by fast oscillations (Class 2). We then assessed their respective roles in network communication using both linear and nonlinear connectivity metrics (Figures 2–3), uncovering a consistent dominance direction originating from Class 1, alongside physiologically dependent differences in inferred connectivity strength and topology (Figure 3). Finally, we examined synchronization geometry in both classes using phase-space embedding methods (Figure 4), revealing differential dominance behavior between networks and a transient bistability that seeds a self-sustaining oscillator — one that maintains stable limit-cycle dynamics while remaining sensitive to amplitude modulation by external input (Figure 5). Together, these findings suggest that bistability — arising from the dynamic interplay between slow and fast oscillatory cell classes — is a fundamental organizing principle of pituitary network function, one that endows the gland with the flexibility to adapt secretory output across physiological demands while raising the question of how this intrinsic computational capacity is reconfigured in pathological states or under sustained hormonal challenge. Hormone secretion is one of the most critical functions of the pituitary gland, and a defining feature of this process is its capacity for intrinsic self-organization [1, 11, 30]. This autonomous architecture allows the pituitary to sustain coordinated hormonal output independent of external input —a property long associated with endocrine cells capable of spontaneous secretion [31, 32]. Yet spontaneity, also could been organized by for example the emerging of structured $[Ca^{2+}]_i$ oscillatory dynamics conserved across pituitary cell types [3, 4], regardless of cell identity; pituitary endocrine cells generate spontaneous action potentials and $[Ca^{2+}]_i$ transients driven by rhythmic Ca^{2+} entry [33]. In this study we addressed how these oscillations are classified, and how their coupling could establish the organization that governs spontaneous secretion at a system level. We combined data from homotypic lactotroph networks —whose specialization across reproductive and lactational demands has been previously evaluated and published [5] —with matched heterotypic interactions analyses, allowing us to determine whether pituitary homotypical versus heterotypical identity network, is reshaped across physiological states. In both network types, we evaluated spontaneous $[Ca^{2+}]_i$ dynamics using ex-vivo calcium imaging and extracted both the periodic and aperiodic signal components, first, where clear differences emerged in phase coupling and aperiodic exponent values depending on physiological stage (Figure 1), motivating a data-driven classification of oscillatory regimes (Figure 2). This revealed two functionally distinct cell classes: one characterized by slow oscillations (Class 1) and one by fast oscillations (Class 2). We then assessed their respective roles in network communication using both linear and nonlinear connectivity metrics (Figures 2–3), uncovering a consistent dominance direction originating from Class 1, alongside physiologically dependent differences in inferred connectivity strength and topology (Figure 3). Finally, we examined synchronization geometry in both classes using phase-space embedding methods (Figure 4), revealing differential dominance behavior between networks and a transient bistability that seeds a self-sustaining oscillator — one that maintains stable limit-cycle dynamics while remaining sensitive to amplitude modulation by external input (Figure 5). Together, these findings suggest that bistability — arising from the dynamic interplay between slow and fast oscillatory cell classes — is a fundamental organizing principle of pituitary network function, one that endows the gland with the flexibility to adapt secretory output across physiological demands while raising the question of how this intrinsic computational capacity is reconfigured in pathological states or under sustained hormonal challenge.

3.1 Cell spontaneous classes intrinsically features

The observation of two functionally distinct oscillatory classes that modulate spontaneous intracellular calcium levels $[Ca^{2+}]_i$ is not new. Previous studies have noted qualitative differences in oscillation waveforms, with particular emphasis on pseudo-plateau bursting. This pattern is characterized by sustained membrane depolarization, upon which bursts of action potentials are superimposed, resulting in high-amplitude $[Ca^{2+}]_i$ transients [13, 34]. For a long time, this bursting regime was thought to be the primary driver of hormone secretion, as larger fluctuations in calcium levels were linked to increased exocytotic output [13, 35]. Further ionic mechanisms have been progressively characterized, including studies involving the blockade of BK channels in somatotrophs. This blockade converts bursting into continuous spiking and significantly reduces the amplitude of $[Ca^{2+}]_i$ oscillations [36]. Additionally, inhibiting voltage-gated calcium channels disrupts calcium influx, which is essential for sustaining both activity patterns [37]. What our data add to this picture is not simply the reidentification of these two regimes by our signal classification, but rather a network-level account of how they interact — and why their balance, rather than the dominance of either class alone, constitutes the operative mechanism of flexible secretory control. Regarding periodic differences between network identities, a notable range of phase-coupled frequencies is already apparent in homotypic interactions under baseline hormonal conditions, suggesting that network communication is functionally specialized even within populations of the same cell type. Nevertheless, it is important to acknowledge that purely homotypic interactions may represent an artificial constraint: given the known spatial intermixing of endocrine cell types across the anterior pituitary, heterotypic proximity is the biological default rather than the exception. In this context, the contribution of neighboring cell types to network organization is not only plausible but expected [10, 11, 38], and this is reflected in our data — the range of phase-coupled frequencies in heterotypic networks is broader than in homotypic ones, consistent with a more integrative communication architecture that is better suited to coordinate across oscillatory regimes. The aperiodic component of the signal offers a complementary window into these interactions. Building on work linking the 1/f spectral exponent to excitation/inhibition (E/I) balance — where a flatter power spectrum decay corresponds to a more excitable network state and a steeper decay to a more inhibition-dominated one — we interpret the aperiodic exponent as a proxy for the local E/I regime within each cell class, while noting that this relationship, though compelling, is not universally invariant and should be understood as a working framework [39–41]. Within homotypic networks, the aperiodic exponent follows a physiologically coherent pattern across conditions: Class 1 cells are associated with the lowest exponent values, consistent with a more excitable or tonically active state, while Class 2 cells exhibit the highest exponents, suggesting a relatively inhibition-biased regime. Importantly, this class-specific aperiodic signature is corroborated by correlation synchrony values, which align with physiological expectations and provide independent mechanistic support for these observations. Heterotypic networks, on the contrary, display distinct features in both the periodic and aperiodic domains, as well as in linear connectivity estimates — with class-specific aperiodic signatures becoming less pronounced. We interpret this as a signature of cooperative signal integration: when cell types such as thyrotrophs and gonadotrophs are included in the network [42, 43], their spontaneous electrical activity and paracrine signaling capacity introduce additional degrees of freedom that partially dissolve the strict class boundaries observed in homotypic settings, reshaping the overall communication topology in a physiologically meaningful way.

3.2 Bistability in geometrical coupling relevance

The emergence of a self-sustaining oscillator from the transient bistability we observe between Class 1 and Class 2 populations is not simply a dynamical curiosity — it represents a functional architecture with clear physiological implications. Limit cycle oscillators exhibit self-sustained oscillations, meaning they intrinsically persist in the absence of external periodic cues, and any perturbation that displaces the system from the cycle is damped out as the system asymptotically returns to its orbit [26, 27, 44]. The fact that the pituitary network operates in this regime — rather than as a noise-driven or input-dependent oscillator — has important consequences: it implies that spontaneous hormone secretion is not a passive response to upstream signals but an intrinsically generated output, the amplitude and timing of which can be modulated by input without depending on input for its existence. This is precisely the logic of pituitary-adrenal ultradian rhythmicity that until this moment have been proposed and explored only theoretically [45, 46]. What our phase-space Procrustes analysis adds is a geometric account of how this limit-cycle behavior is organized across populations. Rather than evaluating synchrony only at the level of pairwise signal correlations, we characterize the shape and alignment of the trajectories that each cell class traces through its reconstructed attractor. This approach is grounded in the neural manifold framework

[47, 48], in which the joint activity of a population is represented as a trajectory in a high-dimensional state space, and the geometry of that trajectory — rather than the activity of individual cells — encodes the computational or functional state of the network. Applied to the pituitary, this framework reveals that the two oscillatory classes do not merely differ Hormone secretion is one of the most critical functions of the pituitary gland, and a defining feature of this process is its capacity for intrinsic self-organization [1, 2, 11]. This autonomous architecture allows the pituitary to sustain coordinated hormonal output independent of external input —a property long associated with endocrine cells capable of spontaneous secretion [31, 32]. Yet spontaneity, also could been organized by for example the emerging of structured $[Ca^{2+}]_i$ oscillatory dynamics conserved across pituitary cell types [3, 4], regardless of cell identity; pituitary endocrine cells generate spontaneous action potentials and $[Ca^{2+}]_i$ transients driven by rhythmic Ca^{2+} entry [33]. In this study we addressed how these oscillations are classified, and how their coupling could establish the organization that governs spontaneous secretion at a system level. We combined data from homotypic lactotroph networks —whose specialization across reproductive and lactational demands has been previously evaluated and published [5] —with matched heterotypic interactions analyses, allowing us to determine whether pituitary homotypical versus heterotypical identity network, is reshaped across physiological states. In both network types, we evaluated spontaneous $[Ca^{2+}]_i$ dynamics using ex-vivo calcium imaging and extracted both the periodic and aperiodic signal components, first, where was clear differences emerged in phase coupling and aperiodic exponent values depending on physiological stage (Figure 1), motivating a data-driven classification of oscillatory regimes (Figure 2). This revealed two functionally distinct cell classes: one characterized by slow oscillations (Class 1) and one by fast oscillations (Class 2). We then assessed their respective roles in network communication using both linear and nonlinear connectivity metrics (Figures 2–3), uncovering a consistent dominance direction originating from Class 1, alongside physiologically dependent differences in inferred connectivity strength and topology (Figure 3). Finally, we examined synchronization geometry in both classes using phase-space embedding methods (Figure 4), revealing differential dominance behavior between networks and a transient bistability that seeds a self-sustaining oscillator — one that maintains stable limit-cycle dynamics while remaining sensitive to amplitude modulation by external input (Figure 5). Together, these findings suggest that bistability — arising from the dynamic interplay between slow and fast oscillatory cell classes — is a fundamental organizing principle of pituitary network function, one that endows the gland with the flexibility to adapt secretory output across physiological demands while raising the question of how this intrinsic computational capacity is reconfigured in pathological states or under sustained hormonal challenge.

4   Limitations

The deep anatomical location of the pituitary gland within the mammalian skull presents a fundamental constraint on the scale at which spontaneous population dynamics can be directly observed in vivo. In this study, we addressed this challenge by operating at a mesoscale level of description, using acute ex vivo pituitary slices that preserve the local cellular architecture and intercellular communication pathways while remaining experimentally accessible. Although this preparation provides a realistic framework for characterizing intra-pituitary cell dynamics, it carries an inherent limitation when interpreting our heterotypic network results: because all spontaneously active cells were imaged collectively without prior cell-type identity labeling, we cannot directly attribute the specific aperiodic or connectivity features of heterotypic interactions to defined endocrine populations such as thyrotrophs or gonadotrophs. The cooperative restructuring of network communication we describe in the heterotypic condition is therefore inferred from aggregate signal properties rather than confirmed at the level of identified cell types. Future work employing transgenic reporter lines — in which specific pituitary cell populations express spectrally distinct fluorescent indicators under hormone-promoter control — combined with multi-laser excitation calcium imaging, would allow simultaneous functional imaging of two or more defined cell types within the same preparation, directly testing whether the class-specific dynamics and dominance relationships we describe segregate along cell-type boundaries or reflect a more general organizational principle independent of hormonal identity. Such an approach would substantially strengthen the cooperative heterotypic communication hypothesis advanced here.

5   Conclusion

The findings presented in this study enhance our understanding of how the pituitary gland organizes spontaneous hormone secretion at a network level. Instead of attributing basal secretory activity to the intrinsic properties of individual cell types in isolation, we show that the collective dynamics of two oscillatory cell classes—characterized by their distinct intracellular calcium ($[Ca^{2+}]_i$) signatures, aperiodic profiles, and phase-space geometry—form a functional architecture. This architecture exhibits an emergent property: a self-sustaining, bistability-seeded limit cycle. Through this mechanism, the anterior pituitary gland can integrate complex signals by coordinating the interactions between different cell populations and signaling pathways, which is essential for maintaining physiological homeostasis. Our results provide a clear and robust dynamical mechanism—supported by low-rank replicability—for this integrative capacity. It is the balance between slow and fast oscillatory regimes, along with the directed coupling that Class 1 cells exert over Class 2 cells, which enables the gland to sustain stable secretory rhythms while remaining responsive to physiological demands. Importantly, this framework offers a new perspective for understanding spontaneity in secretory diseases, such as the disruption of secretory processes in hyperfunctioning adenomas.

6   Resource availability

The source code for processed data and all methods here reported is available on GitHub: `https://github.com/AnaAquiles/GeoSync-Pituitary`. Further information or resource will be send it under considerable. Raw datasets are available from the corresponding author upon reasonable request.

7   Acknowledgements

We gratefully acknowledge the valuable discussion on theoretical foundations with Pierre Fontanaud, and the valuable editing suggestion. The authors have received grants from the Agence Nationale de la Recherche (to P.M, ANR-22-CE14-0001) and the Foundation de la Recherche Medical (to A.A, SPF202309017725).

8   Author contributions

A.A., M.P., F.T., and S.A.Y. contributed to the conception of the study and offered experimental method guidance. A.A.J and L.C performed the experimental procedure of calcium imaging recordings. A.A designed the algorithm, performed formal analysis and visualization, and drafted the manuscript. All authors contributed to the review and editing of the manuscript.

9   Declaration of interest

None declared.

10   Methods

All procedures were conducted in accordance with the Animal Welfare guidelines of the European Community and were approved by the relevant institutional ethics committees.

10.1   Animals

Wild-type C57BL/6 female mice (2 month-old-wild-type) were studied across four reproductive states: virgin ($n = 4$), lactant (8–10 days post-partum; $n = 4$), multiparous (second lactation, 8–10 days post-partum; $n = 4$), and weaned (dams weaned at postnatal day 15; $n = 4$). These conditions span the principal neuroendocrine states associated with prolactin and growth hormone secretion, allowing controlled comparison of pituitary network dynamics across reproductive demand.

10.2   Calcium Imaging Datasets

10.2.1   Re-evaluated datasets

A subset of analyses was performed on previously published and publicly reported calcium imaging datasets [5]. For lactotroph populations, four physiological conditions were included: virgin, lactant (8–10 days post-partum), multiparous (second or third lactation), and weaned (females weaned at postnatal day 21). For somatotroph populations, re-evaluation was restricted to the virgin condition. Full methodological details for these datasets are provided in the original publications [5]; only a brief summary is given here for completeness.

10.2.2   *In vitro* calcium imaging

Female mice at the reproductive stages described above were anaesthetised and decapitated. Pituitary glands were extracted and coronally sectioned into $\sim$200 µm slices using a vibratome; three slices per animal and condition were coated with poly-L-lysine, for imaging.

Slices were loaded with the calcium-sensitive fluorescent dye Fluo-4 AM (22 µM; Invitrogen, Eugene OR, USA) dissolved in dimethyl sulfoxide (DMSO; Sigma, St Louis MO, USA) with 0.02% pluronic acid F-127 (Sigma, St Louis MO, USA) in 1 mL of oxygenated Ringer solution (in mM: NaCl 138, KCl 2.5, NaH_2PO_4 1.25, $NaHCO_3$ 3, HEPES 10, glucose 11, $CaCl_2$ 2, $MgCl_2$ 1). Dye loading was carried out for 45 min at 37C in a humidified atmosphere (5% CO_2 / 95% O_2).

Fluorescence imaging was performed on a fluorescence stereomicroscope (SteREO Discovery V12; Carl Zeiss, Oberkochen, Germany) equipped with a 20× water-immersion objective (WplanApo, 20Xw NA=1, Carl Zeiss), coupled to a scientific sCMOS camera (ORCA-Fire; Hamamatsu Photonics, Hamamatsu, Japan) operating at 4× pixel binning. Calcium signals were acquired at one frame per 200 ms (5 Hz) under constant perfusion with oxygenated Ringer solution at 34C for a total duration of 10 min per slice.

10.3   Time-Series Preprocessing

Raw fluorescence time series were preprocessed in four sequential steps to yield stable, comparable signals across cells and animals.

10.3.1   Baseline normalisation

Raw fluorescence matrices were first transposed to the convention (cells × time). Each cell's resting baseline F_0 was defined as the minimum fluorescence value observed during the first 4,000 time samples, corresponding to an epoch prior to any stimulus-evoked activity. The full trace was then divided by F_0 to produce a dimensionless ratio:

$$\frac{F(t)}{F_0} = \frac{F(t)}{\min_{t \leq 4000} F(t)}, \tag{1}$$

anchoring each cell's signal to its own pre-stimulus baseline and correcting for inter-cell differences in absolute fluorescence intensity.

10.3.2   Exponential drift correction

Slow multiplicative drift in the normalised signal — a characteristic artefact of photobleaching in calcium imaging — was removed by estimating and subtracting a shared exponential trend. A single linear regression was performed on the log-transformed F/F_0 values pooled across all cells, yielding a global slope $\hat{m}$ and intercept $\hat{b}$ that characterise the dominant exponential decay:

$$\log\left[\frac{F(t)}{F_0}\right] \approx \hat{m}\,t + \hat{b}. \tag{2}$$

The fitted exponential $e^{\hat{m}\,t + \hat{b}}$ was then subtracted from each individual trace, producing drift-corrected signals with a stable baseline. Fitting a single trend to the pooled data ensures that the correction captures the component of variance shared across the population (i.e. global photobleaching) while preserving cell-specific dynamics.

10.3.3 Low-pass filtering

Following drift correction, residual high-frequency noise was suppressed by applying a zero-phase Bessel low-pass filter (order 2, normalised cutoff frequency $f_c = 0.3$) along the time axis using forward-backward filtering (`scipy.signal.sosfiltfilt`). The Bessel design was chosen for its maximally flat group delay, which preserves the temporal waveform of calcium transients without introducing phase distortion [49].

10.4 Aperiodic Spectral Analysis

The aperiodic ($1/f$-like) component of the power spectrum was characterised for each cell by fitting a Lorentzian model to the single-sided power spectrum and extracting the aperiodic exponent. Cross-cell variability in log-power space was then summarised via frequency-resolved variance and mean profiles.

10.4.1 Power spectrum estimation

For each cell c, the single-sided power spectrum was obtained from the discrete Fourier transform of the fluorescence trace $x_c(t)$ ($t = 1, \ldots, N_{\text{samples}}$):

$$P_c(f) = \left| \mathcal{F}\{x_c\}(f) \right|^2, \tag{3}$$

where $\mathcal{F}\{\cdot\}$ denotes the real-valued FFT and f ranges over the positive Nyquist grid with resolution $\Delta f = f_s / N_{\text{samples}}$. Frequency bins at or below a threshold $f_{\min} = 0.5\,\text{Hz}$ were discarded to exclude slow drift and DC components, yielding the trimmed spectrum $\{P_c(f)\}_{f > f_{\min}}$.

10.4.2 Lorentzian model of the aperiodic component

The aperiodic background of neural power spectra follows a $1/f^{\alpha}$ power law that, in log-space, is well described by a Lorentzian function [40]:

$$\log P(f) = \beta - \log(k + f^{\alpha}), \tag{4}$$

where β is a log-power intercept, $\alpha > 0$ is the aperiodic exponent governing the spectral slope, and $k \geq 0$ is a knee parameter that captures any low-frequency roll-off. In the limit $k \to 0$ and for large f, Eq. (4) reduces to the pure power law $\log P(f) \approx \beta - \alpha \log f$.

10.4.3 Aperiodic exponent estimation

The three parameters (β, α, k) were estimated for each cell by non-linear least squares minimisation of the residual sum of squares between the observed log-power and the Lorentzian model (Eq. (4)), using the Levenberg–Marquardt algorithm as implemented in `scipy.optimize.curve_fit` [49]. The aperiodic exponent was then defined as the absolute value of the fitted slope:

$$\hat{\alpha}_c = \left| \alpha_c^* \right|, \tag{5}$$

where α_c^* denotes the least-squares estimate for cell c. Taking the absolute value ensures that $\hat{\alpha}_c \geq 0$ regardless of the sign convention returned by the optimiser, with larger values indicating steeper (more coloured) spectra.

10.4.4 Cross-cell log-power variance

To quantify frequency-resolved heterogeneity across the cell population, the power spectra were transformed to $\log_{10}$-space with a small regularisation constant $\varepsilon = 10^{-10}$ added to prevent numerical instability at bins with near-zero power:

$$\tilde{P}_c(f) = \log_{10}(P_c(f) + \varepsilon). \tag{6}$$

The cross-cell mean and variance were then computed at each frequency bin:

$$\bar{P}(f) = \frac{1}{n_{\text{cells}}} \sum_{c=1}^{n_{\text{cells}}} \tilde{P}_c(f), \qquad \sigma_P^2(f) = \frac{1}{n_{\text{cells}}} \sum_{c=1}^{n_{\text{cells}}} \left[\tilde{P}_c(f) - \bar{P}(f) \right]^2. \tag{7}$$

Frequency bins where $\sigma_P^2(f)$ is large indicate spectral frequencies at which cells are most heterogeneous in their power, providing a data-driven criterion for identifying frequency bands of interest for subsequent analyses.

10.5 Phase–Amplitude Coupling Analysis

Phase–amplitude coupling (PAC) quantifies the degree to which the amplitude envelope of a high-frequency oscillation is modulated by the phase of a lower-frequency oscillation within the same signal. We computed PAC using the Modulation Index (MI) of [50], applied cell-by-cell to background-subtracted, exponentially normalised fluorescence traces ($\Delta F/F$, n_{cells} cells).

10.5.1 Frequency decomposition

For each candidate phase frequency f_ϕ and amplitude frequency f_A, two narrow-band signals were obtained by applying a third-order Butterworth bandpass filter with passbands $[f_\phi, f_\phi + \Delta f_\phi]$ and $[f_A, f_A + \Delta f_A]$, respectively, where $\Delta f_\phi = 0.01\,\text{Hz}$ and $\Delta f_A = 0.10\,\text{Hz}$. The instantaneous phase $\phi(t)$ and amplitude envelope $A(t)$ were then extracted via the analytic signal obtained from the Hilbert transform:

$$\phi(t) = \arg\big[\mathcal{H}\{x_\phi(t)\}\big], \qquad A(t) = \big|\mathcal{H}\{x_A(t)\}\big|, \tag{8}$$

where x_ϕ and x_A denote the phase- and amplitude-filtered signals, respectively, and $\mathcal{H}\{\cdot\}$ is the Hilbert transform.

10.5.2 Modulation Index

The full phase cycle $[-\pi, \pi)$ was partitioned into $N = 18$ equal bins. For each bin k the mean amplitude was computed as

$$\bar{A}_k = \frac{1}{|\mathcal{I}_k|} \sum_{t \in \mathcal{I}_k} A(t), \qquad \mathcal{I}_k = \big\{t : \phi(t) \in [\theta_k, \theta_{k+1})\big\}, \tag{9}$$

yielding a discrete amplitude-by-phase distribution $\{\bar{A}_k\}_{k=1}^N$. This distribution was normalised to a probability mass function

$$p_k = \frac{\bar{A}_k}{\sum_{j=1}^N \bar{A}_j}, \tag{10}$$

and its Shannon entropy was compared to the maximum entropy of a uniform distribution to obtain the MI:

$$\text{MI}(f_\phi, f_A) = \frac{\ln N - H(p)}{\ln N}, \qquad H(p) = -\sum_{k=1}^N p_k \ln p_k. \tag{11}$$

MI $= 0$ indicates a uniform amplitude distribution across phases (no coupling), whereas MI $= 1$ indicates complete concentration of amplitude at a single phase bin. A small regularisation constant ($\epsilon = 10^{-12}$) was added inside the logarithm to avoid numerical instability.

10.5.3 Comodulogram

PAC was evaluated across a two-dimensional frequency grid. The phase frequency axis spanned $f_\phi \in [0.005, 0.045]\,\text{Hz}$ in steps of $0.005\,\text{Hz}$; the amplitude frequency axis spanned $f_A \in [0.05, 0.75]\,\text{Hz}$ in steps of $0.05\,\text{Hz}$, yielding a 9×15 comodulogram per cell. The population-level comodulogram was obtained by averaging MI values across all cells:

$$\overline{\text{MI}}(f_\phi, f_A) = \frac{1}{n_{\text{cells}}} \sum_{c=1}^{n_{\text{cells}}} \text{MI}_c(f_\phi, f_A). \tag{12}$$

10.5.4 Marginal profiles

To characterise how spectral power varies across cells as a joint function of frequency ff f and time tt t, principal component analysis (PCA) was applied to the per-cell spectrograms computed as described in Section

$$\overline{\text{MI}}_\phi(f_\phi) = \frac{1}{n_{\text{cells}}} \sum_{c=1}^{n_{\text{cells}}} \Big[\frac{1}{|F_A|} \sum_{f_A} \text{MI}_c(f_\phi, f_A) \Big]. \tag{13}$$

The dominant amplitude-frequency profile $\overline{\mathrm{MI}}_A(f_A)$ was computed analogously by collapsing the phase axis. Error bars in the corresponding figures represent $\pm 1\,\mathrm{SEM}$ across cells.

10.6   Spectral Signal Classification

To characterise how spectral power varies across cells as a joint function of frequency f and time t, principal component analysis (PCA) was applied to the per-cell spectrograms computed as described before.

10.6.1   Spectrogram covariance and PCA

For each cell i, the spectrogram $S_i(f,t)$ was evaluated over $F = 60$ logarithmically spaced frequency bins spanning 0.005–1 Hz and T non-overlapping time windows. The cross-cell covariance matrix was constructed over the joint (frequency $\times$ time) space as:

$$C_{ij} = \frac{1}{T \cdot F} \sum_{t=1}^{T} \sum_{f=1}^{F} \Big[S_i(f,t) - \bar{S}_i(f) \Big] \Big[S_j(f,t) - \bar{S}_j(f) \Big], \tag{14}$$

where $\bar{S}_i(f) = T^{-1} \sum_{t=1}^{T} S_i(f,t)$ is the time-averaged spectrum of cell i at frequency f, and indices i,j run over all recorded cells. The maximum number of significant dimensions is bounded by $F = 60$, corresponding to the number of frequency bins.

The leading principal components — those capturing the largest fraction of total variance — were retained, and each cell's spectrogram was projected onto these components to obtain a low-dimensional spectral embedding.

10.6.2   Unsupervised clustering

Cells were grouped according to their spectral covariance structure by applying k-means clustering to the low-dimensional PCA projections, as implemented in `scikit-learn` [51]. The number of clusters was set to $k = 2$, with $n_{\mathrm{init}} = 10$ independent initialisations to mitigate sensitivity to centroid initialisation; the solution minimising the within-cluster sum of squares was selected. The cluster bandwidth was estimated directly from the data prior to fitting.

10.7   Aperiodic Signal Decomposition and Cluster-Level Analysis

To examine the relationship between spectral cluster membership and aperiodic signal properties at single-cell resolution, a cell-by-cell adjacency matrix was constructed for each combination of cell population (heterotypic and homotypic) and physiological condition (virgin, lactant, weaned, and multiparous). Spectral class indices and per-cell aperiodic exponents were merged at the level of individual cells using the recording index and cell identifier as joint keys; only cells present in both datasets were retained for subsequent analysis.

10.7.1   Cluster size balancing

To prevent asymmetries in matrix structure driven by cluster size imbalance, clusters were downsampled to a common size prior to matrix construction. The minimum cluster size across all clusters within a given condition was determined, and any larger cluster was randomly subsampled without replacement to match this reference size (random seed fixed at 42 for reproducibility). A condition was included in the analysis only if both clusters contained at least one cell after merging.

10.7.2   Aperiodic adjacency matrix

For each condition, an $N \times N$ symmetric adjacency matrix $\mathbf{A}$ was constructed, where N is the total number of retained cells across both clusters. Diagonal entries encode each cell's intrinsic aperiodic exponent:

$$A_{ii} = \hat{\alpha}_i, \tag{15}$$

where $\hat{\alpha}_i$ is the aperiodic exponent of cell i estimated as described before. Off-diagonal entries represent a pairwise aperiodic similarity measure defined as the arithmetic mean of the exponents of cells i and j:

$$A_{ij} = \frac{\hat{\alpha}_i + \hat{\alpha}_j}{2}, \qquad i \neq j. \tag{16}$$

All matrix entries were subsequently z-score normalised across the full set of elements to centre the distribution and express values in units of standard deviation from the population mean:

$$\tilde{A}_{ij} = \frac{A_{ij} - \mu_A}{\sigma_A}, \tag{17}$$

where μ_A and σ_A are the mean and standard deviation computed over all entries of $\mathbf{A}$. The lower triangular portion of $\tilde{\mathbf{A}}$ was visualised as a heatmap using a diverging colormap centred at zero, with rows and columns ordered by cluster identity to allow visual inspection of within- versus between-cluster aperiodic structure. Cluster boundaries were annotated along both axes.

10.7.3 Bootstrap Estimation of Within-Cluster Aperiodic Reliability

To assess the reliability of the mean aperiodic exponent within each spectral cluster, a non-parametric bootstrap procedure. For each combination of cell population, physiological condition, and cluster identity, the observed aperiodic values $\{\hat{\alpha}_c\}$ were resampled with replacement over $B = 1{,}000$ iterations. The mean aperiodic exponent was computed for each bootstrap replicate b:

$$\bar{\alpha}^{(b)} = \frac{1}{n} \sum_{c=1}^{n} \hat{\alpha}_c^{(b)}, \qquad b = 1, \ldots, B, \tag{18}$$

where n is the number of cells in the cluster and $\hat{\alpha}_c^{(b)}$ denotes the c-th draw of the b-th replicate. This procedure yields an empirical sampling distribution of the mean from which the 95% confidence interval was derived as the interval between the 2.5th and 97.5th percentiles:

$$\mathrm{CI}_{95} = \left[\hat{\alpha}^{(0.025)}, \ \hat{\alpha}^{(0.975)} \right], \tag{19}$$

where $\hat{\alpha}^{(q)}$ denotes the q-th quantile of the bootstrap distribution. Results were aggregated across all populations, conditions, and clusters; the observed mean together with its bootstrap confidence interval was retained as the summary statistic for each group.

10.8 Contingency Analysis of Aperiodic Exponent and Shannon Entropy

To examine the joint distribution of aperiodic signal properties and spectral entropy across cluster identities, both continuous measures were discretised and analysed via cross-tabulation at three levels of granularity.

10.8.1 Tertile discretisation

The aperiodic exponent $\hat{\alpha}$ and the Shannon entropy H were each independently binned into population tertiles, partitioning the observed range into three ordered categories — Low, Medium, and High — each containing approximately one-third of the observations:

$$\tilde{x}_i = \begin{cases} \text{Low} & x_i \leq Q_{33}(x), \\ \text{Medium} & Q_{33}(x) < x_i \leq Q_{67}(x), \\ \text{High} & x_i > Q_{67}(x), \end{cases} \tag{20}$$

where Q_{33} and Q_{67} denote the 33rd and 67th sample percentiles, respectively. Tertile boundaries were computed independently for $\hat{\alpha}$ and H, so no assumption of a shared distribution is imposed.

10.8.2 Cross-tabulation

Contingency tables were constructed at three levels of aggregation:

1. **Pooled**: all cell populations and physiological conditions combined, providing a global view of the $\hat{\alpha}$–H–cluster relationship.

2. **Population-level**: a single cell population (e.g. lactotrophs or somatotrophs) with all conditions pooled.

3. **Group-level**: a single cell population within a single physiological condition (e.g. lactotrophs during lactation).

At each level, three tables were derived: (i) the marginal cross-tabulation of $\tilde{\alpha}$ against cluster identity, (ii) the marginal cross-tabulation of $\tilde{H}$ against cluster identity, and (iii) the joint cross-tabulation of $(\tilde{\alpha}, \tilde{H})$ against cluster identity.

10.8.3 Percentage representation

The joint contingency table was further expressed as a percentage of the total number of cells N in the corresponding subset:

$$P_{(a,h),k} = \frac{n_{(a,h),k}}{N} \times 100, \tag{21}$$

where $n_{(a,h),k}$ is the count of cells falling in tertile category ($\tilde{\alpha} = a$, $\tilde{H} = h$) and belonging to cluster k, and $N = \sum_{a,h,k} n_{(a,h),k}$ is the total cell count in the subset. This normalisation allows direct comparison of cell proportions across groups of different sizes and was used as the primary input for figure generation.

10.8.4 Comparison of compositional distributions across conditions

To assess whether the joint aperiodic–entropy cell-state composition differed across physiological conditions, a multivariate permutation test was applied separately for each oscillatory class (Class 1 and Class 2) and each cell population (homotypic and heterotypic). For each recording r, the percentage values $P_{(a,h),k}$ were collapsed across entropy position by summing within each aperiodic tertile, yielding a three-component compositional vector $\mathbf{v}_r = [p_{\mathrm{Low}}, p_{\mathrm{Medium}}, p_{\mathrm{High}}]_r$ that sums to 100% per recording. This compositional structure — whereby the three components are mutually exclusive and exhaustive — precluded the use of standard univariate tests applied independently per bin, as such an approach would violate the sum constraint and introduce spurious correlations between test statistics.

A pseudo-F statistic was used as the test statistic, defined as the ratio of between-group to within-group dispersion in compositional space:

$$F_{\mathrm{pseudo}} = \frac{\sum_c n_c \, \|\bar{\mathbf{v}}_c - \bar{\mathbf{v}}\|^2 \, / \, (C-1)}{\sum_c \sum_{r \in c} \|\mathbf{v}_r - \bar{\mathbf{v}}_c\|^2 \, / \, (N-C)}, \tag{22}$$

where C is the number of conditions, n_c is the number of recordings in condition c, $\bar{\mathbf{v}}_c$ is the centroid of condition c in compositional space, $\bar{\mathbf{v}}$ is the grand centroid, and N is the total number of recordings. The null distribution of F_{pseudo} was obtained by randomly permuting condition labels across recordings and recomputing the statistic over $n_{\mathrm{perm}} = 10{,}000$ iterations. The permutation p-value was defined as the proportion of permuted statistics exceeding or equal to the observed value.

Following the omnibus test, pairwise permutation tests were conducted for all condition pairs using the same pseudo-F statistic restricted to the two groups under comparison. Resulting p-values were adjusted for multiple comparisons using the Bonferroni method ($m = 6$ pairs for four conditions).

10.9 Surrogate-Thresholded Pairwise Correlation Analysis

To identify statistically significant pairwise interactions among cells, a Fourier phase-randomisation surrogate procedure was applied to define a cell-pair-specific significance threshold for the observed Spearman correlation matrix.

10.9.1 Fourier surrogate generation

For each cell i, the discrete Fourier transform of the fluorescence trace was computed and decomposed into its amplitude and phase spectra:

$$\mathcal{F}\{x_i\}(k) = A_i(k) \, e^{i\phi_i(k)}, \tag{23}$$

where $A_i(k) = |\mathcal{F}\{x_i\}(k)|$ and $\phi_i(k) = \arg[\mathcal{F}\{x_i\}(k)]$. For each of $B = 1{,}000$ surrogate resamples, the phase spectrum of every cell was independently replaced by phases $\tilde{\phi}_i(k)$ drawn uniformly from $[-\pi, \pi)$, while the amplitude spectrum was held fixed. Conjugate symmetry was enforced to ensure a real-valued inverse transform:

$$\tilde{\phi}_i(N-k) = -\tilde{\phi}_i(k), \qquad \tilde{\phi}_i(N/2) = 0, \tag{24}$$

where N denotes the number of time points. The surrogate signal was then obtained as

$$\tilde{x}_i = \text{Re}\Big[\mathcal{F}^{-1}\Big\{A_i(k)\,e^{i\tilde{\phi}_i(k)}\Big\}\Big].\tag{25}$$

This procedure preserves the power spectrum of each individual cell while destroying inter-cell phase relationships, thereby generating a null distribution of pairwise correlations under the hypothesis of no genuine synchrony.

10.9.2 Significance threshold and matrix thresholding

For each cell pair (i, j), the element-wise mean μ_{ij} and standard deviation σ_{ij} were computed across the $B = 1{,}000$ surrogate Spearman correlation matrices. An observed correlation ρ_{ij} was deemed significant only if it exceeded the surrogate threshold:

$$|\rho_{ij}| > \mu_{ij} + 2\,\sigma_{ij}.\tag{26}$$

Correlations that did not satisfy this criterion were set to zero, yielding a sparse, thresholded correlation matrix $\hat{\rho}$ that retains only interactions whose magnitude is unlikely under the phase-randomisation null.

10.9.3 Interaction classification

Non-zero entries of $\hat{\rho}$ were partitioned into three interaction classes based on their magnitude and sign relative to the surrogate distribution:

1. **Strongly synchronous**: $\hat{\rho}_{ij} > \mu_{ij} + 2\,\sigma_{ij}$.

2. **Negatively asynchronous**: $\hat{\rho}_{ij} < -\mu_{ij}$ and $\hat{\rho}_{ij} \neq 0$.

3. **Weakly asynchronous**: $|\hat{\rho}_{ij}| < \mu_{ij}$ and $\hat{\rho}_{ij} \neq 0$.

The proportion of positive and negative interactions among all surviving (non-zero) pairs was computed, and the distribution of values within each class was visualised as a boxplot.

10.10 Cluster–Synchrony Association via Random Forest Classification

To assess whether spectral cluster identity is predictive of pairwise synchrony class, a supervised classification analysis was performed. Spectral cluster labels and synchrony labels were merged at the cell level using the recording index, cell identifier, and physiological condition as joint keys; only cells present in both tables were retained.

Synchrony labels were encoded numerically (synchronous = 1, asynchronous = 2). Cluster identity was used as the sole predictor feature and synchrony class as the target variable. A Random Forest classifier (implemented in `scikit-learn` [51] ; random seed fixed at 42) was evaluated by stratified 5-fold cross-validation, with classification accuracy as the performance metric. The mean and standard deviation of accuracy across folds are reported as the summary statistic. A confusion matrix was constructed to visualise the correspondence between cluster identity and synchrony class for each population–condition combination.

10.11 Transfer entropy along electrode clusters

This methodology section was inspired on the previous reported in Aquiles 2025 [23], but further described here. We computed the transfer entropy from baseline in every cluster, ie X_t = signal from cellclass$n$ and Y_t = signal from cellclass m; using a finite set of lags (100, (-50,50))) $L = \{\tau_1, \tau_2, \ldots, \tau_k\}$. For each lag, we define the lagged version of the time series:

$$X_t(\tau) = X_{t+\tau}, \quad Y_t(\tau) = Y_{t+\tau}$$

Then we calculate the Mutual Information (MI) between $X_t(\tau)$ and Y_t as:

$$\text{MI}_{xy}(\tau) = I(X_t(\tau); Y_t), \quad \text{MI}_{yx}(\tau) = I(Y_t(\tau); X_t)$$

Where I corresponds to Shannon mutual information, and H to Shannon entropy:

$$I(A; B) = H(A) + H(B) - H(A, B)$$

Where A and B represent the different cluster sets. To compute directionality, we took the vector $\text{MI}_{xy}(\tau)$ for all $\tau \in L$, where we consider the best delay as the one that maximizes the mutual information:

$$XY = \arg\max \text{MI}_{xy}(\tau)$$

And we follow the same logic in a similar way for the other direction YX.

10.11.1 Multi-Set Directionality Computation

The directionality analysis is extended to multiple signal sets by evaluating all pairwise interactions among them. Each pair of sets is analyzed independently following the same time-lagged MI and permutation-testing procedure, see below.

10.11.2 Directionality Asymmetry Index

To summarize net directional influence between two signal sets, the Directionality Asymmetry Index (DAI) is computed. Total significant information transfer is summed separately for each direction, and the DAI is defined as

$$DAI = \frac{TE_{X \to Y} - TE_{Y \to X}}{TE_{X \to Y} + TE_{Y \to X}}, \tag{27}$$

where $TE_{X \to Y}$ and $TE_{Y \to X}$ denote the total information transfer in each direction. The DAI ranges from -1 to 1, indicating dominant directional influence or balanced interactions.

10.11.3 Permutation-Based Significance Testing of Mutual Information

To determine whether the observed mutual information (MI) at a given lag exceeded chance levels, a permutation null distribution was constructed independently for each lag τ. Specifically, $N = 1{,}000$ surrogate time series $Y_t^{(n)}$ were generated by randomly shuffling the original Y_t (and equivalently X_t for the reverse direction). The MI was recomputed for each surrogate, yielding a null distribution. Only lag values for which the observed MI exceeded the 95^{th} percentile of the null distribution ($p < 0.05$, one-tailed) were retained for subsequent directionality analyses.

10.12 Geometric Synchrony Analysis Between Neural Populations

To quantify the degree to which two pituitary cell populations (PopA and PopB) share geometric structure in their collective dynamics, a manifold-based synchrony analysis was performed. The pipeline comprises five sequential steps: time-delay embedding, dimensionality reduction, Procrustes alignment, and two complementary synchrony metrics evaluated at global and local temporal scales.

10.12.1 Time-delay embedding

Each cell's fluorescence trace $x_i(t)$ was embedded into a pseudo-phase-space attractor using the Takens time-delay construction:

$$\mathbf{v}_i(t) = \big[x_i(t),\ x_i(t + \tau),\ \ldots,\ x_i(t + (m - 1)\tau)\big] \in \mathbb{R}^m, \tag{28}$$

where $m = 2$ is the embedding dimension and $\tau = 1$ sample is the time delay. The embedded trajectories of all cells within a population were concatenated horizontally, yielding a population-level state matrix $\mathbf{X} \in \mathbb{R}^{T \times (n_{\text{cells}} \cdot m)}$, where T is the number of embedded time points.

10.12.2 Dimensionality reduction via PCA

Principal component analysis was applied independently to the state matrix of each population, retaining up to $k = \min(10,\ n_{\text{cells}} \cdot m)$ components:

$$\mathbf{Z}_A = \mathbf{X}_A \mathbf{W}_A, \qquad \mathbf{Z}_B = \mathbf{X}_B \mathbf{W}_B, \tag{29}$$

where $\mathbf{W}_A$ and $\mathbf{W}_B$ are the matrices of leading eigenvectors of the respective sample covariance matrices. This yields low-dimensional latent trajectories $\mathbf{Z}_A, \mathbf{Z}_B \in \mathbb{R}^{T \times k}$ that capture the dominant variance structure of each population in its own coordinate frame.

10.12.3   Procrustes alignment

Because the PCA coordinate frames of PopA and PopB are independently defined, a direct geometric comparison requires a common reference frame. An optimal orthogonal rotation $\mathbf{R}$ was found by solving the orthogonal Procrustes problem:

$$\mathbf{R}^* = \arg \min_{\mathbf{R} \in \mathcal{O}(k)} \|\mathbf{Z}_A - \mathbf{Z}_B \mathbf{R}\|_F, \tag{30}$$

where $\mathcal{O}(k)$ denotes the group of $k \times k$ orthogonal matrices and $\| \cdot \|_F$ is the Frobenius norm. The solution is given by $\mathbf{R}^* = \mathbf{V}\mathbf{U}^\top$, where $\mathbf{U}\mathbf{\Sigma}\mathbf{V}^\top$ is the singular value decomposition of $\mathbf{Z}_B^\top \mathbf{Z}_A$. The aligned trajectory $\mathbf{Z}_B^* = \mathbf{Z}_B \mathbf{R}^*$ then lives in the same coordinate frame as $\mathbf{Z}_A$, enabling direct pointwise comparison.

10.12.4   Geometric synchrony metrics

Pointwise trajectory distance.   The instantaneous divergence between the two population states was quantified by the Euclidean distance between their aligned latent representations at each time point:

$$\Delta(t) = \|\mathbf{Z}_A(t) - \mathbf{Z}_B^*(t)\|_2. \tag{31}$$

Time points where $\Delta(t)$ falls below its temporal mean are interpreted as epochs of high geometric synchrony; time points above the mean indicate divergent (asynchronous) population states.

Global principal subspace angles.   To characterise the overall geometric relationship between the two population subspaces, the principal angles $\theta_1 \leq \cdots \leq \theta_k$ between the column spaces of $\mathbf{Z}_A$ and $\mathbf{Z}_B^*$ were computed via QR decomposition followed by SVD of the cross-product of the orthonormal bases:

$$\cos \theta_j = \sigma_j \big(\mathbf{Q}_A^\top \mathbf{Q}_B \big), \qquad j = 1, \dots, k, \tag{32}$$

where $\mathbf{Q}_A$ and $\mathbf{Q}_B$ are the orthonormal factors from the QR decompositions of $\mathbf{Z}_A$ and $\mathbf{Z}_B^*$, respectively, and $\sigma_j(\cdot)$ denotes the j-th singular value. Angles below 45° indicate shared geometric structure (synchrony); angles exceeding 45° indicate geometrically divergent subspaces (dominance).

Sliding-window subspace alignment score.   To capture the temporal evolution of geometric synchrony, a sliding-window variant of the principal angle analysis was applied. At each time step t, local segments of half-width $w = 20$ samples were extracted from both trajectories, and the mean cosine of the local principal angles was computed:

$$S(t) = \frac{1}{k} \sum_{j=1}^{k} \cos \theta_j \left(\mathbf{Z}_A[t-w:t+w], \, \mathbf{Z}_B^*[t-w:t+w] \right), \qquad w < t < T - w. \tag{33}$$

$S(t)$ ranges from 0 (orthogonal, maximal divergence) to 1 (identical subspaces, perfect alignment). Values above the temporal mean of S are classified as high-alignment (synchronous) epochs; values below the mean as low-alignment (dominant) epochs.

10.12.5   Temporal density estimation of synchrony epochs

To identify at which periods of the recording synchrony and dominance preferentially occur, a kernel density estimate (KDE) with a Gaussian kernel (bandwidth $h = 0.2$, Scott's rule) was fitted separately to the set of time indices classified as high-alignment (top quartile of $S(t)$) and low-alignment (bottom quartile of $S(t)$). The resulting density curves provide a continuous representation of the temporal concentration of each synchrony regime across the recording.

10.13   One-Dimensional Quasi-Potential Landscape and Dominance Angle

To quantify the stability and relative dominance of two coexisting population states, the quasi-potential energy landscape was estimated from the empirical signal distribution of each cell group and physiological condition. This analysis operates on a one-dimensional projection of the population signal — complementing the two-dimensional latent-space landscape described before — and provides an explicit scalar measure of bistability and directional dominance.

10.13.1 Quasi-potential estimation

For each scalar signal distribution $\{x_i\}_{i=1}^N$, the probability density $P(x)$ was estimated by a Gaussian kernel density estimator (KDE):

$$\hat{P}(x) = \frac{1}{Nh}\sum_{i=1}^N K\left(\frac{x-x_i}{h}\right), \qquad h = 0.9\,\hat{\sigma}\,N^{-1/5}, \tag{34}$$

where K is the standard Gaussian kernel and $\hat{\sigma}$ is the sample standard deviation. The quasi-potential was then defined as the negative log-density, regularised to avoid numerical instability at grid points with near-zero probability:

$$U(x) = -\log[\hat{P}(x) + \varepsilon], \qquad \varepsilon = 10^{-10}, \tag{35}$$

evaluated on a uniform grid of 500 points spanning the observed signal range. The landscape was shifted so that its global minimum equals zero: $U \leftarrow U - \min(U)$. Minima of U correspond to preferred (high-probability) population states; the value of U at each minimum is proportional to the negative log-probability of that state.

10.13.2 Well and barrier identification

Local minima and maxima of U were identified using a neighbourhood-based extremum detector with order parameter $q = 15$ grid points (`scipy.signal.argrelmin` and `argrelmax`). A bistable landscape was defined as one containing at least two local minima separated by at least one local maximum. When multiple minima were present, the two deepest (lowest U) were retained as the primary wells; the highest local maximum strictly between these two wells was taken as the energy barrier.

Denoting the left and right wells as A and B:

$$\Delta U_{A\to B} = U_{\mathrm{bar}} - U_A, \tag{36}$$
$$\Delta U_{B\to A} = U_{\mathrm{bar}} - U_B, \tag{37}$$

where U_{bar} is the barrier height. The mean escape time from well P is proportional to $e^{\Delta U_{P\to P'}}$, so a larger barrier corresponds to a longer-lived (more stable) state.

10.13.3 Dominance angle

The relative stability of the two wells was summarised by the dominance angle θ, defined as the signed inclination of the line connecting well A to well B in the (x, U) plane:

$$\theta = \arctan\left(\frac{\Delta U}{\Delta x}\right), \qquad \Delta U = U_B - U_A, \quad \Delta x = x_B - x_A > 0. \tag{38}$$

Because well A is always to the left of well B by construction ($\Delta x > 0$), the sign of θ is determined entirely by ΔU:

- $\theta > 0$: well B is higher than well A; population A occupies the deeper well and is the dominant state.

- $\theta < 0$: well A is higher than well B; population B dominates.

- $\theta \approx 0$: symmetric landscape; neither population is strongly preferred.

The dominant population is formally defined as the one occupying the well with the lower U value (higher probability density). The magnitude $|\theta|$ quantifies the degree of asymmetry: larger angles correspond to stronger dominance and greater stability of the preferred state.

The dominance angle analysis was applied to three signal distributions per recording: the pooled signal across both populations, and the signals of each group separately. Results were aggregated across all physiological conditions (virgin, lactant, weaned, multiparous) and all recordings, yielding a condition-level summary table of θ, ΔU, and the dominant population for each group.

10.14 Bistability Analysis: Mixture Model Selection and State Dwell Times

To characterise the degree of bistability in pituitary endocrine cell populations, the empirical distribution of fluorescence signals was analysed using Gaussian Mixture Model (GMM) selection and per-cell dwell-time statistics. All analyses were applied per recording (sample) and condition, with results aggregated across samples for reporting.

10.14.1 Signal pooling and quasi-potential estimation

For each recording, fluorescence traces were flattened across cells and time to yield three one-dimensional signal distributions: (i) all cells pooled (X_{all}), (ii) Class 1 cells only (X_1), and (iii) Class 2 cells only (X_2). The quasi-potential landscape of each distribution was estimated as described in Section **??**, Eq. (35), using a Gaussian KDE with Silverman's bandwidth on a uniform grid of 500 points spanning the observed signal range. For multi-sample conditions, individual-sample landscapes were interpolated onto a common grid and averaged to obtain a condition-level mean landscape with pointwise standard deviation shown as a shaded band.

10.14.2 Gaussian Mixture Model selection

To test whether the pooled signal distribution X_{all} was consistent with a bimodal (bistable) regime, Gaussian Mixture Models with $n \in \{1, 2, 3\}$ components were fitted by Expectation–Maximisation. Each model was initialised $n_{\text{init}} = 10$ times with random starting parameters (random seed fixed at 42) and the initialisation yielding the highest log-likelihood was retained. Model complexity was penalised using the Bayesian Information Criterion:

$$\text{BIC}(n) = k_n \ln N - 2\,\hat{\ell}_n, \tag{39}$$

where k_n is the number of free parameters for the n-component model, N is the total number of observations, and $\hat{\ell}_n$ is the maximised log-likelihood. The model minimising Eq. (39) was selected as the best fit; a best-fit component count of $n^* = 2$ was taken as evidence of bistability in the signal distribution.

10.14.3 State dwell-time analysis

To characterise the temporal persistence of each oscillatory class (state), per-cell dwell times were extracted from the group-identity label sequence. Each cell was assigned a binary state label at every time point: state 0 for Class 1 cells and state 1 for Class 2 cells. Run-length encoding was applied to the label sequence of each cell to obtain a list of consecutive residence durations in each state. Dwell times were pooled across all cells and recordings within a condition for reporting.

The distribution of dwell times τ in each state was characterised by its mean, median, and maximum, and was compared against an exponential null model. An exponential distribution with rate parameter λ was fitted to the pooled dwell times by maximum likelihood estimation with location fixed at zero:

$$f(\tau; \lambda) = \lambda\, e^{-\lambda \tau}, \qquad \tau \geq 0. \tag{40}$$

Exponentially distributed dwell times are consistent with thermally activated escape from a metastable potential well. Deviations from exponential behaviour — in particular heavy-tailed or multi-modal dwell-time distributions — would indicate more complex transition dynamics. Dwell-time histograms were plotted on a log-density scale to facilitate visual discrimination of exponential from non-exponential tails.

10.15 Low-Rank Recurrent Neural Network Model

To validate the geometric synchrony pipeline and provide a mechanistic interpretation of directed coupling between pituitary cell populations, a rank-2 recurrent neural network (RNN) was simulated under three coupling conditions: directed coupling from population A to B ($g_{AB} > 0$, $g_{BA} = 0$), null coupling ($g_{AB} = g_{BA} = 0$), and reversed coupling from B to A ($g_{AB} = 0$, $g_{BA} > 0$).

10.15.1 Network architecture and dynamics

Each population $P \in \{A, B\}$ consists of N_P leaky-integrator units governed by:

$$\tau \frac{d\mathbf{x}_P}{dt} = -\mathbf{x}_P + \mathbf{W}_P\, \mathbf{r}_P + g_{P'P}\, \mathbf{C}_{P'P}\, \mathbf{r}_{P'} + \boldsymbol{\eta}_P(t), \tag{41}$$

where $\mathbf{x}_P \in \mathbb{R}^{N_P}$ is the state vector, $\mathbf{r}_P = \tanh(\mathbf{x}_P)$ is the firing-rate output, $\mathbf{W}_P \in \mathbb{R}^{N_P \times N_P}$ is the within-population recurrent weight matrix, $\mathbf{C}_{P'P} \in \mathbb{R}^{N_P \times N_{P'}}$ is the rank-1 cross-population coupling matrix, $g_{P'P}$ is the coupling gain, and $\boldsymbol{\eta}_P(t) \sim \mathcal{N}(\mathbf{0}, \sigma^2 \Delta t\, \mathbf{I})$ is additive Gaussian noise. Equation (41) was integrated with the Euler method using time step $\Delta t = 0.5$ and membrane time constant $\tau = 5.0$, giving an effective per-step update factor $\Delta t/\tau = 0.1$.

10.15.2   Low-rank connectivity

The recurrent weight matrix of each population was constructed as a rank-r outer product:

$$\mathbf{W}_P = \frac{1}{\sqrt{N_P}}\mathbf{U}_P\mathbf{V}_P, \qquad \mathbf{U}_P \in \mathbb{R}^{N_P \times r},\ \mathbf{V}_P \in \mathbb{R}^{r \times N_P}, \tag{42}$$

with $r = 2$. The matrix was then rescaled so that its dominant eigenvalue equals the target spectral radius $\rho = 1.15$, placing the network in the oscillatory (weakly chaotic) regime:

$$\mathbf{W}_P \leftarrow \mathbf{W}_P \cdot \frac{\rho}{\max_i |\lambda_i(\mathbf{W}_P)|}. \tag{43}$$

The cross-population coupling matrices were drawn as rank-1 random matrices, $\mathbf{C}_{AB} = \boldsymbol{\xi}\boldsymbol{\zeta}^\top/N_A$ with $\boldsymbol{\xi} \in \mathbb{R}^{N_B}$, $\boldsymbol{\zeta} \in \mathbb{R}^{N_A}$ independently sampled from $\mathcal{N}(0,1)$. Both populations used a shared random seed for $\mathbf{U}$, $\mathbf{V}$, and $\mathbf{C}$, ensuring comparable geometric structure across conditions while the coupling gains g_{AB} and g_{BA} alone broke the symmetry.

10.15.3   Simulation parameters

Populations contained $N_A = 50$ and $N_B = 80$ units. Each simulation ran for $T = 1{,}200$ time steps following random initialisation ($\mathbf{x}_P(0) \sim \mathcal{N}(\mathbf{0}, 0.09\,\mathbf{I})$). Coupling gains were set to $g = 3.0$ for the active direction and $g = 0$ for the suppressed direction. Noise amplitude was $\sigma = 0.015$. All random seeds were fixed for reproducibility (weight seed = 8, initial condition seed = 108).

10.16   Geometric Synchrony Pipeline Applied to the RNN

The geometric synchrony pipeline described in precedent Section, was applied to the RNN trajectories with the following adaptations.

10.16.1   Strict separation of lag estimation and geometric alignment

A key methodological constraint is that Procrustes alignment and cross-correlation lag estimation must not be chained. Applying Procrustes before cross-correlation erases the temporal lag by rotating PopB to maximally overlap PopA simultaneously across all time points, artificially inflating apparent synchrony. Therefore:

1. **Temporal lag** was estimated on sign-aligned raw PCA trajectories ($\mathbf{Z}_A$, $\text{sign}(\mathbf{Z}_B)$) via cross-correlation of the first three principal components, searching over both PCA sign polarities to resolve the sign ambiguity inherent to PCA. The peak lag across components and polarities was retained.

2. **Geometric metrics** (pointwise distance, subspace angles, sliding-window alignment) were computed after Procrustes alignment of $\mathbf{Z}_B$ onto $\mathbf{Z}_A$, as described in Section ??.

The lag convention adopted is: a negative lag (cross-correlation peak at $\tau < 0$) indicates that A leads B (B is a delayed copy of A), while a positive lag indicates that B leads A.

10.16.2   Validation criteria

A condition was considered correctly recovered if the peak lag satisfied: directed ($g_{AB} = 3$): $\tau < -3$ (A leads); null: $|\tau| \le 15$ (no systematic lead); reversed ($g_{BA} = 3$): $\tau > 3$ (B leads).

10.17 Quasi-Potential Energy Landscape

To characterise the geometry of the population attractor and its evolution over time, a quasi-potential energy landscape was estimated from the latent trajectories. The landscape U was defined as the negative log-probability density of the population state in the first two latent dimensions:

$$U(x_1, x_2) = -\log P(x_1, x_2), \tag{44}$$

where P was estimated by a Gaussian kernel density estimator (KDE) with bandwidth parameter $h = 0.35$ (Scott's method):

$$\hat{P}(x_1, x_2) = \frac{1}{T} \sum_{t=1}^{T} K_h(x_1 - z_{t,1}, \ x_2 - z_{t,2}), \tag{45}$$

evaluated on an 80×80 grid spanning the observed latent range with 0.5-unit margins on each side. The estimated landscape was smoothed with a Gaussian filter ($\sigma = 2.0$ grid units) and shifted so that its minimum equals zero. Minima of U correspond to preferred (high-probability) population states; maxima correspond to energy barriers separating distinct attractor basins.

10.17.1 Temporal landscape analysis

To capture the time-varying geometry of the shared attractor, the landscape was re-estimated within sliding windows of half-width $w_\ell = 30$ samples, pooling the trajectories of both populations within each window. Two rolling metrics were derived:

1. **Well depth**: $U_{\max} - U_{\min}$ within the window, quantifying attractor strength; deeper wells correspond to more confined, structured population dynamics.

2. **Centroid separation**: Euclidean distance between the time-windowed mean positions of PopA and PopB in latent space, serving as a proxy for geometric dominance (larger separation indicates that the two populations occupy distinct regions of the attractor).

Snapshot landscapes were also constructed for four equal temporal windows (each of duration $T/4$) and visualised as both three-dimensional surfaces (U versus latent dimensions 1 and 2) and two-dimensional contour maps with time-gradient coloured trajectories.

References

1. Mollard, P., Hodson, D. J., Lafont, C., Rizzoti, K. & Drouin, J. A tridimensional view of pituitary development and function. *Trends in Endocrinology & Metabolism* **23**, 261–269. ISSN: 1043-2760. https://www.sciencedirect.com/science/article/pii/S1043276012000288 (2025) (June 2012).

2. Le Tissier, P. *et al.* An updated view of hypothalamic–vascular–pituitary unit function and plasticity. en. *Nature Reviews Endocrinology* **13**, 257–267. ISSN: 1759-5037. https://www.nature.com/articles/nrendo.2016.193 (2025) (May 2017).

3. Bonnefont, X. & Mollard, P. Electrical activity in endocrine pituitary cells in situ: A support for a multiple-function coding. *FEBS Letters* **548**, 49–52. ISSN: 0014-5793. https://www.sciencedirect.com/science/article/pii/S0014579303007270 (2025) (July 2003).

4. Bonnefont, X., Fiekers, J., Creff, A. & Mollard, P. Rhythmic bursts of calcium transients in acute anterior pituitary slices. eng. *Endocrinology* **141**, 868–875. ISSN: 0013-7227 (Mar. 2000).

5. Hodson, D. J. *et al.* Existence of long-lasting experience-dependent plasticity in endocrine cell networks. en. *Nature Communications* **3**, 605. ISSN: 2041-1723. https://www.nature.com/articles/ncomms1612 (2025) (Jan. 2012).

6. Drummond, J. B., Ribeiro-Oliveira, A. & Soares, B. S. eng. in *Endotext* (eds Feingold, K. R. *et al.*) (MDText.com, Inc., South Dartmouth (MA), 2000). http://www.ncbi.nlm.nih.gov/books/NBK534880/ (2026).

7. Molitch, M. E. Pituitary incidentalomas. *Best Practice & Research Clinical Endocrinology & Metabolism. Pituitary Tumours* **23**, 667–675. ISSN: 1521-690X. https://www.sciencedirect.com/science/article/pii/S1521690X09000487 (2026) (Oct. 2009).

8. Sanchez-Cardenas, C. *et al.* Pituitary growth hormone network responses are sexually dimorphic and regulated by gonadal steroids in adulthood. eng. *Proceedings of the National Academy of Sciences of the United States of America* **107,** 21878–21883. ISSN: 1091-6490 (Dec. 2010).

9. Bonnefont, X. *et al.* Revealing the large-scale network organization of growth hormone-secreting cells. *Proceedings of the National Academy of Sciences* **102,** 16880–16885. https://www.pnas.org/doi/10.1073/pnas.0508202102 (2025) (Nov. 2005).

10. Fauquier, T., Guérineau, N. C., McKinney, R. A., Bauer, K. & Mollard, P. Folliculostellate cell network: A route for long-distance communication in the anterior pituitary. *Proceedings of the National Academy of Sciences* **98,** 8891–8896. https://www.pnas.org/doi/abs/10.1073/pnas.151339598 (2025) (July 2001).

11. Le Tissier, P. R. *et al.* Anterior pituitary cell networks. eng. *Frontiers in Neuroendocrinology* **33,** 252–266. ISSN: 1095-6808 (Aug. 2012).

12. Veldhuis, J. D., Carlson, M. L. & Johnson, M. L. The pituitary gland secretes in bursts: appraising the nature of glandular secretory impulses by simultaneous multiple-parameter deconvolution of plasma hormone concentrations. *Proceedings of the National Academy of Sciences of the United States of America* **84,** 7686–7690. ISSN: 0027-8424. https://pmc.ncbi.nlm.nih.gov/articles/PMC299365/ (2026) (Nov. 1987).

13. Goor, F. V., Zivadinovic, D., Martinez-Fuentes, A. J. & Stojilkovic, S. S. Dependence of Pituitary Hormone Secretion on the Pattern of Spontaneus Voltage-gated Calcium Influx. English. *Journal of Biological Chemistry* **276,** 33840–33846. ISSN: 0021-9258, 1083-351X. https://www.jbc.org/article/S0021-9258(19)34785-4/abstract (2025) (Sept. 2001).

14. Stojilkovic, S. S. Pituitary cell type-specific electrical activity, calcium signaling and secretion. *Biological Research* **39,** 403–423. ISSN: 0716-9760. http://www.scielo.cl/scielo.php?script=sci_abstract&pid=S0716-97602006000300004&lng=es&nrm=iso&tlng=en (2025) (2006).

15. Stojilkovic, S. S. Signaling pathways regulating pituitary functions. *Molecular and cellular endocrinology* **463,** 1–3. ISSN: 0303-7207. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6326166/ (2025) (Mar. 2018).

16. Christian, H. C. en. in *Neurosecretion: Secretory Mechanisms* (eds Lemos, J. R. & Dayanithi, G.) 173–193 (Springer International Publishing, Cham, 2020). ISBN: 978-3-030-22989-4. https://doi.org/10.1007/978-3-030-22989-4_9 (2026).

17. Santiago-Andres, Y., Golan, M. & Fiordelisio, T. Functional Pituitary Networks in Vertebrates. *Frontiers in Endocrinology* **11.** ISSN: 1664-2392. https://www.frontiersin.org/articles/10.3389/fendo.2020.619352 (2023) (2021).

18. Briano, C., Meikle, A., Velazco, J. I. & Quintans, G. Metabolic and hormonal profiles and productive performance in primiparous and multiparous cows grazing different forage allowance in late gestation. *Theriogenology* **227,** 68–76. ISSN: 0093-691X. https://www.sciencedirect.com/science/article/pii/S0093691X24002735 (2026) (Oct. 2024).

19. en-US. in *Maternal-Fetal and Neonatal Endocrinology* 189–205 (Academic Press, Jan. 2020). https://www.sciencedirect.com/science/chapter/edited-volume/abs/pii/B9780128148235000143 (2026).

20. Brake, N. *et al.* A neurophysiological basis for aperiodic EEG and the background spectral trend. en. *Nature Communications* **15,** 1514. ISSN: 2041-1723. https://www.nature.com/articles/s41467-024-45922-8 (2024) (Feb. 2024).

21. Donoghue, T. *A historical overview of the study of aperiodic neural activity* en-us. Nov. 2024. https://osf.io/zrvxa_v1 (2025).

22. Donoghue, T. *A systematic review of aperiodic neural activity in clinical investigations* en. Pages: 2024.10.14.24314925. Oct. 2024. https://www.medrxiv.org/content/10.1101/2024.10.14.24314925v1 (2025).

23. Aquiles, A. *et al. Assessing the degree of cortical dislamination through electrical pattern analysis* en. Pages: 2025.06.16.659942 Section: New Results. June 2025. https://www.biorxiv.org/content/10.1101/2025.06.16.659942v2 (2025).

24. Rombouts, J. & Gelens, L. Dynamic bistable switches enhance robustness and accuracy of cell cycle transitions. *PLoS Computational Biology* **17,** e1008231. ISSN: 1553-734X. https://pmc.ncbi.nlm.nih.gov/articles/PMC7817062/ (2026) (Jan. 2021).

25. Joshi, B. & Shiu, A. A Survey of Methods for Deciding Whether a Reaction Network is Multi-stationary. en. *Mathematical Modelling of Natural Phenomena* **10,** 47–67. ISSN: 0973-5348, 1760-6101. https://www-mmnp-journal-org.insb.bib.cnrs.fr/articles/mmnp/abs/2015/05/mmnp201510p47/mmnp201510p47.html (2026) (2015).

26. Strogatz, S. H. & Stewart, I. Coupled Oscillators and Biological Synchronization. *Scientific American* **269,** 102–109. ISSN: 0036-8733. https://www.jstor.org/stable/24941731 (2023) (1993).

27. Izhikevich, E. M. *Dynamical Systems in Neuroscience: The Geometry of Excitability and Bursting* en. ISBN: 978-0-262-27607-8. https://direct.mit.edu/books/monograph/2589/Dynamical-Systems-in-NeuroscienceThe-Geometry-of (2026) (The MIT Press, July 2006).

28. Mastrogiuseppe, F. & Ostojic, S. Linking Connectivity, Dynamics, and Computations in Low-Rank Recurrent Neural Networks. *Neuron* **99,** 609–623.e29. ISSN: 0896-6273. https://www.sciencedirect.com/science/article/pii/S0896627318305439 (2024) (Aug. 2018).

29. Izhikevich, E. M. *Dynamical Systems in Neuroscience* en. ISBN: 978-0-262-09043-8 (MIT Press, 2007).

30. Le Tissier, P. *et al.* An updated view of hypothalamic-vascular-pituitary unit function and plasticity. eng. *Nature Reviews. Endocrinology* **13,** 257–267. ISSN: 1759-5037 (May 2017).

31. Spontaneous oscillations of intracellular calcium and growth hormone secretion. en-US. *Journal of Biological Chemistry* **263,** 9682–9685. ISSN: 0021-9258. https://www.sciencedirect.com/science/article/pii/S0021925819815715 (2025) (July 1988).

32. Kidokoro, Y. Spontaneous calcium action potentials in a clonal pituitary cell line and their relationship to prolactin secretion. en. *Nature* **258,** 741–742. ISSN: 1476-4687. https://www.nature.com/articles/258741a0 (2025) (Dec. 1975).

33. Kwiecien, R. & Hammond, C. Differential Management of Ca2+ Oscillations by Anterior Pituitary Cells: A Comparative Overview. *Neuroendocrinology* **68,** 135–151. ISSN: 0028-3835. https://doi.org/10.1159/000054360 (2026) (Sept. 1998).

34. Kuryshev, Y. A., Childs, G. V. & Ritchie, A. K. Corticotropin-releasing hormone stimulates Ca2+ entry through L- and P-type Ca2+ channels in rat corticotropes. *Endocrinology* **137,** 2269–2277. ISSN: 0013-7227. https://doi.org/10.1210/en.137.6.2269 (2026) (June 1996).

35. Tsaneva-Atanasova, K., Sherman, A., van Goor, F. & Stojilkovic, S. S. Mechanism of spontaneous and receptor-controlled electrical activity in pituitary somatotrophs: experiments and theory. eng. *Journal of Neurophysiology* **98,** 131–144. ISSN: 0022-3077 (July 2007).

36. Tabak, J., Tomaiuolo, M., Gonzalez-Iglesias, A. E., Milescu, L. S. & Bertram, R. Fast-activating voltage- and calcium-dependent potassium (BK) conductance promotes bursting in pituitary cells: a dynamic clamp study. eng. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* **31,** 16855–16863. ISSN: 1529-2401 (Nov. 2011).

37. Stojilkovic, S. S., Tabak, J. & Bertram, R. Ion Channels and Signaling in the Pituitary Gland. *Endocrine Reviews* **31,** 845–915. ISSN: 0163-769X. https://doi.org/10.1210/er.2010-0005 (2025) (Dec. 2010).

38. Gahete, M. D. *et al.* Understanding the multifactorial control of growth hormone release by somatotropes: lessons from comparative endocrinology. eng. *Annals of the New York Academy of Sciences* **1163,** 137–153. ISSN: 1749-6632 (Apr. 2009).

39. Gao, R., Peterson, E. J. & Voytek, B. Inferring synaptic excitation/inhibition balance from field potentials. *NeuroImage* **158,** 70–78. ISSN: 1053-8119. https://www.sciencedirect.com/science/article/pii/S1053811917305621 (2025) (Sept. 2017).

40. Donoghue, T. *et al.* Parameterizing neural power spectra into periodic and aperiodic components. en. *Nature Neuroscience* **23,** 1655–1665. ISSN: 1546-1726. https://www.nature.com/articles/s41593-020-00744-x (2025) (Dec. 2020).

41. Voytek, B. *et al.* Age-Related Changes in 1/f Neural Electrophysiological Noise. eng. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience* **35,** 13257–13265. ISSN: 1529-2401 (Sept. 2015).

42. He, M.-L., Gonzalez-Iglesias, A. & Stojilkovic, S. Role of Nucleotide P2 Receptors in Calcium Signaling and Prolactin Release in Pituitary Lactotrophs. *The Journal of biological chemistry* **278,** 46270–7 (Dec. 2003).

43. Stojilkovic, S. S., Bjelobaba, I. & Zemkova, H. Ion Channels of Pituitary Gonadotrophs and Their Roles in Signaling and Secretion. *Frontiers in Endocrinology* **8,** 126. ISSN: 1664-2392. https://pmc.ncbi.nlm.nih.gov/articles/PMC5465261/ (2026) (June 2017).

44. Sneyd, J. *et al.* On the dynamical structure of calcium oscillations. *Proceedings of the National Academy of Sciences* **114.** Publisher: Proceedings of the National Academy of Sciences, 1456–1461. https://www.pnas.org/doi/10.1073/pnas.1614613114 (2026) (Feb. 2017).

45. Bertram, R., Tabak, J., Teka, W., Vo, T. & Wechselberger, M. en. in *Mathematical Analysis of Complex Cellular Activity* (eds Bertram, R. *et al.*) 1–52 (Springer International Publishing, Cham, 2015). ISBN: 978-3-319-18114-1. https://doi.org/10.1007/978-3-319-18114-1_1 (2026).

46. Fletcher, P. A. & Li, Y.-X. An Integrated Model of Electrical Spiking, Bursting, and Calcium Oscillations in GnRH Neurons. English. *Biophysical Journal* **96.** Publisher: Elsevier, 4514–4524. ISSN: 0006-3495, 1542-0086. https://www.cell.com/biophysj/abstract/S0006-3495(09)00757-7 (2026) (June 2009).

47. Gallego, J. A., Perich, M. G., Miller, L. E. & Solla, S. A. Neural Manifolds for the Control of Movement. eng. *Neuron* **94,** 978–984. ISSN: 1097-4199 (June 2017).

48. Rossi-Pool, R. & Romo, R. Low Dimensionality, High Robustness in Neural Population Dynamics. English. *Neuron* **103.** Publisher: Elsevier, 177–179. ISSN: 0896-6273. https://www.cell.com/neuron/abstract/S0896-6273(19)30571-9 (2026) (July 2019).

49. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. en. *Nature Methods* **17.** Number: 3, 261–272. ISSN: 1548-7105. https://www.nature.com/articles/s41592%E2%80%90019%E2%80%900686%E2%80%902 (2023) (Mar. 2020).

50. Tort, A. B. L., Komorowski, R., Eichenbaum, H. & Kopell, N. Measuring phase-amplitude coupling between neuronal oscillations of different frequencies. eng. *Journal of Neurophysiology* **104,** 1195–1210. ISSN: 1522-1598 (Aug. 2010).

51. Pedregosa, F. *et al.* Scikit-learn: Machine learning in Python. *the Journal of machine Learning research* **12,** 2825–2830. http://www.jmlr.org/papers/volume12/pedregosa11a/pedregosa11a.pdf?source=post_page (2025) (2011).

Figures

Graphical Abstract. Structural and functional distinctions between homotypic and heterotypic interactions have been widely described in the pituitary endocrine system. However, whether functional differences in intrinsic calcium time-series dynamics are relevant to pulsatile hormone secretion remains unexplored. Here, we classify the spontaneous activity underlying both homotypic and heterotypic interactions and characterise their synchrony. We find that heterotypic interactions exhibit transient bistability, consistent with a Hopf-type oscillator regime, in which slow oscillations drive secretory output according to physiological demand.

Figure 1: Periodic and aperiodic descriptors of spontaneous oscillatory activity across physiological conditions. **(A)** Spontaneous activity recorded from three distinct cell populations under control (Virgin) conditions. Blue-green traces correspond to lactotrophs; green traces represent the full mixed population (see diagram, left panel). The right panel shows a representative phase–amplitude coupling (PAC) comodulogram in the Virgin condition; the colour scale indicates the Modulation Index (MI) normalised within each cell population. Lower sub-panels show the dominant amplitude frequencies (left) and the MI compared across conditions and populations (right; group-level comparison using Kruskal–Wallis for Lactotrophs ($H = 5.9$, $p = 0.12$, n.s.) and pairwise Mann–Whitney U with Bonferroni correction for Heterotypic recordings; Heterotypic MI was significantly higher in Lactant than in Multipara ($p = 0.003$, $**$), with no other significant pairwise differences detected; within the Lactant condition, Heterotypic MI was significantly greater than Lactotroph MI ($p < 0.05$, $*$)) **(B)** Mean log-power spectrum for each population (dark line $\pm$ shaded variance), shown separately for homotypic (lactotroph) and heterotypic populations across all conditions. **(C)** Distribution of aperiodic exponent values across conditions and populations (Kruskal-Wallis + Dunn post-hoc with Bonferroni correction; $* = p < 0.05$, $** = p < 0.01$, $*** = p < 0.001$).

Figure 2: Sensitivity of the aperiodic exponent to spectral signal classification and entropy. **(A)** Left: spectral decomposition-based classification of spontaneous calcium time-series. Inset panels show representative examples of two signal classes: Class 1 (violet) and Class 2 (aqua). The scatter plot projects cell classes onto the first two principal dimensions (shown for multiparous lactotrophs). Top-right: cumulative variance explained across the five principal dimensions, for each cell population and condition (colour code as in Figure 1). **(B)** Group comparison of aperiodic slope as a function of signal class. The triangular matrix indicates the proportion of cells in Class 1 (upper triangle) or Class 2 (lower triangle), with the colourmap encoding the z-scored aperiodic exponent. Square panels display the mean aperiodic exponent $\pm$ 95% confidence intervals per class (colours as in panel A). **(C)** Group-level entropy (H) values across populations and conditions. Distributions are shown separately for Class 1 (top) and Class 2 (bottom), shaded in their respective class colours. Square panels show significant Spearman correlations (***$p < 0.05$) between entropy and the aperiodic exponent, per population and condition. **(D)** Percentage of cells with low, medium, and high aperiodic–entropy co-classification values, stratified by oscillatory class (Class 1, right; Class 2, left) and population. Stacked bar charts show mean percentages across recordings ($n = 4$ per condition, individual values overlaid as points), with blue, gray, and red bars denoting low, medium, and high aperiodic–entropy bins, respectively, for homotypic (top) and heterotypic (bottom) populations across physiological conditions. Significant effect in Class 1 homotypic cells ($F = 4.62$, $p = 0.003$), with no individual pairwise comparison surviving correction. No significant effects were found in Class 1 heterotypic ($F = 1.38$, $p = 0.27$), Class 2 homotypic ($F = 0.88$, $p = 0.55$), or Class 2 heterotypic cells ($F = 1.40$, $p = 0.24$).

Figure 3: Synchronous and asynchronous dynamics, their relationship with signal classes, and directional information dominance. (A) Distribution of pairwise Spearman correlation coefficients across cell populations and conditions. The middle panel shows a representative adjacency matrix for the Virgin condition, with distinct population clusters; positive correlations are shown in red, negative in blue. The right panel compares the proportion of cells displaying positive (synchronous) versus negative (asynchronous) correlations across conditions. **(B)** Mean Spearman correlation coefficient (ρ) across recordings and conditions. **(C)** Confusion matrices from random-forest classification, illustrating the relationship between signal class membership and correlation coefficient values. Entries and colour scale represent the percentage of true-positive cells. The left panel shows mean cross-validation accuracy, ordered as in the confusion matrices. **(D)** Direction Asymmetry Index (DAI) stratified by signal class, condition, and network type. DAI < 0 indicates Class 2 dominance; DAI $= 0$ indicates no dominance; DAI > 0 indicates Class 1 dominance. Individual sample values are shown as dots; the mean DAI per condition and population is indicated by a diamond symbol.

Figure 4: Geometric landscape structure of synchronous dynamical states.
(A) Class synchrony in embedding time. Trajectories of two representative signal classes
are shown for homotypic networks (top) and heterotypic networks (bottom). Grey
shading marks the shared state-space regions visited by both trajectories, illustrating
their subsequent divergence in representative examples from the Lactant, Weaned, and
Multiparous conditions. **(B)** Temporal alignment relative to class trajectories. Green:
probability density of trajectory points indicating high inter-class synchronisation.
Orange: trajectory points exhibiting low synchronisation, associated with dominant-
population activity. **(C)** Instantaneous dominance angles between Classes 1 and 2
across conditions and network types. Boxplots display principal angle values for
significant dimensions, colour-coded by condition, for homotypic (top) and heterotypic
(bottom) networks. The dashed line marks the chance angle of $45°$. The middle
panel shows the spontaneous-activity quasi-potential landscape of a Virgin heterotypic
network, highlighting both fixed-point wells (Class 1 and Class 2) and the energy
barrier separating them. Right boxplots depict barrier height between the Class 1 and
Class 2 wells (arbitrary units). **(D)** Three-dimensional landscape representations for
Class 1 (blue) and Class 2 (red), from a representative Weaned heterotypic network.
Bottom: top-down views of Class 1 and two Class 2 examples illustrating a single
stable fixed point. The bar chart shows the percentage of homotypic and heterotypic
networks exhibiting bistability (more than one stable point) versus unimodality (single
stable point), as determined by a Gaussian mixture test for multimodality.

Figure 5: Low-rank dynamic geometric shift: generative null model.
(A) Model architecture comprising two coupled rank-2 recurrent neural networks
(W_A and W_B) with asymmetric coupling strengths ($A \to B = 3.0$, $B \to A = 3.0$; null
symmetric coupling $= 0.0$). Top: schematic of the coupling between population A (50
units) and population B (80 units). Bottom: construction of the weight matrix from the
rank-1 vectors $\mathbf{m}$ and $\mathbf{n}$. Left inset: three representative synthetic time series generated
by the null model; the pink segment indicates epochs in which population A leads, and
the blue segment indicates epochs in which population B leads. **(B)** Quasi-potential
geometric landscapes for the three coupling cases. Case 1: directed connection $A \to B$
(star marks trajectory onset, cross marks end). Case 2: reversed direction $B \to A$.
Null case: symmetric coupling. The shaded bottom panel shows the quasi-potential
landscape with the dominance angle of population A for Case 1. **(C)** Bootstrap
instantaneous dominance angles from population A to population B. Angle > 0: popu-
lation A dominates; angle < 0: population B dominates. Significance brackets indicate
FDR-corrected Kruskal–Wallis tests per population (ns: non-significant; $^{***}p < 0.05$).
(D) Effect of tonic input (I) to population A on its latent representation within the
population B latent space. Blue square: no input (autonomous oscillation); yellow:
$I = 0.6$; red: $I = 1.2$. Magenta phase-portrait vectors correspond to population B;
blue vectors to population A. Arrow direction and colour encode oscillation direction
and input strength, respectively. **(E)** Two-dimensional phase portrait for Case 1, with
and without tonic input. Background colourmap: energy landscape (magenta = high
energy, yellow = low energy). Red trajectories: population B; blue trajectories: pop-
ulation A. White arrows indicate oscillation direction for $I = 0.3$ and $I = 0$. Right
inset: representative time series projected onto the first principal component over
600 ms, illustrating how population A acts as a leader, increasing the temporal lag
(Δt) between the two populations when $I = 0.6$ (top example).