Strategy Sets the Scene: Genetic architecture of linalool resistance in *Botrytis cinerea*

Melanie Madrigal*[1], Jenna C. Moseley[1], Daniel J. Kliebenstein[2], Jordan A. Dowell[1,3]

Department of Biological Sciences, Louisiana State University, Baton Rouge, Louisiana 70803, USA[1]

Department of Plant Sciences, University of California, Davis, California 95616, USA[2]

Department of Plant Pathology and Crop Physiology, Louisiana State University AgCenter, Baton Rouge, Louisiana 70803, USA[3]

*Correspondence to: madrigal.melanie007@gmail.com

Emails of authors: moseleyjc05@gmail.com; kliebenstein@ucdavis.edu; jordan.dowell@gmail.com

Abstract:

Botrytis cinerea is a necrotrophic fungal pathogen that infects thousands of plant species. During infection, these diverse plant hosts produce different specialized metabolites that can inhibit pathogen growth and shape pathogen fitness. However, the genetic architecture of pathogen resistance toward individual host defense metabolites remains poorly understood. To address this question, we exposed 83 *B. cinerea* isolates to the metabolite linalool and quantified metabolic and structural responses. Exposure revealed extensive phenotypic diversity across isolates. Genome-wide association identified 101 genes of interest associated with membrane transport and stress response regulation. Genetic associations were stronger for morphological traits than for metabolic traits, suggesting that hyphal architecture may have a complex genetic architecture contributing to linalool resistance. Together, these results establish natural variation in linalool

response and provide candidate loci for understanding how generalist pathogens respond to host-

derived chemical defenses.

Article Summary:

To understand how a generalist pathogen responds to host defenses, we asked how *Botrytis cinerea*
responds to linalool, a widespread monoterpene involved in plant defense. We exposed 83 *B. cinerea*
isolates to 1000 µM of linalool for 72 hours and quantified metabolic traits (growth curves and
growth dynamics over time) and morphological traits (hyphal network features). Using GWA, we
linked phenotypic variation to genetic variants. Results indicate substantial natural variation in
linalool resistance and distinct genetic architectures across trait classes: metabolic responses are
driven by a relatively small number of loci with larger effects, whereas structural/morphological
responses appear more polygenic.

38
Introduction

Plant-pathogen interactions are dynamic, often shaped by host and pathogen genetic

factors, as well as by other complex evolutionary and ecological processes. Plants respond to

pathogens through multiple defenses, including changes in metabolism, such as the induction of

the jasmonic or salicylic acid pathways. These signaling pathways coordinate the production of

specialized metabolites (SMs). These SMs vary in structure, abundance, and toxicity across

different plant species, serving as chemical barriers that can inhibit pathogen growth[24]. Among

SMs, volatile organic compounds (VOCs) can mediate information transfer between hosts and

their pathogens, along with directly inhibiting pathogen growth[29]. VOCs are released in response

to herbivore or pathogen attack and abiotic stressors, indicating damaged plant tissue or

compromised immunity. At the same time, these VOCs can prime plants for future attacks, such

that post-exposure plants mount more severe responses when they are attacked [10,15,32]. Thus,

VOCs have been a major target for plant-derived pesticides due to their low molecular weight,

high vapor pressure, low water solubility, and their ability to mediate resistance to multiple

biological targets.[59] Due to the multifaceted effects of VOCs on pathogens and reduced

environmental persistence, they are an attractive alternative to conventional synthetic

pesticides.[59]

As VOC profiles vary with host identity[62] and environment,[25] they provide rich

information about host identity and condition[16]. Microorganisms may perceive these VOCs as a

signal of host vulnerability. For example, the slime mold *Physarum polycephalum* exhibits

chemotaxis toward/away from multiple VOCs [35], with linalool causing *P. polycephalum* to direct

its growth away from the compound. Among plant-associated habitats, such as the rhizosphere,

fungi respond to plant VOCs, which can modulate fungal development and influence regulatory

pathways associated with stress responses and metabolism [27].

The generalist necrotrophic pathogen *Botrytis cinerea* provides an ideal system for

investigating fungal responses to VOCs. *B. cinerea* infects thousands of hosts with few genetic

indicators of host specificity[44] despite evidence that gene expression responses can be host

specific[49]. As such, virulence in *B. cinerea* is polygenic, with small-effect loci contributing to

infection success rather than a few major-effect loci [7,14,55]. Moreover, it relies on a multitude of

broad-spectrum virulence strategies, with genome-wide association studies demonstrating that

variation in virulence genes is distributed across the genome and shows little enrichment for

specific pathways or compartments [14,55]. As a necrotrophic pathogen, *B. cinerea* relies not only

on host tissue degradation but also on its capacity to overcome host defenses through a variety of

mechanisms, including enzymatic detoxification and transporter-mediated efflux [63,66]. Further,

the sheer diversity of hosts suggests that *B. cinerea* individuals may be exposed to a wide range

of lineage-specific and general VOCs, necessitating strategies to deal with an array of complex

SMs and VOCs alike.

Among plant VOCs, linalool is found in over 200 monocotyledonous and dicotyledonous

plant species, belonging to over 50% of plant families [4]. Linalool is an acyclic monoterpene

alcohol that is produced through the methylerythritol phosphate (MEP) pathway, where geranyl

diphosphate (GPP) serves as its immediate precursor and is catalyzed by linalool synthases,

members of the large terpene synthase gene family, and is responsible for the production of other

monoterpenes. Reported concentrations of linalool range from 0.01 ppm to over 0.6 ppm in fruit

tissue and from 1.6 ppm up to 20000 ppm in leaves[21] .

As a VOC, linalool plays several roles, from attraction to defense. Linalool is a major

constituent of the floral scent of many species and is widely used to attract pollinators, which

rely on sensory cues to find suitable hosts[31], and as a cue for nearby plants, informing them of

environmental conditions [27]. Linalool also acts as a direct defense mechanism against pathogens.

For example, in *Pseudomonas fluorescens,* linalool reduces membrane potential, increases

cellular leakage, and decreases protein content [28]. Similar modes of action have been observed in

fungal pathogens, including *Candida*[33], and *B. cinerea*. Specifically, Xu et al. reported that when

exposed to linalool, a *B. cinerea* strain downregulates ergosterol biosynthesis genes and

upregulates genes encoding CYP450 enzymes, a large and diverse family of enzymes involved in

xenobiotic detoxification, specialized metabolism, and lipid modification. Reduced ergosterol

biosynthesis compromises plasma membrane integrity, disrupting membrane-associated

signaling processes and membrane structure, as ergosterol is a vital component of the fungal

plasma membrane. In addition to enzymatic detoxification, other studies have identified major

facilitator superfamily transporters (MFS) as key components of resistance and virulence in *B.*

cinerea. These transporters mediate the efflux of toxic compounds from the cell, reducing the

accumulation of plant defensive SMs or other pesticides. Beyond its direct effects on pathogens,

linalool also acts indirectly by stimulating plant-host hormonal pathways. When *Orzo sativa* was

exposed to linalool, it upregulated defense genes and demonstrated enhanced resistance to the

bacterial pathogen *Xanthomonas oryzae*[58].

102   While there is a large support for the direct and indirect effects of linalool on plant

pathogen interactions, most studies focus on a single or a few isolates [53,65]. Although these

approaches provide valuable mechanistic insight, they capture only an individual genotype's

response and do not necessarily reflect how individuals across a genetically diverse population

respond or the strategies they employ. Given the polygenic architecture of virulence in *B. cinerea*

and high effective population size[13], this suggests that candidate genes for linalool resistance

validated in the model isolate of *B. cinerea* (BO5.10)[65] may not capture the full diversity of

resistance mechanisms employed across the species. While genome-wide association studies

(GWAS) have been successfully applied to virulence traits, few studies have examined the

genetic architecture of *B. cinerea's* responses to common specialized metabolites at the

population level [18]. To address this gap, we exposed a population *of B. cinerea* isolates to

linalool to assess population-level variation in resistance and to investigate underlying trait

genetic architecture.

MATERIALS AND METHODS:

Botrytis Collection:

We leveraged 83 isolates of the haploid *B. cinerea* association mapping population previously

described by Atwell et al., 2015, 2018 and Caseys et al., 2021) . Isolates were obtained from

more than 14 different plant species, of which 90% were isolated in California, and 10% were

worldwide strains regularly used for functional genomic studies [3,60]. Within this population, there

is little evidence of host specialization or local population structure, as Casey et al. found Fst

values of 0.03 between strains isolated in Europe and California, and 0.02 among isolates from

different plant host species [14]. All strains used in this study have been previously whole-genome

sequenced to an average of 164-fold coverage[5,14] and aligned to the genome assembly of B05.10

ASM83294v1[60] , resulting in 217,749 SNPs with a minor allele frequency greater than 0.20.

Linalool Exposure:

The isolates used in this study were grown for 2 weeks on oatmeal agar plates. Conidia

were collected from full grown plates using sterile water, counted using an automated spore

counter pipeline(https://github.com/melaniemadrigal07/Post.ImageProcessing) and diluted to a

final concentration of 10 spores/µL in a gamborg B5 (Sigma Aldrich, St. Louis MO)

supplemented with 41 g/L fructose (Thermo Fisher Scientific, Waltham, Massachusetts) 26.65

g/L glucose (Sigma Aldrich, St. Louis MO), 1.03 g/L sucrose (Fisher Scientific, Pittsburgh, PA),

with pH adjusted to 6.5 using 0.01 KOH. To prevent spore germination, spores were kept on ice

from collection to inoculation. 10 µL of the spore suspension (10 spores/µL) were inoculated

into a 96-well plate with 70 µL of liquid Gamborg B5, 10 µL of 1000 µM Linalool (0.045%

DMSO (Sigma Aldrich, St. Louis, MO), and 10 µL of OZ blue cell viability (OZBiosciences,

San Diego, CA), in an 8-fold replication. The 1000 µM linalool concentration was selected based

on its biological relevance, as reported concentrations range from 0.06 µM to 130,000 µM in

fruit tissues. In line with this range, Xu et al. demonstrated minimum inhibitory concentration

(MIC; the lowest dose that inhibited spore germination) and fungicidal concentrations (MFC; the

lowest dose that inhibited mycelial growth) for *B. cinerea* (MIC=20.95 µM; MFC=2,095 µM),

placing 1000 µM within a physiologically meaningful window that is consistent with host-

derived exposure and below fungicidal thresholds.

The OZ blue cell viability assay relies on mitochondrial metabolic activity in viable cells:

the blue-redox dye is taken up and reduced in the mitochondria, resulting in a detectable color

change. Plates were imaged every 4 hours for 72 hours for a total of 20 time points using a

Cytation Gen5 Plate (Agilent, Santa Clara, CA) coupled to a Biospa (Agilent, Santa Clara, CA).

Quantitative colorimetric data were extracted from the time-lapse images for subsequent analysis

of respiratory rates.

Image Analysis

The Cytation Gen5 generated approximately 336,000 images, each exported as a separate red,

green, and blue (RGB) channel. To organize data for analysis, an ImageJ macro was used to

group and compile images by time point for each plate, for downstream analysis. Using these

images, the RGB channels were converted to the L*a*b* color space, and values were extracted

using a custom Python script https://github.com/melaniemadrigal07/GWASLIN). With the L*

value measuring the shift from light to dark, when L = 0, it indicates a dark appearance, and L =

100 indicates a bright appearance. The a* value quantifies the shift from green to red, with

negative a* indicating green and positive a* indicating a strong red color. The b* value is a proxy

for the shift from blue to yellow, with a negative b* indicating a strong blue component and a

positive b* indicating a strong yellow component.

To accurately extract values from the images, a 10% L-value threshold was applied to

isolate hyphal growth in each well. We then measured values external to hyphae, thereby

enabling quantification of color changes in the media associated with fungal metabolic activity.

After extracting the color values, background correction was performed by subtracting the

average L*, a*, and b* values from blank wells (media + dye without spores) from those of

experimental wells at each time point.

Metabolic Activity Quantification

Generated L*, a*, b* values were combined using the following formula to obtain an ΔE

value: $\sqrt{(L_2 - L_1)^2 + (a_2 - a_1)^2 + (b_2 - b_1{}^2)}$, ΔE serves as a proxy for color change relative

to the first time point. This metric was used to generate a curve representing color change over

time. To capture early metabolic activity, we focused on the early germination phase by using

timepoints one to five (up to 20 hours post-inoculation) for downstream analyses. During this 0-

20 h window, spores initiate germination and begin forming penetrative structures[14,17] (e.g.,

appressoria, infection cushions). In addition to ΔE, b* values were analyzed independently, as

they reflect the reduction of the resazurin dye and provide a more specific measure of redox-

driven metabolic activity.

Several phenotypes were extracted from the ΔE curve (Supplementary Table 1) area

under the curve and maximum ΔE. The area under the curve represents the magnitude of the

color change at the early time points. Max ΔE is the value at which isolates reached their peak

color change. Similarly to the phenotypes extracted from ΔE, the same was done for the b*

value, which represents the shift along the blue-yellow axis, with positive values indicating

yellow and negative values indicating blue. The color change range was incorporated as a

phenotype to represent metabolic variation in response to linalool.

To examine the foraging strategies of the strains, hyphal network images were

skeletonized using SkelPy and subsequently processed with SkelpyR to remove skeletonization

artifacts [40].Network-level traits were extracted, including edge lengths, total hyphal length,

number of tips, number of branches, tip fraction, and branch fraction. Total hyphal length and tip

number provide measures of overall growth and active extension, while tip and branch fraction

capture how growth is allocated between exploratory hyphal tips and network consolidation, and

edge length provides information on the spatial scale of individual hyphal segments.

A principal component analysis (PCA) was used to summarize multivariate patterns of

phenotypic variation among *B. cinerea* isolates. PCA was conducted separately for structural

hyphal network traits and metabolic traits, using the same framework for both datasets. PCAs

were performed on centered and scaled trait matrices using singular value decomposition,

excluding traits with zero variance. K-means clustering was subsequently applied to isolate

eigenvectors corresponding to the first two principal components. The number of clusters was set

to k=2 based on inspection of the within-cluster sum of squares and the elbow method.

Clustering was used to summarize the major axes of phenotypic variation rather than to define

discrete phenotypic classes.

Genome-wide Association Study

To estimate the genetic architecture of resistance to linalool in *B. cinerea,* we performed

genome-wide association using GEMMA's univariate linear mixed model [68]. Population structure

and relatedness among isolates were accounted for using a centered genotype matrix in

GEMMA. This approach was based on the model of Zhang et al. (2010) and further adapted by

Dowell et al.[20] In this framework, Single Nucleotide Polymorphisms (SNPs) with a minor allele

frequency (MAF) of > 0.20 and $< 10\%$ missing data were retained, yielding 271,495 SNPs for

analyses. Due to low linkage disequilibrium in *B. cinerea*[5,8], association significance was

assessed using a Bonferroni correction with the family-wise error rate set at $\alpha = 0.05$ divided by

the number of SNPs tested[23].

Following GWA, we assessed linkage disequilibrium (LD) among significantly

associated SNPs using PLINK [46]. Pairwise LD (R^2) was calculated for SNPs within 10 kb

windows centered on significant SNPs to assess regional associations. SNP effect sizes and

standard error were calculated as the beta divided by the trait range, as *B. cinerea* is haploid [42].

Relative effect sizes were summed across all associated regions for each trait to estimate the

relative effect size of the associated genomic regions. Differences in the distribution of effect

sizes between metabolic and structural traits were assessed using a Wilcox rank sum test in R.

Regions associated with multiple traits were considered colocalized. Gene functional

annotations, when available, were extracted from the fungal genomic resources[9]. SNPs were

further classified by genomic location (coding or non-coding) and mutation type (e.g.,

synonymous, missense, or nonsense).

To prioritize associated loci for downstream interpretation, we identified lead SNPs exceeding

the genome-wide significance thresholds ($P < 0.05/m$, where m is the number of SNPs tested)

and defined associated regions using local linkage disequilibrium patterns around each lead SNP

(pairwise R^2 within 10 kb). Among significant regions, loci were ranked by lead SNP P-value,

standardized effect magnitude, and evidence of colocalization across traits. Candidate genes were

defined as those overlapping each region or as the nearest to the lead SNP. For candidate SNPs in

protein-coding genes, we assessed codon bias among synonymous SNPs by reconstructing

reference and alternate codons for each SNP (using SNP context, CDS position, and gene strand)

using a custom R script(https://github.com/melaniemadrigal07/GWASLIN). Usage was then

compared with the genome-wide codon usage frequencies of B05.10's reference genome [57]. All

analyses were conducted within the R environment[48] (R Core Team 2025, version 4.5.3).

Molecular Docking

To explore the potential role of identified variants that directly interact with linalool,

rather than via indirect effects on gene expression or the overall cellular environment, we used an

in-silico protein-ligand binding approach. For each protein of interest, we used D-I-TASSER[67]

with default settings to predict its structure. D-I-TASSER was selected due to its demonstrated

performance in CASP benchmarking. I-T Model confidence was assessed using the reported C-

score (a confidence score that estimates the quality of predicted models) and the estimated TM-

score (a metric for assessing the topological similarity of protein structures), and the top-ranked

model was used for downstream analyses. To assess the interaction of linalool with predicted

protein transporters, we used CaverDock [41]and AutoDock Vina to estimate the binding channel

for simulation, employing a blind docking approach when proteins lack previously identified and

validated active sites. The protein Bcin07g05680 was selected among genes with significantly

associated SNPs as a candidate for docking analysis because it encodes an MFS transporter and

lacks an identified active site. Given its association with metabolic traits, we used a blind

docking approach to identify potential binding sites for linalool.

RESULTS

Metabolic activity

Across *B. cinerea* isolates, the Area Under the Curve ΔE varied from 3.57 to 109 among

samples, with an average of 16.9.99 ($\pm$2.2 *SE,* supplementary Table 2). The slope from timepoint

5 varied from -0.01 to 2.37, with an average of 0.08 ($\pm$0.01 *SE,*Supplementary Table 2). For

traits related to the shift along the b* axis, the maximum blue varied from -1.15 to 24.69, with an

average of 4.66 ($\pm$0.65 *SE,* Supplementary Table 2). The slope of b* ranged from -0.86 to 6.40

among samples, with an average of 0.92 ($\pm$0.17 *SE,* Supplementary Table 2). Area under the

curve along the b axis varied from -6.47 to 79.89 among samples, with an average of 11

($\pm$1.61 *SE*, Supplementary Table S2).

Principal component analysis (PCA) of the ΔE trajectories and blue-axis traits separated

isolates along PC1 and PC2. The PC1 axis accounted for 76.3% of the total variance and

captured the magnitude of the metabolic response, whereas PC2 accounted for 12.6% of the

variance and described differences in response timing and the shape of the metabolic activity

trajectory (Figure 1B).

Hyphal network traits

Structural traits of *B. cinerea* isolates, such as fractal dimension, varied from 0.192 to

1.44, with an average of 0.72 ($\pm$0.02 *SE*, Supplementary Table 2). The slope of the fractal

dimension varied from -0.11 to 0.19, with an average of 0.05 ($\pm$0.006 *SE*, Supplementary Table

S1). Area under the curve of fractal dimension ranged from 1.01 to 8.73

($\pm$0.18 *SE*, Supplementary Table 2). Cumulative total hyphal length across the time course

ranged from 778 to 144,171 µm with an average of 37,519 µm ($\pm$0.02 *SE*, Supplementary Table

SX). Architectural ratios such as tip fraction ranged from 0.702 to 0.937, with an average of

0.787 ($\pm$0.02 *SE*, Supplementary Table S2). The number of branch points (locations where

hyphae bifurcate) ranged from 0.37 to 403, with an average of 122 ($\pm$10.7 *SE*, Supplementary

Table 2). The number of tips (proxy for active regions of the hyphal region) ranged from 4.6 to

863, with an average of 294 ($\pm$ 20.3 *SE*, Supplementary Table 2). Edges representing the

connectivity of the hyphal network ranged from 2.99 to 1279, with an average of 404

($\pm$ 67.3 *SE*, Supplementary Table 2).

A PCA of structural traits revealed three distinct clusters among *B. cinerea* (Figure 1C). PC1

accounts for 69.7% of the variance and is largely driven by connectivity-related traits (high

AUC, fractal dimension, tips, and edges). PC2 accounts for 12.1% of the variance and is driven

by growth metrics (fractal dimension, slope, and fragmentation index). A PCA of both structural

and metabolic traits revealed a similar pattern; PC1 explained 57.3% of the total variance, and

PC2 explained 17.0% (Figure 1C). PC1 was primarily associated with structural traits such as

tips and edges. PC2 captured secondary variation, dominated by metabolic color-shift dynamics

along the b* axis (Figure 1D).

Genetic architecture of resistance

Genome-wide association analyses using the traits as phenotypes identified 202

significant SNP-trait associations and 101 unique SNPs (Figure 2, Supplementary Table 3).

Significant SNPs were distributed across the genome, with the greatest number on chromosome

2 (n=15; 11 non-coding,4 coding), chromosome 8 (n=13; 12 noncoding,1 coding), chromosomes

4 (n=11; 3 non-coding, 8 coding) and 13 (n=11 noncoding), all remaining chromosomes

contained 8 or fewer significant SNPs (Figure 2, Supplementary Table 3). Of the 101 unique

SNPs, 78 were noncoding, 16 were coding-synonymous, and 7 were coding-missense variants.

Several significant SNPs were mapped within or near four genes on chromosome 7

(*Bcin07g05680, Bcin07g05230, Bcin07g05240, and Bcin07g05210*), two distributed across

chromosomes 2 (*Bcin02g09200 and Bcin02g05460*) and 15 (Bcin15g00890), and 3

(*Bcin03g06600*). Although multiple SNPs supported individual loci, overlap among traits was

limited (Figure 2). When present, trait overlap suggests that a shared locus may contribute to

multiple aspects of the linalool response.

Multi-trait associations were rare, with most SNPs associated with a single phenotype,

suggesting that most SNPs contribute to specific components of the linalool response rather than

broadly affecting multiple traits. Here, a pleiotropy is defined as a SNP associated with multiple

phenotypes. Across metabolic-activity traits (ΔE and b features), several traits shared

associations within/near *Bcin07g05680*, which encodes a major facilitator superfamily

transporter. Notably, most SNPs within this locus were synonymous mutations. Metabolic traits

exhibited slightly larger standardized absolute SNP effect magnitudes than structural traits

(median = 0.131 vs 0.127; Figure 3A). However, structural traits included substantially more

unique SNPs (97 vs 7) and significant SNP-trait associations (181 vs 21) (Figure 3B).

Of the SNPs identified, 78 mapped to non-coding regions and 23 to coding regions.

Within coding regions, 16 were synonymous, and 7 were missense variants (Figure 4A). 6 of the

missense variants mapped to Bcin04g02170, a protein kinase domain-containing protein, while

one variant mapped to the transcription factor Bcifm1. Within Bcin04g02170, three substitutions

were altered amino acid charge, and two altered amino acid size class (Supplementary Table 3).

Notably, one substitution changed arginine to glycine, thereby shifting from a positively charged

residue to a nonpolar residue. Arginine is larger and can maintain its charge, allowing stronger

hydrogen and electrostatic bonding, while glycine is smaller, uncharged, and has a minimal

hydrogen side chain, which could increase flexibility and alter glycine's substrate specificity.

Previous work on protein kinases has shown that hydrostatic and electrostatic balance is a major

determinant of kinase-substrate specificity, and that residues outside the active site can influence

substrate specificity through effects on protein structure and electrostatic environments[36]. The

final missense variant mapped to the translation initiation factor Bcifm1, which changed alanine

to valine. Although both amino acids are nonpolar, valine is bulkier due to its branched side

chain. This increase in steric volume could subtly alter local packing, protein stability, or

interaction surfaces[26].

Among synonymous SNPs, codon usage frequency varied in direction and magnitude

(Figure 4B). Six variants (46.2%) shifted towards rarer codons, one variant (7.7%) shifted

towards the preferred codon, and six variants (46.2%) shifted to codons with approximately

equal usage across the genome. At the gene level, synonymous hits predominantly shifted

toward rarer codons (6/7), suggesting a potential directional bias in codon usage changes across

multiple loci, with the largest magnitude change observed in *Bcin07g05680* (MFS transporter-

encoding gene). In contrast, *Bcin02g08180* (AP complex subunit beta) exhibited a synonymous

shift in which the minor allele encoded a more preferred codon relative to the major allele

(Figure 4B).

Docking Results

Bcin07g05680 was selected as a protein of interest for follow-up docking analysis, based

on the presence of significantly associated synonymous SNPs. This gene was selected as a

candidate because its functionality is expected to be conserved across the population, while its

expression may be more sensitive to transcriptional and/or translational variation arising from

codon usage differences [57]. Putative ligand transport pathways were identified using CaverWeb

to calculate potential access tunnels. CaverWeb generated over 232 pockets of interest. As many

of these predicted pathways were either shallow, discontinuous, or sterically constrained,

subsequent analyses focused on tunnels likely to represent physiologically relevant transport

routes. The primary pocket was chosen based on quantitative criteria associated with functional

substrate channels, such as having the widest bottleneck (2.87 $\pm$ 0.43), the shortest path (11.46 $\pm$

5.79), and a curvature of 1.14 ($\pm$ 0.11). These parameters suggested a more accessible path for

ligand transport than other tunnels.[41] Using Caverweb, linalool was docked to transport through

the first tunnel. Transport through this tunnel exhibited a favorable trajectory, with values

ranging from approximately -4.0 kcal/mol near the internal binding region to -2.6 kcal/mol as it

exited the pocket. This trajectory suggests a low energetic barrier for diffusion through the

tunnel, supporting the possibility that this transporter could facilitate the export of small,

lipophilic compounds such as linalool.

Discussion

Phenotypic Variation

Filamentous fungi exhibit substantial phenotypic plasticity in growth form and

metabolism in response to environmental stressors, and these responses often reflect alternative

strategies that balance exploration, resource acquisition, and damage mitigation[12]. In *B. cinerea,*

exposure to linalool revealed isolate-to-isolate variation in metabolic activity, as quantified by

ΔE and b* dynamics (Figure 1B). Isolates with steep ΔE slopes displayed rapid color shifts,

suggesting rapid metabolic activity. Isolates with gradual slopes may experience a slower

metabolic response or higher susceptibility to linalool toxicity. Linalool has been shown to

disrupt mitochondrial function and downregulate mitochondrial and ergosterol genes, as well as

glutathione synthesis [65], which may underlie the observed differences in ΔE trajectories.

These patterns were recapitulated by PCA of metabolic traits (Figure 1B). PC1 (63.8% of

total variance) captured response magnitude, separating isolates with higher AUC and steep ΔE

slopes (fast responders) from those with more gradual slopes (slow responders). PC2 (16.7%)

captured variation in response timing and trajectory, distinguishing isolates with earlier versus

later dye reduction. Although isolates formed two broad groups with minimal overlap along PC1,

overlap among PC2 suggests that metabolic phenotypes are better described as a continuum

dominated by response strength rather than discrete classes. Together, these patterns suggest that

isolates differ in both the strength and the rate at which they maintain metabolic reducing

capacity under linalool exposure, potentially reflecting variation in tolerance mechanisms.

Similarly, Fusarium responds to linalool exposure through broad transcriptional remodeling, with

enrichment in ABC transporter activity, sterol biosynthesis, and glutathione metabolism[37] ,

supporting a model in which linalool perturbs membrane homeostasis, detoxification, and redox

buffering pathways. While resistance to chemical stress can be expressed not only through

physiological detoxification. Filamentous fungi can also respond by reprogramming growth

through adjustments in hyphal tips, branching, and network connectivity to reallocate growth

under inhibitory conditions. In *Rhizophagus irregularis,* exposure to fungicides reduces hyphal

length and alters anastomosis, demonstrating that chemical environments can reshape hyphal

architecture and interaction behavior [49]

Isolates not only differ in metabolic activity but also in structural traits, reflecting

differences in how each isolate's hyphal network responds to linalool (Figure 1A). Network

features, such as fractal dimension, hyphal tips, and edges, varied across the population,

reflecting differences in network complexity and connectivity (Supplementary Table 1). These

traits reveal that some isolates form compact, highly interconnected networks, whereas others

exhibit more of an exploratory phenotype. In the structural PCA (Figure 1A), isolates are

separated into three broad morphological groupings driven by network size and connectivity

metrics (edges, nodes, tips, branches, and total hyphal length). Similar contrasts have been

described in filamentous fungi under different environmental conditions, where dense, cohesive

networks are associated with nutrient-rich environments and exploratory growth with nutrient-

poor environments [2,22]. In the context of linalool, one possibility is that highly connected

networks increase physiological robustness (i.e., maintaining growth by distributing transport

and reducing power demands across a denser mycelial system), while exploratory architectures

reflect avoidance or slower growth under stronger inhibition. More broadly, the observed

architectural plasticity aligns with evidence that *B. cinerea* can undergo hyphal specialization as

a strategy to overcome host defenses[47]. Infection cushions, specialized multicellular structures

that secrete cell wall-degrading enzymes, exhibit higher tolerance to host defenses[47]. The

structural diversity we observe may mirror this, suggesting that remodeling of the hyphal

network is a key in *B. cinerea's* ability to overcome host defenses and enact disease.

Genetic Architecture of Resistance to Linalool

Linalool exposure is known to elicit broad physiological remodeling in *B. cinerea,*

including disruption of mitochondrial function and changes in sterol and redox processes[37,65]. In

a proteomic time course, brief exposure to linalool led to broad proteomic changes, involving

enrichment of proteins involved in translation, amino acid metabolism, and post-translational

modification[65]. These findings suggest that linalool triggers energetic and metabolic

reprogramming in response to mitochondrial stress and membrane disruption.

In our GWAS, the strongest metabolic response associations localized to *Bcin07g05680*,

an MFS transporter-encoding gene. Most of these variants were synonymous, indicating no

change in the amino acid sequence; however, this mutation could influence protein expression or

conformation via mRNA stability or translation efficiency[1]. This mechanism has been

demonstrated in the multidrug resistance gene MDR1, where a synonymous polymorphism

altered substrate specificity by altering folding and drug-protein interactions[34] . The synonymous

mutations in *Bcin07g05680* resulted in shifts toward the use of non-preferred or uncommon

codons (Figure 4B). Importantly, codon usage mismatch has been shown to have strong

functional consequences in *B. cinerea*[57]. Codon optimization of the geneticin resistance gene

NPTII to match *B. cinerea* codon preferences increases transformation by 30-fold, demonstrating

that *B. cinerea* is sensitive to codon usage in heterologous coding sequences[57]. While a shift

toward rarer codons could be detrimental under baseline conditions due to reduced translational

efficiency, the net fitness effect of these synonymous changes may be context-dependent and

could be favored under linalool exposure[43]. In *Salmonella,* codon usage bias has been linked to

stress and virulence-associated gene regulation, where enrichment towards synonymous codons

is thought to support translation under nutrient-limited conditions and allows for colonization of

different host tissues[11].

Efflux transporters are frequently implicated in chemical stress responses in fungal

pathogens. In Fusarium, a GWAS examining sensitivity to the demethylation inhibitor (DMI)

class of fungicides identified candidate loci mapping to transport-related proteins [56], suggesting

that membrane transport is part of a broader genetic toolkit to cope with chemical-related stress.

In *B. cinerea,* for example, transcriptomic comparisons of fludioxonil-resistant isolates found

MFS to be upregulated in fludioxonil-resistant isolates [38]. In addition, the *MfsG* transporter has

been identified as a virulence factor that confers tolerance to other SMs, such as glucosinolates[61].

To explore whether this transporter could interact with linalool, molecular docking was

performed using the predicted *Bcin07g05680* (MFS transporter) protein structure. Notably, the

variants in/near *Bcin07g05680* (MFS transporter) *were* predominantly synonymous and shifted

codon usage toward rarer codons, providing a plausible translation-linked route by which silent

variation could influence transporter performance. In addition, *Bcin07g05210,* a btb domain-

containing protein, was associated with variation in metabolic response, suggesting a potential

regulatory role in mediating cellular adaptation to linalool exposure.

Several SNPs were linked to variation in structural traits, including the area under the

fractal dimension curve, total hyphal length, and tip fraction (Figure 2). These traits capture

differences in hyphal network connectivity and cohesion. Associated loci were in a mitochondrial

translation initiation factor (*Bcin02g09200*), a Zn-C6 domain containing protein

(*Bcin02g05460*), *Bcin02g08180* (AP complex subunit beta), and *bcltf1*. *Bcin02g09200* may

contribute to variation in hyphal structure by regulating mitochondrial efficiency and availability.

As hyphae extend and grow, the hyphal tips rely on ATP for growth; therefore, small shifts in

mitochondrial activity could alter branching frequency and overall network cohesion. The ZN(II)

C6 domain-containing transcription factor (Bcin02g05460) is likely to regulate downstream

genes involved in carbon metabolism, stress signaling, and cell wall synthesis. In other

filamentous fungi, ZN(II) C6 proteins regulate carbon metabolism and drug sensitivity [39].

Bcin02g08180 encodes an AP complex subunit beta, which is involved in vesicle-mediated

trafficking and cargo sorting. Vesicle trafficking is critical for fungal virulence in *B. cinerea* as it

transports sRNAs to plants via extracellular vesicles[30]. Extracellular vesicles carry a diverse set

of cargo that range from involvement in transport to autophagy[19]. *Bcltf1* has been categorized as

a transcription factor that regulates virulence, with deletion mutants exhibiting hypersensitivity

to oxidative stresses, therefore variation within *Bcltf1* could plausibly influence responses to

oxidative stress from linalool exposure[52].

Our *B. cinerea* population showed that in response to linalool exposure, phenotypic

diversity is driven by metabolic and morphological plasticity. At the population level, metabolic

resistance to linalool appears to be governed by a small number of genes, suggesting an

oligogenic basis for this trait. This pattern contrasts with prior studies showing that variation in

B. cinerea virulence and host is polygenic, involving many loci of small effect that shape lesion

formation and host specificity [7,14,51,54,55]. Correlation of lesion areas across different hosts

highlights that virulence is driven by hundreds of genes scattered throughout the genome [6,14,55].

In contrast, resistance to specific host-derived defenses, such as linalool, may be mediated by a

smaller set of key genes, such as transporters with broad substrate specificity or stress response

regulators that can exert larger individual effects on survival and growth under chemical

challenge. Consistent with this, experimental evolution with specialized metabolites can induce

multidrug resistance in *B. cinerea* via upregulation of ABC and MFS transporter genes [64].

Additionally, work in a Michigan *B. cinerea* population found that fungicide resistance was

consistently associated with detectable population differentiation, supporting the idea that

chemical exposure can act as a strong selective pressure and shape the genetic architecture of

resistance in the *B. cinerea* population[45].

These results provide insight into how *B. cinerea* overcomes host defenses and how it

functions with known virulence mechanisms, demonstrating that its success as a generalist

pathogen arises from a mixed genetic architecture that balances a polygenic basis for virulence

with oligogenic resistance mechanisms to linalool. While our study is one of the first to examine

a population of *B. cinerea* response to linalool, our population represents a subset of the total

genetic diversity and may not capture all allelic variation underlying resistance. Moreover, the

population was tested against a single VOC, whereas during natural host-pathogen interactions,

B. cinerea encounters mixtures of hundreds of specialized metabolites. Future work should

incorporate broader isolate sampling and exposure to mixtures of specialized metabolites, which

is necessary to elucidate the full genetic architecture of resistance.

Data Availability Statement

All code used for image processing and downstream analyses is available at .

(https://github.com/melaniemadrigal07/GWASLIN). Supplementary material is available at

10.5281/zenodo.19425171and provides detailed information supporting the analyses performed

in this study. Supplementary Table 1 provides descriptions of the traits used in this study.

Supplementary Table 2 contains trait means. Supplementary Table 3 contains results of the

genome wide association analyses it includes SNPs, genomic positions, effect sizes, and

statistically significant values for all trait-SNP associations. Supplementary Table 4 provides

descriptions of all annotation fields used in SNP characterization including genomic location,

allele identity, codon context, and functional classification. Supplementary Table 5 contains

location of the SNP and classification of the mutation.

Acknowledgments

J.A.D., M.M., and J.C.M. were funded in part by a USDA grant (2022-67012-43019). M.M. was
also funded by the Future Scholar fellowship at Louisiana State University.

M.M. and J.A.D. designed the study; M.M. and J.C.M. collected the data; M.M, J.C.M., and

J.A.D. analyzed the data; and M.M. and J.A.D. wrote the manuscript with input from D.J.K. and

J.C.M.

Study funding

J.D., M.M., and J.M. were funded in part by a USDA grant (2022-67012-43019). M.M. was also
funded by the Future scholar's fellowship at Louisiana State University.

Conflict of interest

None declared.

Figures

Figure 1. Metabolic and structural trait variation across isolates (A, C, E): PCA score plots
of *B. cinerea* isolates projected onto PC1 and PC2 for metabolic (A), structural (C), and
combined (E) trait datasets. Points are colored by k-means cluster assignment (k=3 for structural,
2 for metabolic, and combined), selected isolates are labeled for reference, including B05.10

(reference and generalist isolate), KatieTomato (highly virulent isolate), DavisNavel (low
virulent isolate). (B, D, F) Corresponding loading plots displaying the top contributing traits for
each dataset. Arrows represent variable loadings on PC1 and PC2. In (B), Traits in numerical
order: 1, time point of maximum blue axis shift; 2, maximum delta E at timepoint 5 (20 hpi); 3,
area under the curve at timepoint 5 (20 hpi); 4 slope of blue axis shift at timepoint 5 (20 hpi); 5
slope of delta E at timepoint 5. (D) Traits in numerical order: 1, tip fraction residuals; 2,
weighted tip fractions; 3, area under the curve of fractal dimension across 72 hours; 4, total
hyphal length; 5, mean total hyphal length; 6, median edge length; 7, number of hyphal tips; 8,
number of hyphal nodes. (F) 1, tip fraction residuals; 2, maximum blue axis shift; 3, range of
blue axis; 4 slope of blue axis shift at timepoint 5 (20 hpi); 5, area under the curve of blue axis at
timepoint 5 (20 hpi); 6, tip fraction weighted; 7, area under the curve of delta eat timepoint 5 (20
hpi) 8, median edge length.

Figure 2. Overlap of significant SNPs and trait associations.

Upset plot showing the intersection of significant SNPs across measured traits. Vertical bars
represent the number of SNPs shared among specific combination of traits (intersection size).
Horizontal bars indicate the total number of SNPs associated with each individual trait (set size)
Connected black dots denote traits included in each intersection.

Figure 3. Distribution and overlap of significant SNP associations across structural and metabolic trait classes. A) Distribution of standardized SNP effect sizes for structural and metabolic traits (Wilcox rank sum test=1233, P<0.01). Violins represent the full distribution of absolute effect sizes across significant SNP-trait associations, and boxplots indicate the interquartile range and median. Sample sizes (total significant associations and unique SNPs are shown below each category. Hits refer to significant SNP-trait associations, while SNP counts represent unique variants after collapsing repeated associations across traits. (B) Upset plot showing the number of unique SNPs associated with structural traits, metabolic traits, or both. Vertical bars represent interaction sizes, and the matrix below indicates the trait class for each intersection. Structural SNPs account for the most significant associations, with limited overlap between structural and metabolic traits.

Figure 4. Functional annotation and codon bias shifts among genome-wide significant SNPs. (A) Distribution of genome-wide associated SNPs partitioned into coding and non-coding

550 categories. Coding SNPs are further subdivided into synonymous or missense variants. (B)
551 Change in codon usage bias for synonymous SNPs in coding regions between the alternate and
552 reference alleles. Positive values indicate shifts toward more preferred codons, whereas negative
553 values indicate shifts towards non-preferred codons. Colors denote the direction of change, and
554 gene labels indicate the trait class.

References

1. A "Silent" Polymorphism in the MDR1 Gene Changes Substrate Specificity | Science.
[accessed 2026 Jan 17]. https://www-science-
org.libezp.lib.lsu.edu/doi/10.1126/science.1135308

2. Aguilar-Trigueros CA, Boddy L, Rillig MC, Fricker MD. Network traits predict ecological
strategies in fungi. ISME Communications. 2022;2(1):1–11. doi:10.1038/s43705-021-
00085-1

3. Amselem J, Cuomo CA, Couloux A, Coutinho PM, de Vries RP, Dyer PS, Fillinger S, Gout
L, Hahn M, Kohn L, et al. Genomic Analysis of the Necrotrophic Fungal Pathogens
Sclerotiniasclerotiorum and Botrytiscinerea.

4. Aprotosoaie AC, Hǎncianu M, Costache I-I, Miron A. Linalool: a review on a key odorant
molecule with valuable biological properties. Flavour and Fragrance Journal.
2014;29(4):193–219. doi:10.1002/ffj.3197

5. Atwell S, Corwin J, Soltis N, Subedy A, Denby K, Kliebenstein DJ. Whole genome
resequencing of Botrytis cinerea isolates identifies high levels of standing diversity.
Frontiers in Microbiology. 2015 [accessed 2026 Jan 18];6.

581 https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2015.00996/full.
582 doi:10.3389/fmicb.2015.00996

583 6. Atwell S, Corwin J, Soltis N, Subedy A, Denby K, Kliebenstein DJ. Whole genome
584 resequencing of Botrytis cinerea isolates identifies high levels of standing diversity.
585 Frontiers in Microbiology. 2015 [accessed 2025 Sept 1];6.
586 https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2015.00996/full.
587 doi:10.3389/fmicb.2015.00996

588 7. Atwell S, Corwin JA, Soltis N, Zhang W, Copeland D, Feusier J, Eshbaugh R, Kliebenstein
589 DJ. Resequencing and association mapping of the generalist pathogen *Botrytis cinerea*.
590 2018 [accessed 2024 Sept 14]. http://biorxiv.org/lookup/doi/10.1101/489799.
591 doi:10.1101/489799

592 8. Atwell S, Corwin JA, Soltis N, Zhang W, Copeland D, Feusier J, Eshbaugh R, Kliebenstein
593 DJ. Resequencing and association mapping of the generalist pathogen *Botrytis cinerea*.
594 2018 [accessed 2024 Sept 10]. http://biorxiv.org/lookup/doi/10.1101/489799.
595 doi:10.1101/489799

596 9. Basenko EY, Pulman JA, Shanmugasundram A, Harb OS, Crouch K, Starns D, Warrenfeltz
597 S, Aurrecoechea C, Stoeckert CJ, Kissinger JC, et al. FungiDB: An Integrated Bioinformatic
598 Resource for Fungi and Oomycetes. Journal of Fungi. 2018;4(1):39. doi:10.3390/jof4010039

599 10. Blande JD, Holopainen JK, Li T. Air pollution impedes plant-to-plant communication by
600 volatiles. Ecology Letters. 2010;13(9):1172–1181. doi:10.1111/j.1461-0248.2010.01510.x

601 11. Bumann D. tRNA modifications enable codon usage signatures to coordinate bacterial
602 virulence in Salmonella.

603 12. Camenzind T, Aguilar-Trigueros CA, Hempel S, Lehmann A, Bielcik M, Andrade-Linares
604 DR, Bergmann J, Dela Cruz J, Gawronski J, Golubeva P, et al. Towards establishing a fungal
605 economics spectrum in soil saprobic fungi. Nature Communications. 2024;15(1):3321.
606 doi:10.1038/s41467-024-47705-7

607 13. Caseys C, Shi G, Soltis N, Gwinner R, Corwin J, Atwell S, Kliebenstein D. Quantitative
608 interactions drive *Botrytis cinerea* disease outcome across the plant kingdom. 2018
609 [accessed 2025 Dec 12]. http://biorxiv.org/lookup/doi/10.1101/507491.
610 doi:10.1101/507491

611 14. Caseys C, Shi G, Soltis N, Gwinner R, Corwin J, Atwell S, Kliebenstein DJ. Quantitative
612 interactions: the disease outcome of Botrytis cinerea across the plant kingdom. G3:
613 Genes|Genomes|Genetics. 2021;11(8):jkab175. doi:10.1093/g3journal/jkab175

614 15. Catola S, Centritto M, Cascone P, Ranieri A, Loreto F, Calamai L, Balestrini R, Guerrieri
615 E. Effects of single or combined water deficit and aphid attack on tomato volatile organic

compound (VOC) emission and plant-plant communication. Environmental and
Experimental Botany. 2018;153:54–62. doi:10.1016/j.envexpbot.2018.05.001

16. Cellini A, Biondi E, Blasioli S, Rocchi L, Farneti B, Braschi I, Savioli S, Rodriguez-Estrada
M t., Biasioli F, Spinelli F. Early detection of bacterial diseases in apple plants by analysis of
volatile organic compounds profiles and use of electronic nose. Annals of Applied Biology.
2016;168(3):409–420. doi:10.1111/aab.12272

17. Choquer M, Rascle C, Gonçalves IR, de Vallée A, Ribot C, Loisel E, Smilevski P, Ferria J,
Savadogo M, Souibgui E, et al. The infection cushion of Botrytis cinerea: a fungal 'weapon'
of plant-biomass destruction. Environmental Microbiology. 2021;23(4):2293–2314.
doi:10.1111/1462-2920.15416

18. Corwin JA, Subedy A, Eshbaugh R, Kliebenstein DJ. Expansive Phenotypic Landscape of
Botrytis cinerea Shows Differential Contribution of Genetic Diversity and Plasticity.
Molecular Plant-Microbe Interactions®. 2016;29(4):287–298. doi:10.1094/MPMI-09-15-
0196-R

19. De Vallée A, Dupuy J-W, Moriscot C, Gallet B, Vanderperre S, Guignard G, Rascle C,
Calvar G, Malbert B, Gillet F-X, et al. Extracellular Vesicles of the Plant Pathogen Botrytis
cinerea. Journal of Fungi. 2023;9(4):495. doi:10.3390/jof9040495

20. Dowell JA, Bowsher AW, Jamshad A, Shah R, Burke JM, Donovan LA, Mason CM.
Historic breeding practices contribute to germplasm divergence in leaf specialized
metabolism and ecophysiology in cultivated sunflower (Helianthus annuus). American
Journal of Botany. 2024;111(11):e16420. doi:10.1002/ajb2.16420

21. Duke JA. Handbook of phytochemical constituents of GRAS herbs and other economic
plants. Boca Raton: CRC Press; 2001. (Herbal reference library).

22. Fricker MD, Heaton LLM, Jones NS, Boddy L. The Mycelium as a Network. Microbiology
Spectrum. 2017;5(3):10.1128/microbiolspec.funk-0033–2017.
doi:10.1128/microbiolspec.funk-0033-2017

23. Gao X, Starmer J, Martin ER. A multiple testing correction method for genetic
association studies using correlated single nucleotide polymorphisms. Genetic
Epidemiology. 2008;32(4):361–369. doi:10.1002/gepi.20310

24. Garagounis C, Delkis N, Papadopoulou KK. Unraveling the roles of plant specialized
metabolites: using synthetic biology to design molecular biosensors. New Phytologist.
2021;231(4):1338–1352. doi:10.1111/nph.17470

25. Goh H-H, Khairudin K, Sukiran NA, Normah M n., Baharum S n. Metabolite profiling
reveals temperature effects on the VOCs and flavonoids of different plant populations.
Plant Biology. 2016;18(S1):130–139. doi:10.1111/plb.12403

26. Gregoret LM, Sauer RT. Tolerance of a protein helix to multiple alanine and valine
substitutions. Folding and Design. 1998;3(2):119–126. doi:10.1016/S1359-0278(98)00017-
0

27. Guo D, Liu Z, Raaijmakers JM, Xu Y, Yang J, Erb M, Zhang J, Zhu Y-G, Xu J, Hu L. Linalool-
triggered plant-soil feedback drives defense adaptation in dense maize plantings. Science.
2025;389(6761):eadv6675. doi:10.1126/science.adv6675

28. Guo F, Chen Q, Liang Q, Zhang M, Chen W, Chen H, Yun Y, Zhong Q, Chen W.
Antimicrobial Activity and Proposed Action Mechanism of Linalool Against Pseudomonas
fluorescens. Frontiers in Microbiology. 2021 [accessed 2025 Sept 14];12.
https://www.frontiersin.org/journals/microbiology/articles/10.3389/fmicb.2021.562094/ful
l. doi:10.3389/fmicb.2021.562094

29. Hammerbacher A, Coutinho TA, Gershenzon J. Roles of plant volatiles in defence
against microbial pathogens and microbial exploitation of volatiles. Plant, Cell &
Environment. 2019;42(10):2827–2843. doi:10.1111/pce.13602

30. He B, Wang H, Liu G, Chen A, Calvo A, Cai Q, Jin H. Fungal small RNAs ride in
extracellular vesicles to enter plant cells through clathrin-mediated endocytosis. Nature
Communications. 2023;14(1):4383. doi:10.1038/s41467-023-40093-4

31. Heuel KC, Raguso RA, Coogan E, Mallick R, Keleher KJ, Ayasse M, Gegear RJ, Burger H.
Spatial partitioning of floral volatiles provides a "chemosensory roadmap" for bumblebee
pollinators. Current Biology. 2025;35(7):1622-1630.e6. doi:10.1016/j.cub.2025.02.010

32. Junker RR, Kuppler J, Amo L, Blande JD, Borges RM, van Dam NM, Dicke M, Dötterl S,
Ehlers BK, Etl F, et al. Covariation and phenotypic integration in chemical communication
displays: biosynthetic constraints and eco-evolutionary implications. New Phytologist.
2018;220(3):739–749. doi:10.1111/nph.14505

33. Khan A, Ahmad A, Akhtar F, Yousuf S, Xess I, Khan LA, Manzoor N. Ocimum sanctum
essential oil and its active principles exert their antifungal activity by disrupting ergosterol
biosynthesis and membrane integrity. Research in Microbiology. 2010;161(10):816–823.
doi:10.1016/j.resmic.2010.09.008

34. Kimchi-Sarfaty C, Oh JM, Kim I-W, Sauna ZE, Calcagno AM, Ambudkar SV, Gottesman
MM. A "Silent" Polymorphism in the *MDR* 1 Gene Changes Substrate Specificity. Science.
2007;315(5811):525–528. doi:10.1126/science.1135308

35. de Lacy Costello BPJ, Adamatzky AI. Assessing the chemotaxis behavior of Physarum
polycephalum to a range of simple volatile organic chemicals. Communicative &
Integrative Biology. 2013;6(5):e25030. doi:10.4161/cib.25030

36. Li L, Shakhnovich EI, Mirny LA. Amino acids determining enzyme-substrate specificity
in prokaryotic and eukaryotic protein kinases. Proceedings of the National Academy of
Sciences. 2003;100(8):4463–4468. doi:10.1073/pnas.0737647100

37. Li X, Wang Q, Li H, Wang X, Zhang R, Yang X, Jiang Q, Shi Q. Revealing the Mechanisms
for Linalool Antifungal Activity against Fusarium oxysporum and Its Efficient Control of
Fusarium Wilt in Tomato Plants. International Journal of Molecular Sciences.
2022;24(1):458. doi:10.3390/ijms24010458

38. Liu M, Peng J, Wang X, Zhang W, Zhou Y, Wang H, Li X, Yan J, Duan L. Transcriptomic
Analysis of Resistant and Wild-Type Botrytis cinerea Isolates Revealed Fludioxonil-
Resistance Mechanisms. International Journal of Molecular Sciences. 2023;24(2):988.
doi:10.3390/ijms24020988

39. MacPherson S, Larochelle M, Turcotte B. A Fungal Family of Transcriptional Regulators:
the Zinc Cluster Proteins. Microbiology and Molecular Biology Reviews. 2006;70(3):583–
604. doi:10.1128/MMBR.00015-06

40. Madrigal M, Moseley AJ, Moseley JC, Dowell JA. SkelPy: A graphic user interface–based
approach for skeletonizing fungal networks. Applications in Plant Sciences.
n/a(n/a):e70049. doi:10.1002/aps3.70049

41. Marques SM, Borko S, Vavra O, Dvorsky J, Kohout P, Kabourek P, Hejtmanek L,
Damborsky J, Bednar D. Caver Web 2.0: analysis of tunnels and ligand transport in dynamic
ensembles of proteins. Nucleic Acids Research. 2025;53(W1):W132–W142.
doi:10.1093/nar/gkaf399

42. Masalia RR, Temme AA, Torralba N de leon, Burke JM. Multiple genomic regions
influence root morphology and seedling growth in cultivated sunflower (Helianthus annuus
L.) under well-watered and water-limited conditions. PLOS ONE. 2018;13(9):e0204279.
doi:10.1371/journal.pone.0204279

43. McInerney JO. 1 Genomic Perplexity and the Evolution of Context-Dependent 2
Function.

44. Muhich AJ, Singh R, Tom C, Caseys C, Srinivas K, Faieta L, Grabbe B, Kliebenstein DJ.
Host-Botrytis co-transcriptomics reveals finely tuned interactions with closely related
legumes. 2026 [accessed 2026 Apr 2]:2026.01.20.700702.
https://www.biorxiv.org/content/10.64898/2026.01.20.700702v1.
doi:10.64898/2026.01.20.700702

45. Naegele RP, DeLong J, Alzohairy SA, Saito S, Abdelsamad N, Miles TD. Population
Genetic Analyses of Botrytis cinerea Isolates From Michigan Vineyards Using a High-
Throughput Marker System Approach. Frontiers in Microbiology. 2021;12:660874.
doi:10.3389/fmicb.2021.660874

46. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P,
De Bakker PIW, Daly MJ, et al. PLINK: A Tool Set for Whole-Genome Association and
Population-Based Linkage Analyses. The American Journal of Human Genetics.
2007;81(3):559–575. doi:10.1086/519795

47. Qian C, Lin Y, Zhu C, Liu H, Zhao J, Li X, Ren D, Li J, Wang Y, Xu L, et al. Dynamic hyphal
specialization remodeling enables Botrytis cinerea to overcome diverse host defenses
during infection. Physiological and Molecular Plant Pathology. 2025 Sept:102966.
doi:10.1016/j.pmpp.2025.102966

48. R Core Team. R: A Language and Environment for Statistical Computing. Vienna,
Austria: R Foundation for Statistical Computing; 2025. https://www.R-project.org/

49. Rodriguez-Morelos VH, Calonne-Salmon M, Declerck S. Anastomosis within and
between networks of Rhizophagus irregularis is differentially influenced by fungicides.
Mycorrhiza. 2023;33(1–2):15–21. doi:10.1007/s00572-023-01103-x

50. Rowe HC, Kliebenstein DJ. Complex Genetics Control Natural Variation in *Arabidopsis*
thaliana Resistance to *Botrytis cinerea*. Genetics. 2008;180(4):2237–2250.
doi:10.1534/genetics.108.091439

51. Rowe HC, Kliebenstein DJ. Elevated Genetic Variation Within Virulence-Associated
Botrytis cinerea Polygalacturonase Loci. Molecular Plant-Microbe Interactions®.
2007;20(9):1126–1137. doi:10.1094/MPMI-20-9-1126

52. Schumacher J, Simon A, Cohrs KC, Viaud M, Tudzynski P. The Transcription Factor
BcLTF1 Regulates Virulence and Light Responses in the Necrotrophic Plant Pathogen
Botrytis cinerea. PLOS Genetics. 2014;10(1):e1004040. doi:10.1371/journal.pgen.1004040

53. Shen Q, Li H, Wang Q, Wang J, Ge J, Yang X, Wang X, Li X, Zhang Y, Zhang R, et al.
Alleviating Effects of Linalool Fumigation on Botrytis cinerea Infections in Postharvest
Tomato Fruits. Horticulturae. 2022;8(11):1074. doi:10.3390/horticulturae8111074

54. Singh R, Caseys C, Kliebenstein DJ. Genetic and molecular landscapes of the generalist
phytopathogen Botrytis cinerea. Molecular Plant Pathology. 2024;25(1):e13404.
doi:10.1111/mpp.13404

55. Soltis NE, Atwell S, Shi G, Fordyce R, Gwinner R, Gao D, Shafi A, Kliebenstein DJ.
Interactions of Tomato and *Botrytis cinerea* Genetic Diversity: Parsing the Contributions of
Host Differentiation, Domestication, and Pathogen Variation. The Plant Cell.
2019;31(2):502–519. doi:10.1105/tpc.18.00857

56. Spanner R, Taliadoros D, Richards J, Rivera-Varas V, Neubauer J, Natwick M, Hamilton
O, Vaghefi N, Pethybridge S, Secor GA, et al. Genome-Wide Association and Selective
Sweep Studies Reveal the Complex Genetic Architecture of DMI Fungicide Resistance in

Cercospora beticola. Genome Biology and Evolution. 2021;13(9):evab209. doi:10.1093/gbe/evab209

57. Tang M, Wang Y, Wang K, Zhou Y, Zhao E, Zhang H, Zhang M, Yu H, Zhao X, Li G. Codon Optimization Enables the Geneticin Resistance Gene to Be Applied Efficiently to the Genetic Manipulation of the Plant Pathogenic Fungus Botrytis cinerea. Plants. 2024;13(2):324. doi:10.3390/plants13020324

58. Taniguchi S, Hosokawa-Shinonaga Y, Tamaoki D, Yamada S, Akimitsu K, Gomi K. Jasmonate induction of the monoterpene linalool confers resistance to rice bacterial blight and its biosynthesis is regulated by JAZ protein in rice. Plant, Cell & Environment. 2014;37(2):451–461. doi:10.1111/pce.12169

59. Thomas G, Rusman Q, Morrison WR, Magalhães DM, Dowell JA, Ngumbi E, Osei-Owusu J, Kansman J, Gaffke A, Pagadala Damodaram KJ, et al. Deciphering Plant-Insect-Microorganism Signals for Sustainable Crop Production. Biomolecules. 2023;13(6):997. doi:10.3390/biom13060997

60. Van Kan JAL, Stassen JHM, Mosbach A, Van Der Lee TAJ, Faino L, Farmer AD, Papasotiriou DG, Zhou S, Seidl MF, Cottam E, et al. A gapless genome sequence of the fungus *Botrytis cinerea*. Molecular Plant Pathology. 2017;18(1):75–89. doi:10.1111/mpp.12384

61. Vela-Corcía D, Aditya Srivastava D, Dafa-Berger A, Rotem N, Barda O, Levy M. MFS transporter from Botrytis cinerea provides tolerance to glucosinolate-breakdown products and is required for pathogenicity. Nature Communications. 2019;10(1):2886. doi:10.1038/s41467-019-10860-3

62. Vivaldo G, Masi E, Taiti C, Caldarelli G, Mancuso S. The network of plants volatile organic compounds. Scientific Reports. 2017;7(1):11050. doi:10.1038/s41598-017-10975-x

63. Westrick NM, Smith DL, Kabbage M. Disarming the Host: Detoxification of Plant Defense Compounds During Fungal Necrotrophy. Frontiers in Plant Science. 2021 [accessed 2025 Dec 17];12. https://www.frontiersin.org/journals/plant-science/articles/10.3389/fpls.2021.651716/full. doi:10.3389/fpls.2021.651716

64. Wu Z, Bi Y, Zhang J, Gao T, Li X, Hao J, Li G, Liu P, Liu X. Multidrug resistance of Botrytis cinerea associated with its adaptation to plant secondary metabolites. mBio. 15(2):e02237-23. doi:10.1128/mbio.02237-23

65. Xu Y, Tong Z, Zhang X, Wang Y, Fang W, Li L, Luo Z. Unveiling the Mechanisms for the Plant Volatile Organic Compound Linalool To Control Gray Mold on Strawberry Fruits. Journal of Agricultural and Food Chemistry. 2019;67(33):9265–9276. doi:10.1021/acs.jafc.9b03103

66. You Y, Suraj HM, Matz L, Herrera Valderrama AL, Ruigrok P, Shi-Kunne X, Pieterse FPJ,
Oostlander A, Beenen HG, Chavarro-Carrero EA, et al. Botrytis cinerea combines four
molecular strategies to tolerate membrane-permeating plant compounds and to increase
virulence. Nature Communications. 2024;15(1):6448. doi:10.1038/s41467-024-50748-5

67. Zheng W, Wuyun Q, Li Y, Liu Q, Zhou X, Peng C, Zhu Y, Freddolino L, Zhang Y. Deep-
learning-based single-domain and multidomain protein structure prediction with D-I-
TASSER. Nature Biotechnology. 2025 May 23:1–13. doi:10.1038/s41587-025-02654-4

68. Zhou X, Stephens M. Genome-wide efficient mixed-model analysis for association
studies. Nature Genetics. 2012;44(7):821–824. doi:10.1038/ng.2310

800