Structural principles of transcriptional collisions

John W. Watters[1,2], Andreas U. Mueller[3], Xiangwu Ju[1], Stephany J. Chuquimarca[1], Hannah J. Ye[1], Seth A. Darst[3], Gregory M. Alushin[2,*], Shixin Liu[1,*,#]

[1]Laboratory of Nanoscale Biophysics and Biochemistry, The Rockefeller University, New York, NY, USA
[2]Laboratory of Structural Biophysics and Mechanobiology, The Rockefeller University, New York, NY, USA
[3]Laboratory of Molecular Biophysics, The Rockefeller University, New York, NY, USA
*Correspondence: galushin@rockefeller.edu & shixinliu@rockefeller.edu
#Lead contact

Abstract

RNA polymerase (RNAP) must navigate crowded genomic tracks to robustly produce transcripts. DNA-bound proteins act as transcription-impeding roadblocks, which are eventually overcome by RNAPs through unclear mechanisms. Here, we use cryo-electron microscopy to visualize actively transcribing *E. coli* RNAP upon collision with an inactivated restriction enzyme (EcoRI*) or with another converging RNAP. Both collisions induce RNAP backtracking into an inactive swiveled state. Swiveling is coupled to DNA deformation through a characteristic structural landscape, mediating RNAP pause stabilization across distinct collision geometries. EcoRI* roadblock bypass efficiency is impacted by factors that modulate RNAP swiveling, backtracking rescue, and roadblock stability. In comparison, head-on RNAP-RNAP collisions feature substantial heterogeneity suggestive of sustained dynamics, with variable inter-RNAP distances that are modulated by nascent transcript hairpins which position bidirectional termination sites. By resolving, to our knowledge, novel structures of actively transcribing RNAP undergoing collisions, we provide a mechanistic framework for interpreting mechanical conflicts during transcription.

Introduction

The genome has been described as a molecular highway where the macromolecular machines that carry out fundamental central dogma processes (e.g. replication and transcription) function in a densely crowded environment[1]. This milieu includes a variety of static roadblocks (e.g. DNA-binding proteins and DNA lesions), signals (regulatory sequences), and other dynamic machines operating on the same substrate, such that collisions *in vivo* are inevitable[1,2]. In addition to interfering with machine function[2], collisions also provide potential for regulation, for example, by defining genomic boundaries or coupling the activities of multiple machines in tandem[3,4], thereby enhancing biological robustness.

Recent studies have emphasized that multiple categories of collisions between DNA machines can have both positive and negative biological consequences. For example, head-on collisions between replication and transcription machinery lead to increased R-loop formation, which is detrimental to genome stability[5–7]. Conversely, local chromatin looping into topologically associated domains is driven by cohesin-CTCF collisions, mediating functional genome organization[8]. Furthermore, co-directional collisions between machines can enhance their processivity, such as translation-transcription coupling in bacteria, which prevents transcriptional arrest[9–15]. Additionally, we recently showed that sequence-specific pausing of bacterial RNA polymerases (RNAPs) positions RNAP-RNAP head-on collisions, thereby programing transcription termination sites to enhance the production of precise transcripts from convergent genes[16,17]. How collisions both disrupt and enhance machine functions at the protein structural level to produce these diverse outcomes remains unknown.

Because of its central role in many genomic collisions, here we focus on the RNAP as a model machine, which operates as a Brownian rachet and can generate strong forces and torques[18–23]. Previous biophysical studies of RNAP collisions have suggested that upon being halted by roadblocks, the complex can bypass them by two mechanisms: (i) passively waiting for roadblocks to dissociate or (ii) actively consuming chemical energy to dislodge them[24], with the relative contributions of these mechanisms dependent on roadblock identity[15,16,24,25]. Additional factors including transcription elongation factors, co-directional transcribing RNAPs, and applied force have also been shown to help RNAP overcome roadblocks[15,16,24]. While RNAP has been extensively characterized by single-molecule biophysics, structural biology, and biochemistry, it remains unclear how roadblocks effectively stall actively transcribing RNAPs and how RNAPs in turn bypass roadblocks. Collisions occur under non-equilibrium, actively energy-dissipating conditions are technically challenging to recapitulate for structural studies. Conceptually, it is interesting to examine how collisions involving a nanoscale machine like RNAP—which operates under the low Reynolds number regime—compare to macroscopic collisions such as a car crash, which produces heterogeneous outcomes in the final motor configuration. However, such microscopic transcriptional collisions have rarely been directly visualized at the atomic level.

In this study, we use cryo-electron microscopy (cryo-EM) to examine actively transcribing *E. coli* RNAPs collided with (i) a static roadblock, namely the catalytically deficient EcoRI E112Q restriction enzyme mutant (hereafter referred to as EcoRI*), and (ii) another RNAP transcribing in the opposite direction, resulting in a head-on collision. By comparing the structural landscapes of these two physiological collision

77　scenarios[24,26,27], we uncover principles for mechanical control of RNAP that can be
78　harnessed to bypass deleterious roadblocks and mediate functional positioning of
79　transcription termination sites.
80
Results
Reconstituting RNAP collision with a stationary roadblock
To capture a collided roadblock complex, we generated a nucleic acid scaffold
containing a pre-formed transcription bubble and loaded the scaffold with an RNAP core
enzyme and a downstream EcoRI* dimer bound to the cognate binding site (5'-GAATTC-
3', K_D ~2.5 fM)[26,27]. Upon ribonucleotide (NTP) addition, RNAP is anticipated to transcribe
for ~20 nucleotides (nt) before encountering the EcoRI* roadblock (**Fig. 1A**). Proper
loading of RNAP and EcoRI* was confirmed via a native gel-shift assay, and the complex
was enriched using gel filtration chromatography (**Fig. 1B, S1**). We used a Cy3-labeled
RNA in the scaffold to evaluate the length of transcription products. In the absence of
EcoRI*, most RNAPs transcribed until the end of the DNA template, generating run-off
transcripts ~70 nt in length (**Fig. 1C**). When EcoRI* was added, shorter RNAs (narrowly
distributed around ~40 nt in length) were observed with minimal run-off products (**Fig.
1C**), suggesting high occupancy of EcoRI* and efficient blockage of RNAP readthrough.
We next used the RNAP-EcoRI* assembly supplemented with 1.5 mM Fos-
Choline-8 detergent for cryo-EM sample preparation. To induce collision, 0.5 mM NTPs
were added and incubated for 60 seconds prior to plunge-freezing (**Fig. 1B**). After data
collection, we performed 3D classification to enrich for particles that contained both RNAP
and EcoRI*. Unexpectedly, we obtained a single three-dimensional (3D) reconstruction
of the RNAP-EcoRI* collided complex with a global Fourier shell correlation (FSC)
resolution of 2.68 Å, indicating that we captured a single metastable state instead of
multiple states along the collision pathway (**Fig. S2, S3A-D**). The resolution of the
downstream EcoRI* roadblock was significantly lower than the RNAP core, suggesting
higher flexibility[26,28,29]. We used 3DFlex in cryoSPARC to improve the resolution of the
flexible downstream DNA and EcoRI*, producing a reconstruction with a global resolution
of 2.83 Å that allowed us to build an atomic model of the entire collided complex (**Fig. 1D,
S3E-H, Table S2**)[30].
108
EcoRI-collided RNAP is backtracked*
We envisioned three possible scenarios for the behavior of RNAP upon colliding
with a roadblock: (1) it could advance to the final position along the DNA template free
from steric hindrance and remain there; (2) it could halt before this final position due to
topological constraints imposed by the DNA; or (3) it could recoil away from the final
position to an energetically more favorable site, due to the ability of RNAP to backtrack
along DNA[31,32], as suggested by previous studies of RNAP transcription in the presence
of an EcoRI* roadblock[15,24]. To assess RNAP backtracking in our system, we took
advantage of the fact that backtracked RNA is sensitive to GreA/B-stimulated
endonucleolytic cleavage[33]. In the absence of GreA/B, the majority of RNA products were
43 nt long (**Fig. 2A),** while in the presence of GreB and, to a lesser extent, GreA, the
distribution of bands was shifted towards shorter lengths. In the presence of both factors,
most RNA products were 39-41 nt in length (**Fig. 2A, S4A, B**), suggesting that the collided

RNAP is backtracked 2-4 nt in our reconstitution system, in agreement with previous
results[15].

During each nucleotide addition cycle, conformational changes in the bridge helix
and trigger loop—structural elements in the RNAP active site—are coordinated to
translocate the DNA register, facilitate nucleotide binding, and accelerate the catalysis of
nucleotide incorporation[34]. We next examined the RNAP active site in our cryo-EM map
and found that it resembles previously reported structures of backtracked RNAP (PDB:
6RI9, 6RIP), featuring an unfolded trigger loop and a straight bridge helix (**Fig. 2B**)[35,36].
Moreover, we observed densities in the secondary channel consistent with extruded,
backtracked RNA (**Fig. 2B**). Two backtracked nucleotides are resolved; the lack of
density from additional backtracked nucleotides is likely attributable to higher levels of
conformational flexibility, as has been observed in previous cryo-EM structures of
backtracked RNAP[35].

These biochemical and structural data suggest that the RNAP transcribes 19 nt of
RNA from its initial position, but it cannot stably reside at that position, potentially due to
steric clashes with the roadblock. After collision, RNAP backtracks by as many as 4 nt,
allowing it to re-equilibrate at a more favorable position. To further evaluate this
hypothesis, we computationally modeled the collision pathway by moving RNAP
structures to each nucleotide position along the collision axis (up to 44 nt, one nt past the
final position transcribed by RNAP in the EcoRI* collision scenario). We then measured
the overlapping volume enclosed by the solvent-excluded surfaces (SES) of RNAP and
EcoRI* as a proxy of steric overlap (**Fig. S4C, D**). The RNAP conformation was modeled
based on a previously reported elongation complex (EC) structure (PDB: 6RH3)[35]. When
the polymerase was placed less than two bases away from the collision interface, the
overlapping SES volume[37] increases substantially, consistent with clashes occurring.
While this analysis is limited by the assumption that the RNAP retains the same
conformation as it approaches the roadblock, it nonetheless indicates that direct steric
hindrance from EcoRI* is a plausible explanation for why RNAP backtracks away from
the collision site.

EcoRI-collided RNAP is swiveled*
We next examined the RNAP structural modules that influence its catalytic activity.
We find that in the collided RNAP-EcoRI* complex, RNAP is in a 'swiveled' conformation,
in which a swivel module (β subunit: AA 1241-1341; β' subunit: AA 1-342, 369-420, 787-
930, 1135-1375) is rotated relative to the RNAP structural core. When RNAP is sufficiently
swiveled, the bridge helix kinks and the trigger loop is unable to fold into the correct
conformation to mediate catalysis, stabilizing long-lived transcriptional pausing[38–40]. In the
active elongation complex (EC; PDB: 6rh3)[35], RNAP is in a non-swiveled state, whereas
pronounced swiveling (~4.5-6° swivel module rotation relative to the active EC) is
observed in various paused and backtracked ECs[35,36,39,41].

In the collided RNAP-EcoRI* complex, the RNAP adopts a moderately swiveled
state with a swivel module rotation of 2.8° relative to the active EC, a lower magnitude
rotation than observed in a previously reported hairpin-stabilized paused EC (4.4°, PDB:
6ASX)[39] (**Fig. 2C, D**). We then analyzed 3DFlex variability trajectories to assess swiveling
heterogeneity in our data. We found that the swivel angle ranged between 2.6° and 3.8°
across frames (**Fig. 2D**)[30]. In contrast, previously resolved backtracked RNAPs populate

either swiveled (3.8°, 63% of population, PDB: 6RI9) or non-swiveled states (0.7°, 37%
of population, PDB: 6RIP), but not in between (**Fig. 2D**)[35]. This result indicates that the
collided RNAP is constrained within a moderately swiveled state, suggesting a novel
configuration imposed by the physical collision.
EcoRI binding to DNA by itself is known to distort DNA from its canonical B-form
at the cognate sequence, introducing multiple kinks that result in a net 25° bending of the
DNA and concomitant increased accessibility of bases in the major groove[28,42]. Given the
ability of DNA to serve as a mechanical transducer, we hypothesized that DNA distortions
could be transmitted to RNAP as a mechanism to indicate the presence of a roadblock.
To investigate coupling between DNA deformation and RNAP structural remodeling, we
once again analyzed 3DFlex trajectories. To assess changes in DNA positioning relative
to the active EC (PDB: 6RH3), a reasonable proxy for the spatial orientation of DNA
relative to the RNAP structural core during catalysis, we established a frame of reference
by subjecting the downstream nucleotides to principal component analysis (PCA). The
displacement of downstream DNA in the RNAP-EcoRI* complex was then measured from
projections onto the PC1 and PC2 orthoplanes after superimposing the structural cores
of the RNAPs. Interestingly, we observed that specific DNA distortions downstream from
the RNAP active site are correlated with changes in the RNAP swiveling angle (**Fig. 2E,
S5, Movie S1**). As the downstream DNA is displaced more along PC2, the magnitude of
RNAP swiveling increases (**Fig. 2F, S5**). This observation suggests that modulation of
DNA substrate geometry can couple inhibitory RNAP rearrangements to the presence of
roadblocks at collision sites.
190
Factors that modulate RNAP backtracking / swiveling impact bypass efficiency
Because RNAP is backtracked and swiveled in the RNAP-EcoRI* complex, we
next examined whether the outcome of this collision (persistent RNAP halting versus
roadblock bypass) can be biased by factors that modulate these aspects of the RNAP
structural landscape. We implemented a bulk biochemical assay to quantitatively evaluate
roadblocking efficiency (E_{RB}), conducting transcription reactions at 37 °C for 30 minutes
with increasing salt concentrations (50 to 500 mM KOAc) to permit measurable levels of
bypass. As anticipated, higher levels of roadblock bypass (lower E_{RB}) were observed at
higher salt concentrations, presumably due to destabilization of the EcoRI*-DNA
interaction (**Fig. 3A, B**). Addition of NusA (a pro-swiveling factor) and NusG (an anti-
swiveling, anti-backtracking factor)[36,43,44] led to decreased and increased bypass,
respectively (normalized E_{RB} = 1.2 ± 0.1 for NusA and 0.8 ± 0.1 for NusG in the presence
of EcoRI*, compared to 1.0 ± 0.1 for EcoRI* alone) (**Fig. 3C**). These results suggest that
biochemically modulating RNAP's swiveling propensity (e.g. through Nus factor
engagement) changes its capacity to overcome roadblocks. Strikingly, the presence of
the transcript cleavage factor GreB, which rescues backtracked RNAP[33], dramatically
reduced roadblocking efficiency (normalized E_{RB} = 0.06 ± 0.03) (**Fig. 3B, C**). This near-
complete roadblock bypass can be explained by repeated collisions between RNAP and
EcoRI* made possible by GreB-induced transcription restart after RNAP backtracking,
consistent with a recently proposed 'battering' model[24].
211
Roadblock structural remodeling facilitates bypass

Having observed substantial structural alterations in RNAP upon collision, we next
examined whether the structure of the EcoRI* roadblock is also impacted. EcoRI*
engages DNA as a homodimer[28,42,45]. The protein's DNA-bound conformation is overall
highly similar between the RNAP-EcoRI* collided complex and a previous crystal
structure in the absence of RNAP (PDB: 1ERI)[42], with the notable exception for the N-
terminal helix (AA 20-31) of the protomer proximal to the collision interface, which flips
~150°, orienting it away from RNAP (**Fig. 3D, Movie S2**). This rearrangement was not
observed in the distal EcoRI* protomer, suggesting it is specifically linked to contact with
RNAP (**Fig. 3D**). We speculate that it occurs at the collision site to reduce steric clashes
with RNAP, becoming kinetically trapped as the RNAP backtracks away. Additionally, this
rearrangement repositions EcoRI*'s N-terminal extension (NTE) residues (5K, 9R, 15K)
in an orientation where they can access DNA (**Fig. 3D**). These amino acids have been
previously shown to enhance DNA binding in biochemical assays, presumably by
mediating electrostatic interactions[26,28,42,45].

In the EcoRI* crystal structure, the N-terminal helix forms hydrophobic contacts
with the protein's structural core[42]. As a protein's potency as a roadblock intuitively
depends on its mechanical resilience, we posited that destabilizing the overall folded
structure of EcoRI* would compromise its roadblocking efficacy. To test this hypothesis,
we introduced alanine substitutions at the positions of hydrophobic residues within the
EcoRI* core (**Fig. 3E**), distal from the DNA binding interface, and assessed their impact
on E_{RB} using our roadblock assay. We found that all point mutations (with the exception
of V166A) led to significantly increased RNAP bypass (decreased E_{RB}) (**Fig. 3F**). Notably,
the two mutations that caused the largest magnitude changes, I21A and F252A, are
internally facing residues adjacent to the predicted collision interface that are likely to
mediate a stabilizing contact between the N-terminal helix and the EcoRI* core (**Fig. 3E,
F**). These results suggest that the structural stability of a roadblock protein—independent
from its affinity to DNA—is linked to its capacity to resist physical impacts from motors
like RNAP; correspondingly, mechanical destabilization of roadblocks can enhance
bypass.

Reconstituting head-on collisions between two RNAPs

To explore the generality of RNAP's collision response, we next investigated
another physiological scenario, when an RNAP runs into another active motor complex
rather than a stationary roadblock. In previous studies, we demonstrated that head-on
collisions between two RNAPs produce long-lived complexes on DNA[16], constituting a
prevalent mechanism for transcription termination of convergent genes in *E. coli*[17]. To
reconstitute a head-on RNAP-RNAP collision in a format suitable for cryo-EM structural
interrogation, we constructed a DNA template with two convergent promoters (**Fig. 4A**).
Each promoter was bound by a σ^{70}-RNAP holoenzyme that was walked to the +18/+20
position by omitting one of the four nucleotides (CTP), followed by purification via size
exclusion chromatography. Transcription was then restarted with addition of 1 mM NTPs
to induce a collision (**Fig. 4B**). We also introduced a native sequence from the *E. coli*
genomic region between the *YnaJ-UspE* gene pair into the template, which encodes an
RNA hairpin containing an 8-bp stem. We have previously shown that this sequence is
important for positioning the collision site between a pair of RNAPs, mediating precise
termination of convergent transcription[16,17].

After biochemically confirming the formation of a head-on RNAP-RNAP collided
complex (**Fig. 4C**), we employed this DNA template (once again supplemented with 1.5
mM Fos-Choline-8 detergent) for cryo-EM sample preparation. Initial 2D class averages
displayed features consistent with two RNAPs in close proximity (**Fig. S6**). Unlike the
RNAP-EcoRI* collided complex, 3D classification of the RNAP-RNAP collided complex
revealed substantial heterogeneity. Notably, the distance between the two RNAP
molecules varied, with the best resolved classes featuring 26, 29, and 31 bp of DNA
between the active sites of the polymerases (**Fig. 4D-F**). Because of the ~10.5-bp helical
pitch of B-form DNA, these three configurations all feature approximate C2 symmetry, a
plausible explanation for why both RNAPs could be effectively aligned and averaged
when separated by these particular distances. We exploited this C2 pseudosymmetry to
perform symmetry expansion and analyzed all RNAP protomers from a given inter-RNAP
distance class within a single frame of reference, facilitating high-resolution refinement
(26 bp, 2.65 Å; 29 bp, 2.87 Å; 31 bp, 2.62 Å global resolution) (**Fig. S7**).
We could not interpret nucleic acid sequences in the density maps (which were
modeled as homopolymers), presumably due to heterogeneity in where collisions occur
along the DNA template. An additional potential source of heterogeneity in our
preparation is the presence of σ^{70} to promote initiation, but the 31-bp class was the only
one that featured weak density (only visible at low map contour levels) for the σ^{70} domain
2 bound to RNAP. Therefore, the variable inter-RNAP spacing we recovered through
classification likely reflects inherent heterogeneity in RNAP-RNAP collisions. We once
again employed 3DFlex to further improve the resolution of these classes and analyzed
their conformational landscapes (**Fig. S7**).
The RNAP-RNAP interface varies considerably among the three classes due to
the differential translational and rotational relative positions of the RNAP pair on the
helical DNA template. Nevertheless, in all three classes, the clamp domain of one RNAP
was engaged with different domains of the other collided RNAP, notably the β lobe/SI2
domain (26 bp class), SI3 domain (29 bp class), and jaw domain (31 bp class) (**Fig. 4G-**
I). We reasoned this could influence RNAP swiveling, as the clamp domain is a
component of the swivel module. Indeed, all of the RNAPs in head-on collided complexes
adopted a swiveled conformation (26 bp, 3.9°; 29 bp, 2.3°; 31 bp, 4.0° relative to PDB:
6RH3) (**Fig. 4J-L**). Additionally, despite sequence ambiguity, all the RNAPs showed clear
density for backtracked RNA in their secondary channel regardless of inter-RNAP
distance (**Fig. 4M-O**). Thus, despite substantial differences at the collision interface,
RNAP-RNAP collisions populate a similar conformational landscape as the RNAP-EcoRI*
collision, suggesting a fundamental RNAP response to roadblocks.
295
RNA hairpin impacts collision positioning and stability
To evaluate the bulk transcriptional output of the paired RNAPs in this system,
which also serves as a proxy for their positioning along the DNA, we performed in vitro
SEnd-seq (simultaneous 5' and 3' end RNA sequencing) to systematically profile end-to-
end RNA products[17,46]. End-biotinylated DNA templates were used in these experiments
to pull down collided complexes, enriching for RNAs that remained bound to RNAP. We
also constructed a template without the hairpin-encoding sequence for comparison. The
sequenced RNAs displayed homogeneous 5' ends (corresponding to initiation from either
promoter) and more heterogeneous 3' ends (**Fig. 5A, B, S8**), as anticipated given the

stochastic nature of RNAP restart[47], which will impact the occurrence and location of
collisions. Samples including the hairpin-encoding sequence in the template featured a
pronounced drop in the RNA length profile at the predicted hairpin sequence boundaries
(**Fig. 5A**), resulting in significantly shorter transcripts for both the forward and reverse
directions compared to the no-hairpin template (**Fig. 5C, S8**). These results indicate that
the hairpin fosters consistent positioning of RNAP-RNAP collisions, mitigating
heterogeneity attributable to the stochastic mechanochemistry of these nanoscale
machines.
We observed that the complex disassembles when incubated for longer time
periods, which could occur when one RNAP is able to overcome the other and push it off
the DNA template. Given the observation that all RNAPs in the collided complex were
backtracked, we reasoned that the hairpin could serve as a barrier against long-range
backtracking, thereby stabilizing the complex. To test this hypothesis, we performed a
time course assay to measure the half-life of the collided complex by halting the
transcription reaction with an EDTA quench at different timepoints post NTP addition.
While the collided complex on the template with the hairpin remained stable for hours ($T_{1/2}$
= 150 ± 40 min), the collided complex on the template without the hairpin disassembled
within minutes ($T_{1/2}$ = 6 ± 3 min) (**Fig. 5D, E**), confirming the hairpin's contribution to
complex stability.
We next sought to examine how the presence of a hairpin in the nascent transcript,
which would form if one of the RNAPs has transcribed past the hairpin-encoding
sequence before collision, influences the structure of the collided complex. No clear RNA
hairpin density was observed in the exit RNA channel for any of the three RNAP-RNAP
complex classes, indicative of flexibility or variable occupancy. We then reprocessed the
data, re-extracting particles using a smaller box size encompassing one RNAP followed
by focused classification at the RNA exit channel (**Fig. S9**). In a single class obtained
using this approach (representing ~7% of the data), we observed hairpin density (**Fig.
5F**). RNAP was still backtracked and swiveled (~3.4°) in this class, suggesting that the
presence of a well-ordered hairpin does not grossly alter RNAP's conformational
response to collision (**Fig. 5G**). To assess the hairpin's impact on collision geometry, we
recentered our particle picks to capture the other collided polymerase, allowing us to
analyze the distribution of inter-RNAP spacings for particles contributing to the hairpin-
containing class versus the distribution for the entire collided population. In the hairpin-
containing particles, we found the class with the largest separation between RNAPs (31
bp) was mildly enriched (~1.3-fold increase), while the classes with shorter inter-RNAP
spacing were less populated, with a lower 26-bp class abundance (~1.5-fold decrease)
and a complete absence of the 29-bp class (**Fig. 5H**). These data are consistent with the
notion that the hairpin reduces the heterogeneity of RNAP-RNAP collisions (**Fig. 5H**,
p<0.001), potentially by restricting the dynamics of the hairpin-associated RNAP and
maintaining its position on DNA.
345
DNA deformation-RNAP swivel coupling depends on inter-RNAP distance
As a physical connection between RNAP and roadblock through the template DNA
is a common element of transcriptional collisions with distinct composition and
geometries, we reasoned this linkage could mediate characteristic RNAP remodeling. To
test this hypothesis, we examined the coupling between DNA deformation and RNAP

swiveling across RNAP-RNAP collision configurations. Swiveling was present in all three
head-on collision orientations. However, interestingly, the degree of RNAP swiveling
varied between the different classes in a manner that was non-monotonic with inter-RNAP
distance, as the 29-bp spaced class featuring a substantially lower swivel angle in the
consensus class (~2.3°) than the 26-bp (~3.9°) and 31-bp (~4.0°) classes (**Fig. 4J-L**). We
speculate this variation is linked to the differential interfaces between the RNAPs, which
likely impact the local mechanical environment. We also found that the degree of DNA
deformation downstream of the collided RNAP varied among classes (**Fig. 6A**). Finally,
we examined whether the RNAP swivel angle and the DNA deformation remain correlated
across different collision scenarios. When compared alongside previous RNAP structures
representative of active EC, backtracked, and swiveled complexes (PDB: 6RH3, 6ASX,
6RI9, 6RIP)[35,39], changes in the DNA PC1-axis did not correlate with changes in swiveling
while changes in the DNA PC2-axis displayed a strong correlation, with the displacement
increasing linearly with the magnitude of the RNAP swivel angle (**Fig. 6B, C**). Additionally,
the extent of PC2 displacement within individual 3DFlex variability trajectories also
correlates with the magnitude of swivel angle change (**Fig. 6D, S10, Movie S1, 3**)[30].
These analyses highlight that while RNAP's conformational landscape varies across
collisions, likely due to variations in the details of the RNAP-roadblock interface restricting
the accessible DNA bending landscape, DNA deformation serves as a common
mechanism to transduce collisions into RNAP swiveling and inactivation.
371
372

Discussion
In this study, we visualized the structural dynamics of RNAP upon collisions with
both a passive (EcoRI*) and an active (another RNAP) roadblock under non-equilibrium,
actively transcribing conditions. Despite variability in the composition and geometry of
these collisions, our results reveal a characteristic RNAP response consisting of
backtracking and swiveling to enter a transcriptionally inactive state (**Fig. 7A**). Our data
highlight how shared mechanical elements across collisions, including roadblock steric
hindrance and DNA deformation, can coordinate this common RNAP response
irrespective of the detailed stereochemistry of a collision.
As we did not directly visualize RNAP at its site of closest contact with roadblocks
(indicated from transcript lengths), we cannot comment on how backtracking is triggered.
We nevertheless speculate that roadblock steric hindrance ultimately limits forward
progress of RNAP, triggering backtracking to a metastable position where clashes are
moderated. At this position, RNAP dynamics can be modulated by coupling between DNA
template deformation and RNAP swiveling, facilitating the maintenance of an inactivated
state. Consistent with a previously proposed battering model[15,24], we find that factors
which reduce RNAP swiveling and enhance transcription restart after backtracking
enhance roadblock bypass (**Fig. 3B-C, 7B**). Our structures also rationalize the previous
observation that applying small opposing forces (~0.2 pN) counterintuitively increases the
rate of EcoRI* bypass by RNAP[24], as this likely promotes RNAP un-swiveling by altering
the downstream DNA deformation profile to more efficiently restart elongation after
backtracked RNA cleavage, leading to eventual bypass. We furthermore uncovered an
additional layer of the battering mechanism, wherein RNAP can physically deform
roadblocks to eventually displace them (**Fig. 3D-F, 7B**). This suggests mechanical

destabilization of deleterious roadblocks plays an instrumental role in their clearance from
the genome.
Our observation that RNAPs in head-on collisions are also backtracked and
swiveled, but feature variable inter-RNAP distances, is compatible with a scenario
featuring intermittent, ping-ponging restart dynamics between the two RNAPs. We
propose that when one RNAP stochastically restarts, it backtracks upon colliding with the
other paused RNAP. This clears the path for either RNAP to once again restart,
generating cycles of randomly alternating transcription and backtracking (**Fig. 7C**). This
model can explain our previous observation that both RNAPs can transcribe through
bidirectional convergent terminators like YnaJ-UspE before termination[17], which is
otherwise incompatible with a static collision. Our analysis also provides a mechanistic
explanation for the positioning effect of the RNA hairpin when bound to RNAP, as we find
its presence is correlated with increased complex stability (**Fig. 5D-E**) and reduced
heterogeneity in inter-RNAP spacing (**Fig. 5H**). This suggests that the hairpin likely
suppresses the motions of the engaged RNAP, thereby stabilizing it in a particular region
along the DNA template and attenuating the variability in transcript length. We did still
observe substantial variability in transcripts even in the presence of the hairpin in our
minimal reconstitution system (**Fig. 5A-C**), suggesting additional cellular factors may
further enhance the robustness of this mechanism, an important topic for future studies.
This work establishes the feasibility of scrutinizing genomic collisions using cryo-
EM, providing a path towards visualizing physiological scenarios of increasing complexity.
An important consideration for future studies is the topology of the template DNA, which
may be addressable by employing closed templates with varying degrees of
supercoiling[48,50]. Additionally, collisions featuring additional machines, such as co-
directionally transcribing RNAP and translating ribosomes (both of which can promote
roadblock bypass[3,10,15]) or head-on/co-directional DNA replication forks (replication-
transcription conflict as a primary source of genome instability[4–6]) are likely accessible by
similar approaches. Such studies will be critical for a mechanistic understanding of how
mechanical conflicts are managed across the crowded genome and functionally
harnessed to enhance the robustness of gene expression.

Methods

Protein purification
E. coli RNA Polymerase core
RNAP was expressed and purified as described previously[49], with an additional step of 3C cleavage of the β'-3C-His10 tag and a reverse Nickel-IMAC (immobilized metal affinity chromatography) column to isolate RNAP core without the His10 tag.

Maltose Binding Protein (MBP)-EcoRI (E112Q) and additional mutants*
EcoRI* E112Q (a gift from Dr. Ruben Gonzalez's lab) was cloned into a pET21 vector with an N-terminal His10-MBP-3C sequence using Gibson assembly. The plasmid was transformed into competent Eco BL21(DE3) by heat shock. The cells were grown in the presence of 50 µg/mL kanamycin to an optical density (OD) at 600 nm of ~0.5-0.6 in a 37 °C shaker. Expression was induced with addition of 0.5 mM Isopropyl β-D-1-thiogalactopyranoside (IPTG) at 16 °C overnight. Cells were harvested by centrifugation, resuspended in lysis buffer [20 mM potassium phosphate, 300 mM sodium chloride (NaCl), 1 mM dithiothreitol (DTT), 0.1 mM ethylenediaminetetraacetic acid (EDTA), 5% glycerol (v/v), 0.15% Triton X-100, 1x protease inhibitor cocktail (Sigma-Aldrich), pH 7.5 at room temperature], and flash frozen in liquid nitrogen prior to storage at -80 °C.

Cell pellets were thawed, then lysed with an Avestin Emulsiflex C5 homogenizer. The lysate was then clarified (2x centrifugation steps, 30 minutes each, 48,000 x *g*) and applied to a 5 mL MBPTrap HP column (Cytiva). The column was washed with MBP Buffer A [20 mM potassium phosphate, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 5% glycerol (v/v), 0.15% Triton X-100, pH 7.5 at room temperature] and eluted with MBP Buffer A supplemented with 10 mM maltose. Peak fraction were pooled, diluted to in Heparin Buffer A [20 mM potassium phosphate, 1 mM DTT, 0.1 mM EDTA, 5% glycerol (v/v), 0.15% Triton X-100, pH 7.5 at room temperature] to 50 mM NaCl final concentration and loaded onto a 10 mL Heparin HP (Cytvia) column equilibrated in 10% Heparin Buffer A/90% Heparin Buffer B [20 mM potassium phosphate, 1 M NaCl,1 mM DTT, 0.1 mM EDTA, 5% glycerol (v/v), 0.15% Triton X-100, pH 7.5 at room temperature]. The bound protein was eluted with a gradient of 10-100% Heparin Buffer B over 10 CVs. Peak fractions of protein without nucleic acid contamination were eluted in a peak centered at ~45 mS/cm conductivity and concentrated with a 30 kDa molecular weight cut-off (MWCO) concentrator (Amicon). Concentrated protein was clarified (5 minutes, 21,000 x *g*) before application onto a HiLoad 16/600 Superdex (Cytiva) gel filtration column equilibrated with gel filtration buffer [20 mM potassium phosphate, 300 mM NaCl, 1 mM tris(2-carboxyethyl)phosphine (TCEP), 0.1 mM EDTA, 5% glycerol (v/v), 0.15% Triton X-100, pH 7.5 at room temperature]. Peak fractions corresponding to the EcoRI* dimer were pooled, concentrated in a 30 kDa MWCO concentrator (Amicon), diluted with gel filtration buffer supplemented with 50% glycerol to a final concentration of 20% glycerol, flash frozen in liquid nitrogen, and stored at −80°C until use. Protein concentration was estimated using a Bradford assay[51] (Bio-Rad Protein Assay Dye Reagent Concentrate) with bovine serum albumin standards.

For single point mutants (I21A, L108A, V166A, F252A, I258A), the same protocol was used with the following substitutions: 1 mM DTT was replaced in all buffers with 1 mM TCEP, and a Superdex 200 Increase (Cytiva) column was used for gel filtration instead of a HiLoad 16/600 Superdex (Cytiva) column.

GreA, GreB, NusA, NusG, σ[70]

GreA, GreB, NusA, NusG, and σ[70] were expressed, purified was performed as described
previously[16,49,52,53].

Endonucleolytic cleavage activity of GreA/B was verified as described previously[16].
In brief, a backtracked RNAP EC scaffold (nucleotide sequences in **Table S1**) was formed
with a short (2 nt RNA backtracked) and long backtracked EC (6 nt RNA backtracked).
Oligonucleotides were high-performance liquid chromatography (HPLC) purified (IDT).
Activity assays were performed at 37 °C for 10 minutes by adding GreB or GreA at a final
concentration of 600 nM in the absence or presence of NTPs. The reactions were stopped
by addition of an equal volume of 2x RNA loading buffer and incubation at 90 °C for 2
minutes. Products were separated on a 25% Urea-PAGE gel, followed by fluorescence
visualization on a Typhoon FLA 7000 scanner. Full-length RNA products indicative of
endonucleolytic cleavage and transcriptional restart were enhanced in the presence of
GreA (short backtracked EC) and GreB (long backtracked EC) as expected[33].

As reported previously, NusA and NusG were verified to be active by bulk in vitro
transcription assays in transcription buffer (25 mM Tris-hydrochloride pH 7.5, 150 mM
potassium chloride, 10 mM magnesium chloride, 1mM DTT). In brief, Rho and NusG in
combination were shown to enhance Rho mediated termination efficiency on the Rho-
dependent λtR1terminator[53]. NusA reduced transcriptional readthrough across the ynaJ-
uspE bidirectional terminator[54]. σ[70] was verified to be active in promoting T7A1 and
LambdaPR dependent promoter complex formation and transcription in in vitro
transcription assays in a transcription buffer (25 mM Tris-HCl pH 7.5, 150 mM KCl, 10
mM MgCl$_2$, 1mM DTT)[55].

RNAP-EcoRI* Scaffold Complex Assembly
DNA oligonucleotide sequences (Polyacrylamide gel electrophoresis (PAGE)-purified
from IDT) and a 5' Cy3-labeled RNA oligonucleotide (Dharmacon) were designed using
a previously used scaffold as a starting point (sequences of template, non-template, and
RNA oligos in **Table S1**)[56]. Template DNA and RNA were annealed in a 1:1 ratio in 1x
IDT annealing buffer (100 mM potassium acetate (KOAc); 30 mM HEPES, pH 7.5) for 5
minutes at 95º C, 5 minutes at 75º C, 5 minutes at 45º C, and 40 cycles of 1º C steps (1
minute hold each step, final temperature 4º C) in a thermocycler. RNAP core desalted
into scaffolding buffer (20 mM Tris pH 7.5 at RT, 150 mM potassium glutamate (KGlu), 5
mM magnesium acetate (Mg(OAc)$_2$), 2.5 mM DTT) was then incorporated and incubated
for 15 minutes at RT. 1.5-fold excess of non-template DNA was then added to the
complex mixture and incubated for 10 minutes at RT. 5-fold excess of EcoRI* was then
added to the mixture (this step was excluded for the -EcoRI* controls) and incubated for
10 minutes at RT before addition of 0.1 mg/ml final concentration of heparin sodium salt
to remove non-specifically bound RNAP and EcoRI* proteins. The fully assembled
complex was purified away from incompletely assembled complexes via gel filtration
chromatography using a Superdex 200 Increase column equilibrated with low salt buffer
(20 mM Tris-acetate (Tris-OAc) pH 7.8 at RT, 50 mM KOAc, 5 mM Mg(OAc)$_2$, 2.5 mM
DTT) containing 0.1 mg/ml heparin (Sigma-Aldrich) to prevent re-binding of non-
specifically loaded RNAP/EcoRI* to the DNA template during the purification. Complex

assembly was monitored via native gel electrophoresis on a 5% polyacrylamide 0.5X TBE
gel.
520
RNAP Head-On Complex Assembly
DNA constructs featuring either two RNAP promoters (T7A1 and λPR) or only one
promoter were generated by PCR, purified via gel filtration chromatography on a
Superdex 200 Increase column equilibrated with DNA gel filtration buffer (20 mM Tris pH
8.0 at RT, 200 mM NaCl, 0.1 mM EDTA), and concentrated into Tris Low EDTA buffer
(10 mM Tris pH 8.0 at RT, 0.1 mM EDTA) for long term storage. PCR oligos and full DNA
sequences of the templates are in **Table S1**.
RNAP holoenzyme was formed by incubating σ^{70} with RNAP core for 20 minutes
at 37º C in their purification buffers, using a 3:1 ratio of σ^{70}:RNAP core. The complex was
then desalted into transcription buffer (25 mM Tris-HCl pH 7.5 at RT, 150 mM KCl, 10
mM $MgCl_2$, 1mM DTT), and its concentration measured using A280 and its calculated
extinction coefficient. To form open promoter complexes, 4-fold excess of holoenzyme
was added to the DNA template (final concentration ~1-2 µM DNA) and incubated for 15
minutes at 37º C. Non-specifically bound RNAPs were removed by incubation with 0.1
mg/mL heparin for 10 minutes at 37º C. 0.5 mM ATP, UTP, and GTP were then added to
the reaction mixture and incubated for 15 minutes at 37º C to walk the promoter bound
RNAPs out to the dual CTP nucleotide deprivation stall site. The fully assembled complex
was purified away from incompletely assembled complexes via gel filtration
chromatography using a Superdex 200 Increase (Cytiva) column equilibrated with low
salt buffer (20 mM Tris-OAc pH 7.8 at RT, 50 mM KOAc, 5 mM $Mg(OAc)_2$, 2.5 mM DTT)
containing 0.1 mg/ml heparin to prevent re-binding of non-specifically loaded RNAP to
the DNA template during the purification. Complex assembly was monitored via native
gel electrophoresis on a 2% agarose 0.5X TBE gel.
544
Cryo-EM
Sample Preparation
After gel filtration, the complex was concentrated using Amicon 100 kDa concentrators
and washed 7x (450 µL added per wash) into low salt buffer without heparin to dilute any
remaining heparin still in solution to a negligible amount. The complex was then diluted
with low salt transcription buffer supplemented with Fos-Choline-8 (Anatrace, final
concentration 1.5 mM). 3 µL of sample was incubated with 0.5 µl of 7x NTP stock solution
(3.5 mM NTPs in low salt transcription buffer plus 1.5 mM Fos-Choline 8) for one minute
(EcoRI* complex) or 30 seconds (RNAP-RNAP) at room temperature before vitrification.
Samples were prepared on glow-discharged (Gatan Solarus, O_2/H_2, 10 seconds) C-Flat
Holey Carbon Film CF-1.2/1.3-4, 400 mesh gold grids at concentrations ranging from 10-
14 µM for RNAP-EcoRI* and ~4 µM for RNAP-RNAP. Grids were loaded into Leica EM
GP apparatus, blotted for 4-5 seconds at 90% humidity at room temperature, and plunge
frozen in liquid ethane.
For the RNAP-EcoRI* complex, the sample concentration was estimated by the
A550 signal (Cy3 extinction coefficient of 150000 M^{-1} cm^{-1}). For the RNAP-RNAP
complex, sample concentration was estimated using A260 and A280 absorbance
measurements using combined extinction coefficients from the individual components.
563

Cryo-EM data acquisition and processing

For RNAP-EcoRI* complex, three different grids were imaged from two different sample preparations of the complex, frozen on different days. Datasets were collected on the same FEI Titan Krios G3i system operating at 300 kV and equipped with a Gatan K3 direct electron detector, BioQuantum energy filter, and a spherical aberration corrector using super-resolution mode. Frame sequences ("movies") were recorded using the SerialEM software[57] suite at a nominal magnification of ×81,000, corresponding to a calibrated pixel size of 0.86 Å at the specimen level (super-resolution pixel size of 0.43 Å per pixel)[58]. Each 1.4 s exposure was dose-fractionated across 35 frames, with a total electron dose of 47.3 $e^- Å^{-2}$ (1.35 $e^- Å^{-2}$ per frame), with defocus values ranging from −0.8 to −2.0 μm underfocus. Beam-image shift was used to collect 43,378 single exposures (4.6k, 17.8k, 23.8k exposures from the different grids) from nine holes in a 3-by-3 grid per each stage translation.

The RNAP-RNAP complex dataset was collected similarly, with the following alterations. Data were collected on a single grid on a FEI Titan Krios G2 operating at 300 kV and equipped with a Gatan K3 direct electron detector and BioQuantum energy filter. Movies were recorded at a nominal magnification of x105,000, corresponding to a calibrated pixel size of 0.847 Å at the specimen level (super-resolution pixel size of 0.4235 Å per pixel)[58]. The total electron dose was 48.79 $e^- Å^{-2}$ (1.39 $e^- Å^{-2}$ per frame). Beam-image shift was used to collect 20,080 single exposures.

Cryo-EM data processing

All processing was performed using cryoSPARC[59]. Movies were aligned with Patch Motion correction (B-factor: 150), and contrast transfer function (CTF) parameters were estimated with Patch CTF using default settings. For each dataset, blob picker was initially used to pick particles from a 256-pixel box, followed by 2D and 3D classification on extracted particles to remove junk particles and RNAPs that did not restart. This initial particle stack was used to generate a preliminary reconstruction that was used to re-pick particles using a template picker approach to yield more particles. All of the picked particles were then re-extracted with a 400-pixel box and separated with multiple rounds of 3D heterogeneous refinement job with candidate 3D volumes as well as junk particle volumes (**Fig. S2, S6**). All models were then further refined via a non-uniform refinement job[60], local CTF refinement, and global CTF refinement. Then low per-particle scale particles were removed, followed by per-particle reference-based motion correction and subsequent local refinement. 3DFlex[30] was then performed (N=5 modes) to improve the density of high flexible regions in the particles and generate variability trajectories for DNA deformation-RNAP swiveling analysis.

For the RNAP-RNAP dataset, the following additional steps were performed. During test dataset processing a 400-pixel box size was used and binned by 4 to enhance processing speed during initial processing steps. Particle box size was increased to 700 pixels (no binning) to accommodate the additional RNAP. After heterogeneous refinement, an additional round of 3D classification was performed using low-resolution volumes as bait volumes (generated by manual docking to mimic RNAPs with 26-36 bp of separation followed by molmap (resolution 20) volume generation from atomic models) using ChimeraX to see if any additional states could be extracted. This resulted in the detection of the 29 bp inter-RNAP class. After non-uniform refinement, models were re-

oriented along their C2 symmetry axis and symmetry expanded before local refinement
and further polishing with a one RNAP sized mask.
To isolate particles that had clear hairpin density in the exit channel, a new set of
particles were extracted from the same micrographs (to remove any potential bias from
the C2 symmetric particles that were able to be resolved to high resolution) and subjected
to a similar processing pipeline to a resolve a single RNAP class (box size 400 pixels)
using a focus mask corresponding to a single RNAP (**Fig. S9**). This class contained
downstream DNA and low-resolution signal further downstream that could correspond to
collided RNAP density being averaged between different states (since the second RNAP
was not included in the focus mask). Taken together, this indicates that these single
RNAPs are still in a post-collision state instead of the alternate hypothesis such as run-
off RNAPs or non-bound RNAPs. These particles were subjected to polishing and
refinement prior to an additional round of focused classification performed with a mask
around the RNAP exit channel that showed one class with clear RNA hairpin density.
Particles contributing to the hairpin-containing class were subjected to final polishing and
refinement. Additionally, to compare the collided states of the hairpin-containing RNAPs,
hairpin-containing particles were re-extracted with a larger box size (700 pixels) and
subjected to 3D classification with a focus mask excluding the high-resolution hairpin-
containing RNAP (N=8 classes). The relative distribution of classes representing either
the 26, 29, 31 bp inter-RNAP distances, or other classes consisting of only a single well-
refined RNAP was then determined via manual inspection of the density maps and
comparison to the 26, 29, and 31 bp maps constructed prior.
632
Model building and refinement
The RNAP-EcoRI* initial model was derived from PDB 6RIP[35]. The model was manually
fit into the cryo-EM density maps using ChimeraX[37], modified in Coot/PyMOL/Isolde[61–63],
and followed by real-space refinement using PHENIX[64]. The nucleic acid sequence was
fixed in the same reference frame as the EcoRI* cognate binding sequence, which
matched the cryo-EM map along the rest of the nucleic acid sequence. For real-space
refinement, rigid body refinement was followed by all-atom refinement with
Ramachandran and secondary structure restraints. For the RNAP-RNAP complexes, the
same refinement methodology was applied with the following modifications. The RNAP
coordinates from the RNAP-EcoRI* complex were used as the starting model for the
RNAP-RNAP complexes. Additionally, since we were unable to determine the nucleic
acid sequence by visual inspection in any of the RNAP-RNAP states (likely due to
heterogeneity in the positioning of collision sites along the template), poly-G and poly-C
sequences were used during modeling building.
647
648 *3DFlex model fitting individual frames*
649 For each frame of the 3DFlex variability trajectories, docking of individual rigid bodies (as
650 defined in **Table S3**) from the consensus model was performed in ChimeraX[37] followed
651 by subsequent unsupervised all-atom refinement in Isolde (5 minutes, 20K, nucleotides
652 restrained by base-pair hydrogen bonding)[63].
653
654 *Swivel domain motion calculation*

The core modules of the different RNAP structures were superimposed in PyMOL by aligning the following the α-carbons: all modeled residues from the α subunit, residues 3:27, 142:152, 445:455, 520:713, 786:828 from the β subunit, residues 343:368, 412:524, 530:552, 569:701, 720:786 from the β' subunit, and all modeled residues from the ω subunit.

RNAP EC (PDB: 6RH3)[35] was used as the reference structure (swivel angle = 0°) for swivel angle measurements. To determine the swivel angle for a given RNAP structure, residues comprising the swivel module from the RNAP structure and the reference were selected (β-subunit AA 1241:1341; β'-subunit AA 1:342, 369:420, 787:930, 1135:1375). The rotation axis and rotation angle transforming one swivel module into the other were then computed using the PyMOL script draw_rotation_axis.py (https://pymolwiki.org/index.php/RotationAxis). For DNA deformation-RNAP swiveling coupling analysis, the same method was applied. For symmetry expanded particles in the head-on collision state, the swivel angle of only the RNAP contained in the refinement mask was analyzed.

DNA displacement calculation
Generation of a frame of reference was necessary to orient downstream DNA displacement on different axes. To accomplish this, the center of masses of twelve downstream nucleotides of the active RNAP EC (PDB: 6rh3)[35] were extracted using PyMOL[62]. PCA was performed on their 3D coordinates to generate the PC1 and PC2 orthoplanes along the long axis of the downstream DNA. Since this definition of PC1 and PC2 is due to inherent variation of the center of mass of the nucleotides downstream of the active site in the reference frame RNAP EC (PDB: 6rh3)[35], the plane definition is arbitrary, meaning that the definition of which axis is most responsible for variance (PC1) is descriptive of the RNAP reference structure variability, not descriptive of the variance between the different structures being compared in this study. For each query structure (consensus structures and 3DFlex frames), the structural cores of the RNAPs were aligned to the structural core of the active RNAP EC (PDB: 6rh3)[35], and displacement of the center of mass of each nucleotide was then projected onto PC1 and PC2, summed across the sequence of twelve nucleotides, and averaged.

Biochemical Assays
RNAP-EcoRI Collided Complex*
<u>Backtracking assays</u>
5' Cy3-labeled RNA oligonucleotides corresponding to zero to four nucleotides of backtracking (based on the RNAP starting position along the nucleic acid sequence) were synthesized (Dharmacon) and used as a ladder to verify the RNA length. For the backtracking assay, the scaffold was assembled as described previously except for omitting the EcoRI* binding step before gel filtration purification. Twenty-fold excess of EcoRI* (if applicable) and 1 μM Gre factors (if applicable) were added to the assembled complex and incubated at RT for 20 minutes before addition of 0.5 mM NTPs for 1 minute in a final volume of 15 μL. To process the RNA samples, 1.5 μL of Turbo DNase (ThermoScientific) was added and incubated for 15 minutes at 37° C, 1.5 μL of proteinase K (New England Biolabs) and 50 mM EDTA for 15 minutes at RT, and finally an equal volume of 2x RNA loading dye (1x components - 47.5% formamide, 0.01% SDS, 0.01%

701 bromophenol blue, 0.5 mM EDTA) was added to the samples. Samples were boiled at
702 95º C for 5 minutes and separated on an 18% TBE-Urea PAGE gel and visualized using
703 Cy3-fluorescence on a Typhoon gel imager. Three biological replicates of this assay were
704 performed on different days.
705
EcoRI* bypass assays
RNAP-EcoRI* containing scaffolds were purified as described previously. Samples were
then diluted into heparin-free buffer to a final concentration of ~200 nM as quantified by
Cy3 RNA fluorescence. Samples were then diluted two-fold into a mixture of low salt
transcription buffer supplemented with varying KOAc concentrations and 0.5 mM NTPs.
These reactions were incubated at 37º C for 30 minutes and then quenched by addition
of 2X RNA loading dye (95% formamide, 0.02% SDS, 0.01% bromophenol blue, 1 mM
EDTA) supplemented with KOAc to equalize the final salt concentration between
conditions. For analysis, the intensity of the roadblock and scaffold band were
background subtracted, measured in FIJI[65], and normalized relative to each other
(roadblock intensity divided by scaffold intensity, and additionally rescaled for each salt
titration gel so that the 50 mM normalized roadblock intensity was set to $E_{RB} = 1$, and the
background intensity was set to $E_{RB} = 0$). E_{RB} calculated at 300 mM KOAc (because that
salt concentration had the largest dynamic range when all the factor addition experiments
were compared) was rescaled so that the average value of the EcoRI* ("normalized E_{RB}")
was equivalent to 1 for each sample condition (factor addition experiments in **Fig. 3C** and
stability mutants experiments in **Fig. 3F**) to adjust for observed batch effects when
different NTP stocks were used.
724
RNAP Head-On Collided Complex
Complex stability assay
RNAP-RNAP head-on complexes were assembled and purified for DNA templates with
and without the YnaJ-UspE hairpin as previously described in low salt transcription buffer
supplemented with 0.1 mg/ml heparin. Samples were concentrated using a 100 kDa
MWCO concentrator (Amicon). Control samples to measure total DNA amount loaded
per well were generated by treating samples with 1 µL proteinase K (New England
Biolabs) per 10 µL reaction volume supplemented with 25 mM EDTA (final concentration)
for over one hour at room temperature. 1 mM NTPs were added to initiate transcription
restart, and aliquots of the reaction were quenched with addition of 25 mM EDTA (final
concentration) at the specified time interval. Samples were run on a 2% 0.5X TBE
agarose gel after being diluted with a 6x stock of 2M sucrose, and the resulting gel was
stained with SYBR safe to visualize the DNA. Lane profiles were extracted using a 0.5x
lane width window. Signal was summed in the 400-1000 bp range, then normalized
between the -NTP control (corresponding to 100% RNAP-RNAP complexes) and the
proteinase K control (corresponding to background signal) to calculate fraction intact
complex. Four different experimental replicates were performed. Bi-exponentials were
fitted through the data using the scipy python module[66] to approximate the half-life of the
collided complex, and 95% confidence intervals are displayed in the plot.
744
SEnd-seq

RNAP-RNAP head-on complexes were assembled and purified as previously described
in low salt transcription buffer supplemented with 0.1 mg/ml heparin. For single promoter
templates, the same general protocol was followed with a polymerase:DNA ratio of 2:1
instead of 4:1 as used for the double promoter templates. Samples were concentrated
using a 100 kDa MWCO concentrator (Amicon). +PK labeled samples were treated with
proteinase K (New England Biolabs) and 25 mM EDTA for over one hour at room
temperature. 1 mM NTPs were added and incubated at RT for 1 minute before quenching
with 25 mM EDTA. For complex pulldown, samples were washed and incubated with
streptavidin M-280 Dynabeads (Thermo Fisher Scientific) before being pulled down via a
magnetic separation rack (Permagen), then washed twice with 200 μL low salt buffer
supplemented with 25 mM EDTA and 0.1 mg/mL heparin. The complexes were then
treated with proteinase K for 1 hour at 37 °C to elute RNAs off the beads followed by
overnight incubation at 4°C. The reaction was purified using 2.2x vol of Agencourt
RNAClean XP beads (Beckman Coulter, A63987). After elution, nucleic acids were
treated with 0.5 μl TURBO DNase (Life Technologies, AM2238) to remove template DNA.
The remaining RNA was further purified via phenol-chloroform-isoamyl alcohol
(Invitrogen, 15593031) extraction (25:24:1, v/v/v) and recovered by ethanol precipitation.
The RNA pellet was resuspended, mixed with 1 μL of spike-in RNA, and subsequently
ligated to a 3' RNA adapter for SEnd-seq library preparation, as previously described[17,46].
Sequencing was performed on an Illumina MiSeq instrument in a paired-end mode (150
nt x2).
Reads were barcode sorted to the correct template and experimental condition.
The separate 5' and 3' end reads of each RNA sequence were aligned to both the forward
and reverse candidate RNA sequence using a local pairwise aligner in the Biopython
module[67]   (match_score=2,   mismatch_score=-1,   open_gap_score=-5,
extend_gap_score=-0.5). Alignment scores were calculated by the summation of the two
best candidate positions. The higher scoring alignment was taken, as long as the distance
between the 5' and 3' ends was greater than 15 nucleotides, as all the RNAs should be
at least 17 nucleotides from the A/G/UTP only transcription initiation walkout before
subsequent purification. Unmapped reads either had the wrong direction of 5' and 3'
alignment or were too short. Furthermore, RNAs that did not initiate from within a 10-
nucleotide range near the transcription start site were filtered out. These lost reads upon
filtering could come from RNAP initiation from the end of the DNA template, RNA
degradation of the 5' end, or non-templated addition at the 3' end[68].
780
781   **Molecular graphics and data analysis**
782   Molecular graphics were prepared with UCSF ChimeraX[37]. Plotting and statistical
783   analysis were performed with Python using the matplotlib module[69]. Python codes were
784   prepared with the assistance of ChatGPT 4.0.
785
786   **Statistical analysis**
787   Conditions were compared via two-tailed, unpaired t-test including Welch's correction. If
788   applicable, a Benjamini-Hochberg correction for multiple hypothesis testing was added.
789   S.D. stands for standard deviation and S.E. stands for standard error when used in the
790   text. P-value asterisk reported as *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.
791

Acknowledgements

We thank Johanna Sotiris, Honkit Ng, and Mark Ebrahim at the Rockefeller University Evelyn Gruss Lipper Cryo-EM Resource Center for their support in data collection. The expression plasmids for Rho, NusA, NusG, GreA and GreB were provided by Rachel Anne Mooney (Landick Laboratory, University of Wisconsin-Madison). NusA, NusG, GreA, and GreB proteins were purified by Rui Gong (Alushin Laboratory, Rockefeller University). S.L. acknowledges support from the National Institute of Health (R01GM149862). S.L. and G.M.A acknowledge support from the Stavros Niarchos Foundation Institute for Global Infectious Disease Research at the Rockefeller University and a joint award from the Alfred P. Sloan Foundation Matter-to-Life Program (G-2024-22709).

Author contributions

J.W.W., S.L., and G.M.A. designed research. J.W.W. purified proteins, collected cryo-EM datasets, and performed biochemical studies. A.U.M. purified proteins and helped with biochemical experimental design. A.U.M. and S.A.D. helped with cryo-EM processing and data interpretation. X.J. performed SEnd-Seq experiments. S.J.C. performed biochemical assays for the RNAP-EcoRI* complex. H.J.Y. performed biochemical assays for the RNAP-RNAP collided complex and collected the RNAP-RNAP collided cryo-EM dataset. J.W.W., S.L., and G.M.A. wrote the paper with input from all other authors.

Data availability

Cryo-EM density maps and corresponding atomic models have been deposited in the PDB and EMDB with the following accession codes: RNAP-EcoRI* Collision (PDB: XXXX, EMDB: YYYYY); Head-on RNAP-RNAP Collision: 26 bp distance (PDB: XXXX, EMDB: YYYYY); Head-on RNAP-RNAP Collision: 29 bp distance (PDB: XXXX, EMDB: YYYYY); Head-on RNAP-RNAP Collision: 31 bp distance (PDB: XXXX, EMDB: YYYYY); Head-on RNAP-RNAP Collision: RNA hairpin (PDB: XXXX, EMDB: YYYYY). 3DFlex volumes and fit models for both consensus and variability trajectory movies have been deposited in zenodo (doi: 10.5281/zenodo.19317100).

Competing interests

The authors declare no competing interests.

References

1. Le, T.T., and Wang, M.D. (2018). Molecular Highways—Navigating Collisions of DNA Motor Proteins. J. Mol. Biol. *430*, 4513–4524. https://doi.org/10.1016/j.jmb.2018.08.006.

2. Sneppen, K., Dodd, I.B., Shearwin, K.E., Palmer, A.C., Schubert, R.A., Callen, B.P., and Egan, J.B. (2005). A Mathematical Model for Transcriptional Interference by RNA Polymerase Traffic in Escherichia coli. J. Mol. Biol. *346*, 399–409. https://doi.org/10.1016/j.jmb.2004.11.075.

3. Klumpp, S., and Hwa, T. (2008). Stochasticity and traffic jams in the transcription of ribosomal RNA: Intriguing role of termination and antitermination. Proc. Natl. Acad. Sci. U. S. A. *105*, 18159–18164. https://doi.org/10.1073/pnas.0806084105.

4. Helmrich, A., Ballarino, M., and Tora, L. (2011). Collisions between Replication and Transcription Complexes Cause Common Fragile Site Instability at the Longest Human Genes. Mol. Cell *44*, 966–977. https://doi.org/10.1016/j.molcel.2011.10.013.

5. Sankar, T.S., Wastuwidyaningtyas, B.D., Dong, Y., Lewis, S.A., and Wang, J.D. (2016). The nature of mutations induced by replication–transcription collisions. Nature 2016 535:7610 *535*, 178–181. https://doi.org/10.1038/nature18316.

6. Prado, F., and Aguilera, A. (2005). Impairment of replication fork progression mediates RNA polII transcription-associated recombination. EMBO Journal *24*, 1267–1276. https://doi.org/10.1038/sj.emboj.7600602.

7. Pomerantz, R.T., and O'Donnell, M. (2008). The replisome uses mRNA as a primer after colliding with RNA polymerase. Nature *456*, 762. https://doi.org/10.1038/nature07527.

8. Zhang, H., Shi, Z., Banigan, E.J., Kim, Y., Yu, H., Bai, X.-C., and Finkelstein Correspondence, I.J. (2023). CTCF and R-loops are boundaries of cohesin-mediated DNA looping. Mol. Cell *83*, 2856-2871.e8. https://doi.org/10.1016/j.molcel.2023.07.006.

9. Epshtein, V., and Nudler, E. (2003). Cooperation between RNA polymerase molecules in transcription elongation. Science (1979). *300*, 801–805. https://doi.org/10.1126/science.1083219.

10. Liang Meng Wee, A., Tong, A.B., Jose Florez Ariza, A., Cañ ari-Chumpitaz, C., Grob, P., Nogales, E., Bustamante, C.J., and Meng Wee, L. (2023). A trailing ribosome speeds up RNA polymerase at the expense of transcript fidelity via force and allostery. Cell *186*, 1244-1262.e34. https://doi.org/10.1016/j.cell.2023.02.008.

11. Wang, C., Molodtsov, V., Firlar, E., Kaelber, J.T., Blaha, G., Su, M., and Ebright, R.H. (2020). Structural basis of transcription-translation coupling. Science (1979). *369*, 1359–1365. https://doi.org/10.1126/SCIENCE.ABB5317.

12. Webster, M.W., Takacs, M., Zhu, C., Vidmar, V., Eduljee, A., Abdelkareem, M., and Weixlbaumer, A. (2020). Structural basis of transcription-translation coupling and collision in bacteria. Science (1979). *369*, 1355–1359. https://doi.org/10.1126/SCIENCE.ABB5036.

13. Zhang, J., Lu, G., Zhou, W., Molodtsov, V., Cheng, Z., Zhang, Y., Li, R., Jiang, Z., Yang, M., Li, L., et al. (2026). Structural basis of translation in transcription-translation coupling. bioRxiv, 2024.03.13.584796. https://doi.org/10.1101/2024.03.13.584796.

14. Proshkin, S., Rachid Rahmouni, A., Mironov, A., and Nudler, E. (2010). Cooperation between translating ribosomes and RNA polymerase in transcription elongation. Science (1979). *328*, 504–508. https://doi.org/10.1126/science.1184939.

15. Epshtein, V., Toulmé, F., Rachid Rahmouni, A., Borukhov, S., and Nudler, E. (2003). Transcription through the roadblocks: the role of RNA polymerase cooperation. EMBO J. *22*, 4719. https://doi.org/10.1093/emboj/cdg452.

16. Wang, L., Watters, J.W., Ju, X., Lu, G., and Liu, S. (2023). Head-on and co-directional RNA polymerase collisions orchestrate bidirectional transcription termination. Mol. Cell *83*, 1153-1164.e4. https://doi.org/10.1016/j.molcel.2023.02.017.

17. Ju, X., Li, D., and Liu, S. (2019). Full-length RNA profiling reveals pervasive bidirectional transcription terminators in bacteria HHS Public Access. Nat Microbiol *4*, 1907–1918. https://doi.org/10.1038/s41564-019-0500-z.

18. Abbondanzieri, E.A., Greenleaf, W.J., Shaevitz, J.W., Landick, R., and Block, S.M. (2005). Direct observation of base-pair stepping by RNA polymerase. Nature *438*, 460. https://doi.org/10.1038/nature04268.

19. Ma, J., Bai, L., and Wang, M.D. (2013). Transcription under torsion. Science (1979). *340*, 1580–1583. https://doi.org/10.1126/science.1235441.

20. Yin, H., Wang, M.D., Svoboda, K., Landick, R., Block, S.M., and Gelles, J. (1995). Transcription against an applied force. Science (1979). *270*, 1653–1657. https://doi.org/10.1126/science.270.5242.1653.

21. Wang, M.D., Schnitzer, M.J., Yin, H., Landick, R., Gelles, J., and Block, S.M. (1998). Force and velocity measured for single molecules of RNA polymerase. Science (1979). *282*, 902–907. https://doi.org/10.1126/science.282.5390.902.

22. Galburt, E.A., Grill, S.W., Wiedmann, A., Lubkowska, L., Choy, J., Nogales, E., Kashlev, M., and Bustamante, C. (2007). Backtracking determines the force sensitivity of RNAP II in a factor-dependent manner. Nature *446*, 820–823. https://doi.org/10.1038/nature05701.

23. Ma, J., Tan, C., Gao, X., Fulbright, R.M., Roberts, J.W., and Wang, M.D. (2019). Transcription factor regulation of RNA polymerase's torque generation capacity. Proc. Natl. Acad. Sci. U. S. A. *116*, 2583–2588. https://doi.org/10.1073/pnas.1807031116.

24. Qian, J., Cartee, A., Xu, W., Yan, Y., Wang, B., Artsimovitch, I., Dunlap, D., and Finzi, L. (2024). Reciprocating RNA Polymerase batters through roadblocks. Nature Communications 2024 15:1 *15*, 1–11. https://doi.org/10.1038/s41467-024-47531-x.

25. Crampton, N., Bonass, W.A., Kirkham, J., Rivetti, C., and Thomson, N.H. (2006). Collision events between RNA polymerases in convergent transcription studied by atomic force microscopy. Nucleic Acids Res. *34*, 5416. https://doi.org/10.1093/NAR/GKL668.

26. Wright, D.J., King, K., and Modrich, P. (1989). The Negative Charge of Glu-11 1 Is Required to Activate the Cleavage Center of EcoRI Endonuclease. Journal of Biological Chemistry *264*, 11816–11821. https://doi.org/10.1016/S0021-9258(18)80138-7.

914   27.   Finkelstein, I.J., Visnapuu, M.-L., and Greene, E.C. (2010). Single-molecule imaging
915   reveals mechanisms of protein disruption by a DNA translocase. Nature *468*, 983–
916   987. https://doi.org/10.1038/nature09561.

917   28.   Frederick, C.A., Grable, J., Melia, M., Samudzi, C., Jen-Jacobson, L., Wang, B.C.,
918   Greene, P., Boyer, H.W., and Rosenberg, J.M. (1984). Kinked DNA in crystalline
919   complex with EcoRI endonuclease. Nature 1984 309:5966 *309*, 327–331.
920   https://doi.org/10.1038/309327a0.

921   29.   Doughty, B., Kazer, S.W., and Eisenthal, K.B. (2011). Binding and cleavage of DNA with
922   the restriction enzyme EcoR1 using time-resolved second harmonic generation. Proc.
923   Natl. Acad. Sci. U. S. A. *108*, 19979–19984.
924   https://doi.org/10.1073/PNAS.1115498108.

925   30.   Punjani, A., and Fleet, D.J. (2023). 3DFlex: determining structure and motion of flexible
926   proteins from cryo-EM. Nat. Methods *20*, 860. https://doi.org/10.1038/S41592-023-
927   01853-8.

928   31.   Komissarova, N., and Kashlev, M. (1997). RNA polymerase switches between
929   inactivated and activated states by translocating back and forth along the DNA and
930   the RNA. Journal of Biological Chemistry *272*, 15329–15338.
931   https://doi.org/10.1074/jbc.272.24.15329.

932   32.   Nudler, E., Kashlev, M., Nikiforov, V., and Goldfarb, A. (1995). Coupling between
933   transcription termination and RNA polymerase inchworming. Cell *81*, 351–357.
934   https://doi.org/10.1016/0092-8674(95)90388-7.

935   33.   Borukhov, S., Sagitov, V., and Goldfarb, A. (1993). Transcript cleavage factors from E.
936   coli. Cell *72*, 459–466. https://doi.org/10.1016/0092-8674(93)90121-6.

937   34.   Zhang, J., and Landick, R. (2009). Substrate Loading, Nucleotide Addition, and
938   Translocation by RNA Polymerase. RNA Polymerases as Molecular Motors, 206–235.
939   https://doi.org/10.1039/9781847559982-00206.

940   35.   Abdelkareem, M., Saint-André, C., Takacs, M., Papai, G., Crucifix, C., Guo, X., Ortiz,
941   J., and Weixlbaumer, A. (2019). Structural Basis of Transcription: RNA Polymerase
942   Backtracking and Its Reactivation. Mol. Cell *75*, 298-309.e4.
943   https://doi.org/10.1016/j.molcel.2019.04.029.

944   36.   Guo, X., Myasnikov, A.G., Chen, J., Takacs, M., Schultz, P., and Weixlbaumer, A.
945   (2018). Structural Basis for NusA Stabilized Transcriptional Pausing. Mol. Cell *69*,
946   816–827. https://doi.org/10.1016/j.molcel.2018.02.008.

947   37.   Meng, E.C., Goddard, T.D., Pettersen, E.F., Couch, G.S., Pearson, Z.J., Morris, J.H.,
948   and Ferrin, T.E. (2023). UCSF ChimeraX: Tools for structure building and analysis.
949   Protein Science *32*, e4792. https://doi.org/10.1002/pro.4792.

950   38.   Kang, J.Y., Mishanina, T. V., Landick, R., and Darst, S.A. (2019). Mechanisms of
951   transcriptional pausing in bacteria. J. Mol. Biol. *431*, 4007.
952   https://doi.org/10.1016/J.JMB.2019.07.017.

953   39.   Kang, J.Y., Mishanina, T. V., Bellecourt, M.J., Mooney, R.A., Darst, S.A., and Landick,
954   R. (2018). RNA polymerase accommodates a pause RNA hairpin by global
955   conformational rearrangements that prolong pausing. Mol. Cell *69*, 802.
956   https://doi.org/10.1016/j.molcel.2018.01.018.

40. Delbeau, M., Omollo, E.O., Froom, R., Koh, S., Mooney, R.A., Lilic, M., Brewer, J.J.,
Rock, J., Darst, S.A., Campbell, E.A., et al. (2023). Structural and functional basis of
the universal transcription factor NusG pro-pausing activity in Mycobacterium
tuberculosis. Mol. Cell *83*, 1474-1488.e8.
https://doi.org/10.1016/j.molcel.2023.04.007.

41. Kang, J.Y., Mishanina, T. V., Bao, Y., Chen, J., Llewellyn, E., Liu, J., Darst, S.A., and
Landick, R. (2023). An ensemble of interconverting conformations of the elemental
paused transcription complex creates regulatory options. Proc. Natl. Acad. Sci. U. S.
A. *120*, e2215945120. https://doi.org/10.1073/PNAS.2215945120.

42. Kim, Y., Grable, J.C., Love, R., Greene, P.J., and Rosenberg, J.M. (1990). Refinement of
Eco RI endonuclease crystal structure: A revised protein chain tracing. Science
(1979). *249*, 1307–1309. https://doi.org/10.1126/science.2399465.

43. Artsimovitch, I., and Landick, R. (2000). Pausing by bacterial RNA polymerase is
mediated by mechanistically distinct classes of signals. Proc. Natl. Acad. Sci. U. S. A.
97, 7090–7095. https://doi.org/10.1073/pnas.97.13.7090.

44. Zhu, C., Guo, X., Dumas, P., Takacs, M., Abdelkareem, M., Vanden Broeck, A., Saint-
André, C., Papai, G., Crucifix, C., Ortiz, J., et al. (2022). Transcription factors modulate
RNA polymerase conformational equilibrium. Nature Communications 2022 13:1 *13*,
1546-. https://doi.org/10.1038/s41467-022-29148-0.

45. Jen-Jacobson, L., Lesser, D., and Kurpiewski, M. (1986). The enfolding arms of EcoRI
endonuclease: Role in DNA binding and cleavage. Cell *45*, 619–629.
https://doi.org/10.1016/0092-8674(86)90294-1.

46. Ju, X., Li, S., Froom, R., Wang, L., Lilic, M., Delbeau, M., Campbell, E.A., Rock, J.M.,
and Liu, S. (2024). Incomplete transcripts dominate the Mycobacterium tuberculosis
transcriptome. Nature 2024 627:8003 *627*, 424–430. https://doi.org/10.1038/s41586-
024-07105-9.

47. Shundrovsky, A., Santangelo, T.J., Roberts, J.W., and Wang, M.D. (2004). A Single-
Molecule Technique to Study Sequence-Dependent Transcription Pausing. Biophys.
J. *87*, 3945. https://doi.org/10.1529/biophysj.104.044081.

48. Zhang, M., Cañari-Chumpitaz, C., Liu, J., Onoa, B., Cleir, S. de, Cheng, E., Requejo,
K.I., and Bustamante, C. (2026). Apical Localization of RNA Polymerases Modulate
Transcription Dynamics and Supercoiling Domains Revealed by Cryo-ET. bioRxiv,
2026.03.25.714350. https://doi.org/10.64898/2026.03.25.714350.

49. Chen, J., Gopalkrishnan, S., Chiu, C., Chen, A.Y., Campbell, E.A., Gourse, R.L., Ross,
W., and Darst, S.A. (2019). E. Coli TraR allosterically regulates transcription initiation
by altering RNA polymerase conformation. Elife *8*.
https://doi.org/10.7554/eLife.49375.

50. Qian, J., Lubkowska, L., Zhang, S., Tan, C., Hong, Y., Jia, X., Fulbright, R.M., Inman, J.T.,
Kay, T.M., Jeong, J., et al. (2025). Chromatin buffers torsional stress during
transcription. Science (1979). https://doi.org/10.1126/science.adv0134.

51. Bradford, M.M. (1976). A rapid and sensitive method for the quantitation of microgram
quantities of protein utilizing the principle of protein-dye binding. Anal. Biochem. *72*,
248–254. https://doi.org/10.1016/0003-2697(76)90527-3.

52. Perederina, A.A., Vassylyeva, M.N., Berezin, I.A., Svetlov, V., Artsimovitch, I., and
Vassylyev, D.G. (2005). Cloning, expression, purification, crystallization and initial
crystallographic analysis of transcription elongation factors GreB from Escherichia
coli and Gfh1 from Thermus thermophilus. urn:issn:1744-3091 *62*, 44–46.
https://doi.org/10.1107/S1744309105040297.

53. Hao, Z., Epshtein, V., Kim, K.H., Proshkin, S., Svetlov, V., Kamarthapu, V., Bharati, B.,
Mironov, A., Walz, T., and Nudler, E. (2021). Pre-termination Transcription Complex:
Structure and Function. Mol. Cell *81*, 281-292.e8.
https://doi.org/10.1016/j.molcel.2020.11.013.

54. Schmidt, M.C., and Chamberlin, M.J. (1987). nusA Protein of Escherichia coli is an
efficient transcription termination factor for certain terminator sites. J. Mol. Biol. *195*,
809–818. https://doi.org/10.1016/0022-2836(87)90486-4.

55. Burgess, R.R., Travers, A.A., Dunn, J.J., and Bautz, E.K.F. (1969). Factor Stimulating
Transcription by RNA Polymerase. Nature 1969 221:5175 *221*, 43–46.
https://doi.org/10.1038/221043a0.

56. Kang, J.Y., Llewellyn, E., Chen, J., Olinares, P.D.B., Brewer, J., Chait, B.T., Campbell,
E.A., and Darst, S.A. (2021). Structural basis for transcription complex disruption by
the Mfd translocase. Elife *10*, 1–86. https://doi.org/10.7554/eLife.62117.

57. Mastronarde, D.N. (2003). SerialEM: A Program for Automated Tilt Series Acquisition
on Tecnai Microscopes Using Prediction of Specimen Position. Microscopy and
Microanalysis *9*, 1182–1183. https://doi.org/10.1017/s1431927603445911.

58. Mastronarde, D.N. (2005). Automated electron microscope tomography using robust
prediction of specimen movements. J. Struct. Biol. *152*, 36–51.
https://doi.org/10.1016/j.jsb.2005.07.007.

59. Punjani, A., Rubinstein, J.L., Fleet, D.J., and Brubaker, M.A. (2017). cryoSPARC:
algorithms for rapid unsupervised cryo-EM structure determination. Nature Methods
2017 14:3 *14*, 290–296. https://doi.org/10.1038/nmeth.4169.

60. Punjani, A., Zhang, H., and Fleet, D.J. (2020). Non-uniform refinement: adaptive
regularization improves single-particle cryo-EM reconstruction. Nature Methods 2020
17:12 *17*, 1214–1221. https://doi.org/10.1038/s41592-020-00990-8.

61. Emsley, P., Lohkamp, B., Scott, W.G., and Cowtan, K. (2010). Features and
development of Coot. Acta Crystallogr. D Struct Biol. *66*, 486–501.
https://doi.org/10.1107/S0907444910007493.

62. Schrödinger LLC (2015). The PyMOL Molecular Graphics System, Version~1.8.

63. Croll, T.I. (2018). ISOLDE: A physically realistic environment for model building into
low-resolution electron-density maps. Acta Crystallogr. D Struct. Biol. *74*, 519–530.
https://doi.org/10.1107/S2059798318002425.

64. Liebschner, D., Afonine, P. V., Baker, M.L., Bunkoczi, G., Chen, V.B., Croll, T.I., Hintze,
B., Hung, L.W., Jain, S., McCoy, A.J., et al. (2019). Macromolecular structure
determination using X-rays, neutrons and electrons: recent developments in Phenix.
Acta Crystallogr. D Struct Biol. *75*, 861–877.
https://doi.org/10.1107/S2059798319011471.

65. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source

platform for biological-image analysis. Nature Methods 2012 9:7 *9*, 676–682.
https://doi.org/10.1038/nmeth.2019.

66.   Virtanen, P., Gommers, R., Oliphant, T.E., Haberland, M., Reddy, T., Cournapeau, D.,
Burovski, E., Peterson, P., Weckesser, W., Bright, J., et al. (2020). SciPy 1.0:
fundamental algorithms for scientific computing in Python. Nature Methods 2020
17:3 *17*, 261–272. https://doi.org/10.1038/s41592-019-0686-2.

67.   Cock, P.J.A., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg, I.,
Hamelryck, T., Kauff, F., Wilczynski, B., et al. (2009). Biopython: freely available
Python tools for computational molecular biology and bioinformatics. Bioinformatics
25, 1422–1423. https://doi.org/10.1093/bioinformatics/btp163.

68.   Galls, D., Mueller, A.U., Greenwald, E., and Fire, A.Z. (2025). A general RNA-templated
RNA extension activity of E. coli RNA polymerase. RNA *31*, 663–678.
https://doi.org/10.1261/rna.080238.124.

69.   Hunter, J.D. (2007). Matplotlib: A 2D graphics environment. Comput. Sci. Eng. *9*, 90–
95. https://doi.org/10.1109/MCSE.2007.55.

1059
1060

1061
1062
Figure 1. Reconstituting transcriptional collision with a stationary roadblock.
[A] Synthetic nucleic acid scaffold design to assemble E. coli RNAP and EcoRI* onto a
DNA template. NT-DNA, non-template DNA strand; T-DNA, template DNA strand.
[B] Scaffold assembly and purification scheme.
[C] Representative denaturing gel showing blocked transcription in the presence of
EcoRI*, suggesting collided complex assembly. Cy3-labeled RNA was visualized.
[D] Cryo-EM density map (top) and atomic model (bottom) of the RNAP-EcoRI* collided
complex.

1071
1072
Figure 2. EcoRI* collided complex is backtracked and moderately swiveled.
[A] Representative gel of in vitro transcription assay in the presence and absence of Gre
factors, with an RNA ladder corresponding to potential transcript lengths within the
collided complex.

[B] View of the active site of the collided complex (cryo-EM map and atomic model).
Resolvable backtracked bases are highlighted in pink.
[C] View of the swivel module of an active elongation complex (PDB: 6RH3) versus that
of the EcoRI* collided complex, superimposed on active EC structure after alignment,
highlighting differential swiveling.
[D] Swivel angle magnitudes of prior RNAP structures versus that of the RNAP-EcoRI*
collided complex. The value for the consensus reconstruction is displayed as a solid dot,
while the range measured across all five 3D flex trajectories for the collided complex is
displayed as a dashed line.
[E] Initial and final frames of the trajectory displayed in **F**, highlighting DNA displacement.
[F] Swivel angle and DNA displacement measured for each frame of the first series
(accounting for the largest conformational movements) of the 3DFlex variability trajectory
for the collided complex.

1090
1091
Figure 3. Factors that affect bypass efficiency.
[A] Representative EcoRI* bypass assay. The RNA lengths under different conditions are
visualized via Cy3 fluorescence.
[B] Fraction of blocked RNAP in the absence and presence of factors (GreB, NusA,
NusG). Raw measurements and their sigmoidal fit for each condition are displayed.
[C] Normalized roadblock efficiencies in the presence of indicated factors at 300 mM
KOAc. Data are shown as mean ± S.D. Individual points are also displayed (N = 6).
[D] Views highlighting the orientation of EcoRI* N-terminal helix in the collided complex
versus in the EcoRI* crystal structure (PDB 1ERI). The helix of the collided EcoRI*
protomer is flipped and re-oriented toward the downstream dsDNA, which may be
stabilized by NTE-DNA interactions, while the distal, non-collided protomer (right) adopts
a conformation similar to the crystal structure.
[E] Positions of single alanine mutations are highlighted on the crystal structure overlay.

[F] Normalized roadblock efficiencies for the destabilizing EcoRI* mutations at 300 mM
KOAc. Data are shown as mean ± S.D. Individual points are also displayed (N = 6).
Conditions were compared via two-tailed, unpaired t test including Welch's correction and
Benjamini-Hochberg correction for multiple testing. n.s., not significant; $*p < 0.05$; $**p <$
0.01; $***p < 0.001$.

1110
1111
Figure 4. Reconstituting head-on RNAP-RNAP collisions.
[A] DNA template featuring two promoters, designed to stall RNAP at cytosine sites. F
RNA, forward RNA; R RNA, reverse RNA. Underlined segments of the YnaJ-UspE hairpin
indicate complementary regions.

[B] Experimental scheme for head-on RNAP-RNAP collided complex assembly and
purification.
[C] Native gel shift assay for evaluating collided complex formation.
[D-O] Columns correspond to 26-bp (left), 29-bp (middle), and 31-bp (right) DNA inter-
RNAP distance complexes. **[D-F]** Top: Cartoons indicating the relative orientation
(arrows) and position of polymerases. Bottom: Surface representations of the
corresponding atomic models. **[G-I]** Zoom-in views of the inter-RNAP interface for each
class. **[J-L]** Views of the swivel module for each class (from symmetry-expanded
reconstructions) superimposed on an active EC structure after alignment. **[M-O]** Zoom-in
views of the backtracked RNA (btRNA). Cryo-EM maps and atomic models are displayed.

1126
1127
Figure 5. RNA hairpin impacts collision positioning and stability.
[A, B] SEnd-seq profiles of forward and reverse RNAs in head-on RNAP-RNAP collided
complexes on a template encoding the YnaJ-UspE hairpin (**A**) or a no-hairpin template
(**B**). The grey box defines the hairpin sequence region or the corresponding scrambled
sequence region.
[C] Quantification of RNA lengths (mapped back to the originating promoter). N = 1789
(hairpin F), 8075 (no hairpin F), 608 (hairpin R), and 5945 (no hairpin R). Arrows indicate
the predicted RNA lengths if transcription terminates right after the hairpin. Conditions
were compared with a two-tailed, unpaired t test with Welch's correction. ***p < 0.001.

[D] Native agarose gels for evaluating collided complex disassembly. PK: Proteinase K
treated. Timepoints indicate time elapsed after NTP addition, followed by EDTA quench.
Left: hairpin-containing template DNA; right, no-hairpin template DNA. DNA was
visualized with SYBR safe.
[E] Quantification of the fraction of intact collided complex under indicated conditions.
Nucleic acid intensity values were normalized for each of four experimental replicates to
the -NTPs and +Proteinase K conditions. Shaded areas indicate 95% confidence interval
of the fits.
[F] View of the hairpin-containing head-on collided RNAP structure obtained through
symmetry expansion.
[G] Views of the hairpin-containing RNAP (cryo-EM map and atomic model) highlighting
the backtracked RNA (top) and rotation of the swivel module relative to the active
elongation complex.
[H] Class distributions of no-hairpin and hairpin-containing collided RNAP complexes.
"Other" is defined as particles that show well-defined density for one RNAP but the density
downstream of the well-defined RNAP is too low of a resolution to confidently model an
RNAP. Significance was calculated via a chi-squared test. ***$p < 0.001$.

1154
1155
Figure 6. Inter-RNAP distance modulates deformation-swivel coupling.
[A] Downstream DNA deformation of each collision scenario compared against active EC
(PDB 6RH3).
[B-C] Plots of swiveling angle versus DNA displacement along **[B]** PC1 and **[C]** PC2 of
consensus collision structures (this study) and previously reported structures of an active
EC (PDB 6RH3), backtracked ECs (unswiveled, PDB 6RI9; swiveled, PDB 6RIP), and his
pause EC (PDB 6ASX). Linear fits are displayed (dashed lines).
[D] Plot of the magnitudes of swiveling and PC2 displacement measured in individual
3DFlex series trajectories for the four collided structures (EcoRI*, 26 bp, 29 bp, 31 bp –
5 series each). A linear fit is displayed (dashed line).

Figure 7. Structural mechanisms of transcriptional collisions.
[A] Common features of all collisions in this study. Regardless of the roadblock identity, RNAP backtracks away from the collision site and swivels into a transcriptionally inactive state, mediated by coupling between template DNA distortion and RNAP swiveling.
[B, C] Unique features of different collisions. **[B]** RNAP collision with a model stationary roadblock (EcoRI*) features roadblock remodeling, a plausible mechanism for how 'battering' (repetitive transcriptional restart and collision) can facilitate roadblock bypass.
[C] Dynamic RNAP-RNAP collisions feature substantial heterogeneity due to the stochastic transcriptional and backtracking activity of both polymerases. These dynamics can be tempered by an RNA hairpin that constrains the collision positions.